Anterior cingulate cortex differently modulates fronto-parietal functional connectivity between

resting-state and working memory tasks

Running title: ACC modulates fronto-parietal connectivity

Xin Di [1,2], Heming Zhang [1], Bharat B Biswal [1,2] *

1, School of Life Sciences and Technology, University of Electronic Science and Technology of China,

Chengdu, China

2, Department of Biomedical Engineering, New Jersey Institute of Technology, Newark, NJ, 07029, USA

* Corresponding author:

Bharat B. Biswal, PhD

607 Fenster Hall, University Height

Newark, NJ, 07102, USA

bbiswal@yahoo.com

Abstract

Fronto-parietal regions and the functional communications between them are critical in supporting

working memory and other executive functions. The functional connectivity between fronto-parietal

regions are modulated by working memory loads, and are also shown to be modulated by a third region in

brain in resting-state. However, it is largely unknown that whether the third-region modulations remain

the same during working memory tasks or were largely modulated by task demands. In the current study,

we collected functional MRI (fMRI) data when the subjects were performing n-back tasks and in resting-

state. We first used a block-designed localizer to define fronto-parietal regions that showed higher

activations in the 2-back than the 1-back condition. Next, we performed physiophysiological interaction

(PPI) analysis using left or right middle frontal gyrus (MFG) and superior parietal lobule (SPL) regions,

respectively, in three continuous-designed runs of resting-state, 1-back, and 2-back conditions. No

regions showed consistent modulatory interactions with the seed pairs in the three conditions. Instead, the

anterior cingulate cortex (ACC) showed different modulatory interactions with the right MFG and SPL

among the three conditions. While increased activity of the ACC was associated with decreased

functional coupling between the right MFG and SPL in resting-state, it was associated with increased

functional coupling between them in the 2-back condition. The observed task modulations support the

functional significance of the modulations of the ACC on fronto-parietal connectivity.

Keywords: anterior cingulate cortex; higher-order brain connectivity; modulatory interaction;

physiophysiological interaction; working memory.

1. Introduction

Working memory involves distributed brain regions, most prominently the bilateral fronto-parietal network (Barch et al., 2013; Mencarelli et al., n.d.; Owen, McMillan, Laird, & Bullmore, 2005). Understanding the functional connectivity among the distributed regions is critical to understand the implementation of working memory in brain. The bilateral fronto-parietal regions showed high correlations even in resting-state, thus forming lateralized fronto-parietal networks when using data driven methods such as independent component analysis (ICA) (Beckmann, DeLuca, Devlin, & Smith, 2005; Biswal et al., 2010; Di & Biswal, 2013). Because of the presence of functional connectivity during resting-state, it would be more critical to investigate the changes of functional connectivity during working memory tasks. Electroencephalogram (EEG) studies typically show increased connectivity in the theta band and reduced connectivity in the alpha band between fronto-parietal regions (Babiloni et al., 2004; Dai et al., 2017; Sauseng, Klimesch, Schabus, & Doppelmayr, 2005). As blood-oxygen-level dependent (BOLD) signals measured by functional MRI (fMRI), the signal synchronizations between some of the fronto-parietal regions were found to be reduced during working memory condition compared with control condition, although these regions were more activated during the higher working memory load condition (Di & Biswal, 2019).

In addition to task modulations, functional connectivity between two regions might also be modulated by a third region (Di & Biswal, 2015a; Friston et al., 1997). In the context of working memory, some executive or distractive signals from other brain region might facilitate or disrupt the functional communications between fronto-parietal regions. This will result in higher order interactions among three brain regions, which can be studied using physiophysiological interaction (PPI) model (Di & Biswal, 2013; Friston et al., 1997) or nonlinear dynamic causal modeling (Stephan et al., 2008). Several studies have been performed to characterize the modulatory interactions in resting-state (Di & Biswal, 2013, 2014, 2015a, 2015b). Specifically, we defined the fronto-parietal regions of interest (ROIs) by using ICA and performed PPI analysis on the left or right fronto-parietal ROIs, respectively (Di & Biswal, 2013). We identified several medial frontal and parietal regions that showed negative modulatory

interaction with the fronto-parietal ROIs, indicating that the increases of activity of these regions are

accompanied by reduced fronto-parietal functional connectivity. However, this analysis was only

performed in resting-state data. It is unclear whether similar effects would be shown in task conditions,

or it could alter significantly upon task demands.

The goal of the current study is to examine whether modulatory interactions of the fronto-parietal

regions are modulated by task demands. We adopted a n-back paradigm with varying working memory

loads where the bilateral fronto-parietal regions are consistently activated (Barch et al., 2013; Owen et al.,

2005). We first used a block-designed localizer to identify the fronto-parietal regions that showed higher

activations during the 2-back than the 1-back condition. We then performed PPI analysis by using the

frontal and parietal ROIs in three separate continuous task conditions, i.e. resting-state, 1-back, and 2-

back conditions. We examined two competing hypotheses. First, there are modulatory interactions of a

third region with the two ROIs, and the effects are consistent across the conditions. In contrast, there may

be modulatory interactions of a third region with the two ROIs, but the effects highly depend on the task

conditions. We performed conjunction analysis to identify brain regions that may fulfill the first

hypothesis, and performed repeated measure one-way ANOVA to find regions that may fulfill the second

hypothesis.

2. Methods

2.1. Subjects

Fifty participants (26 females) were recruited for the current study. The mean age was 22.34 years (19 –

24 years, SD = 1.303). One subject was removed because of large head motion during MRI scan. All

participants reported normal auditory and normal or corrected-to-normal visual acuity, and were free of

neurological or psychiatric problems. All study procedures were carried out with written informed

consent of each subject. Each subject received honorarium of 200 RMB for the participation. The study

was approved by institutional review board.

2.2. Study procedure

At the beginning of the MRI scan session, the participants underwent a resting-state fMRI scan (8 min 30

sec). The participants were instructed to lay still with eyes open and staring at a white cross fixation on a

dark background. Four working memory task runs were then performed with the following order: two

block-designed runs with both 1-back and 2-back condition in each run (3 min 46 sec each), one

continuous run of 1-back condition (5 min 10 sec), and one continuous run of 2-back condition (5 min 10

sec). The participants also underwent a few other tasks, which were not relevant to the current study.

Lastly, a high resolution anatomical T1-weighted MRI was scanned at the end of the MRI session.

2.2.1. N-back task

The N-back task tests the participants' working memory of the spatial locations of the letters presented on

the screen. A white cross fixation was presented at the center of the dark screen throughout the

experiment. A random letter would be presented in 1 of the 4 visual field quadrants around the fixation.

In a n-back task condition (n =1 or 2), participants were asked to press the left button with the left thumb

when the location of the current letter matched with the one presented "n" item(s) back, and pressed the

right button with the right thumb when it didn't match the one presented "n" item(s) back. The letter

stimulus was presented for 500 ms, followed by an interstimulus interval of 2500 ms. One third of the

total trials were "matches". Participants were instructed to focus only on the location of the letter, but not

on the letter itself, and to classify the stimuli as accurately and quickly as possible. Visual stimuli were

presented and responses were collected using E-Prime (Psychology Software Tools).

The N-back task procedures were designed in two ways. First, in the two localizer runs, the n-

back stimuli were presented as separate blocks of 1-back or 2-back conditions. Each run started with a 10

s fixation. Then, each of the block consisted of 8 trials (24 sec), with a 24-s fixation period intercepted

between the task blocks. The orders of task blocks of the two runs were "ABBA" and "BAAB",

respectively. As a result, each run lasted for 3 min and 46 sec. Second, in the two continuous runs, the n-

back trials were presented continuously without long fixation period between them. The 1-back and 2-

back conditions were allocated in two separate runs. Each run started with a 10 s fixation period,

followed by 100 trials. Each run lasted for 5 min and 10 sec.

2.2.2. MRI scanning parameters

MRI data were acquired on a 3T GE Signa Scanner (General Electric Company, Milwawkee, WI), using

an 8-channel head coil. The parameters for the fMRI images were: TR (repetition time) = 2000 ms; TE

(echo time)) = 30 ms; flip angle = 90°; FOV (field of view) = 240×240 mm^2; matrix size = 64×64; axial

slice number = 42 with slice thickness = 3 mm and gap = 0). As a result, each resting-state run was

consisted of 255 images, each block-designed run was consisted of 113 images, and each continuous task

run was consisted of 155 images. Structural T1-weighted images were acquired using the following

parameters: TR = 6 ms; TE = Minimum; TI = 450 ms; flip angle = 12°; FOV = 256×256 mm^2; matrix size

= 256×256; sagittal slice number = 156 with slice thickness = 1 mm.

2.3. FMRI data analysis

2.3.1. Preprocessing

FMRI images were processed using SPM12 (SPM, RRID:SCR_007037;

https://www.fil.ion.ucl.ac.uk/spm/) under MATLAB environment (R2017b). The anatomical image of

each subject was segmented into gray matter (GM), white matter (WM), cerebrospinal fluid (CSF), and

other brain tissue types, and normalized into standard Montreal Neurological Institute (MNI) space. The

first five functional images of each run were discarded from analysis. The remaining images were

realigned to the first image of each run, and coregistered to the anatomical image. The deformation field

images obtained from the segmentation step were used to normalize all the functional images into MNI,

with a resampled voxel size of 3 x 3 x 3 mm^3. All the images were spatially smoothed using an 8 x 8 x 8

mm^3 Gaussian kernel.

We calculated frame-wise displacement for the translation and rotation directions, respectively, to

reflect the amount of head motions (Di & Biswal, 2015a). We adopted the threshold of maximum frame-

wise displacement of 1.5 mm or 1.5 degree (half voxel size), or mean frame-wise displacement of 0.2 mm

or 0.2 degree. The subjects with any of the five runs exceeding the threshold would be removed from the

analysis. As a result, one subject's data were discarded.

2.3.2. Activation analysis of the block-designed runs

We first defined general linear model (GLM) to perform voxel-wise analysis on the block-designed runs

to identify task activations between the 2-back and 1-back conditions. The two runs were modeled

together with their own task regressors, covariates, and constant terms. The 2-back and 1-back conditions

were defined as two box-car functions convolved with canonical hemodynamic response function (HRF).

The first eigenvector of signals in the WM and that in the CSF, 24 head motion regressors (Friston,

Williams, Howard, Frackowiak, & Turner, 1996) were added as covariates. There was also as high-pass

filtering (1/128 Hz) implicitly implemented in the GLM. After model estimation, a contrast of 2-back –

1-back was defined to reflect the differences of activations between the two conditions.

Group level analysis was performed using one sample test GLM with the input of the contrast

images of 2-back vs. 1-back. Activated clusters were first identified using a threshold of $p < 0.001$ of

two-tailed test (Chen et al., 2019), and the cluster extent was thresholded at cluster level false discovery

rate (FDR) of $p < 0.05$. Because we were interested in fronto-parietal regions, we searched the peak

coordinates of the resulting clusters as well as local maxima within large clusters that covered these

regions. As a result, we defined bilateral middle frontal gyrus regions (MNI coordinates: RMFG, 24, 11,

56; LMFG, -24, 8, 50) and superior parietal lobule (MNI coordinates: LSPL, -18, -70, 50; RSPL, 21, -67,

53) as ROIs.

2.3.3. Physiophysiological interaction analysis of the continuous-designed runs

We first defined GLMs for each continuous run and subject to define ROIs. The GLMs did not include

task regressors, but only had the WM/CSF, head motion, and constant regressors. There was also as high-

pass filtering (1/128 Hz) implicitly implemented in the GLM. After model estimation, the time series of

the LMFG, LSPL, RMFG, and RSPL were extracted within spherical ROIs of 6 mm radius centered at

the above mentioned MNI coordinates. All the effects of no-interests, i.e. WM/CSF signals, head motion

parameters, constant, and low-frequency drifts were adjusted during the time series extraction. PPI terms

were calculated for LMFG and LSPL, and RMFG and RSPL, respectively. The time series of the two

ROIs were deconvolved with canonical HRF, multiplied together, and convolved back with canonical

HRF to form a PPI term (Di & Biswal, 2013; Gitelman, Penny, Ashburner, & Friston, 2003).

Next, new GLMs were built with the time series of the two ROIs and the PPI term between them

for each of the ROI pairs and conditions. Other regressors of no-interests as well as the implicit high-pass

filter were also included in the GLMs. The beta estimates corresponding to the interaction term was the

effect of interest, which were used for the group level analysis.

The first goal of the group analysis is to identify regions that show modulatory interaction effects

consistently present in the three conditions. We performed conjunction analysis of the three conditions.

Second-level GLM was built for the LMFG-LSPL and RMFG-RSPL analyses separately using a one-way

analysis of variance (ANOVA) model. First, a t contrast of each condition was defined for both positive

and negative effects. Next, we examined the conjunction effects of the three conditions for the positive

and negative effects, respectively, using a threshold of one-tailed $p < 0.0005$ (corresponding to two-tailed

$p < 0.001$). Cluster level FDR of $p < 0.05$ was used for the cluster extent threshold. Because there were

no clusters survived at the two-tailed $p < 0.001$ threshold, we also explored lower threshold of two-tailed

$p < 0.01$ for potential effects.

The second goal is to identify regions that showed variable modulatory interactions in the three

conditions. Repeated measure one-way ANOVA model was used for this purpose, with the three

conditions as three levels of a factor. The significant results of the repeated measure ANOVA indicate

differences in the PPI effects between any two of the three conditions. The resulting statistical maps were

thresholded at $p < 0.001$ with cluster level FDR at $p < 0.05$.

3. Results

3.1. Task activations in the localizer runs

We observed typical bilateral fronto-parietal regions that showed higher activations during the 2-back

condition compared with 1-back condition (Figure 1 and Table 1). The frontal clusters mainly covered

the bilateral middle frontal gyrus and precentral gyrus. The parietal clusters mainly covered the bilateral

superior parietal lobule and precuneus. The right cerebellum and left basal ganglia were also activated.

There were also reduced activations in the 2-back compared with 1-back condition, mainly in the default

model network and bilateral temporo-opercular regions.

Figure 1 Increased (warm color) and decreased (cold color) activations in the 2-back condition compared

with the 1-back condition. The map was thresholded at p < 0.001 (two-tailed) with cluster-level false

discovery rate of p < 0.05. The surface presentation was made using BrainNet Viewer

(RRID:SCR_009446) (Xia, Wang, & He, 2013).

3.2. Modulatory interactions during different task conditions

We first performed conjunction analysis to identify regions that showed consistent PPI effects across the

three conditions. No statistical significant clusters were found of any sizes at p < 0.001 for both the

LMFG-LSPL and RMFG-RSPL analyses. We further checked the threshold of p < 0.01, and still there

were no clusters of any sizes survived.

Repeated measure one-way ANOVA showed only significant effects on the modulatory

interactions of RMFG and RSPL. As shown in Figure 2 and Table 2, the only cluster mainly covered the

anterior cingulate cortex (ACC). Post-hoc analysis showed that the PPI effect in the ACC was positive in

the 2-back condition but negative during resting-state (Figure 2B). Repeated measure one-way ANOVA

of the modulatory interactions of LMFG and LSPL showed a similar cluster in the ACC. However, the

cluster size could not pass the cluster-level threshold.

Figure 2 A) Regions that showed different modulatory interactions effects with right middle frontal gyrus

(RMFG) and right superior parietal lobule (RSPL) among the three task conditions (repeated measure one

way analysis of variance: ANOVA). The map was thresholded at $p < 0.001$ with cluster level false

discovery rate (FDR) of $p < 0.05$. B) Mean modulatory interactions of the cluster in the in the three

conditions of continuous runs. The center red lines represent the mean effects, and the light red bars and

light blue bars represent 95% confidence interval and standard deviation, respectively. Panel B was made

by using notBoxPlot (https://github.com/raacampbell/notBoxPlot). A.u., arbitrary unit.

`

In order to better interpret the PPI effects in the ACC, we correlated the mean PPI effects in the

ACC cluster with RMFC and RSPL with behavioral measures of mean reaction time and accuracy (Figure

3). The PPI effect showed a very small correlation with reaction time ($r = -0.16$), and a moderate

negative correlation with the accuracy ($r = -0.39$). But it can be seen in Figure 3C that there were

potential outliers near the x axis that might introduce spurious correlations. We therefore performed

bootstrapping for 10,000 times to obtain a 95% confidence interval of the correlation (-0.6352, 0.0046)

(Figure 3D).

Figure 3 Behavioral correlates of the mean modulatory interactions in the ACC with RMFG and RSPL

during the 2-back continuous run. A and B illustrate the scatter plot of correlations between the

modulatory interaction and reaction time and 10,000 bootstrapping distributions of the correlation. C and

D illustrate the scatter plot of correlations between the modulatory interaction and accuracy and 10,000

bootstrapping distributions of the correlation.

Lastly, we also extracted the mean task activations of the ACC in the block-designed runs (Figure

4). The ACC showed reduced activations in both the 1-back and 2-back conditions with reference to the

fixation baseline. But the activations were more negative in the 2-back condition than in the 1-back

condition (paired t test: $t(48) = 4.49, p < 0.001$).

Figure 4 Mean task activations of the cluster in the block-designed runs. The center red lines represent

the mean effects, and the light red bars and light blue bars represent 95% confidence interval and standard

242   deviation, respectively. This figure was made by using notBoxPlot

243   (https://github.com/raacampbell/notBoxPlot). A.u., arbitrary unit.

4. Discussion

By comparing modulatory interactions of two key regions in working memory across three continuously

designed task conditions, the current analysis identified the ACC that showed different modulatory

interactions with the RMFG and RSPL in the resting-state, 1-back, and 2-back conditions. On the other

hand, no regions showed consistent modulatory interactions with the fronto-parietal regions across the

three conditions. The activity in the ACC was positively correlated with the connectivity of RMFG and

RSPL during the 2-back condition, but was negatively correlated with the connectivity of RMFG and

RSPL in resting-state. Due to the nature of regression model, this is impossible to infer the directions of

the modulations (Di & Biswal, 2013). However, the RMFG and RSPL were co-activated by the working

memory task and are also considered part of the same functional network (Biswal et al., 2010; Yeo et al.,

2011), while the ACC showed increased deactivation in the 2-back condition. We prefer to interpret the

results as that the ACC increase the functional connectivity between RMFG and RSPL during the 2-back

condition, and reduce the functional connectivity between the RMFG and RSPL.

Due to the fact that the ACC was negatively activated in the task conditions compared with the

fixation condition (Figure 4), it is likely that the ACC is part of the default mode network (Raichle et al.,

2001). It is consistent with our previous study in resting-state, which also showed some midline regions

from the default mode network having negative modulatory interactions with RMFG and RSPL (Di &

Biswal, 2013). The task positive network including the fronto-parietal regions and the default mode

network are anti-correlated both in resting-state (Fox et al., 2005) and during task executions (Shulman et

al., 1997). The current results together with our previous work (Di & Biswal, 2013) further confirm that

the competing nature of the task positive and default mode networks not only exist in first order

relationships but also in higher order interactions.

More interestingly, current analysis found that the modulatory interactions among ACC, RMFG,

and RSPL were largely modulated by task conditions. In contrast to resting-state, the ACC showed no

significant modulatory interactions in the 1-back condition, and positive modulatory interactions in the 2-

back condition. The task dependent effect is in line with some studies that have demonstrated task

modulated modulatory interactions in other brain systems by using higher order psycho-physio-

physiological interaction models (Gorka, Knodt, & Hariri, 2015; Stamatakis, Marslen-Wilson, Tyler, &

Fletcher, 2005). In neuronal level models, it has also been shown that higher order interactions present

only in certain task conditions (Ganmor, Segev, & Schneidman, 2011; Macke, Opper, & Bethge, 2011).

Taken together, all the evidence conversely suggests that high order interactions may be sensitive to

certain task conditions.

During the 2-back condition with higher working memory loads, the signals from the ACC were

associated with increased functional communications between the fronto-parietal regions. One of the

functions of the ACC is error detection and conflict monitoring (Bush, Luu, & Posner, 2000). Then, the

ACC activity may represent error related signals that would enhance the communications between the

fronto-parietal regions to maintain task performances. The brain-behavioral correlation analysis

supported this interpretation. The modulatory interactions in the 2-back condition were not correlated

with reaction time, but were negatively correlated with accuracy. In other words, the more errors one

made, the larger the modulatory interactions were among ACC, RMFG, and RSPL.

The current study adopted functionally defined ROIs of the MFG and SPL from a localizer for the

PPI analysis. The bilateral MFGs are a little anterior to the premotor regions and posterior to the

dorsolateral prefrontal cortex reported in a meta-analysis of N-back tasks (Owen et al., 2005). And the

bilateral SPLs are superior and posterior to the inferior parietal lobule region reported in (Owen et al.,

2005). The differences may represent discrepancies in task designs and control conditions compared with

other studies. But the fact that these regions showed the highest contrast between the 2-back and 1-back

condition in the current localizer task support the usage of these regions to represent regions that are

involved in working memory process. The fronto-parietal ROIs also do not exactly match those used in

the resting-state study (Di & Biswal, 2013). But similar to this paper, the current analysis showed

negative modulatory interactions in the middle line region of ACC with RMFG and RSPL (Di & Biswal,

2013).

The current analysis adopted a ROI-based approach, with ROIs identified directly from the

working memory task studied. This helped us to focus on specific brain regions that are related to the

task. The whole brain PPI analysis identified a region that are not a part of the fronto-parietal network

nor activated during the working memory tasks. It is reasonable because our previous study has shown

that modulatory interactions are more likely to take place among regions from different brain networks

(Di & Biswal, 2015a). There may be other brain regions that involve higher order interactions with one

of the fronto-parietal regions. But the potential interactions will increase exponentially when considering

the combinations of two brain regions outside the fronto-parietal network, making it difficult to do an

exhaustive search based on the current sample size. Further studies may adopt the whole brain approach

(Di & Biswal, 2015a) to examine the whole brain characterizations of modulatory interactions effects.

In conclusion, the current analysis extended our previous analysis in resting-state and showed that

the modulatory interaction among ACC and right fronto-parietal regions were highly modulated by task

demands. The results may provide new model on how error related signals affecting working memory

process through higher order interactions among brain regions.

Acknowledgements:

This study was supported by grants from National Natural Science Foundation of China (NSFC61871420)

and (US) National Institute of Health (R01 AT009829; R01 DA038895).

Author contributions:

X.D. conceived the idea. H.Z. designed the experiment and collected the fMRI data. X.D. performed the

data analysis and wrote the draft. All authors discussed the results, and contributed to the final manuscript.

Conflict of interest statement:

The authors declare that there is no conflict of interest regarding the publication of this article.

Reference:

Babiloni, C., Babiloni, F., Carducci, F., Cincotti, F., Vecchio, F., Cola, B., … Rossini, P. M. (2004).

Functional Frontoparietal Connectivity During Short-Term Memory as Revealed by High-

Resolution EEG Coherence Analysis. *Behavioral Neuroscience*, *118*(4), 687–697.

https://doi.org/10.1037/0735-7044.118.4.687

Barch, D. M., Burgess, G. C., Harms, M. P., Petersen, S. E., Schlaggar, B. L., Corbetta, M., … Van Essen,

D. C. (2013). Function in the human connectome: Task-fMRI and individual differences in

behavior. *NeuroImage*, *80*, 169–89. https://doi.org/10.1016/j.neuroimage.2013.05.033

Beckmann, C. F., DeLuca, M., Devlin, J. T., & Smith, S. M. (2005). Investigations into resting-state

connectivity using independent component analysis. *Philosophical Transactions of the Royal*

Society of London. Series B, Biological Sciences, *360*(1457), 1001–13.

https://doi.org/10.1098/rstb.2005.1634

Biswal, B. B., Mennes, M., Zuo, X.-N., Gohel, S., Kelly, C., Smith, S. M., … Milham, M. P. (2010).

Toward discovery science of human brain function. *Proceedings of the National Academy of*

Sciences of the United States of America, *107*(10), 4734–9.

https://doi.org/10.1073/pnas.0911855107

Bush, G., Luu, P., & Posner, M. I. (2000). Cognitive and emotional influences in anterior cingulate cortex.

Trends in Cognitive Sciences, *4*(6), 215–222. https://doi.org/10.1016/S1364-6613(00)01483-2

Chen, G., Cox, R. W., Glen, D. R., Rajendra, J. K., Reynolds, R. C., & Taylor, P. A. (2019). A tail of two

sides: Artificially doubled false positive rates in neuroimaging due to the sidedness choice with t-

tests. *Human Brain Mapping*, *40*(3), 1037–1043. https://doi.org/10.1002/hbm.24399

Dai, Z., de Souza, J., Lim, J., Ho, P. M., Chen, Y., Li, J., … Sun, Y. (2017). EEG Cortical Connectivity

Analysis of Working Memory Reveals Topological Reorganization in Theta and Alpha Bands.

Frontiers in Human Neuroscience, *11*. https://doi.org/10.3389/fnhum.2017.00237

Di, X., & Biswal, B. B. (2013). Modulatory interactions of resting-state brain functional connectivity.

PloS One, *8*(8), e71163. https://doi.org/10.1371/journal.pone.0071163

Di, X., & Biswal, B. B. (2014). Modulatory interactions between the default mode network and task

positive networks in resting-state. *PeerJ*, *2*, e367. https://doi.org/10.7717/peerj.367

Di, X., & Biswal, B. B. (2015a). Characterizations of resting-state modulatory interactions in the human

brain. *Journal of Neurophysiology*, *114*(5), 2785–96. https://doi.org/10.1152/jn.00893.2014

Di, X., & Biswal, B. B. (2015b). Dynamic brain functional connectivity modulated by resting-state

networks. *Brain Structure and Function*, *220*(1), 37–46. https://doi.org/10.1007/s00429-013-

0634-3

Di, X., & Biswal, B. B. (2019). Toward Task Connectomics: Examining Whole-Brain Task Modulated

Connectivity in Different Task Domains. *Cerebral Cortex*, *29*(4), 1572–1583.

https://doi.org/10.1093/cercor/bhy055

Fox, M. D., Snyder, A. Z., Vincent, J. L., Corbetta, M., Van Essen, D. C., & Raichle, M. E. (2005). The

human brain is intrinsically organized into dynamic, anticorrelated functional networks.

Proceedings of the National Academy of Sciences of the United States of America, *102*(27),

9673–8.

Friston, K. J., Buechel, C., Fink, G. R., Morris, J., Rolls, E., & Dolan, R. J. (1997). Psychophysiological

and modulatory interactions in neuroimaging. *NeuroImage*, *6*(3), 218–29.

Friston, K. J., Williams, S., Howard, R., Frackowiak, R. S., & Turner, R. (1996). Movement-related

effects in fMRI time-series. *Magnetic Resonance in Medicine : Official Journal of the Society of*

Magnetic Resonance in Medicine / Society of Magnetic Resonance in Medicine, *35*(3), 346–55.

https://doi.org/DOI 10.1002/mrm.1910350312

Ganmor, E., Segev, R., & Schneidman, E. (2011). Sparse low-order interaction network underlies a

highly correlated and learnable neural population code. *Proceedings of the National Academy of*

Sciences, *108*(23), 9679–9684. https://doi.org/10.1073/pnas.1019641108

Gitelman, D. R., Penny, W. D., Ashburner, J., & Friston, K. J. (2003). Modeling regional and

psychophysiologic interactions in fMRI: the importance of hemodynamic deconvolution.

NeuroImage, *19*(1), 200–7.

Gorka, A. X., Knodt, A. R., & Hariri, A. R. (2015). Basal forebrain moderates the magnitude of task-

dependent amygdala functional connectivity. *Social Cognitive and Affective Neuroscience*, *10*(4),

501–507. https://doi.org/10.1093/scan/nsu080

Macke, J. H., Opper, M., & Bethge, M. (2011). Common input explains higher-order correlations and

entropy in a simple model of neural population activity. *Physical Review Letters*, *106*(20),

208102. https://doi.org/10.1103/PhysRevLett.106.208102

Mencarelli, L., Neri, F., Momi, D., Menardi, A., Rossi, S., Rossi, A., & Santarnecchi, E. (n.d.). Stimuli,

presentation modality, and load-specific brain activity patterns during n-back task. *Human Brain*

Mapping, *0*(0). https://doi.org/10.1002/hbm.24633

Owen, A. M., McMillan, K. M., Laird, A. R., & Bullmore, E. (2005). N-back working memory paradigm:

A meta-analysis of normative functional neuroimaging studies. *Human Brain Mapping*, *25*(1),

46–59. https://doi.org/10.1002/hbm.20131

Raichle, M. E., MacLeod, A. M., Snyder, A. Z., Powers, W. J., Gusnard, D. A., & Shulman, G. L. (2001).

A default mode of brain function. *Proceedings of the National Academy of Sciences of the United*

States of America, *98*(2), 676–82. https://doi.org/10.1073/pnas.98.2.676

Sauseng, P., Klimesch, W., Schabus, M., & Doppelmayr, M. (2005). Fronto-parietal EEG coherence in

theta and upper alpha reflect central executive functions of working memory. *International*

Journal of Psychophysiology, *57*(2), 97–103. https://doi.org/10.1016/j.ijpsycho.2005.03.018

Shulman, G. L., Fiez, J. A., Corbetta, M., Buckner, R. L., Miezin, F. M., Raichle, M. E., & Petersen, S. E.

(1997). Common Blood Flow Changes across Visual Tasks: II. Decreases in Cerebral Cortex.

Journal of Cognitive Neuroscience, *9*(5), 648–663. https://doi.org/10.1162/jocn.1997.9.5.648

Stamatakis, E. A., Marslen-Wilson, W. D., Tyler, L. K., & Fletcher, P. C. (2005). Cingulate control of

fronto-temporal integration reflects linguistic demands: A three-way interaction in functional

connectivity. *NeuroImage*, *28*(1), 115–21. https://doi.org/10.1016/j.neuroimage.2005.06.012

Stephan, K. E., Kasper, L., Harrison, L. M., Daunizeau, J., den Ouden, H. E. M., Breakspear, M., &

Friston, K. J. (2008). Nonlinear dynamic causal models for fMRI. *NeuroImage*, *42*(2), 649–62.

https://doi.org/10.1016/j.neuroimage.2008.04.262

Xia, M., Wang, J., & He, Y. (2013). BrainNet Viewer: A network visualization tool for human brain

connectomics. *PloS One*, *8*(7), e68910. https://doi.org/10.1371/journal.pone.0068910

Yeo, B. T. T., Krienen, F. M., Sepulcre, J., Sabuncu, M. R., Lashkari, D., Hollinshead, M., … Buckner, R.

L. (2011). The organization of the human cerebral cortex estimated by intrinsic functional

connectivity. *Journal of Neurophysiology*, *106*(3), 1125–65.

https://doi.org/10.1152/jn.00338.2011

Table 1 Clusters that showed increased activations in the 2-back condition compared with 1-back

condition in the block designed runs. The cluster was defined as two tailed p < 0.001, with cluster level

false discovery rate p < 0.05.

p (cluster FDR)	voxels	Coordinates			peak T	Label
		x	y	z		
< 0.001	2108	24	11	56	11.65	Right middle frontal gyrus
		-24	8	50	10.72	Left middle frontal gyrus
		-48	5	32	9.810	Left precentral gyrus
< 0.001	2897	-6	-61	44	10.73	Precuneus
		-18	-70	50	10.68	Left superior parietal lobule
		21	-67	53	10.44	Right superior parietal lobule
0.004	120	48	5	23	7.00	Right precentral gyrus
0.003	149	27	-61	-37	6.92	Right cerebellum
		9	-73	-31	4.78	Right cerebellum
0.003	136	-18	5	11	5.84	Left caudate
		-30	26	2	5.75	Left anterior insula
0.038	63	-33	50	2	4.20	Left middle frontal gyrus
		-42	50	2	4.02	Left middle frontal gyrus
< 0.001	661	-3	-16	32	-8.73	Middle cingulate gyrus
		0	-37	20	-6.08	Posterior cingulate gyrus
		0	-28	44	-5.42	Posterior cingulate gyrus
< 0.001	660	39	-19	20	-6.54	Right parietal operculum
		36	-16	2	-5.56	Right posterior insula
		39	2	-1	-5.16	Right anterior insula
< 0.001	910	12	59	20	-6.11	Superior frontal gyrus
		-6	62	8	-5.86	Medial superior frontal gyrus
		-9	53	-1	-5.84	Medial superior frontal gyrus
< 0.001	498	-36	-10	-4	-5.39	Left posterior insula
		-63	-25	5	-4.73	Left superior temporal gyrus
		-39	-19	17	-4.64	Left central operculum
0.037	74	21	38	-1	-5.19	Anterior cingulate gyrus

413　FDR, false discovery rate. X, y, and z coordinates are in (Montreal Neurological Institute) MNI space.

415 **Table 2** Clusters that showed different physiophysiiological interaction (PPI) effects with right middle

416 frontal gyrus (RMFG) and right superior parietal lobule (RSPL) among the resting-state, 2-back, and 1-

417 back conditions in the continuous runs (repeated measure one way analysis of variance). The cluster was

418 defined as $p < 0.001$, with cluster level false discovery rate $p < 0.05$.

p (cluster FDR)	voxels	Coordinates			peak F	Label
		x	y	z		
0.005	133	-3	32	14	14.94	Anterior cingulate gyrus
		9	35	5	14.82	Anterior cingulate gyrus
		3	44	-4	8.27	Anterior cingulate gyrus

420 FDR, false discovery rate. X, y, and z coordinates are in (Montreal Neurological Institute) MNI space.