Measurement of Atom Resolvability in CryoEM Maps with Q-scores

Grigore Pintilie[1*], Kaiming Zhang[1], Zhaoming Su[1], Shanshan Li[1], Michael F. Schmid[2], Wah Chiu[1,2*]

1. Department of Bioengineering, James H.Clark Center, Stanford University, Stanford, CA 94305, USA
2. Division of CryoEM and Bioimaging, SSRL, SLAC National Accelerator Laboratory, Stanford University, Menlo Park, CA 94025, USA

* Corresponding authors: gregp@slac.stanford.edu, wahc@stanford.edu

Abstract

CryoEM density maps are now at the point where resolvability of individual atoms can be achieved. However, resolvability is not necessarily uniform throughout the map. We introduce a quantitative parameter to characterize the resolvability of individual atoms in cryoEM maps, the map Q-score. Q-scores can be calculated for atoms in proteins, nucleic acids, water, ligands, and other solvent atoms, using models fitted to or derived from cryoEM maps. Q-scores can also be averaged to represent larger features such as entire residues and nucleotides. In several examples, Q-scores are shown to correlate well to resolvability. Q-scores averaged over entire models are also shown to correlate closely to the estimated resolution of cryoEM maps for both protein and RNA. Assuming the models they are calculated from are well-fitted to the map, Q-scores can thus be used as another measure to indicate the quality or resolvability in entire maps and also of features as small as individual atoms.

Introduction

CryoEM single particle methods strive to create accurate, high-resolution 3D maps of macromolecular complexes. Depending on many factors including imaging apparatus, detector, reconstruction method, structure flexibility, sample heterogeneity, and differential radiation damage, resulting maps have varying degrees of resolvability, or the level at which molecular features can be seen. Accurate quantification of resolvability in cryoEM maps has been a challenge in the field[1]. This task is very important as it can affect the interpretability and insights derived from such maps.

For every cryoEM map, a resolution is commonly reported or estimated, calculated from a
Fourier shell correlation (FSC) plot between two independent reconstructions of the same
complex[2]. It is well recognized that cryoEM maps usually do not have isotropic resolution
throughout, hence a single number may not accurately represent the entire map. Local resolution
can be estimated by most of image processing software (e.g. ResMap[3]), however such
information is not as easy to comprehend in terms of specific residues as in the case with an
atomic model.

Atomic models can be either fitted or built directly into cryoEM maps[4,5]. Map-model scores are
then calculated from the model and map to assess how well the model fits the map[6]. Refinement[7]
or flexible fitting[8,9] can then be applied, while making sure not to distort or overfit to noise[10,11].
The latter is accomplished by applying various stereochemical constraints, e.g. proper bond
lengths, angles, dihedrals, preferred rotamers and van-der Waals distances; additional secondary-
structure constraints (e.g.in the form of hydrogen bonds) can also be applied[7,9,12,13].

Once an atomic model has been fitted to or derived from a cryoEM map, it can then be used to
measure the resolvability of the features in the map . This can be done in several ways, including
a map-model FSC curve, which requires that the model first be converted to a cryoEM-like map
at the same resolution as the map. Occupancies and atomic displacement parameters of residues
or atoms can also be used in this process to make the model-map better match the cryoEM
map[14]. However, the FSC plot reflects the entire map volume. Proper masking may evaluate the
resolvability of smaller features such as individual protein chains[10], however it is impractical to
quantify the resolvability of even smaller features such as a single side chain using this approach.

Two other methods that measure resolvability of such smaller features in a cryoEM map using a
fitted model are EMRinger[15] and Z-scores[16]. EMRinger considers map values near carbon-β
atoms, while Z-scores can be applied to secondary structures or entire side chains. These scores
were shown to correlate with the reported resolution when averaged over an entire map and
model, meaning they can also be used to double-check whether the estimated resolution of the
map is reasonable. Moreover, they can also pinpoint smaller features in the model (e.g.
secondary structures or side chains) which are not well-resolved in the map or not fitted properly
to the map.

CryoEM maps have reached resolutions nearer to atomic-scale, for example apoferritin at 1.54Å
(EMD:9865), 1.62Å (EMD:0144)[17], 1.65Å (EMD:9599), and 1.75Å (EMD: 20026). A new
question now arises as to how resolvability of individual atoms may be assessed. In
crystallography, this is often reflected in the B-factor calculated for each atom[18]. Several
formulations and interpretations of the B-factors are possible[19], and their use in cryoEM has been
suggested in the form of atomic displacement parameters (ADPs)[14]. So far, such formulations
have not been fully characterized in terms of resolvability and resolution of map.

In this paper, we introduce a new score which is calculated from map values around an atom's
position, the Q-score. It aims to be a direct measure of the resolvability of atoms in cryoEM
maps of complexes containing proteins, nucleic acids, and solvent molecules.

Atomic Map Profiles

The basis of the Q-score is the atomic map profile. Atomic map profiles are calculated by
averaging map values at increasing radial distances from the atom's position. The radial
distances range from 0Å to 2.0Å, and only points that are closer to the atom in question than to
any other atoms in the model are considered. Figure 1A shows example atomic profiles in our
two new maps of Apoferritin with resolutions of 1.75Å and 2.32Å, now deposited as
EMD:20026, and EMD:20027. The model is the X-ray model of Apoferritin, (PDB:3ajo), which
was first rigidly fitted to the cryoEM map, and then further refined using the Phenix real-space
refinement procedure[7]. In the examples, atomic profiles have Gaussian-like contours. We
consider a Gaussian equation of the form:

$$y = Ae^{-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2} + B \qquad\qquad (1)$$

Gaussian functions of the form in Eqn.1, where x is the radial distance and y the average map
value, fit extremely well to the atomic profiles shown in Figure 1, up to a distance of 2Å (mean
error of ~0.01Å). Past this distance, observations in various maps indicate that atomic map
profiles become noisy and start to increase. This is likely due to effects from other nearby atoms
and/or solvent.

When the model is well-fitted to the map, the relative height, **A-B**, and width, **σ**, of the Gaussian
function (Eqn.1) fitted to the profile may be considered to be proportional to several factors
including the resolution of the map, and the overall mobility of the atom. It may be impossible to
fully separate such factors based on the observed cryoEM map alone. Regardless of the cause,
the overall Gaussian profile seen in the map represents to what degree the respective atom is
resolved in the map - the more resolved an atom is in the map, the higher (relative to other peaks
in the same map) and narrower (up to a certain point, i.e. the radius of the atom itself) the
Gaussian profile around it would be.

A **1.75Å**, EMD:20026, PDB:3ajo, Leu26

2.3Å, EMD:20027, PDB:3ajo, Leu26

0.5Å   1.0Å   1.5Å   2.0Å

Radial Distance (Å)

B

Radial Distance (Å)

Average Map Value

Figure 1. Atomic map profiles in cryoEM maps of Apoferritin at 1.75Å and 2.3Å resolution. (A) The residue Leu26 in the fitted model (PDB:3ajo) is shown, along with contour surface of the cryoEM map around this residue. Spherical shells of points centered on the CD2 atom are shown at increasing radial distances; only points that are closer to the CD2 atom than to any other atom in the model are used. (B) Average map values at these points are plotted vs. radial distance; these are the atomic map profiles. The dotted lines represent Gaussian functions with parameters *A*, *B* and *σ* which are fitted to each profile.

Q-score

The idea behind the Q-score is to measure how closely the map profile of an atom matches that of the Gaussian-like function we would see if an atom is well-resolved. Thus, to calculate the Q-score, the atomic map profile is compared to a 'reference Gaussian' as given by Eqn. 1, with the following parameters:

$$\mu = 0 \qquad (2)$$
$$A = avg_M + 10\sigma_M \qquad (3)$$
$$B = avg_M - 1\sigma_M \qquad (4)$$
$$\sigma = 0.6\text{Å} \qquad (5)$$

In the above, the mean, μ, is set to 0, as the Gaussian is expected to be centered around the atom's position. The parameters *A* and *B* are obtained using the mean/average across all values in the entire map, avg_M, and the standard deviation of all values around this mean, σ_M. A well

142 resolved atom would be centered on a peak that has a relatively high value in the map, and fall
143 off to a value below the mean, but not necessarily as low as the background noise. The width of
144 the reference gaussian is set as σ=0.6. These parameters in Eqns. 2-5 are chosen to make the
145 reference Gaussian roughly match the atomic profile of a well-resolved atom in the 1.54Å
146 cryoEM map as shown in Figure 2B.

147
148

149
Figure 2. Calculation of Q-scores for an atom in 6 maps at different resolutions, including an X-ray map.
The atom is CD2 from Leu 26 in PDB:3ajo. The atomic profile in each map is marked with the letter u,
while the reference Gaussian is marked with v.

The Q-score is then calculated as a correlation between values in the atomic profile obtained
from the map, u, and values obtained from the reference Gaussian, v, defined in Eqn. 1 and with
parameters in Eqns. 2-5. The following normalized, about the mean, cross-correlation formula is
used:

$$Q(atom) = \frac{\langle u - u_{mean} \rangle \langle v - v_{mean} \rangle}{|u - u_{mean}||v - v_{mean}|} \qquad (6)$$

Several atomic profiles and reference Gaussians are illustrated in Figure 2, for an X-ray map and
5 cryoEM maps at various resolution. At high resolutions, the atomic profiles are more similar to
the reference Gaussian, and hence Q-scores are higher. At lower resolutions, the atomic profile
of the same atom is wider than the reference Gaussian, hence Q-scores are lower. Q-scores
would also be low for atomic profiles that are mostly noise (e.g. random values or a sharp peak).

In some cases when the atom is not well-placed in the map, the Q-score can be negative if the
atomic profile has a shape that increases away from the atom's position.
168
Calculating Q-scores is similar to calculating a cross-correlation between the model and a
cryoEM map, using a simulated map of the model blurred using a Gaussian function with the
parameters in Eqns. 2-5. The main difference is that with Q-scores, the cross-correlation is
performed atom-by-atom, separating out parts of the density that are closest to each atom. The
cross-correlation about the mean is used so that the Q-scores decrease as resolution also
decreases. When not subtracting the mean, this effect would not be ensured[16].
175
176
177
178 **Q-scores of Atoms in Proteins**
179
Figure 3 shows Q-scores for atoms taken from maps of Apoferritin at various resolutions. One of
the maps is an X-ray map at 1.52Å resolution (2fo-fc, PDB:3ajo) as a reference; another is a
recent high-resolution map at 1.54Å (EMD:9599). The other three are new maps we
reconstructed to 1.75Å (EMD:20026), 2.3Å (EMD:20027), and 3.1Å (EMD:20028) with
different numbers of particle images, from the same data set. For the cryoEM maps, the X-ray
model PDB:3ajo was fitted to the density and also refined using Phenix real-space refinement[7].
Q-scores for each atom correlate well with visual resolvability, i.e. the more resolvable an atom,
the higher the Q-score. They also increase as the estimated resolution of the map increases.
188
Resolvability and Q-scores can decrease for some residues faster than others as a function of
resolution. For example, in Figure 3, the Q-score for ASP126 drops more than for ASN25 from
1.52Å to 3.9Å. This effect may be due to several reasons. First, some residue types may be more
susceptible to radiation damage (as previously shown using EMRinger[15]). Also, certain residue
types may be more conformationally dynamic, or occur in environments that are more dynamic
(e.g. solvent accessible), and hence may not resolve as well with a fewer number of particles.
Finally, the interaction of the electron beam with charged side chains may have a weakening
effect on map values around them[14].
197

198
199 Figure 3. Atom Q-scores for three types of residues, taken from Apoferritin maps at various resolutions.
200 Atom Q-scores are shown in purple close to each atom.
201
202
203 **Q-scores for Atoms in Nucleic Acids**
204
205 Q-scores can also be calculated for atoms in models of nucleic acids. In figure 4, we used several
206 maps and models containing RNA from the EMDB at resolutions ranging from 2.5Å to 4.0Å. Q-
207 scores were averaged over atoms in bases, phosphate-sugar backbones, and entire nucleotides.
208 As with proteins, Q-scores decrease with resolvability and estimated map resolution.
209
210 Figure 4 also illustrates a general trend that at ~4Å and lower resolutions, stacked bases from
211 adjacent nucleotides are typically not separable in cryoEM maps, whereas at higher than 4Å
212 resolutions, they usually do become separate at appropriate contour levels.
213

It is also interesting to note that for the examples in Figure 4, at high resolutions (~2.5Å), the
difference in Q-score or resolvability of individual bases is higher than that of the backbone
(0.84 for base vs. 0.73 for backbone). Going towards lower resolutions in this example, bases
become less resolvable (0.45 for bases vs 0.56 for backbone). This may be counter-intuitive as
bases can have higher values in the map (i.e. appear first at a high contour level). However, these
contours may have overall less detail as adjacent stacked bases are not fully separable and merge
together.
221
222

224 Figure 4. Q-scores averaged over entire nucleotides (Q_{nt}) in RNA maps and models from the EMDB at
225 four different resolutions. Q-scores are also averaged for the base (Q_{base}) and phosphate-sugar backbone
226 (Q_{bb}) groups in the nucleotides shown on the top row.
227
228
229 **Q-score vs. Resolution**
230
Q-scores can also be averaged across an entire model to represent an average resolvability
measure for the entire map. Such average Q-scores were plotted as a function of reported
resolution for a number of maps and models obtained from the EMDB. Figure 5 shows these
plots for two sets of maps and models, one set using only protein models, and the other set only
nucleic acids (RNA). The protein set includes the maps used in the EMRinger analysis[15], and
further adding 24 maps of Apoferritin and β-galactosidase at resolutions up to 1.54Å. In the
RNA set, a total of 52 maps and models were used at a range of resolutions ranging from 2.5Å

(the highest resolution of an RNA-containing map to date) to 5.4Å. The full sets are listed in
Tables 1 and 2. In both cases, the average Q-score correlates very strongly to reported resolution,
with R^2 of 0.90 for proteins and 0.89 for RNA. The R^2 quantifies the error in fitting the linear
function to the observed data; it is 1 for perfect correlation and 0 for no correlation. The high
values of R^2 in Figure 5 show that Q-scores closely capture the resolvability of atomic features in
cryoEM maps. Thus, average Q-scores from a properly fitted model may be useful to double-
check whether the reported resolution of the cryoEM map is accurate.
245
246

247
248   Figure 5. Model Q-scores compared to reported resolution for maps and models obtained from EMDB.
249   (A) Average Q-scores for atoms in proteins. (B) Average Q-scores for atoms in RNA.
250
251

Q-scores of Solvent Atoms

253
The X-ray Apoferritin model (PDB:3ajo) contains one protein chain, 229 oxygen (O) atoms
(from water) and 12 Mg atoms. A closeup on the map and model with two Mg and three O atoms
is shown in Figure 6. Q-scores calculated for each of these atoms correlate well with the contours
seen in the map. Of the four maps shown, three of them are cryoEM maps at near-atomic
resolutions (1.54Å, 1.65Å, and 1.75Å). The model used all cases comes from the X-ray map. It is
reassuring to see that some of the solvent atoms placed in the X-ray map can also be observed in
the cryoEM maps (e.g. Mg183, O280, O236). However, some of the solvent atoms (e.g. Mg184),
is not seen equally well in all three maps; for example, in the 1.54Å and 1.65Å maps, Mg184 has

low Q-score (0.12 and 0.03 respectively), and are not seen at the same map contour level where
the other solvent atoms are seen.
264
In this region of the map, the three water molecules shown in Figure 6 have high Q scores and
observable map contours. Along with Mg183, these provide evidence that cryoEM structures can
be used to identify locations of solvent molecules, much like with X-ray crystallography.
However, since Mg184 is only visible and has good Q-scores in only 2 of the 4 maps considered
here, differences between the cryoEM and X-ray maps can also be seen. Such differences may be
due to different affinities at some sites and/or different biochemical conditions across the
different data sets.
272
273

274
275 Figure 6. A close up in Apoferritin models showing solvent atoms (Mg and O from water), along with
276 calculated Q-scores in purple under each atom and nearby residue. The model comes from the X-ray map
277 (PDB:3ajo) shown in A. It was further refined into each of the three cryoEM maps, B-D.
278
279
280

Figure 7A shows distributions of Q-scores for solvent atoms in the X-ray map (PDB:3ajo). Most
solvent atoms have very high Q-scores of 0.9 and higher. Visual inspection confirmed that all
these solvent atoms can be seen in the X-ray map (2fo-fc), e.g. as shown in Figure 6A. Figure
7B,C shows distribution plots for the same model fitted to the cryoEM maps at 1.54Å and 1.75Å
resolution, using the rigidly fitted model and also after refinement (including solvent atoms) of
the rigidly fitted model using Phenix real-space refine[7].
287
For the rigidly fitted model, Q-scores of the solvent atoms are considerably lower than in the X-
ray map (Figure 7B). For example, in the 1.75Å cryoEM map, only 12 O atoms from water have
Q-scores of 0.9 and higher, and 32 have Q-scores of 0.8 to 0.9. In the 1.54Å map, 34 atoms have
Q-scores of 0.9 and higher, and another 34 have Q-scores of 0.8 to 0.9. Thus, water atoms are
less resolved in the cryoEM maps than in X-ray. It is possible that some of the solvent atoms
seen in the X-ray model may not be resolvable in the cryoEM maps or may be in different
positions.
295
To explore whether solvent atoms may have different positions in the cryoEM maps, Q-scores of
the solvent atoms were also calculated in the X-ray model after real-space refinement with
Phenix[7]. This refinement method moves solvent atoms towards higher map values, while keeping
them within reasonable distance of other atoms. The distributions in the Q-scores for solvent
atoms after this procedure are plotted in Figure 7B, C for the two cryoEM maps. Q-scores are
now higher; 142 water atoms in the 1.54Å map and 145 atoms in the 1.75Å map have Q-scores
of 0.8 and higher, compared to 225 water atoms in the X-ray map with Q-scores of 0.8 and
higher.
304
In the 1.54Å map, after refinement, water atoms with Q-scores 0.8 and higher moved between
0.1Å and 2.2Å, on average 0.54Å. In the 1.75Å map, the water atoms with Q-scores of 0.8 and
higher moved between 0.1Å and 1.6Å, on average 0.67Å. Although it is difficult to assess the
exact cause of the movements in these maps, it is reasonable to conclude that the water found in
cryoEM maps are real and potentially within experimental errors of their atom positions in both
X-ray and cryoEM structures.
311
312 In the above analysis, the water molecules were based on those originally observed in the X-ray
313 map. If one studies a *de novo* map, the identification of water molecules would require a protocol
314 used in modeling software, e.g. Phenix and Coot. In addition to such a protocol, Q-scores may be
315 used as an additional validation parameter to assist in the finding of water and ions.
316

319 Figure 7. Distribution of Q-scores for solvent atoms (water and Mg) in X-ray map (PDB:3ajo), and in two
320 cryoEM maps before and after refinement.

323 Radial Plots for Solvent Atoms

Radial plots in Figure 8 further characterize distances between solvent atoms (H_2O and Mg) and
other atoms including other water molecules (H_2O-H_2O), and also O and N atoms in protein,
H_2O-O and H_2O-N respectively. The radial plot for the X-ray model of Apoferritin (PDB:3ajo) is
shown in Figure 8A. This plot shows that H_2O-H_2O distances have a sharp peak at 2.8Å. A
similar peak is seen for distances between O atoms in water and O atoms in protein (H_2O-O).
Distances from Mg atoms to H_2O and to O have smaller peaks (since there are much fewer Mg
atoms in the model) at a distance of 2.2Å.

Radial plots for the X-ray model fitted to the 1.54Å and 1.75Å cryoEM maps are shown in
Figure 8B, C, considering only solvent atoms with Q-scores of 0.8 and higher after refinement. A
wider range H_2O-H_2O can be seen in both cases; instead of a sharp peak at 2.8Å, a broader peak

from ~2.4Å up to ~3.2Å can be seen. This seems to indicate that water-water distances in Cryo-
EM may vary. On the other hand, H_2O-O still have a main peak at 2.8Å after refinement in both
cryoEM maps, matching the peak seen in the X-ray map. Thus, H_2O-O distances are very similar
in X-ray and cryoEM maps in these examples, however the differences in H_2O-H_2O distances
suggests that there may be a difference in water organization around protein in cryoEM vs X-ray
maps.

346 Figure 8. Radial plots of distances from solvent atoms to other types of atoms. Oxygen atoms in water are
347 labeled H_2O, whereas oxygen/nitrogen atoms in protein are labeled O/N respectively.

Q-scores of Solvent Atoms at different resolutions

Finally, we looked at the resolvability and Q-scores of solvent atoms in cryoEM maps of
Apoferritin at different resolutions, as shown in Figure 9. The locations of the solvent atoms are
again taken from the X-ray model (PDB:3ajo). As Figure 9 shows, Mg183 appears resolved at
both 1.75Å and 2.3Å, with separable contours in both maps and high Q-scores (0.93 and 0.80).
However, the contours no longer have a symmetric spherical shape, indicating possibly more
variation in its position. In the 3.1Å map, the contour is no longer separable from that of the
nearby His65 residue, and the Q-score is also considerably lower (0.60). The water atoms are
similarly resolved in the 1.75Å and 2.3Å maps and contours around them can be seen, however
at 3.1Å and 3.9Å they can no longer be seen and Q-scores become very low (-0.44 to 0.38).

At 3.9Å resolution, both Mg atoms still have high Q-scores and thus high map values around
them, and they can be seen at a lower threshold. However, the map contours at these thresholds
do not necessarily separate them fully from the nearby residues (as again for Mg183).
Nevertheless, even at such lower resolutions (3Å-4Å), it appears that the larger solvent atoms
can still significantly influence the cryoEM map values, producing strong though more diffuse

peaks in the map. This may have some implications when creating or refining models in such
maps. Perhaps placement of solvent atoms should be considered for the model to be accurately
created and/or refined.
370
371

372
373 Figure 9. Solvent atoms from X-ray model (PDB:3ajo) in cryoEM maps at resolutions of 1.75Å to 3.9Å.
374 Q-scores are shown in purple below each atom. Nearby residues with Q-scores are also labeled (Q58,
375 E61, E62, H65, E27).
376
377
378 **Conclusions**
379
Q-scores can measure the resolvability of individual atoms in cryoEM maps, using atomic
positions and nearby map values. As was noted, this metric is closely related to the map-model
cross-correlation score, which is already widely used in the field to assess the fit of a model to a
map. However, the Q-score improves in two ways on the cross-correlation score: 1) it is
formulated so that it correlates to the resolution of the map and 2) it makes it applicable to small
features (individual atoms) while avoiding explicit masking. Aside from this, it is important to

note that nothing is assumed about the model itself, e.g. whether it has good stereochemistry; this
could be deduced with other scores such as the Molprobity score[20].
388
Q-scores averaged over entire models were shown to correlate very well with the reported
resolution of cryoEM maps containing both proteins and nucleic acids. Various visualizations
also showed that Q-scores indeed correlate well with the resolvability of individual atoms, and
also groups of atoms such as side chains. However, it still requires a model to first be fitted to or
built based on the cryoEM map. The score can be very useful to analyze the map and its
resolvability in different regions, and also test whether the model accurately interprets the map. It
could thus be useful as a map-model validation metric.
396
In this paper, several quantifiable observations were made with the help of Q-scores. For
example, when applied to atoms in protein side chains, the Q-score showed that resolvability of
certain types of side chains (Asp) drop faster than others (Asn) as a function of resolution. In the
case of nucleic acids, per-nucleotide Q-scores could be used to indicate whether stacked bases
are separable. Finally, Q-scores were also applied to water and other solvent atoms, helping to
confirm that water and other solvent atoms can indeed be resolved and placed in cryoEM maps
much as they are in X-ray crystallography.
404
405
406 **Experimental Methods**
407
408 Data: all the data for the analysis were drawn from EMDB and PDB. The EMDB 20026, 20027,
409 20028 maps were collected in Titan Krios electron microscope (Thermo Fisher) at 300 keV
410 equipped with Bioquantum energy filter and K2 director detector (Gatan). Images were recorded
411 in movie mode and corrected prior to image processing with Relion software[17]. The map
412 resolution was estimated from two independent maps with a total of 70,000 particle images.
413
414 Q-score calculation is implemented as a plugin to UCSF chimera, and is available from the
415 following website: https://cryoem.slac.stanford.edu/ncmi/resources/software.
416
417
418 **Acknowledgements**
419

420 This research has been supported by NIH grants (R01GM079429, P41GM103832, and
421 S10OD021600). Molecular graphics and analyses were performed with the UCSF Chimera
422 package. Chimera is developed by the Resource for Biocomputing, Visualization, and
423 Informatics at the University of California, San Francisco (supported by NIGMS
424 P41GM103311).

Author contributions

G.P. conceived Q-scores, implemented the software and performed all the testing. W.C. came up
with the term "Q-score". K.Z. collected the images and reconstructed the maps (EMDB: 20026,
20027, 20028). Z.S. and S.L. provided additional data (not shown) for testing the Q-score.
M.F.S. and W.C. contributed the discussion during the development. G.P. wrote the manuscript
with inputs from other authors.

Tables

Table 1. Maps from EMDB for which Q-scores of protein components are calculated for the plot
in Figure 5A. The table on the left shows the original maps used in the original EMRinger
analysis[15]. The table on the right contains a new set of maps of Apoferritin and β-galactosidase.

	EMD ID	Resolution	Q-score	EMRinger
1	2273	4.5	0.317	0.13
2	2278	3.5	0.429	3.26
3	2364	4.4	0.315	-0.47
4	2513	3.36	0.559	1.29
5	2677	4.5	0.260	-0.41
6	2762	3.4	0.485	2.09
7	2763	4	0.348	0.54
8	2764	3.75	0.384	0.9
9	2773	3.8	0.287	0.36
10	2787	3.4	0.469	1.85
11	2788	4.7	0.327	1.27
12	5160	3.2	0.515	2.18
13	5256	3.1	0.531	1.54
14	5391	4.9	0.216	0.2
15	5600	4.1	0.339	0.18
16	5623	3.2	0.558	3.05
17	5645	4.6	0.188	-0.05
18	5646	4.7	0.154	0.55
19	5678	4.5	0.357	0.49
20	5764	3.5	0.510	1.95
21	5778	3.27	0.335	0.56

	EMD ID	Resolution	Q-score	EMRinger
1	9865	1.54	0.850	8.22
2	9599	1.62	0.866	8.19
2	0144	1.65	0.854	8.14
3	20026	1.75	0.811	7.09
4	10101	1.84	2.814	8.44
5	0153	1.89	0.723	5.55
6	7770	1.9	0.713	5.41
7	9890	1.9	0.819	7.53
8	9914	2.01	0.843	7.21
9	4905	2.1	0.830	4.18
10	2984	2.2	0.620	3.58
11	4116	2.2	0.691	5.00
12	4415	2.2	0.691	4.38
13	8908	2.2	0.693	5.02
14	20027	2.32	0.750	5.53
15	4414	2.4	0.677	4.16
16	6480	2.6	0.638	3.99
17	4701	2.7	0.674	3.53
18	20227	2.85	0.484	1.55
19	20028	3.08	0.598	3.67
20	3854	3.15	0.661	4.61

22	5830	3.8	0.383	1.05
23	5886	5	0.340	0.8
24	5895	4.7	0.269	0.09
25	5896	5	0.246	0.06
26	5925	3.6	0.390	1.23
27	5995	3.2	0.540	2.04
28	6000	3.8	0.479	2.08
29	6035	3.5	0.461	0.96
30	6187	5	0.189	-0.71
31	6188	5	0.179	-0.16

21	5995	3.2	0.544	3.60
22	0140	3.9	0.482	3.44
23	2824	4.2	0.382	1.30

444 Table 2. Maps from EMDB containing RNA for which Q-scores vs. resolution are plotted in
445 Figure 5B.

	EMD ID	PDB File	Resolution	Q-score
1	7025	6az3-pdb-bundle2	2.5	0.699649
2	7025	6az3-pdb-bundle1	2.5	0.703164
3	8361	5t5h-pdb-bundle1	2.54	0.675016
4	0243	6hma	2.65	0.656016
5	7024	6az1	2.7	0.656345
6	6583	3jcs-pdb-bundle1	2.8	0.573334
7	20173	6ore-pdb-bundle1	2.9	0.615376
8	4638	6qul	3	0.649598
9	0600	6ole-pdb-bundle3	3	0.618584
10	0233	6hiz-pdb-bundle1	3.08	0.655904
11	4560	6qik-pdb-bundle1	3.1	0.606439
1	10068	6rzz-pdb-bundle1	3.2	0.584933
2	0101	6gzq-pdb-bundle1	3.28	0.564434
3	4125	5lze-pdb-bundle1	3.5	0.497689
4	4125	5lze-pdb-bundle2	3.5	0.538745
5	2938	4ug0-pdb-bundle1	3.6	0.54117
6	2938	4ug0-pdb-bundle2	3.6	0.503634
7	6559	3jcj-pdb-bundle1	3.7	0.471255
8	6559	3jcj-pdb-bundle2	3.7	0.421646
9	8620	5uyq-pdb-bundle1	3.8	0.42095
10	8620	5uyq-pdb-bundle2	3.8	0.426834
11	0076	6gwt-pdb-bundle2	3.8	0.417034
12	0076	6gwt-pdb-bundle1	3.8	0.411493

13	0192	6hcf-pdb-bundle2	3.9	0.524457
14	0192	6hcf-pdb-bundle1	3.9	0.506735
15	0192	6hcf-pdb-bundle3	3.9	0.408919
16	8279	5kps-pdb-bundle2	3.9	0.43481
17	8279	5kps-pdb-bundle1	3.9	0.437756
18	8618	5uyn-pdb-bundle2	4	0.383391
19	8618	5uyn-pdb-bundle1	4	0.385951
20	4080	5lmu	4	0.428174
21	2763	3j81_real_space_refined	4	0.453
22	2763	3j81	4	0.402465
23	4350	6g51	4.1	0.426988
24	8280	5kpv-pdb-bundle1	4.1	0.435967
25	8280	5kpv-pdb-bundle2	4.1	0.43031
26	643	6o7k	4.2	0.395112
27	20188	6ost-pdb-bundle1	4.2	0.3981
28	4382	6gc7	4.3	0.339055
29	0083	6gxp-pdb-bundle1	4.4	0.331033
30	4349	6g4w	4.5	0.311581
31	3133	5ady	4.5	0.361631
32	4351	6g53	4.5	0.338915
33	0104	6gzx-pdb-bundle1	4.57	0.356781
34	4083	5lmv	4.9	0.228577
35	3553	5mrf-pdb-bundle1	4.97	0.349647
36	8473	5tzs	5.1	0.183097
37	3661	5no2	5.16	0.326795
38	3662	5no3	5.16	0.310096
39	4122	5lzb-pdb-bundle1	5.3	0.28398
40	4427	6i7o-pdb-bundle1	5.3	0.292207
41	4075	5lmp	5.35	0.282151

447
448
449
450
451 **References**
452

1. Challenges for cryo-EM. *Nat. Methods* **15**, 985–985 (2018).

2. Henderson, R. *et al.* Outcome of the first electron microscopy validation task force meeting.

Struct. Lond. Engl. 1993 **20**, 205–214 (2012).

3. Kucukelbir, A., Sigworth, F. J. & Tagare, H. D. Quantifying the local resolution of cryo-EM

density maps. *Nat. Methods* **11**, 63–65 (2014).

4. Pintilie, G. & Chiu, W. Comparison of Segger and other methods for segmentation and rigid-

body docking of molecular components in cryo-EM density maps. *Biopolymers* **97**, 742–760

(2012).

5. Terwilliger, T. C., Adams, P. D., Afonine, P. V. & Sobolev, O. V. A fully automatic method

yielding initial models from high-resolution cryo-electron microscopy maps. *Nat. Methods*

15, 905 (2018).

6. Joseph, A. P., Lagerstedt, I., Patwardhan, A., Topf, M. & Winn, M. Improved metrics for

comparing structures of macromolecular assemblies determined by 3D electron-microscopy.

J. Struct. Biol. **199**, 12–26 (2017).

7. Afonine, P. V. *et al.* Real-space refinement in PHENIX for cryo-EM and crystallography.

Acta Crystallogr. Sect. Struct. Biol. **74**, 531–544 (2018).

8. Trabuco, L. G. *et al.* Applications of the molecular dynamics flexible fitting method. *J.*

Struct. Biol. **173**, 420–427 (2011).

9. Joseph, A. P. *et al.* Refinement of atomic models in high resolution EM reconstructions

using Flex-EM and local assessment. *Methods San Diego Calif* **100**, 42–49 (2016).

10. Pintilie, G., Chen, D.-H., Haase-Pettingell, C. A., King, J. A. & Chiu, W. Resolution and

Probabilistic Models of Components in CryoEM Maps of Mature P22 Bacteriophage.

Biophys. J. **110**, 827–839 (2016).

11. DiMaio, F., Zhang, J., Chiu, W. & Baker, D. Cryo-EM model validation using independent

map reconstructions. *Protein Sci.* **22**, 865–868 (2013).

12. Trabuco, L. G., Villa, E., Schreiner, E., Harrison, C. B. & Schulten, K. Molecular dynamics

flexible fitting: a practical guide to combine cryo-electron microscopy and X-ray

crystallography. *Methods San Diego Calif* **49**, 174–180 (2009).

13. DiMaio, F., Tyka, M. D., Baker, M. L., Chiu, W. & Baker, D. Refinement of protein

structures into low-resolution density maps using rosetta. *J. Mol. Biol.* **392**, 181–190 (2009).

14. Hryc, C. F. *et al.* Accurate model annotation of a near-atomic resolution cryo-EM map. *Proc.*

Natl. Acad. Sci. 201621152 (2017). doi:10.1073/pnas.1621152114

15. Barad, B. A. *et al.* EMRinger: side chain-directed model and map validation for 3D cryo-

electron microscopy. *Nat. Methods* **12**, 943–946 (2015).

16. Pintilie, G. & Chiu, W. Assessment of structural features in Cryo-EM density maps using

SSE and side chain Z-scores. *J. Struct. Biol.* **204**, 564–571 (2018).

17. Zivanov, J. *et al.* New tools for automated high-resolution cryo-EM structure determination

in RELION-3. *eLife* **7**, e42166 (2018).

18. Dunitz, J. D., Schomaker, V. & Trueblood, K. N. Interpretation of atomic displacement

parameters from diffraction studies of crystals. *J. Phys. Chem.* **92**, 856–867 (1988).

19. Trueblood, K. N. *et al.* Atomic Dispacement Parameter Nomenclature. Report of a

Subcommittee on Atomic Displacement Parameter Nomenclature. *Acta Crystallogr. A* **52**,

770–781 (1996).

20. Chen, V. B. *et al.* MolProbity: all-atom structure validation for macromolecular

crystallography. *Acta Crystallogr. D Biol. Crystallogr.* **66**, 12–21 (2010).

499
500
501