Organoid-derived Duodenum Intestine-Chip enables preclinical drug assessment in a human relevant system

Magdalena Kasendra[1,4,*], Raymond Luc[1,4], Jianyi Yin[2], Dimitris V. Manatakis[1], Athanasia Apostolou[1,3], Laxmi Sunuwar[2], Jenifer Obrigewitch[1], Geraldine A. Hamilton[1], Mark Donowitz[2], and Katia Karalis[1]

[1] Emulate Inc., 27 Drydock Avenue, Boston, MA 02210, USA
[2] Department of Medicine, Division of Gastroenterology, Johns Hopkins University School of Medicine, Baltimore, MD 21205, USA
[3] Graduate Program, Department of Medicine, National and Kapodistrian University of Athens, Greece
[4] Equal first co-authors
*Corresponding author

Abstract

Induction of intestinal drug metabolizing enzymes can complicate the development of new drugs, owing to potential to cause drug-drug interactions (DDIs) leading to changes in pharmacokinetics, safety and efficacy. The development of a human relevant model of the adult intestine that accurately predicts CYP450 induction could help address this challenge as species differences preclude extrapolation from animals. Here, we combined organoids and Organ-Chip technology to create a human Duodenum Intestine-Chip that emulates intestinal tissue architecture and functions, that are relevant for the study of drug transport, metabolism, and DDI. Duodenum Intestine-Chip demonstrates the polarized cell architecture, intestinal barrier function, presence of specialized cell subpopulations, and *in vivo*-relevant expression, localization, and function of major intestinal drug transporters. Notably, in comparison to Caco-2, it displays improved CYP3A4 expression and induction capability. This model could enable improved *in vitro* to *in vivo* extrapolation for better predictions of human pharmacokinetics and risk of DDIs.

Introduction

A challenge in drug development is low oral bioavailability and poor pharmacokinetics caused by drug-drug interactions of orally administered drug. One key driver can be high affinity for drug transporters and activity of metabolic enzymes in the human intestine (Dietrich, Geier et al. 2003, Thummel 2007, Shugarts and Benet 2009, Peters, Jones et al. 2016). After the decades of research, the hepatic drug clearance is well-understood and relatively well predicted by pre-clinical models. While accurate prediction of first-pass extraction of xenobiotics in human intestinal epithelium still remains elusive. This is due to a number of confounding factors that affect oral drug absorption including the properties of the compound (solubility, permeability), physiology of the intestinal tract (transit time, blood flow), and patient phenotype (including age, gender, polymorphism in drug metabolizing enzymes, disease states) (Pang 2003). Species differences in the isoforms, regional abundances, differences in

substrate specificity of drug metabolism enzymes (Martignoni, Groothuis et al. 2006,
Paine, Hart et al. 2006, Komura and Iwaki 2011) and transporters (Tucker, Milne et al.
2012, Groer, Bruck et al. 2013), and mechanism regulating transcriptional activation
(LeCluyse 2001, Mackowiak, Hodge et al. 2018), precludes accurate extrapolation of
the data from animal models to the clinic. In mice, for example, there are 34
cytochromes CY450 in the major gene families involved in drug metabolism, i.e., the
Cyp1a, Cyp2c, Cyp2d, and Cyp3a gene clusters, while in humans, there are only
eight(Nelson, Zeldin et al. 2004). Interestingly three human enzymes, CYP2C9,
CYP2D6, and CYP3A4, account for ~75% of all reactions, with CYP3A4 being the
single most important human CYP450 accounting for ~45% of phase 1 drug
metabolism in humans (Guengerich 2008). In addition, the expression levels of many of
the major human CYP450 enzymes and drug transporter (which determine levels and
variability in drug exposure) are controlled by multiple transcription factors, primarily the
xenosensors constitutive androstane receptor (CAR), pregnane X receptor (PXR), and
aryl hydrocarbon receptor (AhR). These nuclear receptors also exhibit marked species
differences in their activation by drugs and exogenous chemicals (Mackowiak, Hodge et
al. 2018). For example, rifampicin and SR12813 are potent agonists for human PXR
(hPXR) but not for mouse PXR (mPXR), whereas the potent mPXR agonist 5-pregnen-
3β-ol-20-one-16α-carbonitrile (PCN) is a poor agonist for hPXR (Kliewer, Moore et al.
1998). On the other hand, 6-(4-chlorophenyl)imidazo[2,1-b][1,3]thiazole-5-
carbaldehyde-O-(3,4-dichlorobenzyl)oxime (CITCO) is a strong agonist for human CAR
(hCAR) but not mouse CAR (mCAR) (Maglich, Parks et al. 2003), while 1,4-bis-[2-(3,5-
dichloropyridyloxy)]benzene,3,3′,5,5′-tetrachloro-1,4-bis(pyridyloxy)benzene
(TCPOBOP) is more selective for mCAR than hCAR. Such species differences together
with the complex interplay between drug metabolizing enzymes and drug transporters in
the intestine and the liver, as well as, the overlap of substrate and inhibitor specificity
(Shi and Li 2014), make it difficult to predict human pharmacokinetics when there is
intestinal metabolism and drug transport involved due to the lack of predictive pre-
clinical models that reflect the human biology that allow researchers to study it in a
human relevant system .
75
Numerous *in vitro* systems have been developed and applied routinely for
characterization of distribution and prediction of absorption, distribution, metabolism,
and excretion (ADME) of potential drug candidates in humans. Among these is Caco-2
monolayer culture on a transwell insert, which is the one of most widely used across the
pharmaceutical industry as an *in vitro* representation of the human small intestine.
However, inherent limitations, such as lack of *in vivo* relevant three-dimensional
cytoarchitecture, lack of appropriate ratio of cell populations, altered expression profiles
of drug transporters and drug metabolizing enzymes, especially CYP450s, and aberrant
CYP450 induction response, challenge the use of these model for predicting human
responses in the clinic (Sun, Chow et al. 2008).
86
A promising alternative to conventional cell monolayer systems emerged with the
establishment of the protocols for generation of three-dimensional intestinal organoids
(or enteroids) from human biopsy specimens (Sato, Vries et al. 2009, Sato, Stange et

al. 2011, Eglen and Randle 2015, Liu, Huang et al. 2016). Using these methods,
organoids derived from all regions of the intestinal tract can be established (Sato,
Stange et al. 2011, Wang, Yamamoto et al. 2015) and applied into different areas of
research including organ development, disease modeling, and regenerative medicine
(Fatehullah, Tan et al. 2016). However, the characterization of the pharmacokinetic
properties of this system, as well as the its validation for the use in drug discovery and
development, is still very limited (Dekkers, Wiegerinck et al. 2013, van de Wetering,
Francies et al. 2015, Zhang, Zhao et al. 2017, Zhao, Zeng et al. 2017, Vlachogiannis,
Hedayat et al. 2018). One potential limitation could be due to the substantial technical
challenges associated with the use of this technology for ADME applications. The 3D-
spheroidal architecture of the organoid restricts access to its lumen, which is crucial for
assessing intestinal permeability or drug absorption. Indeed, exposure of the apical cell
surface of intestinal organoid to compounds requires the use of time-consuming and
labor-intensive procedures, such as microinjection. The presence of relatively thick gel
of extracellular matrix (Matrigel^{TM}) surrounding organoids, might limit drug penetration.
While heterogeneity of organoids in terms of their size, shape, and viability, can also
impede studies in ADME and robust results (Fatehullah, Tan et al. 2016).

In addition, none of these models have fully recapitulated critical aspects of an organ
microenvironment such as the presence of microvasculature, mechanical forces of fluid
flow (shear stress) and peristalsis, all of which contribute to capturing the complex and
dynamic nature of *in vivo* tissue function (Gayer and Basson 2009). For these reasons,
there is a need for new systems for predicting human ADME and determining risk for
drug-drug interactions mediated by intestinal CYP450s and drug transporters in the
clinic.

We have recently developed a human Duodenum Intestine-Chip that combines healthy
intestinal organoids with our Organs-on-Chips technology (Kasendra, Tovaglieri et al.
2018) to overcome the existing limitations of the current systems. Here, we
demonstrated that the Duodenum Intestine-Chip provides a more human-relevant
model, in respect to organoids, as supported by the comparison of their global RNA
gene expression profiles, and that it can be applied for the study of drug transport, drug
metabolism, and drug-drug interactions. Presence of the mechanical forces, which are
applied into this system in order to recapitulate the blood flow and shear stress, showed
to improve the formation of the right cell cytoarchitecture and the increased appearance
of intestinal microvilli on the apex cell surface. In addition, Duodenum Intestine-Chip
supported successful maturation of all major intestinal cell types in the physiologically
relevant ratios and low paracellular permeability. Importantly for its application into
pharmacokinetic studies, it showed closer to *in vivo* expression of drug uptake and
efflux transporters, as compared to Caco-2-cells based system. Additionally, we were
able to prove that it the system supports the correct luminal localization and functional
activity of MDR1 (P-gp), as well as high expression of cytochrome CYP450 (CYP) 3A4 -
close to the levels observed in human duodenal tissue. Importantly, exposure of the
Duodenum Intestine-Chip to known CYP450 inducers in humans, such as rifampicin
and 1, 25-dihydroxyvitamin D3, resulted in the significant increase of both mRNA and

protein levels of CYP3A4. Our results indicate that the organoid-derived human tissue
combined with Organs-on-Chips technology could provide a robust human-relevant
system the assessment of CYP450 metabolism, drug transporter effects, and risk for
potential drug-drug interactions in preclinical testing.
139
Development of the adult Duodenum Intestine-Chip
We have previously developed a human organoid-based Intestine-Chip, referred to at
the time as the "Small Intestine-on-a-Chip", which combined the use of intestinal
organoids and Organ-Chips (Kasendra, Tovaglieri et al. 2018) and provides a system to
model intestinal biology. Here we sought to establish the adult Duodenum Intestine-
Chip to serve as a model for preclinical assessment of drug transport and metabolism.
In brief, we first established cultures of organoids (Figure 1A; top) derived from the
endoscopic biopsies of three different healthy adult donors; the organoids were then
dissociated into fragments and seeded on the ECM-coated porous flexible
poly(dimethylsiloxane) (PDMS) membrane of the chips (Figure 1B; 1: indicates the
epithelial tissue). Primary human intestinal microvascular endothelial cells (HIMECs,
Cell Biologics), derived from the human small intestine (Figure 1A; bottom), were used
to populate the other surface of the PDMS membrane in the vascular channel (Figure
1B; 5: indicates the endothelial cells). Next, the Duodenum Intestine-Chips were
perfused continuously through luminal and vascular compartment with 30 µl / hr of fresh
media. Once the epithelial monolayers reached confluency they were subjected to cyclic
mechanical strain (10% strain, 0.2 Hz) in order to emulate intestinal peristalsis and the
physiologically relevant mechanical forces associated with it.
158
We then assessed the effect of applying mechanical forces (via fluid flow and stretch)
on the phenotypic characteristics of the human primary intestinal cells in the chips. We
used multiple endpoints including immunofluorescent staining for apical (villin) and
basolateral (E-cadherin) cell surface markers and scanning electron microscopy (SEM)
for the presence of apical microvilli. In line with our previous findings in the Caco-2
Intestine-Chip(Kim, Huh et al. 2012), exposure of the Duodenum Intestine-Chip to flow
for 72 hours resulted in accelerated cell polarization and formation of apical microvilli
(Supplementary Figure 1). We found that cells cultured under static conditions in the
Organ-Chips formed a flat (14.8 ± 2.6 µm) monolayer of epithelial cells of squamous
appearance and were characterized by poorly defined cell-cell junctions
(Supplementary Figure 1A) and sparsely distributed microvilli (Supplementary Figure
1B). In contrast, cells cultured under flow (30 µLh^{-1}) with or without concomitant
application of cyclic stretch (10%, 0.2 Hz), exhibited a well-polarized and cobblestone-
like morphology with increased cell height (27.0 ± 1.3 µm), strongly delineated junctions,
and dense microvilli-like structures. In line with our previous findings with Caco-2 cells,
the application of constant flow (shear stress) was critical for promoting maturation of a
well-polarized epithelium, while short-term application of cyclic strain did not show any
additional effect. In addition, more prolonged exposure of cells to flow and cyclic strain
resulted in the spontaneous development of epithelial undulations ("villi-like structures")
extending into the lumen of the epithelial channel and covered by continuous brush
border (Figure 1C). Immunofluorescence confocal analysis confirmed the establishment

of confluent epithelial and endothelial monolayers across the entire length of the chip
(Figure 1D), with well-defined epithelial tight junctions, as demonstrated by *in vivo*-
relevant ZO-1 protein staining and endothelial adherent junctions visualized using
antibody against VE-cadherin (green). Importantly, these culture conditions resulted in a
time-dependent improvement of intestinal permeability as indicated by the low
permeability coefficient (Papp) of fluorescently labeled dextran recorded in the
Duodenum Intestine-Chips generated from 3 different human donors (Figure 1E). This
data indicates that this model supports the formation of a functional barrier with *in vivo*
relevant cytoarchitecture, cell-cell interactions, and permeability parameters.

To confirm differentiation of the organoid-derived primary enterocytes to the relevant
subpopulations of cells as found *in vivo*, we assessed expression levels of cell type
specific markers, including alkaline phosphatase (ALPI) for absorptive enterocytes,
mucin 2 (MUC2) for goblet cells, chromogranin A (CHGA) for enteroendocrine cells, and
lysozyme (LYZ) for Paneth cells in Duodenum Intestine-Chips from 3 different human
donors. We compared expression levels of these genes to those found in freshly
isolated adult duodenal tissue (Duodenum). As shown (Figure 2A), expression of all the
above genes, with the exception of LYZ, was increased over time in culture. Notably,
markers such as alkaline phosphatase and mucin 2 reached similar levels in Duodenum
Intestine-Chips to those detected in RNA isolated directly from the human duodenum
after 8 days in culture. The opposite trend depicted in the expression of LYZ suggests a
decrease in Paneth cells in line with increased differentiation of the villus positive
epithelium and a decrease in progenitor/stem cells population. In addition,
immunostaining for the characterization of all major differentiated intestinal cell types,
followed by quantification using confocal microscopy, showed physiologically relevant
relative ratios of each cell-type to one another (Figure 2C). The ratios of the cell
populations in the chips were close to those reported following histopathological
analysis of sections from the human duodenum (Figure 2D)(Karam 1999). Taken
together, these results demonstrate the successful establishment of an adult Duodenum
Intestine-Chip that closely recreates the barrier function and multilineage differentiation
of adult human intestinal tissue.

Transcriptomic comparison of the Duodenum Intestine-Chip versus organoids
To further verify whether Duodenum Intestine-Chip faithfully recapitulates human adult
duodenum tissue and to better understand how much it differs from the organoids from
which it's derived, we performed RNA-seq analysis. We compared global RNA
expression data obtained from: i) duodenum organoids (Organoids; n=3) cultured in
conventional plastic-adherent Matrigel™ drop overlaid with growth medium; ii)
Duodenum Intestine-Chips derived from organoids (Duodenum Intestine-Chip; n=3) in
the presence of constant flow and stretch; iii) human adult duodenum tissue (adult
duodenum; n=2; full-thickness samples) (Supplementary Table 1). We annotated
13,735 genes in the genome and performed differential gene expression analysis
(DGE). For the DGE analysis, we used the "limma" R package and applied the widely
accepted thresholds: adjusted p-value<0.05 and |log2FoldChange| > 2 in order to select
the differentially expressed genes(Ritchie, Phipson et al. 2015).

First, we examined the differential gene expression in organoids compared to human
adult duodenum tissue. Out of the 13,735 genes annotated in the genome, 1437 were
found to be significantly differentially regulated among these samples: 562 and 875
genes were respectively up- and downregulated (Supplementary Figure 2A and
Supplementary Table 2). Next a functional enrichment analysis was performed utilizing
the PANTHER classification system to highlight biological processes, i.e., significantly
enriched gene ontology (GO) terms within these gene sets (Ashburner, Ball et al. 2000,
Mi, Muruganujan et al. 2013, The Gene Ontology 2017). The majority of differentially
expressed genes belonged to pathways related to digestion, extracellular matrix
organization, angiogenesis, cell adhesion, tissue development, and cell responses to
drugs and xenobiotics (Supplementary Figure 2A). This comparison allowed us to
identify genes responsible for global transcriptomic differences between organoids
technology and native human tissue and highlight biological functions which could be
affected by the observed differences.
We applied similar type of analysis, to the one described above, to compare global gene
expression profile of Duodenum Intestine-Chip with adult duodenum. This analysis
resulted in the identification of 1023 differentially expressed genes that were
significantly up- (382 genes) or downregulated (641 genes). The fact, that the
significantly smaller number of differentially expressed genes was identified in this
analysis as compared to the DGE performed between organoids and adult duodenum
tissue, suggests that combining organoids with Organs-on-Chips enables a better
emulation of the human duodenum tissue (Supplementary Figure 2B and
Supplementary Table 3). Functional enrichment analysis highlighted biological
processes such as protein synthesis and targeting, cell cycle and cell proliferation
underlying the differences between Duodenum Intestine-Chip model and native human
tissue.
Next, in order to determine which genes are responsible for a closer similarity of
Duodenum Intestine-Chip system to human duodenum tissue in comparison to
organoids, we carried out additional DGE analysis. This time we assessed the
differences between Duodenum Intestine-Chip and organoids, and examined these
differences relative to those that exist when comparing the adult human duodenal tissue
and organoids (Duodenum Intestine-Chip versus organoids and human adult duodenum
versus organoids). We identified genes that were significantly up- and downregulated in
the Duodenum Intestine-Chip relative to organoid culture (Figure 3—source data 1) and
identified the proportion of these genes that were also significantly different in adult
duodenal tissue relative to organoids (Figure 3—source data 2 and 6). We found 305
genes which were common for Duodenum Intestine-Chip and adult duodenum but
different from organoids, representing 39.25% of differentially expressed genes in
Duodenum Intestine-Chip versus organoids comparison, and 21.22% of differentially
expressed genes in human adult duodenal tissue versus organoids comparison (Figure
3A and Figure 3—source data 3). To further explore overlapping genes, we used the
Gene Ontology and Kyoto Encyclopedia of Genes and Genomes (KEGG) Pathway
Analysis to examine biological processes and pathways that are enriched in this gene
list (Kanehisa and Goto 2000, Thomas, Campbell et al. 2003). In addition, we used the
well-established Reduce and Visualize Gene Ontology (REVIGO) tool to reduce

redundancies by clustering semantically related GO terms and outputting them as a
scatter plot for visualization (Eden, Lipson et al. 2007, Supek, Bosnjak et al. 2011). As a
result, the list of 305 overlapping genes was associated with 117 significant GO terms,
which were reduced to 74 unique GO terms in REVIGO (Figure 3—source data 4).
Notably, biological processes enriched in this gene set were associated with important
intestinal functions such as digestion and transport of nutrients and ions, metabolism,
detoxification, as well as tissue and tube development (Fig 3B).
Additionally, KEGG Pathway analysis identified the total of 7 significantly enriched
pathways (threshold FDR p-value <0.05) among the overlapping genes (Figure 3C).
The corresponding fold enrichments of these pathways are shown in Figure 3C.
Importantly, many of these pathways were linked to drug metabolism, such as –
cytochrome CYP450 (associated with 10 genes), metabolism of xenobiotic by
cytochrome CYP450 (associated with 12 genes), and other important pathways related
to nutrients absorption and digestion, such as mineral absorption (associated with 12
genes), fat digestion and absorption (associated with 9 genes), protein digestion and
absorption (associated with 11 genes) (Figure 3C). To more closely examine gene
signatures that belong to these pathways, we used log (FPKM) values generated from
the RNA-seq dataset, and generated heatmaps for differentially expressed genes that
fall under the 7 significantly enriched pathways (Figure 3D). A large number of genes
involved in the nutrient absorption, digestion, as well as, metabolism of xenobiotics was
highly expressed in Duodenum Intestine-Chip (Chip) and adult duodenum tissue (Adult
Duo) but expressed at the low level in organoids.
Cumulatively, these data demonstrated an increased similarity in the global gene
expression profile between Duodenum Intestine-Chip and human adult duodenum
tissue in comparison to the organoids from the same donor.
295
Intestinal drug transporters and MDR1 efflux activity in the Duodenum Intestine-
Chip
Encouraged by the transcriptomic analysis detailed above, that showed a clear
advantage of Duodenum Intestine-Chip over organoids from which it was derived and a
closer to *in vivo* expression of genes related to absorption, transport, and metabolism
we next sought to focus further on the expression, localization and function of main
intestinal drug transporters. We determined their expression and localization in the
Duodenum Intestine-Chips established from the organoids of three independent donors
by qRT-PCR and immunofluorescent imaging. We have compared the observed gene
expression levels with the values obtained for the freshly isolated human duodenal
tissue samples (Duodenum) and the previously described Intestine-Chip model based
on the use of Caco-2 cells (Caco-2 Intestine-Chip) (Figure 4A). Averaged gene
expression levels of efflux (MDR1, BCRP, MRP2, MRP3) and uptake (PepT1,
OATP2B1, OCT1, SLC40A1) drug transporters in the Duodenum Intestine-Chips were
close to those observed in the human duodenal tissue. Similar pattern of gene
expression was revealed for Caco-2 Intestine-Chip, suggesting the existence of a good
correlation between the chip and human *in vivo* tissue. However, in case of the latter
model, Caco-2 Intestine-Chip, a couple of important organic anion and cation
transporters, including OATP2B1 and OCT1, showed a markedly increased expression

in comparison to human duodenum. This confirmed the maintenance of already well-
known differences between the cancer-derived cell line and normal human intestinal
tissue in Intestine-Chip system(Maubon, Le Vee et al. 2007, Olander, Wisniewski et al.
2016). Moreover, although not statistically significant, but notable differences in the
expression of MDR1, BCRP and PEPT1 were present in Caco-2 Intestine-Chip in
comparison to Duodenum. In line with the trends previously reported by others(Sun,
Lennernas et al. 2002, Maubon, Le Vee et al. 2007, Harwood, Achour et al. 2016), a
3.5-fold higher expression of MDR1, 3.6-fold and 24-fold lower expression of BCRP and
PEPT1, respectively, were found in Caco-2 cells compared with human duodenum. On
the other hand, much smaller differences in respect to native human tissue were
observed for Duodenum Intestine-Chip: with ~1.5-fold and ~2.6-fold increased
expression of MDR1 and BCRP, respectively, and no differences noted in the
expression of PEPT1. These results demonstrate that by combining duodenal organoids
with Organ-on-Chips technology we enabled a closer emulation of human *in vivo* tissue
than it was possible with the previously described Caco-2 Intestine-Chip or organoids
alone.
331
We further demonstrated *in vivo* relevant localization of the luminal efflux pumps,
MDR1, more commonly referred to as P-gp or P-glycoprotein (Figure 4B) and BCRP
(Supplementary Figure 3A), as well as the uptake Peptide Transporter 1 (PEPT1) in
organoids-derived Duodenum Intestine-Chips (Supplementary Figure 3B). All three
transporters showed to co-distribute together with villin, a marker specific for apical cell
membrane, at the intestinal cell brush border in Duodenum Intestine-Chip. This was
confirmed on the cross-sectional confocal images of the duodenal epithelium cultured
on chip as a co-localization (merge channel; white) between the fluorescent signal for
MDR1, BCRP or PEPT1 and anti-villin staining. Similarly, plots depicting the distribution
of the fluorescent signal across the cell Z-axis in these cross-sections revealed the
presence of the significant overlap between the fluorescent spectra of each transporter
and villin (Figure 4B and Supplementary Figure 3). Notably, the physiologically-relevant
localization of MDR1, BCRP1 and PEPT1 at the luminal surface of intestinal epithelium
was confirmed at two different time-points of Duodenum Intestine-Chip culture – when
the cells have formed a confluent monolayer (around day 4 of culture) and after the villi-
like structures morphogenesis has occurred (at day 8 of culture). The MDR1 activity
was confirmed by measuring the intracellular accumulation of rhodamine 123 in the
presence and absence of specific MDR1 inhibitor, vinblastine, across Duodenum
Intestine-Chips and was compared to the expression in the Caco-2 Intestine-Chip model
(Figure 4C). Addition of the inhibitor induced an ~2-fold increase in intracellular
accumulation of rhodamine (1.84-fold increase in Duodenum Intestine-Chip and 2.14-
fold increase in Caco-2 cells-based model), confirming active MDR1 efflux pumps in
both cell systems.
355
356 **Drug-mediated CYP3A4 induction in the Duodenum Intestine-Chip**
357 Induction of CYP450 drug metabolizing enzymes in human intestine is a major concern
358 the pharmaceutical industry, as it is known to impact the pharmacokinetics and
359 bioavailability of various orally-administered drugs, as well as, mediate drug-drug

interactions. We therefore, evaluated the capability of our Duodenum Intestine-Chip to be applied for CYP3A4 induction studies and to help identify risk for drug-drug interactions in the clinic. This is not feasible in pre-clinical species such as rat and dog due to marked species differences in the expression and regulation of cytochromes P450 as well as substrate specificity of the nuclear receptors, such as pregnane X receptor (PXR), responsible for the transcriptional regulation of CYP3A4 and several drug transporters. Caco-2 Intestine-Chip has been previously reported to possess an increased activity of the CYP450 enzymes when compared to the conventional static culture of Caco-2 cells on transwell insert (25). However, the gene expression level of CYP3A4 measured in this system is significantly lower than in the adult human intestine (Figure 5A), limiting its application for pharmaceutical research, specifically pharmacokinetic evaluation. In the present study we demonstrate that, in comparison with the Caco-2 cells based system, our Duodenum Intestine-Chip expressed CYP3A4 at the much higher gene (~6000-times higher, p <0.0001) (Figure 5A) and protein level (Figure 5B), reaching the expression similar to the one observed in the adult human duodenum.

In order to assess the drug-mediated CYP3A4 induction potential in the Duodenum Intestine-Chip, we exposed it to rifampicin (RIF) and 1,25-dihidroxyvitamin D3 (VD3), both of which are known prototypical CYP3A4 inducers. Treatment of Duodenum Intestine-Chips from all three independent donors with RIF and VD3 resulted in 5.3-fold and 4.1-fold induction of CYP3A4 expression, respectively, relative to that in the DMSO-treated controls (Figure 5C, left). Alternatively, Caco-2 Intestine-Chips were shown to respond to vitamin D3, but not RIF treatment (Figure 5C, right), which is consistent with the previously published reports (Schmiedlin-Ren, Thummel et al. 1997, Ozawa, Takayama et al. 2015, Negoro, Takayama et al. 2016). Dramatically low baseline expression of CYP3A4 in Caco-2 cells derived system showed to increase by 316-fold in the presence of VD3. However, it did not reach the levels observed in the human tissue and remained undetectable at the protein level as assessed by Western Blot analysis (Figure 5D, right). The differences between drug-mediated CYP3A4 induction potency of these two systems could be attributed to differences in gene expression levels of intestinal nuclear receptors, including PXR and vitamin D receptor (VDR). The gene expression levels of PXR and VDR in Duodenum Intestine-Chip were significantly higher and closer to human duodenal tissue than those in Caco-2 Intestine-Chip. An evident lack of PXR, which is known to be one of the key transcriptional regulators of CYP3A4 induction in humans, in Caco-2 cells may explain the differences observed between the Duodenum Intestine-Chip, and other Caco2–based systems including chips with Caco2. All together our findings demonstrate that the Duodenum Intestine-Chip represents a superior and more appropriate model for predicting CYP3A4-mediated drug-drug interactions in the intestine for oral drugs than Caco-2 cell-based models.

Discussion
A number of *in vitro* and animal models have been developed and routinely applied for characterization of distribution and prediction of absorption, metabolism, and excretion

(ADME) of xenobiotics in humans. Among these, one of the most widely used models is
the conventional Caco-2 monolayer culture, considered as the current "gold standard" in
studying intestinal disposition of drugs *in vitro*. However, inherent limitations, such as
lack of in vivo relevant 3D cytoarchitecture, lack of appropriate ration of cell populations,
altered expression profiles of drug transporters and drug metabolizing enzymes,
especially CYP450s, and aberrant CYP450 induction response, challenge the use of
these model for predicting ADME in the clinic. On the other side of the spectrum, animal
models, while retaining proper physiological conditions, exhibit species differences in
both drug metabolism and drug transport, as well as substrate specificity for nuclear
receptors regulating CYP450s and transporters. For these reasons, there is a need for
new systems for predicting human ADME and determining risk for drug-drug
interactions mediated by intestinal CYP450s and drug transporters.
417
Current preclinical models are not able to fully recapitulate the complex nature and
function of human intestinal tissue, leading to a limited accuracy and poor predictability
in drug development. In this study, we leveraged Organs-on-Chips technology and
primary organoids to emulate multicellular complexity, physiological environment, native
intestinal tissue architecture and functions and to create an alternative human-relevant
model for the assessment of ADME of orally administered drugs. Similarly, to our
previous study (Kasendra, Tovaglieri et al. 2018) morphological analysis of the
Duodenum Intestine-Chip confirmed the establishment of intact tissue-tissue interface
formed by organoids-derived epithelium and small intestinal microvascular endothelial
cells. The application of the mechanical stimulation in the form of continuous luminal
and vascular flow showed to exert the beneficial effect on the intestinal tissue
architecture. Increased epithelial cell height, acquisition of cobblestone-like cell
morphology, formation of well-defined cell-cell junctions, and dense intestinal microvilli
was attributed to the presence of flow. We confirmed the development of comparable
levels of intestinal barrier function to hydrophilic solute (dextran) across Duodenum
Intestine-Chip cultures established from organoids, which were isolated from three
independent donors. This functional feature of intestinal tissue becomes critical when
studying drug uptake, efflux, and disposition in polarized organ systems. Although many
studies have demonstrated that Caco-2 transwell allow good prediction of transcellular
drug absorption, they showed to be unreliable in the assessment of passive diffusion of
polar molecules such as hydrophilic drugs and peptides. This is related to much smaller,
in respect to small intestinal villi, effective area of the monolayer and its higher tight
junctional resistance (Press and Di Grandi 2008, Kumar, Karnati et al. 2010). Organoids
have quickly become a model of interest in drug discovery and development
applications, yet the use of organoids for the studies of intestinal permeability has been
linked with major technical difficulties related to their three-dimensional structure, limited
access to lumen, and the need for the use of sophisticated microinjection techniques for
the application of the compound at the apical cell surface. Therefore, Duodenum
Intestine-Chip that emulates complex intestinal tissue architecture, as evidenced by the
presence of three-dimensional villi-like structures on scanning electron micrographs
acquired on day 8 of fluidic culture, and allows a direct exposure of apical cell surface to
test compounds represents a major technological advantage over Caco-2 transwell and

organoids and might provide a valid alternative for the preclinical studies of intestinal
drug absorption.
452
Immunofluorescent imaging and gene expression analysis demonstrated that
Duodenum Intestine-Chip possess specialized intestinal cell subpopulations, that are
absent in tumor-derived cell lines including Caco-2. Importantly, they are present in the
chip at the physiologically relevant ratios, similar to the ones observed in the native
human duodenum. Expression of the markers specific for the intestinal types that
naturally reside in the villus compartment – alkaline phosphatase for absorptive
enterocytes, mucin 2 for goblet cells, chromogranin A for enteroendocrine cells –
showed to increase up to day 8 of Duodenum Intestine-Chip culture. This has been
accompanied by the decline in the expression of genes specific for crypt-residing cells –
lysozyme for Paneth cells and Lgr5 for stem cells (data not shown) suggesting the
acquisition of terminally differentiated cell phenotype by all of the cells present in the
chip. This is in line with well-known effect of Wnt3a withdrawal on organoids-derived
primary intestinal epithelium (Sato, Vries et al. 2009, Sato, Stange et al. 2011). Because
human intestinal cells do not produce significant amounts of Wnt3A, EGF, or several
other growth factors essential for stem cell division and cell proliferation, removal of
Wnt3A supplementation leads to a loss of LGR5-positive cells, decreased cell
proliferation, appearance of secretory cell lineages, including goblet and
enteroendocrine cells, and the transformation of immature crypt-like enterocytes into
differentiated nutrient-absorptive cells. The presence of differentiated cell
subpopulations is critical for modelling various aspects of intestinal biology and function,
including mucus production, antimicrobial response, host-microbiome interaction,
secretion of intestinal hormones, nutrients digestion and absorption. Given that we have
shown that the Duodenum Intestine-Chip recapitulates faithfully the physiologically
relevant ratios of these cells it could be readily applied beyond ADME applications in
development of new approaches and therapeutics that targets specific cell populations.
For example, it could serve to test to target relevant biology of Paneth cells, which
through a series of genetic studies have been implicated in inflammatory bowel disease
(Xavier and Podolsky 2007, Khor, Gardet et al. 2011, Adolph, Tomczak et al. 2013, Liu,
Gurram et al. 2016).
482
We compared global RNA expression profiles, obtained by RNA-seq, of Duodenum
Intestine-Chip, duodenal organoids and human native duodenal tissue in order to
determine which of the two models better reflects their natural counterpart. Surprisingly,
although chips and organoids were established with the cells of the same origin
(donors, source of the tissue) their transcriptomic profiles were shown to significantly
differ from each other – the total number of 472 up- and downregulated genes were
found in this comparison. Moreover, several different analyses demonstrated that the
transcriptome of Duodenum Intestine-Chip more closely resembles global gene
expression in human adult duodenum than do the organoids, strongly suggesting that
Duodenum Intestine-Chip constitute a more accurate representation of human *in vivo*
tissue. Importantly, the subset of 305 genes that were found to be common for
Duodenum Intestine-Chip and human tissue but different from organoids, showed to be

associated with important biological functions including: digestion, transport of nutrients
and ions, extracellular matrix organization, wound healing, metabolism, detoxification as
well as tissue development. In addition, several genes involved in drug metabolism,
including but not limited to drug metabolism enzymes: CYP3A4, UGT1A4, UGT2A3,
UGT2B7, UGT2B15, UGT2B17, were observed to exhibit similar pattern of expression
in Duodenum Intestine-Chip and adult duodenal tissue while they differed from
organoids, suggesting the improved potential of this system to study biotransformation
of xenobiotics and drug-drug interactions.
503
Intestinal efflux and uptake transporters are key determinants of absorption and
subsequent bioavailability of a large number of orally administered drugs. *In vivo*-like
expression of the major intestinal drug transporters, including clinically relevant MDR1,
BCRP and PEPT1, was demonstrated in Duodenum Intestine-Chip system. Additionally,
we confirmed the apical localization, which is known to be crucial for the unique
gatekeeper function of these proteins in controlling drug access to metabolizing
enzymes and excretory pathways (Shugarts and Benet 2009, Estudante, Morais et al.
2013), within the plasma membrane of intestinal epithelial cells grown on chip.
Functional assessment of MDR1 efflux revealed similar level of activity as observed in
Caco-2 Intestine-Chip and its successful inhibition by vinblastine. Notably, in
comparison to Caco-2 Intestine-Chip the organoids-based chip system showed
improved relative mRNA expression levels of organic anion and cation transporters:
OATP2B1 and OCT1. These transporters are responsible for the uptake of numerous
xenobiotics, including statins, antivirals, antibiotics, and anticancer drugs (Roth, Obaidat
et al. 2012). While significantly higher expression of these proteins, in comparison to
human duodenal tissue, was observed in Caco-2 model, their levels showed to be
closer to *in vivo* in Duodenum Intestine-Chip. Our findings suggest that organoids-
derived Intestine-Chip system could be applied to assess the specific contribution of
efflux transporters to drug disposition, to evaluate the active transport of xenobiotics
across intestinal barrier, as well as, for the modelling of increased absorption by
targeting of specific uptake transporters such as PEPT1 or OATP2B1.
525
Evaluation of the expression and drug-mediated induction of the CYP3A4, a major
enzyme involved in human metabolism of xenobiotics revealed a key advantage of
Duodenum Intestine-Chip over Caco-2 models. Significantly higher and much closer to
human duodenal tissue expression of CYP450 enzyme was observed in Duodenum
Intestine-Chip in comparison to Caco-2 cells-based system. Consistent with the results
of others (Negoro, Takayama et al. 2016) CYP3A4 was undetectable at the protein level
in Caco-2 cells and remained unchanged upon stimulation with RIF (a PXR agonist).
While successful and reproducible modulation of CYP3A4 expression using agonists
specific for PXR and VDR, rifampicin and vitamin D3, respectively, was achieved in
Duodenum Intestine-Chips engineered individually from the organoids of three
independent donors. CYP3A4 induction levels observed were similar to those shown for
intestinal slices (van de Kerkhof, de Graaf et al. 2008), suggesting suitability of this
model for the studies of drug metabolism, that cannot be supported by the previous
Caco2 models.

540
In conclusion, Duodenum Intestine-Chip provides a faithful representation of human
duodenum and represents a potential tool for preclinical drug assessment in human-
relevant system. Moreover, as it's composed of the cells isolated from individual patient
it could in the future be personalized in order to assess interindividual differences in
drug disposition and responses, studies of the effect of genetic polymorphisms on
pharmacokinetics and pharmacodynamics as well as decoupling of the effect of various
factors such as age, sex, disease state and diet on metabolism, clearance and
bioavailability of xenobiotics. This system could also be applied help us to better
understand the basic biology of human intestinal tissue in health and disease state and
potentially enable novel therapeutic development as we further our understanding of
mechanisms driving key disease phenotypes.

552
553 **Materials and Methods:**

554
Human tissue collection, generation, and culture of organoids
Human duodenal organoids cultures were established from biopsies obtained during
endoscopic or surgical procedures utilizing the methods developed by the laboratory of
Dr. Hans Clevers(Sato, Stange et al. 2011). De-identified biopsy tissue was obtained
from healthy adult subjects who provided informed consent at Johns Hopkins University
and all methods were carried out in accordance with approved guidelines and
regulations. All experimental protocols were approved by the Johns Hopkins University
Institutional Review Board (IRB #NA 00038329). Briefly, organoids generated from
isolated intestinal crypts were grown embedded in Matrigel^{TM} (Corning, USA) in the
presence of Expansion Medium consisting of Advanced DMEM F12 supplemented with
50% v/v Wnt3A conditioned medium (produced by L-Wnt3A cell line, ATCC CRL-2647),
20% v/v R-spondin-1 conditioned medium (produced by HEK293T cell line stably
expressing mouse R-spondin1; kindly provided by Dr Calvin Kuo, Stanford University,
Stanford, CA), 10% v/v Noggin conditioned medium (produced by HEK293T cell line
stably expressing mouse Noggin), 10mM HEPES, 0.2 mM GlutaMAX, 1x B27
supplement, 1x N2 supplement, 1 mM n-acetyl cysteine, 50 ng/ml human epithelial
growth factor, 10 nM human [Leu15] -gastrin, 500 nM A83-01, 10 µM SB202190, 100
µg/ml primocin. EM was replaced every other day and supplemented with 10 µM
CHIR99201 and 10 µM Y-27632 during the first 2 days after passaging. Organoids were
passaged every 7 days and used for chip seeding between passage numbers 5 and 30.

575
Duodenum Intestine-Chip
The design and fabrication of Organ-Chips used to develop the Duodenum Intestine-
Chip was based on previously described protocols(Huh, Kim et al. 2013). The chip is
made of a transparent, flexible polydimethylsiloxane (PDMS), an elastomeric polymer.
The chip contains two parallel microchannels (a 1 × 1 mm epithelial channel and a 1 ×
0.2 mm vascular channel) that are separated by a thin (50 µm), porous membrane (7
µm diameter pores with 40 µm spacing) coated with ECM (200 µg/ml collagen IV and
100 µg / ml Matrigel^{TM} at the epithelial side and 200 µg / ml collagen IV and 30 µg / ml
fibronectin at the vascular side). Chips were seeded with intestinal epithelial cells

obtained from enzymatic dissociation of organoids, as described previously(Kasendra,
Tovaglieri et al. 2018), and incubated overnight before being washed with fresh media.
Next day, the chips were connected to the culture module instrument (inside the
incubator), that can hold up to 12 chips and allows for control of flow and stretching
within the chips using pressure driven flow(Vatine, Barrile et al. 2019). The chips are
maintained under constant perfusion of fresh expansion medium at 30 μl / hr through
top and bottom channels of all chips until day 6. Human Intestinal Microvascular
Endothelial Cells (HIMECs; Cell Biologics) were than plated on the vascular side of the
ECM-coated porous membrane in EGM2-MV medium, which contains human epidermal
growth factor, hydrocortisone, vascular endothelial growth factor, human fibroblastic
growth factor-B, R3-Insulin-like Growth Factor-1, Ascorbic Acid and 5% fetal bovine
serum (Lonza Cat. no. CC-3202). At this time the medium supplying the epithelial
channel was switched to differentiation medium. Differentiation medium consisted of the
same components as those of Expansion Medium with 50% less Noggin and R-spondin
and devoid of Wnt3A and SB202190. Media supplying both chip channels were under
continuous flow. Cyclic, peristalsis-like deformations of tissue attached to the membrane
(10% strain; 0.2 Hz) were initiated after the formation of confluent monolayer at ~4 days
in culture.
603
Permeability assays
In order to evaluate the establishment and integrity of the intestinal barrier, 3kDa
Cascade Blue dextran (Sigma, D7132) was added to the epithelial compartment of the
Duodenum Intestine-Chip at 0.1 mg / ml on the day of their connection to flow. Effluents
of the endothelial compartment were sampled every 48 hours to determine the
concentration of dye that had diffused through the membrane. The apparent
paracellular permeability (Papp) was calculated based on a standard curve and using
the following formula:
612

$$Papp \left(\frac{cm}{s}\right) = \frac{C_{output} \left(\frac{mg}{ml}\right) x \; Flow \; rate \; \left(\frac{ml}{s}\right)}{C_{input} \left(\frac{mg}{ml}\right) x \; A \; (cm^2)}$$

613
Where C_{output} is the concentration of dextran in the effluents of the endothelial
compartment, A is the seeded area, and C_{input} is the input concentration of dextran
spiked into the epithelial compartment. The establishment of intestinal barrier function in
Duodenum Intestine-Chips was evaluated in three independent experiments, each
performed using chips established from a different donor of biopsy-derived organoids.
At least three different chips were used per condition.
620
Morphological analysis
Immunofluorescent staining of cells in the Organ-Chip was performed with minor
modifications to previously reported protocols(Kasendra, Tovaglieri et al. 2018) . Cells
were fixed with 4% formaldehyde or cold methanol, and, when required, were
permeabilized, using 0.1% Triton X-100. 5% (v/v). Donkey serum solution in PBS was

used for blocking. Incubation with primary antibodies directed against ZO-1, VE-
cadherin, E-cadherin, villin, mucin 2, lysozyme, chromogranin A, MDR1, BCRP, PEPT1
(see Supplementary Table 2) was performed overnight at 4°C. Chips treated with
corresponding Alexa Fluor secondary antibodies (Abcam) were incubated in the dark for
2 hours at room temperature. Cells were then counterstained with nuclear dye DAPI.
Images were acquired with an inverted laser-scanning confocal microscope (Zeiss LSM
880 with Airyscan).

Chips processed for SEM were fixed in 2.5% glutaraldehyde, treated with 1% osmium
tetroxide in 0.1 M sodium cacodylate buffer, dehydrated in a series of graded
concentrations of ethanol solutions and critical point dried, as described
previously(Kasendra, Tovaglieri et al. 2018). Prior to imaging, samples were coated with
a thin (10 nm) layer of Pt/Pd using a sputter coater.

Measurement of the density of microvilli
Images of microvilli on the surface of cells were captured with a scanning electron
microscope (JSM-5600LV; JEOL). The morphological analysis and quantification of
microvilli was performed using ImageJ. Number of intestinal microvilli per μm^2 were
calculated after applying two image processing techniques, namely binarization and
particle analysis, and Otsu's thresholding method, as described previously (Julio,
Merindano et al. 2008).

MDR-1 efflux pump activity
The transporter activity of MDR-1 was assessed using the MDR1 Efflux Assay Kit
(ECM910, Millipore), as per manufacturer's instructions. Briefly, Duodenum Intestine-
Chips and Caco-2 Intestine-Chips(Kim, Huh et al. 2012, Kim and Ingber 2013) were
perfused apically with rhodamine 123, a fluorescent transport substrate of MDR1.
Intracellular accumulation of dye, was detected by fluorescent imaging (Olympus IX83)
and measured in the presence and absence of MDR-1 specific inhibitor vinblastine (22
μM). Three independent experiments were performed for Caco-2 Intestine-Chips and
Duodenum Intestine-Chips, each using chips established from a different donor of
biopsy-derived organoids. At least three different chips were used per condition. Images
of each were taken at three different fields of view, digitally processed and quantified
using Fuji software.

CYP3A4 induction
Duodenum Intestine-Chips and Caco-2 Intestine-Chips were treated with 100 nM 1,25-
Dihydroxyvitamin D3 (Sigma) or 20 μM rifampicin (Sigma), which are known to induce
CYP3A4, for 48 hours. Controls were treated with DMSO (final concentration 0.1%).
Subsequently, cells in the epithelial channel were harvested either for RNA isolation and
gene expression analysis or for Western blotting in order to assess the level of CYP3A4
induction at the gene and protein level, respectively.

Western blotting

RIPA cell lysis buffer (Pierce) supplemented with protease and phosphatase inhibitors
(Sigma) was used for the extraction of total protein from the chips. The protein
concentration in each sample was determined using the bicinchoninic acid method.
Equal amounts (15 µg) of protein lysates were heat denatured and separated on a 4–
10% Mini-Protean Precast Gel (Bio-Rad), followed by transfer on a nitrocellulose
membrane (Bio-Rad). After blocking with 5% nonfat milk, membranes were probed with
primary antibodies for CYP3A4 (mouse monoclonal, Santa Cruz Biotechnology) and
GAPDH (rabbit polyclonal, Abcam) and incubated overnight at 4°C, followed by
incubation for 1 hour with IRDye-conjugated secondary antibodies against rabbit and
mouse immunoglobulin G (LI-COR), at room temperature. Finally, blots were scanned
using an Odyssey Infrared Imaging System (LI-COR) and the protein bands were
visualized and quantified using Image Studio software (LI-COR). Gluceraldehyde-3-
phosphatase dehydrogenase (GAPDH) was used as the loading control.
683
Gene expression analysis
Total RNA was isolated from the chip using PureLink RNA Mini kit (Fisher Scientific)
and reverse transcribed to cDNA using SuperScript IV Synthesis System (Fisher
Scientific). qRT-PCR was performed using TaqMan Fast Advanced Master Mix (Applied
Biosystems) and TaqMan Gene Expression Assays (see Supplementary Table 3, Fisher
Scientific) in QuantStudio 5 PCR System (Fisher Scientific). Relative expression of gene
was calculated using 2-ΔΔCt method.
691
RNA isolation and sequencing
RNA was extracted using TRIzol (Life Technologies) according to manufacturer's
guidelines. Samples were submitted to GENEWIZ South Plainfield, NJ for next
generation sequencing. After quality control and a complementary DNA library creation,
all samples were sequenced using HiSeq 4000 with 2X150 bp paired-end reads per
sample.
698
RNA sequencing bioinformatics
Pre-processing: raw sequence data (.bcl files) generated from Illumina HiSeq was
converted into fastq files and de-multiplexed using Illumina's bcl2fastq 2.17 software.
Read quality was assessed using FastQC. Adaptor and low-quality (< 15) sequences
were removed using Trimmomatic v.0.36. The trimmed reads were mapped to the
Homo sapiens reference genome available on ENSEMBL using the STAR aligner
v.2.5.2b. The STAR aligner uses a splice aligner that detects splice junctions and
incorporates them to help the alignment of the entire read sequences. BAM files were
generated following this step. Unique gene raw counts were calculated by using feature
Counts from the Subread package v.1.5.2. Only the unique reads that fell within exon
regions were counted.
710
711 Differential Gene Expression Analysis: to combine our gene expression dataset with the
712 publicly available data for adult duodenum gene expression (provided as Fragments Per
713 Kilobase of transcript per Million mapped read (FPKM) values in (Finkbeiner, Hill et al.
714 2015)), we converted the raw counts to FPKMs. Then using the log2(FPKM

expressions of the combined datasets, we applied DE gene analysis using the R package "limma"(Ritchie, Phipson et al. 2015). For each comparison, the thresholds used to identify the DE genes were set to a) adjusted p-values < 0.05 and b) absolute log2 fold change > 2.

GO term enrichment analysis and KEGG Pathway analysis
After expression pattern clustering, the transcripts from specific groups were subjected to functional annotation, including GO (Gene Ontology) functional annotation and KEGG (Kyoto Encyclopedia of Genes and Genomes) pathway annotation. The GO terms and KEGG pathway enrichment was performed using The Database for Annotation, Visualization and Integrated Discovery (DAVID v 6.8, http://david.abcc.ncifcrf.gov).

Statistical analysis
All experiments were performed in triplicates and repeated with organoids from three different human donors. One-way or two-way ANOVA was performed to determine statistical significance, as indicated in the figure legends. The error bars represent standard error of the mean [s.e.m]; P values < 0.05 and above were considered as significant.

Data availability
RNA sequencing data have been deposited in the National Center for Biotechnology Information Gene Expression Omnibus (GEO) under accession number GSE135196.

Figures

Figure 1. Duodenum Intestine-Chip: a microengineered model of the human duodenum.
(a) Brightfield images of human duodenal organoids (top) and human microvascular endothelial cells (bottom) acquired before their seeding into epithelial and endothelial channels of the chip. **(b)** Schematic representation of Duodenum Intestine-Chip, including its top view (left) and vertical section (right) showing: the epithelial (1; blue) and vascular (2; pink) cell culture microchannels populated by intestinal epithelial cells (3) and endothelial cells (4), respectively, and separated by a flexible, porous, ECM-coated PDMS membrane (5). **(c)** Scanning electron micrograph showing complex intestinal epithelial tissue architecture achieved by duodenal epithelium grown on the chip (top) in the presence of constant flow of media (30 µl h^{-1}) and cyclic membrane deformations (10% strain, 0.2 Hz). High magnification of the apical epithelial cell surface with densely packed intestinal microvilli (bottom). **(d)** Composite tile scan fluorescence image (top) showing a fully confluent monolayer of organoids-derived intestinal epithelial cells (magenta, ZO-1 staining) lining the lumen of Duodenum Intestine-Chip and interfacing with microvascular endothelium (green, VE-cadherin staining) seeded in the adjacent vascular channel. Higher magnification views of epithelial tight junctions (bottom left) stained against ZO-1 (magenta) and endothelial adherence junctions visualized by VE-cadherin (green) staining. Cells nuclei are shown in grey. Scale bar, 1000 µm (top) 100 µm (bottom) **(e)** Apparent permeability values of Duodenum Intestine-Chips cultured in the presence of flow and stretch (30 ul/h, 10% 0,2 Hz) for up to 10 days. Papp values were calculated from the diffusion of 3 kDa Dextran from the luminal to the vascular channel. Data represent three independent experiments performed with three different chips/donor, total of 3 donors; Error bars indicate s.e.m.

762
Figure 2. Duodenum Intestine-Chip emulates multi-lineage differentiation of native
human intestine.
(a) Comparison of the relative gene expression levels of markers specific for different intestinal
cell types, including mucin 2 (MUC2) for goblet cells, alkaline phosphatase (ALPI) for absorptive
enterocytes, chromogranin A (CHGA) for enteroendocrine cells, lysozyme (LYZ) for Paneth
cells, leucine-rich repeat-containing G-protein coupled receptor 5 (LGR5) for stem cells and
proliferation marker (KI67) across Duodenum Intestine-Chips established from 3 individual
organoids donors at different times in the chip under continuous flow (day 2, 4, 6, 8, 10) and
RNA isolated directly from the duodenal tissue of other 3 different donors (Duodenum). In each
graph, values represent average gene expression ± s.e.m (error bars) from three independent
experiments, each using different donors of biopsy-derived organoids and at least three different
chips per time point. Values are shown relative to duodenal tissue expressed as 1. EPCAM
expression was used as normalizing control. One-way ANOVA, ****$p<0.0001$, ***$p<0.001$,
$p<0.01$, *$p<0.05$, ns $p>0.05$ **(b) Representative confocal fluorescent micrographs
demonstrating the presence of all major intestinal cell types (green) in Duodenum Intestine-Chip
at day 8 of fluidic culture, including goblet cells stained with anti-Muc2; enteroendocrine cells
visualized with anti-chromogranin A, absorptive enterocytes stained with anti-villin and Paneth
cells labeled with anti-lysozyme. Cell-cell borders are stained with anti-E-cadherin and are
shown in magenta. Scale bar, 10 μm **(c)** Quantification of the different epithelial intestinal cell
types present in Duodenum Intestine-Chip at day 8 and identified by immunostaining, as
described in (b). Percentage of different cell types based on 10 different fields of view (10 FOV)
counted in three individual chips per staining established from three different donors. DAPI
staining was used to evaluate the total cell number. Duodenum values, represent cell ratios
observed in the histological sections and are based on the literature (Karam 1999).
787
Figure 3. Duodenum Intestine-Chip exhibits higher transcriptomic similarity to adult
duodenal tissue than organoid culture (a) Differential gene expression analysis was carried
out to identify genes that are upregulated or downregulated in Duodenum Intestine-Chip
compared to organoids (blue circle) (Figure 3—source data 1) and adult duodenum compared
to organoids (yellow circle) (Figure 3—source data 2). The gene lists were then compared to
determine how many genes overlap between those two comparisons (green) (Figure 3—source
data 3), and the results are shown as a Venn diagram. 305 genes were identified as common
and responsible for the closer resemblance of Duodenum Intestine-Chip to human adult
duodenum than organoids from which chips were derived. Sample sizes were as follows:
Duodenum Intestine-Chip, n = 3 (independent donors); Organoids, n = 3 (independent donors);
Adult duodenum, n = 2 (independent biological specimens). Intestinal crypts derived from the
same three independent donors were used for the establishment of Duodenum Intestine-Chip
and organoid cultures **(b)** The list of overlapping genes was subjected to GO analysis to identify
enriched biological processes (GO terms) (Figure 3—source data 4). The results are shown as
REVIGO scatterplots in which similar GO terms are grouped in arbitrary two-dimensional space
based on semantic similarity. Each circle corresponds to a specific GO term and circle sizes are
proportional to the number of genes included in each of the enriched GO terms. Finally, the
color of a circle indicates the significance of the specific GO term enrichment. GO terms
enriched in the overlapping gene set demonstrate that Duodenum Intestine-Chip is more similar
to human duodenum with respect to important biological functions of the intestine, including
digestion, transport and metabolism. **(c)** The results of the KEGG pathway analysis using the
305 differentially expressed genes showed seven significantly enriched (FDR corrected p-value
<0.05) pathways related to absorption, metabolism, digestion and chemical carcinogenesis. The

size of the bars indicates the fold-enrichment of the corresponding pathways. **(d)** Curated
heatmaps were generated to examine particular genes that belong to the enriched KEGG
pathways and show the expression intensity (log$_2$(FPKM) of these genes across different
samples. The provided results further demonstrate that Duodenum Intestine-Chip is more
similar to adult duodenum than are the organoids. Sample sizes were as follows: Duodenum
Intestine-Chip, n = 3 (independent donors); Organoids, n = 3 (independent donors); Adult
duodenum, n = 2 (independent biological specimens). Intestinal crypts derived from the same
three independent donors were used for the establishment of Duodenum Intestine-Chip and
organoid cultures.
820
Figure 4. Duodenum Intestine-Chip shows the presence of major intestinal drug
transporters and correct localization and function of efflux pump MDR1 (P-gp). (a)
Comparison of the relative average gene expression levels of drug efflux (MDR1, BCRP, MRP2,
MRP3) and uptake (PEPT1, OATP2B1, OCT1, SLC40A1) transporters in Caco-2 based
Intestine-Chips (Intestine-Chip), three donor-specific Duodenum Intestine-Chips and RNA
isolated directly from the duodenal tissue of three independent individuals (Duodenum). The
results show that Duodenum Intestine-Chips express drug transport proteins at the levels close
to human duodenal tissue. Note, that the expression of OATP2B1 and OCT1 in Caco-2 was
significantly higher than in human duodenum while the difference between Duodenum Intestine-
Chip and adult duodenum was not significant. Each value represents average gene expression
± s.e.m (error bars) from three independent experiments, each involving Duodenum Intestine-
Chips established from a tissue of three independent donors (three chips/donor), RNA tissue
from three independent biological specimens, and Caco-2 Intestine-Chips (three chips). Values
are shown relative to the duodenal tissue expressed as 1, Two-way ANOVA, ****p<0.0001,
***p<0.001, **p<0.01. EPCAM expression was used as normalizing control. **(b)** Representative
confocal immunofluorescence micrographs of vertical cross-sections (left) and corresponding to
the plots (right) showing co-distribution of fluorescent signal of efflux transporter MDR1 (green),
and villin (magenta) as visualized by merge channel (white) and fluorescent spectra overlap
across vertical cross-sections of the differentiated epithelium in Duodenum Intestine-Chip. Efflux
transporter MDR1 colocalize precisely with luminal cell surface marker (villin) in the monolayer
of cells closely attached to membrane (monolayer) as well as in the cells lining subsequently
formed villi-like structures (Villi-like Structure). Fluorescent signal representing cell nuclei is
visualized in cyan. Scale bar, 10 µm. See also Figure 4—figure supplement 1 showing luminal
localization of additional efflux (BCRP) and uptake (PEPT1) transporters in Duodenum
Intestine-Chip. **(c)** Activity of efflux pump proteins. The intracellular accumulation of the
fluorescent substrate of MDR1 - Rhodamine 123 was significantly increased in response to the
MDR1 inhibitor vinblastine (black bars) in comparison to vehicle (DMSO) control (grey bars) in
both systems – Caco2 and organoid-derived Duodenum Intestine-Chips. Data are presented as
mean ± s.e.m (error bars) of at least three independent experiments involving chips generated
from organoids of at least three individual donors and Caco-2 cell line. Two-way ANOVA,
**p<0.01, *p<0.05.
852
Figure 5. CYP3A4 expression levels and induction in Duodenum Intestine-Chip and Caco-
2 cells-based Intestine-Chip. (a) Average gene expression levels of CYP3A4 ± s.e.m (error
bars) in Caco-2 Intestine-Chip, Duodenum Intestine-Chip and human duodenum (three
independent biological specimens). All values are shown relative to the adult duodenal tissue
expressed as 1, One-way ANOVA, ****p<0.0001, **p<0.01. EPCAM expression was used as
normalizing control. **(b)** Protein analysis of CYP3A4 in Caco-2 and Duodenum Intestine-Chips
using western blot. **(c)** The CYP3A4 induction in Caco-2 Intestine-Chips and Duodenum

Intestine-Chips treated with solvent (DMSO), 20 µM rifampicin (RIF) or 100 nM 1,25-
dihidroxyvitamin D3 (VD3) for 48 h. The gene expression levels (top) of CYP3A4 were
examined by real-time PCR analysis. On the y axis, the gene expression levels in the DMSO-
treated chips were taken as 1.0. All data are represented as means ± s.e.m. Two-way ANOVA,
****p<0.0001 (compared with DMSO-treated cells). The corresponding CYP3A4 protein
expression levels (bottom) were measured by western blotting analysis. **(d)** Gene expression
analysis of the receptors PXR and VDR in organoids and Caco-2 cell-derived chips examined
by real-time RT-PCR analysis and compared to their expression in adult duodenal tissue
(Duodenum). On the y axis, the gene expression levels in adult tissue were taken as 1.0. All
data are represented as means ± s.e.m. Two-way ANOVA, ***p<0.001, **p<0.01, *p<0.05.

**Supplementary Figure 1. Flow-induced increase in primary intestinal epithelial cells
height and microvilli formation. (a)** Representative confocal images of x-y (top) and x-z
(bottom) optical sections of duodenal organoid-derived epithelial cells cultured under static
(Static) or fluid flow (30 µl h^{-1}; Flow) or flow and stretch (30 µl h^{-1}; 10% strain, 0.2 Hz;
Flow+Stretch) conditions and stained for apical marker villin (green) and basolateral protein E-
cadherin (magenta). Nuclei we counterstained with DAPI (grey). Scale bar, 50 µm. **(b)** The
quantitative analysis of the average cell height measured from Z-stack images as the distance
between apical marker villin (green, a) and PDMS membrane (dotted line, a). The data
represent the mean ± s.e.m; One-way ANOVA, ****p<0.0001. **(c)** Scanning electron microscopy
surface images of duodenal enteroid-derived epithelium cultured under static or flow +/- stretch
conditions. Cells seeded in the top channel of Organ-Chip were maintained with (Flow) or
without (Static) medium perfusion (30µl h^{-1}) in both channels and 10% of mechanical stretch
(0.2 Hz) (Flow+Stretch). Images were captured at the center area of the chamber. Scale bar, 5
µm. **(d)** Quantification of microvilli. Density of microvilli per µm^2 was measured from the SEM
images (100 µm2, 20 FOV) as described in the Methods. The data represent the mean ± s.e.m;
One-way ANOVA, ****p<0.0001.

**Supplementary Figure 2. Differentially expressed genes and enriched Gene Ontology
categories in Organoids or Duodenum Intestine-Chip with respect to Adult Duodenum.**
(a) Volcano plot (left) and functional enrichment analysis (right) of differentially expressed genes
between Organoids and Adult Duodenum (Supplementary Figure 2—source data 1). The red
dots represent genes that are significantly (adj. p-value<0.05) up- or downregulated. The black
dots correspond to the non-differentially expressed genes. The vertical lines correspond to 2.0-
fold up and down and the horizontal line has been drawn at the level of the selected cutoff
adjusted p-value (adj. p-value<0.05). Sample sizes were as follows: Duodenum Intestine-Chip,
n = 3; Organoids, n = 3. All samples were biologically independent (derived from a different
donor). Samples from the same 3 donors were used for the establishment of Duodenum
Intestine-Chip and organoid cultures. Functional enrichment analysis demonstrated over (+) and
under (-) represented biological processes in the GO categories concerning digestion,
extracellular matrix organization, angiogenesis, cell adhesion, tissue development, cell
response to drugs and toxic substances, while nucleic acid metabolic process and RNA
processing were under represented. GO, Gene Ontology. **(b)** Differential gene expression and
functional enrichment analysis between Duodenum Intestine-Chip and adult human tissue
(Supplementary Figure 2—source data 2) demonstrating up- and down-regulated genes
(volcano plot, left) and annotated to them biological processes (table, right) involving but not
limited to protein synthesis and targeting as well as cell cycle and cell proliferation. Red dots:
significant genes (adj. p-value<0.05). Black dots: non-differentially expressed genes. Sample
sizes were as follows: Duodenum Intestine-Chip, n = 3; Adult Duodenum Human tissue, n = 2.

909 Sample sizes were as follows: Duodenum Intestine-Chip, n = 3; Adult Duodenum, n = 2. All
910 samples were biologically independent (derived from a different donor).
911
912 **Supplementary Table 1.** RNAseq Datasets Downloaded from Public Databases
913
914 **Supplementary Table 2.** List of primary antibodies used for immunostaining.
915
916 **Supplementary Table 3.** List of human TaqMan Gene Expression Assays used for qRT-PCR.
917
918 **Figure 3—source data 1**. Differentially Expressed Genes in Duodenum Intestine-Chip vs.
919 Duodenal Organoids, Related to Figure 3.
920
921 **Figure 3—source data 2**. Differentially Expressed Genes in Adult Duodenum vs. Duodenal
922 Organoids, Related to Figure 3.
923
924 **Figure 3—source data 3**. Differentially Expressed Genes Common in Duodenum Intestine-
925 Chip and Adult Duodenum versus Duodenal Organoids, Related to Figure 3.
926
927 **Figure 3—source data 4**. Enriched GO Terms from a List of Differentially Expressed Genes
928 Common in Duodenum Intestine-Chip and Adult Intestine versus Duodenal Organoids, Related
929 to Figure 3.
930
Figure 4—figure supplement 1. Luminal localization of efflux (BCRP) and uptake (PEPT1)
transporters in Duodenum Intestine-Chip.
Representative cross-sectional confocal images of Duodenum Intestine-Chip (left) showing
apical localization of efflux BCRP (**a**; green) and uptake PEPT1 drug transporters (**b**; green) that
co-localize with luminal cell surface marker villin (magenta) at the time of confluent monolayer
formation as well as within successively formed villi-like structures. Plots representing
distribution of the fluorescence signal across epithelial cell z-axis revealed close overlap of
green (transporters; BRCP and PEPT1) and magenta (apical cell marker; villin) signals
confirming co-distribution of these proteins on the luminal cell surface. Cell nuclei are visualized
in cyan. Scale bar, 10 µm.
941
942 **Supplementary Figure 2—source data 1.** Differentially Expressed Genes in Duodenal
943 Organoids vs. Adult Duodenum, Related to Supplementary Figure 2.
944
945 **Supplementary Figure 2—source data 2.** Differentially Expressed Genes in Duodenum
946 Intestine-Chip vs. Adult Duodenum, Related to Supplementary Figure 2.
947
References

Adolph, T. E., M. F. Tomczak, L. Niederreiter, H. J. Ko, J. Bock, E. Martinez-Naves, J.
N. Glickman, M. Tschurtschenthaler, J. Hartwig, S. Hosomi, M. B. Flak, J. L. Cusick, K.
Kohno, T. Iwawaki, S. Billmann-Born, T. Raine, R. Bharti, R. Lucius, M. N. Kweon, S. J.
Marciniak, A. Choi, S. J. Hagen, S. Schreiber, P. Rosenstiel, A. Kaser and R. S.
Blumberg (2013). "Paneth cells as a site of origin for intestinal inflammation." Nature
503(7475): 272-276.

Ashburner, M., C. A. Ball, J. A. Blake, D. Botstein, H. Butler, J. M. Cherry, A. P. Davis,
K. Dolinski, S. S. Dwight, J. T. Eppig, M. A. Harris, D. P. Hill, L. Issel-Tarver, A.
Kasarskis, S. Lewis, J. C. Matese, J. E. Richardson, M. Ringwald, G. M. Rubin and G.
Sherlock (2000). "Gene ontology: tool for the unification of biology. The Gene Ontology
Consortium." Nat Genet **25**(1): 25-29.
Dekkers, J. F., C. L. Wiegerinck, H. R. de Jonge, I. Bronsveld, H. M. Janssens, K. M. de
Winter-de Groot, A. M. Brandsma, N. W. de Jong, M. J. Bijvelds, B. J. Scholte, E. E.
Nieuwenhuis, S. van den Brink, H. Clevers, C. K. van der Ent, S. Middendorp and J. M.
Beekman (2013). "A functional CFTR assay using primary cystic fibrosis intestinal
organoids." Nat Med **19**(7): 939-945.
Dietrich, C. G., A. Geier and R. P. Oude Elferink (2003). "ABC of oral bioavailability:
transporters as gatekeepers in the gut." Gut **52**(12): 1788-1795.
Eden, E., D. Lipson, S. Yogev and Z. Yakhini (2007). "Discovering motifs in ranked lists
of DNA sequences." PLoS Comput Biol **3**(3): e39.
Eglen, R. M. and D. H. Randle (2015). "Drug Discovery Goes Three-Dimensional:
Goodbye to Flat High-Throughput Screening?" Assay Drug Dev Technol **13**(5): 262-
265.
Estudante, M., J. G. Morais, G. Soveral and L. Z. Benet (2013). "Intestinal drug
transporters: an overview." Adv Drug Deliv Rev **65**(10): 1340-1356.
Fatehullah, A., S. H. Tan and N. Barker (2016). "Organoids as an in vitro model of
human development and disease." Nat Cell Biol **18**(3): 246-254.
Finkbeiner, S. R., D. R. Hill, C. H. Altheim, P. H. Dedhia, M. J. Taylor, Y. H. Tsai, A. M.
Chin, M. M. Mahe, C. L. Watson, J. J. Freeman, R. Nattiv, M. Thomson, O. D. Klein, N.
F. Shroyer, M. A. Helmrath, D. H. Teitelbaum, P. J. Dempsey and J. R. Spence (2015).
"Transcriptome-wide Analysis Reveals Hallmarks of Human Intestine Development and
Maturation In Vitro and In Vivo." Stem Cell Reports.
Gayer, C. P. and M. D. Basson (2009). "The effects of mechanical forces on intestinal
physiology and pathology." Cell Signal **21**(8): 1237-1244.
Groer, C., S. Bruck, Y. Lai, A. Paulick, A. Busemann, C. D. Heidecke, W. Siegmund and
S. Oswald (2013). "LC-MS/MS-based quantification of clinically relevant intestinal
uptake and efflux transporter proteins." J Pharm Biomed Anal **85**: 253-261.
Guengerich, F. P. (2008). "Cytochrome p450 and chemical toxicology." Chem Res
Toxicol **21**(1): 70-83.
Harwood, M. D., B. Achour, S. Neuhoff, M. R. Russell, G. Carlson, G. Warhurst and A.
Rostami-Hodjegan (2016). "In Vitro-In Vivo Extrapolation Scaling Factors for Intestinal
P-glycoprotein and Breast Cancer Resistance Protein: Part II. The Impact of Cross-
Laboratory Variations of Intestinal Transporter Relative Expression Factors on Predicted
Drug Disposition." Drug Metab Dispos **44**(3): 476-480.
Huh, D., H. J. Kim, J. P. Fraser, D. E. Shea, M. Khan, A. Bahinski, G. A. Hamilton and
D. E. Ingber (2013). "Microfabrication of human organs-on-chips." Nat Protoc **8**(11):
2135-2157.
Julio, G., M. D. Merindano, M. Canals and M. Rallo (2008). "Image processing
techniques to quantify microprojections on outer corneal epithelial cells." J Anat **212**(6):
879-886.

Kanehisa, M. and S. Goto (2000). "KEGG: kyoto encyclopedia of genes and genomes."
Nucleic Acids Res **28**(1): 27-30.
Karam, S. M. (1999). "Lineage commitment and maturation of epithelial cells in the gut."
Front Biosci **4**: D286-298.
Kasendra, M., A. Tovaglieri, A. Sontheimer-Phelps, S. Jalili-Firoozinezhad, A. Bein and
A. Chalkiadaki (2018). "Development of a primary human small intestine-on-a-chip
using biopsy-derived organoids." Sci Rep **8**.
Kasendra, M., A. Tovaglieri, A. Sontheimer-Phelps, S. Jalili-Firoozinezhad, A. Bein, A.
Chalkiadaki, W. Scholl, C. Zhang, H. Rickner, C. A. Richmond, H. Li, D. T. Breault and
D. E. Ingber (2018). "Development of a primary human Small Intestine-on-a-Chip using
biopsy-derived organoids." Sci Rep **8**(1): 2871.
Khor, B., A. Gardet and R. J. Xavier (2011). "Genetics and pathogenesis of
inflammatory bowel disease." Nature **474**(7351): 307-317.
Kim, H. J., D. Huh, G. Hamilton and D. E. Ingber (2012). "Human gut-on-a-chip
inhabited by microbial flora that experiences intestinal peristalsis-like motions and flow."
Lab Chip **12**(12): 2165-2174.
Kim, H. J. and D. E. Ingber (2013). "Gut-on-a-Chip microenvironment induces human
intestinal cells to undergo villus differentiation." Integr Biol (Camb) **5**(9): 1130-1140.
Kliewer, S. A., J. T. Moore, L. Wade, J. L. Staudinger, M. A. Watson, S. A. Jones, D. D.
McKee, B. B. Oliver, T. M. Willson, R. H. Zetterstrom, T. Perlmann and J. M. Lehmann
(1998). "An orphan nuclear receptor activated by pregnanes defines a novel steroid
signaling pathway." Cell **92**(1): 73-82.
Komura, H. and M. Iwaki (2011). "In vitro and in vivo small intestinal metabolism of
CYP3A and UGT substrates in preclinical animals species and humans: species
differences." Drug Metab Rev **43**(4): 476-498.
Kumar, K. K., S. Karnati, M. B. Reddy and R. Chandramouli (2010). "Caco-2 cell lines in
drug discovery- an updated perspective." J Basic Clin Pharm **1**(2): 63-69.
LeCluyse, E. L. (2001). "Pregnane X receptor: molecular basis for species differences in
CYP3A induction by xenobiotics." Chem Biol Interact **134**(3): 283-289.
Liu, F., J. Huang, B. Ning, Z. Liu, S. Chen and W. Zhao (2016). "Drug Discovery via
Human-Derived Stem Cell Organoids." Front Pharmacol **7**: 334.
Liu, T. C., B. Gurram, M. T. Baldridge, R. Head, V. Lam, C. Luo, Y. Cao, P. Simpson, M.
Hayward, M. L. Holtz, P. Bousounis, J. Noe, D. Lerner, J. Cabrera, V. Biank, M.
Stephens, C. Huttenhower, D. P. McGovern, R. J. Xavier, T. S. Stappenbeck and N. H.
Salzman (2016). "Paneth cell defects in Crohn's disease patients promote dysbiosis."
JCI Insight **1**(8): e86907.
Mackowiak, B., J. Hodge, S. Stern and H. Wang (2018). "The Roles of Xenobiotic
Receptors: Beyond Chemical Disposition." Drug Metab Dispos **46**(9): 1361-1371.
Maglich, J. M., D. J. Parks, L. B. Moore, J. L. Collins, B. Goodwin, A. N. Billin, C. A.
Stoltz, S. A. Kliewer, M. H. Lambert, T. M. Willson and J. T. Moore (2003). "Identification
of a novel human constitutive androstane receptor (CAR) agonist and its use in the
identification of CAR target genes." J Biol Chem **278**(19): 17277-17283.
Martignoni, M., G. M. Groothuis and R. de Kanter (2006). "Species differences between
mouse, rat, dog, monkey and human CYP-mediated drug metabolism, inhibition and
induction." Expert Opin Drug Metab Toxicol **2**(6): 875-894.

Maubon, N., M. Le Vee, L. Fossati, M. Audry, E. Le Ferrec, S. Bolze and O. Fardel
(2007). "Analysis of drug transporter expression in human intestinal Caco-2 cells by
real-time PCR." Fundam Clin Pharmacol **21**(6): 659-663.
Mi, H., A. Muruganujan, J. T. Casagrande and P. D. Thomas (2013). "Large-scale gene
function analysis with the PANTHER classification system." Nat Protoc **8**(8): 1551-1566.
Negoro, R., K. Takayama, Y. Nagamoto, F. Sakurai, M. Tachibana and H. Mizuguchi
(2016). "Modeling of drug-mediated CYP3A4 induction by using human iPS cell-derived
enterocyte-like cells." Biochem Biophys Res Commun **472**(4): 631-636.
Nelson, D. R., D. C. Zeldin, S. M. Hoffman, L. J. Maltais, H. M. Wain and D. W. Nebert
(2004). "Comparison of cytochrome P450 (CYP) genes from the mouse and human
genomes, including nomenclature recommendations for genes, pseudogenes and
alternative-splice variants." Pharmacogenetics **14**(1): 1-18.
Olander, M., J. R. Wisniewski, P. Matsson, P. Lundquist and P. Artursson (2016). "The
Proteome of Filter-Grown Caco-2 Cells With a Focus on Proteins Involved in Drug
Disposition." J Pharm Sci **105**(2): 817-827.
Ozawa, T., K. Takayama, R. Okamoto, R. Negoro, F. Sakurai, M. Tachibana, K.
Kawabata and H. Mizuguchi (2015). "Generation of enterocyte-like cells from human
induced pluripotent stem cells for drug absorption and metabolism studies in human
small intestine." Sci Rep **5**: 16479.
Paine, M. F., H. L. Hart, S. S. Ludington, R. L. Haining, A. E. Rettie and D. C. Zeldin
(2006). "The human intestinal cytochrome P450 "pie"." Drug Metab Dispos **34**(5): 880-
886.
Pang, K. S. (2003). "Modeling of intestinal drug absorption: roles of transporters and
metabolic enzymes (for the Gillette Review Series)." Drug Metab Dispos **31**(12): 1507-
1519.
Peters, S. A., C. R. Jones, A. L. Ungell and O. J. Hatley (2016). "Predicting Drug
Extraction in the Human Gut Wall: Assessing Contributions from Drug Metabolizing
Enzymes and Transporter Proteins using Preclinical Models." Clin Pharmacokinet **55**(6):
673-696.
Press, B. and D. Di Grandi (2008). "Permeability for intestinal absorption: Caco-2 assay
and related issues." Curr Drug Metab **9**(9): 893-900.
Ritchie, M. E., B. Phipson, D. Wu, Y. Hu, C. W. Law, W. Shi and G. K. Smyth (2015).
"limma powers differential expression analyses for RNA-sequencing and microarray
studies." Nucleic Acids Res **43**(7): e47.
Roth, M., A. Obaidat and B. Hagenbuch (2012). "OATPs, OATs and OCTs: the organic
anion and cation transporters of the SLCO and SLC22A gene superfamilies." Br J
Pharmacol **165**(5): 1260-1287.
Sato, T., D. E. Stange, M. Ferrante, R. G. Vries, J. H. Van Es, S. Van den Brink, W. J.
Van Houdt, A. Pronk, J. Van Gorp, P. D. Siersema and H. Clevers (2011). "Long-term
expansion of epithelial organoids from human colon, adenoma, adenocarcinoma, and
Barrett's epithelium." Gastroenterology **141**(5): 1762-1772.
Sato, T., R. G. Vries, H. J. Snippert, M. van de Wetering, N. Barker, D. E. Stange, J. H.
van Es, A. Abo, P. Kujala, P. J. Peters and H. Clevers (2009). "Single Lgr5 stem cells
build crypt-villus structures in vitro without a mesenchymal niche." Nature **459**(7244):
262-265.

Schmiedlin-Ren, P., K. E. Thummel, J. M. Fisher, M. F. Paine, K. S. Lown and P. B.
Watkins (1997). "Expression of enzymatically active CYP3A4 by Caco-2 cells grown on
extracellular matrix-coated permeable supports in the presence of 1alpha,25-
dihydroxyvitamin D3." Mol Pharmacol **51**(5): 741-754.
Shi, S. and Y. Li (2014). "Interplay of Drug-Metabolizing Enzymes and Transporters in
Drug Absorption and Disposition." Curr Drug Metab **15**(10): 915-941.
Shugarts, S. and L. Z. Benet (2009). "The role of transporters in the pharmacokinetics of
orally administered drugs." Pharm Res **26**(9): 2039-2054.
Sun, D., H. Lennernas, L. S. Welage, J. L. Barnett, C. P. Landowski, D. Foster, D.
Fleisher, K. D. Lee and G. L. Amidon (2002). "Comparison of human duodenum and
Caco-2 gene expression profiles for 12,000 gene sequences tags and correlation with
permeability of 26 drugs." Pharm Res **19**(10): 1400-1416.
Sun, H., E. C. Chow, S. Liu, Y. Du and K. S. Pang (2008). "The Caco-2 cell monolayer:
usefulness and limitations." Expert Opin Drug Metab Toxicol **4**(4): 395-411.
Supek, F., M. Bosnjak, N. Skunca and T. Smuc (2011). "REVIGO summarizes and
visualizes long lists of gene ontology terms." PLoS One **6**(7): e21800.
The Gene Ontology, C. (2017). "Expansion of the Gene Ontology knowledgebase and
resources." Nucleic Acids Res **45**(D1): D331-D338.
Thomas, P. D., M. J. Campbell, A. Kejariwal, H. Mi, B. Karlak, R. Daverman, K. Diemer,
A. Muruganujan and A. Narechania (2003). "PANTHER: a library of protein families and
subfamilies indexed by function." Genome Res **13**(9): 2129-2141.
Thummel, K. E. (2007). "Gut instincts: CYP3A4 and intestinal drug metabolism." J Clin
Invest **117**(11): 3173-3176.
Tucker, T. G., A. M. Milne, S. Fournel-Gigleux, K. S. Fenner and M. W. Coughtrie
(2012). "Absolute immunoquantification of the expression of ABC transporters P-
glycoprotein, breast cancer resistance protein and multidrug resistance-associated
protein 2 in human liver and duodenum." Biochem Pharmacol **83**(2): 279-285.
van de Kerkhof, E. G., I. A. de Graaf, A. L. Ungell and G. M. Groothuis (2008).
"Induction of metabolism and transport in human intestine: validation of precision-cut
slices as a tool to study induction of drug metabolism in human intestine in vitro." Drug
Metab Dispos **36**(3): 604-613.
van de Wetering, M., H. E. Francies, J. M. Francis, G. Bounova, F. Iorio, A. Pronk, W.
van Houdt, J. van Gorp, A. Taylor-Weiner, L. Kester, A. McLaren-Douglas, J. Blokker,
S. Jaksani, S. Bartfeld, R. Volckman, P. van Sluis, V. S. Li, S. Seepo, C. Sekhar
Pedamallu, K. Cibulskis, S. L. Carter, A. McKenna, M. S. Lawrence, L. Lichtenstein, C.
Stewart, J. Koster, R. Versteeg, A. van Oudenaarden, J. Saez-Rodriguez, R. G. Vries,
G. Getz, L. Wessels, M. R. Stratton, U. McDermott, M. Meyerson, M. J. Garnett and H.
Clevers (2015). "Prospective derivation of a living organoid biobank of colorectal cancer
patients." Cell **161**(4): 933-945.
Vatine, G. D., R. Barrile, M. J. Workman, S. Sances, B. K. Barriga, M. Rahnama, S.
Barthakur, M. Kasendra, C. Lucchesi, J. Kerns, N. Wen, W. R. Spivia, Z. Chen, J. Van
Eyk and C. N. Svendsen (2019). "Human iPSC-Derived Blood-Brain Barrier Chips
Enable Disease Modeling and Personalized Medicine Applications." Cell Stem Cell
24(6): 995-1005 e1006.

Vlachogiannis, G., S. Hedayat, A. Vatsiou, Y. Jamin, J. Fernandez-Mateos, K. Khan, A.
Lampis, K. Eason, I. Huntingford, R. Burke, M. Rata, D. M. Koh, N. Tunariu, D. Collins,
S. Hulkki-Wilson, C. Ragulan, I. Spiteri, S. Y. Moorcraft, I. Chau, S. Rao, D. Watkins, N.
Fotiadis, M. Bali, M. Darvish-Damavandi, H. Lote, Z. Eltahir, E. C. Smyth, R. Begum, P.
A. Clarke, J. C. Hahne, M. Dowsett, J. de Bono, P. Workman, A. Sadanandam, M.
Fassan, O. J. Sansom, S. Eccles, N. Starling, C. Braconi, A. Sottoriva, S. P. Robinson,
D. Cunningham and N. Valeri (2018). "Patient-derived organoids model treatment
response of metastatic gastrointestinal cancers." Science **359**(6378): 920-926.
Wang, X., Y. Yamamoto, L. H. Wilson, T. Zhang, B. E. Howitt, M. A. Farrow, F. Kern, G.
Ning, Y. Hong, C. C. Khor, B. Chevalier, D. Bertrand, L. Wu, N. Nagarajan, F. A.
Sylvester, J. S. Hyams, T. Devers, R. Bronson, D. B. Lacy, K. Y. Ho, C. P. Crum, F.
McKeon and W. Xian (2015). "Cloning and variation of ground state intestinal stem
cells." Nature **522**(7555): 173-178.
Xavier, R. J. and D. K. Podolsky (2007). "Unravelling the pathogenesis of inflammatory
bowel disease." Nature **448**(7152): 427-434.
Zhang, L., J. Zhao, C. Liang, M. Liu, F. Xu and X. Wang (2017). "A novel biosensor
based on intestinal 3D organoids for detecting the function of BCRP." Drug Deliv **24**(1):
1453-1459.
Zhao, J., Z. Zeng, J. Sun, Y. Zhang, D. Li, X. Zhang, M. Liu and X. Wang (2017). "A
Novel Model of P-Glycoprotein Inhibitor Screening Using Human Small Intestinal
Organoids." Basic Clin Pharmacol Toxicol **120**(3): 250-255.
1154

a

Human duodenal organoids

100 μm

Human microvascular endothelial cells

100 μm

b

① ③ ② ④ ⑤

c

10 μm

5 μm

d

ZO-1 VE-cadherin

1000 μm

ZO-1

100 μm

VE-cadherin

100 μm

e

a Differentially Expressed Genes

Duodenum Intestine-Chip vs. Organoids

Adult Duodenum vs. Organoids

472   305   1132

Overlapping Genes
GO Term and KEGG Pathway Analysis

b Overlapping Genes (GO Terms)

digestion
wound healing
circulatory system development
tube development
response to chemical
response to stress
hormone metabolism
detoxification
tissue development
antibiotic metabolism
cell adhesion
ion transport
transport
extracellular matrix organization
multicellular organismal process
log10 p-value
secretion
carbohydrate transport
vitamin metabolism
cholesterol efflux
small molecule metabolism
steroid metabolism
gene expression

Semantic space y
Semantic space x

c Overlapping Genes (KEGG Pathways)

Mineral absorption
Fat digestion and absorption
Retinol metabolism
Metabolism of xenobiotics by cytochrome P450
Chemical carcinogenesis
Drug metabolism - cytochrome P450
Protein digestion and absorption

FDR
0.0100
0.0075
0.0050
0.0025

Fold Enrichment

d

Mineral absorption
TRPM6, MT1M, MT2A, MT1E, S100G, TRPV6, MT1B, SLC6A19, MT1H, MT1X, MT1G, MT1F
Adult Duo_1, Adult Duo_2, Chip_1, Chip_2, Chip_3, Organoids_1, Organoids_2, Organoids_3

Fat digestion and absorption
APOA4, ABCG8, APOB, APOA1, PNLIPRP2, CD36, ABCG5, PLA2G1B, PLA2G3
Adult Duo_1, Adult Duo_2, Chip_1, Chip_2, Chip_3, Organoids_1, Organoids_2, Organoids_3

Retinol metabolism
CYP3A4, UGT2B17, LRAT, CYP1A1, ADH4, UGT1A4, ADH6, ADH1A, UGT2A3, UGT2B15, UGT2B7
Adult Duo_1, Adult Duo_2, Chip_1, Chip_2, Chip_3, Organoids_1, Organoids_2, Organoids_3

Metabolism of xenobiotics by cytochrome P450
CYP3A4, GSTA2, UGT2B17, CYP1A1, SULT2A1, ADH4, UGT1A4, ADH6, ADH1A, UGT2A3, UGT2B15, UGT2B7
Adult Duo_1, Adult Duo_2, Chip_1, Chip_2, Chip_3, Organoids_1, Organoids_2, Organoids_3

Chemical carcinogenesis
CYP3A4, GSTA2, UGT2B17, CYP1A1, SUL2A1, ADH4, UGT1A4, ADH6, ADH1A, UGT2A3, UGT2B15, UGT2B7
Adult Duo_1, Adult Duo_2, Chip_1, Chip_2, Chip_3, Organoids_1, Organoids_2, Organoids_3

Protein digestion and absorption
KCNN4, COL13A1, COL6A2, SLC7A9, MME, MEP1B, COL6A1, SLC6A19, COL5A2, COL5A1, XPNPEP2
Adult Duo_1, Adult Duo_2, Chip_1, Chip_2, Chip_3, Organoids_1, Organoids_2, Organoids_3