Title: The Human Motor Cortex Contributes to Gravity Compensation to Maintain Posture and During
Reaching

Authors: R. L. Hardesty[1], P. H. Ellaway[2], and V. Gritsenko[1]

Running Head: corticospinal compensation for gravity

Affiliation

[1]   Department of Human Performance, Department of Neuroscience, Rockefeller Neuroscience
Center, West Virginia University, Morgantown, WV, USA

[2]   Department of Medicine, Imperial College London, London, UK

Address for Correspondence

V. Gritsenko, PO Box 9226, One Medical Center Drive, Morgantown, WV, 26505-9226. Phone: (304)
293-7719 vgritsenko@mix.wvu.edu

Abstract

How the neural motor system recruits muscles to support the arm against gravity is a matter of
active debate. It is unknown how the neural motor system compensates for the changing gravity-related
joint moments either when holding a steady-state posture or during movement between postural steady
states, e.g., during reaching. Here we used single-pulse transcranial magnetic stimulation to compare the
roles that the human primary motor cortex plays in the muscle recruitment to compensate for gravity. We
hypothesized that the motor cortex contributes to muscle recruitment to both maintain posture and to
compensate for changes in gravitational passive joint moments during movement. To test this hypothesis,
we used visual targets in virtual reality to instruct five postures and three movements with or against
gravity. We then measured the amplitude and gain of motor evoked potentials in multiple muscles of the
arm at several phases of the reaching motion and during posture maintenance. Stimulation below the
resting motor threshold, calibrated to the biceps muscle, caused motor evoked potentials in all muscles
during all postural and reaching tasks. The amplitude of motor evoked potentials was proportional to the
motoneuronal excitability measured as muscle activity. The coefficient of proportionality was positively
correlated with the postural component of muscle moment during posture and movement. Altogether our
results support the hypothesis. The observed contribution of the motor cortex to the recruitment of
multiple antagonistic muscles suggests a whole-limb strategy for overcoming passive gravity-related
moments with both active muscle moments and muscle co-activation that modulates limb impedance.

New & Noteworthy
Maintaining static posture and producing motion appear to be contradictory tasks for the nervous
system. In contrast to this seeming dichotomy, our results show that the motorcortical control signals play
the same role in both tasks when it is framed in biomechanical terms, i.e., muscle contractions needed to
compensate for gravity.

Keywords
Corticospinal, descending, biomechanics, impedance

Introduction

Posture and movement control are distinct tasks for the neural motor system. Posture requires maintaining a steady state, such as holding the elbow flexed against gravity when typing. In contrast, movement can be viewed as a transition between steady states. Traditionally, the neural control of these tasks has also been subdivided into a focal movement and postural adjustments. An example of the neural command for a focal movement would be the activation of arm muscles to produce a reaching movement, while the postural adjustment would be the activation of trunk muscles that compensate for the change in the body center of mass (Massion 1992). Biomechanically, both movement and posture, including postural adjustments, can be accomplished in two ways, 1) generating active joint moments with muscle contractions, termed muscle torques and 2) changing joint stiffness and viscosity components of mechanical impedance through muscle co-contraction. The forces generated by muscle contraction sum around a joint's degrees of freedom (DoFs) with directions defined by muscle geometry and moment arms. When they create counterbalanced muscle moments that sum up to zero, that defines the stiffness and viscosity components of mechanical impedance (Zajac 1989). Thus, increasing mechanical impedance through the co-contraction of antagonistic muscles can help maintain posture in presence of internal and external perturbations. However, when the muscle forces do not perfectly balance out around the joint, muscle torques arise. When these muscle torques are equal and opposite to the passive torques caused by gravity and any other external or internal force, no motion is produced or the same posture is maintained. Alternatively, when the amplitude of these muscle torques are larger than the passive torques, then active motion happens. Active motion is under the direct control of the nervous motor system through muscle contractions, we termed this component of motor control focal movement. Alternatively, when the amplitude of muscle torques are smaller than the passive torques, then passive motion results, which is not under the direct control of the nervous system. In this case, modulating mechanical impedance can enable the nervous system to keep control over the passive motion of the limb. Based on these biomechanical considerations, several models of the neural mechanisms for the control of posture and movement have been proposed. These models consist of two parallel or serialized controllers, one for generating the focal movement command through active muscle torques and one for posture maintenance through mechanical impedance (Ghez et al. 2007; Lametti et al. 2007; Todorov and Jordan 2002; Yadav and Sainburg 2011).

In this study of reaching, we further subdivide focal movement into posture- and movement-related components. The posture-related component of focal movement is the transition from the starting posture to a new posture. It is accompanied by the changing gravitational moments at arm joints caused by the different orientation of arm segments relative to the gravity vector. The second movement-related component is the residual muscle activity responsible for propulsion, i.e., acceleration and deceleration phases of motion. These posture- and movement-related components of focal movement can be observed in electromyographic (EMG) profiles. The former corresponds to the static component of EMG defined as the change in EMG amplitude between the starting and ending posture with a linear or cosine transition during movement; the latter is the phasic component of EMG we typically observe as bursts (Buneo et al. 1994; Flanders et al. 1996; Olesh et al. 2017). Here we suggest that the posture-related component of focal movement or the corresponding static EMG is generated by the same neural mechanism as that responsible for posture maintenance, specifically that for postural adjustment during focal movement. To test this idea, we hypothesize that the motor cortex contributes to muscle recruitment to both maintain posture and to compensate for changes in gravitational passive joint moments during movement.

Evidence in cats suggest that postural adjustments are initiated and shaped by the output of the primary motor cortex via the corticospinal tract through its collaterals to the reticulospinal tract (Schepens and Drew 2004; Yakovenko and Drew 2009). In humans, the excitability of the corticospinal tract and its collaterals has been studied using single-pulse transcranial magnetic stimulation (TMS). The descending tracts converge on the motoneurons, the final common pathway. Consequently, TMS over the motor cortex activated the corticospinal tract and its collaterals, which in turn activate spinal motoneurons

causing motor evoked potentials (MEPs) that can be observed in EMG. MEPs provide a measure of
instantaneous excitability or recruitment of both pre- and post-synaptic elements of the corticospinal tract
and its collaterals (Bestmann and Krakauer 2015; Di Lazzaro et al. 2004; Reis et al. 2008; Rothwell
1997). TMS at low intensity does not disrupt neural state (Cros et al. 2007), therefore changes in MEP
amplitudes can reveal the changes in the descending contribution to muscle recruitment during posture
and movement. For example, MEP amplitudes has been shown to change with different hand postures
reflecting different corticospinal contribution to the recruitment of muscles supporting those postures
(Perez and Rothwell 2015). The amplitude of a given MEP also depends on the excitability of the
motoneuron pool of the parent muscle (Darling et al. 2006; Groppa et al. 2012; MacKinnon and Rothwell
2000; Taylor 2006). However, the proportional contribution of the presynaptic elements to muscle
recruitment, termed corticospinal gain, can be estimated as the proportion between MEP amplitude and
the amplitude of the recruitment of motoneuronal pools observed with EMG in absence of MEPs
(Gritsenko et al. 2011). During movement the corticospinal gain is modulated in a phase-dependent
manner independently from the profile of EMG in wrist muscles and proximal antigravity muscles
(Lemon et al. 1995). Here we directly compare the corticospinal gain in the same muscles involved in
both posture and movement to determine if the changes in corticospinal gain reflect the changes in gravity
load during both posture and movement.

Materials and Methods

Subjects

The Institutional Review Board approved all procedures in this study (Protocol #1309092800).
Potential participants with any musculoskeletal pathologies or injuries, prior history of seizures, fainting,
or tinnitus, or those taking prescribed psychoactive medications were excluded. We obtained informed
consent prior to the start of experiments. We recruited 10 healthy individuals (6 male, 4 female, 24.3 ±
1.8 years old, 76.3 ± 14.5 kg). All participants reported to be right-hand dominant.

Motion Capture and Tasks

During the experiment, arm postures and reaching goals were defined in virtual reality
environment created using Vizard software (WorldViz) and Oculus headset (Fig. 1A). Spherical virtual
targets 8 cm in diameter were displayed in a sagittal plane and defined a set of postures and reaching tasks
as described in detail in (Hardesty et al. 2020) (Fig. 1B). The movements were similar to the planar
movements in a horizontal plane in Gritsenko et al. (2011) but occurred in a vertical sagittal plane without
external mechanical constraints.

Figure 1 about here

To minimize the inter-subject variability in kinematics during experiments, the locations of
virtual targets were adjusted based on individual segment lengths. The targets were displayed relative to
each individual's shoulder marker on the acromion (Fig. 1B). The experiment was subdivided into two
sessions, Session 1 with static postures and Session 2 with reaching movements. In both sessions the
participants were asked to reach to or hold at the virtual target with their right (dominant) arm without
moving the trunk while keeping the elbow close to the trunk and a neutral, pronated wrist.

During Session 1, participants were asked to reach to a virtual target and hold there for up to 1
minute before moving to the next target and holding there. The starting position for the Resistive task was
the same as the ending position for the Control task, therefore 5 postures were explored (Fig. 1B). While
the participant was holding the arm in one posture, we applied 12 TMS pulses more than 5s apart at 90%
of resting motor threshold (RMT, see below for details). Six or more TMS pulses has been shown to
provide sufficiently reliable estimates of MEPs (Lewis et al. 2014).

During Session 2, each trial started with the appearance of start (green) and stop (red) targets
(Fig. 1C). The colors and target locations did not change until the participants placed their index finger,
indicated with a yellow sphere, into the start target. One second after this occurred, the stop target

changed color from red to green, directing individuals to begin the movement (Fig. 1C). The target pairs for a given task were presented to each individual in the same pseudorandom order generated prior to the study. Each task was repeated 138 times (total of 414 trials). The movement trials were divided into stimulation trials (126 per movement, 378 total) and non-stimulation control trials (12 per movement, 36 total) interspersed randomly. During the stimulation trials TMS was performed at 90% of the RMT. In one half of trials (189 of 378), the TMS was triggered once per trial at a random delay of $0 - 550$ ms after the participant touched the start target. This triggering method ensured that MEPs were observed either preceding or around the movement onset time. In the other half of stimulation trials, the TMS pulses were triggered once per trial after the participant left the start target at a random delay of $0 - 550$ ms. This triggering method ensured that MEPs were observed through the duration of the movement. The timing of each TMS pulse was recorded for post-hoc normalization to the movement phase and binning described below.

Motion capture was performed using the Impulse system (PhaseSpace). Nine light-emitting diode markers were placed on bony landmarks of the arm and trunk using the best practice guidelines (Robertson et al. 2013). Marker coordinates were sampled at 480 Hz using Recap software (PhaseSpace). The marker locations were also streamed to virtual reality and used to represent the three main segments of the arm (hand, forearm, and upper arm) as a stick figure (Fig. 1A). Motion capture, electromyography (EMG, details below), and virtual events were synchronized using custom hardware as described in (Talkington et al. 2015).

Joint torques

After experiments, motion capture data from Session 2 were low pass filtered at a cutoff frequency of 10Hz. The mean residuals of marker triangulation ± standard deviation across individuals were 5.7 ± 0.45 mm. Joint angles were calculated by defining local coordinate systems of four rigid bodies for the trunk, humerus, forearm, and hand using at least 3 markers per rigid body (Fig. 1B). To calculate active joint torques, we used a dynamical model similar to that used for the selection of tasks as described in (Hardesty et al. 2020). The model was customized to individual morphology by scaling segment length and inertia based on individual's height and weight using published average proportions (Winter 2009). Inverse simulations were run with the scaled models using the angular kinematic trajectories averaged per task to obtain applied joint torques for a given individual and task, termed muscle torques. The muscle torques were verified using forward simulations and the simulated angular kinematics was compared to the angular kinematics obtained from motion capture data by calculating a root-mean-squared error (RMSE: 0.059 ± 0.035 rad). Positive rotations defined by local coordinate systems are illustrated in Fig. 1B. Analysis was done on the angles and torques around the X axes, i. e. flexion/extension degrees of freedom at all three joints, because motion was primarily in the sagittal plane and out-of-plane torques were negligible. The postural component of muscle torque, termed postural torque, was extracted by running simulations without gravity and subtracting those from the muscle torques simulated in presence of gravity as described in detail in (Olesh et al. 2017). The postural torque has been shown to follow closely the static component of EMG during similar reaching movements (Olesh et al. 2017).

Electromyography

Muscle activity and responses to TMS were recorded in twelve upper limb muscles using Trigno (Delsys Inc.), a wireless surface EMG system. The recorded muscles included four muscles spanning the shoulder, three muscles spanning both shoulder and elbow, two muscles spanning only the elbow, and three muscles spanning the wrist (Table 1). Muscles were identified based on anatomical landmarks and palpation during contraction; EMG sensors were placed on muscle bellies oriented longitudinally along the muscle fibers EMG signals were sampled at 2 kHz with a gain of 1000. EMG recordings were high pass filtered at 10Hz to remove any signal drift and rectified prior to any EMG/MEP quantification. EMG

180   profiles from trials without TMS were low pass filtered at a cutoff frequency of 20Hz and normalized per
181   individual using a maximum value for each muscle across all tasks.

183   **Table 1: Muscle nomenclature and MEP scaling with background EMG**

Muscle	Abbr.	Joint spanned	static MEP slope	dynamic MEP slope
pectoralis	Pec	shoulder	1.00–1.35–1.53	1.31–1.50–1.61
anterior deltoid	AD	shoulder	0.93–1.76–3.13	1.08–1.80–2.60
posterior deltoid	PD	shoulder	1.15–1.97–3.64	2.11–2.97–4.52
teres major	TM	shoulder	1.13–1.66–2.09	1.43–1.83– 2.92
triceps (long head)	TriLo	elbow, shoulder	1.36–2.31–4.22	1.79–2.36–3.90
triceps (lateral head)	TriLa	elbow	1.47–3.09–5.46	2.56–4.13–6.43
biceps (long head)	BicL	elbow, shoulder	1.56–4.41–6.83	3.08–4.01–5.43
biceps (short head)	BicS	elbow	1.64–3.75–5.90	2.29–2.64–4.60
brachioradialis	Br	elbow, wrist	3.94–5.89–10.10	4.63–5.49–7.61
flexor carpi radialis	FCR	wrist	1.73–4.85–7.44	3.07–5.11–6.19
flexor carpi ulnaris	FCU	wrist	3.43–4.49–9.46	2.99–4.56–9.79
extensor carpi radialis	ECR	wrist	2.84–5.05–5.58	2.93–4.54–5.59

Abbr. indicates abbreviated muscle names. MEP indicates motor evoked potential. Joint spanned lists the
flexion/extension degrees of freedom that were included into the linear regressions for the analysis of the
relationships between MEPs and joint angles and postural torques. Values show 25^{th} percentile – median slope – 75^{th}
percentile across individuals. Static MEP / EMG are the same slopes as B in eq. (2).

Transcranial magnetic stimulation
We assessed the corticospinal contribution to individual muscles using single-pulse TMS
delivered by the Super Rapid stimulator with a figure-of-eight coil (Magstim). The coil was oriented
tangentially to the scalp, oriented at a 45° angle to the midline with the handle pointing posteriorly and
laterally (Fig. 1C). The coil location over the scalp and its orientation was maintained using the Brainsight
neuronavigation system (Rogue Research). This coil positioning has been shown to preferentially activate
pyramidal tract neurons (Kaneko et al. 1996; Di Lazzaro et al. 1998; Nakamura et al. 1996) and to
generate responses proportional to cortical activity

Focal TMS with the figure-of-eight coil causes MEPs in multiple muscles (Roth et al. 1991; Tofts
1990). The amplitudes of these MEPs and which muscles are involved depends on the strength of the
magnetic field generated by the TMS device, the location and orientation of the coil over the scalp, and
individual anatomy. The location of the coil was selected using the hot-spot method (Ellaway et al. 1998;
Traversa et al. 1997), during which the coil was moved over the estimated location of the primary motor
cortex until a location with at least 50 µV motor evoked potential (MEP) in biceps was evoked. This
controlled for the anatomical differences between individuals and defined a consistent stimulation
location on the motor homunculus (Penfield and Rasmussen 1950). This was done with the anticipation
that stimulating the same anatomical location with the lowest corticospinal excitability measured in
biceps would activate the same component of the whole corticospinal tract across individuals and produce
similar responses in other muscles across individuals.

The strength of the magnetic field is typically tailored to individuals using a resting motor
threshold (RMT) method, i.e., identifying the lowest magnetic field strength that evokes MEPs 50% of
the time. This reduces the inter-subject variability in TMS responses. RMT was determined at the hot-
spot location by varying the stimulation intensity until a MEP > 50 µV was evoked 50% of the time in

BicL. This procedure ensured that the stimulation amplitude was adjusted to individual differences in
corticospinal excitability at rest at the time of experiment and thus, further minimized inter-subject
differences in MEP amplitudes. A single TMS pulse was delivered in a given trial and each trial lasted 5
s. This was done to minimize the instances when two TMS pulses would be delivered at > 0.2 Hz
instantaneous frequency to avoid any long-term changes in corticospinal excitability (Chen et al. 1997).
The experiment consisted of two consecutive sessions conducted on the same day. The number of
repetitions of some conditions was limited due to time constraints in efforts of preventing fatigue.

The amplitudes of MEPs produced using the described method are stable across days (Pascual-
Leone et al. 1995; Uy et al. 2002) and have sufficient spatial resolution to differentiate proximal and
distal muscles (Pascual-Leone et al. 1997). Here MEPs were quantified using two methods, peak-to-peak
amplitude and area methods commonly used in other studies (Ellaway et al. 1998; Pascual-Leone et al.
1998; Thickbroom et al. 1999). MEPs in upper limb muscles were observed within the time period lasting
from 10 ms to 50 ms after the TMS pulse (Fig. 2A). The peak-to-peak MEP amplitude was calculated by
subtracting the minimum from the maximum of the EMG trace within this time period (Fig. 2B). The
MEP area was calculated by first rectifying the EMG trace within this time period and then integrating it
(Fig. 2C).

For MEPs collected in Session 1, termed static MEPs, both peak-to-peak and area methods were
included into the analysis. The background EMG was calculated over the 40-ms time window directly
preceding the TMS pulse using the corresponding peak-to-peak or integrated method respectively. We
defined the presence of a MEP as a peak-to-peak amplitude of at least 5 standard deviations above
background EMG amplitude. The probability of evoking MEP was calculated by dividing the number of
detected MEPs by the number of stimuli under the same conditions. To determine the consistency of
MEPs based on MEP area metric, we calculated the coefficient of variation (CV) across repetitions of the
same task per individual per muscle.

237   Figure 2 about here

239   The peak-to-peak MEP amplitude (MEP_{amp}) during isometric contraction has been shown to
240 follow the Boltzmann equation (Darling, 2006; Devanne, 2002):

$$MEP_{amp} = \frac{MEP_{max}}{1 + e^{\left(\frac{S_{50}-S}{k}\right)}} + B \times PreEMG + C_1 \,, \tag{1}$$

242
243 where MEP_{max} is the maximum MEP amplitude, S_{50} is the stimulation intensity to elicit a MEP at 50% of
244 the maximum amplitude, S is the stimulation intensity, and $PreEMG$ is the amplitide of EMG measured
245 directly preceding stimulation, and C_1 is a constant. In this study, we performed all stimulations at a
246 consistent intensity of 90% the RMT, which makes the first term a constant. Therefore equation (1) can
247 be simplified and adapted for examining the linear relationships between EMG and MEP area as follows:
248

$$MEP_{amp} = B \times PreEMG + C_2 \,, \tag{2}$$

249
250 where MEP_{amp} in our experiment is MEP area measured as described above, $PreEMG$ term is the
251 background EMG calculated as describe above, and B is the slope of the linear relationship between MEP
252 and EMG during posture maintenance. C_2 was assumed to be equal to 0 expecting no MEPs in fully

relaxed muscles due to stimulation below RMT. We derived the slope B in static Session 1 and used it to normalize the MEP area values recorded during movement in Session 2.

We calculated MEP latencies using a procedure similar to that used to determine kinematic onsets. Unrectified EMG signals were aligned on the TMS pulse and averaged across repetitions of the same postures. The MEP latency was defined as the time from the TMS pulse to the first maximum of the third derivative of this averaged trace.

For MEPs collected in Session 2, termed dynamic MEPs, both peak-to-peak and area methods were included into the analysis. All trials during which TMS was applied were included into the MEP metrics without determining directly whether a MEP was evoked or not in a given trial. This is because of difficulties distinguishing MEPs from compound motor unit action potentials in presence of changing EMG during movement. MEP area and the CV values were calculated as in Session 1. The MEP area values obtained from single trials were grouped into 5 temporal bins, each corresponded to 20% increments of phase duration from onset to offset of movement (Fig. 2D). This binning procedure grouped MEPs occurring at similar times during movement and provided adequate repetitions to estimate median values. Additionally, TMS responses that occurred up to 20% phase duration prior to movement were grouped into a bin 0 (Fig. 2D). We defined a minimum repetition criterion of 5 MEPs based upon (Lewis et al. 2014). Bins that contained < 5 MEPs were excluded from subsequent analyses. The MEP amplitude values were normalized by the median MEPs in the starting posture for the corresponding task and muscle from Session 1. Thus, MEP area values equal to 1 indicated that the MEPs during movement were equal to those at the starting posture for that movement.

To estimate the changed in corticospinal excitability that were independent from the motoneuronal excitability we calculated MEP gain. For this measure we sampled background EMG in trials without TMS at the same phase as TMS pulses in other trials of Session 2. We selected short chunks of EMG at the corresponding phase of movement as MEPs and processed them the same way as the MEP area or perk-to-peak metric (Fig. 2C). We then used the linear MEP-EMG relationship observed in static postures in Session 1 to predict MEP amplitude (and MEP_{Pred}) for a given background EMG level using the following equation:

$$MEP_{Pred} = B \cdot EMG_{dynamic} \tag{3}$$

where $EMG_{dynamic}$ is the median EMG within a given bin per muscle, task, and individual and slope B is from Eq. (2).

The MEP gain was calculated as the ratio between the observed MEP area (MEP_{Dyn}) and MEP_{Pred} from Eq. (3). The MEP gain equal to 1 means that the median MEP at a given phase is equal to the predicted MEP based on the linear relationship between EMG and MEPs described in eq. (2) with the same slope as that observed in static trials. MEP gain values < 1 indicate a decrease in corticospinal contribution to a given muscle relative to that observed in static trials. By controlling for the linear relationship between EMG and MEP, the MEP gain metric reveals changes in the corticospinal tract that are independent of motoneuronal excitability.

Results in text are reported as mean values ± standard deviation across individuals, unless otherwise indicated.

Statistical Analyses

The amount of shared variance between median MEP area values and the background EMG were calculated using the coefficient of determination (R^2) across all movement phases per task per muscle per individual. For static MEPs, the background EMG was the value of $PreEMG$ from Eq. (2). For dynamic MEPs, the background EMG was the value of $EMG_{dynamic}$ from Eq. (3). The median MEP area was calculated as described above for each session. For data from Session 1, the linear regressions were fitted between MEPs, EMG, and the postural torques across all 5 postures (Fig. 1B). For data from Session 2,

the linear regressions were fitted between MEPs and EMG across movement phases (bins 0 – 5). Significance was determined based on the goodness of fit statistic with Sidak correction for repeated testing setting $\alpha = 0.0043$. The between-task differences in individual R^2 values were normally distributed based on a single-sample Kolmogorov-Smirnov test (kstest function in MATLAB). Therefore, the significance of between-task differences was determined using paired t-tests on individual R^2 across tasks using a ttest function in MATLAB. A separate test was done for each muscle using Sidak-corrected $\alpha = 0.017$.

To determine if MEP gain was changing during movement, the values were divided into early and late groups based on whether MEPs occurred in the acceleration or deceleration phases of reach respectively. The early group comprised MEP gain values from bins 0, 1, and 2 and the late group comprised values from bins 4 and 5 (Fig. 2D). Bins 0 and 1 usually included fewer MEPs, therefore the total number of values in the two groups was comparable, > 40 in each acceleration or deceleration phase per muscle per individual per task. Single-trial MEP gain values were not normally distributed. Therefore, significance was determined using a two-sample Kolmogorov-Smirnov test using kstest2 function in MATLAB. Tests were applied to compare single-trial MEP gain values occurring in the acceleration and deceleration phases of movement for each individual per muscle with Sidak-corrected $\alpha = 0.0051$. In contrast, the differences between median MEP gain values were normally distributed based on a single-sample Kolmogorov-Smirnov test. Therefore, paired t-tests were used to determine if MEP gain was modulated during movement per task per muscle with Sidak-corrected $\alpha = 0.017$.

To quantify the changes in MEP gain between tasks, differences between individual median MEP gain values were calculated between tasks separately for acceleration and deceleration phases for each muscle. Then, corresponding differences in joint angle, muscle torque and its components were calculated for each individual per DOF. The joint angles in the acceleration (bins 0, 1, and 2) and deceleration (bins 5 and 6; Fig. 4A) phases of movement were averaged per individual per DOF and subtracted between tasks. The postural torques in the acceleration (bins 0, 1, and 2) and deceleration (bins 5 and 6; Fig. 4B and C) phases of movement were also averaged per individual per DOF and subtracted between tasks. For the dynamic component, maximal values in the acceleration phase and minimal values in the deceleration phase due to changing sign were calculated per individual per DOF and subtracted between tasks (Fig. 4D). The resulting difference values were normalized to their range across all individuals. Lastly, liner regressions were fitted separately for acceleration and deceleration phases of movement between each of the kinematic and dynamic variables and MEP gain of muscles that span the corresponding joint (Table 1). Sidak correction for repeated regressions was used to set $\alpha = 0.0127$.

Results

Our evidence supports the involvement of the corticospinal tract in posture maintenance in humans. At stimulation amplitudes below RMT, we observed MEPs during static posture, termed static MEPs, in most muscles. For example, biceps muscle participates directly in posture maintenance by holding the elbow and shoulder flexed in the tested postures (Fig. 3A). We found that the probability of evoking a MEP in this muscle was higher than expected from stimulation at the RMT in most individuals (Fig. 3B). Furthermore, the probability of evoking a static MEP was higher than expected in all muscles, even those not directly involved in supporting the arm against gravity (Fig. 3C). Altogether this shows that the overall corticospinal excitability was increased in a consistent manner when the arm was held against gravity in different postures compared to when the arm was relaxed.

Figure 3 about here

The static MEPs were evoked at similar latencies and with consistent amplitudes across individuals. The average latency across muscles was 15 ± 1.9 ms (Fig. 3D). These latencies broadly reflected the proximal to distal distribution of muscles and the associated differences in the distance travelled by the action potentials evoked by TMS.

The MEP amplitudes were highly consistent as evidenced by the CV of static MEPs equal to 0.35
and dynamic MEPs equal to 0.5 (Fig. 4A), which is comparable to reported values recorded at rest and in
relaxed muscles (Ellaway et al. 1998; Jung et al. 2010; Wassermann 2002; Zanette et al. 1995). The larger
CV for the dynamic MEPs compared to the static MEPs is likely driven by the changing motoneuronal
excitability reflected in the changing background EMG and muscle lengths during movement (Day et al.
1991; Kiers et al. 1993; Thickbroom et al. 1999). Furthermore, we observed that the static MEPs were
linearly related to the changes in background EMG across postures (Fig. 4B). This linear relationship
expressed in eq. (2) accounted for 25% of the static MEP variance (mean R^2 = 25% ± 13% across
muscles) with a small mean squared error of $2.2e^{-5} ± 4e^{-5}$. The regression slopes were above unity in all
muscles and individuals (Table 1, static MEP / EMG column). These slopes also broadly reflected the
proximal to distal distribution of muscles with distal muscles trending toward higher slopes than proximal
ones. The dynamic MEPs displayed many similar characteristics to the static MEPs. Dynamic MEPs were
also observed in all reaching tasks and in all muscles despite subthreshold stimulation. Dynamic MEP
latencies were on average 14.2 ± 2.5 ms, similar to static MEPs. The dynamic MEPs were linearly related
to the changes in background EMG during movement (Fig. 4C). In all muscles the ratios between
dynamic MEPs and EMG in corresponding bins were above one and the ratios were similar or in some
muscles above the static slopes (Table 1, dynamic MEP / EMG column).

365   Figure 4 about here

The virtual targets were effective in evoking the desired behavior and standardizing the
movement kinematics and dynamics across all individuals (Fig. 5). Holding the arm against gravity in the
five postures and transitioning between them during reaching was accomplished with postural torques of
different magnitude at the major arm joints (Fig. 5B). The postural torques were larger during Session 1
(Fig. 5B, hold) than at the start and end of movement during Session 2 (Fig. 5B, bin 0 and 5) because of
the additional contribution of the propulsive torques to accelerate and descelerate the arm during
movement. However, the differences between postural torques across tasks were the same in both
conditions. For example, the postural torque about the shoulder in the starting position for the Resistive
condition was smaller than that in the Control and Assistive conditions in both Sessions (Fig. 5B,
Shoulder). The shoulder motion in the Control and Assistive tasks was with gravity, both were
accompanied by decreasing postural torque at the shoulder, while in the Resistive task the shoulder
motion was against gravity and the postural torque was increasing (Fig. 5B Shoulder). In contrast, the
postural torques about the elbow and wrist joints were more alike in the Assistive and Resistive (Fig. 5B,
Elbow & Wrist, blue and red) tasks despite different motion of the elbow in these two tasks (Fig. 5A,
Elbow & Wrist, blue and red). The postural torques at the elbow and wrist joints were equal across all
starting postures and in the beginning of movement; they were diminishing during reaching in the
Assistive and Resistive tasks but not in the Control task. These results suggest that the postural control of
the elbow and wrist can be coupled, while the postural control of the shoulder needs to be decoupled from
that of the elbow and wrist to appropriately compensate for the changing gravity-related passive torques.
This applies to both posture-related component of focal movement and maintaining static posture.

388   Figure 5 about here

The temporal profiles of postural torques during reaching can be used to define the tonic
components of EMG in multiple muscles (Olesh et al. 2017). Here too we observed that in Session 1 the
EMG amplitude in most muscles spanning the shoulder was linearly dependent on the amplitude of

shoulder postural torque across the five postures with shared variance for Pec = 30%, AD = 65%, PD =
38%, TM = 31%, TriLo = 45% at $p < 0.001$ for each (significant $\alpha = 0.004$). In contrast, the EMG
amplitude of muscles spanning the elbow or wrist was weakly dependent on the amplitude of elbow or
wrist postural torques (R^2 for TriLo = 6%, TriLa = 4%, BicL = 1%, BicS = 6%, FCR = 2%, FCU = 7%,
ECR = 5% at $p > 0.07$ for each and BicL = 11% at $p = 0.017$, significant $\alpha = 0.004$). Furthermore, the
static MEPs only in the muscles spanning the shoulder were linearly related to the shoulder postural
torque in Session 1 with high shared variance (R^2 for Pec = 49%, AD = 51%, PD = 43%, TM = 47%,
TriLo = 48% at $p < 0.001$ for each; significant $\alpha = 0.004$). Interestingly, removing the static MEP
dependency on the background EMG by calculating static MEP gain did not fully remove the MEP
dependency on the postural torque. There remained a linear dependency between shoulder postural torque
and static MEP gain in Pec ($R^2 = 19\%$ at $p = 0.002$) and in TriLo ($R^2 = 23\%$ at $p < 0.001$), indicating
modulated descending contribution to the recruitment of these muscles. In the rest of the muscles, the
static MEPs were not correlated with the changes in the elbow or wrist postural torques across the 5 static
postures. This shows that to hold the arm against gravity, the descending contribution to most muscles
was elevated. Furthermore, the descending contribution to the recruitment of Pec and TriLo was
modulated with the shoulder postural torque.

The dynamic MEP gain values during movement varied more between individuals than between
muscles. This suggests that the inter-subject variability may contain biomechanically relevant information
rather than noise. To test this idea, we performed a regression analysis, in which we included only the
dynamic MEP gain values per individual per muscle that were significantly different between tasks in a
corresponding phase of movement. The MEP gain in monoarticular shoulder muscles (Pec, AD, PD, and
TM) was not correlated with the changes in shoulder postural torque at the beginning of movement (Fig.
6A). This shows that change in the descending contribution to the recruitment of monoarticular shoulder
muscles does not serve the role of counteracting the passive postural torque at the shoulder in the
beginning of movement. In contrast, the dynamic MEP gain in the biarticular biceps and triceps muscles
(BicL and TriLo) was increased when the arm was more outstretched, i.e. at larger shoulder flexion angle
and lower elbow flexion angle (Fig. 5, black and blue). The outstretched postures in Assistive and Control
tasks were accompanied by larger shoulder postural torque, while the elbow torque remained unchanged.
Consequently, the dynamic MEP gain in BicL and TriLo was linearly proportional to the increase in
shoulder postural torque between tasks (Fig. 6B). The elbow postural torque was similar between tasks in
the beginning of movement (Fig. 5B), therefore no correlations with MEP gain in the muscles spanning
the elbow were observed (Fig. 6C). The wrist angle and postural torque were similar across tasks in the
beginning of movement (Fig. 5) and thus no correlations with MEP gain in wrist muscles (FCU, FCR,
and ECR) were observed (Fig. 6D). This shows that the inter-subject variability of the dynamic MEP gain
in BicL and TriLo was linearly dependent on the individual differences in the postural torques in the
beginning of reaching tasks. Furthermore, these results suggest that the changes in the descending
contribution to the recruitment of biarticular biceps and triceps muscles in the beginning of movement
compensate for the interaction torques between shoulder and elbow caused by the changing passive
gravity torques at these joints.

433   Figure 6 about here

At the end of movement, the MEP gain in the monoarticular shoulder muscles was proportional to
the shoulder and wrist postural torques respectively (Fig. 7A). In contrast, the dynamic MEP gain in the
biceps and triceps muscles did not depend on either shoulder or elbow postural torque (Fig. 7B, 7C)
despite large differences in these torques between tasks (Fig. 5). Furthermore, the MEP gain in the wrist
muscles and brachioradialis was proportional to the wrist postural torque in absence of motion around the

wrist (Fig. 7D). These results suggest that the postural adjustments to stabilize the arm at the new posture
at the end of movement are driven by modulated descending contribution to the recruitment of multiple
monoarticular shoulder muscles, brachioradialis, and all recorded muscles spanning the wrist. The
changes in the descending contribution to proximal muscles serve the role of counteracting the passive
postural torque at the shoulder, while changes in the descending contribution to distal muscles, including
brachioradialis, serve the role of counteracting the passive postural torque at the wrist. However, the
descending contribution to the recruitment of biceps and triceps muscles at the end of movement does not
serve the role of counteracting the passive postural torque at the joints that these muscles span.

Figure 7 about here

Discussion
Here we used single-pulse TMS to probe corticospinal excitability during unconstrained postural
and reaching tasks. We observed that stimulation below RMT evoked MEPs in all muscles during both
holding arm in static posture and during reaching. This provides evidence that both maintaining a static
posture against gravity and transitioning from one posture to another involve increased motorcortical
contribution to the recruitment of most muscles of the arm. Under all conditions MEPs were linearly
dependent on the background EMG with 25% of variance in static MEP amplitude and even more in
dynamic MEP amplitude captured by this linear relationship (Fig. 4C). This is consistent with previous
work showing a large amount of MEP variance is captured by a simple linear relationship between MEP
amplitude and EMG during isometric voluntary contractions (Darling et al. 2006). The slope of this linear
relationship quantifies a certain level of the motorcortical contribution to the recruitment of the
corresponding muscle. Any changes in this slope or changes in MEP gain reflect the change in the level
of motorcortical contribution to the recruitment of a given motoneuron pool. For example, the slopes of
the linear relationship between MEP and EMG in distal muscles tended to be higher than proximal
muscles (Table 1). This is consistent with prior work showing a greater involvement of the corticospinal
tract in the voluntary activation of distal vs. proximal muscles (Rothwell et al. 1991; Turton and Lemon
1999). In our study the MEP gain in pectoralis and the biarticular triceps muscles changed in proportion
to the shoulder postural torque during posture maintenance. Moreover, in the beginning of movement, the
MEP gain in both biarticular biceps and triceps muscles changed in proportion to shoulder postural
torque. At the end of movement, the MEP gain in the monoarticular shoulder muscles (pectoralis,
posterior deltoid, and teres major) was proportional to the shoulder postural torque and the MEP gain in
wrist muscles (flexor carpi radialis, flexor carpi ulnaris, and extensor carpi radialis) was proportional to
the wrist postural torque. This shows that the motorcortical contribution to the recruitment of more arm
muscles increased in proportion to postural torques during movement compared to that during posture
maintenance. Altogether, our results support the hypothesis that the motor cortex contributes to muscle
recruitment to both maintain posture and to compensate for changes in gravitational passive joint
moments during movement.

The reaching tasks used in our study were selected to maximize the differences in the postural
forces that occur at the major arm joints while minimizing the differences in elbow and wrist kinematics.
Consequently, large differences in the postural components of muscle torques were needed to hold the
arm in different postures or to transition between postures (Fig. 5). The transitions between postures
required modulated motorcortical involvement in the recruitment of muscles forming distinct groups. In
the acceleration phase of movement, the MEP gain in both antagonistic biarticular biceps and triceps
muscles was proportional to the shoulder postural torque (Fig. 6B). This suggests that at the beginning of
movement the motorcortical projections contribute to the coactivation of biceps and triceps muscles (Fig.
8, red projections). This may serve the purpose of compensating for the interaction torques caused by the

changing gravity load at the shoulder through increasing mechanical impedance of shoulder and elbow joints (Gribble and Ostry 1999; Gritsenko et al. 2011). This was followed by the MEP gains in proximal and distal muscles being proportional to the shoulder and wrist postural torque respectively at the end of movement (Fig. 7A, 7D). These results are consistent with earlier observations of phase-specific modulation of corticospinal gain in multiple arm and hand muscles during reach and grasp movement (Lemon et al. 1995, 1996). Our results indicate more specifically that in the deceleration phase of movement the motorcortical projections contribute to the coactivation of subsets of proximal and distal muscles, which likely increased the mechanical impedance of shoulder and wrist joints and stabilized the arm in the new posture (Fig. 8, blue projections). We postulate that the postural torque in both acceleration and deceleration phases of movement reflect postural adjustments that accompany focal movement, as it is precisely the active joint torque needed to support the arm against gravity. Therefore, the observed modulation of MEP gain with the postural torque both in our postural and reaching tasks is likely to involve the reticulospinal tract via collaterals from the corticospinal tract (Keizer and Kuypers 1989). The importance of the reticulospinal tract in the integration between posture and movement, including reach, is well established in cats (Schepens and Drew 2004, 2006) and primates (Baker 2011; Riddle et al. 2009). The TMS methodology used here may help study the maladaptive changes in the neural mechanisms of postural adjustments in humans with neurological conditions and identify avenues of new neuromodulative interventions.

<center>Figure 8 about here</center>

The limitations of study methodology may limit the generality of our conclusions. The first limitation is that the contribution of descending projections from premotor and supplementary motor cortices to motoneuronal excitability cannot be ruled out. In our study we used focal TMS that was targeted and scaled based on responses observed in biceps EMG. This method is designed to limit the volume of current density that is sufficient to stimulate cells located in the arm area of the primary motor and possibly primary sensory motor cortices (Pascual-Leone et al. 1997; Pascual-Leone and Walsh 2002). Therefore, the descending projections from premotor areas were likely not stimulated and their contribution to muscle recruitment and the compensation for passive torques not quantified in the present study. The second limitation is the possibility that using a different muscle for determining the RMT would alter the modulation of MEP gain in other muscles. It seems unlikely for proximal muscles, because the spatial resolution of TMS is rather low. Stimulation over scalp locations 1-2 cm apart has been shown to crudely distinguish between proximal or distal muscle groups (Pascual-Leone et al. 1997). Here we have targeted a proximal muscle group by determining our hot-spot and RMT based on MEPs in biceps. If we choose another proximal muscle for thresholding, the hot-spot method will likely zero in on the same scalp location but a different magnetic field strength. Increasing or decreasing the magnetic field strength over the same spatial location in order to target another proximal muscle is likely to activate more or less respectively all the same muscles recorded here. The underlying cortical activity that drives the modulation of MEP gain would be the same. However, if we choose a distal muscle for thresholding, the hot-spot method will likely zero in on a different scalp location and a different magnetic field strength. In this case, proximal muscles are not likely to have MEPs due to their generally higher threshold (Rothwell et al. 1991; Turton and Lemon 1999). Therefore, the activity of the different underlying cortical area may cause different MEP amplitudes and gains across all muscles compared to those caused by the scalp area chosen based on the biceps hot-spot. Consequently, our results generalize to represent a subset of cortical contribution to the control of reaching, but they may not generalize to the more dexterous corticospinal control of distal muscles. The third limitation is that the MEPs produced by TMS during movement may saturate in some muscles at some phases of movement. This possibility is based on observations of similar or lower MEP amplitudes despite increasing background EMG from 10% to 40% of maximal voluntary effort for the former or 25% to >50% of maximal voluntary effort for the latter (Devanne et al. 1997; Kamen 2004). Here, we did observe a linear relationship between background EMG and MEP amplitude with similar slopes during both postural and reaching tasks (Table 1). Importantly,

during reaching tasks the range of EMG changes is larger than during the different postures, yet we did
not observe signs of saturation that would be evident from reduced slope of the regression during reaching
tasks. However, the slopes for distal muscles (Br, FCR, FCU, and ECR) were rather high, which suggests
a very high recruitment of these projections by TMS. Therefore, the potential saturation of corticospinal
contribution to the recruitment of distal muscles probed with TMS in our study cannot be completely
ruled out.

Endnote

543   All data relevant for assessing the conclusions of this study are presented in the manuscript and
544   figures.

Acknowledgements

546   We thank the study participants for generously giving their time; we acknowledge the data
547   collection and analyses contributions of Dr. W. Talkington; we acknowledge the technical support
548   expertly provided by B. Pollard; we are thankful for insightful and critical comments of Dr. S.
549   Yakovenko. The current affiliation of RLH is Albany Stratton VA Medical Center, Albany, NY.

Funding sources

552   RLH was supported by a training grants T32GM081741 and T32AG052375 from the National
553   Institute of General Medical Sciences. VG was supported by grants P20GM109098 and P30GM103503
554   from the National Institute of General Medical Sciences (https://www.nigms.nih.gov) and 1007978 from
555   the National Science Foundation. The funders had no role in study design, data collection and analysis,
556   decision to publish, or preparation of the manuscript.

Figure Legends

Figure 1. Experimental paradigm. A. Participant's view of the tasks in virtual reality. Black lines
connect virtual representation of LED markers and show the arm position and orientation in virtual reality
relative to targets. The yellow sphere represents the fingertip LED on the index finger, which the
participants were instructed to move to the center of the green target. B. Schematic of the dynamic model
used to calculate joint angles and torques from motion capture and representative endpoint trajectories
from the three tasks. Three segments (pink ovals) were used to simulate inertial properties of upper arm,
forearm, and hand; centers of mass are indicated by black and white circles. Positive directions for each
joint (DOF) are shown with black circular arrows. The start and stop target locations are shown for the
Control, Resistive, and Assistive tasks (black, red, and blue, respectively). The motion capture marker
locations are shown as yellow circles. C. Visual representation of the experimental paradigm. D. Example
timeline of events during a single trial of a movement task. RT stands for reaction time; GO represents the
color cue to start the movement.

Figure 2. Quantifying MEP amplitude. A. Biceps EMG trace aligned on TMS pulse (red line) during
reaching trials from the Control task in one individual. Grey area shows the window used to quantify
MEP amplitude and MEP area. B. Averaged EMG from BicL muscle in one individual performing the
Control task. EMG was aligned on TMS pulse (red line) and peak-to-peak MEP amplitude was calculated
with the shaded area. C. Averaged rectified EMG from BicL muscle in one participant performing the

Control task. EMG was aligned on TMS pulse (red line) and MEP area was calculated by integrating the
shaded area under the curve. D. EMG from FCU muscle in one participant performing the Assistive task.
MEP area values were binned according to the phase of movement, where 0 denotes the kinematic onset
of movement and 1 denotes the kinematic offset. Background EMG is from trials without TMS; predicted
MEPs were calculated using equation (3).

Figure 3. Static MEPs in BicL. A. EMG traces of the biceps muscle during static postures are displayed
for all participants. The black line corresponds to the time of TMS pulse (stim). B. Each bar corresponds
to the probability of a MEP occurring for each participant across all postures. The solid black line denotes
the probability of MEP occurrence at RMT. The dashed line corresponds to the increase in probability
that could be detected given the statistical power of our study design ($\beta = 0.9$).

Figure 4. A. Coefficient of variation (CV) across all muscles and postures or movement per individual.
CV for static MEPS are shown in red. CV from published studies are shown in black. CV for dynamic
MEP gain values are shown in blue calculated as described in Methods, i.e. by dividing recorded MEP by
predicted MEP from the corresponding bins (Fig. 2D). B. An example of the "best" (left) and "worst"
(right) linear fit in BicL from different individuals. Lines are mean trajectories; error bars are standard
deviations across individuals. C. Variance of dynamic MEPs explained by background EMG in individual
muscles across tasks. Horizontal lines show medians and bars show quartile ranges across individuals.
The horizontal red line and shaded area represent the mean variance of static MEPs explained by
background EMG.

Figure 5. Movement kinematics and dynamics. A. Joint angles calculated from motion-capture B. The
component of muscle torque that compensates for gravity load on the joints, termed postural torque.

Figure 6. Modulation of MEP gain between tasks in the beginning of movement. Circles show MEP
gain differences between tasks per individual per muscle. Thick black lines show regressions fitted to all
values in the plot with goodness of fit statistic (p and R^2) above each plot. Plots with significant linear
dependencies at alpha < 0.0127 have grey background. A. Differences in MEP gain between tasks in
pectoralis (Pec), posterior deltoid (PD), and teres major (TM) relative to the differences between tasks in
shoulder angle and in shoulder postural torque. B. Differences in MEP gain between tasks in triceps (long
head, TriLo) and biceps (long head, BicL) relative to the differences between tasks in shoulder angle and
shoulder postural torque. C. Differences in MEP gain between tasks in triceps (long and lateral heads,
TriLo and TriLa respectively) and biceps (long and should heads, BicL and BicS respectively) relative to
the differences between tasks in elbow angle and elbow postural torque. D. Differences in MEP gain
between tasks in brachialis (Br), flexor carpi ulnaris (FCU), flexor carpi radialis (FCR), and extensor
carpi radialis (ECR) relative to the differences between tasks in wrist angle and wrist postural torque.

Figure 7. Modulation of MEP gain between tasks at the end of movement. Circles show MEP gain
differences between tasks per individual per muscle. Thick black lines show regressions fitted to all
values in the plot with goodness of fit statistic (p and R^2) above each plot. Plots with significant linear
dependencies at alpha < 0.0127 have grey background. A. Differences in MEP gain between tasks in
pectoralis (Pec), posterior deltoid (PD), and teres major (TM) relative to the differences between tasks in
shoulder angle and in shoulder postural torque. B. Differences in MEP gain between tasks in triceps (long
head, TriLo) and biceps (long head, BicL) relative to the differences between tasks in shoulder angle and
shoulder postural torque. C. Differences in MEP gain between tasks in triceps (long and lateral heads,
TriLo and TriLa respectively) and biceps (long and should heads, BicL and BicS respectively) relative to
the differences between tasks in elbow angle and elbow postural torque. D. Differences in MEP gain
between tasks in brachialis (Br), flexor carpi ulnaris (FCU), flexor carpi radialis (FCR), and extensor
carpi radialis (ECR) relative to the differences between tasks in wrist angle and wrist postural torque.

Figure 8. Concept of the control schema for postural adjustments during reaching. A. Motorcortical
outputs for counteracting the changing gravity load on the arm. Grey outlines the human body and arm;

spindles show the basic anatomy of major muscle groups. Diamonds indicate motoneuron pools of the corresponding muscles. They are colored to indicate common input from the pyramidal cells in the primary motor cortex (triangles). Colors of the pyramidal contributions indicate changes in MEP gain in the acceleration phase of movement (red) and in the deceleration phase of movement (blue).

Extended Data 1. Software and data are included in two folders and described in the readme.txt file. The code for inverse kinematic and dynamic simulations is in the Simulation Functions folder. To obtain joint angles from a given set of marker coordinates, run getEuler_ARM_140205.m. To obtain joint torques from a given set of joint angles, run MODEL_getTorques.m. This function runs a Simulink model of the arm KINARM_Dyn_Inv.mdl. The code for plotting figures and statistics is in the Manuscript tests and plots folder. Run plot_RSquaredAnalysis.m to plot figures 6 C and D from data saved in mat files. Run getDynamicMEP_plot_EarlyLateTest.m to plot figures 7, 8, and 9 with associated statistics using data saved in mat files. This script also contains a power test for one of the regressions.

References

Baker SN. The primate reticulospinal tract, hand function and functional recovery. *The Journal of Physiology* 589: 5603–5612, 2011.

Bestmann S, **Krakauer JW**. The uses and interpretations of the motor-evoked potential for understanding behaviour. *Experimental Brain Research* 233: 679–689, 2015.

Buneo CA, **Soechting JF**, **Flanders M**. Muscle activation patterns for reaching: the representation of distance and time. *Journal of Neurophysiology* 71: 1546–1558, 1994.

Chen R, **Classen J**, **Gerloff C**, **Celnik P**, **Wassermann EM**, **Hallett M**, **Cohen LG**. Depression of motor cortex excitability by low-frequency transcranial magnetic stimulation. *Neurology* 48: 1398–1403, 1997.

Cros D, **Soto O**, **Chiappa KH**. Transcranial magnetic stimulation during voluntary action: directional facilitation of outputs and relationships to force generation. 1185: 103–116, 2007.

Darling WG, **Wolf SL**, **Butler AJ**. Variability of motor potentials evoked by transcranial magnetic stimulation depends on muscle activation. *Experimental Brain Research* 174: 376–385, 2006.

Day BL, **Riescher H**, **Struppler A**, **Rothwell JC**, **Marsden CD**. Changes in the response to magnetic and electrical stimulation of the motor cortex following muscle stretch in man. *The Journal of Physiology* 433: 41–57, 1991.

Devanne H, **Lavoie BA**, **Capaday C**. Input-output properties and gain changes in the human corticospinal pathway: *Exp Brain Res* 114: 329–338, 1997.

Di Lazzaro V, **Oliviero A**, **Pilato F**, **Saturno E**, **Dileone M**, **Mazzone P**, **Insola A**, **Tonali PA**, **Rothwell JC**. The physiological basis of transcranial motor cortex stimulation in conscious humans. *Clin Neurophysiol* 115: 255–266, 2004.

Ellaway PH, **Davey NJ**, **Maskill DW**, **Rawlinson SR**, **Lewis HS**, **Anissimova NP**. Variability in the amplitude of skeletal muscle responses to magnetic stimulation of the motor cortex in man. *Electroencephalography and Clinical Neurophysiology - Electromyography and Motor Control* , 1998. doi:10.1016/S0924-980X(98)00007-1.

Flanders M, **Pellegrini JJ**, **Geisler SD**. Basic features of phasic activation for reaching in
vertical planes. *Exp Brain Res* 110: 67–79, 1996.

Ghez C, **Scheidt R**, **Heijink H**. Different learned coordinate frames for planning trajectories and
final positions in reaching. *J Neurophysiol* 98: 3614–3626, 2007.

Gribble PL, **Ostry DJ**. Compensation for Interaction Torques During Single- and Multijoint Limb
Movement. *Journal of Neurophysiology* 82: 2310–2326, 1999.

Gritsenko V, **Kalaska JF**, **Cisek P**. Descending Corticospinal Control of Intersegmental
Dynamics. *Journal of Neuroscience* 31: 11968–11979, 2011.

Groppa S, **Schlaak BH**, **Münchau A**, **Werner-Petroll N**, **Dünnweber J**, **Bäumer T**, **van**
Nuenen BFL, **Siebner HR**. The human dorsal premotor cortex facilitates the excitability of
ipsilateral primary motor cortex via a short latency cortico-cortical route. *Human Brain Mapping*
33: 419–430, 2012.

Hardesty RL, **Boots MT**, **Yakovenko S**, **Gritsenko V**. Computational evidence for nonlinear
feedforward modulation of fusimotor drive to antagonistic co-contracting muscles. *Sci Rep* 10:
10625, 2020.

Jung NH, **Delvendahl I**, **Kuhnke NG**, **Hauschke D**, **Stolle S**, **Mall V**. Navigated transcranial
magnetic stimulation does not decrease the variability of motor-evoked potentials. *Brain*
Stimulation 3: 87–94, 2010.

Kamen G. Reliability of motor-evoked potentials during resting and active contraction
conditions. *Med Sci Sports Exerc* 36: 1574–1579, 2004.

Kaneko K, **Kawai S**, **Fuchigami Y**, **Morita H**, **Ofuji A**. The effect of current direction induced
by transcranial magnetic stimulation on the corticospinal excitability in human brain.
Electroencephalography and clinical neurophysiology supplement 101: 478–482, 1996.

Keizer K, **Kuypers HGJM**. Distribution of corticospinal neurons with collaterals to the lower
brain stem reticular formation in monkey (Macaca fascicularis). *Exp Brain Res* 74: 311–318,
1989.

Kiers L, **Cros D**, **Chiappa KH**, **Fang J**. Variability of motor potentials evoked by transcranial
magnetic stimulation. *Electroencephalography and clinical Neurophysiology* 89: 415–423, 1993.

Lametti DR, **Houle G**, **Ostry DJ**. Control of Movement Variability and the Regulation of Limb
Impedance. *Journal of Neurophysiology* 98: 3516–3524, 2007.

Di Lazzaro V, **Oliviero A**, **Profice P**, **Saturno E**, **Pilato F**, **Insola A**, **Mazzone P**, **Tonali P**,
Rothwell JC. Comparison of descending volleys evoked by transcranial magnetic and electric
stimulation in conscious humans. *Electroencephalography and clinical neurophysiology*
supplement 109: 397–401, 1998.

Lemon R, **Johansson R**, **Westling G**. Corticospinal control during reach, grasp, and precision
lift in man. *J Neurosci* 15: 6145–6156, 1995.

Lemon RN, **Johansson RS**, **Westling G**. Modulation of corticospinal influence over hand
muscles during gripping tasks in man and monkey. 74: 12, 1996.

Lewis GN, **Signal N**, **Taylor D**. Reliability of lower limb motor evoked potentials in stroke and
healthy populations: How many responses are needed? *Clinical Neurophysiology* 125: 748–
754, 2014.

MacKinnon CD, **Rothwell JC**. Time-varying changes in corticospinal excitability accompanying
the triphasic EMG pattern in humans. *The Journal of Physiology* 528: 633–645, 2000.

Massion J. Movement, posture and equilibrium: Interaction and coordination. *Progress in
Neurobiology* 38: 35–56, 1992.

Nakamura H, **Kitagawa H**, **Kawaguchi Y**, **Tsuji H**. Direct and indirect activation of human
corticospinal neurons by transcranial magnetic and electrical stimulation. *Neuroscience Letters*
210: 45–48, 1996.

Olesh E V, **Pollard BS**, **Gritsenko V**. Gravitational and Dynamic Components of Muscle
Torque Underlie Tonic and Phasic Muscle Activity during Goal-Directed Reaching. *Frontiers in
human neuroscience* 11: 1–12, 2017.

Pascual-Leone A, **Grafman J**, **Cohen LG**, **Roth BJ**, **Hallett M**. Transcranial magnetic
stimulation. A new tool for the study of higher cognitive functions in humans. In: *Handbook of
Neuropsychology*, edited by Grafman, Boller. 1997, p. 267–290.

Pascual-Leone A, **Nguyet D**, **Cohen LG**, **Brasil-Neto JP**, **Cammarota A**, **Hallett M**.
Modulation of muscle responses evoked by transcranial magnetic stimulation during the
acquisition of new fine motor skills. *Journal of Neurophysiology* 74: 1037–1045, 1995.

Pascual-Leone A, **Tormos JM**, **Keenan J**, **Tarazona F**, **Cañete C**, **Catalá MD**. Study and
Modulation of Human Cortical Excitability With Transcranial Magnetic Stimulation: *Journal of
Clinical Neurophysiology* 15: 333–343, 1998.

Pascual-Leone A, **Walsh V**. Transcranial Magnetic Stimulation. In: *Brain Mapping The
Methods*. Elsevier Science (USA), 2002.

Penfield W, **Rasmussen T**. *The cerebral cortex of man; a clinical study of localization of
function*. Oxford, England: Macmillan, 1950.

Perez MA, **Rothwell JC**. Distinct Influence of Hand Posture on Cortical Activity during Human
Grasping. *J Neurosci* 35: 4882–4889, 2015.

Reis J, **Swayne OB**, **Vandermeeren Y**, **Camus M**, **Dimyan MA**, **Harris-Love M**, **Perez MA**,
Ragert P, **Rothwell JC**, **Cohen LG**. Contribution of transcranial magnetic stimulation to the
understanding of cortical mechanisms involved in motor control: TMS and motor control. *The
Journal of Physiology* 586: 325–351, 2008.

Riddle CN, **Edgley SA**, **Baker SN**. Direct and Indirect Connections with Upper Limb
Motoneurons from the Primate Reticulospinal Tract. *J Neurosci* 29: 4993–4999, 2009.

Robertson G, **Caldwell G**, **Hamill J**, **Kamen G**, **Whittlesey S**. *Research Methods in Biomechanics, 2E*. Human Kinetics, 2013.

Roth BJ, **Saypol JM**, **Hallett M**, **Cohen LG**. A theoretical calculation of the electric field induced in the cortex during magnetic stimulation. *Electroencephalography and Clinical Neurophysiology/Evoked Potentials Section* 81: 47–56, 1991.

Rothwell JC. Techniques and mechanisms of action of transcranial stimulation of the human motor cortex. *J Neurosci Methods* 74: 113–122, 1997.

Rothwell JC, **Thompson PD**, **Day BL**, **Boyd S**, **Marsden CD**. Stimulation of the human motor cortex through the scalp. *Experimental Physiology* 76: 159–200, 1991.

Schepens B, **Drew T**. Independent and Convergent Signals From the Pontomedullary Reticular Formation Contribute to the Control of Posture and Movement During Reaching in the Cat. *Journal of Neurophysiology* 92: 2217–2238, 2004.

Schepens B, **Drew T**. Descending Signals From the Pontomedullary Reticular Formation Are Bilateral, Asymmetric, and Gated During Reaching Movements in the Cat. *Journal of Neurophysiology* 96: 2229–2252, 2006.

Talkington WJ, **Pollard BS**, **Olesh E V**, **Gritsenko V**. Multifunctional Setup for Studying Human Motor Control Using Transcranial Magnetic Stimulation, Electromyography, Motion Capture, and Virtual Reality. *Journal of visualized experiments : JoVE* e52906, 2015.

Taylor J. Stimulation at the cervicomedullary junction in human subjects. *Journal of Electromyography and Kinesiology* 16: 215–223, 2006.

Thickbroom GW, **Byrnes ML**, **Mastaglia FL**. A model of the effect of MEP amplitude variation on the accuracy of TMS mapping. *Clinical Neurophysiology* , 1999. doi:10.1016/S1388-2457(98)00080-7.

Todorov E, **Jordan MI**. Optimal feedback control as a theory of motor coordination. *Nat Neurosci* 5: 1226–1235, 2002.

Tofts PS. The distribution of induced currents in magnetic stimulation of the nervous system. *Phys Med Biol* 35: 1119–1128, 1990.

Traversa R, **Cicinelli P**, **Bassi A**, **Rossini PM**, **Bernardi G**. Mapping of motor cortical reorganization after stroke. A brain stimulation study with focal magnetic pulses. *Stroke* 28: 110–117, 1997.

Turton A, **Lemon RN**. The contribution of fast corticospinal input to the voluntary activation of proximal muscles in normal subjects and in stroke patients. *Experimental Brain Research* 129: 0559–0572, 1999.

Uy J, **Ridding MC**, **Miles TS**. Stability of Maps of Human Motor Cortex Made with Transcranial Magnetic Stimulation. *Brain Topography* 14: 5, 2002.

Wassermann EM. Variation in the response to transcranial magnetic brain stimulation in the general population. *Clinical Neurophysiology* 113: 1165–1171, 2002.

Winter DA. *Biomechanics and Motor Control of Human Movement*. 4th ed. Hoboken, New
Jersey: John Wiley & Sons, 2009.

Yadav V, **Sainburg RL**. Motor lateralization is characterized by a serial hybrid control scheme.
Neuroscience 196: 153–167, 2011.

Yakovenko S, **Drew T**. A Motor Cortical Contribution to the Anticipatory Postural Adjustments
That Precede Reaching in the Cat. *Journal of Neurophysiology* 102: 853–874, 2009.

Zajac FE. Muscle and tendon: properties, models, scaling, and application to biomechanics and
motor control. *Critical Reviews in Biomedical Engineering* 17: 359–411, 1989.

Zanette G, **Bonato C**, **Polo A**, **Tinazzi M**, **Manganotti P**, **Fiaschi A**. Long-lasting depression of
motor-evoked potentials to transcranial magnetic stimulation following exercise. *Exp Brain Res*
107: 80–86, 1995.

Figure 1

Figure 2

Figure 3

Figure 4

A. CV of MEPs

B. Examples of static MEPs vs. EMG in BicL

C. Shared variance between MEPs and EMG

Figure 5

Figure 6

Figure 7

Figure 8