Extended-spectrum beta-lactamase (ESBL)-producing and non-ESBL-

producing *Escherichia coli* isolates causing bacteremia in the

Netherlands (2014 – 2016) differ in clonal distribution, antimicrobial

resistance gene and virulence gene content

Denise van Hout*[1], Tess D. Verschuuren[1], Patricia C.J. Bruijning-Verhagen[1,2], Thijs Bosch[2],

Anita C. Schürch[3], Rob J.L. Willems[3], Marc J.M. Bonten[1-3], Jan A.J.W. Kluytmans[1,4]

Author affiliations:

[1] Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht,

University Utrecht, Utrecht, The Netherlands

[2] Center for Infectious Disease Control, National Institute for Public Health and the Environment,

Bilthoven, The Netherlands

[3] Department of Medical Microbiology, University Medical Center Utrecht, Utrecht, University

Utrecht, The Netherlands

[4] Microvida Laboratory for Medical Microbiology and Department of Infection Control, Amphia

Hospital, Breda, The Netherlands

* Corresponding author:

Denise van Hout

UMC Utrecht; Julius Center

Huispost nr. STR 6.131

P.O. Box 85500

3508 GA Utrecht

+31887567407

D.vanHout-3@umcutrecht.nl

26
27   Word count manuscript: 3325
28

ABSTRACT

Background: Knowledge on the molecular epidemiology of *Escherichia coli* causing *E. coli*
bacteremia (ECB) in the Netherlands is mostly based on extended-spectrum beta-lactamase-
producing *E. coli* (ESBL-Ec). We determined differences in clonality and resistance and
virulence gene (VG) content between non-ESBL-producing *E. coli* (non-ESBL-Ec) and ESBL-Ec
blood isolates with different epidemiological characteristics.

Materials/methods: A random selection of non-ESBL-Ec isolates as well as all available ESBL-
Ec blood isolates was obtained from two Dutch hospitals between 2014 and 2016. Whole
genome sequencing was performed to infer sequence types (STs), serotypes, acquired
antibiotic resistance genes and VG scores, based on presence of 49 predefined putative
pathogenic VG.

Results: ST73 was most prevalent among the 212 non-ESBL-Ec (N=26, 12.3%) and ST131
among the 69 ESBL-Ec (N=30, 43.5%). Prevalence of ST131 among non-ESBL-Ec was 10.4%
(N=22, *P* value < 0.001 compared to ESBL-Ec). O25:H4 was the most common serotype in both
non-ESBL-Ec and ESBL-Ec. Median acquired resistance gene counts were 1 (IQR 1 – 6) and 7
(IQR 4 – 9) for non-ESBL-Ec and ESBL-Ec, respectively (*P* value < 0.001). Among non-ESBL-
Ec, acquired resistance gene count was highest among blood isolates from a primary gastro-
intestinal focus (median 4, IQR 1 – 8). Median VG scores were 13 (IQR 9 – 20) and 12 (IQR 8 –
14) for non-ESBL-Ec and ESBL-Ec, respectively (*P* value = 0.002). VG scores among non-
ESBL-Ec from a primary urinary focus (median 15, IQR 11 – 21) were higher compared to non-
ESBL-Ec from a primary gastro-intestinal (median 10, IQR 6 – 13) or hepatic-biliary focus
(median 11, IQR 5 – 18) (*P* values = 0.007 and 0.036, respectively). VG content varied between
different *E. coli* STs.

Conclusions: Non-ESBL-Ec and ESBL-Ec blood isolates from two Dutch hospitals differed in
clonal distribution, resistance gene and VG content. Also, resistance gene and VG content
differed between non-ESBL-Ec from different primary foci of ECB.

INTRODUCTION

Despite advances in medical healthcare and in contrast to the decline in other infectious diseases, the annual incidence of Gram-negative bacteremia in Europe is increasing [1–4]. *Escherichia coli* is the leading causative pathogen in Gram-negative bacteremia and is associated with 30-day mortality up to 18% [1,4–6]. Antibiotic treatment options of *E. coli* bacteremia (ECB) are getting compromised by the pandemic presence of extended-spectrum beta-lactamases (ESBLs) [1–4]; enzymes that confer resistance to antibiotics commonly used for ECB treatment such as third-generation cephalosporins. Genes encoding ESBLs are encoded on mobile genetic elements and can be exchanged between strains by horizontal gene transfer. In some European countries, the incidence of ECB with antibiotic-resistant strains seems to increase faster than ECB caused by susceptible strains [1–4]. The individual patient and financial burden is increased for ECB episodes that are caused by resistant *E. coli*. Yet, ECB due to susceptible strains is far more common and therefore determines the major part of the ECB disease burden in the population [1–4]. The majority of ECBs is of community onset and is preceded by an infection in the urinary tract, but other sources, such as the hepatic-biliary tract, also comprise important primary foci [4,7]. These clinical characteristics of ECB episodes are important because they can indicate different target populations for prevention. More insight in the molecular epidemiology of ESBL-negative as well as ESBL-positive ECB with different clinical characteristics is needed to help identify key targets for the development of future preventive strategies such as *E. coli* vaccines, which are currently being developed [8]. Up to now, the molecular epidemiology of ECB in the Netherlands has been mainly described in single-center studies [9] or among antimicrobial resistant isolates only [10]. Dutch studies combining patient characteristics with high-resolution genetic data of *E. coli* isolates are limited, specifically for ECB, with its potential severe clinical consequences.

In this study, we aimed to analyze the current population structure of ECB in the

Netherlands, with special attention to differences in antimicrobial resistance and virulence gene

content and clonal and serotype distribution between isolates with different clinical

epidemiological characteristics and between non-ESBL-producing *E. coli* (non-ESBL-Ec) and

ESBL-producing *E. coli* (ESBL-Ec) blood isolates.

METHODS

Study design

Details of the study design, clinical epidemiological data collection and laboratory methods (i.e.

phenotypic ESBL detection) are described elsewhere [11]. In short, patients with ECB were

retrospectively identified from medical microbiological records in the University Medical Center

Utrecht (UMCU), a 1,042-bed tertiary care center and the Amphia Hospital in Breda, an 837-bed

teaching hospital. In each hospital, we selected a random sample of 40 isolates of unique

patients per year for the years 2014, 2015 and 2016, comprising ~24% of all first bacteremic *E.*

coli isolates in a year. In addition to this random sample, all ESBL-Ec blood isolates from 2014 –

2016 were selected from the two hospitals. Whole genome sequencing (WGS) was performed

by The Netherlands National Institute for Public Health and the Environment (RIVM) using the

Illumina HiSeq 2500 (BaseClear, Leiden, the Netherlands). All generated raw reads were

submitted to the European Nucleotide Archive (ENA) of the European Bioinformatics Institute

(EBI) under the study accession number PRJEB35000. De novo assembly was performed using

SPAdes genome assembler v.3.6.2 and the quality of assembles was assessed using QUAST

[12]. Only genomes with an estimated genome size between 3 MB and 6 MB and number of

contigs not exceeding 1,000 were included in further analyses. Baseline clinical epidemiological

characteristics were compared between the non-ESBL-Ec and ESBL-Ec ECB episodes. ESBL-

production was defined as confirmed phenotypic ESBL-positivity, unless described otherwise

[11]. Baseline characteristics were compared by the Fisher's Exact or Pearson χ^2 test for

categorical variables and by Mann-Whitney U test for continuous variables when applicable. A

two-tailed *P* value <0.05 was considered statistically significant.

This study does not fall under the scope of the Medical Research Involving Human

Subjects Act. The Medical Research Ethics Committee of the UMCU has therefore waived the

need for official approval by the Ethics Committee (IRB number 18/056) and informed consent

was not obtained. All statistical analyses were performed with Statistical Package for Social

Sciences V.25.0 (SPSS, Chicago, Illinois, USA) and R Version 3.4.1. Boxplots were made with

R packages *ggplot2* and *ggpubr* and bar charts were made with Graphpad Prism Version 8.0.1.

Multi-locus sequence types (MLST)

Multi-locus   sequence   types   (STs)   were   determined   using   mlst2.0

(https://github.com/tseemann/mlst) by scanning contig files against the *E. coli* PubMLST typing

scheme (updated May 12th, 2018). Clonal (i.e. ST) distribution was presented stratified for non-

ESBL-Ec and ESBL-Ec isolates and by epidemiological subgroup (i.e. community versus

hospital onset; different primary foci of ECB). Genotype (ST) diversity was analysed by

Simpson's diversity index [13].

Serotyping

We assigned serotypes by using the web-tool SerotypeFinder 2.0 from the Center for Genomic

Epidemiology   at   the   Danish   Technical   University,   Lyngby,   Denmark

(https://cge.cbs.dtu.dk/services/SerotypeFinder) [14]. Simpson's index for serotype diversity was

calculated for non-ESBL-Ec and ESBL-Ec isolates. Serotype distribution among non-ESBL-Ec

and ESBL-Ec was compared to two current *E. coli* vaccine candidates [8,15], excluding those

isolates in which no definitive serotype could be defined, and the occurrence of serotypes was

described by primary focus of ECB.

Antimicrobial resistance genes

Abricate (https://github.com/tseemann/abricate) version 0.8.13 was used for mass screening of contigs for antimicrobial resistance genes using the ResFinder 3.1.0 database (acquired resistance genes only), date of download 24 January 2019 [16]. The thresholds for coverage length and sequence identity were 80% and 95%, respectively. A resistance gene count was made per isolate, which was defined as the total number of identified acquired resistance genes. In case of double detection of identical resistance genes within a single isolate, they were only counted once. The resistance gene counts were compared between non-ESBL-Ec and ESBL-Ec with the non-parametric Wilcoxon rank sum test (for this comparison only, the scores of the ESBL-Ec isolates were corrected for presence of the ESBL gene). Resistance gene scores were then analysed for non-ESBL-Ec and ESBL-Ec separately and were compared between isolates with different epidemiological characteristics and different STs using Kruskal-Wallis one-way ANOVA. In case of an overall ANOVA P value <0.05, post-hoc pairwise comparisons were made and the Holm-Bonferroni P value correction was applied to account for multiple testing. For pairwise comparisons, the non-parametric Wilcoxon rank sum test was used.

Virulence genes

The presence of putative virulence factor genes (VG) was identified using abricate version 0.8.13 for BLAST against the VFDB database (http://www.mgc.ac.cn/VFs), date of download 8 February 2019, with minimal coverage length and sequence identity 80% and 95% [17]. We searched for 49 putative VG that were previously described as extra-intestinal pathogenic *E. coli* (ExPEC)-associated VG [18–22]. If any of the predefined VG were not included in VFDB, BLAST against the ecoli_VF_collection database was performed (date 8 February 2019), a repository that contains known VG from VFDB supplemented with additional *E. coli* VG that have been reported in literature [23]. The *kpsM*, *afa/dra* and *sfa/foc* operons were considered

present if any of the corresponding genes or allelic variants were identified. A virulence score

was made per isolate and was defined as the total number of pre specified VG, adjusted for

multiple detection of the *afa/dra* (Afa/Dr adhesins), *pap* (P fimbrial adhesins), *sfa/foc* (S and F1C

fimbrial adhesins) and *kpsM* (group 2 and III capsule) operons, as described previously [20]. If a

VG was detected multiple times within a single isolate (i.e. with different quality measures), it

was only counted once. These virulence scores were then compared between isolates with

different epidemiological characteristics and between different STs using Kruskal-Wallis one-

way ANOVA. In case of an overall ANOVA *P* value <0.05, post-hoc pairwise comparisons were

made with the non-parametric Wilcoxon rank sum test and the Holm-Bonferroni *P* value

correction was applied to account for multiple testing.

RESULTS

Patient characteristics

The isolate collection consisted of 212 phenotypic non-ESBL-Ec and 69 ESBL-Ec blood isolates

(Fig. 1). Distribution of age, sex, onset of infection and primary foci were comparable between

non-ESBL-Ec and ESBL-Ec bacteremia episodes (Table 1). As compared to non-ESBL-Ec,

ECB episodes with ESBL-Ec were less often of community onset (63.8% versus 81.1%, *P* value

= 0.003). Crude 30-day and 1-year mortality were higher in ECB episodes caused by ESBL-Ec

(27.5% and 50.7%, respectively) compared to ECB episodes caused by non-ESBL-Ec (11.3%

and 29.2%, respectively) (both *P* values = 0.001).

Clonal distribution

Among non-ESBL-Ec, ST73 was the most frequently observed ST (N = 26, 12.3%), followed by

ST131 (N = 22, 10.4%). Isolates of ST73, 95, 127, 141, 80 and 1193 were solely identified

among non-ESBL-Ec (Fig. 2). ST131 was dominant among ESBL-Ec (N = 30, 43.5%) and

prevalence was higher than among non-ESBL-Ec (*P* value < 0.001). Simpson's index for clonal

diversity was 95.6% (95% CI 94.4% – 96.8%) and 80.6% (95% CI 70.9% – 90.4%) for non-

ESBL-Ec and ESBL-Ec, respectively. The occurrence of different STs did not differ between

nosocomial and community onset ECB (S1 Appendix). ST131 was the dominant ST among

ESBL-positive ECB episodes with a primary urinary (63%) and gastro-intestinal focus (57%),

which was higher as compared to other primary foci of ESBL-positive ECB (i.e. 21% among

primary hepatic-biliary focus, see S1 Appendix).

Figure 1. Flowchart of selection of *E. coli* blood isolates

185
186

187   **Table 1.** Baseline epidemiological characteristics of *E. coli* bacteremia episodes

	Non-ESBL-Ec[a] N = 212		ESBL-Ec[a] N = 69		*P* value[b]
Median age, years (IQR)	69	(59 – 77)	69	(56 – 76)	0.802
Female sex (%)	102	(48.1)	32	(46.4)	0.802
Community onset (%)	172	(81.1)	44	(63.8)	*0.003*
Primary focus of ECB (%)					
Urinary tract	103	(48.6)	30	(43.5)	0.785
Hepatic-biliary	46	(21.7)	14	(20.3)	
Gastro-intestinal	23	(10.8)	7	(10.1)	
Other	10	(4.7)	5	(7.2)	
Unknown	30	(14.2)	13	(18.8)	
Urinary catheter (%)	69	(32.5)	28	(40.6)	0.223
Ward (%)					
Non-ICU	182	(85.8)	58	(84.1)	0.714
ICU	30	(14.2)	11	(15.9)	
Mortality (%)					
30-day	24	(11.3)	19	(27.5)	*0.001*
1-year	62	(29.2)	35	(50.7)	*0.001*

ECB, *E. coli* bacteremia; ESBL, extended-spectrum beta-lactamase; ESBL-Ec, ESBL-producing *E. coli*; ICU, intensive care unit; IQR, interquartile range; non-ESBL-Ec, non-ESBL-producing *E. coli*
[a]ESBL-positivity based on phenotype.
[b]*P* value of comparison between non-ESBL-Ec versus ESBL-Ec, calculated with Pearson's χ^2, Fisher's exact, or Mann-Whitney U test when applicable. *P* values in italic represent *P* values <0.05.

188   **Figure 2.** ST distribution among non-ESBL-Ec versus ESBL-Ec[a] in order of frequency[b]

189
190   ESBL-Ec, ESBL-producing *E. coli*; non-ESBL-Ec, non-ESBL-producing *E. coli*; ST, sequence type
191   [a]ESBL-positivity based on phenotypic ESBL production.
192   [b]Missing STs and STs that occurred ≤3 times are grouped in "Other". STs that only occurred once are grouped in
193   "Singletons". The height of each individual bars represents the proportion of the ST within the group of non-ESBL-Ec
194   and ESBL-Ec, respectively. The numbers represent the absolute numbers of occurrence.

Serotypes

The most common serotype O25:H4 was identified in 19 (9.0%) non-ESBL-Ec and 24 (34.8%) ESBL-Ec isolates, which largely reflected the prevalence of ST131 in each group (Table 2). Multiple serotypes only occurred among non-ESBL-Ec, such as O6:H1 and O6:H31. ST73 was most often of serotype O6:H1 (16 / 26, 61.5%). Simpson's index for serotype diversity was 96.7% (95% CI 95.8% – 97.6%) and 83.8% (95% CI 76.9% – 90.6%) for non-ESBL-Ec and ESBL-Ec, respectively. Non-ESBL-Ec and ESBL-Ec isolates from ECB episodes with a primary focus in the urinary tract were most often of O serotype O6 (15 / 103, 14.6%) and O25 (17 / 30, 56.7%), respectively (S2 Appendix). For ECB episodes with a primary focus in the hepatic-biliary tract, O25 was the most prevalent O serotype among non-ESBL-Ec (7 / 46, 15.2%) and O8 (4 / 14, 28.6%) among ESBL-Ec isolates (S2 Appendix).

53 (25.0%) non-ESBL-Ec and 25 (36.2%) ESBL-Ec isolates belonged to either O1, O2, O6 or O25, the serotypes of the 4-valent *E. coli* vaccine that has reached phase 2 development stage [8,24], whereas the majority of non-ESBL-Ec (N = 113; 53.3%) and ESBL-Ec isolates (N = 35; 50.7%) belonged to one of the O serotypes of the new 10-valent conjugant *E. coli* vaccine (ExPEC-10V) that is currently in development [15].

Antimicrobial resistance genes

In total, 69 unique acquired resistance genes were identified (S3 Appendix). ESBL-genes were detected in 65 (94.2%) of 69 *E. coli* blood isolates with phenotypic ESBL production. $bla_{CTX-M-15}$ was the most prevalent ESBL gene (N = 28, 43.1%), followed by $bla_{CTX-M-9}$ (N = 14, 21.5%) and $bla_{CTX-M-27}$ (N = 9, 13.8%). Assemblies of the phenotypic ESBL-Ec isolates in which no ESBL-gene was identified with the used bioinformatics pipeline (N = 4) were individually uploaded on the DTU Resfinder 3.1.0 website (date 11 March 2019, thresholds for coverage length 80% and sequence identity 95%); these isolates remained genotypically ESBL-negative. One of these isolates was positive for bla_{CMY-2} (AmpC gene).

The median acquired resistance gene count for non-ESBL-Ec versus ESBL-Ec was 1

(IQR 1 – 6) versus 7 (IQR 4 – 9) (*P* value < 0.001). Among non-ESBL-Ec, acquired resistance

gene counts were not different between community and hospital-onset ECB episodes (S3

Appendix). Among non-ESBL-Ec, there were statistically significant differences in resistance

gene count for different primary foci of ECB, but absolute differences were small: median

resistance gene count from ECB with a primary hepatic-biliary focus was 1 (IQR 1 – 1), whereas

for a primary urinary focus this was 2 (IQR 1 – 6) (*P* value ≤0.001), for a primary gastro-

intestinal focus this was 4 (IQR 1 – 8) (*P* value ≤0.01 for comparison to hepatic-biliary focus)

and for an unknown primary focus this was 2 (IQR 1 – 7) (*P* value ≤0.0001 for comparison to

hepatic-biliary focus) (S3 Appendix). Among ESBL-Ec isolates, there were no statistical

significant differences in acquired resistance gene counts between community and hospital-

onset ECB or different primary foci of ECB (S3 Appendix). There were no statistically significant

differences observed in resistance gene count among non-ESBL-Ec or ESBL-Ec isolates of

different clonal background (Fig. 3 and S3 Appendix).

 Table 2. Serotype distribution among *E. coli* blood isolates, stratified for ESBL-positivity

	Non-ESBL-Ec N = 212 (%)	ESBL-Ec[a] N = 69 (%)
O25:H4 (%)	19 (9.0)	24 (34.8)
O6:H1 (%)	16 (7.5)	-
O2/O50:H6 (%)	10 (4.7)	-
O6:H31 (%)	9 (4.2)	-
O15:H18 (%)	7 (3.3)	2 (2.9)
O17/O44/O77:H18 (%)	8 (3.8)	-
O4:H5 (%)	7 (3.3)	1 (1.4)
O75:H5 (%)	8 (3.8)	-
O8:H9	5 (2.4)	2 (2.9)
O16:H5 (%)	3 (1.4)	3 (4.3)
O86:H18	1 (0.5)	4 (5.8)
O4:H1 (%)	5 (2.4)	-
O1:H7	4 (1.9)	-
O117:H4	4 (1.9)	-
O2/O50:H1	4 (1.9)	-
O23:H16	2 (0.9)	2 (2.9)
O25:H1	4 (1.9)	-
O18/O18ac:H7	3 (1.4)	-
O2/O50:H7	3 (1.4)	-
O45:H7	3 (1.4)	-
O75:H7	3 (1.4)	-
O8:H17	3 (1.4)	-
O9:H17	-	2 (2.9)
O9/O104:H9	-	2 (2.9)
O13/O135:H4	2 (0.9)	-
O18:H1	2 (0.9)	-
O18:H5	2 (0.9)	-
O22:H1	2 (0.9)	-
O24:H4	2 (0.9)	-
O8:H10	2 (0.9)	-
O8:H25	2 (0.9)	-
O8:H30	2 (0.9)	-
Singletons	45 (21.2)	13 (18.8)
Unknown	20 (9.4)	14 (20.3)

[a]ESBL-positivity based on phenotypic ESBL production.
ESBL, extended-spectrum beta-lactamase; ESBL-Ec, ESBL-producing *E. coli*, non-ESBL-Ec, non-ESBL-producing *E. coli*

ᵃESBL-positivity based on phenotypic ESBL production.
Boxplots display median resistance gene count and inter quartile range (IQR); every dot represents a single isolate.
Only STs that occurred >5% within non-ESBL-Ec or ESBL-Ec were grouped into main groups, the rest was
categorized as "Other". Results of the pairwise comparisons between STs can be found in S3 Appendix.

Virulence genes

Of the 49 predefined ExPEC-associated VG, 44 (89.8%) were detected in at least one *E. coli*

blood isolate (S4 Appendix). The median VG score was 13 (IQR 9 – 20) for non-ESBL-Ec and

12 (IQR 8 – 14) for ESBL-Ec blood isolates (*P* value = 0.002). In one non-ESBL-Ec isolate no

predefined ExPEC-associated VG was detected, while a maximum VG score of 25 was found in

two non-ESBL-Ec isolates.

For non-ESBL-Ec and ESBL-Ec isolates, there was no significant difference in the VG

score between isolates that caused community or hospital onset ECB (S4 Appendix). Non-

ESBL-Ec isolates that caused ECB with a primary gastro-intestinal focus and hepatic-biliary

focus had lower VG scores (median 10, IQR 6 – 13 and median 11, IQR 5 – 18, respectively) as

compared to isolates with a primary focus in the urinary tract (median 15, IQR 11 – 21) (*P* value

= 0.007 and *P* value = 0.036, respectively) (see S4 Appendix). Among non-ESBL-Ec and

ESBL-Ec, there were no statistical significant differences in VG scores between isolates of

patients without versus with a urinary catheter, between patients alive or deceased after 30

days or between patients admitted to the intensive care unit (ICU) versus a non-ICU ward (S4

Appendix).

There was heterogeneity in VG scores between non-ESBL-Ec of different STs, this was

less pronounced for ESBL-Ec isolates (Fig. 4 and S4 Appendix). ESBL-negative ST38 had the

lowest average VG score (median 7, IQR 6 – 7) and ESBL-positive ST12 had the highest VG

score (median 23, IQR 23 – 23). Median VG score of both ESBL-negative and ESBL-positive

ST131 isolates was 13 (IQR 12 – 15). All pairwise comparisons between ESBL-negative STs

yielded Holm-Bonferroni adjusted *P* values < 0.05, except for the comparison ST12 versus

ST73 and all pairwise comparisons that included ST38.

Figure 4. ExPEC-associated VG score in different STs, stratified for ESBL-positivity[a]

267
[a]ESBL-positivity based on phenotypic ESBL production.
Boxplots display median VG score and inter quartile range (IQR); every dot represents a single isolate. Only STs that
occurred >5% within non-ESBL-Ec or ESBL-Ec were grouped into main groups, the rest was categorized as "Other".
Results of pairwise comparisons between STs can be found in S4 Appendix.

DISCUSSION

In this study, we found that ESBL-producing *E. coli* blood isolates were different from non-

ESBL-producing *E. coli* causing bacteraemia in terms of clonal distribution, serotype distribution,

antimicrobial resistance gene count and VG scores.

In line with previous research, the clonal distribution among ESBL-Ec blood isolates was

less diverse as compared to non-ESBL-Ec [25–27]. This was mainly caused by the

predominance of ST131 within ESBL-Ec, as has been described before [28,29]. In contrast,

ST73, a ST that is known for its susceptibility to antibiotics [28], was only identified among non-

ESBL-Ec blood isolates. The association between ESBL phenotype and STs in *E. coli*, which is

repeatedly found, implies that the genetic make-up of strains contributes to the ability to acquire

and subsequently maintain plasmids carrying ESBL genes. Indeed, a recent large-scale study

that compared the pan-genomes of invasive *E. coli* isolates, including ST131 and ST73,

suggested that due to ongoing adaptation to long term human intestinal colonisation and

consequent evolutionary gene selection, ST131 might have become able to reduce the fitness

costs of long term plasmid maintenance [30,31]. It has been hypothesized that this is also true

for other *E. coli* lineages that are associated with multi-drug resistance. Reducing the fitness

costs of replicating plasmids encoding multi-drug resistance will result in having competitive

advantage over other intestinal strains [32].

We hypothesized that the clonal distribution and resistance gene and VG content would

differ between ECB episodes of community and hospital onset and between different primary

foci, as a result of adaptive evolution of intestinal *E. coli*. We observed some statistical

significant differences in resistance gene count and VG scores among non-ESBL-Ec from

different primary foci of ECB, such as higher VG scores of blood isolates from a primary urinary

focus as compared to isolates from a primary focus in the gastro-intestinal or hepatic-biliary

tract. However, absolute differences in median gene counts were small and the clinical

significance remains unclear. In the current study, we found that differences in molecular

content mostly depended on phenotypic ESBL-production and STs. This confirms the findings

from a recent study that was performed in Scotland [33]. In that study, there were combinations

of VGs as well as a particular accessory gene composition that differentiated between STs

rather than between epidemiological factors. The association between ST69 and community

onset ECB, as found in the Scottish study, was not identified in the current study. Other

differences were the large proportion of *E. coli* isolates from ECB episodes that were deemed

hospital-acquired (62%) as compared to our study (18.4% for ESBL-negative and 36.2% for

ESBL-positive ECB) and in that study, analyses were not stratified for ESBL-positivity.

Interestingly, in our study, isolates that belonged to ST73 had low resistance gene

content but relatively high VG scores as compared to other STs. Furthermore, the average VG

score among non-ESBL-Ec was slightly higher in comparison to ESBL-Ec blood isolates, which

demonstrates that ESBL-positivity in *E. coli* is not necessarily related to an increased VG

content. These findings do not support the theory that increased virulence of resistant strains

causes the increased incidence of resistant ECB as compared to sensitive ECB. This theory has

been suggested for other pathogens, such as MRSA [1,34,35].

We identified serotype O25:H4 as the most prevalent serotype causing ESBL-negative

as well as ESBL-positive ECB in The Netherlands, followed by O6:H1. The serotype distribution

among non-ESBL-Ec was more heterogeneous as compared to ESBL-Ec, similar to the

differences in clonal diversity between these two groups [36]. A large recent European

surveillance study that included 1,110 *E. coli* blood isolates from adults between 2011 and 2017

showed that there is heterogeneity in serotype distribution among different countries, which

highlights the need for country specific data, such as provided in the current study [15]. We

showed that the coverage of the new potential 10-valent vaccine was higher as compared to the

4-valent vaccine and was actually doubled for non-ESBL-Ec bacteraemia. Findings of the

current study can be used for future studies and can help further evaluation and implementation

of *E. coli* vaccines.

Strengths of the current study are the multicenter design and combination of

epidemiological characteristics and highly discriminatory genetic data. There are also important

limitations. Firstly, *E. coli* is a heterogeneous species, of which the seven MLST genes only

constitute a small proportion of the entire gene content. Because we also only investigated a

small fraction of the genes that are commonly part of the accessory genome, such as VGs and

acquired resistance genes, we may have missed genomic differences that could have

importantly contributed to ecological specialization in the different clinically relevant primary foci.

Secondly, we selected *E. coli* isolates from a tertiary care center and teaching hospital from the

Netherlands from two different regions, which we considered to be representative of the

Netherlands. The description of strains that were identified here might not be entirely

generalizable to other countries since there could be differences between circulating *E. coli*

strains, dependent on local population characteristics and antimicrobial resistance levels.

Thirdly, many pairwise comparisons between subgroups were performed, which increases the

risk of false-positive findings (i.e. type I errors). Even though we applied a strict *P* value

correction for multiple testing, this naturally does not eliminate the risk of false-positive findings.

The analyses on resistance gene and VG content should therefore be viewed as hypothesis

generating.

In conclusion, associations between clinical characteristics of ECB episodes and

molecular content of *E. coli* isolates were limited. However, we did identify important differences

in clonality, serotypes, antimicrobial resistance genes and VG scores between non-ESBL-Ec

and ESBL-Ec blood isolates that reached beyond their phenotypic ESBL-positivity. Future

studies that aim to describe the molecular epidemiology of ECB should therefore preferably

focus on *E. coli* without preselection on ESBL-positivity, to limit the risk of inferring

characteristics of resistant *E. coli* to the *E. coli* population as a whole. Furthermore, a more

thorough understanding of the molecular epidemiology of ECB and specifically studies that

further disclose targets for surveillance or infection-prevention will help to reduce the occurrence

of this invasive infectious disease with its severe potential consequences.

FUNDING

This study was internally funded by the Netherlands National Institute for Public Health and the Environment (RIVM) and the University Medical Center Utrecht (UMCU) and the authors received no specific funding for this work.

ACKNOWLEDGMENTS

We would sincerely like to thank Kim van der Zwaluw, Carlo Verhulst and Judith Vlooswijk for their contributions in the laboratory execution of the study. Preliminary results of this paper were presented at the 29th European Congress of Clinical Microbiology & Infectious Diseases, Amsterdam, the Netherlands, 13-16 April 2019 (P1450).

AUTHOR CONTRIBUTIONS

Author	Roles
D. van Hout	Conceptualization, data curation, formal analysis, investigation, methodology, project administration, software, validation, visualization, writing – original draft preparation, writing – review & editing
T.D. Verschuuren	Conceptualization, data curation, formal analysis, investigation, methodology, project administration, software, validation, writing – review & editing
P.C.J. Bruijning-Verhagen	Conceptualization, formal analysis, methodology, supervision, validation, writing – review & editing
T. Bosch	Conceptualization, data curation, formal analysis, investigation, methodology, funding acquisition, resources, software, supervision, writing – review & editing
A.C. Schürch	Methodology, software, resources, supervision, validation, writing – review & editing
R.J.L. Willems	Methodology, software, resources, supervision, writing – review & editing
M.J.M. Bonten	Conceptualization, formal analysis, funding acquisition, resources, supervision, writing – review & editing
J.A.J.W. Kluytmans	Conceptualization, formal analysis, resources, supervision, writing – review & editing

REFERENCES

1.   de Kraker MEA, Jarlier V, Monen JCM, Heuer OE, van de Sande N, Grundmann H. The changing epidemiology of bacteraemias in Europe: trends from the European Antimicrobial Resistance Surveillance System. Clin Microbiol Infect. 2013;19: 860–868. doi:10.1111/1469-0691.12028

2.   Van Der Steen M, Leenstra T, Kluytmans JAJW, Van Der Bij AK. Trends in expanded-spectrum
cephalosporin-resistant Escherichia coli and Klebsiella pneumoniae among Dutch clinical isolates,
from 2008 to 2012. PLoS One. 2015;10: e0138088. doi:10.1371/journal.pone.0138088

3.   Schlackow I, Stoesser N, Walker AS, Crook DW, Peto TEA, Wyllie DH. Increasing incidence of
Escherichia coli bacteraemia is driven by an increase in antibiotic-resistant isolates: electronic
database study in Oxfordshire 1999-2011. J Antimicrob Chemother. 2012;67: 1514–1524.
doi:10.1093/jac/dks082

4.   Vihta K-D, Stoesser N, Llewelyn MJ, Quan TP, Davies T, Fawcett NJ, et al. Trends over time in
Escherichia coli bloodstream infections, urinary tract infections, and antibiotic susceptibilities in
Oxfordshire, UK, 1998-2016: a study of electronic health records. Lancet Infect Dis. 2018;18:
1138–1149. doi:10.1016/S1473-3099(18)30353-0

5.   Abernethy JK, Johnson AP, Guy R, Hinton N, Sheridan EA, Hope RJ. Thirty day all-cause
mortality in patients with Escherichia coli bacteraemia in England. Clin Microbiol Infect. 2015;21:
251.e1-8. doi:10.1016/j.cmi.2015.01.001

6.   Fitzpatrick JM, Biswas JS, Edgeworth JD, Islam J, Jenkins N, Judge R, et al. Gram-negative
bacteraemia; a multi-centre prospective evaluation of empiric antibiotic therapy and outcome in
English acute hospitals. Clin Microbiol Infect. 2016;22: 244–251. doi:10.1016/j.cmi.2015.10.034

7.   Abernethy J, Guy R, Sheridan EA, Hopkins S, Kiernan M, Wilcox MH, et al. Epidemiology of
Escherichia coli bacteraemia in England: results of an enhanced sentinel surveillance programme.
J Hosp Infect. 2017;95: 365–375. doi:10.1016/j.jhin.2016.12.008

8.   Huttner A, Hatz C, van den Dobbelsteen G, Abbanat D, Hornacek A, Frolich R, et al. Safety,
immunogenicity, and preliminary clinical efficacy of a vaccine against extraintestinal pathogenic
Escherichia coli in women with a history of recurrent urinary tract infection: a randomised, single-
blind, placebo-controlled phase 1b trial. Lancet Infect Dis. 2017;17: 528–537. doi:10.1016/S1473-
3099(17)30108-1

9.   Overdevest ITMA, Bergmans AMC, Verweij JJ, Vissers J, Bax N, Snelders E, et al. Prevalence of
phylogroups and O25/ST131 in susceptible and extended-spectrum β-lactamase-producing
Escherichia coli isolates, the Netherlands. Clin Microbiol Infect. 2015;21: 570.e1-e4.
doi:10.1016/j.cmi.2015.02.020

10.   Van Der Bij AK, Peirano G, Goessens WHF, Van Der Vorm ER, Van Westreenen M, Pitout JDD.
Clinical and molecular characteristics of extended-spectrum-β- lactamase-producing Escherichia
coli causing bacteremia in the Rotterdam Area, Netherlands. Antimicrob Agents Chemother.
2011;55: 3576–3578. doi:10.1128/AAC.00074-11

11.   van Hout D, Verschuuren TD., Bruijning-Verhagen PCJ., Bosch T., Reuland EA., Fluit AC., et al.
Design of the EPIGENEC Study: Assessing the EPIdemiology and GENetics of Escherichia coli in
the Netherlands. Preprints. 2019; 2019020066. doi:10.20944/preprints201902.0066.v1

12.   Gurevich A, Saveliev V, Vyahhi N, Tesler G. QUAST: quality assessment tool for genome
assemblies. Bioinformatics. 2013;29: 1072–1075. doi:10.1093/bioinformatics/btt086

13.   Simpson EH. Measurement of diversity. Nature. 1949;163.

14.   Joensen KG, Tetzschner AMM, Iguchi A, Aarestrup FM, Scheutz F. Rapid and Easy In Silico
Serotyping of Escherichia coli Isolates by Use of Whole-Genome Sequencing Data. J Clin
Microbiol. 2015;53: 2410–2426. doi:10.1128/JCM.00008-15

15.   Geurtsen J, Weerdenburg E, Davies T, Go O, Spiessens B, Geet G Van, et al. Extraintestinal
pathogenic Escherichia coli surveillance study to determine O-serotype prevalence and antibiotic
resistance in blood isolates collected in Europe, 2011–2017. ECCMID Conf #P1451.

16.   Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, et al. Identification of
415   acquired antimicrobial resistance genes. J Antimicrob Chemother. 2012;67: 2640–2644.
416   doi:10.1093/jac/dks261

417   17.   Liu B, Zheng D, Jin Q, Chen L, Yang J. VFDB 2019: a comparative pathogenomic platform with an
418   interactive web interface. Nucleic Acids Res. 2019;47: D687–D692. doi:10.1093/nar/gky1080

419   18.   Johnson JR, Stell AL. Extended Virulence Genotypes of Escherichia coli Strains from Patients with
420   Urosepsis in Relation to Phylogeny and Host Compromise. J Infect Dis. 2000;181: 261–72.
421   doi:10.1086/315217

422   19.   Johnson JR, Johnston BD, Porter S, Thuras P, Aziz M, Price LB. Accessory Traits and
423   Phylogenetic Background Predict Escherichia coli Extraintestinal Virulence Better Than Does
424   Ecological Source. J Infect Dis. 2019;219: 121–132. doi:10.1093/infdis/jiy459

425   20.   Johnson JR, Porter S, Johnston B, Kuskowski MA, Spurbeck RR, Mobley HLT, et al. Host
426   Characteristics and Bacterial Traits Predict Experimental Virulence for Escherichia coli
427   Bloodstream Isolates From Patients With Urosepsis. Open Forum Infect Dis. 2015;2: ofv083.
428   doi:10.1093/ofid/ofv083

429   21.   Dale AP, Woodford N. Extra-intestinal pathogenic Escherichia coli (ExPEC): Disease, carriage and
430   clones. J Infect. 2015;71: 615–626. doi:10.1016/j.jinf.2015.09.009

431   22.   Dale AP, Pandey AK, Hesp RJ, Belogiannis K, Laver JR, Shone CC, et al. Genomes of
432   Escherichia coli bacteraemia isolates originating from urinary tract foci contain more virulence-
433   associated genes than those from non-urinary foci and neutropaenic hosts. J Infect. 2018;77: 534–
434   543. doi:10.1016/j.jinf.2018.10.011

435   23.   Leimbach A. ecoli_VF_collection: V.0.1. Zenodo. [Internet]. 2016. Available:
436   http://dx.doi.org/10.5281/zenodo.56686.

437   24.   Frenck RW, Ervin J, Chu L, Abbanat D, Spiessens B, Go O, et al. Safety and immunogenicity of a
438   vaccine for extra-intestinal pathogenic Escherichia coli (ESTELLA): a phase 2 randomised
439   controlled trial. Lancet Infect Dis. 2019;19: 631–640. doi:10.1016/S1473-3099(18)30803-X

440   25.   Mathers AJ, Peirano G, Pitout JDD. The role of epidemic resistance plasmids and international
441   high-risk clones in the spread of multidrug-resistant Enterobacteriaceae. Clin Microbiol Rev.
442   2015;28: 565–591. doi:10.1128/CMR.00116-14

443   26.   Hertz FB, Nielsen JB, Schonning K, Littauer P, Knudsen JD, Lobner-Olesen A, et al. "Population
444   structure of drug-susceptible,-resistant and ESBL-producing Escherichia coli from community-
445   acquired urinary tract". BMC Microbiol. 2016;16: 63. doi:10.1186/s12866-016-0681-z

446   27.   Manges AR, Geum HM, Guo A, Edens TJ, Fibke CD, Pitout JDD. Global Extraintestinal
447   Pathogenic Escherichia coli (ExPEC) Lineages. Clin Microbiol Rev. 2019;32.
448   doi:10.1128/CMR.00135-18

449   28.   Kallonen T, Brodrick HJ, Harris SR, Corander J, Brown NM, Martin V, et al. Systematic
450   longitudinal survey of invasive Escherichia coli in England demonstrates a stable population
451   structure only transiently disturbed by the emergence of ST131. Genome Res. 2017;27: 1437–
452   1449. doi:10.1101/gr.216606.116

453   29.   Nicolas-Chanoine M-H, Bertrand X, Madec J-Y. Escherichia coli ST131, an intriguing clonal group.
454   Clin Microbiol Rev. 2014;27: 543–574. doi:10.1128/CMR.00125-13

455   30.   McNally A, Oren Y, Kelly D, Pascoe B, Dunn S, Sreecharan T, et al. Combined Analysis of
456   Variation in Core, Accessory and Regulatory Genome Regions Provides a Super-Resolution View
457   into the Evolution of Bacterial Populations. PLoS Genet. 2016;12: e1006280.

458   doi:10.1371/journal.pgen.1006280

459   31.   McNally A, Kallonen T, Connor C, Abudahab K, Aanensen DM, Horner C, et al. Diversification of
460   Colonization Factors in a Multidrug-Resistant Escherichia coli Lineage Evolving under Negative
461   Frequency-Dependent Selection. MBio. 2019;10: e00644-19. doi:10.1128/mBio.00644-19

462   32.   Dunn SJ, Connor C, McNally A. The evolution and transmission of multi-drug resistant Escherichia
463   coli and Klebsiella pneumoniae: the complexity of clones and plasmids. Curr Opin Microbiol.
464   2019;51: 51–56. doi:10.1016/j.mib.2019.06.004

465   33.   Goswami C, Fox S, Holden M, Connor M, Leanord A, Evans TJ. Genetic analysis of invasive
466   Escherichia coli in Scotland reveals determinants of healthcare-associated versus community-
467   acquired infections. Microb genomics. 2018;4. doi:10.1099/mgen.0.000190

468   34.   Laupland KB, Lyytikäinen O, Sgaard M, Kennedy KJ, Knudsen JD, Ostergaard C, et al. The
469   changing epidemiology of Staphylococcus aureus bloodstream infection: a multinational
470   population-based surveillance study. Clin Microbiol Infect. 2013;19: 465–471. doi:10.1111/j.1469-
471   0691.2012.03903.x

472   35.   Mostofsky E, Lipsitch M, Regev-Yochay G. Is methicillin-resistant Staphylococcus aureus
473   replacing methicillin-susceptible S. aureus? J Antimicrob Chemother. 2011;66: 2199–2214.
474   doi:10.1093/jac/dkr278

475   36.   Ciesielczuk H, Jenkins C, Chattaway M, Doumith M, Hope R, Woodford N, et al. Trends in ExPEC
476   serogroups in the UK and their significance. Eur J Clin Microbiol Infect Dis. Germany; 2016;35:
477   1661–1666. doi:10.1007/s10096-016-2707-8

480   **SUPPORTING INFORMATION**

482   **S1 Appendix**

483   **S2 Appendix**

484   **S3 Appendix**

485   **S4 Appendix**

486