Natural brown adipose expansion and remission of hyperglycemia in obese SM/J mice

Caryn Carson[1,*], Mario A Miranda[1,*], Juan F Macias-Velasco[1], Subhadra Gunawardana[2], Jing Hughes[3], Sakura Oyama[1], Marcus Kunzmann[1], Heather Schmidt[1], Jessica P Wayhart[1], Heather A Lawson[1,#]

[1]Department of Genetics, Washington University School of Medicine, 660 South Euclid Ave, Saint Louis, MO, USA
[2]Department of Cell Biology and Physiology, Washington University School of Medicine, 660 South Euclid Ave, Saint Louis, MO, USA
[3]Department of Medicine, Washington University School of Medicine, 660 South Euclid Ave, Saint Louis, MO, USA

*Indicates co-first authors

[#]Corresponding author
660 South Euclid Ave
Campus Box 8232
Saint Louis, MO, 63110
ph: 314-362-7269, fax: 314-362-7855

Keywords: glycemic control, brown adipose, islets, obesity, metabolism, mouse model, transcriptome, extracellular matrix, cytokines

Summary

On a high fat diet, obese SM/J mice initially develop metabolic dysfunction, including impaired glucose tolerance and elevated fasting glucose. These abnormalities resolve spontaneously by 30 weeks of age despite persistence of obesity. The mice dramatically expand their brown adipose depots as they resolve glycemic dysfunction. They also expand their pancreatic islet populations and improve beta cell function. When the brown adipose depot is removed from normoglycemic high fat-fed mice, fasting blood glucose and glucose tolerance revert to unhealthy levels. This occurs naturally and spontaneously on a high fat diet, with no temperature or genetic manipulation. We identified 267 genes whose expression changes in the brown adipose when the mice resolve their unhealthy glycemic parameters, and find the expanded tissue has a 'healthier' expression profile. Understanding the physiologic and genetic underpinnings of this phenomenon in SM/J mice will open the door for innovative therapies aimed at improving glycemic control.

Introduction

An estimated 10-30% of obese individuals maintain glycemic control (Karelis, 2008) and some longitudinal studies suggest their risk of developing type II diabetes is no greater than matched lean individuals (Meigs et al., 2006). No causative factors underlying glycemic control in obesity have been discovered, however the strongest predictors of impaired glycemic control in obesity are increased visceral fat mass and adipose tissue dysfunction (Goossens, 2017; Klöting et al., 2010). Thus research efforts have focused on understanding the genetic and physiological mechanisms of action of adipose (Rosen and Spiegelman, 2014). Recent research reveals that brown adipose activity is associated with anti-diabetic properties (Chechi et al., 2013; Cypess et al., 2009; Hanssen et al., 2016; van Marken Lichtenbelt et al., 2009a; Saito, 2013; Saito et al., 2009; Stanford et al., 2013; Virtanen et al., 2009). Cold exposure in both obese and lean individuals causes increased uptake of fatty acids and glucose into brown adipose tissue, which is associated with an increase in total energy expenditure (van Marken Lichtenbelt et al., 2009b; Ouellet et al., 2012; Saito et al., 2009). Further, increased brown adipose activity has been shown to improve glucose homeostasis and insulin sensitivity in adults (Chondronikola et al., 2014). The role of brown adipose in healthy metabolism is generally thought to be a function of its role in non-shivering thermogenesis. However recent research indicates brown adipose also has a secretory role, releasing adipokines into circulation that contribute to glycemic control, including fibroblast growth factor 21 (FGF21) (Stanford et al., 2013) and neuregulin 4 (NRG4) (Wang et al., 2014). Transplantation of brown adipose tissue into mouse models of both type I and type II diabetes greatly improves glucose parameters, including fasting glucose levels and response to a glucose challenge (Gunawardana and Piston, 2012, 2015), and this is thought to be due to brown adipose secreted factors (Villarroya et al., 2017). Thus there is significant interest in identifying the brown adipokines responsible for these anti-diabetic properties. While there are a variety of obese and diabetic mouse models, there are no mouse models for understanding the relationship between brown adipose and glycemic control in obesity.

The SM/J inbred mouse strain has long been used for studying interactions between diet and metabolism, and more recently has started to help uncover the genetic architecture underlying diet

induced obesity and glucose homeostasis (Cheverud et al., 2011; Lawson and Cheverud, 2010; Lawson

et al., 2010, 2011b, 2011a; Nikolskiy et al., 2015). It has previously been shown that fed a high fat diet,

SM/J mice display many of the characteristics of a diabetic-obese mouse: obesity, hyperglycemia,

glucose intolerance, and deficient insulin production at 20 weeks of age (Ehrich, 2003). Here, we report

that SM/J mice undergo a remarkable transformation between 20 and 30 weeks of age. Despite

persistence of the obese state, these mice enter into diabetic remission: returning to normoglycemia,

reestablishing glucose tolerance, and increasing insulin production without loss of insulin sensitivity.

Contemporary with this remission of glycemic parameters is a dramatic expansion of the interscapular

brown adipose depot. This study describes the morphological, physiological, and transcriptomic changes

that occur during this transition, and establishes the SM/J mouse as a unique model for understanding

the relationship between brown adipose and glycemic control in obesity. Understanding this relationship

in a natural model of glycemic resolution will set the stage for identifying novel, potentially therapeutic

targets for the improvement of glycemic control in obesity.

Results

SM/J mice improve glucose parameters without weight loss

The SM/J inbred mouse strain is used to explore dietary obesity and glycemic control (Lawson

and Cheverud, 2010). When fed a high fat diet (**Supplemental Table 1)**, SM/J mice develop obesity,

hyperglycemia, and impaired glucose tolerance by 20 weeks of age (Ehrich et al., 2003). We find that by

30 weeks of age, high fat-fed SM/J mice resolve their hyperglycemia and impaired glucose tolerance to

levels indistinguishable from low fat-fed controls, despite persistence of the obese state (**Figure 1A-D**).

Thirty-week-old high fat-fed SM/J mice experience a 2.5-fold increase in fasting insulin levels compared

to 20 week-old high fat-fed mice, and a 1.9-fold increase in fasting insulin compared to 30 week-old low

fat-fed mice (**Figure 1E**). This is reflected in a similar trend in total pancreatic insulin content (**Figure 1F**),

with no loss of insulin sensitivity, suggesting enhanced insulin secretion contributes to the resolution of

glucose parameters (**Figure 1G and H**).

Figure 1. Obese SM/J mice improve glucose parameters between 20 and 30 weeks of age. A High fat-fed mice weigh significantly more than low fat-fed mice, and SM/J mice gain weight between 20 and 30 weeks of age on both diets, n = 48-131 mice per cohort. **B** 30 week-old high fat-fed mice have significantly lower fasting glucose levels than at 20 weeks, which is no different than low fat-fed controls, n = 22-47 mice per cohort. **C and D** 30 week-old high fat-fed mice have improved glucose tolerance relative to 20 weeks, n = 22-47 mice per cohort. **E** Lower fasting glucose and improved glucose tolerance corresponds with increased serum and **F** pancreatic insulin, n = 10-20 mice per cohort. **G and H** This increase in insulin does not correspond to changes in insulin sensitivity in 30 week-old high fat-fed mice, n = 22-47 animals per cohort. Equal numbers of males and females represented; * p<0.05, ** p< 0.01, *** p <0.001

High fat-fed C57BL/6J mice also show a reduction in fasting glucose that is accompanied by increased insulin with age (Ahren, 2004). However, the difference in circulating glucose between the high fat- and low fat-fed controls remain significantly different over time. Moreover, high fat-fed C57BL/6J mice show marked glucose intolerance that does not resolve with age. We observe a similar trend in the LG/J strain of mice, where high fat-fed animals maintain higher fasting glucose levels and impaired glucose tolerance relative to low fat-fed controls as they age (**Supplemental Figure 1**). The unique remission of hyperglycemia and improved glucose tolerance observed in the high fat-fed SM/J strain indicates a genetic basis.

Increased islet cell replication underlies increased insulin production and improved islet function

To understand the potential causes of the increased insulin secretion in high fat-fed SM/J mice, we analyzed the morphology and function of pancreatic islets. The resolution of glucose parameters correlates with increased pancreatic islet mass of high fat-fed mice at 30 weeks compared to 20 weeks or to low fat-fed controls (**Figure 2A**). This increase in islet mass is driven by sustained mitosis of cells within the islet, with a substantial increase in beta cell mass between 20 and 30 week high fat-fed mice (**Figure 2B; Supplemental Figure 2**). A high fat diet and obesity are usually accompanied by a progressive decline in beta cell function (Kahn et al., 2001; Da Silva Xavier, 2018), however a static glucose-stimulated insulin secretion assay reveals that islets from high fat-fed SM/J mice improve insulin secretion efficiency between 20 and 30 weeks, transitioning from non-glucose sensitive to glucose sensitive (**Figure 2C**). Because high fat-fed SM/J islets are proliferative at 20 weeks, the non-existent

glucose-stimulated insulin secretion at 20 weeks suggests the nascent beta cells are not yet functionally

mature (**Figure 2B and C**). This contrasts with high fat-fed C57Bl/6J mice, which have increased

pancreatic islet size and reduced beta cell function (Roat et al., 2014), further underscoring the genetic

basis of this phenomenon.

Figure 2. 30 week-old high fat-fed SM/J mice increase islet replication and improve insulin
secretion. A Increased islet cell mass in high relative to low fat-fed mice, and in 30 relative to 20 week-
old high fat-fed mice. **B** Islet cells in high fat-fed mice have elevated levels of phosphohistone-H3, a
marker of mitosis. **C** Islets isolated from 30 week-old high fat-fed mice show restored insulin secretion
capacity in the presence of high glucose compared to 20 week-old high fat-fed mice. Equal numbers of
males and females represented, panels A and B: n = 12-16 mice per cohort; panel C: n = 6-8 mice per
cohort; * p<0.05, ** p< 0.01, *** p<0.001
174

High fat-fed SM/J mice expand their interscapular brown adipose tissue depots

In conjunction with the resolution of glycemic parameters and improved insulin secretion, high fat-

fed SM/J mice dramatically expand their interscapular brown adipose depots, which is not seen in low

fat-fed control mice (**Figure 3A-C**). This has never been described in another mouse strain, and we do

not observe the phenomenon in the LG/J strain of mice on the same diets at any age (**Supplemental**

Figure 3). To understand whether the tissue mass expansion is due to increased size of individual cells

or to increased number of total cells, we quantified adipocyte cell size and percent of phosphohistone-

H3 positive cells. There are no significant differences in average cell size in high fat-fed mice between

20 and 30 weeks, or relative to low fat-fed controls (**Supplemental Figure 4A**). Mice on both diets

undergo altered adipocyte area profiles between 20 and 30 weeks of age, however the low fat tissue

develops a profile significantly trending towards larger adipocytes at 30 weeks ($p=6.4^{-07}$) whereas the

high fat tissue develops a profile significantly trending towards smaller adipocytes at 30 weeks ($p=2.2^{-16}$)

(**Figure 3D and E**). This suggests that the expansion of the brown adipose depot in high fat-fed mice is

the result of increased proliferation of adipocytes, as newer adipocytes are smaller due to less lipid

accumulation. This is supported by quantification of brown adipose cells stained positive for the mitotic

marker phosphohistone H3, which trends towards higher mitosis in the brown adipose of high fat-fed

animals (**Supplemental Figure 4B**).

Obesity has been associated with structural and functional "whitening" of brown adipose depots

in rodents (Lapa et al., 2017; Roberts-Toler et al., 2015; Shimizu and Walsh, 2015; Shimizu et al., 2014).

Histological analysis of the fat depot taken from high fat-fed SM/J mice at 30 weeks of age confirms the

adipocytes in this expansion are brown adipocytes, with small multilocular lipid droplets and high UCP1

staining (**Supplemental Figure 5**). Expression of canonical brown adipose genes *Ucp1* and *Cidea* do

not change between 20 and 30 weeks (**Figure 3F-G).** Further, expression of *Tbx1*, a marker specific for

beige adipocytes (Wu et al., 2012), indicates that neither brown nor white adipose is "beiging" (**Figure**

3H). Finally, there is no significant difference in brown adipose tissue mitochondrial content between the

diets or ages (**Figure 3I**). There is no difference in core body temperature or circulating free fatty acids

between high and low fat-fed cohorts or between 20 and 30 weeks of age (**Supplemental Figure 6A**

and B). Additionally, while there are diet-dependent differences in the catecholamines norepinephrine

and epinephrine, which activate UCP1-mediated leak respiration and non-shivering thermogenesis, there

is no change in levels between ages in the high fat-fed mice (**Supplemental Figure 6C and D**). Thus,

the interscapular adipose depot in high fat-fed SM/J mice maintains a brown adipose identity after

expansion that is not dependent on whole-animal beiging, and is also not associated with altered

thermogenesis.

Figure 3. Brown adipose expansion in 30 week-old high fat-fed SM/J mice. Representative pictures of **A** 20 week and **B** 30 week-old high fat-fed female mice. **C** Quantification of interscapular brown adipose depot as a proportion of total fat mass, n = 16-25 mice per cohort. **D and E** Cell area density graphs for high fat and low fat-fed cohorts. Data are plotted on a log10 scale for visualization, n = 4 mice per cohort. **F-H** Gene expression levels quantified in three adipose depots of high fat-fed mice: interscapular brown adipose (iBAT), inguinal white adipose (ingWAT), and reproductive white adipose (repWAT), n = 6-10 mice per cohort and tissue. Canonical brown adipose genes (**F)** *Ucp1* and (**G)** *Cidea* show high expression in iBAT and no difference between 20 and 30 week-old mice. Beige adipose marker (**H)** *Tbx1* is not expressed in any depot. **I** Mitochondrial copy number was significantly higher in brown adipose tissue than in inguinal white adipose tissue at both 20 and 30 week time points with no significant difference between high or low fat-fed mice, n = 6-7 mice per cohort and tissue. Equal numbers of males and females represented. * $p<0.05$, ** $p< 0.01$

RNA sequencing reveals enrichment of differentially expressed cytokines and genes affecting extra

cellular matrix

Since the brown adipose tissue expansion is unique to high fat-fed SM/J mice, we anticipated that there would be corresponding unique transcriptomic changes in the brown adipose. Indeed, we identified 267 genes whose expression significantly and uniquely changes between 20 and 30 weeks of age in high fat-fed SM/J brown adipose tissue (at a 5% FDR, out of 13,253 total genes expressed; **Supplemental Table 2**). These expression changes occur when the mice resolve their unhealthy glycemic parameters and expand their brown adipose depots. These genes are not differentially expressed in white adipose tissue taken from the same animals or in low fat-fed SM/J controls (**Figure 4A**). Additionally, they are not differentially expressed in the LG/J strain of mouse, again underscoring the genetic basis of the phenomenon (**Supplemental Figure 7; Supplemental Table 3**).

Over-representation analysis indicates these genes are enriched for those involved in cytokine-cytokine receptor interactions ($p=3.23e^{-06}$), signaling receptor activity ($p = 5.70e^{-06}$), cell surface receptor signaling ($p=2.04e^{-07}$), and extracellular matrix components ($p = 7.93 e^{-13}$), among others (**Figure 4B; Supplemental Table 4**). These are intriguing results because brown adipose has been identified as a source of cytokines that influence beta cell health and glucose homeostasis (Villarroya et al., 2017; Wang et al., 2015), and extracellular matrix changes are essential for tissue expansion, cellular signaling, and regulation of growth factor bioavailability (Frantz et al., 2010).

Several genes belonging to these biological categories have evidence for their involvement in

glucose homeostasis and change expression in a direction that is associated with improved metabolic

health in high fat-fed SM/J mice between 20 and 30 weeks of age (**Figure 4C**). In particular, the direction

of expression change reveals that the expansion of brown adipose is associated with a decrease in

inflammatory (e.g. interleukin 7 receptor, *Il7r*) (Kim et al., 2014) and fibrotic markers (e.g. collagen type

VIII alpha 1 chain, *Col8a1*; semaphorin 3C, *Sema3c*) (Mejhert et al., 2013; Sun et al., 2013), and changes

in extracellular matrix components (e.g. matrix metallopeptidase 12, *Mmp12*; procollagen c-

endopeptidase enhancer, *Pcolce*) (Huang et al., 2011; Lee et al., 2014) and cytokine activity (e.g.

coagulation factor VII, *F7*; leptin, *Lep*; secreted frizzled-related protein 1, *Sfrp1*) (D'souza et al., 2017;

Edén et al., 2015; Gauger et al., 2013) (**Supplemental Figure 8**). Other mouse models of diet-induced

obesity develop unhealthy brown adipose transcriptomes characterized by increased expression of pro-

inflammatory genes and fibrotic markers (McGregor et al., 2013; Alcalá et al., 2017). The direction of

expression change in our brown adipose tissue supports the uniqueness of the high fat-fed SM/J mice.

Figure 4. High fat-fed SM/J mice have unique brown adipose differential expression between 20 and 30 weeks of age. A Venn diagram illustrating the number of genes differentially expressed between high and low fat-fed 20 and 30 week-old SM/J interscapular brown or reproductive white adipose tissues. No genes are differentially expressed in more than one diet-by-tissue cohort. **B** Enriched terms colored by class (Gene Ontology Biological Process (GO_BP), Cellular Component (GO_CC), Molecular Function (GO_MF), and KEGG Pathway. **C** Heatmap of differentially expressed brown adipose tissue genes between high fat-fed 20 and 30 week-old mice belonging to cytokine, extracellular matrix, or both gene ontologies. Equal numbers of males and females represented, n = 8 animals per age-by-diet cohort.

30 week old high fat-fed SM/J brown adipose has a healthier co-expression profile

Because variation in glucose homeostasis is complex and the result of many interacting genes, we examined the co-expression profile of genes belonging to the enriched cytokine and extracellular matrix (ECM) biological categories (**Figure 4B and C**). We find that the structure of the differentially expressed ECM and cytokine genes is significantly different between 20 and 30 week-old high fat-fed

animals (p=0.01). To determine if the co-expression structure of these genes in 30 week-old high fat-fed

animals' brown adipose is more similar to the 20 week-old high fat-fed or to the low fat-fed animals', we

compared the overall co-expression correlation structure between the diet and age cohorts for these

genes. Remarkably, we find the 30 week-old high fat-fed small brown adipose ECM and cytokine co-

expression profile is most similar to the 20 week-old low fat-fed animals' (probability of difference between

high fat-fed 30 weeks and low fat-fed 20 weeks = 0.07; probability of difference between high fat-fed 30

weeks and low fat-fed 30 weeks = 0.04) (**Figure 5A**). Thus, the brown adipose cytokine and ECM gene

co-expression profile appears 'healthier' in 30 week-old high fat-fed animals after expansion and

remission of the diabetic phenotype. This is illustrated in **Figure 5B**.

B

Figure 5. High fat-fed 30 week-old SM/J mice have a cytokine and ECM gene co-expression profile most similar to low fat-fed 20 week-old mice. A The median change in correlation structure is plotted as a vertical line against the null model. High fat-fed 30 week-old SM/J mice have a co-expression profile significantly different from high fat-fed 20 week-old SM/J mice, and not significantly different from low fat-fed 20 week-old mice. **B** Heatmap of the gene expression correlation matrices for each cohort. HF30 = high fat-fed 30 week-old; HF20 = high fat-fed 20 week-old; LF30 = low fat-fed 30 week-old; LF20 = low

284 fat-fed 20 week-old. Equal numbers of males and females represented, n = 8 animals per age-by-diet
285 cohort.
286

Glucose parameters revert to an unhealthy state in SM/J mice when the brown adipose depot is removed

If the brown adipose expansion is directly related to the glycemic resolution of the high fat-fed

SM/J mice, removing that expansion should revert the glucose parameters to their unhealthy state. To

test this, we removed the interscapular brown adipose depots from normoglycemic 30 week-old mice.

After recovery, at 35 weeks of age, we measured basal glucose levels and performed a glucose tolerance

test. We find that glucose parameters revert to unhealthy, 20 week-old measurements when the brown

adipose depot is removed (**Figure 6 A-B**), indicating that the expanded brown adipose tissue is

necessary for the observed remission of unhealthy glycemic parameters in high fat-fed SM/J mice.

Figure 6. Glycemic parameters revert to unhealthy levels when the brown adipose depot is
removed. A Blood glucose assessed after a 4 hour fast. **B** A glucose tolerance test indicates that glucose
tolerance is reduced after the brown adipose depot is removed. n = 5 excision animals, representing 4
males and 1 female.
300

Discussion

Obesity (body-mass index [BMI] ≥30 kg.m^2) is associated with serious metabolic complications,

including type II diabetes, cardiovascular disease, cancer, and stroke (Abdullah et al., 2010; Kenchaiah

et al., 2002; Rauscher et al., 2000; Reeves et al., 2007; Strazzullo et al., 2010). Currently, 38% of

American adults are classified as obese (Flegal et al., 2016), 9.4% have type II diabetes, and an

additional 34% are pre-diabetic, costing 327 billion dollars in annual medical costs (Centers for Disease

Control and Prevention, 2017; Yang et al., 2018). Obesity and diabetes are tightly linked; obesity raises

the risk of developing type II diabetes 27—76 fold, while approximately 60% of diabetics are obese

(Abdullah et al., 2010; Centers for Disease Control and Prevention, 2017; Chatterjee et al., 2017; Colditz

et al., 1995). The obesity-diabetes axis centers on glycemic control, where excess caloric intake and

body fat demands increased beta cell insulin secretion to maintain normoglycemia. In many individuals,

the prolonged obese state results in insulin resistance, beta cell death, loss of insulin production, and

chronic hyperglycemia. Though weight loss is *the* gold standard for treating glycemic dysfunction in

obesity, many obese people are unable to achieve long-term weight loss (Dulloo and Montani, 2015;

Tomiyama et al., 2013). Currently, metformin and thiazolidinediones are prescribed to prevent the

development of diabetes in obese individuals, but pharmacological therapy has been shown to have only

modest protective effects (Chatterjee et al., 2017; Nathan et al., 2015). Greater understanding of the

relationship between obesity and glycemic control is needed to develop more effective preventative

measures for obese patients.

Here we describe morphological, physiological, and transcriptomic changes that occur during

brown adipose expansion and remission of hyperglycemia in SM/J mice. The SM/J strain was derived

from a pool of seven inbred strains and selected for small body size at 60 days (Beck et al., 2000;

MacArthur, 2002; Nikolskiy et al., 2015). The strain has been used extensively in genetic studies of

complex traits related to growth and metabolism (Kenney-Hunt et al., 2006; Lawson et al., 2010, 2011a,

2017; Norgard et al., 2011), particularly because SM/J mice are strongly responsive to high fat diet-

induced obesity. These studies all used mice aged 20 weeks or less, when SM/J's develop the classic

hallmarks of obese-diabetic mice. We discovered that by 30 weeks of age, coinciding with a dramatic

expansion of interscapular brown adipose tissue, their hyperglycemia, impaired glucose tolerance, and

deficient insulin secretion in response to glucose stimulation goes into remission despite persistence of

obesity. Dissecting the genetic basis of this phenomenon has the potential to uncover novel relationships

among brown adipose, glucose homeostasis, and obesity.

We identified 267 genes whose expression significantly and uniquely changes between 20 and

30 weeks of age in high fat-fed SM/J brown adipose tissue. We hypothesize that these genes affect

brown adipose function and contribute to the phenomenon we observe. The expression changes occur

when the mice resolve their unhealthy glycemic parameters and expand their brown adipose depots.

These genes are not differentially expressed in white adipose tissue taken from the same animals. These

genes are not differentially expressed between low fat-fed intra-strain controls at the same ages or in a

unique mouse strain that does not resolve glycemic parameters or expand brown adipose tissue. Further,

30 week-old high fat-fed SM/J brown adipose has an overall 'healthier' expression profile, supported by

the analysis of the correlation structure among the age-by-diet cohorts for cytokines and ECM genes

(**Figure 5A**).

We focus on genes associated with ECM and cytokine activity because both biological categories

are enriched in the set of genes that significantly change expression in brown adipose during the

remission of glycemic parameters. Brown adipose is a source of endocrine signals with anti-diabetic

properties (Stanford et al., 2013; Wang et al., 2014) and is involved in extensive cross-talk with other

organs (Poekes et al., 2015). It secretes cytokines that influence whole-body glucose homeostasis and

insulin sensitivity including IGF1, FGF21, NRG-3 and NRG-4 (Kajimura et al., 2015; Wang et al., 2015).

ECM changes are essential for cellular signaling, regulation of growth factor bioavailability, and

accompany healthy adipose expansion. However, extreme changes in ECM protein levels are associated

with adipose dysfunction in obesity; thus a fine balance between tissue remodeling and excessive

accumulation of ECM proteins must be achieved to maintain adipose tissue homeostasis (Hasegawa et

al., 2018; Sun et al., 2013).

We highlight 8 cytokines and ECM genes that significantly change expression in a direction

associated with improved metabolic health in previous studies. *Il7r*, which was found to be one of the

highest ranking genes in the white adipose tissue inflammatory response pathway (Moreno-Viedma et

al., 2016b), decreases expression between 20 and 30 weeks of age in high fat-fed SM/J brown adipose.

Col8a1 and *Sema3C* are both associated with adipose tissue fibrosis (Hasegawa et al., 2018; Mejhert et

al., 2013). Increased adipose tissue fibrosis is a signature of dysfunctional adipose and is associated with

impaired glucose homeostasis and insulin resistance (Sun et al., 2013). Both *Col8a1* and *Sema3c*

expression decrease between 20 and 30 weeks in high fat-fed SM/J brown adipose. *Mmp12* is an enzyme

that contributes to adipose tissue remodeling (Maquoi et al., 2002). Increased *Mmp12* expression is

associated with white adipose tissue inflammation and insulin resistance and *Mmp12*[-/-] mice are more

insulin sensitive than wildtype controls on a high fat diet (Lee et al., 2014). Its expression decreases in

30 week old high fat-fed SM/J brown adipose. *Pcolce* encodes a glycoprotein that regulates collagen

processing at the ECM (Raz et al., 2013). Mice with defects in ECM collagen are glucose intolerant,

hyperglycemic, and insulin resistant (Huang et al., 2011). PCOLCE is one of 15 key drivers that

collectively account for 22% of GWAS hits for type II diabetes in a recent multiethnic meta-analysis (Shu

et al., 2017). *Pcolce* expression is significantly increased in 30 week old high fat-fed SM/J brown adipose.

F7, *Lep*, and *Sfrp1* are each secreted proteins. Increased *F7* plays a role in the pathogenesis of obesity

(Takahashi et al., 2015). In particular it has been shown to induce beta cell death and impaired islet

glucose-stimulated insulin secretion (Edén et al., 2015). Increased *Lep* can dramatically lower blood

glucose levels in diabetic rodent models (D'souza et al., 2017). In brown adipose, leptin has been shown

to stimulate glucose uptake (Denroche et al., 2016). *Sfrp1* is dysregulated in obesity and *Sfrp1*[-/-] mice

have elevated blood glucose and impaired glucose tolerance when fed a high fat diet (Gauger et al.,

2013; Lagathu et al., 2010). *F7* expression is decreased and *Lep* and *Sfrp1* are increased in 30 week old

high fat-fed SM/J brown adipose tissue. Most of what is known about the role of these 8 genes in adipose

comes from studies of white adipose tissue, but none of these genes are differentially expressed in SM/J

white adipose. Many additional genes likely contribute to the observed phenomenon, however little, if

anything, is known about their role in brown adipose tissue. The 267 differentially expressed genes we

identified represent a set of actionable candidates for further functional studies of their role in brown

adipose and glucose homeostasis.

There is great interest in harnessing the potential of brown adipose to treat obesity and diabetes,

either through the calorie burning action of non-shivering thermogenesis or the endocrine action of

adipokines. Research into the effects of brown adipose on systemic metabolism is in its infancy, and the

community needs appropriate animal models to interrogate its physiological roles and identify potentially

druggable targets. We present the SM/J mouse strain as a unique model to address this need. High fat-

fed obese SM/J mice revert to normoglycemic at 30 weeks of age. High fat-fed normoglycemic SM/J mice

have dramatically expanded brown adipose depots and improved islet glucose-stimulated insulin

secretion. When the brown adipose depot is removed from normoglycemic high fat-fed SM/J mice, fasting

blood glucose and glucose tolerance revert to unhealthy levels. This occurs naturally and spontaneously

on a high fat diet, with no temperature or genetic manipulation. To our knowledge this has never been

described in another mouse strain and our transcriptomic studies indicate the phenomenon is genetic.

The SM/J mouse provides a tractable system in which to understand the relationship between brown

adipose and glycemic control in obesity. Understanding this relationship in the SM/J mouse will open

doors for identifying novel, potentially druggable targets for the improvement of glycemic control in

humans.

Methods

Animal Husbandry and Phenotyping

SM/J and LG/J mice were obtained from The Jackson Laboratory (Bar Harbor, ME). Experimental

animals were generated at the Washington University School of Medicine and all experiments were

approved by the Institutional Animal Care and Use Committee in accordance with the National Institutes

of Health guidelines for the care and use of laboratory animals. Pups were weaned at 3 weeks and reared

in same-sex cages of 3-5 animals until necropsy. At weaning, mice were randomly placed on a high fat

diet (42% kcal from fat; Teklad TD88137) or an isocaloric low fat diet (15% kcal from fat; Research Diets

D12284) (**Supplemental Table 1**). Feeding was *ad libitum*. The animal facility operates on a 12 hour

light/dark cycle with a constant ambient temperature of 21°C. Animals were weighed weekly until

sacrifice. At 18 and 28 weeks of age, animals were subject to an intraperitoneal glucose tolerance test

after a 4 hour fast. At 19 and 29 weeks of age animals were subject to an intraperitoneal insulin tolerance

test. At 20 or 30 weeks of age, body composition was determined by MRI and temperature was measured

with a rectal thermometer. After a 4 hour fast, at 20 or 30 weeks of age, animals were given an overdose

of sodium pentobarbital and blood was collected via cardiac puncture. Euthanasia was achieved by

cardiac perfusion with phosphate-buffered saline. After cardiac perfusion, tissues were collected and

flash frozen in liquid nitrogen and stored at -80°C, or processed according to protocols for histology and

other assays.

Blood plasma assays

Fasting blood glucose was measured using a GLUCOCARD Vital glucometer (Arkay, MN USA).

ELISAs measuring plasma levels of insulin (ALPCO 80-INSMR-CH01) and free fatty acids (Wako Life

Sciences 995-34693) were quantified according to manufacturer's protocol. Catecholamines were

assayed through the Vanderbilt University Medical Center's Hormone Assay and Analytical Services

Core (www.vumc.org/hormone/assays; NIH grants DK059637 (MMPC) and DK020593 (DRTC)).

Pancreatic insulin content

Whole pancreas was homogenized in acid ethanol and incubated at 4°C for 48 hours, shaking.

Homogenate was centrifuged at 2500 rpm for 30 min at 4°C. Supernatant was collected and stored at -

20°C. Protein content was measured using Pierce BCA Protein Assay kit (Thermo Scientific) according

to manufacturer's instructions, and read at 562 nm on the Synergy H1 Microplate Reader (Biotek). Insulin

content was measured with ALPCO ultrasensitive Insulin ELISA (ALPCO 80-INSMR-CH01) according to

manufacturer's instructions.

Glucose-stimulated insulin secretion

Pancreas was removed and placed in 8mL HBSS buffer on ice. Pancreas was then thoroughly

minced. 120mg Collagenase P (Roche) was dissolved in HBSS and aliquoted into 20 x 500ul tubes. One

tube (500ul) was added to the minced pancreas in 8ml HBSS. Mixture was then shaken in a 37°C water

bath for 12 minutes. Mixture was spun at 2000 rpm for 1 minute. The pellet was washed twice with HBSS,

spinning in between. The pellet was re-suspended in HBSS and transferred a petri dish. Hand-selected

islets where placed in sterile-filtered RPMI with 11mM glucose, 5% pen/strep, and 10% Fetal Bovine

Serum. Islets were rested overnight in a cell culture incubator set to 37°C with 5% C02. The following

day, islets were equilibrated in KRBH buffer containing 2.8 mM glucose for 30 minutes at 37°C. 5 Islets

were hand selected and placed in 150ul KRBH containing either 2.8 or 11mM glucose. Tubes were placed

in a 37°C water bath for 45 min. Islets were then spun at 2000RPM, hand-picked with a pipette, and

transferred from the secretion tube and placed in the content tube with acid ethanol. The content and

secretion tubes were stored at -20°C overnight. Each condition was performed in duplicate for each

individual.

ALPCO Ultrasensitive ELISA (80-INSMU-E01) was performed according to manufacturer's

instructions, with the secretion tubes diluted 1:5, and content tubes diluted 1:100. Normalized insulin

secretion was calculated by dividing the secreted value by the content value. Glucose stimulated insulin

secretion was calculated by dividing the normalized insulin secretion at 11mM by the normalized insulin

secretion at 2.8mM. Each sample was measured in duplicate.

Islet histology and analyses

At the time of tissue collection, whole pancreas was placed in 3 mL of neutral buffered formalin.

These samples were incubated at 4C while gently shaking for 24 hours. Immediately afterwards, samples

were placed into plastic cages and acclimated to 50% EtOH for 1 hour. Samples were then processed

into paraffin blocks using a Leica tissue processor with the following protocol: 70% EtOH for 1 hour x 2,

85% EtOH for 1 hour, 95% EtOH for 1 hour x 2, 100% EtOH for 1 hour x 2, Xylenes for 1 hour x 2, paraffin

wax. Pancreas blocks were sectioned into four sections 4 μm thick at least 100 μm from each other.

Slides were incubated at 60C for 1 hour, then placed in xylenes to remove remaining paraffin wax.

Slides were then rehydrated using successive decreasing EtOH concentrations (xylenes x 2, 50% EtOH

in xylenes, 100% EtOH x 2, 95% EtOH, 70% EtOH, 50% EtOH, H2O). Slides were incubated in sodium

citrate (pH 6) at 85C for 30 minutes, then submerged in running water for 5 minutes. Slides were washed

with 0.025% Triton X-100 in TBS and blocked in 10% normal donkey serum for 1 hour (Abcam ab7475),

followed by incubation with primary antibody overnight at 4C. [Primary antibodies: rat anti-insulin (1:100,

R&D MAB1417), mouse anti-glucagon (1:100, abcam ab10988), and rabbit anti-phospho-histone H3

(1:100, Sigma SAB4504429)]. After an additional wash, secondary antibody was applied for 1 hour at

room temperature. [Secondary antibodies: donkey anti-rabbit 488 (1:1000, abcam ab150061), donkey

anti-mouse 647 (1:1000, abcam ab150107), and donkey anti-rat 555 (1:1000, abcam ab 150154)].

Fluoroshield Mounting Medium with DAPI (Abcam) was applied to seal the coverslip and slides were

stored at 4C. Imaging was performed using the Zeiss AxioScan .Z1 at 20X magnification and 94.79%

laser intensity.

Background was subtracted from DAPI, insulin, glucagon, and PHH3 channels using ImageJ.

DAPI channel was used to identify total nuclei in CellProfiler. Insulin and glucagon channels were

combined and overlaid on the DAPI image to identify islet nuclei. Islet nuclei images were overlaid with

PHH3 stain to identify mitotic islet nuclei. Total nuclei, islet nuclei, and mitotic nuclei were summed across

all 4 slides for each individual, 12-16 individuals per cohort. Islet mass is reported as islet nuclei divided

by total nuclei. Mitotic islet index is reported as mitotic islet nuclei divided by islet nuclei.

Brown adipose histology

At the time of tissue collection, small portions of interscapular brown and reproductive white

adipose tissues were placed in 1 mL of neutral buffered formalin. These samples were incubated at 4C

while gently shaking for 24 hours. Immediately afterwards, samples were placed into plastic cages and

processed into paraffin blocks using a Leica tissue processor with the following protocol: 70% EtOH for

1 hour x 2, 85% EtOH for 1 hour, 95% EtOH for 1 hour x 2, 100% EtOH for 1 hour x 2, Xylenes for 1 hour

x 2, paraffin wax. Adipose blocks were sectioned into 6 µm sections, with 2-4 slices on each slide.

H&E Staining

Slides were incubated at 60C for 1 hour, then placed in xylenes to remove remaining paraffin wax. Slides were then rehydrated using successive decreasing EtOH concentrations (xylenes x 2, 100% EtOH x 2, 95% EtOH, 70% EtOH, H2O). Slides were incubated in hematoxylin (Leica Surgipath 3801570), Define (3803590), Blue Buffer 8 (3802915), and eosin (3801616), and dehydrated (95% EtOH, 100% EtOH, xylene x 2). Imaging was performed using the Zeiss AxioPlan2 microscope and Olympus DP software. Analysis of adipocyte size was performed using ImageJ. Images were converted to black and white and skeletonized to reveal only the cell wall outlines. Cell area was calculated from outlines with a lower limit of 50 um and upper limit of 700 um to reduce noise. All cells from a cohort (4-7 images each from 4 animals per cohort, equal numbers of males and females) were pooled for cell area density analysis. A Welch's unequal variances t-test was performed between ages in each diet to determine significant differences.

Immunofluorescence

Slides were incubated at 60C for 1 hour, then placed in xylenes to remove remaining paraffin wax. Slides were then rehydrated using successive decreasing EtOH concentrations (xylenes x 2, 50% EtOH in xylenes, 100% EtOH x 2, 95% EtOH, 70% EtOH, 50% EtOH, 0.3% H2O2 in MeOH, H2O). Slides were washed with TBS and blocked in 10% normal donkey serum (Abcam ab7475) for 1 hour, followed by incubation with primary antibody overnight at 4C. [Primary antibodies: rabbit anti-Ucp1 (1:100, Sigma U6382) and mouse anti-PHH3 (1:100, Invitrogen MA5-15220)]. After an additional wash, secondary antibody was applied for 1 hour at room temperature [Secondary antibodies: donkey anti-rabbit 488 (1:1000, Abcam ab150061) and donkey anti-mouse 647 (1:200, Abcam ab150107)]. Fluoroshield Mounting Medium with DAPI (Abcam) was applied to seal the coverslip and slides were stored at 4C.

Imaging was performed using the Zeiss Confocal microscope and Zen Lite imaging program. PHH3

analysis was performed using the CellProfiler program. Background was subtracted from DAPI and PHH3

channels using ImageJ. DAPI channel was used to identify total nuclei in CellProfiler. Adipose nuclei

images were overlaid with PHH3 stain to identify mitotic adipose nuclei. Mitotic nuclei were summed

across all 4 slides for each individual. Mitotic adipose index is reported as mitotic adipose nuclei divided

by adipose nuclei multiplied by 100%.

Quantitative rt-PCR

Total RNA was extracted from brown, inguinal, and reproductive adipose samples using the

Qiagen RNeasy Lipid Kit. High-Capacity cDNA Reverse Transcription Kit (Thermofisher) was used for

reverse transcription. Quantitative-rtPCR was performed to assess expression levels of target genes with

an Applied Biosystems (USA) QuantStudio 6 Flex instrument using SYBR Green reagent. Results were

normalized to *L32* expression, which was experimentally determined to not be differentially expressed

across diet and age cohorts. cDNA products were analyzed using the ΔC_T method. Primers used: *L32*

forward TCCACAATGTCAAGGAGCTG, reverse GGGATTGGTGACTCTGATGG; *Cidea* forward

TGCTCTTCTGTATCGCCCAGT, reverse GCCGTGTTAAGGAATCTGCTG; *Tbx1* forward

GGCAGGCAGACGAATGTTC, reverse TTGTCATCTACGGGCACAAAG; *Ucp1* forward

CCTCTCCAGTGGATGTGGTAA, reverse AGAAGCCACAAACCCTTTGA.

Mitochondrial DNA quantification

DNA was extracted from brown and inguinal adipose tissues using the Qiagen DNeasy Blood and

Tissue Kit. Briefly, 40mg of tissue was homogenized in 10% proteinase K through vortexing and

incubation at 56°C. DNA was precipitated with ethanol, collected in a spin column, and eluted in 150mL

of buffer. DNA concentration was quantified on a Nanodrop, and 50ng was used in a qPCR reaction to

quantify the amount of *h19* (nuclear gene) and *CytB* (mitochondrial gene). Mitochondrial content was

calculated   as   the   ratio   of   mtDNA   to   nucDNA.   Primers   used: *Cytb*   forward

TCTACGCTCAATCCCCAATAAAC,   reverse   TTAGGCTTCGTTGCTTTGAGGT;   *h19*   forward

TATGTGCCATTCTGCTGCGA, reverse AAGGTTTAGAGAGGGGGCCT.

RNA sequencing and analyses

Sixty-four LG/J and SM/J mice were used for sequencing analysis, representing 4 males and 4

females from each diet (high and low fat) and age (20 and 30 weeks). Total RNA was isolated from

interscapular brown and reproductive white adipose tissues using the RNeasy Lipid Tissue Kit (QIAgen).

RNA concentration was measured via Nanodrop and RNA quality/integrity was assessed with a

BioAnalyzer (Agilent). RNAseq libraries were constructed using the RiboZero kit (Illumina) from total RNA

samples with RIN scores >7.5. Libraries were checked for quality and concentration using the DNA

1000LabChip assay (Agilent) and quantitative PCR, according to manufacturer's protocol. Libraries were

sequenced at 2x100 paired end reads on an Illumina HiSeq 4000. After sequencing, reads were de-

multiplexed and assigned to individual samples.

FASTQ files were filtered to remove low quality reads and aligned against LG/J and SM/J custom

genomes using STAR (Dobin et al., 2013; Nikolskiy et al., 2015). Briefly, LG/J and SM/J indels and SNVs

were leveraged to construct strain-specific genomes using the GRC38.72-mm10 reference as a template.

This was done by replacing reference bases with alternative LG/J and SM/J bases using custom python

scripts. Ensembl R72 annotations were adjusted for indel-induced indexing differences for both genomes

(Macias-Velasco et al., 2019). Read counts were normalized via upper quartile normalization and a

minimum normalized read depth of 10 was required. Alignment summaries are provided in **Supplemental**

Table 5 and Supplemental Figure 9. Library complexity was assessed and differential expression

between each age cohort for each strain-by-diet comparison was determined after TMM normalization in

EdgeR (Chen et al., 2015) (**Supplemental Figure 10**).

Functional enrichment of differentially expressed genes was tested by over-representation

analysis in the WEB-based Gene Set Analysis Toolkit v2019 (Zhang et al., 2005). We performed analyses

of gene ontologies (biological process, cellular component, molecular function), pathway (KEGG), and

phenotype (Mammalian Phenotype Ontology). For each tissue, the list of all unique differentially

expressed genes was analyzed against the background of all unique genes expressed in that tissue

(Supplemental Tables 2 and 3). A Benjamini-Hochberg FDR-corrected p-value ≤ 0.05 was considered

significant.

Correlation structure

568   Co-expression was assessed for the set of 62 differentially expressed cytokines and ECM genes

569   by correlating expression of each gene with the expression of the other 61 genes in each diet-by-age

570   cohort. Each pair of genes then had their correlations correlated (R_g), where gene: G.

571
$$R_{g,G\in(i,j)} = cor\left(\begin{bmatrix} cor(G_i,G_1) \\ \vdots \\ cor(G_i,G_n) \end{bmatrix}, \begin{bmatrix} cor(G_j,G_1) \\ \vdots \\ cor(G_j,G_n) \end{bmatrix}\right)$$

573   Gene-pair-correlations were then compared between the high fat-fed 30 week-old cohort and the other

574   three cohorts (high fat-fed 30 weeks to high fat-fed 20 weeks, high fat-fed 30 weeks to low fat-fed 30

575   weeks, high fat-fed 30 weeks to low fat-fed 20 weeks) to obtain the ΔR_g between a pair of cohorts, where

576   cohort: K.

577
$$\Delta R_{g,G\in(i,j),K_1,K_2} = \left|R_{g,G\in(i,j),K_1} - R_{g,G\in(i,j),K_2}\right|$$

579   The median change in correlation ($M\Delta R_g$) was calculated and permutation was employed to identify the

580   background of expected $M\Delta R_g$ values. Permutation was performed by randomly selecting 2 groups of 8

581   animals from any cohort 10,000 times.

582
$$M\Delta R_{g,K_1,K_2} = median\left(\Delta R_{g,G\in(i,j),K_1,K_2}\right)$$

$M\Delta R_g$ was determined for the 2 randomized groups (rK_1, rK_2) for all 10,000 permutations to generate a

null model. Log transformation was performed to approximate normality, which was determined by Wilks-

Shapiro test and Q-Q plot. Significance was drawn from the cumulative normal null model to test if the

difference in correlation structure between each pair of cohorts was greater than by chance under the

randomized null model.

$$p_{K_1,K_2} = P\big(X \geq M\Delta R_{g,K_1,K_2}\big) \sim \mathcal{N}(\mu_r, \sigma_r)$$

Brown adipose excision

Interscapular brown adipose tissue depots were removed from 30 week-old high fat-fed SM/J

mice. A small longitudinal incision was made between the shoulder blades. All interscapular adipose

tissue was carefully removed, and a cauterizing wand used to stop excessive bleeding when necessary.

Surgeries were performed under general anesthesia by IP injection of ketamine/ xylazine (100/200

mg/Kg) and mice were maintained in the surgical plane by isofluorane/oxygen for the duration of the

procedure. Incisions were closed with 5-0 nonabsorbable sutures. Ketoprofen (2-5 mg/Kg) was provided

post-procedure and topical antibiotic was applied to the incision for up to 3 days as necessary. Animal

health and well-being was monitored daily. Sutures were removed at 10 days post-surgery. Four weeks

after surgery, mice underwent a glucose tolerance test and an insulin tolerance test one week later. After

an additional week of recovery, animals were sacrificed and serum and multiple tissues harvested

(reproductive and inguinal adipose depots, liver, heart, soleus, pancreas, hypothalamus) as described

above.

Statistics

Data within individual cohorts were assessed for normality using a Wilks-Shapiro test. Islet mass

and mitotic islet cell numbers were log10 transformed to achieve a normal distribution. Outliers were

identified by a Grubbs test ($p < 0.05$) and removed. Data were tested for significant differences among

cohorts by ANOVA with a Tukey's post-hoc correction. The sex-by-diet-by-age term was not significant

for any phenotype so males and females were pooled for analyses. P-values <0.05 were considered

significant. All statistical analyses were performed using the R software package.

References

Abdullah, A., Peeters, A., de Courten, M., and Stoelwinder, J. (2010). The magnitude of association

between overweight and obesity and the risk of diabetes: A meta-analysis of prospective cohort

studies. Diabetes Res. Clin. Pract.

Ahren, W. and (2004). The High-Fat Diet–Fed Mouse. Diabetes.

Beck, J.A., Lloyd, S., Hafezparast, M., Lennon-Pierce, M., Eppig, J.T., Festing, M.F.W., and Fisher,

E.M.C. (2000). Genealogies of mouse inbred strains. Nat. Genet.

CANNON, B. (2004). Brown Adipose Tissue: Function and Physiological Significance. Physiol. Rev.

Centers for Disease Control and Prevention (2017). National Diabetes Statistics Report. US Dep. Heal.

Hum. Serv.

Chatterjee, S., Khunti, K., and Davies, M.J. (2017). Type 2 diabetes. Lancet.

Chechi, K., Carpentier, A.C., and Richard, D. (2013). Understanding the brown adipocyte as a

contributor to energy homeostasis. Trends Endocrinol. Metab.

Chen, Y., Mccarthy, D., Robinson, M., and Smyth, G.K. (2015). edgeR : differential expression analysis

of digital gene expression data User's Guide.

Cheverud, J.M., Lawson, H.A., Fawcett, G.L., Wang, B., Pletscher, L.S., R Fox, A., Maxwell, T.J.,

Ehrich, T.H., Kenney-Hunt, J.P., Wolf, J.B., et al. (2011). Diet-dependent genetic and genomic

imprinting effects on obesity in mice. Obesity (Silver Spring). *19*, 160–170.

Chondronikola, M., Volpi, E., Børsheim, E., Porter, C., Annamalai, P., Enerbäck, S., Lidell, M.E., Saraf,

M.K., Labbe, S.M., Hurren, N.M., et al. (2014). Brown adipose tissue improves whole-body glucose

homeostasis and insulin sensitivity in humans. Diabetes.

Colditz, G.A., Willett, W.C., Rotnitzky, A., and Manson, J.E. (1995). Weight gain as a risk factor for

clinical diabetes mellitus in women. Ann. Intern. Med.

Cypess, A.M., Lehman, S., Williams, G., Tal, I., Rodman, D., Goldfine, A.B., Kuo, F.C., Palmer, E.L.,

Tseng, Y.-H., Doria, A., et al. (2009). Identification and importance of brown adipose tissue in adult

humans. N. Engl. J. Med. *360*, 1509–1517.

D'souza, A.M., Neumann, U.H., Glavas, M.M., and Kieffer, T.J. (2017). The glucoregulatory actions of

leptin. Mol. Metab.

Denroche, H.C., Kwon, M.M., Glavas, M.M., Tudurí, E., Philippe, M., Quong, W.L., and Kieffer, T.J.

(2016). The role of autonomic efferents and uncoupling protein 1 in the glucose-lowering effect of leptin

therapy. Mol. Metab.

Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., and

Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner. Bioinformatics *29*, 15–21.

Dulloo, A.G., and Montani, J.P. (2015). Pathways from dieting to weight regain, to obesity and to the

metabolic syndrome: An overview. Obes. Rev.

Edén, D., Siegbahn, A., and Mokhtari, D. (2015). Tissue factor/factor VIIa signalling promotes cytokine-

induced beta cell death and impairs glucose-stimulated insulin secretion from human pancreatic islets.

Diabetologia.

Ehrich, T.H., Kenney, J.P., Vaughn, T.T., Pletscher, L.S., and Cheverud, J.M. (2003). Diet, obesity, and

hyperglycemia in LG/J and SM/J mice. Obes. Res.

Flegal, K.M., Kruszon-Moran, D., Carroll, M.D., Fryar, C.D., and Ogden, C.L. (2016). Trends in Obesity

Among Adults in the United States, 2005 to 2014. JAMA *315*, 2284–2291.

Frantz, C., Stewart, K.M., and Weaver, V.M. (2010). The extracellular matrix at a glance. J. Cell Sci.

Gauger, K.J., Bassa, L.M., Henchey, E.M., Wyman, J., Bentley, B., Brown, M., Shimono, A., and

Schneider, S.S. (2013). Mice deficient in Sfrp1 exhibit increased adiposity, dysregulated glucose

metabolism, and enhanced macrophage infiltration. PLoS One.

Goossens, G.H. (2017). The Metabolic Phenotype in Obesity: Fat Mass, Body Fat Distribution, and

Adipose Tissue Function. Obes. Facts.

Gunawardana, S.C., and Piston, D.W. (2012). Reversal of type 1 diabetes in mice by brown adipose

tissue transplant. Diabetes.

Gunawardana, S.C., and Piston, D.W. (2015). Insulin-independent reversal of type 1 diabetes in

nonobese diabetic mice with brown adipose tissue transplant. Am. J. Physiol. Endocrinol. Metab. *308*,

E1043-55.

Hanssen, M.J.W., van der Lans, A.A.J.J., Brans, B., Hoeks, J., Jardon, K.M.C., Schaart, G., Mottaghy,

F.M., Schrauwen, P., van Marken Lichtenbelt, W.D., Lichtenbelt, W. van M., et al. (2016). Short-term

Cold Acclimation Recruits Brown Adipose Tissue in Obese Humans. Diabetes *65*, 1179–1189.

Hasegawa, Y., Ikeda, K., Chen, Y., Alba, D.L., Stifler, D., Shinoda, K., Hosono, T., Maretich, P., Yang,

Y., Ishigaki, Y., et al. (2018). Repression of Adipose Tissue Fibrosis through a PRDM16-GTF2IRD1

Complex Improves Systemic Glucose Homeostasis. Cell Metab.

Huang, G., Ge, G., Wang, D., Gopalakrishnan, B., Butz, D.H., Colman, R.J., Nagy, A., and Greenspan,

D.S. (2011). α3(V) Collagen is critical for glucose homeostasis in mice due to effects in pancreatic islets

and peripheral tissues. J. Clin. Invest.

Kahn, S.E., Prigeon, R.L., Schwartz, R.S., Fujimoto, W.Y., Knopp, R.H., Brunzell, J.D., and Porte, D.

(2001). Obesity, body fat distribution, insulin sensitivity and Islet beta-cell function as explanations for

metabolic diversity. J. Nutr.

Kajimura, S., Spiegelman, B.M., and Seale, P. (2015). Cell Metabolism Review Brown and Beige Fat:

Physiological Roles beyond Heat Generation. Cell Metab.

Karelis, A.D. (2008). Metabolically healthy but obese individuals. Lancet (London, England) *372*, 1281–

1283.

Kenchaiah, S., Evans, J.C., Levy, D., Wilson, P.W.F., Benjamin, E.J., Larson, M.G., Kannel, W.B., and

Vasan, R.S. (2002). Obesity and the Risk of Heart Failure. N. Engl. J. Med.

Kenney-Hunt, J.P., Vaughn, T.T., Pletscher, L.S., Peripato, A., Routman, E., Cothran, K., Durand, D.,

Norgard, E., Perel, C., and Cheverud, J.M. (2006). Quantitative trait loci for body size components in

mice. Mamm. Genome.

Kim, D., Kim, J., Yoon, J.H., Ghim, J., Yea, K., Song, P., Park, S., Lee, A., Hong, C.P., Jang, M.S., et al. (2014). CXCL12 secreted from adipose tissue recruits macrophages and induces insulin resistance in mice. Diabetologia.

Klöting, N., Fasshauer, M., Dietrich, A., Kovacs, P., Schön, M.R., Kern, M., Stumvoll, M., and Blüher, M. (2010). Insulin-sensitive obesity. Am. J. Physiol. Metab.

Lagathu, C., Christodoulides, C., Tan, C.Y., Virtue, S., Laudes, M., Campbell, M., Ishikawa, K., Ortega, F., Tinahones, F.J., Fernández-Real, J.M., et al. (2010). Secreted frizzled-related protein 1 regulates adipose tissue expansion and is dysregulated in severe obesity. Int. J. Obes.

Lapa, C., Arias-Loza, P., Hayakawa, N., Wakabayashi, H., Werner, R.A., Chen, X., Shinaji, T., Herrmann, K., Pelzer, T., and Higuchi, T. (2017). Whitening and Impaired Glucose Utilization of Brown Adipose Tissue in a Rat Model of Type 2 Diabetes Mellitus. Sci. Rep.

Lawson, H.A., and Cheverud, J.M. (2010). Metabolic syndrome components in murine models. Endocr. Metab. Immune Disord. Drug Targets *10*, 25–40.

Lawson, H. a, Zelle, K.M., Fawcett, G.L., Wang, B., Pletscher, L.S., Maxwell, T.J., Ehrich, T.H., Kenney-Hunt, J.P., Wolf, J.B., Semenkovich, C.F., et al. (2010). Genetic, epigenetic, and gene-by-diet interaction effects underlie variation in serum lipids in a LG/JxSM/J murine model. J. Lipid Res.

Lawson, H.A., Lee, A., Fawcett, G.L., Wang, B., Pletscher, L.S., Maxwell, T.J., Ehrich, T.H., Kenney-Hunt, J.P., Wolf, J.B., Semenkovich, C.F., et al. (2011a). The importance of context to the genetic architecture of diabetes-related traits is revealed in a genome-wide scan of a LG/J × SM/J murine model. Mamm. Genome *22*, 197–208.

Lawson, H.A., Cady, J.E., Partridge, C., Wolf, J.B., Semenkovich, C.F., and Cheverud, J.M. (2011b). Genetic effects at pleiotropic loci are context-dependent with consequences for the maintenance of genetic variation in populations. PLoS Genet. *7*, e1002256.

Lawson, H.A., Zayed, M., Wayhart, J.P., Fabbrini, E., Love-Gregory, L., Klein, S., and Semenkovich, C.F. (2017). Physiologic and genetic evidence links hemopexin to triglycerides in mice and humans. Int. J. Obes. *41*.

Lee, J.T., Pamir, N., Liu, N.C., Kirk, E.A., Averill, M.M., Becker, L., Larson, I., Hagman, D.K., Foster-

Schubert, K.E., Van Yserloo, B., et al. (2014). Macrophage metalloelastase (MMP12) regulates adipose

tissue expansion, insulin sensitivity, and expression of inducible nitric oxide synthase. Endocrinology.

MacArthur, J.W. (2002). Genetics of Body Size and Related Characters. I. Selecting Small and Large

Races of the Laboratory Mouse. Am. Nat.

Macias-Velasco, J., St. Pierre, C., Wayhart, J., Yin, L., Spears, L., Miranda, M., Funai, K., Cheverud, J.,

Semenkovich, C., and Lawson HA (2019). Epistatic networks associated with parent-of-origin effects on

metabolic traits. BioRxiv.

Maquoi, E., Munaut, C., Colige, A., Collen, D., and Roger Lijnen, H. (2002). Modulation of adipose

tissue expression of murine matrix metalloproteinases and their tissue inhibitors with obesity. Diabetes.

van Marken Lichtenbelt, W.D., Vanhommerig, J.W., Smulders, N.M., Drossaerts, J.M.A.F.L., Kemerink,

G.J., Bouvy, N.D., Schrauwen, P., and Teule, G.J.J. (2009a). Cold-Activated Brown Adipose Tissue in

Healthy Men. N. Engl. J. Med.

van Marken Lichtenbelt, W.D., Vanhommerig, J.W., Smulders, N.M., Drossaerts, J.M.A.F.L., Kemerink,

G.J., Bouvy, N.D., Schrauwen, P., and Teule, G.J.J. (2009b). Cold-activated brown adipose tissue in

healthy men. N. Engl. J. Med.

Meigs, J.B., Wilson, P.W.F., Fox, C.S., Vasan, R.S., Nathan, D.M., Sullivan, L.M., and D'Agostino, R.B.

(2006). Body mass index, metabolic syndrome, and risk of type 2 diabetes or cardiovascular disease. J.

Clin. Endocrinol. Metab. *91*, 2906–2912.

Mejhert, N., Wilfling, F., Esteve, D., Galitzky, J., Pellegrinelli, V., Kolditz, C.I., Viguerie, N., Tordjman, J.,

Näslund, E., Trayhurn, P., et al. (2013). Semaphorin 3C is a novel adipokine linked to extracellular

matrix composition. Diabetologia.

Moreno-Viedma, V., Amor, M., Sarabi, A., Bilban, M., Staffler, G., Zeyda, M., and Stulnig, T.M. (2016a).

Common dysregulated pathways in obese adipose tissue and atherosclerosis. Cardiovasc. Diabetol.

Moreno-Viedma, V., Amor, M., Sarabi, A., Bilban, M., Staffler, G., Zeyda, M., and Stulnig, T.M. (2016b).

Common dysregulated pathways in obese adipose tissue and atherosclerosis. Cardiovasc. Diabetol.

Nathan, D.M., Barrett-Connor, E., Crandall, J.P., Edelstein, S.L., Goldberg, R.B., Horton, E.S., Knowler,

W.C., Mather, K.J., Orchard, T.J., Pi-Sunyer, X., et al. (2015). Long-term effects of lifestyle intervention

or metformin on diabetes development and microvascular complications over 15-year follow-up: The

Diabetes Prevention Program Outcomes Study. Lancet Diabetes Endocrinol.

Nikolskiy, I., Conrad, D.F., Chun, S., Fay, J.C., Cheverud, J.M., and Lawson, H.A. (2015). Using whole-

genome sequences of the LG/J and SM/J inbred mouse strains to prioritize quantitative trait genes and

nucleotides. BMC Genomics *16*, 415.

Norgard, E.A., Lawson, H.A., Pletscher, L.S., Wang, B., Brooks, V.R., Wolf, J.B., and Cheverud, J.M.

(2011). Genetic factors and diet affect long-bone length in the F34 LG,SM advanced intercross. Mamm.

Genome *22*, 178–196.

Ouellet, V., Labbé, S.M., Blondin, D.P., Phoenix, S., Guérin, B., Haman, F., Turcotte, E.E., Richard, D.,

and Carpentier, A.C. (2012). Brown adipose tissue oxidative metabolism contributes to energy

expenditure during acute cold exposure in humans. J. Clin. Invest.

Poekes, L., Lanthier, N., and Leclercq, I.A. (2015). Brown adipose tissue: a potential target in the fight

against obesity and the metabolic syndrome. Clin. Sci.

Rauscher, G.H., Mayne, S.T., and Janerich, D.T. (2000). Relation between body mass index and lung

cancer risk in men and women never and former smokers. Am. J. Epidemiol.

Raz, V., Sterrenburg, E., Routledge, S., Venema, A., van der Sluijs, B.M., Trollet, C., Dickson, G., van

Engelen, B.G.M., van der Maarel, S.M., and Antoniou, M.N. (2013). Nuclear entrapment and

extracellular depletion of PCOLCE is associated with muscle degeneration in oculopharyngeal

muscular dystrophy. BMC Neurol.

Reeves, G.K., Pirie, K., Beral, V., Green, J., Spencer, E., and Bull, D. (2007). Cancer incidence and

mortality in relation to body mass index in the Million Women Study: Cohort study. Br. Med. J.

Roat, R., Rao, V., Doliba, N.M., Matschinsky, F.M., Tobias, J.W., Garcia, E., Ahima, R.S., and Imai, Y.

(2014). Alterations of pancreatic islet structure, metabolism and gene expression in diet-induced obese

C57BL/6J mice. PLoS One.

Roberts-Toler, C., O'Neill, B.T., and Cypess, A.M. (2015). Diet-induced obesity causes insulin

resistance in mouse brown adipose tissue. Obesity.

Rosen, E.D., and Spiegelman, B.M. (2014). What we talk about when we talk about fat. Cell.

Saito, M. (2013). Brown adipose tissue as a regulator of energy expenditure and body fat in humans.

Diabetes Metab. J.

Saito, M., Okamatsu-Ogura, Y., Matsushita, M., Watanabe, K., Yoneshiro, T., Nio-Kobayashi, J.,

Iwanaga, T., Miyagawa, M., Kameya, T., Nakada, K., et al. (2009). High incidence of metabolically

active brown adipose tissue in healthy adult humans: Effects of cold exposure and adiposity. Diabetes.

Shimizu, I., and Walsh, K. (2015). The Whitening of Brown Fat and Its Implications for Weight

Management in Obesity. Curr. Obes. Rep.

Shimizu, I., Aprahamian, T., Kikuchi, R., Shimizu, A., Papanicolaou, K.N., MacLauchlan, S., Maruyama,

S., and Walsh, K. (2014). Vascular rarefaction mediates whitening of brown fat in obesity. J. Clin.

Invest.

Shu, L., Chan, K.H.K., Zhang, G., Huan, T., Kurt, Z., Zhao, Y., Codoni, V., Trégouët, D.A., Yang, J.,

Wilson, J.G., et al. (2017). Shared genetic regulatory networks for cardiovascular disease and type 2

diabetes in multiple populations of diverse ethnicities in the United States. PLoS Genet.

Da Silva Xavier, G. (2018). The Cells of the Islets of Langerhans. J. Clin. Med.

Skrbic, B., Engebretsen, K.V.T., Strand, M.E., Lunde, I.G., Herum, K.M., Marstein, H.S., Sjaastad, I.,

Lunde, P.K., Carlson, C.R., Christensen, G., et al. (2015). Lack of collagen VIII reduces fibrosis and

promotes early mortality and cardiac dilatation in pressure overload in mice. Cardiovasc. Res.

Stanford, K.I., Middelbeek, R.J.W., Townsend, K.L., An, D., Nygaard, E.B., Hitchcox, K.M., Markan,

K.R., Nakano, K., Hirshman, M.F., Tseng, Y.-H., et al. (2013). Brown adipose tissue regulates glucose

homeostasis and insulin sensitivity. J. Clin. Invest. *123*, 215–223.

Strazzullo, P., D'Elia, L., Cairella, G., Garbagnati, F., Cappuccio, F.P., and Scalfi, L. (2010). Excess

body weight and incidence of stroke: Meta-analysis of prospective studies with 2 million participants.

Stroke.

Sun, K., Tordjman, J., Clément, K., and Scherer, P.E. (2013). Fibrosis and adipose tissue dysfunction.

Cell Metab.

Takahashi, N., Yoshizaki, T., Hiranaka, N., Kumano, O., Suzuki, T., Akanuma, M., Yui, T., Kanazawa,

K., Yoshida, M., Naito, S., et al. (2015). The production of coagulation factor VII by adipocytes is

enhanced by tumor necrosis factor-α or isoproterenol. Int. J. Obes.

Tomiyama, A.J., Ahlstrom, B., and Mann, T. (2013). Long-term Effects of Dieting: Is Weight Loss

Related to Health? Soc. Personal. Psychol. Compass.

Villarroya, F., Cereijo, R., Villarroya, J., and Giralt, M. (2017). Brown adipose tissue as a secretory

organ. Nat. Rev. Endocrinol.

Virtanen, K.A., Lidell, M.E., Orava, J., Heglind, M., Westergren, R., Niemi, T., Taittonen, M., Laine, J.,

Savisto, N.-J., Enerbäck, S., et al. (2009). Functional Brown Adipose Tissue in Healthy Adults. N. Engl.

J. Med.

Wang, G.-X., Zhao, X.-Y., Meng, Z.-X., Kern, M., Dietrich, A., Chen, Z., Cozacov, Z., Zhou, D.,

Okunade, A.L., Su, X., et al. (2014). The brown fat-enriched secreted factor Nrg4 preserves metabolic

homeostasis through attenuation of hepatic lipogenesis. Nat. Med. *20*, 1436–1443.

Wang, G.X., Zhao, X.Y., and Lin, J.D. (2015). The brown fat secretome: Metabolic functions beyond

thermogenesis. Trends Endocrinol. Metab.

Wu, J., Boström, P., Sparks, L.M., Ye, L., Choi, J.H., Giang, A.H., Khandekar, M., Virtanen, K.A.,

Nuutila, P., Schaart, G., et al. (2012). Beige adipocytes are a distinct type of thermogenic fat cell in

mouse and human. Cell.

Yang, W., Dall, T.M., Beronjia, K., Lin, J., Semilla, A.P., Chakrabarti, R., Hogan, P.F., and Petersen,

M.P. (2018). Economic costs of diabetes in the U.S. in 2017. Diabetes Care.

Zhang, B., Kirov, S., and Snoddy, J. (2005). WebGestalt: An integrated system for exploring gene sets

in various biological contexts. Nucleic Acids Res.

RNA sequencing count data available for download at: http://lawsonlab.wustl.edu/data/