Hybridization and diversification are positively correlated across vascular plant families

Nora Mitchell[1*], Kenneth D. Whitney[1]

[1]Department of Biology, University of New Mexico, Albuquerque NM, 87131 United States

[*] Present address: Department of Biology, University of Wisconsin - Eau Claire, Eau Claire, WI,

54702 United States

E-mail for correspondence: nmitchell@unm.edu; nora.c.mitchell@gmail.com

Short running title: Hybridization and diversification

Author contributions: K.D.W. and N.M. conceived of the study. N.M. performed the analyses.

N.M. and K.D.W. wrote the manuscript. All authors contributed to revisions.

Acknowledgments: We thank Jennifer Rudgers, Kellen Paine, Steven Poe, and John Wiens for

valuable discussion and feedback on this manuscript, and the UNM Center for Advanced

Research Computing for access to supercomputing resources used for the diversification

analyses. Funding for the study was provided by NSF DEB 1257965 and UNM (both to

K.D.W.).

Data accessibility statement: All data and code generated for this work will be available via the

Open Science Framework upon publication.

ABSTRACT

Hybridization has experimental and observational ties to evolutionary processes and outcomes

such as adaptation, speciation, and radiation. Although it has been hypothesized that

hybridization and diversification are correlated, this idea remains largely untested empirically.

Here, we use a hybridization database on 195 plant families, life history information, and a time-

calibrated family-level phylogeny to test for phylogenetically-corrected associations between

hybridization and diversification rates, while also taking into account life-history traits that may

be correlated with both processes. We show that diversification and hybridization are positively

correlated using three different methods to estimate diversification rates and two different

metrics of hybridization. Moreover, the relationship remains detectable when incorporating the

correlations between diversification and two other life history characteristics, perenniality and

woodiness. We further provide potential mechanisms for this association under three different

scenarios: hybridization may drive diversification, diversification may drive hybridization, or

both hybridization and diversification may jointly be driven by other factors. We suggest future

studies to disentangle the causal structure.

Keywords: adaptive radiation; extinction; perenniality; speciation; traits; woodiness

INTRODUCTION

Hybridization is a biological phenomenon whereby different species mate, typically forming

offspring that are genetically and often phenotypically distinctive. The prevalence of natural

hybridization is becoming increasingly recognized in animals (involving up to 25% of species in

some groups (Mallet 2005), fungi (reviewed in Albertin and Marullo 2012), and in plants, with

hybridization occurring in 40% of plant families (Whitney et al. 2010a) and involving up to 25%

of species (Mallet 2005).

Hybridization increases the availability of genetic variation on which natural selection can act

(Schluter et al. 2004; Barrett and Schluter 2008) and can result in novel phenotypic traits or in

novel combinations of traits, e.g. via transgressive segregation (Rieseberg et al. 1999; Bell and

Travis 2005; Dittrich-Reed and Fitzpatrick 2013). Moreover, introgressive hybridization, or

repeated backcrossing from one lineage into another, can result in the transfer of adaptive traits

between lineages that confer advantages in novel environments, explored in (Anderson 1953;

Dowling and Secor 1997; Suarez-Gonzalez et al. 2018).

Both theory and empirical observations support the idea that hybridization can promote increased

rates of evolution and/or speciation (Anderson and Stebbins 1954; Stelkens et al. 2014; Marques

et al. 2019; Taylor and Larson 2019). Hybridization has been linked to multiple evolutionary and

ecological processes, such as adaptation, e.g. (Lewontin and Birch 1966; Campbell and Snow

2007; Hovick et al. 2012; Stankowski and Streisfeld 2015; Mitchell et al. 2019a), speciation, e.g.

(Rieseberg 2003; Mallet 2007; Rieseberg et al. 2007; Soltis and Soltis 2009; Abbott et al. 2013),

and evolutionary radiation, e.g. (Anderson and Stebbins 1954; Stebbins 1959; Barton 2001;

Seehausen 2004; Yakimowski and Rieseberg 2014; Berner and Salzburger 2015; Marques et al.

2019). For example, on a microevolutionary scale, hybridization has resulted in the adaptive

introgression of traits related to both abiotic and biotic factors in Texas sunflowers (*Helianthus*

annuus ssp. *texanus*) (Whitney et al. 2006, 2010b). Experimental hybridization in this system

increased the speed of adaptive evolution over eight generations when compared to non-hybrid

controls (Mitchell et al. 2019a). Hybridization has also led to speciation within this same genus.

Historical interbreeding between two sunflowers (*Helianthus annuus* and *H. petiolaris*)

generated three species of hybrid origin that are adapted to novel and extreme environments in a

repeatable fashion (reviewed in Rieseberg et al. 2007). On a macroevolutionary scale,

hybridization has been linked to rapid speciation and evolutionary radiations. For example, in the

Hawaiian silverswords, an ancient hybrid founder may have provided the evolutionary novelty

necessary to promote adaptive radiation (Barrier et al. 1999).

Based on the association between hybridization and evolutionary change at different scales, it

has been hypothesized that hybridization may be linked to overall net diversification (Dowling

and Secor 1997; Seehausen 2004), yet this idea remains untested (but see Tank et al. 2015;

Landis et al. 2018 for related work linking polyploidy to diversification). Importantly, the

direction of causality between hybridization and diversification could go either way; high rates

of hybridization could result in high rates of diversification, or high rates of diversification could

result in the increased prevalence of hybridization within a lineage. For instance, in the latter

scenario, high rates of diversification may mean that species have low amounts of genetic

divergence between them, allowing for higher rates of hybridization.

Net diversification, the collective result of speciation minus extinction, can be estimated from

time-calibrated phylogenies (Ricklefs 2007). Diversification is an emergent property of a given

lineage and is influenced by individual, population, or species-specific factors that can increase

or decrease the likelihood of either speciation or extinction (Barraclough 1998; Langerhans and

DeWitt 2004; Bouchenak-Khelladi et al. 2015). It is therefore necessary to investigate the

relative effects of different factors on diversification in a multivariate context, while

acknowledging that additional yet unmeasured factors (and their interactive effects) are likely at

play and that the explanatory power of any one factor is likely small.

Here, we present a test of the hypothesized correlation between hybridization and diversification

in vascular plants. Both diversity (Davies et al. 2004) and rates of hybridization (Ellstrand et al.

1996; Whitney et al. 2010a; Beddows and Rose 2018; Mitchell et al. 2019b) are unevenly

distributed across plant lineages. Moreover, both diversification and hybridization have been

linked to numerous aspects of plant biology, including life form and life history. For instance,

plant groups dominated by herbaceous life forms or short generation times tend to have faster

diversification rates (Eriksson and Bremer 1992; Dodd et al. 1999), while both perenniality and

woodiness are positively associated with hybridization (Ellstrand et al. 1996; Beddows and Rose

2018; Mitchell et al. 2019b).

Using a database containing two measures of hybridization and life history information on 195

vascular plant families, as well as a time-calibrated family-level phylogeny, we estimate family-

level diversification rates using three different methods and ask 1) Are diversification rates and

hybridization rates related across vascular plants?, and 2) Does hybridization remain predictive

of diversification when accounting for other correlated traits?

MATERIALS AND METHODS

Hybridization rates, perenniality, and woodiness. For hybridization rates in vascular plant

families, as well as two life history characteristics (perenniality and woodiness) known to be

positively correlated with hybridization rates (Mitchell et al. 2019b), we used the hybrid and trait

database previously examined in Mitchell et al. (2019b). This database includes data from eight

regional floras: the Great Plains of the U.S. (McGregor and Barkley 1986) the British Isles

(Stace 1997); Hawai'i (Wagner et al. 1999); the Intermountain Region of the western U.S.

(Cronquist et al. 1972); the Northeastern U.S. (Magee and Ahles 1999); California (Hickman

1993); Europe (Tutin et al. 1964); and Victoria, Australia (Walsh and Entwisle 1994). In

summary, for each plant family in each flora the number of non-hybrids and interspecific hybrids

was assessed as in Whitney et al. (2010). For counting purposes, a "hybrid" was defined as a

hybrid type derived from a unique combination of two parental species (as in Ellstrand et al.

1996). Thus, in each flora, each pair of hybridizing species was counted as generating a single

hybrid, even if there was evidence that the pair had hybridized multiple times. Recognition of an

interspecific hybrid does not imply that it was formally or taxonomically recognized in the flora

(though some were). From these data, two metrics of hybridization (hybridization propensity and

hybrid ratio were estimated). Hybridization propensity (HybProp) is calculated as

$$Hybridization\ propensity = 100 \times \left(\frac{number\ of\ hybrids}{\frac{n(n-1)}{2}} \right) \text{ eq.1}$$

where n = the number of nonhybrid species in the family. It thus represents the percentage of

possible hybrid combinations that have been actually realized in nature. Hybrid ratio (HybRatio)

is calculated as

$$Hybrid\ ratio = \frac{number\ of\ hybrids}{n} \quad \text{eq. 2}$$

and has been used in previous studies examining patterns of hybridization (e.g. Mitchell et al.

2019b; Beddows and Rose 2018). Note the scale difference: by convention, hybridization

propensity is a percentage bounded between 0 and 100, while hybrid ratio is unbounded (in the

Mitchell et al. 2019b dataset, it ranges from 0 – 0.15 with outliers up to 1.2). Overall, the dataset

contains hybridization data on 282 plant families, and after eliminating genera where a single

non-hybrid species was observed in a single flora (thus having no opportunity for hybridization),

195 plant families were retained for subsequent analyses. See Mitchell et al. (2019b) for details

on collection of these data.

We used the values of perenniality and woodiness estimated by Mitchell et al. (2019b). Briefly,

the number of annual, biennial, and perennial species, and the number of herbaceous vs. woody

species were counted in the floras. Each species was assigned a score (for perenniality: 0 if

annual, 0.5 if biennial, 1 if perennial; for woodiness: 0 if herbaceous, 1 if characterized by

aboveground woody biomass; with intermediates split between categories). Perenniality and

woodiness were then estimated as percentages of species within a family possessing the

characteristic.

Vascular plant phylogeny. We used the family-level seed plant phylogeny of Qian and Zhang

(2014) to assess diversification rates (while incorporating family-level species richness, see

below) and account for phylogenetic non-independence of lineages. Briefly, the Qian and Zhang

(2014) phylogeny was built by extracting a single species from each family from the Zanne et al.

(2014) species-level phylogeny, updating the taxonomy, correcting errors, and including six

additional families to produce a phylogeny with tips representing 437 seed plant families with

branch lengths proportional to time in millions of years. The phylogeny was downloaded from

Appendix S3 of Qian and Zhang (2014). We checked taxonomy according to The Plant List

(<www.theplantlist.org> , as of March 2019) and trimmed families from the phylogeny that are

no longer recognized, resulting in a seed plant phylogeny of 413 families. To incorporate ferns

and their allies, we used the time-calibrated phylogeny of Testo and Sundue (2016) .We trimmed

the fern phylogeny to a random species from each family and bound the seed plant phylogeny to

the family-level fern phylogeny. We then ultrametricized the product using the

force.ultrametric() function and the "nnls" method in the R package phytools (Revell 2012),

resulting in a vascular plant phylogeny with 459 families. The phylogeny and additional metrics

were visualized using the R package ggtree (Yu et al. 2017).

Species richness. We created a taxonomic richness matrix based on the number of species in

each family listed in The Plant List (<www.theplantlist.org> , as of March 2019).

Diversification rates. We used the family-level phylogeny to assess diversification rates using

three different methods: Method-of-moments estimation, Medusa, and BAMM.

Method-of-moments (MS). We used the stem-group method-of-moments estimation (MS) of

diversification using the stem ages of each family (tip edge lengths from the family-level

phylogeny) and family species richness to calculate a measure of net diversification rates

(Rohatgi 1976; Magallon and Sanderson 2001). We calculated MS using three different relative

extinction rates (epsilon, ϵ = 0, 0.5, or 0.9) and the equation $r = \ln(n (1 - \epsilon) + \epsilon)/t$, where r =

the net diversification rate, n = the extant species richness and t = the stem group age (in millions

of years).

Medusa. We used Medusa (Alfaro et al. 2009) to model diversification using stepwise AIC using

piecewise birth-death models. We used the medusa() function in the R package geiger (Harmon

et al. 2008) using the family-level phylogeny and family species richness. The program

compares stepwise models until the improvement, measured using the sample-size corrected

Akaike information criterion (AICc), does not exceed the internally computed threshold (for a

phylogeny with 459 tips, threshold = 8.089111). Because results are highly dependent on

threshold value (May and Moore 2016), we ran Medusa using several threshold values (8.089, 4,

and 2) to compare outputs; note that a previous version of Medusa used a default threshold value

of 4. We extracted the net diversification rates associated with each tip (family) from each of

these threshold runs.

BAMM. We used BAMM (Bayesian Analysis of Macroevolutionary Mixture Models) (Rabosky

2014) to model speciation and extinction rates using reversible jump Markov chain Monte Carlo

to explore model space. Prior values were obtained using the BAMMtools R package (Rabosky

et al. 2014). We tested four values for the expected number of shifts (25, 50, 100, and 200), and

accounted for richness by providing a sampling fraction for each family calculated as 1 over the

number of species in the family. To account for the fact that the "sampling fraction" was often

extremely low, we set extinctionProbMax = 0.9999999. We ran BAMM on the family-level seed

plant phylogeny on the high-performance computing facilities maintained by the UNM Center

for Advanced Research Computing for 50 million generations sampled every thousand

generations. We processed the output in BAMMtools and used a burnin of 20%, leaving 40,001

samples. Convergence was checked visually using the R package coda (Plummer et al. 2006); we

ensured that the effective sample sizes for the number of shifts and log-likelihoods were over

200. We then extracted net diversification rates for each family.

Phylogenetic Generalized Least Squares (PGLS). We used phylogenetic generalized least

squares regression (PGLS) (Grafen 1989; Martins and Garland Jr 1991) to detect associations

between diversification rates and rates of hybridization in plant families while accounting for

evolutionary history. We trimmed the family-level phylogeny to include only the families for

which we had hybridization data, resulting in a phylogeny of 195 families. We standardized all

variables to make results comparable between analyses. We ran univariate PGLS using the pgls()

function in the R package caper (Orme et al. 2013), modeling diversification rates as outcomes

and using each hybridization metric as our predictor with estimated lambda values and kappa and

delta set to 1.

For each combination of diversification and hybridization metrics, we also ran multivariate

regressions using standardized data to detect associations between diversification rates and

hybridization metrics when including two other predictors: perenniality and woodiness.

Sister-clade comparison. We further examined the relationship between hybridization and

diversification using the more traditional approach of sister-clade comparisons (Slowinski and

Guyer 1993; Barraclough 1998) to ask whether families with higher rates of hybridization also

had higher diversification rates. We identified sister families present in the 459-taxon phylogeny

and consistent with relationships on the Angiosperm Phylogeny Website

<http://www.mobot.org/MOBOT/research/APweb/, accessed 23 April 2019>. We restricted

comparisons to sister pairs for which we had data and where both hybridization propensity and

hybridization ratio differed consistently (i.e., the same family had higher values for both

metrics), as there were some cases where the metrics were inconsistent in identifying the more

hybridization-prone family. This resulted in 11 sister-clade comparisons. We then asked whether

the diversification rates differed between families in the same direction that hybridization

differed using a non-parametric one-tailed sign-test.

RESULTS

Hybridization rates. Hybridization rates across families in this dataset have been presented and

discussed previously in Mitchell et al. (2019b). Briefly, out of 195 plant families, 112 contained

hybrids. The average family-level hybridization propensity was 2.55 and the average

hybridization ratio was 0.086. See Mitchell et al. (2019b) Fig. 2 and Table S2 for details.

Diversification rates. We used three different methods to estimate diversification rates across

the family-level vascular plant phylogeny: method-of-moments (MS), Medusa, and BAMM.

We estimated the MS net diversification rates using relative extinction fractions of $\square = 0.0, 0.5,$

and 0.9. Although we are unable to use model comparison since the response measures are

different, diversification rates were highly correlated across values of ☐ (Table S1). We chose to

examine rates computed with the moderate value of ☐ = 0.5 for presentation in the main

analyses. The mean net diversification rate was 0.050 species per Ma (range = 0.000 – 0.255)

across all 459 families in the phylogeny; the mean rate was 0.077 species per Ma across the 195

families for which we had hybridization data.

Using the computed AICc threshold value of 8.089 for our 459-taxon phylogeny, Medusa

detected 48 rate shifts in diversification, while the previously standard threshold value of four

detected 82 rate shifts and a threshold of two detected 104 rate shifts. Rate shifts for Medusa runs

using different thresholds were highly correlated (Table S2). We chose to examine rates

computed using the threshold of four for presentation in the main results. The mean net

diversification rate was 0.072 species per Ma (range = 0.000 – 0.280) across all 459 families in

the phylogeny and the mean rate was 0.088 species per Ma across the 195 families for which we

had hybridization data.

The number of rate shifts detected by BAMM was largely robust to the prior on the expected

number of shifts, where the priors of 25, 50, 100, and 200 resulted in 74.2, 77.5, 79.8, and 79.8

mean rate shifts, respectively (Fig. S1). The diversification rates estimated using the four

different priors are highly correlated with each other (Table S3). We chose to examine rates

estimated using a prior of 100 expected shifts for presentation in the main results, as the posterior

estimate was consistent between values of 100 and 200 and differed slightly between 50 and 100.

The mean net diversification rate was 0.044 species per Ma (range = -0.069 – 0.162) across all

459 families in the phylogeny and the mean rate was 0.057 species per Ma across the 195

families for which we had hybridization data.

Net diversification rates across the three different methods were highly correlated across the full

459-taxon dataset (MS – Medusa: r = 0.808, p = 0.000; MS – BAMM: r = 0.817, p = 0.000;

Medusa – BAMM: r = 0.878, p = 0.000) (Figs. 1, 2).

Some evidence for phylogenetic correlation between hybridization and diversification.

Univariate phylogenetically generalized least squares (PGLS) regressions of hybridization

measures on diversification rates were all positive, meaning that families with higher

hybridization rates also had higher diversification rates when accounting for phylogenetic

nonindependence (Table 1, Fig. 3A). Using HybProp, hybridization was weakly associated (0.05

$< p < 0.10$) with increased diversification rates estimated using all three methods. Using

HybRatio, hybridization was positively associated with the MS and Medusa diversification rates

($p < 0.05$) and hybridization was weakly associated with the BAMM diversification rates (0.05

$< p < 0.10$) (Table 1, Fig. 3A). The relationships were weak (see adjusted-R^2 values in Table 1),

though the results were largely robust to the use of alternate parameters, thresholds, or priors for

estimating diversification rates (see Tables S4, S5, S6 for details of supplementary modeling

results).

Correlations between hybridization and diversification are not driven by perenniality or

woodiness. Multivariate PGLS regressions incorporating family-level perenniality and

woodiness measures reinforced the trends observed in the univariate analyses. Relationships

between diversification rates and hybridization measures were consistently positive, though the

models overall had very low R^2 values (Table 2). In the combination of HybProp and BAMM-

estimated diversification, the hybridization – diversification relationship was positive and

significant at $p < 0.05$; but when diversification was estimated by MS or Medusa, the

hybridization – diversification relationships were weaker but still trended positive ($0.05 < p <$

0.10). All three diversification measures were significantly positively associated with HybRatio

($p < 0.05$) (Table 2, Fig. 3B).

Diversification rates tended to be negatively associated with woodiness (Table 2, Fig. 3B), but

the relationships were only marginally significant ($0.05 < p < 0.10$) using the MS and BAMM

diversification measures. Relationships between woodiness and Medusa diversification rates, and

relationships between perenniality and all diversification rates, were not significant. The

standardized partial regression coefficients were similar in magnitude for hybridization metrics

and woodiness, while the effects of perenniality were much lower and non-significant (Table 2,

Fig. 3B). The best models (as measured by the corrected Akaike Information Criterion, AICc)

included either hybridization alone or both hybridization and woodiness, although multiple

models fell within two AICc of the best model (Table S7). Multivariate results using alternate

parameters, threshold, and priors for estimating diversification rates were qualitatively similar

(results not shown).

For completeness, we also examined correlations between hybridization and diversification

without accounting for phylogenetic nonindependence. None of these raw correlations were

significant, although Medusa diversification rates and HybProp had a trend toward being

negative (0.05 < p < 0.10) (Table S8). The estimated relationships with HybProp were all

negative, and for HybRatio were all positive, which differed from both the univariate and

multivariate PGLS analyses (Fig. 3, Table 1, Table 2), demonstrating the need for phylogenetic

context.

Sister-clade comparisons support a positive hybridization - diversification relationship. We

identified 11 pairs of sister families in our dataset with consistent differences in hybridization

rates (Table S9). There were no differences in diversification rates estimated by Medusa in any

of these pairs (within each pair, Medusa estimated the same net diversification rate for each

family), consistent with its recognition of far fewer rate shifts across the phylogeny relative to

the other methods. For diversification estimates using both MS and BAMM, there were nine

cases where the diversification rates were higher for the family with higher hybridization rates

and two cases where the diversification rates were lower for the family with higher hybridization

rates. This result was significant using a non-parametric one-tailed sign-test (n = 11, p = 0.033).

DISCUSSION

Hybridization and diversification are correlated, but weakly so. The idea that hybridization

can promote processes such as adaptation, speciation, and radiation has long been hypothesized,

but direct links between hybridization and diversification have been largely untested. Here, we

find a positive association between hybridization rates and diversification rates among vascular

plant families using three measures of diversification and two measures of hybridization, though

these relationships were not always statistically significant, and when significant, models

accounted for only a small amount of the observed variation in diversification rates (Fig. 3A,

Table 1). Sister-clade comparisons also supported the idea that plant families exhibiting more

hybridization tend to diversify faster, as the family with higher hybridization rates also had more

rapid diversification (estimated by MS and BAMM) than its sister family in nine of our sample

of eleven comparisons (Table S9).

Potential explanations for relationship between hybridization and diversification. The

correlative associations presented here do not necessarily imply that hybridization causes

increased diversification. The relationship could be due to causation in either direction or due to

correlation with an additional unmeasured or untested variable. Here, we discuss potential

mechanisms underlying the association, emphasizing that these mechanisms are speculative and

cannot be tested with our observational data.

Scenario 1: Increased hybridization leads to faster diversification. Interbreeding between species

results in increased genetic and phenotypic diversity which can promote adaptation and

speciation. Hybridization can lead to increased genetic diversity (Stebbins 1950, 1959; Hedrick

2013; Suarez-Gonzalez et al. 2018; Marques et al. 2019) and trait diversity, e.g. transgressive

segregation (Rieseberg et al. 1999; Seehausen 2004; Kerbs et al. 2017). Hybridization has been

shown experimentally to speed up adaptive evolution in annual sunflowers (Mitchell et al.

2019a) and can result in hybrid speciation (for instance in sunflowers, reviewed in Rieseberg

2006) and the Louisiana irises (Arnold et al. 2012), reviewed more broadly in (Mallet 2007;

Soltis and Soltis 2009; Arnold et al. 2012; Abbott et al. 2013; Marques et al. 2019). Marques et

al. (2019) recently synthesized the idea that the reassembly of old genetic variation into new

combinations can provide a substrate for speciation. In contrast to novel mutations arising within

a lineage, alleles incorporated from other lineages have previously been tested in other genomic

and ecological backgrounds (Mitchell et al. 2019a), and thus may be more likely to promote

adaptation. Such adaptation in turn could reduce the chance for extinction and/or promote

speciation.

It has also been postulated that hybridization events can lead to adaptive radiation, broadly

defined as "the evolution of ecological and phenotypic diversity within a rapidly multiplying

lineage", via similar mechanisms (Seehausen 2004, 2013). For instance, Barrier et al. (1999)

infer that the founder of the Hawaiian silversword alliance (an iconic plant radiation) was a

hybrid and suggest that enhanced genetic diversity in the founder may have promoted adaptive

radiation. Additionally, hybridization can facilitate ecological speciation by providing increased

genetic variation and multiple genotypes that can fill available niches, allowing for colonization

of novel environments or habitats (Schluter 2000; Seehausen 2004).

In contrast, there is evidence that hybridization could act to decrease diversification. In certain

scenarios, hybridization can result in "extinction-by-hybridization" (Campbell et al. In Press;

Rhymer and Simberloff 1996; Wolf et al. 2010; Todesco et al. 2016) in which the genetic

material of one species is replaced by that of another. At present little data exists to assess how

common this process is relative to the above-described processes by which hybridization could

enhance diversification.

Scenario 2: Faster diversification leads to increased hybridization. Rapid speciation within a

lineage results in many closely related species and may also result in incomplete reproductive

isolation (e.g., Baldwin and Sanderson 1998; Barrier et al. 1999). Most hybridization takes place

within genera (Whitney et al. 2010a), so if fast diversification rates result in many congeneric

species then there is enhanced opportunity for hybridization due to the number of potential

partners and possibly their geographic proximity. Additionally, taxa that are more closely related

have weaker reproductive barriers than distantly related taxa (Coyne and Orr 1989, 1997; Moyle

and Nakazato 2010), and these reduced barriers should result in increased interspecific mating

(Ellstrand et al. 1996).

Scenario 3: Diversification and hybridization are jointly driven by other factors not tested or

described here. A correlation between two variables can arise in the absence of a direct causal

relationship between them, if they are both driven by a third factor or set of factors. We discuss

two of these potential factors (woodiness and perenniality) below.

Hybridization, diversification, and life history/life form. Previous work has linked both

hybridization and diversification to other aspects of plant biology, such as life form and life

history. Plant groups with woody growth forms and perennial life histories tend to contain more

hybrids (Grant 1958, 1981; Stebbins 1959; Ellstrand et al. 1996; Beddows and Rose 2018;

Mitchell et al. 2019b), while plant groups with herbaceous life forms and faster generation times

(e.g., annual life histories) have faster diversification (Eriksson and Bremer 1992; Baker et al.

2014) or speciation rates (Dodd et al. 1999). We incorporated perenniality and woodiness into

multivariate analyses to ask whether hybridization still had a positive association with

diversification rates when these traits are included, and to compare the effect sizes of these

potential correlates. We found that hybridization remains positively associated with

diversification in these analyses, while woodiness is generally negatively associated with

diversification and perenniality had little to no effect (Fig. 3B, Table 2, Table S7). The best

model always contained hybridization (Table S7) and the effect of hybridization was always

positive and at least weakly significant (Fig. 3B, Table 2). Moreover, the association between

hybridization and diversification (measured as the standardized partial regression coefficients)

was equivalent to the magnitude of the woodiness – diversification associations, and was

stronger than the perenniality – diversification association (Fig. 3B, Table 2). The relationships

between woodiness and diversification were in the expected direction (woody groups tended to

have lower diversification rates). Interestingly, the best models did not include perenniality

(Table S7) and perenniality was not significant (Fig. 3B, Table 2), contrary to expectations

(Eriksson and Bremer 1992; Dodd et al. 1999).

Other correlates of diversification. Although we find detectable associations between

hybridization and diversification, our models explain only a small amount of the variation in

diversification rates among plants (low adjusted-R^2 values, Table 1, Table 2). Diversification is

an integrative property of a lineage, which is influenced by multiple dimensions of life history,

environmental factors, and physiology of the organisms within a lineage. Diversification rates or

species richness in plants have been tied to numerous attributes (in addition to life history).

These include aspects of the genome such as polyploidy (Vamosi and Dickinson 2006; Wood et

al. 2009; Tank et al. 2015; Landis et al. 2018) or genome size (Puttick et al. 2015); dispersal

mode or geography such as biotic vs. abiotic dispersal (Larson-Johnson 2016), latitudinal

gradients (Davies et al. 2004; Jansson and Davies 2008), biome or habitat differences (Moore

and Donoghue 2007; Valente et al. 2010; Goldberg et al. 2011; Onstein et al. 2016), geographic

area (Vamosi and Vamosi 2010, 2011), and geographic isolation (Baldwin and Sanderson 1998);

species interactions such as defense mutualisms (Weber and Agrawal 2014) and pollination

mode (Eriksson and Bremer 1992; Hodges and Arnold 1995; Hodges 1997; Dodd et al. 1999;

Vamosi and Vamosi 2010, 2011); and other reproductive aspects such as heterostyly (de Vos et

al. 2014), self-incompatibility (Goldberg et al. 2010), and dioecy (Käfer et al. 2014). Future

multivariate analyses or phylogenetic path analyses (van der Bijl 2018) with more complete

datasets are needed to disentangle and weight the relative contributions of these traits, hopefully

resulting in models with greater explanatory power.

Choice of hybridization and diversification metrics. Our two measures of hybridization

(HybProp and HybRatio) and three methods used to estimate diversification rates (MS, Medusa,

and BAMM) yield very similar results. Previous work on the correlates of hybridization also

found similar results for the two hybridization metrics (Beddows and Rose 2018; Mitchell et al.

2019b). Our results with hybridization propensity and hybrid ratio, though analyses using hybrid

ratio tend to show slightly stronger relationships with diversification (Fig. 3, Table 1, Table 2).

There is much debate surrounding the methods of estimation of diversification rates (May and

Moore 2016; Moore et al. 2016; Rabosky et al. 2017; Meyer and Wiens 2018; Meyer et al. 2018;

Rabosky 2018). For our purposes, the tight correlations among the methods (Fig. 2) and

consistency in results across methods (Fig. 3, Table 1, Table 2) suggest a robust relationship

between hybridization and diversification. The differences in significance in these methods may

be due to the fact that both BAMM and Medusa estimate rate shifts across the phylogeny (Alfaro

et al. 2009; Rabosky 2014), meaning that multiple closely-related families can share a

diversification rate, while the method-of-moments (MS) estimation is based on branch lengths

and richness (Magallon and Sanderson 2001), resulting in a uniquely-estimated diversification

rate for each family. Our results suggest that the choice of method to estimate diversification is

not crucial to our understanding of the overall direction of relationships between hybridization

and diversification, but is important in determining whether the relationship is statistically

significant.

Conclusions. Here, we present evidence that hybridization and diversification rates are

positively related across vascular plant families. Although the explanatory power of

hybridization is weak, the relationship is detectable even when other factors (perenniality,

woodiness) collinear with hybridization rates and previously linked to diversification rates are

accounted for. The magnitude of the correlation between diversification and hybridization is on

par with that of woodiness and greater than that of perenniality. Although we cannot determine

the directionality of causation via data at this taxonomic scale, our results are consistent with

experimental evidence indicating that hybridization has the potential to speed up adaptive

evolution (Mitchell et al. 2019a), so hybridization may enable more rapid diversification. More

likely, hybridization and diversification may have positive effects on each other through a variety

of mechanisms. The use of hypothesis-driven phylogenetic path analysis (van der Bijl 2018) may

be one way to disentangle causality and incorporate both direct and indirect mechanisms related

to diversification rates overall. Detailed evidence suggests that hybridization events may trigger

evolutionary radiations, such as in the Hawaiian silverswords (Barrier et al. 1999), Galapagos

finches (Grant et al. 2005), Hawaiian crickets (Shaw 2002) or African cichlids (Genner and

Turner 2012; Keller et al. 2013; Meier et al. 2017). Exploration of hybridization at the base of

additional radiations and incorporation of ancient or cryptic hybridization events are needed to

provide further confirmation of this hypothesis. Moreover, a deeper theoretical, empirical,

model-based, and experimental knowledge of hybridization, diversification, other factors, and

the interplay between them is necessary to more fully understand these evolutionary phenomena.

REFERENCES

Abbott, R., D. Albach, S. Ansell, J. W. Arntzen, S. J. E. Baird, N. Bierne, J. Boughman, A.

Brelsford, C. A. Buerkle, R. Buggs, R. K. Butlin, U. Dieckmann, F. Eroukhmanoff, A.

Grill, S. H. Cahan, J. S. Hermansen, G. Hewitt, A. G. Hudson, C. Jiggins, J. Jones, B.

Keller, T. Marczewski, J. Mallet, P. Martinez-Rodriguez, M. Möst, S. Mullen, R.

Nichols, A. W. Nolte, C. Parisod, K. Pfennig, A. M. Rice, M. G. Ritchie, B. Seifert, C.

M. Smadja, R. Stelkens, J. M. Szymura, R. Väinölä, J. B. W. Wolf, and D. Zinner. 2013.

Hybridization and speciation. Journal of Evolutionary Biology 26:229–246.

Albertin, W., and P. Marullo. 2012. Polyploidy in fungi: evolution after whole-genome

duplication. Proceedings of the Royal Society of London B: Biological Sciences

rspb20120434.

Alfaro, M. E., F. Santini, C. Brock, H. Alamillo, A. Dornburg, D. L. Rabosky, G. Carnevale, and

L. J. Harmon. 2009. Nine exceptional radiations plus high turnover explain species

diversity in jawed vertebrates. Proceedings of the National Academy of Sciences

106:13410–13414.

Anderson, E. 1953. Introgressive hybridization. Biological Reviews 28:280–307.

Anderson, E., and G. L. Stebbins. 1954. Hybridization as an evolutionary stimulus. Evolution

8:378–388.

Arnold, M. L., E. S. Ballerini, and A. N. Brothers. 2012. Hybrid fitness, adaptation and

evolutionary diversification: lessons learned from Louisiana Irises. Heredity 108:159.

Baker, T. R., R. T. Pennington, S. Magallon, E. Gloor, W. F. Laurance, M. Alexiades, E.

Alvarez, A. Araujo, E. J. Arets, and G. Aymard. 2014. Fast demographic traits promote

high diversification rates of Amazonian trees. Ecology Letters 17:527–536.

Baldwin, B. G., and M. J. Sanderson. 1998. Age and rate of diversification of the Hawaiian

silversword alliance (Compositae). Proceedings of the National Academy of Sciences

95:9402–9406.

Barraclough, T. G. 1998. Sister-group analysis in identifying correlates of diversification. Evol

Ecol 12:751–754.

Barrett, R. D. H., and D. Schluter. 2008. Adaptation from standing genetic variation. Trends in

Ecology & Evolution 23:38–44.

Barrier, M., B. G. Baldwin, R. H. Robichaux, and M. D. Purugganan. 1999. Interspecific hybrid

ancestry of a plant adaptive radiation: allopolyploidy of the Hawaiian silversword

alliance (Asteraceae) inferred from floral homeotic gene duplications. Mol Biol Evol

16:1105–1113.

Barton, N. H. 2001. The role of hybridization in evolution. Mol. Ecol. 10:551–568.

Beddows, I., and L. E. Rose. 2018. Factors determining hybridization rate in plants: A case study

in Michigan. Perspectives in Plant Ecology, Evolution and Systematics 34:51–60.

Bell, M. A., and M. P. Travis. 2005. Hybridization, transgressive segregation, genetic

covariation, and adaptive radiation. Trends in Ecology & Evolution 20:358–361.

Berner, D., and W. Salzburger. 2015. The genomics of organismal diversification illuminated by

adaptive radiations. Trends in Genetics 31:491–499.

Bouchenak-Khelladi, Y., R. E. Onstein, Y. Xing, O. Schwery, and H. P. Linder. 2015. On the

complexity of triggering evolutionary radiations. New Phytologist 207:313–326.

Campbell, L. G., C. M. Blanchette, and E. Small. In Press. Gene flow and introgression from

pharmaceutical plants into their wild relatives. Botanical Review.

Campbell, L. G., and A. A. Snow. 2007. Competition alters life history and increases the relative

fecundity of crop–wild radish hybrids (*Raphanus* spp.). New Phytologist 173:648–660.

Coyne, J. A., and H. A. Orr. 1989. Patterns of speciation in Drosophila. Evolution 362–381.

Coyne, J. A., and H. A. Orr. 1997. "Patterns of speciation in *Drosophila*" revisited. Evolution

51:295–303.

Cronquist, A., S. Holmgren, N. Holmgren, J. Reveal, and P. Holmgren. 1972. Intermountain

Flora vols. 1, 2B, 3A, 3B, 4, 5, 6. Hafner, New York.

Davies, T. J., T. G. Barraclough, M. W. Chase, P. S. Soltis, D. E. Soltis, and V. Savolainen.

2004. Darwin's abominable mystery: insights from a supertree of the angiosperms.

Proceedings of the National Academy of Sciences 101:1904–1909.

de Vos, J. M., C. E. Hughes, G. M. Schneeweiss, B. R. Moore, and E. Conti. 2014. Heterostyly

accelerates diversification via reduced extinction in primroses. Proceedings of the Royal

Society B: Biological Sciences 281:20140075.

Dittrich-Reed, D. R., and B. M. Fitzpatrick. 2013. Transgressive hybrids as hopeful monsters.

Evolutionary biology 1–6.

Dodd, M. E., J. Silvertown, and M. W. Chase. 1999. Phylogenetic analysis of trait evolution and

species diversity variation among angiosperm families. Evolution 53:732–744.

Dowling, T. E., and C. L. Secor. 1997. The role of hybridization and introgression in the

diversification of animals. Annual review of Ecology and Systematics 28:593–619.

Ellstrand, N. C., R. Whitkus, and L. H. Rieseberg. 1996. Distribution of spontaneous plant

hybrids. Proceedings of the National Academy of Sciences 93:5090–5093.

Eriksson, O., and B. Bremer. 1992. Pollination systems, dispersal modes, life forms, and

diversification rates in angiosperm families. Evolution 46:258–266.

Genner, M. J., and G. F. Turner. 2012. Ancient hybridization and phenotypic novelty within Lake Malawi's cichlid fish radiation. Mol Biol Evol 29:195–206.

Goldberg, E. E., J. R. Kohn, R. Lande, K. A. Robertson, S. A. Smith, and B. Igić. 2010. Species selection maintains self-incompatibility. Science 330:493–495.

Goldberg, E. E., L. T. Lancaster, and R. H. Ree. 2011. Phylogenetic inference of reciprocal effects between geographic range evolution and diversification. Systematic Biology 60:451–465.

Grafen, A. 1989. The phylogenetic regression. Philosophical Transactions of the Royal Society of London. B, Biological Sciences 326:119–157.

Grant, P. R., B. R. Grant, and K. Petren. 2005. Hybridization in the recent past. The American Naturalist 166:56–67.

Grant, V. 1981. Plant speciation. Columbia University Press,.

Grant, V. 1958. The regulation of recombination in plants. Cold Spring Harb Symp Quant Biol 23:337–363.

Harmon, L. J., J. T. Weir, C. D. Brock, R. E. Glor, and W. Challenger. 2008. GEIGER: investigating evolutionary radiations. Bioinformatics 24:129–131.

Hedrick, P. W. 2013. Adaptive introgression in animals: examples and comparison to new mutation and standing variation as sources of adaptive variation. Molecular Ecology 22:4606–4618.

Hickman, J. C. 1993. The Jepson Manual. University of California Press, Berkeley, CA.

Hodges, S. A. 1997. Floral nectar spurs and diversification. International Journal of Plant Sciences 158:S81–S88.

Hodges, S. A., and M. L. Arnold. 1995. Spurring plant diversification: are floral nectar spurs a

key innovation? Proceedings of the Royal Society of London B: Biological Sciences

262:343–348.

Hovick, S. M., L. G. Campbell, A. A. Snow, and K. D. Whitney. 2012. Hybridization alters early

life-history traits and increases plant colonization success in a novel region. The

American Naturalist 179:192–203.

Jansson, R., and T. J. Davies. 2008. Global variation in diversification rates of flowering plants:

energy vs. climate change. Ecology Letters 11:173–183.

Käfer, J., H. J. de Boer, S. Mousset, A. Kool, M. Dufay, and G. A. B. Marais. 2014. Dioecy is

associated with higher diversification rates in flowering plants. Journal of evolutionary

biology 27:1478–1490.

Keller, I., C. E. Wagner, L. Greuter, S. Mwaiko, O. M. Selz, A. Sivasundar, S. Wittwer, and O.

Seehausen. 2013. Population genomic signatures of divergent adaptation, gene flow and

hybrid speciation in the rapid radiation of Lake Victoria cichlid fishes. Molecular ecology

22:2848–2863.

Kerbs, B., J. Ressler, J. K. Kelly, M. E. Mort, A. Santos-Guerra, M. J. S. Gibson, J. Caujapé-

Castells, and D. J. Crawford. 2017. The potential role of hybridization in diversification

and speciation in an insular plant lineage: insights from synthetic interspecific hybrids.

AoB Plants 9.

Landis, J. B., D. E. Soltis, Z. Li, H. E. Marx, M. S. Barker, D. C. Tank, and P. S. Soltis. 2018.

Impact of whole-genome duplication events on diversification rates in angiosperms.

American Journal of Botany 105:348–363.

Langerhans, R. B., and T. J. DeWitt. 2004. Shared and unique features of evolutionary

diversification. The American Naturalist 164:335–349.

Larson-Johnson, K. 2016. Phylogenetic investigation of the complex evolutionary history of

dispersal mode and diversification rates across living and fossil Fagales. New Phytologist

209:418–435.

Lewontin, R. C., and L. C. Birch. 1966. Hybridization as a source of variation for adaptation to

new environments. Evolution 20:315–336.

Magallon, S., and M. J. Sanderson. 2001. Absolute diversification rates in angiosperm clades.

Evolution 55:1762–1780.

Magee, D. W., and H. E. Ahles. 1999. Flora of the northeast: a manual of the vascular flora of

New England and adjacent New York. Univ of Massachusetts Press.

Mallet, J. 2007. Hybrid speciation. Nature 446:279–283.

Mallet, J. 2005. Hybridization as an invasion of the genome. Trends in Ecology & Evolution

20:229–237.

Marques, D. A., J. I. Meier, and O. Seehausen. 2019. A combinatorial view on speciation and

adaptive radiation. Trends in Ecology & Evolution 34:531–544.

Martins, E. P., and T. Garland Jr. 1991. Phylogenetic analyses of the correlated evolution of

continuous characters: a simulation study. Evolution 45:534–557.

May, M. R., and B. R. Moore. 2016. How well can we detect lineage-specific diversification-rate

shifts? A simulation study of sequential AIC methods. Systematic Biology 65:1076–

1084.

McGregor, R. L., and T. M. Barkley. 1986. Flora of the great plains. University Press of Kansas.

Meier, J. I., D. A. Marques, S. Mwaiko, C. E. Wagner, L. Excoffier, and O. Seehausen. 2017.

Ancient hybridization fuels rapid cichlid fish adaptive radiations. Nature communications

8:14363.

Meyer, A. L., C. Román-Palacios, and J. J. Wiens. 2018. BAMM gives misleading rate estimates

in simulated and empirical datasets. Evolution 72:2257–2266.

Meyer, A. L., and J. J. Wiens. 2018. Estimating diversification rates for higher taxa: BAMM can

give problematic estimates of rates and rate shifts. Evolution 72:39–53.

Mitchell, N., G. L. Owens, S. M. Hovick, L. H. Rieseberg, and K. D. Whitney. 2019a.

Hybridization speeds adaptive evolution in an eight-year field experiment. Scientific

reports 9:6746.

Mitchell, N., L. G. Campbell, J. R. Ahern, K. C. Paine, A. B. Giroldo, and K. D. Whitney.

2019b. Correlates of hybridization in plants. BioRxiv:726323.

Moore, B. R., and M. J. Donoghue. 2007. Correlates of diversification in the plant clade

Dipsacales: geographic movement and evolutionary innovations. The American

Naturalist 170:S28–S55.

Moore, B. R., S. Höhna, M. R. May, B. Rannala, and J. P. Huelsenbeck. 2016. Critically

evaluating the theory and performance of Bayesian analysis of macroevolutionary

mixtures. Proceedings of the National Academy of Sciences 113:9569–9574.

Moyle, L. C., and T. Nakazato. 2010. Hybrid incompatibility "snowballs" between *Solanum*

species. Science 329:1521–1523.

Onstein, R. E., G. J. Jordan, H. Sauquet, P. H. Weston, Y. Bouchenak-Khelladi, R. J. Carpenter,

and H. P. Linder. 2016. Evolutionary radiations of Proteaceae are triggered by the

interaction between traits and climates in open habitats. Global Ecology and

Biogeography, doi: 10.1111/geb.12481.

Orme, D., R. Freckleton, G. Thomas, and T. Petzoldt. 2013. The caper package: comparative

analysis of phylogenetics and evolution in R. R package version 5:1–36.

Plummer, M., N. Best, K. Cowles, and K. Vines. 2006. CODA: Convergence diagnosis and

output analysis for MCMC. R news 6:7–11.

Puttick, M. N., J. Clark, and P. C. J. Donoghue. 2015. Size is not everything: rates of genome

size evolution, not C-value, correlate with speciation in angiosperms. Proceedings of the

Royal Society B: Biological Sciences 282:20152289.

Qian, H., and J. Zhang. 2014. Using an updated time-calibrated family-level phylogeny of seed

plants to test for non-random patterns of life forms across the phylogeny. Journal of

Systematics and Evolution 52:423–430.

Rabosky, D. L. 2014. Automatic detection of key innovations, rate shifts, and diversity-

dependence on phylogenetic trees. PloS one 9:e89543.

Rabosky, D. L. 2018. BAMM at the court of false equivalency: a response to Meyer and Wiens.

Evolution 72:2246–2256.

Rabosky, D. L., M. Grundler, C. Anderson, P. Title, J. J. Shi, J. W. Brown, H. Huang, and J. G.

Larson. 2014. BAMM tools: an R package for the analysis of evolutionary dynamics on

phylogenetic trees. Methods in Ecology and Evolution 5:701–707.

Rabosky, D. L., J. S. Mitchell, and J. Chang. 2017. Is BAMM flawed? Theoretical and practical

concerns in the analysis of multi-rate diversification models. Systematic biology 66:477–

498.

Revell, L. J. 2012. phytools: an R package for phylogenetic comparative biology (and other

things). Methods in Ecology and Evolution 3:217–223.

Rhymer, J. M., and D. Simberloff. 1996. Extinction by hybridization and introgression. Annual

review of ecology and systematics 27:83–109.

Ricklefs, R. E. 2007. Estimating diversification rates from phylogenetic information. Trends in

Ecology & Evolution 22:601–610.

Rieseberg, L. H. 2006. Hybrid speciation in wild sunflowers. mobt 93:34–48.

Rieseberg, L. H. 2003. Major ecological transitions in wild sunflowers facilitated by

hybridization. Science 301:1211–1216.

Rieseberg, L. H., M. A. Archer, and R. K. Wayne. 1999. Transgressive segregation, adaptation

and speciation. Heredity 83:363–372.

Rieseberg, L. H., S.-C. Kim, R. A. Randell, K. D. Whitney, B. L. Gross, C. Lexer, and K. Clay.

2007. Hybridization and the colonization of novel habitats by annual sunflowers.

Genetica 129:149–165.

Rohatgi, V. K. 1976. An introduction to probability theory and mathematical statistics. Wiley,

New York.

Schluter, D. 2000. The ecology of adaptive radiation. Oxford University Press.

Schluter, D., E. A. Clifford, M. Nemethy, and J. S. McKinnon. 2004. Parallel evolution and

inheritance of quantitative traits. The American Naturalist 163:809–822.

Seehausen, O. 2013. Conditions when hybridization might predispose populations for adaptive

radiation. Journal of Evolutionary Biology 26:279–281.

Seehausen, O. 2004. Hybridization and adaptive radiation. Trends Ecol. Evol. 19:198–207.

Shaw, K. L. 2002. Conflict between nuclear and mitochondrial DNA phylogenies of a recent

species radiation: what mtDNA reveals and conceals about modes of speciation in

Hawaiian crickets. Proceedings of the National Academy of Sciences 99:16122–16127.

Slowinski, J. B., and C. Guyer. 1993. Testing whether certain traits have caused amplified

diversification: an improved method based on a model of random speciation and

extinction. The American Naturalist 142:1019–1024.

Soltis, P. S., and D. E. Soltis. 2009. The role of hybridization in plant speciation. Annual Review

of Plant Biology 60:561–588.

Stace, C. 1997. New flora of the British Isles, ed. 2.

Stankowski, S., and M. A. Streisfeld. 2015. Introgressive hybridization facilitates adaptive

divergence in a recent radiation of monkeyflowers. Proceedings of the Royal Society B:

Biological Sciences 282:20151666.

Stebbins, G. L. 1959. The role of hybridization in evolution. Proceedings of the American

Philosophical Society 231–251.

Stebbins, G. L. 1950. Variation and evolution in plants. Geoffrey Cumberlege.; London.

Stelkens, R. B., M. A. Brockhurst, G. D. Hurst, and D. Greig. 2014. Hybridization facilitates

evolutionary rescue. Evolutionary applications 7:1209–1217.

Suarez-Gonzalez, A., C. Lexer, and Q. C. B. Cronk. 2018. Adaptive introgression: a plant

perspective. Biology Letters 14:20170688.

Tank, D. C., J. M. Eastman, M. W. Pennell, P. S. Soltis, D. E. Soltis, C. E. Hinchliff, J. W.

Brown, E. B. Sessa, and L. J. Harmon. 2015. Nested radiations and the pulse of

angiosperm diversification: increased diversification rates often follow whole genome

duplications. New Phytologist 207:454–467.

Taylor, S. A., and E. L. Larson. 2019. Insights from genomes into the evolutionary importance

and prevalence of hybridization in nature. Nat Ecol Evol 3:170–177.

Testo, W., and M. Sundue. 2016. A 4000-species dataset provides new insight into the evolution

of ferns. Molecular Phylogenetics and Evolution 105:200–211.

Todesco, M., M. A. Pascual, G. L. Owens, K. L. Ostevik, B. T. Moyers, S. Hübner, S. M.

Heredia, M. A. Hahn, C. Caseys, D. G. Bock, and L. H. Rieseberg. 2016. Hybridization

and extinction. Evol Appl 9:892–908.

Tutin, T. G., V. H. Heywood, N. A. Burges, D. M. Moore, D. H. Valentine, S. M. Walters, and

D. A. Webb. 1964. Flora Europaea, vols. 1-5. Cambridge University Press, Cambridge.

Valente, L. M., G. Reeves, J. Schnitzler, I. P. Mason, M. F. Fay, T. G. Rebelo, M. W. Chase, and

T. G. Barraclough. 2010. Diversification of the African genus *Protea* (Proteaceae) in the

Cape biodiversity hotspot and beyond: equal rates in different biomes. Evolution 64:745–

759.

Vamosi, J. C., and T. A. Dickinson. 2006. Polyploidy and diversification: a phylogenetic

investigation in Rosaceae. International Journal of Plant Sciences 167:349–358.

Vamosi, J. C., and S. M. Vamosi. 2011. Factors influencing diversification in angiosperms: at the

crossroads of intrinsic and extrinsic traits. American Journal of Botany 98:460–471.

Vamosi, J. C., and S. M. Vamosi. 2010. Key innovations within a geographical context in

flowering plants: towards resolving Darwin's abominable mystery. Ecology Letters

13:1270–1279.

van der Bijl, W. 2018. phylopath: Easy phylogenetic path analysis in R. PeerJ 6.

Wagner, W. L., D. R. Herbst, and S. H. Sohmer. 1999. Manual of the Flowering Plants of

Hawai'i, Vols. 1 and 2. University of Hawai'i and Bishop Museum Press.

Walsh, N. G., and T. J. Entwisle. 1994. Flora of Victoria, vols. 2-4. Inkata Press, Melbourne,

Australia.

Weber, M. G., and A. A. Agrawal. 2014. Defense mutualisms enhance plant diversification.

Proceedings of the National Academy of Sciences 111:16442–16447.

Whitney, K. D., J. R. Ahern, L. G. Campbell, L. P. Albert, and M. S. King. 2010a. Patterns of

hybridization in plants. Perspectives in Plant Ecology, Evolution and Systematics

12:175–182.

Whitney, K. D., R. A. Randell, and L. H. Rieseberg. 2010b. Adaptive introgression of abiotic

tolerance traits in the sunflower *Helianthus annuus*. New Phytologist 187:230–239.

Whitney, K. D., R. A. Randell, and L. H. Rieseberg. 2006. Adaptive introgression of herbivore

resistance traits in the weedy sunflower *Helianthus annuus*. The American Naturalist

167:794–807.

Wolf, J. B., T. Bayer, B. Haubold, M. Schilhabel, P. Rosenstiel, and D. Tautz. 2010. Nucleotide

divergence vs. gene expression differentiation: comparative transcriptome sequencing in

natural isolates from the carrion crow and its hybrid zone with the hooded crow.

Molecular ecology 19 Suppl 1:162–75.

Wood, T. E., N. Takebayashi, M. S. Barker, I. Mayrose, P. B. Greenspoon, and L. H. Rieseberg.

2009. The frequency of polyploid speciation in vascular plants. Proceedings of the

national Academy of sciences 106:13875–13879.

Yakimowski, S. B., and L. H. Rieseberg. 2014. The role of homoploid hybridization in

evolution: A century of studies synthesizing genetics and ecology. American Journal of

Botany 101.

Yu, G., D. K. Smith, H. Zhu, Y. Guan, and T. T.-Y. Lam. 2017. ggtree: an R package for

visualization and annotation of phylogenetic trees with their covariates and other

associated data. Methods in Ecology and Evolution 8:28–36.

Zanne, A. E., D. C. Tank, W. K. Cornwell, J. M. Eastman, S. A. Smith, R. G. FitzJohn, D. J.

McGlinn, B. C. O'Meara, A. T. Moles, P. B. Reich, D. L. Royer, D. E. Soltis, P. F.

Stevens, M. Westoby, I. J. Wright, L. Aarssen, R. I. Bertin, A. Calaminus, R. Govaerts,

F. Hemmings, M. R. Leishman, J. Oleksyn, P. S. Soltis, N. G. Swenson, L. Warman, and

J. M. Beaulieu. 2014. Three keys to the radiation of angiosperms into freezing

environments. Nature 506:89.

FIGURE LEGENDS

Figure 1. Phylogeny containing 195 vascular plant families analyzed in this study. Rings depict

the standardized values of the log-transformed rates for (moving from inside to outside) the

hybridization measures (HybProp and HybRatio) and the net diversification rates (MS, Medusa,

and BAMM). Phylogenetically corrected analyses typically find a positive association between

hybridization and diversification (Fig. 3, Tables 1 and 2), but non-phylogenetically corrected

analyses find no relationship (Table S8). Note that the value for the standardized log-transformed

HybRatio in Genseriaceae was very high (5.953) and was not plotted (replaced with an "X") to

improve readability of the remaining data. Blue = relatively slow diversification rates / low

hybridization rates, orange = relatively fast diversification rates / high hybridization rates.

Figure 2. Raw net diversification rates calculated using three different methods are correlated for

(A) the full 459 family dataset and (B) the subset 195 family dataset.

Figure 3. PGLS results for (A) univariate PGLS regressions or (B) multivariate PGLS

regressions between either hybridization propensity or hybrid ratio and net diversification as

estimated by MS, medusa, or BAMM. Multivariate regressions include the percent of species

within each family that are perennial (Perenniality) and that are woody (Woodiness) as

additional covariates. Circle size corresponds to the absolute value of the partial regression

coefficient, color indicates the direction of the relationship (blue = positive, pink = negative) and

transparency indicates the signficance of the relationship (lightest = not significant, medium =

$0.05 < p < 0.10$, and dark = $p < 0.05$).

Table 1. Univariate PGLS modeling results.

Method	Hybrid Metric	Adj-R^2	Estimate	Std. Error	T-value	P-value
MS	HybProp	0.011	0.116	0.066	1.754	0.081
	HybRatio	0.023	0.135	0.057	2.372	0.019
Medusa	HybProp	0.013	0.102	0.055	1.859	0.065
	HybRatio	0.016	0.090	0.044	2.021	0.045
BAMM	HybProp	0.009	0.080	0.047	1.691	0.093
	HybRatio	0.012	0.066	0.036	1.817	0.071

Table 2. Multivariate PGLS modeling results

Method	Hybrid Metric	Adj-R^2	Covariate	Estimate	Std. Error	T-value	P-value
MS	HybProp	0.019	Hybridization	0.131	0.067	1.949	0.053
			Perenniality	0.056	0.067	0.844	0.400
			Woodiness	-0.140	0.074	-1.891	0.060
	HybRatio	0.033	Hybridization	0.148	0.058	2.573	0.011
			Perenniality	0.050	0.066	0.748	0.455
			Woodiness	-0.143	0.073	-1.949	0.053
Medusa	HybProp	0.005	Hybridization	0.108	0.056	1.932	0.055
			Perenniality	-0.013	0.053	-0.254	0.800
			Woodiness	-0.027	0.060	-0.450	0.653
	HybRatio	0.009	Hybridization	0.097	0.046	2.123	0.035
			Perenniality	-0.016	0.053	-0.303	0.762
			Woodiness	-0.030	0.060	-0.499	0.619
BAMM	HybProp	0.018	Hybridization	0.095	0.048	1.981	0.049
			Perenniality	0.002	0.043	0.055	0.956
			Woodiness	-0.083	0.050	-1.665	0.098
	HybRatio	0.022	Hybridization	0.081	0.037	2.191	0.030
			Perenniality	0.001	0.043	0.035	0.973
			Woodiness	-0.088	0.050	-1.755	0.081