Improved chemistry restraints for crystallographic refinement by integrating
the Amber force field into Phenix

Authors

Nigel W. Moriarty[a]*, Pawel A. Janowski[b1], Jason M. Swails[b], Hai Nguyen[b], Jane S.

Richardson[c], David A. Case[a] and Paul D. Adams[ad]

[a]Molecular Biosciences and Integrated Bioimaging, Lawrence Berkeley National Laboratory,

Berkeley, California, 94720-8235, USA

[b]Department of Chemistry & Chemical Biology, Rutgers University, Piscataway, NJ, 08854, USA

[c]Department of Biochemistry, Duke University, Durham, NC, 27710, USA

[d]Department of Bioengineering, University of California at Berkeley, Berkeley, CA, 94720, USA

Correspondence email: NWMoriarty@LBL.Gov

[1]Currently at Microsoft

Funding information   National Institutes of Health (grant No. GM122086 to David A. Case; grant No.

P01GM063210 to Paul D. Adams, Jane S. Richardson); Department of Energy (grant No. DE-AC02-

05CH11231 to Lawrence Berkeley National Laboratory).

Synopsis   The full Amber force field has been integrated into Phenix as an alternative refinement

target. With a slight loss in speed, it achieves improved stereochemistry, fewer steric clashes and

better hydrogen bonds.

Abstract   The refinement of biomolecular crystallographic models relies on geometric restraints to

help address the paucity of experimental data typical in these experiments. Limitations in these

restraints can degrade the quality of the resulting atomic models. Here we present an integration of the

full all-atom Amber molecular dynamics force field into Phenix crystallographic refinement, which

enables a more complete modeling of biomolecular chemistry. The advantages of the force field

include a carefully derived set of torsion angle potentials, an extensive and flexible set of atom types,

Lennard-Jones treatment of non-bonded interactions and a full treatment of crystalline electrostatics.

The new combined method was tested against conventional geometry restraints for over twenty-two

thousand protein structures. Structures refined with the new method show substantially improved

model quality. On average, Ramachandran and rotamer scores are somewhat better; clash scores and

MolProbity scores are significantly improved; and the modelling of electrostatics leads to structures

that exhibit more, and more correct, hydrogen bonds than those refined with traditional geometry

restraints. We find in general that model improvements are greatest at lower resolutions, prompting

plans to add the Amber target function to real-space refinement for use in electron cryo-microscopy.

This work opens the door to the future development of more advanced applications such as Amber-

based ensemble refinement, quantum mechanical representation of active sites and improved

geometric restraints for simulated annealing.

Keywords: Amber refinement target; H-bond quality; Amber in Phenix; Cβ deviations; peptide
orientations

1. Introduction

Accurate structural knowledge lies at the heart of our understanding of the biomolecular function and

interactions of proteins and nucleic acids. With close to 90% of structures in the Protein Data Bank

(Berman *et al.*, 2000) solved via x-ray diffraction methods, crystallography is currently the pre-

eminent method for determining biomolecular structure. Crystal structure refinement is a

computational technique that plays a key role in post-experiment data interpretation. Refinement of

atomic coordinates entails solving an optimization problem to minimize the residual difference

between the experimental and model structure factor amplitudes (Jack & Levitt, 1978; Agarwal, 1978;

Murshudov *et al.*, 1997). However, due to inherent experimental limitations and a typically low data

to parameter ratio, the employment of additional restraints, commonly referred to as geometry or

steric restraints, is key to successful structural refinement (Waser, 1963). These restraints, which can

be thought of as a prior in the Bayesian sense, provide additional observations in the optimization

target and reduce the danger of overfitting. Their use leads to higher quality, more chemically

accurate models.

Most current refinement programs (Afonine *et al.*, 2012; Murshudov *et al.*, 2011; Sheldrick, 2008;

Bricogne *et al.*, 2011) employ a set of covalent-geometry restraints first proposed by Engh & Huber in

1991 and later augmented and improved in 2001 (Engh & Huber, 1991, 2001). This set of restraints is

based on a survey of accurate high-resolution small molecule crystal structures from the Cambridge

Structural Database (Groom *et al.*, 2016) and includes restraints on interatomic bond lengths, bond

angles and ω torsion angles. In addition, parameters are added to enforce proper chirality and

planarity; multiple-minimum targets for backbone and side chain torsion angles; and repulsive terms

to prevent steric overlap between atoms. Those terms are defined from small-molecule and high-

resolution macromolecular crystal structure data and from interaction-specified van der Waals radii.

They are very similar but not identical between refinement programs.

The Engh & Huber restraints function reasonably well, while the additional terms have been gradually

improved, but a number of limitations have been identified over the years. Some of these limitations

include: a lack of adjustability to differences in local conformation, protonation, and hydrogen

bonding and to their changes during refinement; incomplete or inaccurate atom types and parameters

for ligands, carbohydrates, and covalent modifications; use only of repulsive and not attractive steric

terms; omission of explicit hydrogen atoms and their interactions; misleading targets resulting from

experimental averaging artifacts; inaccurate dihedral restraints; and lack of awareness of electrostatic

and quantum dispersive interactions with a consequent lack of accounting for hydrogen bonding

cooperativity (Priestle, 2003; Touw & Vriend, 2010; Davis *et al.*, 2003; Moriarty *et al.*, 2014; Tronrud

et al., 2010).

Phenix (Adams *et al.*, 2010) includes a built-in system for defining ligand parameters (Moriarty *et al.*,

2009) that by default restrains the explicit hydrogen atoms at electron-cloud-center positions for X-ray

and optionally at nuclear positions for neutron crystallography (Williams, Headd *et al.*, 2018).

Addition of the Conformation Dependent Library (CDL) (Moriarty *et al.*, 2014), which makes

backbone bond lengths and angles dependent on φ,ψ values, has improved the models obtained from

refinement at all resolutions, and thus is the default in Phenix refinement (Moriarty *et al.*, 2016).

Similarly, Phenix uses ribose-pucker and base-type dependent torsional restraints for RNA (Jain *et al.*,

2015). For bond lengths and angles, protein side chains continue to use standard Engh & Huber

restraints while RNA/DNA use early values (Gelbin *et al.*, 1996; Parkinson *et al.*, 1996) with a few

modifications. This use of combined restraints is here designated CDL/E&H.

An alternative approach is the use of geometry restraints based on all-atom force fields used for

molecular dynamics studies. This is not a novel idea. In fact, some of the earliest implementations of

refinement programs employed molecular mechanics force fields (Jack & Levitt, 1978; Brünger *et al.*,

1987, 1989). However, at the time, restraints derived from coordinates of ideal fragments (Tronrud *et*

al., 1987; Hendrickson & Konnert, 1980) were found to provide better refinement results. The

insufficiency of molecular mechanics-based restraints was mainly attributed to two factors: inaccurate

representation of chemical space because of too few atom types, and biases in conformational

sampling resulting from unshielded electrostatic interactions. Subsequently, however, the methods of

molecular dynamics and corresponding force fields have seen significant development and

improvement. Current force fields contain more atom types and are easily adjustable as needed. They

are typically parameterized against accurate quantum mechanical calculations, not feasible just a few

years ago, as well as using more representative experimental results. Significant methodological

advances, such as the development of Particle Mesh Ewald (York *et al.*, 1993; Darden *et al.*, 1993) for

accurate calculation of crystalline electrostatics and improved temperature and pressure control

algorithms, have greatly increased accuracy. Modern force fields have been shown to agree well with

experimental data (Zagrovic *et al.*, 2008; van Gunsteren *et al.*, 2008; Showalter & Brüschweiler,

2007; Grindon *et al.*, 2004; Bowman *et al.*, 2011), including crystal diffraction data (Cerutti *et al.*,

2009; Janowski *et al.*, 2013; Cerutti *et al.*, 2008; Liu *et al.*, 2015; Janowski *et al.*, 2015).

We have made it possible to use of the Amber molecular mechanics force field as an alternative

source of geometry restraints to those of CDL/E&H. Here we present an integration of the Phenix

software package for crystallographic refinement, *phenix.refine* (Afonine *et al.*, 2012) and the Amber

software package (Case *et al.*, 2018) for molecular dynamics. We present results of paired refinements

for 22,544 structures and compare Amber to traditional refinement in terms of model quality,

chemical accuracy and agreement with experimental data, studied both for overall statistics and for

representative individual examples. We also describe the implementation and discuss future

directions.

2. Methods

2.1. Code preparation

The integration of the Amber code into *phenix.refine* uses a thin client. Amber provides a python API

to its *sander* module, so that a simple "import sander" python command allows Phenix to obtain

Amber energies and forces through a method call. At each step of coordinate refinement, Phenix

expands the asymmetric unit coordinates to a full unit cell (as required by *sander*), combines energy

gradients returned from Amber (in place of those from its internal geometric restraint routines) with

gradients from the X-ray target function, and uses these forces to update the coordinates, either by

minimization or by simulated annealing molecular dynamics. Alternate conformers take advantage of

the "locally-enchanced-sampling" (LES) facility in *sander*: atoms in single-conformer regions interact

with multiple-copy regions via the average energy of interaction, while different copies of the same

group do not interact among themselves (Roitberg & Elber, 1991; Simmerling *et al.*, 1998).

The Amber files required are created by a preliminary *AmberPrep* program that takes a PDB file as

input. It creates both a parameter-topology (*prmtop*) file used by Amber and a new PDB file

containing a complete set of atoms (including hydrogens and any missing atoms) needed to do force

field calculations. Alternate conformers, if present in the input PDB file, are translated into *sander*

LES format. For most situations, *AmberPrep* does not require the user to have any experience with

Amber or with molecular mechanics; less-common situations (described below) require some

familiarity with Amber. All the code required for both the *AmberPrep* and *phenix.refine* steps is

included in the current major release, 1.16-3549 and subsequent nightly builds of Phenix.

2.2. Structure selection and overall refinement protocol

To compare refinements using Amber against traditional refinements with CDL/E&H restraints,

structures were selected from the Protein Data Bank (Burley *et al.*, 2019) using the following criteria.

Entries must have untwinned experimental data available that are at least 90% complete. Each entry's

R_{free} was limited to a maximum of 35%, R_{work} to 30% and the ΔR (R_{free}-R_{work}) to a minimum of 1.5%.

The lowest resolution was set at 3.5Å. Entries containing nucleic acids were excluded.

Coordinate and experimental data files were obtained directly from the Protein Data Bank (PDB) and

inputs prepared via the automated *AmberPrep* program (see section 2.1 above). Entries containing

complex ligands were included if the file preparation program *AmberPrep* was able to automatically

generate and include the ligand geometry data. Details of the internals of *AmberPrep* will be

described elsewhere. Resolution bins (set at 0.1Å) with less than 10 refinement pairs were eliminated

to reduce noise caused by limited statistics. Complete graphs are included in the supplemental

material. The resulting 22,000+ structures had experimental data resolutions between 0.5Å and 3.2Å,

with most of the structures in the 1.0-3.0 Å range (see figure 1).

Each model was then subjected to 10 macrocyles of refinement using the default strategy in

phenix.refine for reciprocal space coordinate refinement, with the exception that real space refinement

was turned off. By default, the first macrocycle uses a least-squares target function and the rest use

maximum likelihood. Other options included optimization of the weight between the experimental

data and the geometry restraints. This protocol was performed in parallel, once using CDL/E&H and

once using Amber geometry restraints. In addition, Cβ pseudo-torsion restraints were not included in

the restraints model. Only one copy of each alternate conformation was considered initially (i.e.

alternative location A). The quality of the resulting models was assessed numerically using

MolProbity (Williams, Headd *et al.*, 2018) available in Phenix (Adams *et al.*, 2010), by *cpptraj* (Roe

& Cheatham, 2013) available in *AmberTools* (Case *et al.*, 2018) and by visual inspection with electron

density and validation markup in KiNG (Chen *et al.*, 2009) . All-atom dots for figure 10 were counted

in Mage (Richardson & Richardson, 2001) and figures 5-9 were made in KiNG. To avoid

typographical ambiguity, PDB codes are given here with lower case for all letters except L (e.g.,

1nLs).

2.3. Weight factor details

157   The target function optimized in *phenix.refine* reciprocal space atomic coordinate refinement is of the

158   general form:

$$T_{xyz} = w * T_{exp} + T_{xyz\_restraints}$$

159   where all the terms are functions of the atomic coordinates, T_{xyz} is the target residual to be minimized,

160   T_{exp} is a residual between the observed and model structure factors and quantifies agreement with

161   experimental data, $T_{xyz\_restraints}$ is the residual of agreement with the geometry restraints and w is a scale

162   factor that modulates the relative weight between the experimental and the geometry restraint terms.

163   In traditional refinement $T_{xyz\_restraints}$ is calculated using the set of CDL/E&H restraints:

$$T_{xyz} = w * T_{exp} + T_{CDL/E\&H}$$

164   To implement Phenix-Amber we substitute this term with the potential energy calculated using the

165   Amber force field:

$$T_{xyz} = w * T_{exp} + E_{AmberFF}$$

where the Amber term is intentionally represented now by an *E* to emphasize that we directly

incorporate the full potential energy function calculated in Amber using the ff14SB (Maier *et al.*,

2015) force field.

In a standard default Phenix refinement, the weight, w, is a combination of a value based on the ratio

of gradient norms (Brünger *et al.*, 1989; Adams *et al.*, 1997) and a scaling factor that defaults to ½.

This initial weight can be optimized using a procedure described previously (Afonine *et al.*, 2011).

This procedure uses the results of ten refinements with a selection of weights, considering the bond

and angle rmsd, the R-factors and validation statistics to determine the best weight for the specific

refinement at each of the ten macrocycles. The same procedure was used to estimate an optimal

weight for the Phenix-Amber refinements. (If faster fixed-weight refinements are desired, we have

found that a scaling factor of 0.2, rather than 0.5, scales the Amber gradients to be close to those from

the CDL/E&H restraints, allowing the simpler, default, weighting scheme in *phenix.refine* to be used.)

178   3. Results

179   3.1. Full-dataset score comparisons

On average, the Phenix-Amber combination produced slightly higher R-work and R-free (figure 2)

but higher quality models (figure 3). The increase in R-factors is most pronounced in the 1.5–2.5Å

range. This is a result of the weight optimisation procedure having different limits for optimal weight

in this resolution range. The increase was less for R-free than R-work such that the R-delta is less for

refinements using Amber gradients. The Phenix-Amber refinements exhibited improved (lower)

MolProbity scores and contained fewer clashes between atoms. Plots show the mean of the values in

the 0.1Å resolution bin as well as the 95% confidence level of the standard error of the mean (SEM).

MolProbity clashscores are particularly striking: for refinement using CDL/E&H restraints,

clashscores steadily increase as resolution worsens, often resulting in very high numbers of steric

clashes. On the other hand, the mean clash-score with Amber restraints appears to be nearly

independent of resolution and remains consistent at about 2.5 clashes per 1000 atoms across all

resolution bins. The SEM range is non-overlapping for worse than 1Å indicating that the Amber force

field is producing better geometries at mid to low resolution. There are more favored Ramachandran

points (backbone ϕ,ψ) and fewer Ramachandran outliers for the Phenix-Amber refinements. This

difference is most marked for resolutions worse than 2Å. Phenix-Amber refinement also improves

(lowers) the number of rotamer outliers but doesn't differentiate via the SEM, and increases the

proportion of hydrogen bonds. While the rotamer outlier results remain similar, the hydrogen bonding

results have a large difference at worse than 1.5Å resulting in nearly double the bonds near 3Å.

Common to all the plots is a change near 1.5Å, where the weight optimisation procedure common to

both CDL/E&H and Amber refinement loosens the weight on geometry restraints somewhat, to allow

more deviations at resolutions where the data is capable of unambiguously showing them. Bond and

angle rmsd comparison are less pertinent as the force fields do not have ideal values for

parameterisations and comparing the Phenix-Amber bonds and angles to the CDL/E&H values is not

a universal metric. The curious can see the plots in figure S1. Overall, improvement with Amber is
substantial in the lower resolution refinements.

Models refined with Phenix-Amber are more likely to exhibit electrostatic interactions such as H-
bonds and salt links, as well as better van der Waals contacts. Though the resulting atom movements
are generally small, these changes can be meaningful, especially when interpreting H-bonding
networks or interaction distances at active sites.

One validation metric that is worse for Phenix-Amber refinements is the number of outliers of the Cβ
positions. Both the mean and the SEM show clear differentiation. The Cβ deviation gives a combined
measure of distortion in the tetrahedron around the Cα atom and with traditional E&H restraints it is
quite robustly sensitive to incompatibility between how the backbone and side chain conformations
have been modelled (Lovell *et al.*, 2003). For CDL/E&H refinements, however, the percentage of Cβd
outliers (>0.25Å) is negligible for low and mid resolutions, only increasing to 0.2% at higher
resolutions (see figure 4). This is in line with the CDL/E&H providing tight geometrical restraints out
to Cβ at most resolutions, but loosened somewhat at better than 1.5Å resolution where there is enough
experimental information to move an angle away from ideal. Note that explicit Cβ restraints were
turned off for all Phenix refinements and that the Amber force field does not have an explicit Cβ term;
however, if all angles around the Cα are kept ideal then the Cβ position will also be ideal even if it is
incorrectly positioned in the structure. The following section analyses specific local examples where
output structures show differences for either the positive or the negative trends seen in the overall
comparisons, in order to understand their nature, causes and meaning across resolution ranges.

3.2. Examination of individual examples

As noted above, in comparison with the CDL/E&H restraint refinements, the Phenix-Amber
refinements have much higher percentages of Cβ deviation outliers, increasing at the low-resolution
end to more than 1% of Cβ atoms. Amber refinement also has more bond length and angle outliers.
The following examines a sample of cases at high, mid and lower resolutions to understand the
starting-model characteristics and refinement behavior that produce these differences.

3.2.1. High resolution: waters, alternates, Cβd outliers and atoms in the wrong peak

In the high-resolution range (better than 1.7Å), it appears that the commonest problems not easily
correctable by refinement are caused either by modeling the wrong atom into a density peak or by
incorrect modeling, labeling, or truncation of alternate conformations. Such problems are usually
flagged in validation either by all-atom clashes, by Cβ deviations and sometimes by bad bond lengths
and angles.

Figure 5a shows a case where a water molecule had been modeled in an electron density peak that
should really be a nitrogen atom of the Arg guanidinium. CDL/E&H refinement (figure 5b) corrected

the bad geometry at the cost of moving the guanidinium even further out of density; Amber

refinement changed the guanidinium orientation but made no overall improvement (figure 5c); all

three versions have a bad clash. If the water were deleted, then either refinement method would

undoubtedly do an excellent job (figure 5d). This type of problem is absent at low resolution where

waters are not modeled but occurs quite often at both high and mid resolution, for other branched side

chains, for Ile Cδ (for example, 3js8 195) and even occasionally for Trp (e.g. 1qw9 B170).

Cβ deviation outliers (≥0.25Å) are often produced by side chain alternates with quite different Cβ

positions but no associated alternates defined along the backbone. Since the tetrahedron around Cα

should be nearly ideal, that treatment almost guarantees bad geometry. The rather simple solution,

implemented in Phenix, is to define alternates for all atoms until the i+1 and i-1 Cα atoms – as in the

"backrub" motion; (Davis *et al.*, 2006). PDB codes 1dy5, 1gwe and 1nLs each have a number of such

cases. Figure 6a,b show 1nLs Ser 215, initially with an outlier Cβd, 0.49Å distance between the two

Cβ atoms and a single Cα. CDL/E&H refinement pulls the Cβ atoms to be only 0.23Å apart, avoiding

a Cβd with only slightly worse fit to the density; Amber reduces the Cβd only slightly, but it does

keep this flag of an underlying problem. When alternates are defined for the backbone peptides, both

systems improve.

Worse cases occur where one or both alternates have been fit incorrectly as well as not being

expanded along the backbone appropriately. Figure 6c shows Thr 196, with a huge Cβd of 0.88Å (not

shown) and very poor geometry, because altB was fit incorrectly (just as a shift of altA rather than as

a new rotamer). This time even CDL/E&H refinement produces a Cβd outlier, but smaller than for

Amber. Figure 6d shows the excellent Amber result after the misfit of altB was approximately

corrected.

3.2.2. Mid resolution: backward side chains and rare conformations

An even commoner case at both high and mid resolutions where the wrong atom is fit into a density

peak is a backward-fit Cβ-branched residue, well illustrated by a very clear Thr example in 1bkr at

1.1Å (figure 7a). Thr 101 is a rotamer outlier (gold) on a regular α-helix with a Cβd of 0.63Å. The

deposited Thr 101 also has a bond-angle deviation of 13.5σ; clashes at the Cγ methyl; its Cβ is out of

density; Oγ is in the lower peak; and Cγ is in the higher peak. It is shown in figure 7 with 1.6σ and 4σ

$2mF_o\text{-}DF_c$ contours (but without Cβ deviation and angle markups for clarity). This mistake was not

obvious because anisotropic B's were used too early in the modeling resulting in the Thr Cβ being

refined to a 6:1 aniso-axis ratio that covered both the modeled atom and the real position. The figures

show the density as calculated with isotropic B factors.

Given this difficult problem for automated refinement, each of the two target functions reacts very

differently. Both refinements still have the Cγ methyl clashing with a helix backbone CO in good

density, very diagnostic of a problem with the Cγ. It is indeed the wrong atom to have in that peak, as

shown also by the relative peak heights. The CDL/E&H refinement (figure 7b) achieves tight

geometry and a good rotamer, moving the Cβ into its correct density peak, but pays the price for not

correcting the underlying problem by swinging the Oγ out of density. The Amber refinement (figure

7c) achieves an atom in each of the three side chain density peaks, but pays the price for not

correcting the underlying problem by having the wrong chirality at the Cβ atom. It still also has bond-

angle outliers, which may be a sign of unconverged refinement.

The original PDB entry, the CDL/E&H refinement and the Amber refinement structures for Thr 101

are all very badly wrong, but each in an entirely different way. The deposited model, 1bkr, looks very

poor by traditional model validation, but has a misleadingly good density correlation, given the

extremely anisotropic Cβ B-factor. The CDL/E&H output looks extremely good on traditional

validation except for the clashes and would show a lowered but still reasonable density correlation;

however, it is the most obviously wrong upon manual inspection. The Amber output has clashes and

currently has modest bond-angle outliers, but it fits the density very closely making it difficult to

identify as incorrect by visual inspection. The problem could be recognized automatically by a simple

chirality check. Shown in figure 7d, Thr 101 was rebuilt quickly in KiNG, with the **p** rotamer and a

small backrub motion. Either Phenix-CDL/E&H or Phenix-Amber refinement would do a very good

job from such a rough refit with the correct atoms near the right places.

At mid resolution, there are also other rotamers and backbone conformations fit into the wrong local

minimum and thus difficult to correct by minimization refinement methods, but not always flagged by

Cβ deviations or other outliers. Some of these, such as *cis*-nonPro peptides (Williams, Videau *et al.*,

2018) or very rare rotamers (Hintze *et al.*, 2016) can be avoided by considering their highly

unfavorable prior probabilities. Others would require explicit sampling of the multiple minima.

3.2.3. Lower resolution: peptide orientations with CaBLAM and Cβd outliers

At low resolution (2.5–4Å), no waters or alternates are modeled. All other problems continue, but an

additional set of common local misfittings occur because the broad electron density is compatible

with significantly different models. 1xgo at 3.5Å is an excellent case for testing in this range, because

it was solved independently from the 1.75Å 1xgs structure – the same molecule in a different space

group. CDL/E&H refinement shows no Cβd outliers, but Amber refinement has six. Comparison with

1xgs shows that each of the Cβd residues has either the side chain or the backbone or both in an

incorrect local-minimum conformation uncorrectable by minimization refinement methods

(Richardson & Richardson, 2018). For example, figure 8 shows Leu 253 on a helix, with a Cβd from

Amber (panel c) and the different, correct 1xgs Leu rotamer in panel d. Those Cβd outliers are thus a

feature, not a bug, in Amber: they serve their designed validation function of flagging genuine fitting

problems. However, the lack of Cβd outliers in the CDL/E&H refinement is also not a defect, because

the tight CDL/E&H geometry is on average quite useful at low resolution.

The 1xgo-vs-1xgs comparison also illustrates many of the ways in which Amber refinement is

superior at low resolution. In figure 8, Amber corrects a Ramachandran outlier in the helix and shows

a helix backbone shape much closer to the ideal geometry of 1xgs than either the deposited or the

CDL/E&H versions.

Since the backbone CO direction cannot be seen at low resolution, the commonest local misfitting is a

misoriented peptide (Richardson *et al.*, 2018). Those can be flagged by the new MolProbity validation

called CaBLAM, which tests whether adjacent CO directions are compatible with the local Cα

backbone conformation (Williams, Headd *et al.*, 2018). Ten such cases were identified in 1xgo, for

isolated single or double CaBLAM outliers surrounded by correct structure as judged in1xgs. For six

of those 10 cases, neither CDL/E&H nor Amber refinement corrected the problem: His62, Thr70,

Gly163, Gly193, Ala217, Glu286 (see stereo figure S2). In two cases CDL/E&H had fewer other

outliers than Amber, but did not actually reorient the CO: for Gly193 and for the Gly163 case shown

in figure S3. In three of the 10 cases Amber did a complete fix, while CDL/E&H did not improve

(Asp88, Gly125, Pro266). For example, in figure 9, 1xgo residues 86-91 (panel a) have a CaBLAM

outlier (magenta lines), uncorrected by CDL/E&H refinement (panel b). But Amber refinement (panel

c) manages to shift several CO orientations by modest amounts (red balls), enough to fix the

CaBLAM outliers and match extremely closely the better backbone conformation of 1xgs (panel d).

The Gly 125 example is shown in figure S4. Finally, in one especially interesting case (Lys22) Amber

turned the CO about halfway up to where it should be, while CDL/E&H made no improvement. The

Amber model still has geometry outliers and further runs moved most of the way up and removed

those outliers, showing that Amber refinement had not yet fully converged in 10 macrocycles (see

Supplement text and figure S5).

Amber refinement is especially good at optimizing hydrogen-aware all-atom sterics, as calculated by

the Probe program (Word, Lovell, LaBean *et al.*, 1999) with H atoms added and optimized by Reduce

(Word, Lovell, Richardson *et al.*, 1999). This is illustrated in figure 10 for 3g8L at 2.5Å resolution.

The deposited structure of the Asn 182 helix N-cap region, which has many outliers of all kinds

(panel a), is improved a great deal by CDL/E&H refinement (panel b). However, the Amber

refinement (panel c) is noticeably better, with more H-bonds and better van der Waals contacts as well

as fewer clashes. These improvements are plotted quantitatively in figure 11, as measured by a

dramatic drop in unfavorable clash spikes (red) and small overlaps (yellow), with a dramatic increase

in favorable H-bonds (green) and van der Waals contacts (blue).

4. Discussion

The idea of including molecular mechanics force fields into crystallographic refinements is not a new

one, with precedents dating back to early work by (Jack & Levitt, 1978) and the XPLOR program

(Brünger & Karplus, 1991) developed in the 1980's. The notion that a force field could (at least in

principle) encode "prior knowledge" about protein structure continues to have a strong appeal and

efforts to replace conventional "geometric restraints", which are very local and uncorrelated, with a

more global assessment of structural quality have been explored repeatedly (e. g., Moulinier *et al.*,

2003; Schnieders *et al.*, 2009). Distinguishing features of the current implementation include

automatic preparation of force fields for many types of biomolecules, ligands and solvent components

as well as close integration with Phenix, a mature and widely used platform for refinement. This has

enabled parallel refinements on more than 22,000 protein entries in the PDB and allows

crystallographers to test these ideas on their own systems by simply adding flags to an existing

phenix.refine command line or adding the same information via the Phenix GUI. Indeed, we expect

most users to "turn on" Amber restraints after having carried out a more conventional refinement to

judge for themselves the significance and correctness of structural differences that arise. As noted in

Section 3.2, an Amber refinement will often flag residues that need manual refitting in ways

complementary to the cues provided by more conventional refinement.

The results presented here show that structures with improved local quality (as monitored by

MolProbity criteria and hydrogen bond analysis) can be obtained by simple energy minimization, with

minimal degradation in agreement with experimental structure factors and with no changes to a

current-generation protein force field. Nevertheless, one should keep in mind that the Amber-refined

structures obtained here are not very different from those found with more conventional refinement.

Both methods require that most local misfittings to be corrected in advance. The hope is that either

sampling of explicit alternatives or else optimization using more aggressive conformational search,

such as with simulated annealing or torsion-angle dynamics, may find the correct low-energy

structures with good agreement with experimental data.

It is likely that further exploration of relative weights between "X-ray" and "energy" terms (beyond

the existing and heuristic weight-optimization procedure employed here) and even within the energy

terms, will become important. In principle, maximizing the joint probability arising from "prior

knowledge" (using a Bolztmann distribution, $\exp(-E_{AmberFF}/k_BT)$ for some effective temperature) and a

maximum likelihood target function (based on a given model and the observed data) is an attractive

approach that effectively establishes an appropriate relative weighting. More study will be needed to

see how well this works in practice, especially in light of the inevitable limitations of current force

fields.

The integration of Amber's force field into the Phenix software for crystallography also paves the way

for the development of more sophisticated applications. The force field can accommodate alternate

conformers by using the locally enhanced sampling (LES) approach (Roitberg & Elber, 1991;

Simmerling *et al.*, 1998); a few examples are discussed here whilst details will be presented

elsewhere. Ensemble refinement (Burnley *et al.*, 2012) could now be performed using a full molecular

dynamics force field, thus avoiding poor quality individual models in the ensemble. Similarly,

simulated annealing could now be performed with an improved physics-based potential. Extension of

the ideas presented to real-space refinement within Phenix is underway, opening a path to new

applications to cryo-EM and low-resolution X-ray structures. These developments would all

contribute significantly to the future of macromolecular crystallography, reinforcing the transition

from a single static-structure-dominated view of crystals to one where dynamics and structural

ensembles play a central important role in describing molecular function (Furnham *et al.*, 2006; van

den Bedem & Fraser, 2015; Wall *et al.*, 2014).

5. Conclusions

We have presented refinement results obtained by integrating Phenix with the Amber software

package for molecular dynamics. Our refinements of over 22,000 crystal structures show that

refinement using Amber's all atom molecular mechanics force field outperforms CDL/E&H restraint

refinement in many respects. An overwhelming majority of Amber-refined models display notably

improved model quality. The improvement is seen across most indicators of model quality including

clashes between atoms, side chain rotamers and peptide backbone torsion angles. In particular,

Phenix-Amber consistently outperforms standard Phenix refinement in clashscore, number of

hydrogen bonds and MolProbity score. It also consistently outperforms standard refinement for

Ramachandran and rotamer statistics at low resolutions and obtains approximately equal results at

high (better than 2.0Å) resolutions. Amber does run somewhat more slowly (generally 20-40%

longer) and may take more cycles to converge completely if it is making any large local changes (see

text for supplementary figure S5). It should be noted that standard refinement consistently

outperforms Phenix-Amber in eliminating Cβ deviation and other covalent-geometry outliers across

all resolutions, but in many cases the Amber outliers serve to flag a real problem in the model.

As the quality of experimental data decreases with resolution, the improvement in model quality

obtained by using Amber, as opposed to CDL/E&H restraints, increases. This improvement is

especially striking in the case of clashscores, which appear to be nearly independent of experimental

data resolution for Amber refinements. Additional improvement is seen in the modelling of

electrostatic interactions, H-bonds and van der Waals contacts, which are currently ignored by

conventional restraints. Improving lower-resolution structures is very important, since they include a

large fraction of the most exciting and biologically important current structures such as the

protein/nucleic acid complexes of big, dynamic molecular machines.

No minimization refinement method, including CDL/E&H and Amber, can in general correct local

misfittings that were modeled in an incorrect local-minimum conformation, especially at relatively

high resolutions. At lower resolution where the barriers are softer, Amber sometimes can manage

such a change, while CDL/E&H still does not. It is, therefore, important and highly recommended that

validation flags be consulted for the initial model and as many as feasible of the worst cases be fixed,

before starting the cycles of automated refinement with either target.

Software distribution Amber was implemented in *phenix.refine* and is available in the 1.16-3549

version of Phenix and later. Instructions for using the *phenix.refine* Amber implementation are

available in the version-specific documentation available with the distribution.

Acknowledgements   JSR thanks David Richardson for help with some aspects of the individual-

example analyses. The content is solely the responsibility of the authors and does not necessarily

represent the official views of the National Institutes of Health, NIGMS, or DOE.

References

Adams, P. D., Afonine, P. V, Bunkóczi, G., Chen, V. B., Davis, I. W., Echols, N., Headd, J.
        J., Hung, L.-W., Kapral, G. J., Grosse-Kunstleve, R. W., McCoy, A. J., Moriarty, N.
        W., Oeffner, R., Read, R. J., Richardson, D. C., Richardson, J. S., Terwilliger, T. C. &
        Zwart, P. H. (2010). *Acta Crystallogr Sect D*. **66**, 213–221.

Adams, P. D., Pannu, N. S., Read, R. J. & Brünger, A. T. (1997). *Proc. Natl. Acad. Sci.* **94**,
        5018–5023.

Afonine, P. V., Echols, N., Grosse-Kunstleve, R. W., Moriarty, N. W. & Adams, P. D.
        (2011). *Comput. Crystallogr. Newsl.* **2**, 99–103.

Afonine, P. V, Grosse-Kunstleve, R. W., Echols, N., Headd, J. J., Moriarty, N. W.,
        Mustyakimov, M., Terwilliger, T. C., Urzhumtsev, A., Zwart, P. H. & Adams, P. D.
        (2012). *Acta Crystallogr Sect D*. **68**, 352–367.

Agarwal, R. C. (1978). *Acta Crystallogr. Sect. A*. **34**, 791–809.

van den Bedem, H. & Fraser, J. S. (2015). *Nat. Methods*. **12**, 307–318.

Berman, H. M., Westbrook, J., Feng, Z., Gilliland, G., Bhat, T. N., Weissig, H., Shindyalov,
        I. N. & Bourne, P. E. (2000). *Nucleic Acids Res.* **28**, 235–242.

Bowman, G. R., Voelz, V. A. & Pande, V. S. (2011). *J. Am. Chem. Soc.* **133**, 664–667.

Bricogne, G., Blanc, E., Brandl, M., Flensburg, C., Keller, P., Paciorek, W., Roversi, P.,
        Sharff, A., Smart, O. S., Vonrhein, C. & Womack, T. O. (2011).

Brünger, A. T. & Karplus, M. (1991). *Acc. Chem. Res.* **24**, 54–61.

Brünger, A. T., Karplus, M. & Petsko, G. A. (1989). *Acta Crystallogr. Sect. A*. **45**, 50–61.

Brünger, A. T., Kuriyan, J. & Karplus, M. (1987). *Science*. **235**, 458–460.

Burley, S. K., Berman, H. M., Bhikadiya, C., Bi, C., Chen, L., Costanzo, L. D., Christie, C.,
Duarte, J. M., Dutta, S., Feng, Z., Ghosh, S., Goodsell, D. S., Green, R. K.,
Guranovic, V., Guzenko, D., Hudson, B. P., Liang, Y., Lowe, R., Peisach, E.,
Periskova, I., Randle, C., Rose, A., Sekharan, M., Shao, C., Tao, Y.-P., Valasatava,
Y., Voigt, M., Westbrook, J., Young, J., Zardecki, C., Zhuravleva, M., Kurisu, G.,
Nakamura, H., Kengaku, Y., Cho, H., Sato, J., Kim, J. Y., Ikegawa, Y., Nakagawa,
A., Yamashita, R., Kudou, T., Bekker, G.-J., Suzuki, H., Iwata, T., Yokochi, M.,
Kobayashi, N., Fujiwara, T., Velankar, S., Kleywegt, G. J., Anyango, S., Armstrong,
D. R., Berrisford, J. M., Conroy, M. J., Dana, J. M., Deshpande, M., Gane, P.,
Gáborová, R., Gupta, D., Gutmanas, A., Koča, J., Mak, L., Mir, S., Mukhopadhyay,
A., Nadzirin, N., Nair, S., Patwardhan, A., Paysan-Lafosse, T., Pravda, L., Salih, O.,
Sehnal, D., Varadi, M., Vařeková, R., Markley, J. L., Hoch, J. C., Romero, P. R.,
Baskaran, K., Maziuk, D., Ulrich, E. L., Wedell, J. R., Yao, H., Livny, M. &
Ioannidis, Y. E. (2019). *Nucleic Acids Res.* **47**, D520–D528.

Burnley, B. T., Afonine, P. V, Adams, P. D. & Gros, P. (2012). *ELife*. **1**, e00311.

Case, D. A., Ben-Shalom, I. Y., Brozell, S. R., Cerutti, D. S., Cheatham, III, T. E., Cruzeiro,
V. W. D., Darden, T. A., Duke, R. E., Ghoreishi, D., Gilson, M. K., Gohlke, H.,
Goetz, A. W., Greene, D., Harris, R., Homeyer, N., Izadi, S., Kovalenko, A.,
Kurtzman, T., Lee, T. S., LeGrand, S., Li, P., Lin, C., Liu, J., Luchko, T., Luo, R.,
Mermelstein, D. J., Merz, K. M., Miao, Y., Monard, G., Nguyen, C., Nguyen, H.,
Omelyan, I., Onufriev, A., Pan, F., Qi, R., Roe, D. R., Roitberg, A., Sagui, C., Schott-
Verdugo, S., Shen, J., Simmerling, C. L., Smith, J., Salomon-Ferrer, R., Swails, J.,
Walker, R. C., Wang, J., Wei, H., Wolf, R. M., Wu, X., Xiao, L., York, D. M. &
Kollman, P. A. (2018). AMBER 18 University of California, San Francisco.

Cerutti, D. S., Le Trong, I., Stenkamp, R. E. & Lybrand, T. P. (2008). *Biochemistry*. **47**,
12065–12077.

Cerutti, D. S., Le Trong, I., Stenkamp, R. E. & Lybrand, T. P. (2009). *J. Phys. Chem. B*. **113**,
6971–6985.

Chen, V. B., Davis, I. W. & Richardson, D. C. (2009). *Protein Sci. Publ. Protein Soc.* **18**,
2403–2409.

Darden, T., York, D. M. & Pedersen, L. (1993). *J. Chem. Phys.* **98**, 10089.

Davis, A. M., Teague, S. J. & Kleywegt, G. J. (2003). *Angew. Chem. Int. Ed Engl.* **42**, 2718–
2736.

Davis, I. W., Arendall, W. B., Richardson, D. C. & Richardson, J. S. (2006). *Struct. Lond.*
Engl. 1993. **14**, 265–274.

Engh, R. A. & Huber, R. (1991). *Acta Crystallogr Sect A*. **47**, 392–400.

Engh, R. A. & Huber, R. (2001). *International Tables for Crystallography. Volume F:*
Crystallography of Biological Macromolecules, Vol. edited by M.G. Rossman & E.
Arnold, pp. 382–392. Dordrecht: Kluwer.

Furnham, N., Blundell, T. L., DePristo, M. A. & Terwilliger, T. C. (2006). *Nat. Struct. Mol.*
Biol. **13**, 184–185.

Grindon, C., Harris, S., Evans, T., Novik, K., Coveney, P. & Laughton, C. (2004). *Philos.*
Trans. R. Soc. Lond. Ser. Math. Phys. Eng. Sci. **362**, 1373–1386.

Groom, C. R., Bruno, I. J., Lightfoot, M. P. & Ward, S. C. (2016). *Acta Crystallogr. Sect. B*
Struct. Sci. Cryst. Eng. Mater. **72**, 171–179.

van Gunsteren, W. F., Dolenc, J. & Mark, A. E. (2008). *Curr. Opin. Struct. Biol.* **18**, 149–
153.

Hendrickson, W. A. & Konnert, J. H. (1980). *Computing in Crystallography*, Vol. edited by
R. Diamond, S. Ramaseshan & K. Venkatesan, pp. 13.01–13.26. Bangalore: Indian
Academy of Sciences.

Hintze, B. J., Lewis, S. M., Richardson, J. S. & Richardson, D. C. (2016). *Proteins-Struct.*
Funct. Bioinforma. **84**, 1177–1189.

Jack, A. & Levitt, M. (1978). *Acta Crystallogr. Sect. A*. **34**, 931–935.

Janowski, P. A., Cerutti, D. S., Holton, J. M. & Case, D. A. (2013). *J. Am. Chem. Soc.* **135**,
7938–7948.

Janowski, P. A., Liu, C., Deckman, J. & Case, D. A. (2015). *Protein Sci. Publ. Protein Soc.*

Liu, C., Janowski, P. A. & Case, D. A. (2015). *Biochim. Biophys. Acta*. **1850**, 1059–1071.

Lovell, S. C., Davis, I. W., Adrendall, W. B., de Bakker, P. I. W., Word, J. M., Prisant, M.
G., Richardson, J. S. & Richardson, D. C. (2003). *Proteins Struct. Funct. Bioinforma.*
50, 437–450.

Maier, J. A., Martinez, C., Kasavajhala, K., Wickstrom, L., Hauser, K. & Simmerling, C.
(2015). *J. Chem. Theory Comput.* **11**, 150707155125009.

Moriarty, N. W., Grosse-Kunstleve, R. W. & Adams, P. D. (2009). *Acta Crystallogr. Sect. -*
Biol. Crystallogr. **65**, 1074–1080.

Moriarty, N. W., Tronrud, D. E., Adams, P. D. & Karplus, P. A. (2014). *FEBS J.* **281**, 4061–
4071.

Moriarty, N. W., Tronrud, D. E., Adams, P. D. & Karplus, P. A. (2016). *Acta Crystallogr.*
Sect. -Biol. Crystallogr. **72**, 176–179.

Moulinier, L., Case, D. A. & Simonson, T. (2003). *Acta Crystallogr. D Biol. Crystallogr.* **59**,
2094–2103.

Murshudov, G. N., Skubák, P., Lebedev, A. A., Pannu, N. S., Steiner, R. A., Nicholls, R. A.,
Winn, M. D., Long, F. & Vagin, A. A. (2011). *Acta Crystallogr. D Biol. Crystallogr.*
67, 355–367.

Murshudov, G. N., Vagin, A. A. & Dodson, E. J. (1997). *Acta Crystallogr. Sect. D*. **53**, 240–
255.

Priestle, J. P. (2003). *J. Appl. Crystallogr.* **36**, 34–42.

Richardson, D. C. & Richardson, J. S. (2001). *Crystallography of Biological*
Macromolecules, Vol. *F*, edited by M.G. Rossmann & E. Arnold, p. Dortrecht:
Kluwer Academic Press.

Richardson, J. S. & Richardson, D. C. (2018). *Comput. Crystallogr. Newsl.* **9**, 21–24.

Richardson, J. S., Williams, C. J., Videau, L. L., Chen, V. B. & Richardson, D. C. (2018). *J.*
Struct. Biol. **204**, 301–312.

Roe, D. R. & Cheatham, T. E. (2013). *J. Chem. Theory Comput.* **9**, 3084–3095.

Roitberg, A. & Elber, R. (1991). *J. Chem. Phys.* **95**, 9277–9287.

Schnieders, M. J., Fenn, T. D., Pande, V. S. & Brunger, A. T. (2009). *Acta Crystallogr. D*
Biol. Crystallogr. **65**, 952–965.

Sheldrick, G. M. (2008). *Acta Crystallogr. Sect. A*. **64**, 112–122.

Showalter, S. A. & Brüschweiler, R. (2007). *J. Chem. Theory Comput.* **3**, 961–975.

Simmerling, C., Fox, T. & Kollman, P. A. (1998). *J. Am. Chem. Soc.* **120**, 5771–5782.

Touw, W. G. & Vriend, G. (2010). *Acta Crystallogr Sect D*. **66**, 1341–1350.

Tronrud, D. E., Berkholz, D. S. & Karplus, P. A. (2010). *Acta Crystallogr. D Biol.*
Crystallogr. **66**, 834–842.

Tronrud, D. E., Ten Eyck, L. F. & Matthews, B. W. (1987). *Acta Crystallogr. Sect. A*. **43**,
489–501.

Wall, M. E., Adams, P. D., Fraser, J. S. & Sauter, N. K. (2014). *Struct. Lond. Engl. 1993*. **22**,
182–184.

Waser, J. (1963). *Acta Crystallogr.* **16**, 1091–1094.

Williams, C. J., Headd, J. J., Moriarty, N. W., Prisant, M. G., Videau, L. L., Deis, L. N.,
Verma, V., Keedy, D. A., Hintze, B. J., Chen, V. B., Jain, S., Lewis, S. M., Arendall,
W. B., Snoeyink, J., Adams, P. D., Lovell, S. C., Richardson, J. S. & Richardson, D.
C. (2018). *Protein Sci.* **27**, 293–315.

Williams, C. J., Videau, L. L., Hintze, B. J., Richardson, D. C. & Richardson, J. S. (2018).
BioRxiv. 324517.

Word, J. M., Lovell, S. C., LaBean, T. H., Taylor, H. C., Zalis, M. E., Presley, B. K.,
Richardson, J. S. & Richardson, D. C. (1999). *J. Mol. Biol.* **285**, 1711–1733.

Word, J. M., Lovell, S. C., Richardson, J. S. & Richardson, D. C. (1999). *J. Mol. Biol.* **285**,
1735–1747.

York, D. M., Darden, T. A. & Pedersen, L. G. (1993). *J. Chem. Phys.* **99**, 8345–8348.

Zagrovic, B., Gattin, Z., Lau, J. K.-C., Huber, M. & van Gunsteren, W. F. (2008). *Eur.*
Biophys. J. **37**, 903–912.

Figure 1 Distribution of refined structures across resolution bins.

Figure 2 R-factors of optimized weight refinements and Rfree-Rwork (R_Δ), versus resolution

(values averaged in each resolution bin). Vertical axes are in % with R_Δ axis on the left. E&H/CDL

values are plotted in dark blue and Amber in burnt orange.

Figure 3 Comparison plots of model quality measures vs resolution, for Amber vs CDL/E&H refinements with error bars depicting the 95% confidence level of the standard error of the mean. MolProbity score is a combination of all-atom clashscore, Ramachandran favored and rotamer outliers, weighted to approximate the expected score at the structure's resolution. The hydrogen bond fraction is calculated using *cpptraj* per 1000 atoms in the model. For all 6 plots, Amber (burnt orange) differs in the better direction.

Figure 4   Fraction of Cβ deviations (in %) per Cβ atoms as a function of resolution, for the
CDL/E&H (dark blue) and Amber (burnt orange) refinements. Values are averaged in each bin of
resolution, with the error bars showing the 95% confidence level of the standard error of the mean.

576 **Figure 5** Differing responses of CDL/E&H versus Amber refinement to the misfitting of a water

577 into what should be a side chain N atom in an Arginine. Neither result here is acceptable, but if the

578 incorrect water is deleted (panel d), both methods do a very good job of moving the guanidinium

579 correctly back into its density.

Figure 6 At high resolution, Cβ deviation outliers are most often due to problems with alternate conformations. a) Amber refinement using the original Ser 215 alternates in PDB file 1nLs, which have widely separated positions for Cβ but only a single Cα atom. b) Amber refinement after the definition of alternates has been spread to include the Cα and both adjoining peptides. c) Amber refinement of the original Thr 196 of 1nLs, where alternate B had been fit backward; there is bad covalent geometry and a huge Cβd of 0.88Å (ball not shown). d) Good Amber result after altB was refit in the correct rotamer, so that all atoms match the density.

Figure 7 Unacceptable ways to get rid of a Cβ deviation without fixing the actual problem. a) 1bkr

Thr 101 as deposited, with a huge Cβd of 0.63Å (not shown as a ball because it obscures the side

chain), clashes, a rotamer outlier, the heavier Oγ branch in the lower electron-density peak and the Cβ

out of density -- all caused by modeling the side chain χ1 180° backwards. b) CDL/E&H makes the

geometry perfect but puts the Oγ far out of density. c) Amber gets all 3 side chain atoms into peaks by

making the chirality at Cβ incorrect. d) A refit in the correct rotamer replaces clashes with H-bonds,

has no outliers and puts each atom into its correct density peak.

Figure 8 A Cβ deviation in the Amber results at 3.5Å, but not for either the original or the CDL

results. a) 1xgo Leu 253 on a quite distorted helix, with many clashes and a Ramachandran outlier; the

Leu rotamer is incorrect, as shown by the 1xgs structure at 1.75Å. b) CDL/E&H refinement fixes the

clashes, but not the rotamer or Ramachandran outliers or the helix distortion. c) Amber refinement

fixes the clashes and the Ramachandran outlier, flags the incorrect Leu rotamer with a Cβd outlier and

moves the helix conformation closer to ideal. d) Leu 253 in 1xgs at 1.75Å, with a clearly correct

rotamer on an ideal helix and no outliers besides one clash.

Figure 9 Two misoriented peptides in 1xgo, flagged by Ramachandran and CaBLAM outliers (magenta outlines on the CO virtual dihedrals). a) Residues 86-91 in the deposited 1xgo structure. b) CDL/E&H result, with unchanged conformation and outliers. c) Amber result, with several peptide orientations changed by modest amounts (red balls on CO), removing the backbone outliers and very closely matching the conformation for 1xgs shown in panel d.

Figure 10   Amber refinement produces better H-bonds and van der Waals contacts as well as
removing somewhat more steric clashes. a) The Asn 182 helix-cap region in PDB file 3g8L at 2.5Å,
with numerous clashes and other outliers. b) CDL/E&H refinement makes large improvements,
removing most clashes and all other outliers. c) Amber refinement does even better, removing all
clashes and most small overlaps (yellow) and optimizing to produce more H-bonds and favorable van
der Waals contacts (green and blue dots).

Figure 11   CDL/E&H versus Amber improvements in steric contacts for the 3g8L helix-cap,
quantified by all-atom contact dot or spike counts measured in Mage (Richardson 2001), normalized
relative to the counts in the deposited 3g8L structure. Amber changes farthest, in the right direction,
for all four contact types.

Supporting information

S1. Full-dataset comparisons

Bond and angle rmsd comparisons (see figure S1) show that the bond rmsd values are numerically different but are smaller than the average sigma of 0.02Å (2pm) applied to protein bond restraints. Furthemore the Amber angle rmsd values are approximately 2° across all resolutions – also lower than the average of ~3° applied to protein angle restraints. The increased CDL/E&H rmsd values at high resolution may be result of the looser rmsd limit used past 1.5Å for the weight optimisation process. Comparing the means of the CDL/E&H and Amber rmsd values is not valid as force fields use more complex energetics rather than harmonic targets to ideal values.

S2. Response to Bad Peptide Orientations

S2.1. Background

The low-resolution analysis of Cβ deviations in the main text made use of comparing the 1xgo structure at 3.5Å (Tahirov 1998) versus 1xgs at 1.75Å from the same paper. All six Cβ deviations in the Amber results versus none from CDL/E&H were compared, finding that in each case that Cβd was flagging an underlying problem: either a misfit side chain or an incompatibility between backbone and side chain.

For the issue of bad peptide orientations, however, only one example was illustrated (Figure 9). These problems are common at resolutions worse than 2.5Å, because the backbone CO direction is no longer seen (Richardson *et al.*, 2018). Misoriented peptides are best diagnosed by CaBLAM (Williams 2018). CaBLAM uses virtual dihedral angles of successive Cαs and of successive COs to test whether the orientations of successive CO groups are compatible with the surrounding Cα trace. It flags outliers graphically in magenta on the CO-CO virtual dihedral. Since typically there is an energy barrier between widely different peptide orientations, the presumption is that refinement cannot easily correct these cases. However, that presumption needs to be tested.

S2.2. Most are not correctable by refinement

Ten cases were identified in 1xgo, for isolated single or double CaBLAM outliers (usually with other outliers also), surrounded by correct structure as judged in the same molecule at 1.75Å resolution (1xgs). For 6 of those 10 cases, neither CDL/E&H nor Amber refinement corrected the problem (His62, Thr70, Gly163, Gly193, Ala217, Glu286).

For example, figure S2 shows stereo images of the Glu286-Lys287 hairpin-loop case, where the CaBLAM outlier in 1xgo is accompanied by clashes, Ramachandran and rotamer outliers. Both

CDL/E&H and Amber conformations are essentially identical to the original 1xgo, with no peptide

improvement. They both remove all the clashes (clusters of hotpink spikes) and remove one of the six

side chain outliers (gold) but not into the correct rotamer. In contrast, the high-resolution 1xgs, with

very clear electron density (bottom panel), shows the Lys Cα and the two peptide carbonyl oxygens

(red balls) differently placed by large distances and dihedral angles, forming a well H-bonded β-

hairpin with no outliers of any kind.

S2.3. Other Outliers Often Better

In two cases the CDL/E&H results had fewer other outliers than Amber, although it did not actually

reorient the peptide CO (Gly163, Gly193). The Gly163 case is shown in stereo in figure S3, for an S-

shaped loop between non-adjacent β-strands, with two CaBLAM flags (magenta) and many other

outliers. Both refinements remove the clashes, one of the rotamer outliers and one of the

Ramachandran outliers (green). The CDL/E&H results in addition removed one of the CaBLAM

outliers and the Cα-geometry outlier (red). However, neither refinement could manage the large

rotation needed to correct the 163-164 peptide orientation, as judged by the more convincing

conformation of the high-resolution 1xgs at bottom.

S2.4. Amber Sometimes Corrects Well

In three cases Amber managed a complete fix, while in contrast CDL/E&H did not improve (Asp88,

Gly125, Pro266). The Asp88-Gly89 tight turn example is shown in Figure 9 of the main text.

Here in figure S4, the Gly125 loop example in a helix-helix connection is shown in stereo, to allow

clear visualization of the CO orientation changes. 1xgo residues 121-126 (figure S3a) have two

CaBLAM outliers (magenta dihedral lines) unchanged by CDL/E&H refinement (panel b). However,

Amber refinement (panel c) manages to shift several CO orientations by up to 80° (red balls), enough

to fix the CaBLAM outliers and to match extremely closely the better backbone conformation of 1xgs

(panel d).

S2.5. A Partial Correction, Unconverged

Finally, in one especially interesting case (Lys22, in Figure S5a for 1xgo) Amber turned the CO (red

circles) about halfway up to where it should be (panels b vs c), while CDL/E&H made no

improvement to the peptide. The Amber model eliminated the Ramachandran and one of the

CaBLAM outliers, but still had geometry outliers (a bond angle and a Cβ deviation). It seemed likely

that Amber refinement had not fully converged and might move the CO all the way if run longer.

A 30-cycle Amber run had earlier been done for 1xgo, without any major changes noticed beyond the

10-cycle. From that endpoint, two further runs were done, first of 30 cycles ("Amber60"), then a

further 10 cycles ("Amber70").

Figure S5d shows the fan of CO positions for all 7 of the deposits and refinements, progressively

rotating counterclockwise from 1xgo to 1xgs. Indeed, both Amber60 and Amber70 successfully

rotated the Lys22 peptide almost all the way to the good helical position seen in the high-resolution

1xsg (panel e), eliminating both the CaBLAM outlier and the intermediate-stage bond-angle outliers,

presumably having crossed an energy barrier in the process.

One other CaBLAM-outlier peptide was corrected in Amber70 as well (Thr71). But for the Ala217

outlier, the wrong peptide was rotated, seduced by H-bonding to an Arg side chain in the wrong

position.

In these long refinements, both R-factors and match to electron density suffer somewhat. In the cases

examined, this often seems due to incorrect side chain rotamers (almost never correctable by

refinement) pushing an otherwise-good backbone conformation a bit out of density (translated

upward, for 1xgo Lys22). Future work will try to guide early correction of as many problems as

feasible, for the faster and more successful refinement afterward that we now know is possible.

S2.6. Discussion

In summary, it is indeed true that refinement cannot usually correct a peptide orientation that is off by

a large amount. The very tight geometry restraints in the CDL/E&H system presumably raise the

barriers to peptide rotation. Amber is rather better at that, and about 1/3 of the time managed a good

correction, although convergence can be very slow for such large changes. We feel it is crucial to try

correcting problems such as flipped peptides in the initial model before refining it, however, crosstalk

between backbone and side chains further complicates that process. However, we are enthusiastic

about use of the Amber target to realistically improve conformation and especially sterics, once the

model is mostly in the right local minima.

S2.7. References

Richardson J, Richardson D (2018) Cβ deviations and other aspects in Amber versus CDL
refinements, *Comp. Cryst. Newsletter* **9**: 21-24

Tahirov TH, Oki H, Tsukihara T, Ogasahara K, Yutani K, Ogata K, Izu Y, Tsunasawa S, Kato I
(1998) Crystal structure of methionine aminopeptidase from hyperthermophile *Pyrococcus*
furiosus, J. Mol. Biol. **284**: 101-124 [1gxo, 1gsx]

Richardson JS, Williams CJ, Videau LL, Chen VB, Richardson DC (2018) "Assessment of detailed
conformations suggests strategies for improving cryoEM models: helix at lower resolution,
ensembles, pre-refinement fixups, and validation at a multi-residue length scale", *J Struct. Biol.*
204: 301-312

Williams CJ, Hintze BJ, Headd JJ, Moriarty NW, Chen VB, Jain S, Prisant MG Lewis SM, Videau
LL, Keedy DA, Deis LN, Arendall WB III, Verma V, Snoeyink JS, Adams PD, Lovell SC,
Richardson JS, Richardson DC (2018) MolProbity: More and better reference data for improved
all-atom structure validation, *Protein Sci.* **27**: 293–315 [CaBLAM]

Figure S1 Bond and angle rmsd values for CDL/E&H (dark blue) and Amber (burnt orange) plotted

against resolution.

1xgo
3.5Å

CDL/E&H

Amber

1xgs
1.75Å

Figure S2 Stereo images of uncorrected CaBLAM problems for the beta-hairpin loop at Glu 286 -

Lys 287 in 1xgo at 3.5Å resolution. a) As deposited, with outliers for CaBLAM (magenta lines on the

CO dihedral), CaBLAM Cα-geometry (red lines on Cα trace), Ramachandran (green lines along

backbone), rotamer (gold sidechains), and all-atom clash (clusters of hot-pink spikes) evaluations. b)

As refined by Phenix CDL/E&H and c) as refined by Phenix Amber, both of which remove the

clashes but do not correct the underlying conformation. d) In the 1xgs structure at 1.75Å resolution,

showing a classic, outlier-free beta hairpin conformation with good backbone H-bonding and

substantial corrections in peptide orientation and sidechain placement. The 286 and 287 peptide

oxygens that move most are circled in red.

Figure S3   Partial correction of an S-shaped loop at 159-164 in 1xgo. a) As deposited, with many

types of outliers. b) CDL/E&H corrects all but two backbone outliers. c) Amber corrects all clashes

but few other outliers, and neither refinement changes the poor underlying conformation. d) The 1xgs

structure achieves an outlier-free, well H-bonded conformation by shifting 4 peptide orientations (red

ball on carbonyl O atoms), especially at Gly 163.

Figure S4 Successful Amber CaBLAM corrections in the helix-helix loop at 1xgo 121-126. a) As

deposited, with clashes and two CaBLAM outliers. a) CDL/E&H corrects the clashes but not the

backbone conformation. b) Amber reorients 3 successive peptides (red balls on peptide Os) by up to

80°, removing both CaBLAM outliers and matching extremely closely the conformation seen at high

resolution in panel d.

Figure S5 Gradual correction of the helix C-cap at 1xgo Lys 22. a) As deposited, with double
CaBLAM outliers, clashes, and Ramachandran outlier. CDL/E&H refinement fixes clashes but leaves
conformation unchanged. b) Amber refinement moves the crucial Lys 22 CO partway up toward α-
helical orientation, relieving one of the CaBLAM outiers. c) Helical, outlier-free conformation of the
C-cap region in 1xgs at high resolution. d) Superposition in side view, showing all Lys 22 CO
orientations between 1xgo outlier and 1xgs α-helical: longer Amber refinement progressively corrects

770   the orientation, converging close to the 1xgs orientation although with a translational shift we believe

771   is an effect of incorrect sidechain rotamers.