Pedigree-based measurement of the de novo mutation rate in the gray mouse lemur

reveals a high mutation rate, few mutations in CpG sites, and a weak sex bias

Authors: C. Ryan Campbell[1*], George P. Tiley[1], Jelmer W. Poelstra[1], Kelsie E. Hunnicutt[1,†],

Peter A. Larsen[1,‡], Mario dos Reis[2], & Anne D. Yoder[1]

[1]Department of Biology, Duke University, Durham, NC 27708, USA

[2]School of Biological and Chemical Sciences, Queen Mary University of London, Mile End

Road, London E1 4NS, UK

[†]Present address: Department of Biological Sciences, University of Denver, Denver, CO 80208

[‡]Present address: Department of Veterinary and Biomedical Sciences, University of

Minnesota, St. Paul, MN 55108

*Author for Correspondence: C. Ryan Campbell, Department of Evolutionary Anthropology,

Duke University, Durham, USA, c.ryan.campbell@duke.edu

Abstract

Spontaneous germline mutations are the raw material on which evolution acts and knowledge of their frequency and genomic distribution is therefore crucial for understanding how evolution operates at both long and short timescales. At present, the rate and spectrum of de novo mutations have been directly characterized in only a few lineages, and it is therefore critical to examine a wide range of species to determine the generality of patterns that have been identified so far. Our study provides the first direct mutation rate estimate for strepsirrhine primates (i.e., the lemurs and lorises) which comprise nearly half of the primate clade. Using high-coverage linked-read sequencing of a family pedigree (n = 8) of gray mouse lemurs (*Microcebus murinus*), we estimate the mutation rate to be 1.64×10^{-8} (95% CI: 1.41×10^{-8} to 1.98×10^{-8}) mutations per basepair per generation. This estimate is higher than those for most previously characterized mammals, including other primates. Contrary to expectation, we found only a modest overrepresentation of mutations at CpG-sites and of paternal mutations. Comparing mutation rates across taxa, we show that expectations based on the drift barrier hypothesis are met at a broad phylogenetic scale but not within primates. Finally, we compared pedigree-based mutation rates with phylogenetically-based substitution rate estimates for mouse lemurs and six other primate lineages and found considerable differences between the two rate estimates. This finding has implications for divergence-time estimation and calls for further study.

Introduction

Spontaneous germline mutations (SGMs) are single basepair errors that occur in DNA transmission as it is passed from parent to offspring in sexually reproducing organisms. The accrual of these errors provides not only the raw material for evolution but can also serve as a measure of the evolutionary time across phylogeny (Zuckerkandl and Pauling 1965; Kimura and Ohta 1971; Langley and Fitch 1974). The rate at which these mutations are introduced into genomes is thus a crucial metric of evolution at the genomic level as well as a measure of fundamental biological processes (Kondrashov and Kondrashov 2010). And by further parsing this measure by genomic regions, we gain an intimate perspective of intragenomic evolutionary patterns. Finally, by quantifying the variation in mutation rates across the tree of life, we can develop increasingly refined hypotheses regarding the relationship between mutation rates and life history characteristics.

Approaches for estimating mutation rates in vertebrates generally fall into one of two categories: phylogenetically-based (indirect) versus pedigree-based (direct) estimation. While phylogenetically-based methods were the standard before the genomics revolution of the past decade, new sequencing technologies have enabled pedigree-based mutation rate measurements for previously uncharacterized species. By comparing the genomes of individuals with known genealogical relationships – typically, parent to offspring – investigators are putatively able to quantify mutations in real time (Scally and Durbin 2012; Smeds, et al. 2016; Feng, et al. 2017; Pfeifer 2017; Thomas, et al. 2018; Koch, et al. 2019). Phylogenetically-based approaches, on the other hand, are dependent on external calibrations to obtain substitution rates in units of absolute time. In a typical study, divergence between homologous genomic regions across a phylogeny is calibrated with fossils or other external criteria at specific nodes in order to determine substitution rates (Thorne, et al. 1998; Sanderson 2002; Thorne and Kishino 2002; Drummond, et al. 2006; Yang 2006), relying on the assumption that the rate at which substitutions accumulate between species is equal to the mutation rate for neutral sites (Kimura 1983). Thus, in principle, phylogenetically-based and pedigree-based methods should produce equivalent estimates of the mutation rate.

Phylogenetically-based methods are known to suffer from various sources of uncertainty including violation of the molecular clock, inaccuracies in external calibration points, incomplete lineage sorting, and the difficulties of recovering multiple overlapping changes (i.e., "multiple hits") at any given site (dos Reis, et al. 2018). Although a number of solutions to these problems have been proposed (Heath, et al. 2014; Ogilvie, et al. 2017), some limitations are difficult to overcome (Magallóan and Sanderson 2005; Near, et al. 2005; Herrera and Dávalos 2016). Pedigree-based mutation rate estimates are not affected by the same uncertainties, however, and allow for the characterization of mutation rate variation across the genome and the various biases in the mutation spectrum (Ségurel, et al. 2014; Harris, et al. 2017). Previously, these estimates have relied on well-assembled genomes available only to model organisms (Scally and Durbin 2012; Venn, et al. 2014; Uchimura, et al. 2015; Jónsson, et al. 2017), and have therefore been limited in taxonomic scope (for a list of mammals, see `Table 1`). Fortunately, recent genome assembly strategies (Dudchenko, et al. 2017) have increasingly enabled chromosome-level assemblies of non-model organisms (Larsen, et al. 2017) and pedigree-based mutation rate estimation is now feasible for virtually any species, as long as related individuals with known pedigrees are available (Smeds, et al. 2016; Feng, et al. 2017; Pfeifer 2017; Martin, et al. 2018; Koch, et al. 2019). Pedigree-based mutation rate estimates from across the tree of life have the potential to enhance our understanding of mutation rate evolution (Lynch 2010), including the relationship between mutation rate and genome architecture (Kim, et al. 2006).

Even so, pedigree-based studies are not without challenges. Perhaps most challenging for estimating the number of de novo mutations in offspring is that the mutation rate is several orders of magnitude lower than the sequencing error rate, even for the most accurate sequencing methods. Thus, pedigree-based estimates of the SGM rate must employ stringent bioinformatic filters to distinguish between true biological mutations and sequencing error. And herein lies one of the method's most significant challenges: stringency can be so vigorously applied that many true SGMs are not called (i.e., false negatives are common), and somewhat counterintuitively, the mutation rate can be under- rather than overestimated.

Here, we utilize two strategies for keeping false negative rates low while ensuring the validity of our called mutations. First, linked short reads from 10x Genomics (Weisenfeld, et al. 2017) provide increased mapping scores and confidence in calling individual variants when coupled with mutation rate estimation methods (Long, et al. 2016; Winter, et al. 2018), especially in repeat-rich mammalian genomes (Chaisson, et al. 2015). Additionally, the phasing information provided by linked reads can determine the parent of origin through just two generations of sequencing. Phased haplotypes with known parental origin then allow us to assign individual mutations as coming from either the maternal or paternal germline. To control for false negatives, we explicitly evaluate the callable proportion of the genome by introducing known false mutations to the sequencing data for one individual and subsequently testing how accurate our bioinformatic pipeline was in recovering these errors (Keightley, et al. 2014; Xie, et al. 2016). We refer to this approach as "allele drop", which can be used to correct observed SGM rates regardless of the variant calling pipeline.

We applied these advances in sequencing technology and computational methods to produce the first pedigree-based estimated mutation rate for the gray mouse lemur (*Microcebus murinus*), the first strepsirrhine primate to be characterized in a pedigree-based SGM study. The gray mouse lemur is a member of a radiation of morphologically cryptic primates distributed throughout Madagascar (Hotaling, et al. 2016), for which an accurate temporal context is needed to, for instance, for comparing the timing of speciation with historical climatic events. Previous divergence time studies have had to rely on either distant external fossil calibrations (Yang and Yoder 2003; dos Reis, et al. 2018) or by using pedigree-based mutation rate estimates from distant relatives (Yoder, et al. 2016).

By measuring the mutation rate in mouse lemurs with a pedigree-based approach, we aim to simultaneously expand our knowledge of mutation rate variation across lineages and to facilitate the estimation of divergence times within this specific radiation. To do so, we deeply sequenced a family pedigree of gray mouse lemurs (*Microcebus murinus*, n = 8), including a mother, father, and two offspring, to accurately identify and count SGMs and to assign the parent-of-origin of de novo mutations.

Results and Discussion

Mutation Rate

To identify de novo mutations, we used DeNovoGear (Ramu, et al. 2013) and validated the inferred mutations using an independent mutation identification analysis with VarScan2 (Koboldt, et al. 2009). In total, we assessed 4,542,770 potential variants across eight related individuals to discover 134 de novo mutations in two focal offspring, one male and one female (*Fig 1*). Among these 134 mutations, 125 (71 in the female and 63 in the male) were located on autosomes and nine (seven in the female and two in the male) were located on the X chromosome. The average depth of coverage for the 134 mutations was 170 reads (170.2, SD=79.95). We estimate the mutation rate in this family quartet to be 1.64×10^{-8} single nucleotide mutations per basepair per generation per individual. The single-base mutation-rate estimate is a weighted average of the number of mutations on the autosomes (125 mutations) and X chromosome (nine mutations) divided by the number of sites where a mutation was detectable given our methods (Long, et al. 2016; Winter, et al. 2018).

To estimate the number of detectable sites, we artificially generated 1,000 mutations placed randomly across the entire genome within the sequence data of the offspring, similar to previous efforts to account for false negative results (Keightley, et al. 2014; Xie, et al. 2016). On the autosomes, we detected 801 of 952 generated mutations, while 39 of 48 generated mutations were detected on the X chromosome. Therefore, we estimate our detection rate to be 84.1% on autosomes and 81.3% on the X chromosome, which yields a total of 2.088 billion detectable sites (out of a total of 2.487 billion).

To calculate the confidence interval around the estimated mutation rate, we assumed that the total number of mutations inherited by an offspring follows a Poisson distribution (*Supplemental Text 1*). Given that an average of 67 mutations were found in each of the focal offspring, the 95% confidence interval is 56.5 to 79.1 mutations per genome, or 1.41×10^{-8} to 1.98×10^{-8} mutations per site per generation. To calculate the expected number of false positive and false negative results, we sequenced the male offspring twice and compared variant presence and absence across the technical replicates (*Supplemental Text 2*). We

calculated 6.46 false positives and 37.51 false negatives from the total of 134 de novo

mutations and 2.088 billion potential mutation sites.

Relatively low numbers of mutations at CpG sites

CpG sites have consistently been found to have higher mutation rates relative to other site

classes, a pattern discovered using sequence comparisons made several decades ago (Bird

1980) and ascribed to the frequent deamination of methylated cytosines (Friedberg, et al.

2005). We found a total of six mutations at parental CpG sites, constituting 4.5% of all de novo

mutations, a more than two-fold overrepresentation given that 1.9% of the genome consists of

CpG sites. This pattern is much less pronounced than in other studies, were CpG site

mutations have been shown to contribute 12-25% of total mutations, representing an

overrepresentation of least ten-fold given the number of CpG sites in the genome (Venn, et al.

2014; Smeds, et al. 2016; Thomas, et al. 2018; Besenbacher, et al. 2019; Gao, et al. 2019).

Given the limited sample size of our study, and the small fraction of the genome this

represents, however, caution is warranted in interpreting this result. Confidence intervals

calculated following the genome-wide Poisson distribution example (as in *Supplemental Text*

1) suggest that the true number of CpG site mutations in the mouse lemur genome could be as

high as 10.9 (over 8% of total mutations and a 4.25-fold increase). Additionally, if the true CpG

site mutation rate in mouse lemurs was such that 10% of total mutations were found there, the

values measured in this study would still fall within the 95% confidence interval (13.4 CpG site

mutations, 95% confidence interval 6.0 to 21).

The elevated mutation rate at CpG sites is linked to methylation, and is therefore not expected

in regions of the genome with high GC content (CpG islands), where CpG sites are much less

likely be methylated. As expected, none of the 134 de novo mutations were found within CpG

islands, which constitute roughly 4% of the *Microcebus murinus* genome.

The discrepancy in the relative frequency of mutations at CpG sites between mouse lemurs

and other studied species could have arisen from a high proximity of CpG sites to genic

regions, where mutations are more likely to be detrimental. A lower substitution rate in

proximity to genes has been observed empirically (Narang and Wilson Sayres 2016).

Accordingly, we investigated this possibility, but found a normal distance between genes and

CG sites (*Fig S1*). Also to be considered, the mouse lemur genomes here compared are of

lesser overall quality than human or chimpanzee, and a lower mapping quality in CG-rich

regions could be affecting our ability to detect CpG site mutations.

Finally, the low number of CpG site mutations has the statistical effect of impacting additional

mutational metrics: transition-transversion ratio, strong-to-weak and weak-to-strong ratio, and

potentially, sex-bias.

Mutation spectrum

We characterized two aspects of the mutation spectrum in mouse lemurs. First, the ratio of

transitions to transversions (Ti:Tv) was estimated to be 1.03 (68 transitions and 66

transversions). Second, the ratio of strong-to-weak mutations (SW; C/G > A/T) to weak-to-

strong mutations (WS; A/T > C/G), SW:WS, was also estimated to be 1.03 (31 SW and 30 WS

mutations). The most common two categories of de novo mutation type were G>A and C>T

(*Fig 2*).

Our estimates of the Ti:Tv and SW:WS are lower than values found in other animals. For

instance, values of Ti:Tv in previous pedigree-based studies varied between 1.97 and 2.67

(Agier and Fischer 2011; Kong, et al. 2012; Venn, et al. 2014; Smeds, et al. 2016; Assaf, et al.

2017; Thomas, et al. 2018; Besenbacher, et al. 2019). Relatively low values for these ratios

are expected in light of the low CpG mutation rate reported above, given that C>T mutations at

CpG sites are strong-to-weak transitions. For instance, without an elevation in the mutation

rate at CpG sites, the Ti:Tv and SW:WS ratios would drop to 1.55 and 1.33 respectively, in a

study of chimpanzees (Venn, et al. 2014). Thus, it is difficult to differentiate the effect of

reduced CpG site mutation on the other metrics of the mutation spectrum.

Reduced Male Mutational Bias

A male mutational bias has long been hypothesized for diploid sexually reproducing organisms

based on the idea that the increased number of cell divisions in sperm versus egg should lead

to higher numbers of mutations in the male germline than the female germline (Haldane 1946;

Kong, et al. 2012; Lindsay, et al. 2018). Indeed, a strong male mutation rate bias has been

observed in the vast majority of pedigree-based mutation rate estimates to date (Venn, et al. 2014; Rahbari, et al. 2015; Lindsay, et al. 2018; Thomas, et al. 2018; Gao, et al. 2019) and in many studies of phylogenetically-based rates (Shimmin, et al. 1993; Ellegren and Fridolfsson 1997; Axelsson, et al. 2004; Zhang 2004; Goetting-Minesky and Makova 2006).

Using the long phasing blocks generated by the linked-read method, we were able to determine the parent of origin for 94 out of 134 (70%) de novo mutations. Thus, the number of mutations confidently assigned to a parent are notably higher in our analysis compared to previous studies that used short read sequencing alone, such as 35% (Venn, et al. 2014) or 38% (Thomas, et al. 2018). In our study, among the assigned mutations, 54% (n = 51) were found on the offspring's paternal haplotype while the remaining 46% (n = 43) were found on the offsprings' maternal haplotype, a ratio of male-to-female mutations of approximately 1.2.

This ratio of male-to-female mutations in gray mouse lemur is therefore considerably less pronounced than that observed in other primate species (2.1-5.5) (Venn, et al. 2014; Rahbari, et al. 2015; Thomas, et al. 2018; Besenbacher, et al. 2019) and mouse (2.7) (Lindsay, et al. 2018). However, it is identical to the ratio found in collared flycatchers (Smeds, et al. 2016). One of the driving factors of the male mutational bias is likely the time of first reproduction after puberty (Ségurel, et al. 2014), and given that mouse lemurs experience puberty and time of first reproduction nearly simultaneously (Blanco 2011; Zohdy, et al. 2014; Blanco, et al. 2015), this may explain our low ratio of male-to-female mutations. Additionally, there are stark differences in the methylation process within male and female germline cells, with male cells experiencing markedly more methylation (Reik and Dean 2001; Kobayashi, et al. 2013). This discrepancy yields more methylation-related mutations in males than females as mammals age (Gao, et al. 2019). Thus, in the absence of these methylation-focused mutations, and with a relatively short time to puberty, the mouse lemur would have multiple factors pointing to a more balanced mutational sex ratio, as was observed. On the other hand, however, mouse lemurs have exaggerated symptoms of sperm competition (Eberle and Kappeler 2004), large testes size (Wong 2014), which might be expected to lead to elevated male mutation rates, thus further complicating the interpretation the a low male bias evident in our study.

Relationship of Mutation Rate Estimates to Life History Traits

The drift barrier hypothesis states that effective population size may explain some variation in mutation rates across species due to a larger efficiency of selection acting on DNA replication fidelity in larger populations, especially across large phylogenetic distances (Lynch 2010; Sung, et al. 2012). Using an MSMC analysis (Schiffels and Durbin 2014), we estimated the effective population size of gray mouse lemur across the last 2 million years to be approximately 41,000 (*Fig S2*). When analyzed in a broad phylogenetic context, we find that the mouse lemur lies comfortably along the regression that shows a negative correlation between effective population size and per-generation mutation rate across animals (*Fig 3*). In stark contrast, this relationship is not upheld in a comparison with other primates with the two species with the largest N_e (orangutan and mouse lemur) also showing the highest mutation rate (*Fig 3*). It is possible that the drift barrier hypothesis does not operate across these relatively small timescales and differences in population size. Moreover, mouse lemurs are the only strepsirrhine primate represented by a pedigree-based analysis of mutation rate and are considerably diverged from the other primates analyzed with a shared common ancestor ≥ 60 million years ago (dos Reis, et al. 2018). Further insight into the relationship between mutation rate, effective population size, and other life history traits within primates thus awaits mutation rate estimates of additional species, especially within the strepsirrhine clade.

A negative correlation between per-year mutation rates and generation time has been long-observed in studies of molecular evolution (Wu and Li 1985; Ohta 1993; Li, et al. 1996; Goetting-Minesky and Makova 2006) and underlies the classic hominid slow-down hypothesis (Goodman 1962; Moorjani, et al. 2016). Compared to other primates, mouse lemurs have both a high per-year mutation rate and a short generation time (3.75 y) (Yoder, et al. 2016), consistent with this general expectation. Across primates, the relationship between per-year mutation rates and generation time is present but weak (*Fig 4*).

Discrepancies between pedigree-based and phylogenetically-based estimates

One revelation in the rapidly developing literature on SGM rates has been that the estimated de novo rate in humans is less than half that predicted by phylogenetic studies, prompting speculation that mutation rates may have changed rapidly among primates (Scally and Durbin

2012). This observation has created some alarm, especially with regard to the impacts on

divergence age estimates for hominids (Besenbacher, et al. 2019). Therefore, we compared

pedigree-based estimates of mutation rates for mouse lemurs and five other primates (humans

Homo sapiens, chimpanzee *Pan troglodytes*, gorilla *Gorilla gorilla*, orangutan *Pongo abeli*,

green monkey *Chlorocebus sabaeus*, and owl monkey *Aotus nancymaae*) with

phylogenetically-based estimates from a recent analysis of primate divergence times (dos

Reis, et al. 2018) (*Table S1*, *Fig 4*).

Phylogenetic substitution rates are estimated per-year, so we rescaled them by generation

time for comparison with pedigree-based per-generation estimates of mutation rates. Although

such estimates introduce uncertainty from generation time estimation as well as fossil

calibration error, there are two notable observations from the comparison: 1) point estimates

for pedigree-based and phylogenetically-based estimates differ considerably for all species,

with pedigree-based estimates falling outside of the confidence intervals of phylogenetically-

based estimates in 4 out of 7 cases; 2) the direction of disagreement varies across lineages:

whereas pedigree-based estimates are higher than phylogenetically-based estimates for

mouse lemur, owl monkey, green monkey, and orangutan, the opposite is true for human,

chimp, and gorilla. Thus, were it not for the result observed in orangutans, it would be tempting

to speculate that phylogenetically-based methods have overestimated mutation rates for

hominids.

There are several potential explanations for the widespread disagreement between pedigree-

based and phylogenetically-based estimates. Phylogenetically-based estimates may be biased

downwards if substitutions are not fully neutral. Even mutations at third-codon positions may

be affected by weak negative selection and may therefore be less likely to be fixed at the

population level. Indeed, previous studies have found evidence for low phylogenetically-based

compared to pedigree-based estimates (Denver, et al. 2000; Howell, et al. 2003; Winter, et al.

2018). For pedigree-based estimates, the degree to which somatic mutations and/or inter-

individual variation might impact these estimates is as yet unclear (Ségurel, et al. 2014).

Additional analyses will be needed to reconcile the differences between pedigree-based and

phylogenetically-based estimates of the mutation rate making this an area ripe for study.

Effect of mutation rate on divergence time mutation rate estimates

Using the estimated de novo mutation rate, we have recalculated branch lengths in absolute

time for a genus-level phylogeny of mouse lemurs (Yoder, et al. 2016). Applying the newly

estimated rate of 1.64×10^{-8}, divergence time estimates are calculated to be considerably

more recent compared to previous estimates (*Fig 5*), and the smaller confidence interval

reduces uncertainty in age estimates (*Table S2*). Similarly, a recent investigation of great ape

divergence times revealed that lineage-specific mutation rate estimates for chimp, gorilla, and

orangutan led to more recent estimates of common ancestor ages that more closely agreed

with fossil evidence than if the human mutation rate was assumed for all lineages

(Besenbacher, et al. 2019). Given the endangered status of mouse lemurs and virtually all

other strepsirrhine species, our ability to provide a temporal context to speciation and to

estimate demographic parameters such as effective population size may yield critical

information for directing conservation policy and efforts.

Conclusions

In this study, we provide a pedigree-based estimate of the de novo mutation rate for gray

mouse lemurs. The results underscore the importance of measuring mutation rate with linked-

read sequencing technology that increases mapping confidence, provides haplotypes for

determining parent-of-origin, thereby eliminating the need to sequence across more than two

generations. Furthermore, we have used allele-drop simulations to determine the appropriate

denominator for mutation rate calculations. Even so, despite our best efforts to develop an

accurate method for determining the de novo mutation rate, our results are not immune to

problems presented by small sample sizes, in particular those relating to the accurate recovery

of the mutation spectrum.

Our study also emphasizes the importance of measuring mutation rate across the tree of life in

order to study the nature and causes of mutation rate evolution. As noted above, this is the first

pedigree-based mutation rate estimate for a strepsirrhine primate, and as such, it is not clear

whether the unusually high mutation rate, low CpG mutation rate, and weak sex bias may be

representative of strepsirrhines. We show that N_e correlates with mutation rate at broad

344 phylogenetic scales but does not appear to do so within primates. Finally, we demonstrate

345 widespread disagreement between phylogenetically-based and pedigree-based estimates of

346 the mutation rate, suggesting that reconciling these methods should be a focus of future work.

347 It is our hope that the details of our study will provide an encouraging pathway for that work.

Acknowledgments

We thank the Duke Lemur Center staff, especially Erin Ehmke, Bobby Schopler, and Cathy Williams, for providing the *Microcebus murinus* and *Mirza zaza* tissue samples. CRC thanks the members of his dissertation committee, Charlie Nunn, Tom Mitchell Olds, Jeff Thorne, and Marcy Uyenoyama for their insightful comments and suggestions for improvement throughout the course of this project as well as Jonathan Pritchard, Matthew Hahn, Priya Moorjani, and Felix Wu for helpful feedback during a poster presentation of this research. We thank Simon Gregory's lab for preparing the 10X Genomics libraries and are grateful for the support of Duke Research Computing and the Duke Data Commons (NIH 1S10OD018164-01). This study was funded by a National Science Foundation Grant DEB-1354610 and Duke University startup funds to ADY and she gratefully acknowledges support from the John Simon Guggenheim Foundation and the Alexander von Humboldt Foundation during the writing phase of this project. This is Duke Lemur Center publication no. XXX.

Materials and Methods

Samples

Eight individuals were selected from the DLC's mouse lemur colony consisting of a focal family of four, two parents and two offspring from separate litters, an additional half-sibling to the offspring, and three other individuals in the maternal lineage. Four of the eight selected samples were colony founders, which had been transferred from the mouse lemur colony in Brunois, Paris, France in 2003.

Blood and tissue samples were collected from these individuals during annual veterinary check-ups. High molecular weight DNA was extracted with the Qiagen MagAttract kit (Qiagen, Germantown, MD, USA) and 10X Genomics library preparation was performed at the Duke Molecular Genomics Core.

Sequencing

The eight individuals were used to produce nine sequencing libraries; every individual was sequenced once, and the focal paternal sample was prepared twice and sequenced as two separate samples. Libraries were sequenced at the Duke Center for Genomic Computational

Biology (GCB) Sequencing and Genomic Technology Shared Resource across two flowcells

(nine total lanes) of a HiSeq 4000. The run was 150 basepairs, paired end with an average

insert size of 554bp (range: 527-574bp). A single lane was run as a test of the 10x Genomics

LongRanger analysis software and was analyzed to confirm successful indexing and

preparation of the samples. After confirmation, the additional eight lanes (an entire flowcell)

were run. All data was combined for downstream analyses (SRA). 933,337,210,328 bases

were generated across nine samples and nine lanes.

10x Genomics Pipeline

Basecall (.bcl) files were demultiplexed and analysed using 10x Genomic's LongRanger v2.2.1

pipeline. Average genomic coverage after filtering was 34.5x across the nine samples. The

LongRanger pipeline was run on the Duke Scalable Computing Resource for approximately

2,000 hours (10 days of compute time per sample). Sequences were aligned to the latest gray

mouse lemur genome assembly (version 3.0, GCF_000165445.3) and variant calling was

handled by the Genomic Analysis Toolkit (v3.8), implemented within LongRanger v2.2.1 (Van

der Auwera, et al. 2013; Weisenfeld, et al. 2017). The mean N50 scaffold length, across

samples, generated within the 10x Genomics LongRanger alignment pipeline was 1.18Mb.

DeNovoGear

LongRanger alignments were used to find de novo mutations within the two focal family

offspring. Several methods were used to in finding mutations. First, the software package

DeNovoGear (Ramu, et al. 2013) was used to analyze the LongRanger variant call files (.vcf)

with default settings. As a follow-up, the software package VarScan2 was used with

LongRanger binary alignment files (.bam) to confirm that the de novo mutations found with

DeNovoGear could be confirmed.

Finally, de novo mutation hits were confirmed by checking independently with each replicate of

the sire. Hits were also checked for absence in all other pedigree individuals as well as

additionally sequenced samples from a diversity panel of gray mouse lemurs. The final list of

mutations was filtered for quality by offspring de novo quality (min DNQ of 100), offspring map

quality (min MQ of 50), and depth of coverage (min coverage of 20) in the parents.

Detectable Mutation Test

We conducted a test to determine the true number of sites at which a mutation would be detectable, hereafter referred to as an "allele drop test" (Long, et al. 2016; Winter, et al. 2018). This test increases the accuracy of the number of base pairs used in the denominator of the mutation rate calculation. It estimates the proportion of sites at which a de novo mutation would have been discoverable by our methods. This test consisted of adding 1,000 fake de novo mutations into the pedigree with the software BAMsurgeon (Ewing, et al. 2015). These mutations were added as heterozygotes, by changing half the aligned bases in the bam file at a site to the non-reference allele. After this manipulation, the same methods were used to find de novo mutations, and the results were mined for the 1,000 sites. By conducting this "allele drop test", we were able to estimate in which fraction of the genome de novo mutations would have been found, were they to have occurred, or detectable sites. Going forward with the number of detectable sites as the denominator in the mutation rate calculation yields a more accurate measure of the rate than using total bases, total autosomal bases, or even calculations based on a minimum depth of coverage because it produces an accurate picture of where mutations can be found specifically via our methods.

CpG estimates

CpG islands were identified by two independent methods and compared to measure the number of mutations within them. First, the software EMBOSS:cpgplot (Chojnacki, et al. 2017) was used on the latest gray mouse lemur genome (3.0, GCF_000165445.3) to identify regions that met the threshold of a CpG island (200 bp, over 50% CG content). Then, to confirm these annotations a fasta file of CpG-annotated regions of gray mouse lemur genome 2.0 (GCF_000165445.2) was downloaded from the UCSC genome browser. A blast (Altschul, et al. 1990) database of the latest gray mouse lemur genome (3.0, GCF_000165445.3) was created and the annotations were queried to determine the coordinates of the 2.0 CpG annotations in the 3.0 genome release used for mapping and assembly. While the concordance between these methods was over 95% out of an abundance of caution only the CpG islands (a total of 67,673 annotations) confirmed via both methods were used in the analyses.

With these coordinates in-house bash were used to determine the location of each mutation

with respect to CpG islands and coding regions. To measure distribution of mutations relative

to the null expectation, a permutation test was conducted of the 1,000 false mutations

generated for the allele drop test. A set of de novo mutations were randomly sampled, without

replacement, to compare the distance to CpG islands with the true set.

Parent-of-origin

Custom in-house bash scripts were used in combination with the phased variant call files

produced by LongRanger to assign the mutations to a maternal or paternal chromosome. In

brief, these methods took input of the three family individuals and a mutation location. The

surrounding haplotype that contained the mutation was directly compared to the parental

haplotypes at the same location to determine a match. As these individuals are all related

members of a colony, dam and sire often shared similar haplotypes. Caution was used when

the mutation-bearing haplotype was found in both parents, resulting in less than 100% parent-

of-origin assignment of mutations.

BPP

Using BPP (Yang 2015) we reran data from a previous study (Yoder, et al. 2016) to measure

the effect of mutation rate estimates on divergence time estimation for the phylogeny of mouse

lemurs. We have written an R package, bppr (available at https://github.com/dosreislab/bppr),

for calibrating BPP estimated phylogenies to geological time using estimates of the mutation

rate. Using bppr, we estimated mouse lemur divergence times twice: (i) using the mutation rate

prior of Yoder et al. (2016), which was based on estimates of mouse and human mutation

rates, and (ii) using our new estimates of the de novo rate. While in the new version of BPP

(v4.0) the theta prior has changed from a gamma distribution to an inverse-gamma distribution,

the resulting tree has the same topology and essentially the same branch lengths as prior

publications (Yoder, et al. 2016).

References

Agier N, Fischer G. 2011. The Mutational Profile of the Yeast Genome Is Shaped by
Replication. Molecular Biology and Evolution 29:905-913.

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool.
Journal of Molecular Biology 215:403-410.

Assaf ZJ, Tilk S, Park J, Siegal ML, Petrov DA. 2017. Deep sequencing of natural and
experimental populations of Drosophila melanogaster reveals biases in the spectrum of new
mutations. Genome research 27:1988-2000.

Axelsson E, Smith NGC, Sundström H, Berlin S, Ellegren H. 2004. Male-Biased Mutation Rate
and Divergence in Autosomal, Z-Linked and W-Linked Introns of Chicken and Turkey.
Molecular Biology and Evolution 21:1538-1547.

Besenbacher S, Hvilsom C, Marques-Bonet T, Mailund T, Schierup MH. 2019. Author
Correction: Direct estimation of mutations in great apes reconciles phylogenetic dating. Nature
Ecology & Evolution 3:859-859.

Bird AP. 1980. DNA methylation and the frequency of CpG in animal DNA. Nucleic Acids
Research 8:1499-1504.

Blanco MB. 2011. Timely Estrus in Wild Brown Mouse Lemur Females at Ranomafana
National Park, Southeastern Madagascar. American Journal of Physical Anthropology
145:311-317.

Blanco MB, Rasoazanabary E, Godfrey LR. 2015. Unpredictable Environments, Opportunistic
Responses: Reproduction and Population Turnover in Two Wild Mouse Lemur Species
(Microcebus Rufus and M-Griseorufus) From Eastern and Western Madagascar. American
Journal of Primatology 77:936-947.

Chaisson MJP, Wilson RK, Eichler EE. 2015. Genetic variation and the de novo assembly of
human genomes. Nature Reviews Genetics 16:627.

Chojnacki S, Cowley A, Lee J, Foix A, Lopez R. 2017. Programmatic access to bioinformatics
tools from EMBL-EBI update: 2017. Nucleic Acids Research 45:W550-W553.

Denver DR, Morris K, Lynch M, Vassilieva LL, Thomas WK. 2000. High Direct Estimate of the
Mutation Rate in the Mitochondrial Genome of Caenorhabditis elegans.
Science 289:2342.

dos Reis M, Gunnell GF, Barba-Montoya J, Wilkins A, Yang Z, Yoder AD. 2018. Using
Phylogenomic Data to Explore the Effects of Relaxed Clocks and Calibration Strategies on
Divergence Time Estimation: Primates as a Test Case. Systematic Biology 67:594-615.

Drummond AJ, Ho SYW, Phillips MJ, Rambaut A. 2006. Relaxed Phylogenetics and Dating
with Confidence. PLOS Biology 4:e88.

Dudchenko O, Batra SS, Omer AD, Nyquist SK, Hoeger M, Durand NC, Shamim MS, Machol
I, Lander ES, Aiden AP, et al. 2017. De novo assembly of the Aedes aegypti genome using Hi-
C yields chromosome-length scaffolds. Science 356:92-95.

Eberle M, Kappeler PM. 2004. Sex in the dark: determinants and consequences of mixed male
mating tactics in Microcebus murinus, a small solitary nocturnal primate. Behavioral Ecology
and Sociobiology 57:77-90.

Ellegren H, Fridolfsson A-K. 1997. Male–driven evolution of DNA sequences in birds. Nature
genetics 17:182-184.

Ewing AD, Houlahan KE, Hu Y, Ellrott K, Caloian C, Yamaguchi TN, Bare JC, P'ng C, Waggott
D, Sabelnykova VY, et al. 2015. Combining tumor genome simulation with crowdsourcing to
benchmark somatic single-nucleotide-variant detection. Nature Methods 12:623.

Feng C, Pettersson M, Lamichhaney S, Rubin CJ, Rafati N, Casini M, Folkvord A, Andersson
L. 2017. Moderate nucleotide diversity in the Atlantic herring is associated with a low mutation
rate. Elife 6.

Friedberg EC, Walker GC, Siede W, Wood RD. 2005. DNA repair and mutagenesis: American
Society for Microbiology Press.

Gao Z, Moorjani P, Sasani TA, Pedersen BS, Quinlan AR, Jorde LB, Amster G, Przeworski M.
2019. Overlooked roles of DNA damage and maternal age in generating human germline
mutations. Proceedings of the National Academy of Sciences 116:9491.

Goetting-Minesky MP, Makova KD. 2006. Mammalian Male Mutation Bias: Impacts of
Generation Time and Regional Variation in Substitution Rates. Journal of molecular evolution
63:537-544.

Goodman M. 1962. Immunochemistry of the Primates and Primate Evolution. Annals of the
New York Academy of Sciences 102:219-234.

Haldane JBS. 1946. The Mutation Rate of the Gene for Haemophilia, and its Segregation
Ratios In Males and Females. Annals of Eugenics 13:262-271.

Harris K, Pritchard JK, Kimura M. 2017. Rapid evolution of the human mutation spectrum
On the evolutionary adjustment of spontaneous mutation rates. Elife 6:e24284.

Heath TA, Huelsenbeck JP, Stadler T. 2014. The fossilized birth–death process for coherent
calibration of divergence-time estimates. Proceedings of the National Academy of Sciences
111:E2957.

Herrera JP, Dávalos LM. 2016. Phylogeny and Divergence Times of Lemurs Inferred with
Recent and Ancient Fossils in the Tree. Systematic Biology 65:772-791.

Hotaling S, Foley ME, Lawrence NM, Bocanegra J, Blanco MB, Rasoloarison R, Kappeler PM,
Barrett MA, Yoder AD, Weisrock DW. 2016. Species discovery and validation in a cryptic
radiation of endangered primates: coalescent-based species delimitation in Madagascar's
mouse lemurs. Molecular Ecology DOI: 10.1111/mec.13604.

Howell N, Smejkal CB, Mackey DA, Chinnery PF, Turnbull DM, Herrnstadt C. 2003. The
Pedigree Rate of Sequence Divergence in the Human Mitochondrial Genome: There Is a
Difference Between Phylogenetic and Pedigree Rates. The American Journal of Human
Genetics 72:659-670.

Jónsson H, Sulem P, Kehr B, Kristmundsdottir S, Zink F, Hjartarson E, Hardarson MT,
Hjorleifsson KE, Eggertsson HP, Gudjonsson SA, et al. 2017. Parental influence on human
germline de novo mutations in 1,548 trios from Iceland. Nature 549:519.

Keightley PD, Pinharanda A, Ness RW, Simpson F, Dasmahapatra KK, Mallet J, Davey JW,
Jiggins CD. 2014. Estimation of the Spontaneous Mutation Rate in Heliconius melpomene.
Molecular Biology and Evolution 32:239-243.

Kim S-H, Elango N, Warden C, Vigoda E, Yi SV. 2006. Heterogeneous Genomic Molecular
Clocks in Primates. PLOS Genetics 2:e163.

Kimura M. 1983. The neutral theory of molecular evolution: Cambridge University Press.

Kimura M, Ohta T. 1971. On the rate of molecular evolution. Journal of molecular evolution
1:1-17.

Kobayashi H, Sakurai T, Miura F, Imai M, Mochiduki K, Yanagisawa E, Sakashita A, Wakai T,
Suzuki Y, Ito T, et al. 2013. High-resolution DNA methylome analysis of primordial germ cells
identifies gender-specific reprogramming in mice. Genome research 23:616-627.

Koboldt DC, Larson DE, Mardis ER, Weinstock GM, Chen K, Ding L, McLellan MD, Wilson RK,
Wylie T. 2009. VarScan: variant detection in massively parallel sequencing of individual and
pooled samples. Bioinformatics (Oxford, England) 25:2283-2285.

Koch EM, Schweizer RM, Schweizer TM, Stahler DR, Smith DW, Wayne RK, Novembre J.
2019. De Novo Mutation Rate Estimation in Wolves of Known Pedigree. Molecular Biology and
Evolution.

Kondrashov FA, Kondrashov AS. 2010. Measurements of spontaneous rates of mutations in
the recent past and the near future. Philosophical Transactions of the Royal Society B:
Biological Sciences 365:1169-1176.

Kong A, Frigge ML, Masson G, Besenbacher S, Sulem P, Magnusson G, Gudjonsson SA,
Sigurdsson A, Jonasdottir A, Jonasdottir A, et al. 2012. Rate of de novo mutations and the
importance of father's age to disease risk. Nature 488:471.

Langley CH, Fitch WM. 1974. An examination of the constancy of the rate of molecular
evolution. Journal of molecular evolution 3:161-177.

Larsen PA, Harris RA, Liu Y, Murali SC, Campbell CR, Brown AD, Sullivan BA, Shelton J,
Brown SJ, Raveendran M, et al. 2017. Hybrid de novo genome assembly and centromere
characterization of the gray mouse lemur (Microcebus murinus). BMC Biology 15:110.

Li W-H, Ellsworth DL, Krushkal J, Chang BHJ, Hewett-Emmett D. 1996. Rates of Nucleotide
Substitution in Primates and Rodents and the Generation–Time Effect Hypothesis. Molecular
Phylogenetics and Evolution 5:182-187.

Lindsay SJ, Rahbari R, Kaplanis J, Keane T, Hurles ME. 2018. Striking differences in patterns
of germline mutation between mice and humans. bioRxiv:082297.

Long H, Winter DJ, Chang AYC, Sung W, Wu SH, Balboa M, Azevedo RBR, Cartwright RA,
Lynch M, Zufall RA. 2016. Low Base-Substitution Mutation Rate in the Germline Genome of
the Ciliate Tetrahymena thermophila. Genome Biology and Evolution 8:3629-3639.

Lynch M. 2010. Evolution of the mutation rate. Trends Genet 26:345-352.

Magallóan SA, Sanderson MJ. 2005. Angiosperm Divergence Times: The Effect of Genes,
Codon Positions, and Time Constraints. Evolution 59:1653-1670.

Martin HC, Batty EM, Hussin J, Westall P, Daish T, Kolomyjec S, Piazza P, Bowden R,
Hawkins M, Grant T, et al. 2018. Insights into Platypus Population Structure and History from
Whole-Genome Sequencing. Molecular Biology and Evolution 35:1238-1252.

Moorjani P, Amorim CEG, Arndt PF, Przeworski M. 2016. Variation in the molecular clock of
primates. Proceedings of the National Academy of Sciences 113:10607.

Narang P, Wilson Sayres MA. 2016. Variable Autosomal and X Divergence Near and Far from
Genes Affects Estimates of Male Mutation Bias in Great Apes. Genome Biology and Evolution
8:3393-3405.

Near TJ, Bolnick DI, Wainwright PC. 2005. Fossil Calibrations and Molecular Divergence Time
Estimates in Centrarchid Fishes (Teleostei: Centrarchidae). Evolution 59:1768-1782.

Ogilvie HA, Bouckaert RR, Drummond AJ. 2017. StarBEAST2 Brings Faster Species Tree
Inference and Accurate Estimates of Substitution Rates. Molecular Biology and Evolution
34:2101-2114.

Ohta T. 1993. Amino acid substitution at the Adh locus of Drosophila is facilitated by small
population size. Proceedings of the National Academy of Sciences 90:4548.

Pfeifer SP. 2017. Direct estimate of the spontaneous germ line mutation rate in African green
monkeys. Evolution 71:2858-2870.

Rahbari R, Wuster A, Lindsay SJ, Hardwick RJ, Alexandrov LB, Al Turki S, Dominiczak A,
Morris A, Porteous D, Smith B, et al. 2015. Timing, rates and spectra of human germline
mutation. Nature genetics 48:126.

Ramu A, Noordam MJ, Schwartz RS, Wuster A, Hurles ME, Cartwright RA, Conrad DF. 2013.
DeNovoGear: de novo indel and point mutation discovery and phasing. Nature Methods
10:985.

Reik W, Dean W. 2001. DNA methylation and mammalian epigenetics. ELECTROPHORESIS
22:2838-2843.

Sanderson MJ. 2002. Estimating Absolute Rates of Molecular Evolution and Divergence
Times: A Penalized Likelihood Approach. Molecular Biology and Evolution 19:101-109.

Scally A, Durbin R. 2012. Revising the human mutation rate: implications for understanding
human evolution. Nature Reviews Genetics 13:745-753.

Schiffels S, Durbin R. 2014. Inferring human population size and separation history from
multiple genome sequences. Nature genetics 46:919-925.

Ségurel L, Wyman MJ, Przeworski M. 2014. Determinants of Mutation Rate Variation in the
Human Germline. Annual Review of Genomics and Human Genetics 15:47-70.

Shimmin LC, Chang BH-J, Li W-H. 1993. Male-driven evolution of DNA sequences. Nature
362:745-747.

Smeds L, Qvarnström A, Ellegren H. 2016. Direct estimate of the rate of germline mutation in a
bird. Genome research 26:1211-1218.

Sung W, Ackerman MS, Miller SF, Doak TG, Lynch M. 2012. Drift-barrier hypothesis and
mutation-rate evolution. Proc Natl Acad Sci U S A 109:18488-18492.

Thomas GWC, Wang RJ, Puri A, Harris RA, Raveendran M, Hughes DST, Murali SC, Williams
LE, Doddapaneni H, Muzny DM, et al. 2018. Reproductive Longevity Predicts Mutation Rates
in Primates. Current Biology 28:3193-3197.e3195.

Thorne JL, Kishino H. 2002. Divergence Time and Evolutionary Rate Estimation with
Multilocus Data. Systematic Biology 51:689-702.

Thorne JL, Kishino H, Painter IS. 1998. Estimating the rate of evolution of the rate of molecular
evolution. Molecular Biology and Evolution 15:1647-1657.

Uchimura A, Higuchi M, Minakuchi Y, Ohno M, Toyoda A, Fujiyama A, Miura I, Wakana S,
Nishino J, Yagi T. 2015. Germline mutation rates and the long-term phenotypic effects of
mutation accumulation in wild-type laboratory mice and mutator mice. Genome research
25:1125-1134.

Van der Auwera GA, Carneiro MO, Hartl C, Poplin R, del Angel G, Levy-Moonshine A, Jordan
T, Shakir K, Roazen D, Thibault J, et al. 2013. From FastQ Data to High-Confidence Variant
Calls: The Genome Analysis Toolkit Best Practices Pipeline. Current Protocols in
Bioinformatics 43:11.10.11-11.10.33.

Venn O, Turner I, Mathieson I, de Groot N, Bontrop R, McVean G. 2014. Strong male bias
drives germline mutation in chimpanzees. Science 344:1272-1275.

Weisenfeld NI, Kumar V, Shah P, Church DM, Jaffe DB. 2017. Direct determination of diploid
genome sequences. Genome research 27:757-767.

Winter DJ, Wu SH, Howell AA, Cartwright RA, Zufall RA, Azevedo RBR. 2018. accuMUlate: a
mutation caller designed for mutation accumulation experiments. Bioinformatics (Oxford,
England) 34:2659-2660.

Wong A. 2014. Covariance between Testes Size and Substitution Rates in Primates.
Molecular Biology and Evolution 31:1432-1436.

Wu CI, Li WH. 1985. Evidence for higher rates of nucleotide substitution in rodents than in
man. Proceedings of the National Academy of Sciences 82:1741.

Xie Z, Wang L, Wang L, Wang Z, Lu Z, Tian D, Yang S, Hurst Laurence D. 2016. Mutation rate
analysis via parent–progeny sequencing of the perennial peach. I. A low rate in woody
perennials and a higher mutagenicity in hybrids. Proceedings of the Royal Society B: Biological
Sciences 283:20161016.

Yang Z. 2006. Computational molecular evolution: Oxford University Press.

Yang ZH. 2015. The BPP program for species tree estimation and species delimitation.
Current Zoology 61:854-865.

Yang ZH, Yoder AD. 2003. Comparison of likelihood and Bayesian methods for estimating
divergence times using multiple gene loci and calibration points, with application to a radiation
of cute-looking mouse lemur species. Systematic Biology 52:705-716.

Yoder AD, Campbell CR, Blanco MB, dos Reis M, Ganzhorn JU, Goodman SM, Hunnicutt KE,
Larsen PA, Kappeler PM, Rasoloarison RM, et al. 2016. Geogenetic patterns in mouse lemurs
(genus Microcebus) reveal the ghosts of Madagascar's forests
past. Proceedings of the National Academy of Sciences 113:8049.

Zhang J. 2004. Evolution of DMY, a Newly Emergent Male Sex-
Determination Gene of Medaka Fish. Genetics 166:1887.

Zohdy S, Gerber BD, Tecot S, Blanco MB, Winchester JM, Wright PC, Jernvall J. 2014. Teeth,
sex, and testosterone: aging in the world's smallest primate. PLoS One 9:e109528.

Zuckerkandl E, Pauling L. 1965. Molecules as documents of evolutionary history. Journal of
Theoretical Biology 8:357-366.
673

674
Figure 1. Focal family quartet with parents (P), offspring (labeled O) and sex (subscript m –
male, or f - female). Lines represent familial relationships as in a traditional pedigree, with
thickness and color reflecting the number and source of *de novo* mutations passed down (red
is from male parent, blue from female parent, gray is undetermined origin). Color of line
represents source and shading represents destination (lighter shading to O_m, darker to O_f).
Numbers within bars show mutation counts and the rate of each individual offspring is listed
below.

683
Figure 2. *de novo* mutation spectrum of the Gray Mouse Lemur, broken down basepair
change (x-axis) and CpG site (color).

Figure 3. The linear relationship between mutation rate and effective population size for
animals (A) and primates (B) with directly estimated mutation rates (for data see Table S3).
Groups and species are identified by color, plots are log-log.

Figure 4. The relationship between mutation rate and substitution rate, (A) per year and (B) per generation, among primates. For the primate species with directly measured mutation rate, plotted against the yearly substitution rate with 95% credible intervals for each measure.

697
698 **Figure 5.** Densitree plot of trees from BPP analysis and bppr plot of previously published
699 mouse lemur RADseq data. Divergence times estimated with either the previous rate estimate
700 (red) or the directly measured rate (blue). Data is also displayed in Table S2, matched by node
701 labels (A-E).
702
703 **Table 1.** Directly estimated mammalian mutation rates.
704

Species	Method	Year	Author	Rate
P. abelii	Pedigree	2019	Besenbacher	1.66×10^{-8}
P. troglodytes	Pedigree	2017	Tatsumoto	1.48×10^{-8}
H. sapiens	Pedigree/Large Scale	2017	Jonsson	1.29×10^{-8}
C. sabaeus	Pedigree	2017	Pfieffer	9.4×10^{-8}
A. nancymaae	Population	2018	Thomas et al.	8.1×10^{-8}
M. musculus	Pedigree/Experimentation	2015	Uchimura	5.4×10^{-9}

706
707 **Figure S1.** Proximity of CG sites to genic regions in four species of mammals (Hsap – *Homo*
708 *sapiens*, Mmur – *Microcebus murinus*, Mmus – *Mus musculus*, Ptro – *Pan troglodytes*). No
709 differences were observed in proximity of CG sites to genic regions.

Figure S2. MSMC plot of gray mouse lemur samples in this study, used to estimate the effective population size of gray mouse lemurs. The harmonic mean of the estimated population size of all samples was taken across the last 2 million years, which yielded an N_e of 41,000.

Table S1. Mutation (various pubs) and substitution rates (dos Reis et al. 2018) calculated per year and per generation.

Species	Per Year		Per Generation		Gen Time
	Mutation	Substitution	Mutation	Substitution	
Chlorocebus sabaeus	1.1E-09	8.5E-10	9.4E-09	7.2E-09	8.5
Pan troglodytes	5.2E-10	7.1E-10	1.3E-08	1.8E-08	24.6
Homo sapiens	4.1E-10	7.5E-10	1.2E-08	2.2E-08	29
Gorilla gorilla	5.8E-10	8.5E-10	1.1E-08	1.6E-08	19.3
Pongo abelii	8.3E-10	3.6E-10	1.7E-08	7.1E-09	20
Aotus nancymaae	2.7E-09	5.4E-10	8.1E-09	1.6E-09	3
Microcebus murinus	4.4E-09	1.7E-09	1.6E-08	6.5E-09	3.75

724 **Table S2.** Ancestral effective population size (N$_e$) and estimated divergence times as
725 determined by BPP and bppr (see Materials and Methods). Table includes results for the
726 previous estimate (Yoder, et al. 2016) and the directly measure rate, with node labels
727 corresponding to Figure 5.
728

Directly Estimated Rate				
Node	**N$_e$ (×10^3)**	**95% CI**	**Divergence Time (ka)**	**95% CI**
E	155	129, 182	275	198, 357
D	10.8	5.29, 14.5	157	117, 201
A	39.8	25.9, 53.1	26.4	4.79, 54.8
C	2.01	1.19, 2.67	146	107, 187
B	44.5	30.8, 57.7	8.84	0.30, 26.0
Previous Rate Estimate				
Node	**N$_e$ (×10^3)**	**95% CI**	**Divergence Time (ka)**	**95% CI**
E	307	200, 427	545	321, 796
D	21.5	8.56, 34.5	313	188, 452
A	79.0	42.3, 118	52.4	7.42, 112
C	3.98	1.92, 6.14	290	174, 421
B	88.4	51.5, 130	17.5	0.51, 52.1

739 **Table S3.** Mutation rates (various pubs) and effective population sizes plotted in Figure 3.
740

Directly Estimated Animal Mutation Rates					
Species	**Mutation Rate**	**Ne**	**Author**	**Year**	**Category**
Human	1.20E-08	1.00E+04	Many	2017	Primate
Chimp	1.27E-08	1.10E+04	Besenbacher	2019	Primate
Orangutan	1.66E-08	2.68E+04	Besenbacher	2019	Primate
Green Monkey	9.40E-09	1.20E+04	Pfeifer	2017	Primate
Owl Monkey	8.10E-09	0.00E+00	Thomas	2018	Primate
Mouse Lemur	1.64E-08	4.10E+04	This Study	2019	Primate
Mouse	5.40E-09	7.50E+04	Uchimara	2015	Rodent
Platypus	7.00E-09	2.00E+04	Martin	2018	Monotreme
Cow	9.70E-09	3.70E+04	Harland	2017	Bovid
Flycatcher	4.60E-09	3.00E+05	Smeds	2016	Bird
Herring	2.00E-09	4.00E+05	Feng	2017	Fish
Bee	6.80E-09	4.73E+05	Yang	2015	Insect
Butterfly	2.90E-09	2.00E+06	Keightley	2015	Insect
Fly	2.80E-09	1.40E+06	Keightley	2014	Insect

The uncertainty of the mutation rate estimate from pedigree-sequencing studies

Let λ_m and λ_p be the maternal and paternal mutation rates per genome. The total mutation rate is $\lambda = \lambda_m + \lambda_p$. For example, if we expect an offspring to inherit 30 and 40 mutations form their mother and father respectively, then $\lambda_m = 30$, $\lambda_p = 40$, and $\lambda = 70$. If we assume that the mutations inhereted form the mother and father have a Poisson distribution, then the total number of mutations inherited also follows a Poisson distribution.

Let x_i be the total number of mutations observed in an individual, and suppose we observe mutations in n individuals. We can use the Bayesian method to estimate the mutation rate, l . Note that the gamma distribution is the conjugate prior of the Poisson, and thus we use a gamma prior on l with parameters α (shape) and β (rate). The posterior distribution is

$$f_\lambda(\lambda \mid x_1,\ldots,x_n) \propto \lambda^{\alpha-1} e^{-\beta\lambda} \times \prod_{i=1}^{n} \frac{\lambda^{x_i} e^{-\lambda}}{x_i!},$$

$$\propto \lambda^{\alpha+\Sigma x_i - 1} e^{-(\beta+n)\lambda}.$$

Thus, the posterior of λ is a gamma distribution with parameters $\alpha + \Sigma x_i$ (shape) and $\beta + n$ (rate).

In the mouse lemur pedigree-sequencing results, we see 71 and 63 mutations in two individuals respectively. Let the prior on λ be gamma with $\alpha = 2$ and $\beta = 0.02$. This is a diffuse prior with mean $\alpha/\beta = 100$ mutations. Thus the estimate of λ is gamma distributed with shape $2 + 71 + 63 = 136$ and rate $0.02 + 2 = 2.02$. Thus the posterior mean is $\tilde{\lambda} = 67.3$, posterior standard deviation $\tilde{s}$ $\tilde{s} = \sqrt{136}/2.02 = 5.77$. Note this estimate of λ is given as mutations per genome. To get the rate per nucleotide site, we simply divide by the diploid genome size, g. For example, assuming this is $g = 4\times10^9$ in mouse lemurs, we get $\tilde{\lambda}$ $=1.68\times10^{-8}$ and $\tilde{s}=1.44\times10^{-9}$. The 95% credibility interval for the estimate between 56.5 and 79.1 mutations per genome, or between 1.41×10^{-8} and 1.98×10^{-8} mutations per nucleotide site.

Supplemental Text 2

Variant Calling's Effect on *de novo* Mutation Rate Estimation

An attempt to, given a duplicate sample providing a known variant call error rate, quantify the effects of incorrect variant calls on the estimation of de novo *mutations.*

Our knowns are as follows, the number of sites in the genome with sufficient coverage to make *de novo* calls (sites, s), the number of variants (variants, v) in the entire pedigree (v_p), the mother's sample (v_m), the father's sample (v_f), and the offspring sample (v_o). Also the number of unique variants (unique variants, w) in the entire pedigree (w_p), the mother's sample (w_m), the father's sample (w_f), and the offspring sample (w_o). We also know the error rate of the replicated sample, presented here as a fraction of variants called (called error rate, c). This is an average rate of variants that are present in one replicate and absent in the other. We also assume the transmission probability of any given site is 0.5 (transmission probability, t).

Finally we'll add a single unknown variable (e), which is the fraction of erroneous variant calls that are false positives. This variable ranges from 0 to 1, with 1 representing the scenario where all erroneous calls are false positives. We have no reason to expect that this fraction differs across the replicates. This variable will allow us to explore a range of possible error effects. It will also allow us to be conservative (taking the value of e that yields the most error) when reaching a final conclusion regarding error rates.

The probability that a *de novo* mutation call is a false positive is simply the probability of a false positive variant within the offspring that is not already a variant in the pedigree or a parental false negative at a site that is unique to the individual and transmitted.

$$P(\mu_{f+}) = P(unique\ offspring\ false\ positive) + P(shared\ paternal\ false\ negative) + P(shared\ maternal\ false\ negative)$$

$$P(unique\ offspring\ false\ positive) = c * e * \frac{w_o}{v_o}$$

$$P(shared\ paternal\ false\ negative) = \frac{c * v_p}{s} * (1 - e) * \frac{w_o}{v_o}$$

$$P(shared\ maternal\ false\ negative) = \frac{c * v_m}{s} * (1 - e) * \frac{w_o}{v_o}$$

So, substituting the these into the original equation:

$$P(\mu_{f+}) = c * e * \frac{w_o}{v_o} + \frac{c * v_m}{s} * (1 - e) * \frac{w_o}{v_o} + \frac{c * v_f}{s} * (1 - e) * \frac{w_o}{v_o}$$

$$P(\mu_{f+}) = \frac{c * w_o}{v_o} * (e + \frac{v_p * v_m * (1 - e)}{s})$$

The probability that a genomic site with sufficient coverage is an uncalled false negative *de novo* mutation call is essentially the converse of the false positive. It is the probability of a false negative within the offspring that is not already a variant in the pedigree or a parental false positive at a site that is unique to the individual and transmitted.

$$P(\mu_{f-}) = P(unique\ offspring\ false\ negative) + 2 * P(shared\ parental\ false\ positive)$$

$$P(unique\ offspring\ false\ negative) = (1 - \frac{w_o}{s}) * \frac{c}{s} * (1 - e)$$

$$P(shared\ parental\ false\ positive) = c * e * \frac{w_o}{s}$$

So, substituting the these into the original equation:

$$P(\mu_{f-}) = (1 - \frac{w_o}{s}) * \frac{c}{s} * (1 - e) + c * e * \frac{w_o}{s} + c * e * \frac{w_o}{s}$$

$$P(\mu_{f-}) = \frac{c}{s} * ((1 - \frac{w_o}{s}) * (1 - e) + 2 * w_o * e)$$

$$P(\mu_{f-}) = c * v_o * ((1 - e) * (s - v_p) + 2 * e * w_o)$$

Using as known values the following rough estimates:

$$c = 0.025$$
$$e = 0.5$$
$$s = 1.968 * 10^9$$
$$\mu_{dn} = 134$$
$$v_{all} = 30,000$$
$$w_{all} = 1,500$$

The probabilites of false negative and false positives are as follows:

$$P(\mu_{f+}) = 4.83 * 10^{-2}$$

$$P(\mu_{f-}) = 1.91 * 10^{-8}$$

Given 134 *de novo* mutations from 1,968,000,000 sites of sufficient coverage, we expect:

$$E(\mu_{f+}) = \mu_{dn} * P(\mu_{f+})$$
$$E(\mu_{f+}) = 6.46$$

$$E(\mu_{f-}) = s * P(\mu_{f-})$$
$$E(\mu_{f-}) = 37.51$$