Pedigree-based measurement of the *de novo* mutation rate in the gray mouse lemur

reveals a high mutation rate, few mutations in CpG sites, and a weak sex bias

*C. Ryan Campbell[1,2], *George P. Tiley[1], Jelmer W. Poelstra[1], Kelsie E. Hunnicutt[1,†], Peter A.

Larsen[1,‡], Hui-Jie Lee[3], Jeffrey L. Thorne[4], Mario dos Reis[5], Anne D. Yoder[1]

[1]Department of Biology, Duke University, Durham, NC 27708, USA

[2]Department of Evolutionary Anthropology, Duke University, Durham, NC 27708, USA

[3]Department of Biostatistics and Bioinformatics, Duke University, Durham, NC 27708, USA

[4]Bioinformatics Research Center, North Carolina State University, Raleigh, NC 27695, USA

[5]School of Biological and Chemical Sciences, Queen Mary University of London, Mile End

Road, London E1 4NS, UK

[†]Present address: Department of Biological Sciences, University of Denver, Denver, CO 80208

[‡]Present address: Department of Veterinary and Biomedical Sciences, University of

Minnesota, St. Paul, MN 55108

*These authors contributed equally

Author for Correspondence: anne.yoder@duke.edu

Word count for Main Text: 6690

Abstract

Spontaneous germline mutations are the raw material on which evolution acts, and knowledge

of their frequency and genomic distribution is crucial for understanding how evolution operates

at both long and short timescales. At present, the rate and spectrum of *de novo* mutations

have been directly characterized in only a few lineages. It is therefore critical to expand the

phylogenetic scope of these studies to gain a more general understanding of observed

mutation rate patterns. Our study provides the first direct mutation rate estimate for a

strepsirrhine (i.e., the lemurs and lorises), which comprise nearly half of the primate clade.

Using high-coverage linked-read sequencing for a focal quartet of gray mouse lemurs

(*Microcebus murinus*), we estimated the mutation rate to be 1.64×10^{-8} (95% credible interval:

1.41×10^{-8} to 1.98×10^{-8}) mutations/site/generation. This estimate is higher than those

measured for most previously characterized mammals. Further, we found an unexpectedly low

count of paternal mutations, and only a modest overrepresentation of mutations at CpG-sites.

Given the surprising nature of these observations, we conducted an independent analysis of

context-dependent substitution types for gray mouse lemur and five additional primate species.

This analysis yielded patterns consistent with the mutation spectrum from the pedigree

mutation-rate analysis, which provides confidence in our ability to accurately identify *de novo*

mutations with our data and bioinformatic filters.

Introduction

Spontaneous germline mutations are errors that occur as DNA is transmitted from parent to offspring in sexually reproducing organisms. The accrual of these errors, often referred to as *de novo* mutations, provides not only the raw material for evolution but can also serve as a measure of the evolutionary time along phylogenies (Kimura and Ohta, 1971; Langley and Fitch, 1974; Zuckerkandl and Pauling, 1965). The rate at which these mutations are introduced into genomes is thus a crucial metric of evolution at the genomic level as well as a measure of fundamental biological processes (Kondrashov and Kondrashov, 2010). And by characterizing mutation rate variation across the genome and between generations, we may be able to disentangle distinct biological processes such as potential sex and parental age biases. Ultimately, by quantifying the variation in *de novo* mutation rates across the tree of life, we can refine hypotheses regarding the relationship between mutation rates and life history characteristics.

Approaches for estimating rates of genomic change in vertebrates generally fall into one of two categories: phylogenetic (indirect) versus pedigree-based (direct) estimation. While phylogenetic methods were the standard before recent developments in sequencing technology that have made whole-genome sequencing widely accessible, pedigree-based mutation rates are now increasingly being estimated for non-model species. By comparing the genomes of individuals with known genealogical relationships – typically, parent to offspring – investigators can count mutations as they appear (Feng *et al*, 2017; Koch *et al*, 2019; Pfeifer, 2017; Scally and Durbin, 2012; Smeds *et al*, 2016; Thomas *et al*, 2018). Phylogenetic approaches, on the other hand, use external calibrations such as fossils or geological events

to obtain substitution rates in units of absolute time (Drummond *et al*, 2006; Sanderson, 2002;

Thorne and Kishino, 2002; Thorne *et al*, 1998). Phylogenetic studies work from the assumption

that the rate at which substitutions accumulate between species at putatively neutral sites is

equal to the *de novo* mutation rate (Kimura, 1983). If this assumption holds, pedigree-based

and phylogenetic methods should in principle produce equivalent estimates of genomic

evolution.

Phylogenetic methods for estimating rates of evolution are known to suffer from various

sources of uncertainty, however, including violation of the molecular clock, inaccuracies in

external calibration points, incomplete lineage sorting, and the difficulties of recovering multiple

overlapping changes (i.e., "multiple hits") at any given site (dos Reis *et al*, 2018). Although a

number of solutions to these problems have been proposed (Heath *et al*, 2014; Ogilvie *et al*,

2017), some limitations such as sampling biases or an absence of fossils are difficult to

overcome (Herrera and Davalos, 2016; Magallon and Sanderson, 2005; Near *et al*, 2005).

Pedigree-based mutation rate estimates are not affected by the same uncertainties and can

help characterize variation among different types of mutations (Harris and Pritchard, 2017) or

among different regions of the genome (Segurel *et al*, 2014). Previously, these estimates have

relied on well-assembled genomes available only in model organisms (Jonsson *et al*, 2017;

Scally and Durbin, 2012; Uchimura *et al*, 2015; Venn *et al*, 2014), and have therefore been

limited in taxonomic scope. For example, mutation rate estimates within mammals are

represented mostly by primates (Table 1). Fortunately, recent genome assembly strategies

(Dudchenko *et al*, 2017) have enabled chromosome-level assemblies of non-model organisms

(Larsen *et al*, 2017) and pedigree-based mutation rate estimation is now feasible for virtually

any species, as long as related individuals with known pedigrees are available (Feng *et al*,

2017; Koch *et al*, 2019; Martin *et al*, 2018; Pfeifer, 2017; Smeds *et al*, 2016).

These advantages notwithstanding, pedigree-based studies also face substantial challenges.

Perhaps foremost among them is the fact that mutation rates are orders of magnitude lower

than the sequencing error rate, even for the most accurate sequencing methods. Therefore,

the number of *de novo* mutations produced in a single generation can be difficult to

differentiate from erroneous variant calls – a challenge that is typically addressed with

computational methods. As a consequence, stringency in bioinformatic filters can be so

vigorously applied that many true mutations are not called (i.e., false negatives are common),

and the mutation rate can be under- rather than overestimated.

In this study, we utilize two strategies for keeping both false negative and false positive rates

low. First, linked short reads from 10x Genomics (Weisenfeld *et al*, 2017) provide improved

mapping and increased accuracy of individual variant calls (Long *et al*, 2016; Winter *et al*,

2018), especially in repeat-rich mammalian genomes (Chaisson et al, 2015). Additionally, the

phasing information provided by linked reads can determine the parent-of-origin with just two

generations of sequencing. Phased haplotypes with known parental origin then allow individual

mutations to be assigned to either the maternal or paternal germline. To control for false

negatives, we explicitly estimate the callable proportion of the genome by introducing synthetic

mutations to the sequencing data for one individual and subsequently testing the accuracy of

our bioinformatic pipeline in recovering these mutations (Keightley *et al*, 2015; Xie *et al*, 2016).

We refer to this approach as "allele drop", which can be used to correct observed mutation

rates regardless of the variant calling pipeline.

We applied these advances in sequencing technology and computational methods to produce

the first pedigree-based mutation rate estimate for a strepsirrhine primate, the gray mouse

lemur (*Microcebus murinus*). Mouse lemurs comprise a radiation of morphologically cryptic

primates distributed throughout Madagascar (Hotaling *et al*, 2016). Numerous studies have

suggested that their rapid speciation dynamics may reflect climatic change through time in

Madagascar (Andriatsitohaina *et al*, 2020; Setash *et al*, 2017) and that their unique life history

characteristics make them an ideal genetic model organism (Ezran *et al*, 2017; Hozer *et al*,

2019). Thus, an accurate mutation rate estimate for these organisms can potentially yield

valuable insight into both geological and biological phenomena. Even though previous

divergence time studies exist, they have had to rely on either phylogenetic methods wherein

only distantly-related external fossil calibrations are available (dos Reis *et al*, 2018; Yang and

Yoder, 2003) or on pedigree-based mutation rate estimates from distant relatives (Yoder *et al*,

2016). Notably, the two approaches have yielded highly divergent age estimates.

By measuring the mutation rate in mouse lemurs with a pedigree-based approach, we aim to

simultaneously expand our knowledge of mutation rate variation across lineages and to

facilitate the estimation of divergence times within the mouse lemur radiation specifically. To

do so, we deeply sequenced eight individuals from a family pedigree of gray mouse lemurs,

including a mother, father, and two offspring, to accurately identify *de novo* mutations and to

assign their parent-of-origin. We found a relatively high mutation rate, an unexpectedly low

rate of transitions at CpG sites, and a weak paternal sex bias compared to other primates.

Given the surprising nature of these results, we take care to discuss the potential

methodological caveats that may possibly yield misleading results in pedigree-based studies,

including this one. Importantly, we validate the estimated patterns observed in the *de novo*

mutation rate spectrum with phylogenetically derived substitution rate patterns, finding that the

latter supports the former. We therefore conclude that though unexpected, the results of our

pedigree analysis offer a true representation of the *de novo* mutation rate in mouse lemurs.

Materials and Methods

Samples

Eight individuals were selected from the Duke Lemur Center's (DLC) mouse lemur colony

consisting of a focal family of two parents and two offspring from separate litters, an additional

half-sibling to the offspring, and three other individuals in the maternal lineage (Fig S1). Four of

the eight selected samples were colony founders, which had been transferred in 2003 from the

CNRS mouse lemur colony in Brunois, Paris, France. Blood and tissue samples were collected

from all individuals during annual veterinary check-ups. High molecular weight DNA was

extracted with the Qiagen MagAttract kit (Qiagen, Germantown, MD, USA) and 10X Genomics

library preparation was performed at the Duke Molecular Genomics Core.

Sequencing

Nine sequencing libraries were produced from the eight individuals; every individual was

sequenced once except the focal paternal sample that was prepared twice and sequenced as

two separate libraries to serve as a biological replicate. Libraries were sequenced at the Duke

Center for Genomic Computational Biology (GCB) Sequencing and Genomic Technology

Shared Resource across nine lanes of a HiSeq 4000. Paired-end sequencing of 150 basepair

reads was performed with an average insert size of 554bp (range: 527-574bp). A single lane

was run as a test of the 10x Genomics LongRanger analysis software and was analyzed to

confirm successful indexing and preparation of the samples. Next, the remaining eight libraries

were multiplexed across eight lanes of a single flowcell. 933 337 210 328 bases were

generated across nine libraries and nine lanes. Sequencing data are available through NCBI's

SRA database (SRR10130788-SRR10130796).

10x Genomics Pipeline

Basecall files were demultiplexed and analyzed using 10x Genomic's LongRanger v2.2.1

pipeline. Average genomic coverage after filtering was 34.5x across the nine samples.

Sequences were aligned to the reference gray mouse lemur genome assembly (mmur3.0,

GCF_000165445.3) and variant calling was performed using GATK v3.8 (McKenna *et al*, 2010;

Van der Auwera *et al*, 2013), implemented within LongRanger v2.2.1 (Weisenfeld *et al*, 2017).

The mean N50 scaffold length, across samples, generated by the 10x Genomics LongRanger

alignment pipeline was 1.18Mb.

DeNovoGear

LongRanger alignments were used to find *de novo* mutations within the offspring in the focal

family. Several methods were used to find mutations. First, DeNovoGear v1.1.1 (Ramu *et al*,

2013) was used to analyze the LongRanger variant call files with default settings. VarScan2

v2.4.3 was run with the LongRanger binary alignment files and the resulting variants were

intersected with the *de novo* mutations found with DeNovoGear. Only mutations found by both

approaches were retained.

De novo mutations were inferred separately with each replicate library from the sire, and

mutations that differed by sire replicate were used to estimate false positive rates (See

Supplementary Material: *Error Rates from Technical Replicates*). Finally, we checked whether

alleles produced by the inferred *de novo* mutations were absent in the remainder of the

pedigree and in existing data from a sequenced diversity panel of gray mouse lemurs (NCBI

SRA:SRP045300). The final list of mutations was filtered for *de novo* quality in the offspring

(*DNQ* of at least 100), offspring mapping quality (*MQ* of at least 50), and for at least 20x depth

of coverage in both parents. The total number of mutations in each offspring was used to

estimate a credible interval for our per-generation mutation rate (See Supplementary Methods:

Mutation Rate Credible Intervals).

Detectable Mutation Test

We conducted a test to determine the number of sites at which a mutation was detectable

given our sequencing data and methods, hereafter referred to as an "allele-drop test". This test

consisted of adding 1,000 synthetic mutations into the pedigree with the software BAMsurgeon

v1.0.0 (Ewing *et al*, 2015). These mutations were added as heterozygotes by changing half of

the aligned bases in the bam file at a given site to the non-reference allele. Next, we again

applied our pipeline to find *de novo* mutations and examined the results for the 1,000 synthetic

sites. By conducting this allele-drop test, we were able to estimate the fraction of the genome

for which *de novo* mutations should have been found. It is this proportion of the genome that

we refer to as "callable sites" and that is used as the denominator for calculating the genome-

wide mutation rate as opposed to quality metric-based callable site estimates (Krasovec *et al*,

2019). We prefer the allele-drop test to estimate callable sites because it jointly considers the

data and uncertainty from our bioinformatic pipelines rather than read quality and depth alone.

Parent-of-Origin

Phased variant call files produced by LongRanger were used to assign the mutations to a

maternal or paternal chromosome. In brief, these methods took input of the three family

individuals and a mutation location. The surrounding haplotype that contained the mutation

was directly compared to the parental haplotypes at the same location to determine a match.

As these individuals are all genetically related members from a single colony, dam and sire

often shared similar haplotypes. When the mutation-bearing haplotype was found in both

parents, a parent-of-origin was not assigned, resulting in less than 100% parent-of-origin

assignment of mutations.

CpG Islands and CpG Mutation Rates

CpG islands were identified by two independent methods and compared to measure the

number of mutations within them. First, the EMBOSS cpgplot tool (Chojnacki *et al*, 2017) was

run with the latest gray mouse lemur genome (mmur3.0, GCF_000165445.3) to identify

regions that met the threshold of a CpG island (200 bp, over 50% CG content). Then, to

confirm these annotations, a fasta file of CpG island annotations from the gray mouse lemur

genome 2.0 (GCF_000165445.2) was downloaded from the UCSC genome browser. A blast

(Altschul, et al. 1990) database of the mmur3.0 genome was created and the mmur2.0 CpG

islands were queried to determine their coordinates in the genome used for mapping and

assembly. Only the CpG islands identified with both methods (a total of 67,673 annotations)

were used to determine whether a mutation at a CpG site were contained in a CpG island.

Context-Dependent Substitution Rate Estimation

Because the mutation spectrum we observed in mouse lemur differed from that observed in

other primates and was complicated by the challenges of robust mutation rate estimation from

a single pedigree, we performed additional analyses to estimate substitution rates across the

primate phylogeny. To do so, we used molecular clock methods that allow rates to differ by

substitution type, including C>T transitions at non-CpG and CpG sites. First, we downloaded

high-coverage mammalian whole-genome alignments from Ensembl

(ftp://ftp.ensembl.org/pub/current_emf/ensemblcompara/multiple_alignments/46_mammals.ep

o/; last accessed February 2020). Analyses only used alignments that included these seven

taxa: *Mus musculus*, *Microcebus murinus*, *Callithrix jacchus*, *Chlorocebus sabaeus*, *Pongo*

abelii, *Pan troglodytes*, *Homo sapiens*. The *M. murinus* reference genome used in the whole-

genome alignment was the same version used for calling mutations (Larsen *et al*, 2017). Sites

that mapped to protein-coding genes and CpG islands based on human gene features were

removed. Data processing was done with Perl scripts available through Dryad. We randomly

sampled ten one-megabase lengths of concatenated alignment to keep analyses

computationally tractable.

We first estimated context-independent substitution rates. Branch lengths were optimized by

Maximum Likelihood with the baseml program in PAML v4.8j (Yang, 2007) using the HKY +

gamma model. The approximate likelihood method (dos Reis and Yang, 2011) was used to

estimate absolute rates of evolution with fossil calibrations on all nodes (Table S1) that follow

"calibration strategy A" from dos Reis et al. (2018). For each sub-sample, we ran four MCMC

chains that discarded the first 50 million generations as burn-in and kept 10000 posterior

samples for every 50000 generations. Input alignments, control files, and the species tree are

available through Dryad. Posteriors were analyzed in R v3.6.3 with the package CODA

(Plummer *et al*, 2006).

The same sub-sampled alignments were used to estimate substitution rates for nine context-

dependent substitution types following the method in (Lee *et al*, 2015). This method

characterizes dinucleotide sites by integrating over uncertainty in substitution history for each

site based on a sample of stochastic character maps. Substitution histories for each site were

generated with PhyloBayes MPI v1.8 (Lartillot *et al*, 2013) under the CAT-GTR model (Lartillot

and Philippe, 2004). 1100 samples were collected for two chains for each sub-sampled

alignment while sampling every five generations. The first 100 samples were discarded as

burn-in. A total of 15 stochastic mappings were collected for each site. These were used to

compute the variance-covariance matrices for the nine substitution types and approximate the

likelihood surface of Bayesian relaxed-clock model. MULTIDIVTIME (Thorne *et al*, 1998) was

then used to estimate absolute rates of evolution for each substitution type under an

autocorrelated model (Thorne and Kishino, 2002) with calibrations in Table S1.

MULTIDIVTIME analyses collected 10,000 posterior samples for two chains, sampling every

10,000 generations after a 10 million generation burn-in. Rate posteriors were evaluated for

convergence and combined.

Divergence Time Estimation

Using BPP v4.0 (Yang, 2015), we re-evaluated divergence time estimates from a previous

study (Yoder *et al*, 2016) using the pedigree-based mutation rate recovered by this study. We

have written an R package, bppr (available at https://github.com/dosreislab/bppr), for

calibrating node heights estimated by BPP to geological time using estimates of the mutation

rate. Using bppr, we estimated mouse lemur divergence times twice: 1) using the mutation rate

prior of Yoder et al. (2016), which was based on estimates of mouse and human mutation

rates, and 2) using our new estimates of the *de novo* rate.

Results

Estimating the Gray Mouse Lemur Mutation Rate

We assessed 4,542,770 potential variants across eight related individuals to discover 134 *de*

novo mutations in two focal offspring, one male and one female (Fig 1). Among these 134

mutations, 125 (71 in the female and 63 in the male) were located on autosomes and nine

(seven in the female and two in the male) were located on the X chromosome. The average

depth of coverage in the quartet for the 134 mutations was 170 reads (170.2, SD=79.95). We

estimate the mutation rate in this family quartet to be 1.64×10^{-8} (95% credible interval: $1.41 \times$

10^{-8} to 1.98×10^{-8}) single nucleotide mutations/site/generation. The single-base mutation-rate

estimate is a weighted average of the number of mutations on the autosomes (125 mutations)

and X chromosome (nine mutations) divided by the number of callable sites.

To estimate the number of callable sites, we artificially generated 1,000 mutations placed

randomly across the entire genome within the sequence data of the offspring, similar to

previous efforts to account for false negative results (Keightley *et al*, 2015; Xie *et al*, 2016). On

the autosomes, we detected 801 of 952 generated mutations, while 39 of 48 generated

mutations were detected on the X chromosome. Therefore, we estimate our detection rate to

be 84.1% on autosomes and 81.3% on the X chromosome, which yields a total of 2.088 billion

callable sites (out of a total of 2.487 billion).

To calculate the credible interval around the estimated mutation rate, we assumed that the

total number of mutations inherited by an offspring follows a Poisson distribution (See

Supplementary Methods: *Mutation Rate Credible Intervals*). Given that an average of 67

mutations were found in each of the focal offspring, the 95% credible interval is 56.5 to 79.1

mutations per genome, or 1.41×10^{-8} to 1.98×10^{-8} mutations/site/generation. To calculate

the expected number of false positives and false negatives, we sequenced the male offspring

twice and compared variant presence and absence across the technical replicates (See

Supplementary Methods: *Error Rates from Technical Replicates*). We calculated 6.46 false

positives and 37.51 false negatives from the total of 134 *de novo* mutations and 2.088 billion

callable mutation sites.

The Mouse Lemur Mutation Spectrum

The ratio of transitions to transversions (Ti:Tv) was estimated to be 1.03 (68 transitions and 66

transversions). The ratio of strong-to-weak mutations (SW; C/G > A/T) to weak-to-strong

mutations (WS; A/T > C/G), SW:WS, was also estimated to be 1.03 (31 SW and 30 WS

mutations). The most common two categories of *de novo* mutation type were A>G and C>T

(Fig 2a). Eleven mutations were detected at parental CpG sites, constituting 8.2% of all *de*

novo mutations. This represents a roughly four-fold enrichment given that 1.9% of the genome

consists of CpG sites. Because the elevated mutation rate at CpG sites is linked to methylation

(Bird, 1980), mutations are typically not expected in regions of the genome with high GC

content (CpG islands), where CpG sites are much less likely be methylated (Bird, 1986; Molaro

et al, 2011). As anticipated, none of the 134 *de novo* mutations were found within CpG islands,

which constitute roughly 4% of the *M. murinus* genome.

The mutation spectrum in mouse lemur was further investigated with an independent approach

based on absolute substitution rates (substitutions/site/year; s/s/y) and fossil-calibrated

relaxed-clock models. All clock model parameters (Supplementary Figure S2) converged

across ten one-megabase replicates (Supplementary Figures S3-S13) and revealed a higher

global substitution rate in mouse lemurs compared to apes and Old World monkeys

(Supplementary Figure S14). We then estimated context-dependent substitution rates for the

same alignments (Lee *et al*, 2015). All rate parameters converged (Supplementary Figures

S15-S24) and transitions at CpG sites (Group 9) were the only substitution type to clearly

break from the pattern expected by not partitioning across substitution types (Supplementary

Figure S14). Mouse lemur had the lowest rate of C>T transitions at CpG sites of all primates

(Supplementary Figures S25-S34). Notably, in mouse lemur, the rate of C>T transitions at

CpG sites is slightly lower than the rate of C>T transitions at non-CpG sites (Group 5),

whereas the converse is true for all other primates across all ten sub-sampled alignments

(Supplementary Figures S25-S34). Specifically, the mean rate estimate for C>T transitions at

CpG sites is 99.7% of the rate of C>T transitions at non-CpG sites (1.233×10^{-11} s/s/y vs

1.236×10^{-11} s/s/y) in mouse lemur. The C>T transition rate is 2.94, 3.1, 2.49, 1.79, and 1.75

times higher for CpG versus non-CpG sites in human, chimp, orangutan, Old World monkey,

and New World monkey respectively. The pattern of rate variation across substitution types

generally agrees with the observed mutation spectrum from our focal quartet (Fig 2b) and

corroborates the low rate of CpG mutations in the gray mouse lemur relative to other primates

(Fig 3)

Discrepancies of Magnitude when Comparing Pedigree-Based Mutation Rates and

Phylogenetic Substitution Rates

We compared pedigree-based estimates of the mutation rate for mouse lemurs together with

published mutation rate estimates from other primates (Table 1) with substitution rates

estimated from a recent relaxed-clock analysis of the same species (dos Reis *et al*, 2018).

Phylogenetic substitution rates are estimated per-year, so we rescaled them by generation

time (Table S3) for direct comparison with per-generation mutation rates from pedigrees.

(Table S1, Fig 4. There are two notable observations: 1) Pedigree-based point estimates fall

outside of the highest posterior density intervals of phylogenetic estimates in four out of seven

cases and 2) substitution rates are not consistently lower than mutation rates across species.

Whereas pedigree-based estimates are higher than phylogenetic estimates for *Microcebus*

murinus, *Aotus nancymaae*, *Chlorocebus sabaeus*, and *Pongo abelii,* the opposite is true for

Homo sapiens, Pan troglodytes, and *Gorilla gorilla* (Fig 4).

Impacts for Divergence Time Estimation

We recalculated branch lengths in absolute time for a genus-level phylogeny of mouse lemurs

(Yoder *et al*, 2016) based on the new mutation rate estimate of 1.64×10^{-8}

mutations/site/generation derived from this study. Previously, the mutation rate was modelled

on a gamma distribution from mouse (Uchimura *et al*, 2015) and human (Scally and Durbin,

2012) estimates, with a mean of 0.87×10^{-8} mutations/site/generation. The higher mutation

rate calculated here yields considerably more recent divergence times (Fig 5) with reduced

uncertainty compared to the previously wide gamma distribution (Table S4).

Sex Bias

Using the long phasing blocks generated by the linked-read method, we were able to

determine the parent-of-origin for 94 out of 134 (70%) *de novo* mutations. The number of

mutations confidently assigned to a parent are notably higher in our analysis compared to

previous studies that used short read sequencing alone, such as 35% (Venn *et al*, 2014) or

38% (Thomas *et al*, 2018). Among the assigned mutations, 54% (n = 51) were found on the

offsprings' paternal haplotype while the remaining 46% (n = 43) were found on the offsprings'

maternal haplotype; a ratio of male-to-female mutations of approximately 1.2.

Relationship of Mutation Rate Estimates to Life History Traits

The drift barrier hypothesis states that effective population size (N_e) may explain some

variation in mutation rates across species due to a larger efficiency of selection acting on DNA

replication fidelity in larger populations, especially across large phylogenetic distances (Lynch,

2010; Sung *et al*, 2012). Using an MSMC analysis (Schiffels and Durbin, 2014), we estimated

the N_e of gray mouse lemur across the last 2 million years to be approximately 41,000 (Fig

S35). When analyzed in a broad phylogenetic context, we find that the mouse lemur lies

comfortably along the regression that shows a negative relationship between N_e and per-

generation mutation rate across animals (Fig 6a; Table S5). However, this relationship is not

apparent when considering primates only in which the two species with the largest N_e

(orangutan and mouse lemur) also have the highest mutation rate (Fig 6b).

Discussion

A High Mutation Rate in Mouse Lemurs

In this study, we provide the first pedigree-based estimate of the *de novo* mutation rate in a

strepsirrhine primate. Our mean mutation rate estimate was calculated to be 1.64×10^{-8}

mutations/site/generation, which is surprisingly high compared to previously characterized

primates with the exception of orangutan. We took several measures to ensure accurate

mutation rate estimation, including the use of allele-drop simulations to determine the

appropriate denominator for mutation rate calculations. Any point estimate of the mutation rate

should be interpreted with caution as there numerous variables that can impact rate estimates,

including biological factors such as variability among pedigrees (Smith *et al*, 2018). Moreover,

any mutation rate estimation is a direct result of accumulated study-design decisions made

regarding available animals, experimental planning, and data quality thresholds. The rate we

present is a product of these decisions. Some mistakes are inevitable, as represented by the

estimations of the number of false positives and false negatives in our dataset. To change any

of these inputs can yield a change in the final resulting mutation rate. Even so, a change in

assignment of 10% of the *de novo* mutation calls (roughly 13 mutations, twice the estimated

number of false positives), in either direction would only marginally alter the final published rate

and accompanying credible intervals. Changing the raw number of mutations discovered by

10% would leave the calculated rate, and credible intervals, within expectations based on the

rates recently measured in other primates (Fig 7).

We used linked-read sequencing technology that improves mapping accuracy to produce high

quality variants for the *de novo* mutations identified here. The linked reads also allowed us to

recover parental haplotypes and subsequently the parent-of-origin for observed mutations in

offspring (Fig. 1). The number of mutations with an assigned parent-of-origin is higher (70%) in

the present study than in analyses that only used standard short reads (Thomas *et al*, 2018;

Venn *et al*, 2014). And although a number of factors such as sequencing depth,

heterozygosity, and recombination rate may vary across investigations and limit the value of

cross-study comparisons, the prospect of successfully phasing more mutations while also

eliminating the need to sequence across more than two generations with linked-read data is

appealing. Even more surprising was the inferred mutation spectrum revealing low paternal

bias and a low frequency of CpG mutations, though several lines of evidence support these

results.

Low Numbers of Mutations at CpG Sites

CpG sites have generally been found to have higher mutation rates relative to other site

classes, a pattern discovered several decades ago using sequence comparisons (Bird, 1980)

and ascribed to the frequent deamination of methylated cytosines (Friedberg *et al*, 2005). Only

a four-fold enrichment of mutations at CpG sites (11 mutations, 8.2% of all mutations) was

found in mouse lemur, which is less than the at least ten-fold enrichment (12-25% of total

mutations) found in other primate studies (Besenbacher *et al*, 2019; Gao *et al*, 2019; Thomas

et al, 2018; Venn *et al*, 2014). It is thus reassuring that findings from our relaxed-clock

analyses of different substitution types are consistent with the observed mutation spectrum

(Fig. 2). Notably, the rate of C>T transitions at CpG sites breaks from the pattern expected

without partitioning (Supplementary Figure S14). This includes C>T transitions at non-CpG

sites (Supplementary Figures S25-S34) in mouse lemurs showing a higher substitution rate

than great apes and Old World monkeys but a lower rate than New World monkeys. That is,

mouse lemurs have the lowest rate of C>T transitions at CpG sites of all primates analyzed

here (Figs S25-S34). This leads to the hypothesis that methylation of CpG sites in mouse

lemur germ cell lines may actually be lower relative to that in other primates (Rahbari *et al*,

2016) thus ultimately contributing less to their mutation spectrum (Fig 2; Fig 3).

A lowered rate of C>T transitions at CpG sites is unprecedented for primate studies. Because

these mutations are caused by deamination of methylated cytosines, they are expected to

adhere to a strict clock. Previous studies of relative substitution rates using similar whole-

genome alignments have found that transitions at CpG sites are much more clock-like than

transitions at non-CpG sites when comparing great apes to Old World monkeys or New World

monkeys (Moorjani *et al*, 2016a). The same analyses of context-dependent substitution rates

also demonstrated clock-like behavior of C>T transitions at CpG sites across anthropoids (Lee

et al, 2015). In both cases, a single stepsirrhine (*Otolemur garnetii*) was treated as an outgroup

and rates within strepsirrhines were not estimated. However, earlier approaches for estimating

context-dependent substitution rates on a 1.7Mb region across mammals (Hwang and Green,

2004) also discovered lowered relative C>T transition rates at CpG sites in lemurs and their

common ancestor when compared to anthropoids, although we also found a notably elevated

rate in New World monkeys (*Callithrix jacchus*; Fig. 3). New World monkeys have been shown

to have rates of transitions at CpG sites approximately 20% higher than great apes (Moorjani

et al, 2016a), but past analyses with context-dependent substitution rates on a 0.15Mb

alignment have suggested much more clock-like behavior (Lee *et al*, 2015). We anticipate that

future analyses with denser sampling of New World monkeys and strepsirrhines will be

necessary to rigorously test clock-like behavior of C>T transitions at CpG sites in primates.

The Mouse Lemur Mutation Spectrum

Our estimates of the Ti:Tv and SW:WS ratios at 1.03 each are lower than values found in other

animals. For instance, Ti:Tv ratio in previous pedigree-based studies varied between 1.97 and

2.67 (Agier and Fischer, 2012; Assaf *et al*, 2017; Besenbacher *et al*, 2019; Kong *et al*, 2012;

Smeds *et al*, 2016; Thomas *et al*, 2018; Venn *et al*, 2014). The finding of a lower Ti:Tv ratio is

likely a consequence of the relatively low number of C>T transitions, especially at CpG sites.

For example, C>T transitions are twice as frequent as A>G transitions in human, chimp, and

owl monkey (Thomas *et al*, 2018; Venn *et al*, 2014), but these two mutation classes occur in

equal frequency in mouse lemur (Fig. 2a). These findings also explain the SW:WS ratio near 1,

since C>T mutations are strong-to-weak transitions. For instance, without an elevation in the

mutation rate at CpG sites, the Ti:Tv and SW:WS ratios would drop from 2.06 and 2.11 to 1.46

and 1.33 respectively, in a study of chimpanzees (Venn *et al*, 2014). Thus, reduced numbers

of C>T transitions at CpG sites can explain several metrics of the mouse lemur mutation

spectrum that deviate from previous studies of primate mutation rates.

Reduced Male Mutational Bias

A paternal mutational bias has long been hypothesized for diploid sexually reproducing

organisms based on the idea that the increased number of cell divisions in sperm versus egg

should lead to higher numbers of mutations in the male germline than the female germline

(Haldane, 1947; Kong *et al*, 2012; Lindsay *et al*, 2019). Indeed, a strong paternal mutation rate

bias has been observed in the vast majority of pedigree-based mutation rate estimates to date

(Gao *et al*, 2019; Lindsay *et al*, 2019; Rahbari *et al*, 2016; Thomas *et al*, 2018; Venn *et al*,

2014) and in many studies of phylogenetically-based rates (Axelsson *et al*, 2004; Ellegren and

Fridolfsson, 1997; Goetting-Minesky and Makova, 2006; Shimmin *et al*, 1993; Zhang, 2004).

The 1.2 ratio of paternal-to-maternal mutations in gray mouse lemur observed here is

considerably lower than the range between 2.1 and 5.5 observed in other primate species

(Gao *et al*, 2019; Lindsay *et al*, 2019; Rahbari *et al*, 2016; Thomas *et al*, 2018; Venn *et al*,

2014) and 2.7 in mouse (Lindsay *et al*, 2019). It is identical, however, to the ratio found in

collared flycatchers (Smeds *et al*, 2016), which suggests that the low sex bias ratio observed in

the gray mouse lemur is not unreasonable in the larger context of vertebrates. One of the

driving factors of the paternal mutational bias is likely the time of first reproduction after puberty

(Segurel *et al*, 2014), and mouse lemurs experience puberty and time of first reproduction

nearly simultaneously (Blanco, 2011; Blanco *et al*, 2015; Zohdy *et al*, 2014). Mouse lemurs are

capable of reproduction in one year, and the sire for our focal quartet was 4.1 and 5 years old

at the time of conception of the male and female offspring respectively (Fig. 1; Fig. S1). This

may not be enough time for a strong male mutational bias to manifest relative to longer-lived

species where more mutations in the male germline would be anticipated (Kong *et al*, 2012;

Thomas *et al*, 2018). Additionally, there are differences in the methylation process within male

and female germline cells, with male cells experiencing markedly more methylation (Kobayashi

et al, 2013; Reik and Dean, 2001). This discrepancy yields more methylation-related mutations

in males than females as mammals age (Gao *et al*, 2019). Thus, fewer methylation-related (i.e.

CpG) mutations, and a short time to puberty in mouse lemurs may in combination lead to the

observed, limited sex bias. As a potential caveat, mouse lemurs have exaggerated symptoms

of sperm competition (Eberle and Kappeler, 2004), and large testes size that is correlated with

their high substitution rate (Wong, 2014), but it is unclear if these traits and sperm competition

should lead to elevated male mutation rates.

Mutation Rates and Life History Traits

The drift barrier hypothesis states that N_e may explain variation in mutation rates across

species due to a larger efficiency of selection acting on DNA replication fidelity in larger

populations, especially across large phylogenetic distances (Lynch, 2010; Sung *et al*, 2012).

While such trends can be observed across eukaryotes (Fig. 6a), it is not apparent among

primates (Fig. 6b), in which, for example, mouse lemurs have both a relatively high mutation

rate and high N_e. It is possible that the drift barrier hypothesis does not operate across these

relatively small timescales and differences in population size. Though mouse lemurs are

diverged from the other primates compared here, with a shared common ancestor that

predates the Cretaceous-Paleogene boundary approximately 65 Ma (dos Reis *et al*, 2018), this

is relatively insignificant with respect to the greater than 1 Ga age of the eukaryote common

ancestor (Betts *et al*, 2018). Clearly, further insight into the relationship between mutation rate,

N_e, and other life history traits within primates will require mutation rate estimates from

additional species, especially within the strepsirrhine clade which represents nearly half of all

living primate species.

Mutation and substitution rates

There is disagreement between the magnitude of mutation rates and phylogenetic substitution

rates when attempting to scale the two similarly (Fig. 4). Several sources of uncertainty

underlie both, especially error in generation time estimation which is needed to scale

phylogenetic substitution rates to per-generation units or scale pedigree-based mutation rates

to per-year units. Additionally, pedigree-based mutation rates offer only a sample of the

present, and both mutation rate and generation time may have varied through time (Moorjani

et al, 2016b). For example, one revelation in the rapidly developing literature on *de novo*

mutation rates has been that the estimated rate in humans is less than half that predicted by

phylogenetic studies, suggesting that the mutation rate has slowed down over time in humans

and that rates can change rapidly among primates (Scally and Durbin, 2012).

Phylogenetically based estimates may be biased downwards if substitutions are not fully

neutral. Substitution rates used for comparison with generation times and mutation rates were

based on third-codon positions from a supermatrix of different data types (dos Reis *et al*, 2018;

Springer *et al*, 2012) and may be under weak purifying selection (Fig 4). Indeed, previous

studies have found evidence for low phylogenetically based compared to pedigree-based

estimates (Denver *et al*, 2000; Howell *et al*, 2003; Winter *et al*, 2018). For pedigree-based

estimates, the degree to which somatic mutations and/or inter-individual variation might impact

these estimates is not clear (Segurel *et al*, 2014). Additional data and analyses will be needed

to reconcile the differences between pedigree-based and phylogenetic estimates of the

mutation rate.

Mutation Rates and Divergence Time Estimates

Application of the pedigree-based mutation rate estimate observed in this study leads to more

recent divergence times among mouse lemur species than previously inferred (Fig 5; Table

S4). These divergence times are obtained by rescaling coalescent units to absolute time given

a mutation rate and generation time (Burgess and Yang, 2008) as opposed to relaxed-clock

phylogenetic methods that estimated older species divergences within mouse lemurs (dos

Reis *et al*, 2018; Yang and Yoder, 2003). A previous analysis made assumptions regarding

mutation rate in mouse lemurs (Yoder *et al*, 2016) that resulted in divergence times twice as

old as those presented here (Fig 5; Table S4). Although such assumptions regarding mutation

rates are reasonable in the absence of data, direct mutation rates from pedigrees can arguably

produce more accurate divergence time estimates.

Similarly, a recent investigation of great ape divergence times revealed that lineage-specific

mutation rate estimates for chimp, gorilla, and orangutan led to more recent estimates of

common ancestor ages compared to those that assumed that all species share the human

mutation rate (Besenbacher *et al*, 2019). Moreover, divergence time estimates obtained with

these lineage-specific mutation rates agreed more closely with available fossil evidence.

Unfortunately, a complete lack of lemuriform fossils means that we cannot similarly evaluate

the accuracy of divergence time estimates for mouse lemurs in the context of the fossil record.

The findings from great apes would nevertheless suggest that divergence times in mouse

lemurs have been previously overestimated, and that these cryptic primates have diversified

more recently than previously appreciated. Given the endangered status of many mouse lemur

species, and virtually all other strepsirrhine species, an enhanced ability to provide a temporal

context to speciation and to estimate demographic parameters such as N_e may yield critical

information for directing ongoing conservation policy and efforts.

Conclusions

Our study emphasizes the importance of increased sampling across the tree of life for gaining

insight into the nature and causes of mutation rate evolution. This is the first pedigree-based

mutation rate estimate for a strepsirrhine primate, and as such, it is not clear whether the high

mutation rate, low CpG mutation rate, and weak sex bias is specific to mouse lemurs or may

be representative of strepsirrhines more generally. We showed that N_e correlates with mutation

rate at broad phylogenetic scales but does not appear to do so within primates. These

comparisons are presently restricted to relatively few primate species, however, and new

trends between mutation rates and life history traits are likely to be revealed as more data

become available, especially as the range of species includes more of the extensive variation

in life history strategies among primates. Reconciling the disparity in magnitude between

mutation rates from pedigrees and substitution rates from phylogenetic methods will be a focus

of future work as more pedigree-based mutation rates become available. As demonstrated by

this study in mouse lemurs, *de novo* mutation rate estimates stand to drastically revise

divergence times, especially in recent evolutionary radiations.

Acknowledgments

We thank the Duke Lemur Center staff, especially Erin Ehmke, Bobby Schopler, and Cathy

Williams, for providing tissue samples. Priya Moorjani, Susanne Pfeifer, Jonathan Pritchard,

Molly Przeworski, and Meredith Yeager all provided helpful discussions in the development of

this project. We especially thank Jonathan Pritchard for his suggestion that substitution rate

analysis could be useful for verifying the observed mutation rate spectrum. Simon Gregory's

lab prepared the 10X Genomics libraries and we are grateful for the support of Duke Research

Computing and the Duke Data Commons (NIH 1S10OD018164-01). This study was funded by

a National Science Foundation Grant DEB-1354610 and Duke University startup funds to ADY

and she gratefully acknowledges support from the John Simon Guggenheim Foundation and

the Alexander von Humboldt Foundation during the writing phase of this project. JLT was

supported by National Science Foundation Grant DEB-1754142. This is Duke Lemur Center

publication no. XXXX.

Competing Interests

The authors declare no competing interests.

Data Archiving

Data are available from the Dryad Digital Repository https://doi.org/10.5061/dryad.8pk0p2njx.
Raw sequence data are available through NCBI under BioProject PRJNA512515.
608

References

Agier N, Fischer G (2012). The mutational profile of the yeast genome is shaped by replication.

Mol Biol Evol **29**(3)**:** 905-913.

Andriatsitohaina B, Ramsay MS, Kiene F, Lehman SM, Rasoloharijaona S, Rakotondravony R

et al (2020). Ecological fragmentation effects in mouse lemurs and small mammals in

northwestern Madagascar. *Am J Primatol* **82**(4)**:** e23059.

Assaf ZJ, Tilk S, Park J, Siegal ML, Petrov DA (2017). Deep sequencing of natural and

experimental populations of Drosophila melanogaster reveals biases in the spectrum of new

mutations. *Genome Res* **27**(12)**:** 1988-2000.

Axelsson E, Smith NG, Sundstrom H, Berlin S, Ellegren H (2004). Male-biased mutation rate

and divergence in autosomal, z-linked and w-linked introns of chicken and Turkey. *Mol Biol*

Evol **21**(8)**:** 1538-1547.

Besenbacher S, Hvilsom C, Marques-Bonet T, Mailund T, Schierup MH (2019). Direct

estimation of mutations in great apes reconciles phylogenetic dating. *Nat Ecol Evol* **3**(2)**:** 286-

292.

Betts HC, Puttick MN, Clark JW, Williams TA, Donoghue PCJ, Pisani D (2018). Integrated

genomic and fossil evidence illuminates life's early evolution and eukaryote origin. *Nat Ecol*

Evol **2**(10)**:** 1556-1562.

Bird AP (1980). DNA methylation and the frequency of CpG in animal DNA. *Nucleic Acids Res*
8(7)**:** 1499-1504.

Bird AP (1986). CpG-rich islands and the function of DNA methylation. *Nature* **321**(6067)**:** 209-
213.

Blanco MB (2011). Timely estrus in wild brown mouse lemur females at Ranomafana National
Park, southeastern Madagascar. *Am J Phys Anthropol* **145**(2)**:** 311-317.

Blanco MB, Rasoazanabary E, Godfrey LR (2015). Unpredictable environments, opportunistic
responses: Reproduction and population turnover in two wild mouse lemur species
(Microcebus rufus and M. griseorufus) from eastern and western Madagascar. *Am J Primatol*
77(9)**:** 936-947.

Burgess R, Yang Z (2008). Estimation of hominoid ancestral population sizes under bayesian
coalescent models incorporating mutation rate variation and sequencing errors. *Mol Biol Evol*
25(9)**:** 1979-1994.

Chaisson MJ, Wilson RK, Eichler EE (2015). Genetic variation and the de novo assembly of
human genomes. *Nat Rev Genet* **16**(11)**:** 627-640.

Chojnacki S, Cowley A, Lee J, Foix A, Lopez R (2017). Programmatic access to bioinformatics

tools from EMBL-EBI update: 2017. *Nucleic Acids Res* **45**(W1)**:** W550-W553.

Denver DR, Morris K, Lynch M, Vassilieva LL, Thomas WK (2000). High direct estimate of the

mutation rate in the mitochondrial genome of Caenorhabditis elegans. *Science* **289**(5488)**:**

2342-2344.

dos Reis M, Gunnell GF, Barba-Montoya J, Wilkins A, Yang Z, Yoder AD (2018). Using

Phylogenomic Data to Explore the Effects of Relaxed Clocks and Calibration Strategies on

Divergence Time Estimation: Primates as a Test Case. *Syst Biol* **67**(4)**:** 594-615.

dos Reis M, Yang Z (2011). Approximate likelihood calculation on a phylogeny for Bayesian

estimation of divergence times. *Mol Biol Evol* **28**(7)**:** 2161-2172.

Drummond AJ, Ho SY, Phillips MJ, Rambaut A (2006). Relaxed phylogenetics and dating with

confidence. *PLoS Biol* **4**(5)**:** e88.

Dudchenko O, Batra SS, Omer AD, Nyquist SK, Hoeger M, Durand NC *et al* (2017). De novo

assembly of the Aedes aegypti genome using Hi-C yields chromosome-length scaffolds.

Science **356**(6333)**:** 92-95.

Eberle M, Kappeler PM (2004). Sex in the dark: determinants and consequences of mixed

male mating tactics in Microcebus murinus, a small solitary nocturnal primate. *Behavioral*

Ecology and Sociobiology **57:** 77-90.

Ellegren H, Fridolfsson AK (1997). Male-driven evolution of DNA sequences in birds. *Nat*

Genet **17**(2)**:** 182-184.

Ewing AD, Houlahan KE, Hu Y, Ellrott K, Caloian C, Yamaguchi TN *et al* (2015). Combining

tumor genome simulation with crowdsourcing to benchmark somatic single-nucleotide-variant

detection. *Nat Methods* **12**(7)**:** 623-630.

Ezran C, Karanewsky CJ, Pendleton JL, Sholtz A, Krasnow MR, Willick J *et al* (2017). The

Mouse Lemur, a Genetic Model Organism for Primate Biology, Behavior, and Health. *Genetics*

206(2)**:** 651-664.

Feng C, Pettersson M, Lamichhaney S, Rubin CJ, Rafati N, Casini M *et al* (2017). Moderate

nucleotide diversity in the Atlantic herring is associated with a low mutation rate. *Elife* **6**.

Friedberg EC, Walker GC, Siede W, Wood RD (2005). *DNA repair and mutagenesis*. American

Society for Microbiology Press.

Gao Z, Moorjani P, Sasani TA, Pedersen BS, Quinlan AR, Jorde LB *et al* (2019). Overlooked

roles of DNA damage and maternal age in generating human germline mutations. *Proc Natl*

Acad Sci U S A **116**(19)**:** 9491-9500.

Goetting-Minesky MP, Makova KD (2006). Mammalian male mutation bias: impacts of

generation time and regional variation in substitution rates. *J Mol Evol* **63**(4)**:** 537-544.

Haldane JB (1947). The mutation rate of the gene for haemophilia, and its segregation ratios in

males and females. *Ann Eugen* **13**(4)**:** 262-271.

Harris K, Pritchard JK (2017). Rapid evolution of the human mutation spectrum. *Elife* **6**.

Heath TA, Huelsenbeck JP, Stadler T (2014). The fossilized birth-death process for coherent

calibration of divergence-time estimates. *Proc Natl Acad Sci U S A* **111**(29)**:** E2957-2966.

Herrera JP, Davalos LM (2016). Phylogeny and Divergence Times of Lemurs Inferred with

Recent and Ancient Fossils in the Tree. *Syst Biol* **65**(5)**:** 772-791.

Hotaling S, Foley ME, Lawrence NM, Bocanegra J, Blanco MB, Rasoloarison R *et al* (2016).

Species discovery and validation in a cryptic radiation of endangered primates: coalescent-

based species delimitation in Madagascar's mouse lemurs. *Mol Ecol* **25**(9)**:** 2029-2045.

Howell N, Smejkal CB, Mackey DA, Chinnery PF, Turnbull DM, Herrnstadt C (2003). The

pedigree rate of sequence divergence in the human mitochondrial genome: there is a

difference between phylogenetic and pedigree rates. *Am J Hum Genet* **72**(3)**:** 659-670.

Hozer C, Pifferi F, Aujard F, Perret M (2019). The Biological Clock in Gray Mouse Lemur:

Adaptive, Evolutionary and Aging Considerations in an Emerging Non-human Primate Model.

Front Physiol **10:** 1033.

Hwang DG, Green P (2004). Bayesian Markov chain Monte Carlo sequence analysis reveals

varying neutral substitution patterns in mammalian evolution. *Proc Natl Acad Sci U S A*

101(39)**:** 13994-14001.

Jonsson H, Sulem P, Kehr B, Kristmundsdottir S, Zink F, Hjartarson E *et al* (2017). Parental

influence on human germline de novo mutations in 1,548 trios from Iceland. *Nature* **549**(7673)**:**

519-522.

Keightley PD, Pinharanda A, Ness RW, Simpson F, Dasmahapatra KK, Mallet J *et al* (2015).

Estimation of the spontaneous mutation rate in Heliconius melpomene. *Mol Biol Evol* **32**(1)**:**

239-243.

Kimura M (1983). *The neutral theory of molecular evolution*. Cambridge University Press.

Kimura M, Ohta T (1971). On the rate of molecular evolution. *J Mol Evol* **1**(1)**:** 1-17.

Kobayashi H, Sakurai T, Miura F, Imai M, Mochiduki K, Yanagisawa E *et al* (2013). High-

resolution DNA methylome analysis of primordial germ cells identifies gender-specific

reprogramming in mice. *Genome Res* **23**(4)**:** 616-627.

Koch E, Schweizer RM, Schweizer TM, Stahler DR, Smith DW, Wayne RK *et al* (2019). De

novo mutation rate estimation in wolves of known pedigree. *Mol Biol Evol*.

Kondrashov FA, Kondrashov AS (2010). Measurements of spontaneous rates of mutations in

the recent past and the near future. *Philos Trans R Soc Lond B Biol Sci* **365**(1544)**:** 1169-

1176.

Kong A, Frigge ML, Masson G, Besenbacher S, Sulem P, Magnusson G *et al* (2012). Rate of

de novo mutations and the importance of father's age to disease risk. *Nature* **488**(7412)**:** 471-

475.

Krasovec M, Sanchez-Brosseau S, Piganeau G (2019). First Estimation of the Spontaneous

Mutation Rate in Diatoms. *Genome Biol Evol* **11**(7)**:** 1829-1837.

Langley CH, Fitch WM (1974). An examination of the constancy of the rate of molecular

evolution. *J Mol Evol* **3**(3)**:** 161-177.

Larsen PA, Harris RA, Liu Y, Murali SC, Campbell CR, Brown AD *et al* (2017). Hybrid de novo

genome assembly and centromere characterization of the gray mouse lemur (Microcebus

murinus). *BMC Biol* **15**(1)**:** 110.

Lartillot N, Philippe H (2004). A Bayesian mixture model for across-site heterogeneities in the

amino-acid replacement process. *Mol Biol Evol* **21**(6)**:** 1095-1109.

Lartillot N, Rodrigue N, Stubbs D, Richer J (2013). PhyloBayes MPI: phylogenetic

reconstruction with infinite mixtures of profiles in a parallel environment. *Syst Biol* **62**(4)**:** 611-

615.

Lee HJ, Rodrigue N, Thorne JL (2015). Relaxing the Molecular Clock to Different Degrees for

Different Substitution Types. *Mol Biol Evol* **32**(8)**:** 1948-1961.

Lindsay SJ, Rahbari R, Kaplanis J, Keane T, Hurles ME (2019). Similarities and differences in

patterns of germline mutation between mice and humans. *Nat Commun* **10**(1)**:** 4053.

Long H, Winter DJ, Chang AY, Sung W, Wu SH, Balboa M *et al* (2016). Low Base-Substitution

Mutation Rate in the Germline Genome of the Ciliate Tetrahymena thermophil. *Genome Biol*

Evol **8**(12)**:** 3629-3639.

Lynch M (2010). Evolution of the mutation rate. *Trends Genet* **26**(8)**:** 345-352.

Magallon SA, Sanderson MJ (2005). Angiosperm divergence times: the effect of genes, codon

positions, and time constraints. *Evolution* **59**(8)**:** 1653-1670.

Martin HC, Batty EM, Hussin J, Westall P, Daish T, Kolomyjec S *et al* (2018). Insights into

Platypus Population Structure and History from Whole-Genome Sequencing. *Mol Biol Evol*

35(5)**:** 1238-1252.

McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A *et al* (2010). The

Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA

sequencing data. *Genome Res* **20**(9)**:** 1297-1303.

Molaro A, Hodges E, Fang F, Song Q, McCombie WR, Hannon GJ *et al* (2011). Sperm

methylation profiles reveal features of epigenetic inheritance and evolution in primates. *Cell*

146(6)**:** 1029-1041.

Moorjani P, Amorim CE, Arndt PF, Przeworski M (2016a). Variation in the molecular clock of

primates. *Proc Natl Acad Sci U S A* **113**(38)**:** 10607-10612.

Moorjani P, Gao Z, Przeworski M (2016b). Human Germline Mutation and the Erratic

Evolutionary Clock. *PLoS Biol* **14**(10)**:** e2000744.

Near TJ, Bolnick DI, Wainwright PC (2005). Fossil calibrations and molecular divergence time

estimates in centrarchid fishes (Teleostei: Centrarchidae). *Evolution* **59**(8)**:** 1768-1782.

Ogilvie HA, Bouckaert RR, Drummond AJ (2017). StarBEAST2 Brings Faster Species Tree

Inference and Accurate Estimates of Substitution Rates. *Mol Biol Evol* **34**(8)**:** 2101-2114.

Pfeifer SP (2017). Direct estimate of the spontaneous germ line mutation rate in African green

monkeys. *Evolution* **71**(12)**:** 2858-2870.

Plummer M, Best N, Cowles K, Vines K (2006). CODA: Convergence diagnosis and output

analysis for MCMC. *R News* **6:** 7-11.

Rahbari R, Wuster A, Lindsay SJ, Hardwick RJ, Alexandrov LB, Turki SA *et al* (2016). Timing,

rates and spectra of human germline mutation. *Nat Genet* **48**(2)**:** 126-133.

Ramu A, Noordam MJ, Schwartz RS, Wuster A, Hurles ME, Cartwright RA *et al* (2013).

DeNovoGear: de novo indel and point mutation discovery and phasing. *Nat Methods* **10**(10)**:**

985-987.

Reik W, Dean W (2001). DNA methylation and mammalian epigenetics. *Electrophoresis*

22(14)**:** 2838-2843.

Sanderson MJ (2002). Estimating absolute rates of molecular evolution and divergence times:

a penalized likelihood approach. *Mol Biol Evol* **19**(1)**:** 101-109.

Scally A, Durbin R (2012). Revising the human mutation rate: implications for understanding

human evolution. *Nat Rev Genet* **13**(10)**:** 745-753.

Schiffels S, Durbin R (2014). Inferring human population size and separation history from

multiple genome sequences. *Nat Genet* **46**(8)**:** 919-925.

Segurel L, Wyman MJ, Przeworski M (2014). Determinants of mutation rate variation in the

human germline. *Annu Rev Genomics Hum Genet* **15:** 47-70.

Setash CM, Zohdy S, Gerber BD, Karanewsky CJ (2017). A biogeographical perspective on

the variation in mouse lemur density throughout Madagascar. *Mammal Review* **47**(3)**:** 212-229.

Shimmin LC, Chang BH, Li WH (1993). Male-driven evolution of DNA sequences. *Nature*

362(6422)**:** 745-747.

Smeds L, Qvarnstrom A, Ellegren H (2016). Direct estimate of the rate of germline mutation in

a bird. *Genome Res* **26**(9)**:** 1211-1218.

Smith TCA, Arndt PF, Eyre-Walker A (2018). Large scale variation in the rate of germ-line de

novo mutation, base composition, divergence and diversity in humans. *PLoS Genet* **14**(3)**:**

e1007254.

Springer MS, Meredith RW, Gatesy J, Emerling CA, Park J, Rabosky DL *et al* (2012).

Macroevolutionary dynamics and historical biogeography of primate diversification inferred

from a species supermatrix. *PLoS One* **7**(11)**:** e49521.

Sung W, Ackerman MS, Miller SF, Doak TG, Lynch M (2012). Drift-barrier hypothesis and

mutation-rate evolution. *Proc Natl Acad Sci U S A* **109**(45)**:** 18488-18492.

Thomas GWC, Wang RJ, Puri A, Harris RA, Raveendran M, Hughes DST *et al* (2018).

Reproductive Longevity Predicts Mutation Rates in Primates. *Curr Biol* **28**(19)**:** 3193-3197

e3195.

Thorne JL, Kishino H (2002). Divergence time and evolutionary rate estimation with multilocus

data. *Syst Biol* **51**(5)**:** 689-702.

Thorne JL, Kishino H, Painter IS (1998). Estimating the rate of evolution of the rate of

molecular evolution. *Mol Biol Evol* **15**(12)**:** 1647-1657.

Uchimura A, Higuchi M, Minakuchi Y, Ohno M, Toyoda A, Fujiyama A *et al* (2015). Germline

mutation rates and the long-term phenotypic effects of mutation accumulation in wild-type

laboratory mice and mutator mice. *Genome Res* **25**(8)**:** 1125-1134.

Van der Auwera GA, Carneiro MO, Hartl C, Poplin R, Del Angel G, Levy-Moonshine A *et al*

(2013). From FastQ data to high confidence variant calls: the Genome Analysis Toolkit best

practices pipeline. *Curr Protoc Bioinformatics* **43:** 11 10 11-11 10 33.

Venn O, Turner I, Mathieson I, de Groot N, Bontrop R, McVean G (2014). Nonhuman genetics.

Strong male bias drives germline mutation in chimpanzees. *Science* **344**(6189)**:** 1272-1275.

Weisenfeld NI, Kumar V, Shah P, Church DM, Jaffe DB (2017). Direct determination of diploid

genome sequences. *Genome Res* **27**(5)**:** 757-767.

Winter DJ, Wu SH, Howell AA, Azevedo RBR, Zufall RA, Cartwright RA (2018). accuMUlate: a

mutation caller designed for mutation accumulation experiments. *Bioinformatics* **34**(15)**:** 2659-

2660.

Wong A (2014). Covariance between testes size and substitution rates in primates. *Mol Biol*

Evol **31**(6)**:** 1432-1436.

Xie Z, Wang L, Wang L, Wang Z, Lu Z, Tian D *et al* (2016). Mutation rate analysis via parent-

progeny sequencing of the perennial peach. I. A low rate in woody perennials and a higher

mutagenicity in hybrids. *Proc Biol Sci* **283**(1841).

Yang Z (2007). PAML 4: phylogenetic analysis by maximum likelihood. *Mol Biol Evol* **24**(8)**:**

1586-1591.

Yang Z (2015). The BPP program for species tree estimation and species delimitation. *Current*

Zoology **61:** 854-865.

Yang Z, Yoder AD (2003). Comparison of likelihood and Bayesian methods for estimating

divergence times using multiple gene Loci and calibration points, with application to a radiation

of cute-looking mouse lemur species. *Syst Biol* **52**(5)**:** 705-716.

Yoder AD, Campbell CR, Blanco MB, Dos Reis M, Ganzhorn JU, Goodman SM *et al* (2016).

Geogenetic patterns in mouse lemurs (genus Microcebus) reveal the ghosts of Madagascar's

forests past. *Proc Natl Acad Sci U S A* **113**(29)**:** 8049-8056.

Zhang J (2004). Evolution of DMY, a newly emergent male sex-determination gene of medaka

fish. *Genetics* **166**(4)**:** 1887-1895.

Zohdy S, Gerber BD, Tecot S, Blanco MB, Winchester JM, Wright PC *et al* (2014). Teeth, sex,

and testosterone: aging in the world's smallest primate. *PLoS One* **9**(10)**:** e109528.

Zuckerkandl E, Pauling L (1965). Molecules as documents of evolutionary history. *J Theor Biol*

8(2)**:** 357-366.

Figures

Figure 1 – Focal family quartet. Parents (P) and offspring (O) are subscripted as male (m) or

female (f). Lines represent familial relationships as in a traditional pedigree, with thickness and

color reflecting the number and source of *de novo* mutations passed down (red is from male

parent, blue from female parent, gray is undetermined origin). Color of line represents source

and shading represents destination (lighter shading to O_m, darker to O_f). Numbers within bars

show mutation counts and the rate of each individual offspring is listed below.

Figure 2 – Mutation spectrum of the Gray Mouse Lemur. (A) Counts of *de novo* mutations

from the pedigree analysis. Mutation types are broken down by weak-to-strong transversions

(A>C and T>G), weak-to-strong transitions (A>G and T>C), weak-to-weak transversions (A>T

and T>A), strong-to-weak transversions (C>A and G>T), strong-to-strong transversions (C>G

and G>C), and strong-to-weak transitions (C>T and G>A). Complementary mutation types are

shown together. (B) Context-dependent substitution rate estimates. Nine possible substitution

type parameters are shown for the gray mouse lemur terminal branch, which are categorized

similarly as the *de novo* mutation spectrum. Error bars on substitution rates represent 95%

highest posterior densities.

Figure 3 – Context-dependent relaxed-clock analysis shows low rates of C>T

substitution rates at CpG sites in the Gray Mouse Lemur. C>T substitution rate estimates

at non-CpG versus CpG sites are compared for six species of primate, including the gray

mouse lemur (*M. murinus*). Note that with the exception of *M. murinus*, all primates examined

show significantly higher CpG rates than non-CpG rates. The C>T substitution rates at non-

CpG and CpG sites are nearly identical in *M. murinus*. Error bars represent 95% highest

posterior densities.

Figure 4 – Difference between mutation rates and substitution rates among primates.

Error bars around substitution rates are 95% highest posterior density intervals from a

Bayesian relaxed-clock analysis. Credible intervals are given for mutation rates where

available from published data. Data are given in Table S3.

Figure 5 – Estimated divergence times among mouse lemur species. Trees are posterior

samples from BPP based on a fixed previously published topology. The directly estimated

mutation rate (blue) is nearly twice as high as the previously assumed rate (red). Divergence

times estimated with the new mutation rate are nearly half of the previous estimates. Summary

statistics are given in Table S4, matched by node labels (A-E).

Figure 6 – Relationship between mutation rate and effective population size. (A) There is

a negative correlation between mutation rate and N_e for animals, but (B) this trend is not

present when considering primates alone. Mutation rate *and N_e* data are given in Table S5.

Both x- and y-axes are on a log scale.

Figure 7 – Effect of filtering thresholds on mutation rate estimation. A) The mutation rate

of mouse lemur (purple) with upper and lower confidence intervals in red and blue, when

calculated for differing numbers of discovered mutations. The highlighted dots reflect an

increase and decrease of discovered mutations by 10%. B) The directly estimated mutation

rate and confidence intervals of other species of primates. Although it is difficult to make

predictions on how changes to experimental design or methods will affect mutation rate

estimates, it is possible to examine how discovering more or less mutations would affect the

rate.

978 **Tables**

979 **Table 1 – Directly estimated mammalian mutation rates.**

Species	Common Name	Citation	Rate
Pongo abelii	Orangutan	Besenbacher et al. 2019	1.66×10^{-8}
Pan troglodytes	Chimpanzee	Tatsumoto et al. 2017	1.48×10^{-8}
Homo sapiens	Human	Jonsson et al. 2017	1.29×10^{-8}
Chlorocebus sabaeus	Green Monkey	Pfieffer 2017	9.4×10^{-9}
Aotus nancymaae	Owl Monkey	Thomas et al. 2018	8.1×10^{-9}
Mus musculus[†]	House Mouse	Uchimura et al. 2015	5.4×10^{-9}
Canis lupus	Wolf	Koch et al. 2019	4.5×10^{-9}

[†]Mutation Accumulation lines were used instead of parent-progeny comparisons