The evolution of paternal care: a role for microbes?

Authors

Yael Gurevich, Ohad Lewin-Epstein, Lilach Hadany

Affiliations

School of Plant Sciences and Food Security, Tel-Aviv University, Tel-Aviv, 6997801, Israel

Correspondence: Lilach Hadany (lilach.hadany@gmail.com)

Abstract

Paternal care is an evolutionary mystery. Despite extensive research, both theoretical and experimental, the reasons for its ubiquity remain unclear. Common explanations include kin selection and limited accuracy in parentage assessment. However, these explanations do not cover the breadth of circumstances in which paternal care has been observed, particularly in cases of uncertain paternity. Here we propose that microbes may play a key role in the evolution of paternal care among their hosts. Using computational models, we demonstrate that microbes associated with increased paternal care could be favoured by natural selection. We find that microbe-induced paternal care could evolve under wider conditions than suggested by genetic models. Moreover, we show that microbe-induced paternal care is more likely to evolve when considering paternal care interactions that increase microbial transmission, such as feeding and grooming. Our results imply that factors affecting the composition of host microbiome may also alter paternal behaviour.

Keywords

Microbiome, paternal care, mathematical model, extra-pair mating, sexual conflict, nongenetic inheritance

Introduction

When would a father benefit more from caring for its offspring, rather than looking for additional mating opportunities? This question has been broadly addressed both theoretically and experimentally. Paternal care was frequently observed among avian species[1] (~85%), and was also found in mammalian species[2] (~5%), amphibians[3], and many species of fish[4]. Paternal care is most commonly observed alongside maternal care, while exclusive paternal care is rare[4,5]. A male may demonstrate care for its offspring with several types of interactions[4,5], such as feeding, grooming, or guarding against predators. It can also provide spousal care for the female while she cares for the young[6,7].

A commonly proposed explanation for the prevalence of paternal care is kin selection[8,9]. This explanation suggests that paternal care would be favoured whenever the paternal contribution to offspring survival increases paternal fitness more than pursuing additional mating opportunities[10,11]. In some settings, such as caring for unrelated young, this explanation is insufficient, and alternative explanations are suggested[12,13]. Interestingly, studies relating paternal effort to certainty of paternity obtained mixed results[14–20], and paternal care has been observed even in cases of very high probability of extra-pair paternity (e.g. in avian species[21] where extra-pair parenthood can range up to 95% in fairywrens[22]).

Here we consider the potential role of the microbes in host paternal care. The microbiome is a significant agent affecting host health and behaviour[23–27], through pathways such as the 'gut-brain axis'[28]. Numerous studies have demonstrated a possible association between microbes and social behaviour[29–34], and certain species of microbiome have been showed to alleviate symptoms of anxiety and depression[32] and improve social interactions[35]. Microbes are highly heritable, through gestation/incubation[36,37] or parental care[38–40]. Microbes can also be transmitted horizontally in a social setting[41], through interactions such as feeding, grooming

and copulation[40]. The effect of microbes on host behaviour has given rise to the idea that host

manipulation by microorganisms may be driven by natural selection on the microbes[28].

Selection could drive such an effect when the induced behaviour increases microbial fitness,

for example by increasing the rate of microbial transmission or proliferation[28], including host

proliferation. Previous theoretical studies suggested that by encouraging host sociality[42] or

altruistic behaviour[29], the microbes can help their own propagation.

We integrated the notion of microbe-associated behaviour into mathematical models for the

evolution of paternal care. A family is a unit with a high probability of microbial

transmission[43], since the members of the family partake in frequent and profound

interactions. Caring for the young presents an excellent opportunity from microbial

perspective, since providing care both increases odds of offspring survival[10] and establishes a

higher transmission probability. Therefore, a microbial gene that is associated with host intra-

family caring behaviour could be favoured by natural selection even when encouraging care

towards genetically unrelated young individuals. The propagation of microbes carrying these

genes may have driven the evolution of paternal care even in the absence of paternity.

Potential candidate microbes, such as Lactobacillus[44] and Bifidobacteria[45], have been shown

to be impact mood or emotional behaviour and could be associated with paternal care.

Results

68 We examine two possible family structures. In every family, there is one female, who is

69 mother to all offspring in the family. In the first family structure, all offspring are fathered by

70 a single male, making them full siblings. In the second family type, we allow mixed paternity

71 within the brood, while still maintaining a social pair structure.

Model I: Broods of full siblings

In this model, males adopt one of two pure strategies, either paternal care or lack thereof. Offspring fitness is increased by paternal care[10], due to provision and protection from predation, by a factor of $s > 0$. The fitness of an offspring whose father does not provide paternal care is $\omega_\beta = 1$, while an offspring that receives paternal care has increased fitness $\omega_\alpha = 1 + s$. Females mate with only one male. A male who provides paternal care can mate with one female and form a social pair. A male who does not provide paternal care can mate with more than one female and is limited only by female availability and receptivity[11]. The two types of males are subject to competition and sexual selection, wherein the caring males suffer a competitive disadvantage relative to the non-caring males (see[46] for discussion). The expected number of matings for both types of males is frequency-dependent[47], governed by the initial male composition in the mating pool. Only a fraction of the males of each type reach mating at all[48], thus the expected number of matings for a young male of the caring type is usually less than one. Total male mating opportunities are bound by the Fisher condition[49], with a mean of 1. We assume that there is a cost to paternal care[9], and that the expected number of matings for a caring male decreases with paternal care[11]. In both models, the number of matings for a caring male is given by $n_\alpha = \exp(-\gamma_c * s * (1 - p))$.

First, we examine a population of non-caring males where the paternal behaviour is genetically driven. A rare mutant providing paternal care will have a higher fitness than a non-caring male when (at $p = 0$):

$$1 + s > \frac{n_\beta}{n_\alpha} \tag{1}$$

Where n_β = expected number of matings for a non-caring male, n_α = expected number of matings for a caring male, and s = increase in offspring fitness due to paternal care.

Now, we extend the model to include microbes as a reproductive unit that can affect paternal

care behaviour. For simplicity, we neglect the effect of host genetic background in the

microbe model and assume that host paternal behaviour is determined by its microbes. Let us

consider microbes of type α, which are associated with paternal care behaviour, and microbes

of type β, which have no effect on paternal care behaviour. Microbes can be transmitted to

the offspring from the mother, with probability T_v, or from the father, with probability T_c

when the father cares for the offspring. If parental microbes fail to establish in the offspring,

it can adopt microbes horizontally by interacting with the general population[50], with

probability determined by the population frequencies. In many species, mate nurturing

behaviour is more common from father to mother than vice versa. In our model, microbes can

also be transmitted from the father to the mother during mating with probability T_m and

possibly through nurture of the mother, with probability T_n. For simplicity, we assume that

females carry the microbe, but it does not affect their behaviour. We assume that each host is

inhabited by a single type of microbe at a given time. A transmission probability thus

includes the probabilities that a microbe transmits to a new individual, establishes, and

replaces the resident microbe, encompassing the competition dynamics between different

microbial strains. The transmission pathways and transmission probabilities of the two

microbes are illustrated in Fig. 1. We consider a model where the mother cares for the

offspring[51,52], and additionally can transmit microbes during gestation[37] and natally[36], so

overall maternal transmission is higher than paternal transmission ($T_v > T_c$). We also assume

that paternal care involves more interaction – and potential for microbe transmission – than a

singular mating encounter ($T_c > T_m$). Since the probability of transmitting microbes during

mating is asymmetric between the sexes, with a higher probability for male-to-female

transmission[53], we neglect the probability of female-to-male transmission. We assume a delay

in the effect of the microbes on behaviour and neglect the possibility of a male altering its

paternal behaviour due to contracting different microbes at the mating stage. We initially

assume that males have full paternity in their brood and relax that assumption later (see Fig.

3).

Figure 1. Model 1 (full siblings): Illustration of microbe transmission pathways within the family. (a) where the father carries microbes of type α, inducing paternal care. (b) where the father carries microbes of type β, that have no effect on behaviour. Males carrying β do not care for the offspring and can be involved in n additional matings (illustrated is the case $n = 1$). T_v – vertical transmission probability through maternal influence (prenatal and postnatal). T_c – probability of transmission through paternal care. T_m - probability of male-to-female microbe transmission during mating. T_n – probability of transmission through male-to-female nurture.

123   We assume the following order of events within the reproductive process: transmission via

124   mating occurs first, second is maternal transmission, and finally transmission via paternal

125   care, if exists.

126   The condition for evolution of microbe type α (see Supplementary for full derivation) is

127   given by:

128
$$1 + s > \frac{n_\beta}{n_\alpha} \cdot \left(\frac{T_m \cdot T_v}{T_c + T_n \cdot T_v \cdot (1 - T_c)} \right) \tag{2}$$

129   Where n_β = expected number of matings for a non-caring male (at $p = 0$), n_α = expected

130   number of matings for a caring male (at $p = 0$), s = increase in offspring fitness due to

131   paternal care, T_v – vertical transmission probability through maternal. T_c – probability of

132   transmission through paternal care. T_m - probability of male-to-female microbe transmission

133   during mating. T_n – probability of transmission through male-to-female nurture.

Figure 2. Model I (full siblings): Microbes can expand the conditions for the evolution of paternal care. T_c – probability of microbe transmission from father to offspring through paternal care. T_m – probability of transmission from male to female during mating. The area below each graph represents the conditions allowing paternal care to evolve in the population. A microbe associated with paternal care behaviour can widen the range of conditions where paternal care prevails, and the effect increases with the transmission probability of the paternal microbes to the offspring during care. Other parameters: $T_n = 0$, $T_v = 0.8$. (i) $T_m = 0.01$, (ii) $T_m = 0.1$, (iii) $T_m = 0.2$.

Fig. 2 presents the range of fitness costs of caring imposed on the male that allows for the

evolution of paternal care in the model. When $p = 0$, then the number of matings for a

mutant of type α is $n_\alpha^{p=0} = exp(-\gamma_c \cdot s)$, while the number of matings for type β is $n_\beta^{p=0} =$

1. We define the cost of caring, $C = n_\beta^{p=0} - n_\alpha^{p=0} = 1 - exp(-\gamma_c \cdot s)$, as the loss of mating

success for a rare caring mutant of type α, where γ_c is a factor governing the cost of caring.

In the genetic case, the cost-benefit isocline is given by Eq. 1. The range of conditions where

a gene for paternal care evolves is shown by the grey area (Fig. 2). The conditions where a

microbe inducing paternal care evolves can be much wider, shown by the areas below the

green lines. The range widens with T_c, the probability of microbe transmission through

paternal care and narrows with T_m, the transmission probability during mating (Fig. 2). Even

when transmission through mating is larger than transmission through care, microbe-induced

paternal care fares better than paternal care driven by a host gene (see Supplementary, Figure

S1). This is in large part due to the maternal transmission of microbes to the offspring, the

probability of which is assumed to be stronger than that of maternal genes ($T = 0.5$).

However, reduced maternal transmission also allows microbe-induced paternal care to evolve

quite easily (see Supplementary Figure S2). Counter intuitively, the results also demonstrate

that microbial genes inducing paternal care behaviour can evolve even in the paradoxical case

where paternal "care" decreases offspring fitness (see Supplementary, Figure S3).

Overall, if the ratio of transmission probability through paternal care to the probability

through mating is sufficiently large, microbe-induced paternal care widens the range of

conditions that allows for the evolution of paternal care behaviour. This is more significant

when paternal care carries a substantial cost to the benefactor in terms of mating success or

does not provide enough benefit to the beneficiary.

Model II: Mixed brood

Now we consider a different social structure, where both males and females can engage in

extra-pair mating, but offspring are brought up by social pairs[21,54–56]. Females mate with one

extra-pair male besides their social mate, and a fraction P_e of their brood are sired by the

extra-pair mate. This factor is affected by mate guarding, sperm competition, and cryptic

female choice[57]. We assume that male mating success as an extra-pair sire reduces with

paternal care of that male[11]. The more the father invests in its offspring, the fitter they will be,

but the father may have fewer extra-pair progeny. The fitness of an offspring cared for by its

social father is increased by a factor of $1 + s$. The fitness of an extra-pair offspring is

increased by a factor of $1 + b$, due to direct or indirect benefits gained from extra-pair

mating[58,59]. Note that this extra-pair offspring can still benefit from the care of its social

father.

Figure 3. Model II (mixed brood): Illustration of microbe transmission pathways within families with extra-pair mating. Males carrying microbes of type β do not care for the offspring, while males carrying microbes of type α care for the offspring in their social nest. All males and females engage in extra-pair mating. T_v – vertical transmission probability through maternal influence (prenatal and postnatal). T_c – probability of transmission through paternal care. T_m – probability of male-to-female transmission during mating. T_n – probability of transmission through male-to-female nurture. Offspring sired by an extra-pair mate are $1 + b$ times more fit than offspring sired by the social mate. Offspring that receive paternal care are $1 + s$ times more fit than offspring that do not receive it.

Let ω_{xyz} be the fitness of an offspring with a social father of type x, a mother of type y, and a

biological father of type z (denoted by ω_{xy} if the social father x is also the biological father).

From our assumptions $\omega_{\alpha\beta\alpha} = \omega_{\alpha\beta\beta} = \omega_{\alpha\alpha\alpha} = \omega_{\alpha\alpha\beta} = (1 + b) * (1 + s)$, $\omega_{\beta\alpha\alpha} = \omega_{\beta\alpha\beta} =$

$\omega_{\beta\beta\alpha} = \omega_{\beta\beta\beta} = (1 + b)$, while $\omega_{\alpha\alpha} = \omega_{\alpha\beta} = (1 + s)$, and $\omega_{\beta\alpha} = \omega_{\beta\beta} = 1$.

We denote the expected number of extra-pair matings for a caring and non-caring male as

n_α, n_β respectively. We define $n_\alpha = \exp\left(-\gamma_c * s * (1 - p)\right)$, where γ_c is a factor governing

the cost of caring, s is the increase in offspring fitness from paternal care, p is the population

frequency of α, the microbe associated with paternal care. Thus, when a rare mutant microbe

inducing paternal care emerges in a population of males carrying microbes of type β, it will

obtain $\exp(-\gamma_c * s)$ matings. On the other hand, if the population is comprised only of caring

males, they will obtain, on average, one extra-pair mating opportunity.

We find the conditions for evolution of an α host gene, coding for paternal care, and similarly

for the evolution of microbes of type α, inducing host paternal care (see Supplementary for

mathematical derivations).

Figure 4. Model II (mixed brood): The evolution of paternal care in face of extra-pair offspring in brood. The figure represents the maximal fraction of extra-pair offspring in brood, P_e, that allows for the evolution of paternal care induced by either genes or microbes. The solid lines represent the genetic case and the dashed and dotted lines represents the microbial cases. Generally, in the microbial case, paternal care evolves under wider conditions. The range narrows with an increase in extra-pair paternity in both the genetic and microbial cases. However, the effect is reduced when the transmission probability through paternal care (T_c) is high. The different plots (i),(ii),(iii) represent different values of transmission through mating (T_m). In both the microbial and the genetic cases, as s increases, this allows for paternal care under higher degrees of extra-pair paternity. Other parameters: $b = 0.5$, $T_n = 0$, $T_v = 0.8$, $C = 0.9$, (i) $T_m = 0.01$, (ii) $T_m = 0.1$, (iii) $T_m = 0.2$.

Fig. 4 shows the maximal fraction of extra-pair offspring in brood, P_e, that allows for a gene

or a microbe of type α (paternal care) to evolve. A high degree of extra-pair paternity in the

population has a dual effect in the same direction. First, it allows for more opportunities to

breed as an extra-pair sire. Males of type β have greater mating success as extra-pair sires

than males of type α, thus could potentially gain more from the benefits to offspring

stemming from extra-pair mating. We examine the effect of extra-pair mating benefits (b) in

Supplementary Figure S4. Second, it reduces the genetic relatedness of the social father to the

offspring in its nest, and thus the fitness benefits it receives from paternal care. This effect is

stronger in the genetic case, since from microbial perspective, paternal care for a genetically

unrelated young individual contributes the same fitness benefits as for a genetically related

one.

The increase in offspring fitness due to paternal care (s) affects the evolution of paternal care

in an intricate manner, as both the benefits and the costs are increased with paternal

investment. When paternal contribution is sufficiently high, microbe-induced paternal care

can evolve even when paternal care is subject to substantial costs (see Supplementary, Figure

S5 for an example of diminished costs, where evolution of microbe-induced paternal care is

unlimited by paternity or paternal contribution).

Figure 5. Model 2 (mixed brood): The evolution of paternal care in face of extra-pair offspring in brood, asymmetric transmission. Here, microbes of type α have a lower transmission probability than microbes of type β in mating interactions and through maternal transmission. When the fitness benefit of paternal care (s) or transmission probability through care (T_c) are sufficiently high, microbe-induced paternal care allows for a wider range of costs than when driven by host genes. Otherwise, the substantially diminished success of microbes of type α via extra-pair behaviour hinders their evolution. The different plots represent different values of transmission through mating (T_m). Other parameters: $b = 0.5$, $T_n = 0$, $T_v = 0.8$, $C = 0.9$, (i) $T_m = 0.01$, (ii) $T_m = 0.1$, (iii) $T_m = 0.2$. $T_\alpha = 0.9 \cdot T_\beta$.

The dynamics between the two microbe types (α and β) are strongly affected by the ratio

between transmission probability through paternal care (T_c) and transmission probability

through mating (T_m). Generally, a higher T_c allows for a wider range of conditions in which

microbe-induced paternal care can evolve (see Supplementary Figure S6 for extreme values

of T_m). Fig. 5 demonstrates the asymmetric contagiousness case, when microbes of type α

have a lower transmission probability than microbes of type β in mating interactions ($T_m^{\alpha} <$

T_m^{β}) and through maternal transmission ($T_v^{\alpha} < T_v^{\beta}$). In this case, when considering the extra-

pair behaviour, microbes of type α have diminished success both due to the males' reduced

mating opportunities and due to the lower transmission probability through mating. Hence,

when the fraction of extra-pair offspring in the brood (P_e) is high, the disparity in fitness

increases in favour of microbe of type β. To counteract these costs and allow the care-

inducing microbe to evolve, paternal care must provide a significant benefit. When the fitness

benefit of paternal care (s) or transmission probability through care (T_c) are sufficiently high,

microbe-induced paternal care allows for a wider range of costs than when driven by host

genes. We examined microbe-induced paternal care with transmission disadvantage for

Model 1 as well (see Supplementary, Figure S7).

In conclusion, microbe-induced paternal care allows for evolution of paternal care behaviour

even in face of loss of paternity due to extra-pair mating, and when paternal investment

results in significantly reduced mating success.

Discussion

In this work, we present an alternative explanation to a long-standing evolutionary

conundrum: the ubiquity of paternal care despite its costs, such as a possible reduction in a

father's fitness due to expenditure of time and resources into offspring care, and the risk of

caring for unrelated young due to uncertainty in paternity. Our findings suggest that paternal

care evolves more easily when induced by the microbes' genes than by the host genes. Our

model predicts that microbe-induced paternal care would more easily evolve when parent-

offspring interactions lead to a high probability of microbial transmission[50,60–63] (high T_c,

probability of paternal transmission of microbes via care, resulting from interactions

corresponding to feeding, grooming). We demonstrate that microbe-induced paternal care can

explain stable levels of paternal care even under high levels of extra-pair mating, when cost

of caring is high, and even in some cases when the microbe inducing paternal care has a

transmission disadvantage.

Previous work has discussed possible explanations for the broad existence of paternal care.

Suggested explanations[64] include gaining practice in caring for young, resulting in higher

chances for successful parenthood in the following year[13]; increasing indirect fitness[65,66]; and

increasing chances of mating or territory acquisition[67,68]. Additionally, since female choice is

a key factor in determining male reproductive success, sexual selection may act directly to

favour paternal care[9,66,69]. It has been suggested that males overestimate the likelihood of

paternity[70,71], which helps preserve a stable level of paternal care[14,17,72]. However, when the

cost of caring is high, and the expected level of paternity is low, selection is expected to

favour more suspicious males that reduce their paternal investment with increased risk of

extra-pair mating[8,17,72–75]. Nevertheless, paternal care has been demonstrated to prevail even

in these cases in natural systems[21,22,55,76]. We expect that microbe-induced paternal care

could play a significant role in circumstances where genetic relatedness falls short of

explaining the observed degree of paternal investment, such as adoption[77], species where

extra-pair paternity is common[21] and the well-known behaviour of cooperative breeding and

eusociality[64,78].

Microbial evolutionary interests may explain stable levels of extra-pair mating. The benefits

of extra-pair mating for females[56,58,79,80] may be obtaining a higher quality or more

compatible sire[79] and bet-hedging by increasing the genetic diversity of offspring[81]. Extra-

pair mating can also result in significant costs to the female. The possible costs include loss

of care by social mate[82], male sexual aggression[83], increased sibling competition[84,85], and the

risk of contracting sexually transmitted pathogens[86]. As demonstrated by our results,

microbe-induced care by the social mate prevails under a wider range of paternity levels in

comparison to care driven by host genes. Within-brood aggression between half-siblings[84]

could also be mitigated by microbes, since relatedness among the microbes of the sibling is

expected to be significant even if their genetic relatedness is not high[87,88].

Our model can be extended in several ways. We examined two extremes, paternal care

governed exclusively by host genes or exclusively by microbial genes. However, evolution of

paternal care is likely driven by selection on reproductive units in both levels, possibly

leading to intermediate results. Additionally, it is possible to consider that when host genes

and microbial genes experience conflicting selective pressures, selection on the host would

drive the evolution of resistance genes to the microbial influence. In this case, we expect the

host-microbe coevolution to generate oscillatory rock-paper-scissors evolutionary dynamics,

that can allow the long-term maintenance of paternal care. Similar dynamics have been found

by some of us with respect to microbe-induced cooperation and host resistance[89]. Another

extension would be allowing more female strategies. We assumed a constant level of

maternal care. Yet, studies show females may reduce their care if the male provides

sufficiently intensive care or increase their care to compensate for lack of male care[15,90,91].

This behaviour lowers the return on paternal investment in terms of benefits to offspring

fitness, effectively increasing the cost/benefit ratio. We may consider maternal care to be

driven by microbial genes as well. We predict that in many cases microbial evolutionary

interests would be to induce high degrees of maternal investment, as the offspring is very

likely to carry microbes of the same type as his mother, especially in cases where paternal

involvement is meagre or lacking.

Our model joins the rank of previous models concerning the role of different nongenetic

elements in the evolution of social traits[29,92–94]. Recent evidence suggests that microbes hold a

significant role in shaping host evolution[24,29,95,96]. However, it is worth noting that the

assumptions presented here are not limited to the microbiome and apply to any class of

nongenetic elements that are capable of both vertical and horizontal/oblique[97,98] transfer and

of influencing complex behavioural phenotypes. Examples of such elements may include

epigenetic states[99,100] and culture[92,101–104]. The general outline of our model can also be

applied to genetic relatives with varying degrees of relatedness and to alloparenting.

Our theoretical results demonstrate a possible explanation, involving microbial regulation, to

a widely studied and unresolved question in evolutionary biology – why males care for the

offspring despite significant costs. Previous studies demonstrated hormonal regulation of

paternal care[105–108], yet the mechanism by which microbes may regulate paternal behaviour is

still to be experimentally validated. Our results call for empirical testing of our predictions:

that microbes are involved in the regulation of paternal behaviour, and that factors that affect

the composition of host microbiome dramatically (e.g., antibiotics[109,110]) may also alter

paternal behaviour.

References

1. Cockburn, A. Prevalence of different modes of parental care in birds. *Proc. R. Soc. B Biol. Sci.* (2006). doi:10.1098/rspb.2005.3458
2. Kleiman, D. . & Malcolm, J. . The Evolution of Male Parental Investment in Mammals. in *Parental Care in Mammals* (1970). doi:10.1007/978-1-4613-3150-6
3. Brown, J. L., Morales, V. & Summers, K. A Key Ecological Trait Drove the Evolution of Biparental Care and Monogamy in an Amphibian. *Am. Nat.* (2010). doi:10.1086/650727
4. Ridley, M. Paternal care. *Anim. Behav.* **26**, 904–932 (1978).
5. Clutton-Brock, T. H. *The evolution of parental care.* (Princeton University Press, 1991).
6. Shellman-Reeve, J. Dynamics of biparental care in the dampwood termite, Zootermopsis nevadensis (Hagen): response to nitrogen availability. *Behav. Ecol. Sociobiol.* **26**, 389–397 (1990).
7. Freeman, H. Behavior in adult pairs of captive snow leopards (Panthera uncia). *Zoo Biol.* **2**, 1–22 (1983).
8. Hamilton, W. D. The genetical evolution of social behaviour. II. *J. Theor. Biol.* **7**, 17–52 (1964).
9. Trivers, R. L. Parental investment and sexual selection. in *Sexual Selection & the Descent of Man* 136–179 (1972). doi:10.1002/ajpa.1330400226
10. Kölliker, M. *The Evolution of Parental Care*. (Oxford University Press, 2012). doi:10.1093/acprof:oso/9780199692576.001.0001
11. Magrath, M. J. L. & Komdeur, J. Is male care compromised by additional mating opportunity? *Trends Ecol. Evol.* **18**, 424–430 (2003).
12. Thomas, L. K. & Manica, A. Intrasexual competition and mate choice in assassin bugs with uniparental male and female care. *Anim. Behav.* **69**, 275–281 (2005).
13. Rosenbaum, S., Vigilant, L., Kuzawa, C. W. & Stoinski, T. S. Caring for infants is associated with increased reproductive success for male mountain gorillas. *Sci. Rep.* **8**, 15223 (2018).
14. Westneat, D. F. & Sherman, P. W. Parentage and the evolution of paternal care. *Behav. Ecol.* **4**, 66–77 (1993).
15. Kokko, H. Cuckoldry and the stability of biparental care. *Ecol. Lett.* **2**, 247–255 (1999).
16. Sheldon, B. C. Relating paternity to paternal care. *Philos. Trans. R. Soc. B Biol. Sci.* **357**, 341–350 (2002).
17. Griffin, A. S., Alonzo, S. H. & Cornwallis, C. K. Why Do Cuckolded Males Provide Paternal Care? *PLoS Biol.* **11**, (2013).
18. Richardson, P. R. K. & Coetzee, M. Mate desertion in response to female promiscuity in the socially monogamous aardwolf Proteles cristatus. *South African J. Zool.* **23**, 306–308 (1988).
19. Westneat, D. F. & Sargent, R. C. Sex and parenting: The effects of sexual conflict and parentage on parental strategies. *Trends in Ecology and Evolution* (1996). doi:10.1016/0169-5347(96)81049-4
20. Alonzo, S. H. & Heckman, K. L. The unexpected but understandable dynamics of mating, paternity and paternal care in the ocellated wrasse. *Proc. R. Soc. B Biol. Sci.* **277**, 115–122 (2010).
21. Petrie, M. & Kempenaers, B. Extra-pair paternity in birds: Explaining variation between

species and populations. *Trends Ecol. Evol.* **13**, 52–57 (1998).

22. Mulder, R. A., Dunn, P. O., Cockburn, A., Lazenby-Cohen, K. A. & Howell, M. J. Helpers liberate female fairy-wrens from constraints on extra-pair mate choice. *Proc. R. Soc. B Biol. Sci.* **255**, 223–229 (1994).

23. Cryan, J. F. & O'Mahony, S. M. The microbiome-gut-brain axis: From bowel to behavior. *Neurogastroenterol. Motil.* (2011). doi:10.1111/j.1365-2982.2010.01664.x

24. Ezenwa, V. O., Gerardo, N. M., Inouye, D. W., Medina, M. & Xavier, J. B. Animal behavior and the microbiome. *Science* (2012). doi:10.1126/science.1227412

25. Sampson, T. R. & Mazmanian, S. K. Control of brain development, function, and behavior by the microbiome. *Cell Host and Microbe* (2015). doi:10.1016/j.chom.2015.04.011

26. Lyte, M. Microbial Endocrinology in the Microbiome-Gut-Brain Axis: How Bacterial Production and Utilization of Neurochemicals Influence Behavior. *PLoS Pathog.* (2013). doi:10.1371/journal.ppat.1003726

27. Neufeld, K. M., Kang, N., Bienenstock, J. & Foster, J. A. Reduced anxiety-like behavior and central neurochemical change in germ-free mice. *Neurogastroenterol. Motil.* (2011). doi:10.1111/j.1365-2982.2010.01620.x

28. Johnson, K. V.-A. & Foster, K. R. Why does the microbiome affect behaviour? *Nat. Rev. Microbiol.* **16**, 647–655 (2018).

29. Lewin-Epstein, O., Aharonov, R. & Hadany, L. Microbes can help explain the evolution of host altruism. *Nat. Commun.* **8**, (2017).

30. Mulle, J. G., Sharp, W. G. & Cubells, J. F. The Gut Microbiome: A New Frontier in Autism Research. *Curr. Psychiatry Rep.* **15**, 337 (2013).

31. Sgritta, M. *et al.* Mechanisms Underlying Microbial-Mediated Changes in Social Behavior in Mouse Models of Autism Spectrum Disorder. *Neuron* **101**, 246-259.e6 (2019).

32. Foster, J. A. & McVey Neufeld, K.-A. Gut–brain axis: how the microbiome influences anxiety and depression. *Trends Neurosci.* **36**, 305–312 (2013).

33. Davis, D. J. *et al.* Lactobacillus plantarum attenuates anxiety-related behavior and protects against stress-induced dysbiosis in adult zebrafish. *Sci. Rep.* **6**, 33726 (2016).

34. Desbonnet, L. *et al.* Effects of the probiotic Bifidobacterium infantis in the maternal separation model of depression. *Neuroscience* **170**, 1179–1188 (2010).

35. Bharwani, A., Mian, M. F., Surette, M. G., Bienenstock, J. & Forsythe, P. Oral treatment with Lactobacillus rhamnosus attenuates behavioural deficits and immune changes in chronic social stress. *BMC Med.* **15**, 7 (2017).

36. Mändar, R. & Mikelsaar, M. Transmission of mother's microflora to the newborn at birth. *Neonatology* (1996). doi:10.1159/000244275

37. Funkhouser, L. J. & Bordenstein, S. R. Mom Knows Best: The Universality of Maternal Microbial Transmission. *PLoS Biol.* (2013). doi:10.1371/journal.pbio.1001631

38. Banning, J. L. *et al.* Antifungal skin bacteria, embryonic survival, and communal nesting in four-toed salamanders, Hemidactylium scutatum. *Oecologia* (2008). doi:10.1007/s00442-008-1002-5

39. Gunderson, A. R., Forsyth, M. H. & Swaddle, J. P. Evidence that plumage bacteria influence feather coloration and body condition of eastern bluebirds Sialia sialis. *J. Avian Biol.* (2009). doi:10.1111/j.1600-048X.2008.04650.x

40. Kulkarni, S. & Heeb, P. Social and sexual behaviours aid transmission of bacteria in birds.

Behav. Processes (2007). doi:10.1016/j.beproc.2006.10.005

41. Koch, H. & Schmid-Hempel, P. Socially transmitted gut microbiota protect bumble bees against an intestinal parasite. *Proc. Natl. Acad. Sci.* (2011). doi:10.1073/pnas.1110474108

42. Stilling, R. M., Bordenstein, S. R., Dinan, T. G. & Cryan, J. F. Friends with social benefits: host-microbe interactions as a driver of brain evolution and development? *Front. Cell. Infect. Microbiol.* **4**, 147 (2014).

43. Rothschild, D. *et al.* Environment dominates over host genetics in shaping human gut microbiota. *Nature* (2018). doi:10.1038/nature25973

44. Bravo, J. A. *et al.* Ingestion of Lactobacillus strain regulates emotional behavior and central GABA receptor expression in a mouse via the vagus nerve. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 16050–16055 (2011).

45. Schmidt, C. Mental health: Thinking from the gut. *Nature* **518**, S12–S15 (2015).

46. Stiver, K. A. & Alonzo, S. H. Parental and Mating Effort: Is There Necessarily a Trade-Off? *Ethology* **115**, 1101–1126 (2009).

47. KOKKO, H. & JENNIONS, M. D. Parental investment, sexual selection and sex ratios. *J. Evol. Biol.* **21**, 919–948 (2008).

48. Ahnesjö, I., Kvarnemo, C. & Merilaita, S. *Using potential reproductive rates to predict mating competition among individuals qualified to mate*. *Behavioral Ecology* **12**,

49. Jennions, M. D. & Fromhage, L. Not all sex ratios are equal: The Fisher condition, parental care and sexual selection. *Philosophical Transactions of the Royal Society B: Biological Sciences* **372**, (2017).

50. Tung, J. *et al.* Social networks predict gut microbiome composition in wild baboons. *Elife* **4**, (2015).

51. Smiseth, P. T. & Moore, A. J. Behavioral dynamics between caring males and females in a beetle with facultative biparental care. *Behav. Ecol.* (2004). doi:10.1093/beheco/arh053

52. Hunt, J. & Simmons, L. W. Behavioural dynamics of biparental care in the dung beetle Onthophagus taurus. *Anim. Behav.* (2002). doi:10.1006/anbe.2002.3036

53. White, J. *et al.* Sexually transmitted bacteria affect female cloacal assemblages in a wild bird. *Ecol. Lett.* (2010). doi:10.1111/j.1461-0248.2010.01542.x

54. Westneat, D. F., Sherman, P. W. & Morton, M. L. The ecology and evolution of extra-pair copulations in birds. *Curr. Ornithol.* **7**, 331–369 (1990).

55. Griffith, S. C., Owens, I. P. F. & Thuman, K. A. Extra pair paternity in birds: a review of interspecific variation and adaptive function. *Mol. Ecol.* **11**, 2195–2212 (2002).

56. Westneat, D. F. & Stewart, I. R. K. Extra-Pair Paternity in Birds: Causes, Correlates and Conflict. *Annu. Rev. Ecol. Evol. Syst.* **34**, 365–396 (2003).

57. Eberhard, W. G. *Female control : sexual selection by cryptic female choice*. (Princeton University Press, 1996).

58. Forstmeier, W., Nakagawa, S., Griffith, S. C. & Kempenaers, B. Female extra-pair mating: Adaptation or genetic constraint? *Trends Ecol. Evol.* **29**, 456–464 (2014).

59. JENNIONS, M. D. & PETRIE, M. Why do females mate multiply? A review of the genetic benefits. *Biol. Rev. Camb. Philos. Soc.* **75**, 21–64 (2000).

60. Ezenwa, V. O., Ghai, R. R., McKay, A. F. & Williams, A. E. Group living and pathogen infection revisited. *Curr. Opin. Behav. Sci.* **12**, 66–72 (2016).

61. Moeller, A. H. *et al.* Social behavior shapes the chimpanzee pan-microbiome. *Sci. Adv.* **2**, e1500997 (2016).

62. Song, S. J. *et al.* Cohabiting family members share microbiota with one another and with

their dogs. *Elife* **2**, 458 (2013).

63. VanderWaal, K. L., Atwill, E. R., Isbell, L. A. & McCowan, B. Linking social and pathogen transmission networks using microbial genetics in giraffe (Giraffa camelopardalis). *J. Anim. Ecol.* **83**, 406–414 (2014).

64. Cockburn, A. Evolution of helping behavior in cooperatively breeding birds. *Annu. Rev. Ecol. Syst.* **29**, 141–177 (1998).

65. Bart, J. & Tornes, A. Importance of monogamous male birds in determining reproductive success. *Behav. Ecol. Sociobiol.* **24**, 109–116 (1989).

66. Moller, A. P. Male parental care, female reproductive success, and extrapair paternity. *Behav. Ecol.* **11**, 161–168 (2000).

67. Marlowe, F. Male care and mating effort among Hadza foragers. *Behav. Ecol. Sociobiol.* **46**, 57–64 (1999).

68. Kerhoas, D. *et al.* Mother-male bond, but not paternity, influences male-infant affiliation in wild crested macaques. *Behav. Ecol. Sociobiol.* **70**, 1117–1130 (2016).

69. Kokko, H., Brooks, R., Jennions, M. D. & Morley, J. The evolution of mate choice and mating biases. *Proc. R. Soc. B Biol. Sci.* **270**, 653–664 (2003).

70. Neff, B. D. Decisions about parental care in response to perceived paternity. *Nature* **422**, 716–719 (2003).

71. Lissåker, M. & Svensson, O. Cannibalize or care? The role of perceived paternity in the sand goby, Pomatoschistus minutus. *Behav. Ecol. Sociobiol.* **62**, 1467–1475 (2008).

72. Kokko, H. Should advertising parental care be honest? *Proc. R. Soc. London. Ser. B Biol. Sci.* **265**, 1871–1878 (1998).

73. Manica, A. Parental fish change their cannibalistic behaviour in response to the cost-to-benefit ratio of parental care. *Anim. Behav.* **67**, 1015–1021 (2004).

74. Houston, A. I. & McNamara, J. M. A self-consistent approach to paternity and parental effort. *Philos. Trans. R. Soc. B Biol. Sci.* **357**, 351–362 (2002).

75. Houston, A. I. Parental effort and paternity. *Anim. Behav.* **50**, 1635–1644 (1995).

76. Bouwman, K. M., Lessells, C. M. & Komdeur, J. Male reed buntings do not adjust parental effort in relation to extrapair paternity. *Behav. Ecol.* **16**, 499–506 (2005).

77. Riedman, M. L. The Evolution of Alloparental Care and Adoption in Mammals and Birds. *Q. Rev. Biol.* **57**, 405–435 (1982).

78. Clutton-Brock, T. Breeding together: kin selection and mutualism in cooperative vertebrates. *Science* **296**, 69–72 (2002).

79. Foerster, K., Delhey, K., Johnsen, A., Lifjeld, J. T. & Kempenaers, B. Females increase offspring heterozygosity and fitness through extra-pair matings. *Nature* **425**, 714–717 (2003).

80. Kempenaers, B. & Dhondt, A. A. Why do females engage in extra-pair copulations? A review of hypotheses and their predictions. *Belg. J Zool.* **123**, 93–103 (1993).

81. Fox, C. W. & Rauter, C. M. *Bet-hedging and the evolution of multiple mating*. (2003).

82. Arnqvist, G. & Kirkpatrick, M. The evolution of infidelity in socially monogamous passerines: the strength of direct and indirect selection on extrapair copulation behavior in females. *Am. Nat.* **165 Suppl 5**, S26-37 (2005).

83. Valera, F., Hoi, H. & Krištín, A. Male shrikes punish unfaithful females. *Behav. Ecol.* **14**, 403–408 (2003).

84. Briskie, J. V., Naugler, C. T. & Leech, S. M. Begging intensity of nestling birds varies with sibling relatedness. *Proc. R. Soc. B Biol. Sci.* **258**, 73–78 (1994).

85.   Brante, A., Fernández, M. & Viard, F. Non-Random Sibling Cannibalism in the Marine Gastropod Crepidula coquimbensis. *PLoS One* **8**, e67050 (2013).

86.   Sheldon, B. C. Sexually transmitted disease in birds: occurrence and evolutionary significance. *Philos. Trans. R. Soc. London. Ser. B Biol. Sci.* **339**, 491–497 (1993).

87.   Mosites, E. *et al.* Microbiome sharing between children, livestock and household surfaces in western Kenya. *PLoS One* **12**, e0171017 (2017).

88.   Brito, I. L. *et al.* Transmission of human-associated microbiota along family and social networks. *Nature Microbiology* **4**, 964–971 (2019).

89.   Lewin-Epstein, O. & Hadany, L. Host-microbiome coevolution promotes cooperation in a rock-paper-scissor dynamic. *bioRxiv* 689299 (2019). doi:10.1101/689299

90.   Houston, A. I., Székely, T. & McNamara, J. M. Conflict between parents over care. *Trends Ecol. Evol.* **20**, 33–38 (2005).

91.   Harrison, F. *et al.* How is sexual conflict over parental care resolved? A meta-analysis. *J. Evol. Biol.* **22**, 1800–1812 (2009).

92.   Feldman, M. W., Cavalli-Sforza, L. L. & Peck, J. R. Gene-culture coevolution: models for the evolution of altruism with cultural transmission. *Proc. Natl. Acad. Sci. U. S. A.* **82**, 5814–8 (1985).

93.   Bonduriansky, R. & Day, T. *Extended Heredity: A New Understanding of Inheritance and Evolution*. (Princeton University Press, 2018).

94.   Bonduriansky, R. & Day, T. Nongenetic inheritance and the evolution of costly female preference. *J. Evol. Biol.* **26**, 76–87 (2013).

95.   Morimoto, J., Simpson, S. J. & Ponton, F. Direct and trans-generational effects of male and female gut microbiota in Drosophila melanogaster. *Biol. Lett.* **13**, (2017).

96.   Vuong, H. E., Yano, J. M., Fung, T. C. & Hsiao, E. Y. The Microbiome and Host Behavior. *Annu. Rev. Neurosci.* **40**, 21–49 (2017).

97.   Ram, Y., Liberman, U. & Feldman, M. W. Evolution of vertical and oblique transmission under fluctuating selection. *Proc. Natl. Acad. Sci.* **115**, E1174–E1183 (2018).

98.   Ram, Y., Liberman, U. & Feldman, M. W. Vertical and oblique cultural transmission fluctuating in time and in space. *Theor. Popul. Biol.* **125**, 11–19 (2019).

99.   Danchin, É. *et al.* Beyond DNA: integrating inclusive inheritance into an extended theory of evolution. *Nat. Rev. Genet.* **12**, 475–486 (2011).

100.   Jablonka, E., Lamb, M. J. & Zeligowski, A. *Evolution in four dimensions, revised edition*. *MIT press* (2014).

101.   Bartsch, C., Weiss, M. & Kipper, S. Multiple song features are related to paternal effort in common nightingales. *BMC Evol. Biol.* **15**, 115 (2015).

102.   Murray, C. M., Gilby, I. C., Mane, S. V. & Pusey, A. E. Adult Male Chimpanzees Inherit Maternal Ranging Patterns. *Curr. Biol.* **18**, 20–24 (2008).

103.   Cavalli-Sforza, L. L. & Feldman, M. *Cultural Transmission and Evolution: A Quantitative Approach*. (Princeton University Press, 1981).

104.   Rushton, J. P., Littlefield, C. H. & Lumsden, C. J. Gene-culture coevolution of complex social behavior: human altruism and mate choice. *Proc. Natl. Acad. Sci. U. S. A.* **83**, 7340–3 (1986).

105.   Pradhan, D. S., Solomon-Lane, T. K., Willis, M. C. & Grober, M. S. A mechanism for rapid neurosteroidal regulation of parenting behaviour. *Proc. R. Soc. B Biol. Sci.* **281**, 20140239–20140239 (2014).

106.   O'Connell, L. A., Matthews, B. J. & Hofmann, H. A. Isotocin regulates paternal care in a

monogamous cichlid fish. *Horm. Behav.* **61**, 725–733 (2012).

107. Martínez, A. *et al.* Paternal behavior in the Mongolian gerbil (Meriones unguiculatus): Estrogenic and androgenic regulation. *Horm. Behav.* **71**, 91–95 (2015).

108. Mota, M. T. da S., Franci, C. R. & de Sousa, M. B. C. Hormonal changes related to paternal and alloparental care in common marmosets (Callithrix jacchus). *Horm. Behav.* **49**, 293–302 (2006).

109. Jakobsson, H. E. *et al.* Short-Term Antibiotic Treatment Has Differing Long-Term Impacts on the Human Throat and Gut Microbiome. *PLoS One* **5**, e9836 (2010).

110. Manichanh, C. *et al.* Reshaping the gut microbiome with bacterial transplantation and antibiotic intake. *Genome Res.* **20**, 1411–9 (2010).

Acknowledgements and funding

This project was supported by the Israeli Science Foundation 2064/18 (L.H.) and by the Minerva Center on Lab Evolution (L.H.).

Author Contributions

Y.G. and L.H. designed the study and formulated the model. Y.G. and O.L.-E. derived the analytical equations and implemented the code. Y.G. and L.H. analysed the results and wrote the manuscript.

Competing interests

The authors declare no competing interests.