Title: Methyl-Metabolite Depletion Elicits Adaptive Responses to Support Heterochromatin

Stability and Epigenetic Persistence

Spencer A. Haws[1,2], Deyang Yu[3,4,5], Cunqi Ye[6], Coral K. Wille[1,6], Long C. Nguyen[8,9], Kimberly A.

Krautkramer[1,2], Jay L. Tomasiewicz[3], Shany E. Yang[3,4], Blake R. Miller[3,4], Wallace H. Liu[1,2],

Kazuhiko Igarashi[8,9], Rupa Sridharan[1,6], Benjamin P. Tu[6], Vincent L. Cryns[3,5,10], Dudley W.

Lamming[3,4,5], and John M. Denu[1,2]*.

[1]Department of Biomolecular Chemistry, SMPH, University of Wisconsin, Madison, WI, 53706,

USA

[2]Wisconsin Institute for Discovery, University of Wisconsin, Madison, WI, 53715, USA

[3]Department of Medicine, SMPH, University of Wisconsin, Madison, WI, 53705, USA

[4]William S. Middleton Memorial Veterans Hospital, Madison, WI, 53705, USA

[5]Molecular & Environmental Toxicology Center, SMPH, University of Wisconsin, Madison, WI,

53705, USA

[6]Department of Biochemistry, University of Texas Southwestern Medical Center, Dallas, TX,

75390, USA

[7]Department of Cell and Regenerative Biology, University of Wisconsin, Madison, WI, 53705,

USA

[8]Department of Biochemistry, Tohoku University Graduate School of Medicine, Sendai, 980-

8575, JPN

[9]Center for Regulatory Epigenome and Diseases, Tohoku University Graduate School of

Medicine, Sendai, 980-8575, JPN

[10]University of Wisconsin Carbone Cancer Center, Madison, WI, 53792, USA

*Correspondence: John M. Denu, john.denu@wisc.edu

Summary

S-adenosylmethionine (SAM) is the methyl-donor substrate for DNA and histone

methyltransferases that regulate cellular epigenetic states. This metabolism-epigenome link

enables the sensitization of chromatin methylation to altered SAM abundance. However, a

chromatin-wide understanding of the adaptive/responsive mechanisms that allow cells to

actively protect epigenetic information during life-experienced fluctuations in SAM availability

are unknown. We identified a robust response to SAM depletion that is highlighted by

preferential cytoplasmic and nuclear *de novo* mono-methylation of H3 Lys 9 (H3K9) at the

expense of global losses in histone di- and tri-methylation. Under SAM-depleted conditions, *de*

novo H3K9 mono-methylation preserves heterochromatin stability and supports global

epigenetic persistence upon metabolic recovery. This unique chromatin response was robust

across the mouse lifespan and correlated with improved metabolic health, supporting a

significant role for epigenetic adaptation to SAM depletion *in vivo*. Together, these studies

provide the first evidence for active epigenetic adaptation and persistence to metabolic stress.

Keywords: metabolism, methionine, SAM, epigenetics, methylation, chromatin, histone, aging,

persistence

Introduction

Metabolism and the epigenome are connected by central metabolites that act as co-substrates for chromatin modifying enzymes, enabling fluctuations in metabolite availability to directly tune a cell's ability to 'write' and 'erase' chromatin post-translational modifications (PTMs) (Fan et al., 2015; Li et al., 2018). Chromatin methylation has been shown to be particularly susceptible to metabolic perturbations as methyltransferases require a single methyl-donor cofactor, S-adenosylmethionine (SAM) (Ducker and Rabinowitz, 2016; Mentch and Locasale, 2017). For example, accumulation of SAM in *S. cerevisiae* results in global increases in histone methylation levels (Ye et al., 2017). On the contrary, reduced SAM availability has been linked with site-specific losses in histone methylation as well as global depression of DNA methylation in several organisms (Shiraki et al., 2014; Ding et al., 2015; Kera et al., 2013; Tang et al., 2017; Shyh-Chang et al., 2013; Mentch et al., 2015; Tang et al., 2015; Hayashi et al., 2018; Towbin et al., 2012; Strekalova et al., 2019; Wang et al., 2019).

Intracellular SAM abundance is dependent on the catalysis of its obligatory precursor metabolite, the essential amino acid methionine (Met). Consumption of popular diets containing low quantities of Met (fish, vegetarian, and vegan) have been shown to significantly correlate with decreased plasma Met concentrations (Farmer, 2014; Schmidt et al., 2016). This link provides an opportunity for diet to directly influence intracellular SAM availability. Furthermore, abundance of Met and SAM have both been shown to fluctuate naturally in tune with circadian rhythms (Krishnaiah et al., 2017). Therefore, individuals consuming adequate amounts of dietary Met likely still experience periods of decreased methyl-metabolite availability. While both dietary intake and circadian regulation are capable of influencing Met and SAM abundance, adaptive mechanisms that allow cells to actively respond to – and recover their functional epigenomes from – such metabolic perturbations are unknown. Here, we define the ability of a cell and/or organism to re-establish the epigenome that was present prior to onset of an environmental perturbation or metabolic stress as *epigenetic persistence*.

Adaptive mechanisms that respond to methyl-metabolite depletion likely support critical cellular functions under these conditions. Interestingly, dietary Met-restriction is associated with beneficial metabolic reprogramming and lifespan extension in mammals (Orentreich et al., 1993; Yu et al., 2018; Miller et al., 2005; Brown-Borg et al., 2018; Lees et al., 2014; Green and Lamming, 2018). These phenotypes run counter to negative pathologies associated with dysregulated chromatin methylation (i.e., intellectual disability syndromes, cancers, and premature as well as natural aging) (Greer and Shi, 2012). This suggests cells possess

responsive mechanisms capable of actively maintaining regulation of site specific and/or global

chromatin methylation events during SAM depletion. The goal of this study is to identify and

characterize such mechanisms to better understand how organisms actively protect their

epigenome in response to fluctuations in central metabolite availability.

Here, we identify a robust, conserved chromatin response to metabolically-depleted

SAM levels. This cellular response is characterized by global decreases in higher-state histone

methylation (di- and tri-methyl) and simultaneous coordinated activation of *de novo* H3K9 mono-

methylation involving newly-synthesized and chromatin-bound histone H3. Under extreme

conditions of both reduced SAM levels and inhibited methylation that would yield H3K9me1,

heterochromatin instability and de-repression of constitutively silenced DNA elements is

exacerbated. Acute inhibition of *de novo* H3K9 mono-methylation also leads to the

dysregulation of global histone PTM states upon metabolic recovery, disrupting epigenetic

persistence to decreased SAM availability. Importantly, chronic metabolic depletion of SAM in

both young and old mice recapitulates these *in vitro* findings, revealing a conserved, age-

independent chromatin response. These results implicate *de novo* H3K9 mono-methylation as

an indispensable mechanism to support heterochromatin stability and global epigenetic

persistence in response to SAM depletion.

Results

<u>Methionine Restriction Stimulates Dynamic Histone PTM Response</u>

To comprehensively investigate how disruption of methyl-donor metabolism affects

chromatin methylation states, global histone proteomics and DNA methylation (5mC) analyses

were performed on two distinct *in vitro* and *in vivo* systems. These included Met-restricted

HCT116 human colorectal cancer cells and C57BL/6J mouse liver. Methionine metabolism was

severely repressed in both systems, highlighted by significant reductions in Met and SAM

abundance (Figure 1A and Figure S1A-S1D). Reduced methyl-metabolite availability did not

affect global DNA 5mC abundance in either system (Figure 1B-1C). However, LC-MS/MS

analysis of more than 40 unique histone H3 peptide proteoforms revealed dynamic PTM

responses to Met-restriction in both HCT116 cells and C57BL/6J liver (Figure 1D-1E). In

HCT116 cells, Met-restriction stimulated distinct biphasic changes in global PTM profiles (Figure

1D). Phase I (0 hr-45 min) was rapid and marked by a trending upregulation of H3K4me2/3,

PTMs known to mark transcriptionally active promoters. Phase II (90 min-24 hrs) was

characterized by global decreases in di- and tri- histone methylation. Decreased levels of di- and tri-methylated peptides were accompanied by increases in acetylated and unmodified peptide species. Similarly in C57BL/6J mice, histone PTM responses were marked by significant decreases in histone di- and tri-methylation that essentially matched the patterns found in HCT116 cells during the prolonged Phase II response (Figure 1E and Figure S1E-S1J). Decreased higher state (di- and tri-) histone methylation both *in vitro* and *in vivo* highlights a decreased methylation capacity resulting from prolonged Met and/or SAM depletion. Together, these observations suggest perturbed methyl-donor metabolism is capable of stimulating significant changes in histone, but not global 5mC, methylation abundance. Furthermore, similarities between the metabolic and epigenetic responses to Met-restriction across both systems support the use of *in vitro* Met-restriction as a model for mechanistic follow-up studies.

Decreased SAM Availability Drives Robust Histone Methylation Response

Dramatic reduction of intracellular Met and SAM correlated with onset of the *in vitro* Phase I and II histone PTM changes, respectively (Figure 1D and Figure S1A-S1B). This implies depletion of individual methyl-metabolites may be capable of stimulating distinct histone modifying pathways. To determine if the global losses in di- and tri- histone methylation *in vitro* are driven specifically by SAM depletion, two alternative approaches were employed to deplete cellular SAM independent of Met (Figure 2A). The first approach utilized RNAi knockdown of the mammalian SAM synthetase MAT2A (methionine adenosyltransferase II alpha). Knockdown via MAT2A-RNAi treatment decreased MAT2A transcript abundance by greater than 97%, resulting in a near complete deprivation of SAM availability with no effect on Met levels (Figure 2B-2D). Histone proteomics analysis of MAT2A-RNAi treated cells identified site-specific histone PTM changes that mirrored those identified in HCT116 cells under Phase II Met-restriction conditions (r=0.778, p=8.59e$_{-10}$) (Figure 2E). Using a complementary approach, SAM levels were depleted by overexpression of a major SAM consumer, PEMT (phosphatidylethanolamine N-methyltransferase). Ye et al. 2017 have previously shown PEMT overexpression significantly reduces SAM availability in HEK293T human embryonic kidney cells. Histone proteomics analysis of PEMT overexpressing HEK293T cells identified site-specific PTM changes that paralleled those of Met-restricted controls (r=0.803, p=3.30e$_{-8}$) (Figure 2F). Together, two separate means of reducing SAM levels demonstrate SAM reduction alone is sufficient to inhibit the methylation capacity of a cell, inducing dynamic changes in histone PTM abundance.

Furthermore, these findings support the use of prolonged Met-restriction as a model of SAM

depletion.

The similar changes in global histone PTM abundance across four SAM depletion

systems were analyzed for the presence of methyl PTMs whose abundances either increased

or were unaffected by this metabolic perturbation. Such PTMs would be candidate mediators of

an adaptive response to SAM depletion. Unexpectedly, all four SAM depletion systems

exhibited identical changes in H3K9 methylation (Figure 2G-2J). SAM depletion stimulated a

decrease in H3K9me2/3 with a corresponding increase in H3K9ac/14ac and unmodified

peptides while net H3K9me1 abundance was unchanged. Preservation of net H3K9me1 levels

in response to SAM depletion was reproducible in Met-restricted MCF7, HEK-293, Panc1, and

HepG2 cell lines, as well as in MAT2A-RNAi treated HepA1 cells (Figure S2A-S2F). The robust

conservation of H3K9me1 abundance in diverse systems marked by reduced methylation

capacity suggest this PTM may be supported by adaptive, active mechanisms.

H3K9me1 Abundance under SAM Depletion is Driven by *de novo* Methylation

Global levels of H3K9me1 could be maintained through three distinct mechanisms

during SAM depletion: 1.) protection from PTM turnover, 2.) de-methylation of H3K9me2/3,

and/or 3.) *de novo* methylation of unmodified H3K9. The decrease in H3K9me2/3 and increase

in unmodified H3K9 peptide levels across SAM depletion systems suggest H3K9 can be

subjected to facile demethylation while *de novo* methylation appears enzymatically unfavorable.

However, *de novo* H3K9 mono-methylation would constitute an active, adaptive response to this

metabolic stress that is capable of supporting critical chromatin functions.

To investigate the contribution of *de novo* methylation to global H3K9me1 levels, RNAi-

mediated repression of five H3K9 HMTs was first performed in Met-replete conditions to

determine which enzymes catalyzed a majority of H3K9 mono-methylation reactions in HCT116

cells. Repression of HMT expression had significant yet restricted effects, given technical

limitations, on H3K9 PTM abundance and suggested Ehmt1/Ehmt2 were the primary H3K9

mono-methyltransferases in this system (Figure S3A-S3D). Because repression of HMT

expression was intended to be coupled with Met-restriction, RNAi treatment could not be

extended to elicit larger changes in PTM abundance as HCT116 cells no longer proliferate

beyond 24 hours of Met-restriction (Figure S3E). Therefore, small molecule (UNC0642)

inhibition of H3K9 HMTs Ehmt1 and Ehmt2 was used as the best approach to acutely block

nuclear H3K9 mono-methylation (Liu F et al., 2013). UNC0642 treatment of Met-replete cells

reduced H3K9me1 abundance to a greater extent than RNAi repression of EHMT1/2 (Figure 3A

and Figure S3D). Coupling UNC0642 treatment with Met-restriction also stimulated a significant

decrease in H3K9me1 levels (Figure 3B and Figure S3F). Comparatively, these data suggest

Ehmt1/Ehmt2 retain nearly 60% of their H3K9 mono-methylation activity during severe *in situ*

SAM deprivation and that this activity is critical for preserving global H3K9me1 abundance

(Figure 3C). Furthermore, H3K9me2 abundance decreased significantly in Met-replete cells

upon UNC0642 treatment but was unaffected when UNC0642 was coupled with Met-restriction

(Figure 3A-3B). This indicates only the mono-methylation activity of Ehmt1 and Ehmt2 is

preferentially supported during SAM depletion.

In addition to methylation of H3K9 by nuclear enzymes, mono-methylation is known to

be enzymatically catalyzed on cytoplasmic histones prior to nuclear import and deposition onto

chromatin (Loyola A et al., 2006; Pinheiro et al., 2012). Subcellular fractionation of Met-

restricted HCT116 cells coupled with western blot analyses indicated that cytoplasmic

H3K9me1 levels increase 3-fold after 24 hours of Met-restriction, with a similar increase in total

H3 protein levels (Figure 3D-3E). To determine the contribution of cytoplasmic H3K9me1 to the

chromatin-bound nuclear pool after 24 hours of Met-restriction, a pulse-chase, SILAC

experiment was performed (Figure 3F). L-Arg$_{+10}$ isotope incorporation prior to Met-restriction

was determined to be nearly complete with average incorporation for H3K9 peptides greater

than 98% (Figure S3H). Newly synthesized H3 was calculated to comprise 24% of the

chromatin bound histone pool and contributed 18% of the global H3K9me1 signal after 24 hours

of Met-restriction (Figure 3G-3H). UNC0642 treatment did not significantly lessen the

contribution of cytoplasmic H3K9me1 to the total pool, suggesting cytoplasmic H3K9me1 may

be largely protected from enzyme catalyzed turnover once deposited onto chromatin under

these conditions (Figure 3H). Modest, insignificant decreases in L-Arg$_{+0}$ H3K9 PTM signal after

UNC0642 treatment can be attributed to reduced incorporation of newly synthesized H3 onto

chromatin. (Figure 3G) Reduced H3 incorporation in UNC0642-treated cells is likely the

consequence of a slower growth rate under these conditions (Figure S3E). Together, these

results suggest ~40% of global H3K9me1 abundance is driven by *de novo* methylation during

Met-restriction with nearly equal contributions from both the cytoplasm and nucleus. The

remaining ~60% of H3K9me1 levels are likely protected from turnover and/or are generated via

H3K9me2/3 de-methylation during SAM depletion (Figure 3I).

The significant upregulation of cytoplasmic H3K9me1 and total H3, coupled with their

incorporation onto chromatin during SAM depletion, were unexpected. These findings suggest

cells utilize scarcely available Met and SAM to support the protein abundance and methylation

of histone H3 under these conditions. Shotgun proteomics was performed on cytoplasmic
protein fractions of the previously described SILAC experiment to determine if histone protein
abundance is uniquely responsive to SAM depletion. Of the more than 2,400 identified proteins,
only 8 possessed significant incorporation of the light L-Arg$_{+0}$ isotope and also increased in
absolute abundance by a log$_2$ fold-change value greater than 2.0 (Figure 3J). Frequency of Met
residues in the primary amino acid sequence did not affect L-Arg$_{+0}$ isotope incorporation (Figure
S3I). Remarkably, all four core histone subunits, Mat2A, and the methionine importer SLC3A2
(Bröer et al., 2001) were among the 8 proteins (Figure 3J). These results suggest proteins
critical for chromatin stability and methyl-donor metabolism are preferentially generated during
metabolically-induced SAM depletion. To our knowledge, this is also the first example of a
metabolic stress stimulating increased histone protein abundance.
226
227   H3K9me1 is Redistributed Over Repetitive and Transposable Elements During SAM Depletion
228
Multiple, non-redundant mechanisms for H3K9me1 maintenance suggest strong
biological pressure exists to retain this PTM under SAM depleted conditions. One critical
function for H3K9me1 is to serve as a primer for H3K9me2/3 methylation as methyltransferases
that yield these higher methylation states require a mono-methyl substrate (Peters et al., 2001;
Loyola et al., 2006; Loyola et al., 2009). Accordingly, installment of H3K9me2/3 for constitutive
heterochromatin-mediated repression of telomeric, pericentromeric, and centromeric
repetitive/transposable elements requires readily available H3K9me1 (Rea et al., 2000;
Bannister et al., 2001; Lachner et al., 2000).
To determine if H3K9me1 is being maintained globally or redistributed to regions of
susceptible heterochromatin during SAM depletion, ChIP-sequencing was performed. Met-
restriction stimulated a 35% loss in total H3K9me3 peak abundance (Figure 4A). A higher
percentage of H3K9me3 peaks were lost at LTR, intergenic, and intron annotated loci relative to
the total, while H3K9me3 enrichment at satellite, simple repeat, and SINE annotated loci were
slightly less sensitive to SAM depletion (Figure 4A). Interestingly, H3K9me1 enrichment at
repetitive and transposable loci were either largely unaffected by SAM depletion or increased
(Figure 4B). Only the percent change for intergenic and intron annotated H3K9me1 peaks was
lost to a greater extent than the total peak number, suggesting H3K9me1 is more susceptible to
de-methylation at these genomic loci.
Maintained or increased levels of H3K9me1 at constitutively repressed regions during
SAM depletion could result from the replacement of H3K9me1 for lost H3K9me3. To investigate

this possibility, H3K9me1 reads were mapped to nearly 16e$_4$ regions determined to lose

H3K9me3 peak enrichment upon SAM depletion. This analysis determined H3K9me1

enrichment increases at 66.4% of these sites (Figure 4C-4D). A higher percentage of these

H3K9me1 enriched sites were annotated for LINE, LTR, simple repeat, and satellite elements

relative to those experiencing decreased H3K9me1 read enrichment (Figure 4E). Figure 4F

provides a representative image of H3K9me3 replacement by H3K9me1 at a pericentric,

repetitive loci. Together, these ChIP-sequencing analyses suggest SAM depletion stimulates

the redistribution and targeting of H3K9me1 methylation patterns to repetitive and transposable

genomic loci which are susceptible to higher state H3K9 demethylation.

De novo H3K9 Mono-methylation Preserves Heterochromatin Stability

261   Replacement of H3K9me3 by H3K9me1 may function to preserve heterochromatin

stability and pericentric chromatin repression during SAM depletion by providing the primer for

site-specific H3K9me2/3 methylation and/or by preventing H3K9 acetylation. An MNase

accessibility assay was used to evaluate global heterochromatin stability under SAM depleted

conditions as MNase can more readily digest euchromatic DNA compared to heterochromatic

DNA. Coupling Met-restriction with UNC0642 treatment resulted in a significantly greater loss

and accumulation of di- and mono-nucleosome species, respectively, compared to Met-

restriction alone (Figure 5A-5B). This indicates inhibition of *de novo* H3K9 mono-methylation

during SAM depletion has a more detrimental effect on global heterochromatin stability than

SAM depletion in isolation. To determine if global decreases in heterochromatin stability

resulted in de-repression of transposable DNA elements, RT-qPCR was employed. Transcript

abundance of *LINE 1*, *HERV-K*, and *HERV-R* were all significantly elevated after SAM

depletion. Coupling SAM depletion with UNC0642 treatment resulted in the further elevation of

LINE1 and *HERV-K* transcript abundance while *HERV-R* expression was unaffected (Figure 5C-

5E). Because SAM depletion-induced losses in heterochromatin stability and retrotransposon

repression are exacerbated with inhibition of nuclear Ehmt1/Ehmt2 activity, these data suggest

stimulation of *de novo* H3K9 mono-methylation functions, in part, to preserve heterochromatin

stability.

Adaptive H3K9 Mechanisms Support Epigenetic Persistence Upon Metabolic Recovery

Dysregulation of adaptive epigenetic mechanisms during SAM depletion may produce

aberrant changes in histone PTMs that compromise long-term epigenetic persistence upon

metabolic recovery. To determine if *de novo* H3K9me1 is required for epigenetic persistence to

SAM depletion, HCT116 cells were allowed to recover in Met-replete media after being

subjected to Met-restriction coupled with acute DMSO or UNC0642 treatment (Figure 6A).

Histone proteomics analysis confirmed that inhibition of Ehmt1/Ehmt2 activity during SAM

depletion results in a significant loss of H3K9me1 abundance (Figure 6B-6C). Upon 5 and 24

hours of Met-repletion, DMSO control cells largely regained their original histone PTM state

(Figure 6B). However, histone PTM states of cells acutely treated with UNC0642 only during

Met-restriction remained dysregulated after both 5 and 24 hours of Met-repletion (Figure 6B).

Although significant global dysregulation was apparent, the H3K9 peptide proteoforms displayed

the most dynamic differences (Figure 6D-6E). Interestingly, the signature H3K9 PTM response

to SAM depletion described in Figure 2 was largely retained upon Met-repletion in cells acutely

treated with UNC0642 (Figure 6D-6E). H3K9me3 abundance was an exception, recovering to

levels consistent with regained *LINE 1* and *HERV-K/R* repression (Figure 5C-5E).

Met-replete cells allowed to recover from acute UNC0642 treatment also experienced

prolonged changes in H3K9 PTM abundance after inhibitor removal (Figure S4A). As Ehmt1/2

inhibition by UNC0642 was determined to be reversible (Figure SB-S4C), these results suggest

disrupted persistence to small molecule treatment – in either isolation from or conjunction with

SAM depletion – is a direct consequence of acute losses in Ehmt1/Ehmt2 activity.

To assess the transcriptional consequences of lost epigenetic persistence, RNA-

sequencing was performed on cells which recovered in Met-replete media after Met-restriction

coupled with acute DMSO or UNC0642 treatment. Principle Component Analysis (PCA) of

normalized Transcripts Per Million (TPM) values produced 3 distinct clusters, suggesting altered

epigenomes correlate with altered transcript profiles in these cells (Figure 6F). A multiple

condition comparison analysis was used to determine which transcripts distinguished the

clusters from one another (Figure 6G). While 68.4% of identified genes were ubiquitously

expressed across samples (Pattern 1), 27.7% were uniquely expressed in control cells relative

to both recovery groups (Pattern 4) (Figure 6H). This result supports the PCA findings as the

pre-restriction control group is most distinct from both recovery groups. Unique gene expression

profiles for UNC0642 (Pattern 2) and DMSO (Pattern 3) recovery cells contained 1.9% and

0.7%, respectively, of all identified genes. Therefore, these subsets of genes are responsible for

the unique clustering of both Met-restriction recovery groups (Figure 6H). Gene Set Enrichment

Analysis (GSEA) was performed on both Pattern 2 and 3 gene lists to determine functional

differences between each gene set. GSEA of Pattern 2 genes identified 71 significantly enriched

gene ontology (GO) terms possessing an FDR q-value less than 0.25, while an identical

analysis of Pattern 3 genes did not identify any significantly enriched GO terms (Figure 6I-6J).

Of the significantly enriched GO terms in Pattern 2, 66 possessed a positive Normalized

Enrichment Score (NES) when comparing UNC0642 to DMSO control recovery cells (Figure

S5A-S5B). Figure 6K is a representative GSEA running enrichment score figure of a GO term

possessing a positive NES. Only 5 of the significantly enriched GO terms in Pattern 2

possessed a negative NES when comparing UNC0642 to DMSO control recovery cells.

Notably, a majority of the genes assigned to these GO terms were zinc finger binding proteins

(Figure S5C-S5D). Figure 6L is a representative GSEA running enrichment score figure of a GO

term possessing a negative NES.

These data suggest adaptive regulation of H3K9 PTMs in response to SAM depletion is

required for long-term epigenetic persistence upon metabolic recovery. Furthermore, inhibited

epigenetic persistence results in significant transcriptional consequences that impact a diverse

set of cellular functions.

Adaptive and Persistent Epigenetic Responses to SAM Depletion are Robust *in vivo,*

Independent of Age

To determine if adaptive and persistent responses to SAM depletion exist *in vivo* and are

age-independent, 6- and 22-month C57BL/6J mice were subjected to 3 weeks of Met-restriction

followed by 5 weeks of Met-repletion (Figure 7A). Several previous reports have associated

decreased H3K9me2/3 abundance with aging in numerous organisms (Wood et al., 2010;

Larson et al., 2012; Ni et al., 2012; Tvardovskey et al., 2017; Scaffidi and Misteli, 2006). Age-

dependent dysregulation of heterochromatin has been associated with de-repression of

repetitive DNA and transposable elements, as well as DNA damage, factors which may

contribute to disease progression (Wood et al., 2016; De Cecco et al., 2013; De Cecco et al.,

2019; Oberdoerffer and Sinclair, 2007; Booth and Brunet, 2016). Therefore, conservation of

both adaptive and persistent responses to SAM depletion across the lifespan would further

support the functional importance of both mechanisms as an "older" chromatin environment is

thought to be inherently dysregulated.

Remarkably, histone proteomics analysis of liver tissue revealed H3K9 was the only

residue on which PTMs responded similarly to SAM depletion in both 6- and 22-month mice

relative to age-matched controls (Figure 7B). This PTM response at H3K9 phenocopied those

identified in the previously described *in vitro* and whole-organism SAM depletion studies, further

highlighting the robust regulation of this residue (Figure 7C). Significant age-dependent

responses to SAM depletion were identified on all remaining histone residues (Figure 7B and

S6A-S6E). Furthermore, significantly elevated *LINE 1*, *SINE B1*, and *SINE B2* transcript

abundance in both 6- and 22-month mice suggest the chromatin instability phenotype

characterized in Figure 5 also occurs during metabolic depletion of SAM *in vivo* (Figure 7D-7E).

After 5 weeks of dietary Met-reintroduction, both young and old mice displayed the ability to re-

establish an epigenome similar to that possessed by age-matched controls (Figure 7B).

Interestingly, 22-month mice appeared to more rapidly adopt the control histone PTM state,

while the 6-month mice still exhibited residual losses in H3K9me3 and increases in H3K9me1

after 5 weeks of recovery (Figure 7F). Consistent with the general recovery of chromatin states,

both age groups largely reestablished repression of *LINE 1*, *SINE B1*, and *SINE B2* expression

(Figure 7G-7H).

Metabolic changes in young and old mice tracked with H3K9 PTMs. Both 6- and 22-

month mice in the Met-restriction group experienced significant losses in total body weight

relative to age matched controls (Figure S6F). Decreases in overall body weight were

comprised of significant losses in both lean and fat mass (Figure S6F). Altered body-weight and

compositions were accompanied by significantly improved glucose tolerance in both young and

old mice (Figure S6G). Upon Met-repletion, mice reacquired their initial metabolic state in an

age-independent manner. All animals experienced significant increases in total body weight,

comprised of elevated lean and fat mass, as well normalized glucose tolerance relative to age-

matched controls (Figure S6H-S6I).

Together, strong association of the H3K9 chromatin response to SAM depletion with

metabolic reprogramming in young and old mice suggest H3K9 regulatory mechanisms are not

only robust across lifespan but may be required to support these metabolic phenotypes.

Furthermore, these data show the ability of epigenetic states to persist upon recovery from a

metabolic challenge is an inherent property of both isolated cells and complex mammals.

Discussion

Here, we provide a comprehensive investigation into the adaptive regulation of

chromatin during a metabolic stress. A highly conserved, robust histone methylation response to

SAM depletion was discovered, characterized by global losses in histone H3 di- and tri-

methylation while H3K9me1 is actively maintained to safeguard against exacerbated

heterochromatin instability and support epigenetic persistence upon metabolic recovery. These mechanisms involve the enrichment of H3K9me1 in susceptible regions of heterochromatin through *de novo* methylation of both chromatin-bound and cytoplasmic H3. To our knowledge, this is the first identified mechanism in which scarce, diet-derived nutrients are funneled toward the preservation of specific chromatin PTM states.

Zee et al., 2010 have previously determined H3K9me1 has the shortest half-life of 17 measured PTMs, highlighting the importance of *de novo* methylation for supporting global H3K9me1 abundance in a SAM depleted environment. Here, *de novo* methylation was found to be critical for more than simply sustaining global H3K9me1 levels under these conditions. ChIP-sequencing analyses determined H3K9me1 patterns experience a significant redistribution across heterochromatin that results in H3K9me1 enrichment at repetitive and transposable DNA elements, particularly in regions where H3K9me3 is lost. Targeting of H3K9me1 to these regions is likely critical for preserving heterochromatin as prevention of *de novo* H3K9 mono-methylation resulted in global losses in heterochromatin stability and site-specific retrotransposon derepression. We hypothesize H3K9me1 facilitates repression of retrotransposons during SAM depletion through two simultaneous mechanisms: 1.) acting as the primed substrate for HMTs which deposit di- and tri-methyl PTMs onto H3K9 (Peters et al., 2001; Loyola et al., 2006; Loyola et al., 2009; Rea et al., 2000; Bannister et al., 2001; Lachner et al., 2000) and 2.) preventing the acetylation of newly demethylated H3K9 residues. Increased H3K9 acetylation through impaired H3K9 deacetylation pathways is associated with global genomic instability as well as derepression of LINE1, providing support for the second mechanism (Mostoslavsky et al., 2006; Van Meter et al., 2014; Simon et al., 2019). Ensuring sufficient amounts of H3K9me1 are available to facilitate heterochromatin stability is not limited to nuclear derived H3K9me1. Pinheiro et al., 2012 have previously shown inhibition of cytoplasmic H3K9 mono-methylation through shRNA knockdown of *PRDM3* and *PRDM16* stimulates pericentric heterochromatin loss and disruption of the nuclear lamina. Our study demonstrates cytoplasmic derived H3K9me1 comprises a significant portion (18%) of the total chromatin bound pool after 24 hours of Met-restriction *in vitro*. Adaptation of H3K9 mono-methylation to SAM depletion is also critical for long-term epigenetic persistence to this metabolic stress. Prevention of *de novo* H3K9 mono-methylation during SAM depletion resulted in sustained dysregulation of global histone PTMs upon metabolic recovery, significantly influencing gene expression patterns. This requirement for an active, adaptive histone PTM response to support epigenetic persistence upon metabolic recovery is the first identified mechanism of its kind.

Replication of *in vitro* results *in vivo*, independent of age, strongly supports the discovery
of a universal adaptive and persistent response to SAM depletion. Here, we demonstrate that
the adaptive mechanisms required to support H3K9me1-mediated heterochromatin stability are
inducible in both 6- and 22-month mice after dietary Met-restriction. Furthermore, H3K9me1
maintenance under SAM depleted conditions uniquely correlated with beneficial metabolic
reprogramming (i.e. reduced fat mass and improved glucose tolerance) in both young and old
mice. This suggests the robust regulation of H3K9 PTMs in response to SAM depletion may be
required to support the metabolic reprogramming stimulated by dietary Met-restriction. Global
epigenetic persistence to metabolic SAM depletion was observed in both 6- and 22-month mice.
Validation of this phenomenon *in vivo*, independent of age, suggests similar mechanisms critical
for maintaining proper regulation of the epigenome during life-experienced fluctuations in the
availability of other metabolic cofactors may exist as well. Such fluctuations could be stimulated
by periods of prolonged fasts, chronic intake of foods lacking essential cofactors (i.e. vegan and
vegetarian dietary patterns), and circadian regulated changes in metabolite availability (Farmer,
2014; Schmidt et al., 2016; Krishnaiah et al., 2017).

Notably, we did not observe an age-associated decrease in H3K9 methylation between
6- and 22-month control diet mice (Table S1). Similar regulation of H3K9 PTMs and mirrored
metabolic responses to SAM depletion across age groups suggest our 22-month mice, the
equivalent to a 60-70 year old human, have not yet experienced a decline in health-span
(Flurkey et al., 2007). It will be interesting to determine if the adaptive and persistent epigenetic
mechanisms revealed here are functional in very old animals (i.e. >30 months) where H3K9
methylation is thought to be inherently dysregulated. Disruption of these mechanisms later in an
organism's lifespan could support age-associated derepression of transposable elements
(Wood et al., 2016; De Cecco et al., 2013; Oberdoerffer and Sinclair, 2007; Booth and Brunet,
2016), contributing to negative phenotypes that accelerate the aging process such as
inflammation (De Cecco et al., 2019; Simon et al., 2019).

Future studies will be needed to determine how H3K9 mono-methyltransferases are able
to maintain their catalytic activity in SAM depleted environments. Favorable steady-state kinetic
constants for H3K9me1 relative to H3K9me2/3 HMTs is one possibility. However, it is difficult to
address this hypothesis by comparing existing literature on H3K9 HMT enzymology, due to
disparate substrates and assays. Direct shuttling of SAM to H3K9me1 HMTs by Mat2A could
also support H3K9me1 HMT catalytic activity. As SAM is energetically expensive to produce,
due to the 1:1 requirement of Met and ATP molecules for its synthesis, in addition to being
highly unstable (Wu et al., 1983), it would be advantageous for the cell to possess mechanisms

which permit preferential shuttling of SAM during methyl-metabolite depletion. Directed SAM

production and utilization have been proposed in other metabolic contexts (Katoh et al., 2011;

Kera et al., 2013; Li et al., 2015).

Acknowledgements

This research was supported in part by grants from the NIH (S.A.H. –T32 DK007665; K.A.K.

T32 DK007665; W.H.L. T32 DK007665; J.M.D. – R37 GM059785; D.W.L. – AG041765,

AG050135, AG051974, AG056771, AG062328, a New Investigator Program Award (D.W.L.)

and a Collaborative Health Sciences Program Award (V.L.C.) from the Wisconsin Partnership

Program, the V. Foundation for Cancer Research (V.L.C.), and a Glenn Foundation Award for

Research in the Biological Mechanisms of Aging (D.W.L.), as well as startup funds from the

UW-Madison School of Medicine and Public Health and the UW-Madison Department of

Medicine (V.L.C. and D.W.L.). This research was conducted while D.W.L. was an AFAR

Research Grant recipient from the American Federation for Aging Research. D.Y. is supported

in part by a fellowship from the American Heart Association (17PRE33410983). The Lamming

laboratory is supported in part by the U.S. Department of Veterans Affairs (I01-BX004031), and

this work was supported using facilities and resources from the William S. Middleton Memorial

Veterans Hospital. This work does not represent the views of the Department of Veterans

Affairs or the United States Government. We would like to thank Alexis J. Lawton for the

bioinformatics analysis presented in Figure S3I. We would also like to thank the University of

Wisconsin-Madison Biotechnology Center and the University of Wisconsin-Madison Center for

High Throughput Computing for their help in generating and analyzing the ChIP-sequencing

data presented in Figure 5, respectively.

Author Contributions

Conceptualization, S.A.H., D.W.L, V.L.C., and J.M.D.; Methodology, S.A.H., C.Y., C.K.W.,

L.N.C., W.H.L., K.I., B.P.T., D.W.L., and J.M.D.; Validation, S.A.H., D.Y., C.Y., L.N.C., K.A.K.,

K.I., B.P.T., D.W.L., and J.M.D.; Formal Analysis, S.A.H., D.Y., C.K.W., K.A.K., D.W.L., and

J.M.D; Investigation, S.A.H., D.Y., C.Y., L.N.C., K.A.K., J.L.T., S.E.Y., and B.R.M.; Resources,

K.I., B.P.T., V.L.C., D.W.L., and J.M.D.; Writing – Original Draft, S.A.H. and J.M.D.; Writing –

Review & Editing, S.A.H., D.Y., C.Y., C.K.W., W.H.L., K.I., R.S., B.P.T., V.L.C., D.W.L., and

J.M.D.; Visualization, S.A.H., D.Y., and D.W.L.; Supervision and Project Administration, K.I.,

R.S., B.P.T., D.W.L., and J.M.D.; Funding Acquisition, S.A.H., K.I., R.S., B.P.T., V.L.C., D.W.L.,

and J.M.D.

489   Declaration of Interests

491   J.M.D. is a consultant for FORGE Life Sciences and co-founder of Galilei BioSciences. D.W.L

492   has received funding from, and is a scientific advisory board member of, Aeonian

493   Pharmaceuticals, which seeks to develop novel, selective mTOR inhibitors for the treatment of

494   various diseases. Remaining authors declare no competing interest.

References

1. Arriola Apelo, S.I., Neuman, J.C., Baar, E.L., Syed, F.A., Cummings, N.E., Brar, H.K., Pumper, C.P., Kimple, M.E., and Lamming, D.W. (2016) Alternative rapamycin treatment regimens mitigate the impact of rapamycin on glucose homeostasis and the immune system. Aging Cell. 15, 28-38.

2. Bannister, A.J., Zegerman, P., Pardridge, J.F., Miska, E.A., Thomas, J.O., Allshire, R.C., and Kouzarides, T. (2001) Selective recognition of methylated lysine 9 on histone H3 by the HP1 chromo domain. Nature. 410, 120-124.

3. Bolger, A.M., Lohse, M., and Usadel, B. (2014) Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics. 30, 2114-2120.

4. Booth, L.N. and Brunet, A. (2016) The aging epigenome. Mol. Cell. 62, 728-744.

5. Bröer, A., Friedrich, B., Wagner, C.A., Fillon, S., Ganapathy, V., Lang, F., and Bröer, S. (2001) Association of 4F2hc with light chains LAT1, LAT2 or y+LAT2 requires different domains. Biochem J. 355, 735-731.

6. Brown-Borg, H.M., Rakoczy, S., Wonderlich, J.A., Borg, K.E., and Rojanathammanee, L. (2018) Metabolic adaptation of short-living growth hormone transgenic mice to methionine restriction and supplementation. Ann. N.Y. Acad. Sci. 1418, 118-136.

7. Clasquin, M.F., Melamud, E., and Rabinowitz, J.D. (2012) LC-MS processing with MAVEN: a metabolomic analysis and visualization engine. Curr. Protoc. Bioinformatics. Chapter 14, Unit 14.11. doi:10.1002/0471250953.bi1411s37.

8. Cox, J., Hein, M.Y., Luber, C.A., Paron, I., Nagaraj, N., and Mann, M. (2014) Accurate proteome-wide label-free quantification by delayed normalization and maximal peptide ratio extraction, termed MaxLFQ. Mol. Cell Proteomics. 13, 2513-2526.

9. De Cecco, M., Criscione, S.W., Peckham, E.J., Hillenmeyer, S., Hamm, E.A., Manivannan, J., Peterson, A.L., Kreiling, J.A., Neretti, N., and Sedivy, J.M. (2013) Genomes of replicatively senescent cells undergo global epigenetic changes leading to gene silencing and activation of transposable elements. Aging Cell. 12, 247-256.

10. De Cecco, M., Ito, T., Petrashen, A.P., Elias, A.E., Skivr, N.J., Criscione, S.W., Caligiana, A., Brocculi, G., Adney, E.M., Boeke, J.D., Le, O., Beauséjour, C., Ambati, J., Ambati, K., Simon, M., Seluanov, A., Gorbunova, V., Slagboom, P.E., Helfand, S.L., Neretti, N., and Sedivy, J.M. (2019) L1 drives IFN senescent cells and promotes age-associated inflammation. Nature. 566, 73-78.

11. Ding, W., Smulan, L.J., Hou, N.S., Taubert, S., Watts, J.L., and Walker, A.K. (2015) S-
adenosylmethionine levels govern innate immunity through distinct methylation-
dependent pathways. Cell Metab. 22, 633-645.

12. Ducker, G.S. and Rabinowitz, J.D. (2017) One-carbon metabolism in health and
disease. Cell Metab. 25, 27-42.

13. Fan, J., Krautkramer, K.A., Feldman, J.L., and Denu, J.M. (2015) Metabolic regulation of
histone post-translational modifications. ACS Chem. Biol. 10, 95-108.

14. Farmer, B. (2014) Nutritional adequacy of plant-based diets for weight management:
observations from the NHANES. Am. J. Clin. Nutr. 100 Suppl 1, 365S-368S.

15. Flurkey, K., Currer, J.M., and Harrison, D.E. (2007) The Mouse in Aging Research. In
the Mouse in Biomedical Research 2nd Edition. Fox, J.G., et al., editors. American
College Laboratory Animal Medicine (Elsevier), pp. 637-671.

16. Green, C.L. and Lamming, D.W. (2018) Regulation of metabolic health by essential
dietary amino acids. Mech. Ageing Dev. S0047-6374(18)30079-4. doi:
0.1016/j.mad.2018.07.004.

17. Greer, E.L. and Shi, Y. (2012) Histone methylation: a dynamic mark in health, disease,
and inheritance. Nat. Rev. Genet. 13, 343-357.

18. Hardies, S.C., Wang, L., Zhou, L., Zhao, Y., Casavant, N.C., and Huang, S. (200) LINE-
1 (L1) lineages in the mouse. Mol. Biol. Evol. 17, 616-628.

19. Hayashi, T., Teruya, T., Chaleckis, R., Morigasaki, S., and Yanagida, M. (2018) S-
adenosylmethionine synthetase is required for cell growth, maintenance of G0 phase,
termination of quiescence in fission yeast. iScience 5, 38-51.

20. Heinz, S., Benner, C., Spann, N., Bertolino, E., Lin, Y.C., Laslo, P., Cheng, J.X., Murre,
C., Singh, H., and Glass, C.K. (2010) Simple combinations of lineage-determining
transcription factors prime cis-regulatory elements required for macrophage and B cell
identities. Mol. Cell. 38, 576-589.

21. Kang, Y.K., Park, J.S., Lee, C.S., Yeom, Y.I., Chung, A.S., and Lee, K.K. (1999) Efficient
integration of short interspersed element-flanked foreign DNA via homologous
recombination. J. Biol. Chem. 274, 36585-36591.

22. Katoh, Y., Ikura, T., Hoshikawa, Y., Tashiro, S., Ito, T., Ohta, M., Kera, Y., Noda, T., and
Igarashi, K. (2011) Methionine adenosyltransferase II serves as a transcriptional
corepressor of Maf oncoprotein. Mol. Cell. 41, 554-566.

23. Kera, Y., Katoh, Y., Ohta, M., Matsumoto, M., Takano-Yamamoto, T., and Igarashi, K.

(2013) Methionine adenosyltransferase II-dependent histone H3K9 methylation at the

COX-2 gene locus. J. Biol. Chem. 288, 13592-13601.

24. Klawitter, S., Fuchs, N.V., Upton, K.R., Upton, K.R., Muñoz-Lopez, M., Shukla, R.,

Wang, J., Garcia-Cañadas, M., Lopez-Ruiz, C., Gerhardt, D.J., Sebe, A., Grabundzija, I.,

Merkert, S., Gerdes, P., Pulgarin, J.A., Bock, A., Held, U., Witthuhn, A., Haase, A.,

Sarkadi, B., Löwer, J., Wolvetang, E.J., Martin, U., Ivics, Z., Izsvák, Z., Garcia-Perez,

J.L., Faulkner, G.J., and Schumann, G.G. (2016) Reprogramming triggers endogenous

L1 and Alu retrotransposition in human induced pluripotent stem cells. Nat. Commun.

7:10286. doi: 10.1038/ncomms10286.

25. Konze, K.D., Ma, A., Li, F., Barsyte-Lovejoy, D., Parton, T., Macnevin, C.J., Liu, F., Gao,

C., Huang, X.P., Kuznetsova, E., Rougie, M., Jiang, A., Pattenden, S.G., Norris, J.L.,

James, L.I., Roth, B.L., Brown, P.J., Frye, S.V., Arrowsmith, C.H., Hahn, K.M., Wang,

G.G., Vedadi, M., and Jin, J. (2013) An orally bioavailable chemical probe of lysine

methyltransferases EZH2 and EZH1. ACS Chem. Biol. 8, 1324-1334.

26. Krishnaiah, S.Y., Wu, G., Altman, B.J., Growe, J., Rhoades, S.D., Coldren, F.,

Venkataraman, A., Olarerin-George, A.O., Francey, L.J., Mukherjee, S., Girish, S.,

Selby, C.P., Cal, S., Er, U., Sianati, B., Sengupta, A., Anafi, R.C., Kavakli, I.H., Sancar,

A., Baur, J.A., Dang, C.V., Hogenesch, J.B., and Weljie, A.M. (2017) Clock regulation of

metabolites reveals coupling between transcription and metabolism. Cell Metab. 35,

961-974.

27. Lachner, M., O'Carroll, D., Rea, S., Mechtler, K., and Jenuwein, T. (2001) Methylation of

histone H3 lysine 9 creates a binding site for HP1 proteins. Nature. 410, 116-120.

28. Larson, K., Yan, S.J., Tsurumi, A., Liu, J., Zhou, J., Gaur, K., Guo, D., Eickbush, T.H.,

and Li, W.X. (2012) Heterochromatin formation promotes longevity and represses

ribosomal RNA synthesis. PLoS Genet. 8, e1002473. doi:

10.1371/journal.pgen.1002473.

29. Lees, E.K., Król, E., Grant, L., Shearer, K., Wyse, C., Moncur, E., Bykowska, A.S.,

Mody, N., Gettys, T.W., and Delibegovic, M. (2014) Methionine restriction restores a

younger metabolic phenotype in adult mice with alterations in fibroblast growth factor 21.

Aging Cell. 13, 817-827.

30. Leng, N., Dawson, J.A., Thomson, J.A., Ruotti, V., Rissman, A.I., Smits, B.M., Haag,

J.D., Gould, M.N., Stewart, R.M., and Kendziorski, C. (2013) EBSeq: an empirical Bayes

hierarchical model for inference in RNA-seq experiments. Bioinformatics. 29, 1035-1043.

31. Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G.,
Abecasis, G., Durbin, R., and 1000 Genome Project Data Processing Subgroup. (2009)
The sequence alignment/map format and SAMtools. Bioinformatics. 25, 2078-2079.

32. Li, B. and Dewey, C.N. (2011) RSEM: accurate transcript quantification from RNA-Seq
data with or without a reference genome. BMC Bioinform. doi: 10.1186/1471-2105-12-
323.

33. Li, S., Swanson, S.K., Gogol, M., Florens, L., Washburn, M.P., Workman, J.L, and
Suganuma, T. (2015) Serine and SAM responsive complex SESAME regulates histone
modification crosstalk by sensing cellular metabolism. Mol. Cell. 60, 408-421.

34. Li, X., Egervari, G., Wang, Y., Berger, S.L., and Lu, Z. (2018) Regulation of chromatin
and gene expression by metabolic enzymes and metabolites. Nat. Rev. Mol. Cell Biol.
19, 563-578.

35. Liu, F., Barsyte-Lovejoy, D., Li, F., Xiong, Y., Korboukh, V., Huang, X.P., Allali-Hassani,
A., Janzen, W.P., Roth, B.L., Frye, S.V., Arrowsmith, C.H., Brown, P.J., Vedadi, M., and
Jin, J. (2013) Discovery of an in vivo chemical probe of the lysine methyltransferases
G9a and GLP. J. Med. Chem. 56, 8931-8942.

36. Loyola, A., Bonaldi, T., Roche, D., Imhof, A., and Almouzni, G. (2006) PTMs on H3
variants before chromatin assembly potentiate their final epigenetic state. Mol. Cell. 24,
309-316.

37. Loyola, A., Tagami, H., Bonaldi, T., Roche, D., Quivy, J.P., Imhof, A., Nakatani, Y., Dent,
S.Y. and Almouzni, G. (2009) The HP1α-CAF1-SETDB1-containing complex provides
H3K9me1 for Suv39-mediated K9me3 in pericentric heterochromatin. Embo Rep. 10,
769-775.

38. MacLean, B., Tomazela, D.M., Shulman, N., Chamber, M., Finney, G.L., Frewen, B.,
Kern, R., Tabb, D.L., Liebler, D.C., and MacCoss, M.J. (2010) Skyline: an open source
document editor for creating and analyzing targeted proteomics experiments.
Bioinformatics. 26, 966-968.

39. Mauger, O., Klinck, R., Chabot, B., Muchardt, C., Allemand, E., and Batské, E. (2015)
Alternative splicing regulates the expression of G9A and SUV39H2 methyltransferases,
and dramatically changes SUV39H2 functions. Nucleic Acids Res. 43, 1869-1882.

40. Melamud, E., Vastag, L, and Rabinowitz, J.D. (2010) Metabolomic analysis and
visualization engine for LC-MS data. Anal. Chem. 82, 9818-9826.

41. Mentch S.J. and Locasale J.W. (2016) One-carbon metabolism and epigenetics:
understanding the specificity. Ann N. Y. Acad. Sci. 1363, 91-98.

42. Mentch, S.J., Mehrmohamadi, M., Huang, L., Liu, X., Gupta, D., Mattocks, D., Gómez
Padilla, P., Ables, G., Bamman, M.M., Thalacker-Mercer, A.E. Nichenametla, S.N., and
Locasale, J.W. (2015) Histone methylation dynamics and gene regulation occur through
the sensing of one-carbon metabolism. Cell Metab. 22,861-873.

43. Miller, R.A., Buehner, G., Chang, Y., Harper, J.M., Sigler, R., and Smith-Wheelock, M.
(2005) Methionine-deficient diet extends mouse lifespan, slows immune and lens aging,
alters glucose, T4, IGF-I and insulin levels, increases hepatocyte MIF levels and stress
resistance. Aging Cell. 4, 119-125.

44. Mootha, V.K., Lindgren, C.M., Eriksson, K.F., Subramanian, A., Sihag, S., Lehar, J.,
Puigserver, P., Carlsson, E., Ridderstråle, M., Laurila, E., Houstis, N., Daly, M.J.,
Patterson, N., Mesirov, J.P., Golub, T.R., Tamayo, P., Spiegelman, B., Lander, E.S.,
Hirschhorn, J.N., Altshuler, D., and Groop, L.C. (2003) PGC-1alpha responsive genes
involved in oxidative phosphorylation are coordinately downregulated in human diabetes.
Nat. Genet. 34, 267-273.

45. Mostoslavsky, R., Chua, K.F., Lombard, D.B., Pang, W.W., Fischer, M.R., Gellon, L.,
Liu, P., Mostoslavsky, G., Franco, S., Murphy, M.M., Mills, K., Patel, P., Hsu, J.T., Hong,
A.L., Ford, E., Cheng, H.L., Kennedy, C., Nunez, N., Bronson, R., Frendewey, D.,
Auerbach, W., Valenzuela, D., Karow, M., Hottiger, M.O., Hursting, S., Barret, J.C.,
Guarente, L., Mulligan, R., Demple, B., Yancopoulos, G.D., and Alt, F.W. (2006)
Genomic instability and aging-like phenotype in the absence of mammalian SIRT6. Cell.
124, 315-329.

46. Ni, Z., Ebata, A., Alipanahiramandi E., and Lee, S.S. (2012) Two SET domain containing
genes link epigenetic changes and aging in Caenorhabditis elegans. Aging Cell. 11, 315-
325.

47. Northcott, P.A., Nakahara, Y., Wu, X., Feuk, L., Ellison, D.W., Croul, S., Mack, S.,
Kongkham, P.N., Peacock, J., Dubuc, A., Ra, Y.S., Zilberberg, K., Mcleod, J., Scherer,
S.W., Rao, J.S., Eberhart, C.G., Grajkowska, W., Gillespie, Y., Lach, B., Grundy, R.,
Pollack, I.F., Hamilton, R.L., Van Meter, T., Carlotti, C.G., Boop, F., Bigner, D.,
Gilbertson, R.J., Rutka, J.T., and Taylor, M.D. (2009) Multiple recurrent genetic events
converge on control of histone lysine methylation in medulloblastoma. Nat. Genet. 41,
465-472.

48. Oberdoerffer, P. and Sinclair, D.A. (2007) The role of nuclear architecture in genome
instability and ageing. Nat. Rev. Mol. Cell Biol. 8, 692-702.

49. Orentreich, N., Matias, J.R., DeFelice, A., and Zimmerman, J.A. (1993) Low methionine
ingested by rats extends life span. J. Nutr. 123, 269-274.

50. Peters, A.H., O'Carroll, D., Scherthan, H., Mechtler, K., Sauer, S., Schöfer, C.,
Weipoltshammer, K., Pagani, M., Lachner, M., Kohlmaier, A., Opravil, S., Doyle, M.,
Sibilia, M., and Jenuwein, T. (2001) Loss of the Suv39h histone methyltransferase
impairs mammalian heterochromatin and genome stability. Cell. 107, 323-327.

51. Pinheiro, I., Margueron, R., Shukeir, N., Eisold, M., Fritzsch, C., Richter, F.M., Mittler, G.,
Genoud, C., Goyama, S., Kurokawa, M., Son, J., Reinberg, D., Lachner, M., and
Jenuwein, T. (2012) Prdm3 and Prdm16 are H3K9me1 methyltransferases required for
mammalian heterochromatin integrity. Cell. 150, 948-960.

52. Qi, F., Shang, K., Zhang, J., Chuai, S., Kong, D., Zhou, T., Fu, S., Liang, Y., Li, C.,
Chen, Z., Zhao, Y., Yu, Z., Huang, Z., Hu, M., Ying, H., Chen, Z., Zhang, Y., Xing, F.,
Zhu, J., Xu, H., Zhao, K., Lu, C., Atadja, P., Xia, Z.X., Li, E., and Shou, J. (2015) Histone
methyltransferase SETDB1 regulates liver cancer cell growth through methylation of
p53. Nat. Commun. 6, 8651.

53. Quinlan, A.R. and Hall, I.M. (2010) BEDTools: a flexible suite of utilities for comparing
genomic features. Bioinformatics. 26, 841-842.

54. Rea S., Eisenhaber, F., O'Carroll, D., Strahl, B.D., Sun, Z.W., Schmid, M., Opravil, S.,
Mechtler, K., Pontin, C.P., Allis, C.D., and Jenuwein, T. (2000) Regulation of chromatin
structure by site-specific histone H3 methyltransferases. Nature. 406, 593-599.

55. Rhyu, D.W., Kang, Y.J., Ock, M.S., Eo, J.W., Choi, Y.H., Kim, W.J., Leem, S.H., Yi, J.M.,
Kim, H.S., and Cha, H.J. (2014) Expression of human endogenous retrovirus env genes
in the blood of breast cancer patients. Int. J. Mol. Sci. 15, 9173-9183.

56. Scaffidi, P. and Misteli, T. (2006) Lamin A-dependent nuclear defects in human aging.
Science. 312, 1059-1063.

57. Schmidt, J.A., Rinaldi, S., Scalber, A., Ferrari, P., Achaintre, D., Gunter, M.J., Appleby,
P.N., Key, T.J., and Travis, R.C. (2016) Plasma concentrations and intakes of amino
acids in male meat-eaters, fish-eaters, vegetarians and vegans: a cross-sectional
analysis in EPIC-Oxford cohort. Eur. J. Clin. Nutr. 70, 306-312.

58. Shiraki, N., Shiraki, Y., Tsuyama, T., Obata, F., Miura, M., Nagae, G., Aburatani, H.,
Kume, K., Endo, F., and Kume, S. (2014) Methionine metabolism regulates maintenance
and differentiation of human pluripotent stem cells. Cell Metab. 19, 780-794.

59. Shyh-Chang, N., Locasale, J.W., Lyssiotis, C.A., Zheng, Y., Teo, R.Y.,
Ratanasirintrawoot, S., Zhang, J., Onder, T., Unternaehrer, J.J., Zhu, H., Asara, J.M.,

Daley, G.Q., and Cantley, L.C. (2013) Influence of threonine metabolism on S-
adonsylmethionine and histone methylation. Science. 339, 222-226.

60. Simon, M., Van Meter, M., Ablaeva, J., Ke, Z., Gonzalez, R.S., Taguchi, T., De Cecco,
699 M., Leonova, K.I., Kogan, V., Helfand, S.L., Neretti, N., Roichman, A., Cohen, H.Y.,
700 Meer, M.V., Gladyshev, V.N., Antoch, M.P., Gudkov, A.V., Sedivy, J.M., Seluanov, A.,
701 and Gorbunova, V. (2019) LINE1 derepression in aged wild-type and SIRT6-deficient
702 mice drives inflammation. Cell Metab. 29, 871-885.

61. Strekalova, E., Malin, D., Weisenhorn, E.M.M., Russell, J.D., Hoelper, D., Jain, A., Coon,
704 J.J., Lewis, P.W., and Cryns, V.L. (2019) S-adenosylmethionine biosynthesis is a
705 targetable metabolic vulnerability of cancer stem cells. Breast Cancer Res. Treat. 175,
706 39-50.

62. Subramanian, A., Tamayo, P., Mootha, V.K., Mukherjee, S., Ebert, B.L., Gillette, M.A.,
708 Paulovich, A., Pomeroy, S.L., Golub, T.R., Lander, E.S., and Mesirov, J.P. (2005) Gene
709 set enrichment analysis: a knowledge-based approach for interpreting genome-wide
710 expression profiles. Proc. Natl. Acad. Sci.102, 15545-15550.

63. Tang, S., Fang, Y., Huang, G., Xu, X., Padilla-Banks, E., Fan, W., Xu, Q., Sanderson,
712 S.M., Foley, J.F., Dowdy, S., McBurney, M.W., Fargo, D.C., Williams, C.J., Locasale,
713 J.W., Guan, Z., and Li, X. (2017) Methionine metabolism is essential for SIRT1-regulated
714 mouse embryonic stem cell maintenance and embryonic development. EMBO J. 36,
715 3175-3193.

64. Tang, X., Keenan, M.M., Wu, J., Lin, C.A., Dubois, L., Thompson, J.W., Freedland, S.J.,
717 Murphy, S.K., and Chi, J.T. (2015) Comprehensive profiling of amino acid response
718 uncovers unique methionine-deprived response dependent on intact creatine
719 biosynthesis. PLoS Genet. 11, e1005158. doi: 10.1371/journal.pgen.1005158.

65. Towbin, B.D., González-Aguilera, C., Sack, R., Gaidatzis, D., Kalck, V., Meister, P.,
721 Askjaer, P., and Gasser, S.M. (2012) Step-wise methylation of histone H3K9 positions
722 heterochromatin at the nuclear periphery. Cell. 150, 934-937.

66. Tvardovskiy, A., Schwämmle, V., Kempf, S.J., Rogowska-Wrzesinska, A., and Jensen,
724 O.N. (2017) Accumulation of histone variant H3.3 with age is associated with profound
725 changes in histone methylation landscape. Nucleic Acids Res. 45, 9272-9289.

67. Van Meter, M., Kashyap, M., Rezazadeh, S., Geneva, A.J., Morello, T.D., Seluanoc, A.,
and Gorbunova, V. (2014) SIRT5 represses LINE1 retrotransposons by ribosylating
KAP1 but this repression fails with stress and age. Nat. Commun. Article number 5011.

68. Wang, Z., Yip, L.Y., Lee, J.H.J., Wu, Z., Chew, H.Y., Chong, P.K.W., Teo, C.C., Ang,
H.Y.K., Peh, K.L.E., Yuan, J., Ma, S., Choo, L.S.K., Basri, N., Jiang, X., Yu, Q., Hillmer,
A.M., Lim, W.T., Lim, T.K.H., Takano, A., Tan, E.H., Tan, D.S.W., Ho, Y.S., Lim, B., and
Tam, W.L. (2019) Methionine is a metabolic dependency of tumor-initiating cells. Nat.
Med. 25, 825-837.
69. Watson, G.W., Wickramasekara, S., Palomera-Sanchez, Z., Black, C., Maier, C.S.,
Williams, D.E., Dashwood, R.H., and Ho, E. (2014) SUV39H1/H3K9me3 attenuates
sulforaphane-induced apoptotic signaling in PC3 prostate cancer cells. Oncogenesis. 3,
e131.
70. Wood, J.G., Hillenmeyer, S., Lawrence, C., Chang, C., Hosier, S., Lightfoot, W.,
Mukherjee, E., Jiang, N., Schorl, C., Brodsky, A.S., Neretti, N., and Helfand, S.L. (2010)
Chromatin remodeling in the aging genome of Drosophila. Aging Cell. 9, 971-978.
71. Wood, J.G., Jones, B.C., Jiang, N., Chang, C., Hosier, S., Wickremesinghe, P., Garcia,
M., Harnett, D.A., Burhenn, L., Neretti, N., and Helfand, S.L. (2016) Chromatin-modifying
genetic interventions surpress age-associated transposable element activation and
extend life span in *Drosophila*. Proc. Natl. Acad. Sci. 113, 11277-11282.
72. Wu, S.E., Huskey, W.P., Borchardt, R.T., and Schowen, R.L. (1983) Chiral instability at
the sulfur of S-adenosylmethionine. Biochemistry. 22, 2828-2832.
73. Ye, C., Sutter, B.M., Wang, Y., Kuang, Z., and Tu, B.P. (2017) A metabolic function for
phospholipid and histone methylation. Mol. Cell 66, 180-193.
74. Yu, D., Yang, S.E., Miller, B.R., Wisinski, J.A., Sherman, D.S., Brinkman, J.A.,
Tomasiewicz, J.L., Cummings, N.E., Kimple, M.E., Cryns, V.L., and Lamming, D.W.
(2018) Short-term methionine deprivation improves metabolic health via sexually
dimorphic, mTORC-1 independent mechanisms. FASEB J. 32, 3471-3482.
75. Zhang, Y., Liu, T., Meyer, C.A., Eeckhoute, J., Johnson, D.S., Bernstein, B.E.,
Nusbaum, C., Myers, R.M., Brown, M., Li, W., and Liu, X.S. (2008) Model-based
analysis of ChIP-Seq (MACS). Genome Biol. 9, R137. doi: 10.1186/gb-2008-9-9-r137.
76. Zee, B.M., Levin, R.S., Xu, B., Leroy, G., Wingreen, N.S., and Garcia, B.A. (2010) In
vivo residue-specific histone methylation dynamics. J. Biol. Chem. 285, 3341-3350.
758

Main Figure Titles and Legends

Figure 1. Methionine Restriction Stimulates Dynamic Histone PTM Response

(A) Metabolic pathway diagram illustrating changes in Met-cycle metabolite abundance after 24 hours and 5 weeks of Met-restriction in HCT116 cells and C57BL/6J liver, respectively. n=5, error bars represent SD, *p<0.05 (Welch's t-Test). (*B-C*) Plots illustrating global, relative 5mC DNA methylation levels during Met-restriction. n≥3, error bars represent SD, *p<0.05 (Welch's t-Test) (*D-E*) Hierarchichal clustered heatmaps of LC-MS/MS generated $\log_2$ fold-change stoichiometric histone H3 peptide proteoform values relative to 0-hour or chow diet controls in HCT116 cells and C57BL/6J liver, respectively. n≥3, *p<0.05, **p<.01 (Welch's t-Test). See also Table S1 and Figure S1.

Figure 2. Decreased SAM Availability Drives Robust Histone Methylation Response

(*A*) Pathway diagram illustrating experimental approaches to depleting intracellular SAM availability. (*B*) Bar graph of relative *MAT2A* mRNA abundance in HCT116 cells as measured by RT-qPCR. n≥2, error bars represent SD, *p<0.05 (Welch's t-Test). (*C-D*) Dot plots of relative Met and SAM levels in HCT116 cells. n≥4, *p<0.05 (Welch's t-test). (*E-F*) Correlation plot of LC-MS/MS generated $\log_2$ fold-change stoichiometric values for individual histone H3 peptide proteoforms. n=3. (*G*-J) Bar graphs illustrating LC-MS/MS generated $\log_2$ fold-changes for individual H3K9 PTMs. n=3 *p<0.05 (Welch's t-test) See also Table S1, Table S2, and Figure S2.

Figure 3. H3K9me1 Abundance under SAM Depletion is Driven by *de novo* Methylation

(*A-B*) Bar graphs illustrating LC-MS/MS generated $\log_2$ fold-changes for individual H3K9 PTMs in HCT116 cells. n≥2, error bars represent SD, *p<0.05 (Welch's t-Test). (*C*) Bar graph illustrating percent loss in global H3K9me1, as measured by LC-MS/MS, in response to 5μm UNC0642 treatment. n≥2, error bars represent SD, *p<0.05 (Welch's t-Test) (*D*) Scatter plot of relative, total H3 and H3K9me1 cytoplasmic protein abundance in HCT116 cells as measured by western blot. Error bars represent SD, n=3, *p<0.05 (Welch's t-Test). (*E*) Representative western blot images of those used for the quantification presented in Figure 3E. Alpha-Tubulin is used as a cytoplasm marker to confirm efficient sub-cellular fractionation. See Document S1 for REVERT Total Protein Stain images used for normalization. (*F*) Diagram illustrating the

SILAC experimental design. (*G*) Bar graph illustrating contribution of newly synthesized histone H3 to the total chromatin bound pool after 24 hours of Met-restriction in HCT116 cells as measured by LC-MS/MS. n≥2, error bars represent SD, *p<0.05 (Welch's t-test). (*H*) Bar graph illustrating ratio of light:heavy L-Arg isotope presence in H3K9 proteoforms after 24 hours of Met-restriction in HCT116 cells as measured by LC-MS/MS. n≥2, Error bars represent SD, *p<0.05 (Welch's t-Test). (*I*) Pie chart depicting sources of H3K9me1 during SAM depletion. (*J*) Scatter plot of 373 cytoplasmic proteins identified by LC-MS/MS with significant incorporation of L-Arg $_{+0}$ that were also significantly upregulated after 24 hours of Met-restriction in 0.1% DMSO treated HCT116 cells. n=4, Statistical significance of p<0.05 (Welch's t-Test). See also Table S1, Table S3, and Figure S3.

Figure 4. H3K9me1 is Redistributed Over Repetitive and Transposable Elements During SAM Depletion

(*A-B*) Bar graphs illustrating percent changes in H3K9 PTM peak number at various annotated loci after 24 hours of Met-restriction in HCT116 cells. (*C-D*) Plots depicting changes in normalized H3K9 PTM ChIP-sequencing reads to loci of lost H3K9me3 after 24 hours of Met-restriction in HCT116 cells. (*E*) Bar graphs illustrating percent changes in loci annotation for regions of lost H3K9me3 experiencing increased relative to decreased H3K9me1 ChIP read enrichment after 24 hours of Met-restriction in HCT116 cells. (*F*) Representative ChIP-sequencing track image.

Figure 5. *De novo* H3K9 Mono-methylation Preserves Heterochromatin Stability

(*A*) Bar graph illustrating the percent abundances of 5 distinct nucleosome species following MNase digestion of HCT116 cells. n≥3, error bars represent SD, *p<0.05 (Welch's t-Test). (*B*) Representative DNA agarose gel image of those used for the quantification presented in Figure 5A. (*C-E*) Bar graphs illustrating relative mRNA abundances of *LINE* 1, *HERV-K,* and *HERV-R* in HCT116 cells as measured by RT-qPCR. n≥4, error bars represent SD, *p<0.05 (Welch's t-Test). See also Table S2.

Figure 6. Adaptive H3K9 Mechanisms Support Epigenetic Persistence Upon Metabolic Recovery

(*A*) Diagram illustrating the experimental design for replenishing Met and SAM availability after

24 hours of Met-restriction coupled with acute 5 μM UNC0642 or 0.1% DMSO treatment in

HCT116 cells. (*B*) Heatmap of $\log_2$ fold-change stoichiometric values for histone H3 and H4

peptides as measured by LC-MS/MS. n=4, *p<0.05, **p<0.01 (Welch's t-Test). (*C-E*) Bar graphs

illustrating $\log_2$ fold-changes for individual H3K9 PTMs in HCT116 cells as measured by LC-

MS/MS. n=4, error bars represent SD, *p<0.05 (Welch's t-Test). (*F*) PCA diagram of TPM

values. (*G*) EBseq multiple condition comparison diagram. (*H*) Pie chart illustrating percent

allocation of genes identified via RNA-sequencing to all EBseq patterns. (*I-J*) Scatter plots

illustrating NES and FDR q-value for GSEA identified GO terms. (*K-L*) Representative GSEA

running enrichment score figures. See also Table S1 and Figure S4.

<u>Figure 7. Adaptive and Persistent Epigenetic Responses to SAM Depletion are Robust *in vivo*,</u>

<u>Independent of Age</u>

(*A*) Diagram depicting the experimental design for *in vivo* Met-restriction of 6- and 22-month

C57BL/6J mice followed by dietary Met-reintroduction. (*B*) Heatmap of $\log_2$ fold-change

stoichiometric values for histone H3 and H4 peptides in C57BL/6J liver relative to age-matched

controls as measured by LC-MS/MS. n$\geq$5, *p<0.05, **p<0.01 (Welch's t-Test). (*C*) Bar graph

illustrating $\log_2$ fold-changes for individual H3K9 PTMs in liver from Met-restricted C57BL/6J

mice relative to age-matched controls as measured by LC-MS/MS. n$\geq$5, error bars represent

SD, *p<0.05 (Welch's t-Test). (*D-E*) Bar graphs illustrating relative mRNA abundances of *LINE*

1, *SINE B1*, and *SINE B2* in C57BL/6J liver as measured by RT-qPCR. n$\geq$5, error bars

represent SD, *p<0.05 (Student's t-Test). (*F*) Bar graph illustrating $\log_2$ fold-changes for

individual H3K9 PTMs in liver from C57BL/6J relative to age-matched controls as measured by

LC-MS/MS. n$\geq$5, error bars represent SD, *p<0.05, (Welch's t-Test). (*G-H*) Bar graphs

illustrating relative mRNA abundances of *LINE 1*, *SINE B1*, and *SINE B2* in C57BL/6J liver as

measured by RT-qPCR. n$\geq$5, error bars represent SD, *p<0.05 (Student's t-Test). See also

Table S1, Table S2, and Figure S5.

Supplemental Figure Titles and Legends

Figure S1, Related to Figure 1

(*A-D*) Scatter plots of relative abundance values for key Met-cycle metabolites in HCT116 cells as measured by LC-MS. n=5, error bars represent SD, *p<0.05 (Welch's t-Test). (*E-J*) Correlation plots of LC-MS/MS generated $\log_2$ fold-change stoichiometric values for individual histone H3 peptide proteoforms. n≥3. See also Table S1.

Figure S2, Related to Figure 2

(*A*) Diagram illustrating the experimental design for determining the conservation of H3K9 regulation in response to SAM depletion across various biological systems. (*B-F*) Bar graphs illustrating $\log_2$ fold-changes for individual H3K9 in response to *in vitro* SAM depletion. n=3, error bars represent SD, *p<0.05 (Welch's t-Test). See also Table S1.

Figure S3, Related to Figure 3

(*A-C*) Scatter plots of relative HMT mRNA abundance in HCT116 cells as measured by RT-qPCR. n=4, error bars represent SD, *p<0.05 (Welch's t-Test). (*D*) Heatmap of $\log_2$ fold-change stoichiometric values for unique histone H3K9 peptide proteoforms in HCT116 cells. n=4, *p<0.05 (Welch's t-Test). (*E*) Scatter plot of relative cell counts for attached HCT116 cells. n=4, error bars represent SD, *p<0.05 (Welch's t-Test). (F) Bar graph of relative H3K9me1 abundance in HCT116 cells as measured by western blot with accompanied representative blot image. See Document S1 for REVERT Total Protein Stain image used for normalization. Error bars represent SD, n≥3, *p<0.05 (Welch's t-Test). (*G*) Bar graphs illustrating $\log_2$ fold-changes for individual H3K9 PTMs. n≥3, *p<0.05 (Welch's t-Test). (*H*) Bar graph showing average incorporation of L-Arg$_{+10}$ in H3K9 peptide proteoforms. n=4, error bars represent SD. (*I*) Density plot illustrating the percentage of Met residue frequency in the primary amino acid sequence of all identified cytoplasmic proteins as well as those determined to be preferentially translated. See also Table S1, Table S2, and Table S3.

Figure S4, Related to Figure 6

(*A*) Bar graphs illustrating $\log_2$ fold-changes for individual H3K9 PTMs 24 hours following

UNC0642 removal. n=4, error bars represent SD, *p<0.05 (Welch's t-Test). *(B-C)* Bar graphs

illustrating relative recombinant enzyme activity in the presence of saturating amounts

UNC0642 and following 1000x or 10,000x overnight dialysis. n=4, error bars represent SD,

*p<0.05 (Welch's t-Test).

Figure S5, Related to Figure 6

(*A*) GSEA heatmap of relative gene expression across positive NES GO terms for Pattern 2. (B)

GSEA set-to-set heatmap illustrating gene enrichment overlap across positive NES GO terms

for Pattern 2. (*C*) GSEA heatmap of relative gene expression across negative NES GO terms

for Pattern 2. (D) GSEA set-to-set heatmap illustrating gene enrichment overlap across negative

NES GO terms for Pattern 2.

Figure S6, Related to Figure 7

(*A-E*) Bar graphs illustrating $\log_2$ fold-changes for individual H3K4, H3K18/K23, H3K27, H3K36,

and H4 PTMs in liver from Met-restricted C57BL/6J mice relative to age-matched controls as

measured by LC-MS/MS. n≥5, error bars represent SD, *p<0.05 (Welch's t-Test). (F) Bar graph

illustrating changes in fat, lean mass, and total body mass (BW) after 3 weeks of control or Met-

restricted diet feeding. n=14, error bars represent SEM, *p<0.05 (two-way ANOVA). (*G*)

Glucose tolerance test in 6- and 22-month C57BL/6J mice after 2 weeks of control or Met-

restricted diet feeding. n=14, error bars represent SEM, *p<0.05 (two-way ANOVA). (*H*) Bar

graph illustrating changes in fat, lean mass, and total body mass (BW) after 5 weeks of control

diet feeding following either 3 weeks of control or Met-restricted diet feeding. n≥6, error bars

represent SEM, *p<0.05 (two-way ANOVA). (*I*) Glucose tolerance test in 6- and 22-month

C57BL/6J mice after 4 weeks of control diet feeding following either 3 weeks of control or Met-

restricted diet feeding. n≥7, error bars represent SEM, *p<0.05 (two-way ANOVA). See also

Table S1 and Table S2.

Star Methods Text

Contact for Reagent and Resource Sharing:

Further information and requests for resources and reagents should be directed to and will be

fulfilled by the Lead Contact, John M. Denu (john.denu@wisc.edu).

<u>Experimental Model and Subject Details</u>

Animals and Diets:

All experiments involving animals were approved by the Institutional Animal Care and Use

Committee of the William S. Middleton Memorial Veterans Hospital, Madison WI. To test the

effect of Met deprivation in the context of young mice, 9-week-old male C57BL/6J mice were

purchased from The Jackson Laboratory (Bar Harbor, ME, USA) and placed on either an amino

acid defined Control diet (TD.01084, Envigo, Madison, WI, USA) or a Met-deficient diet

(TD.140119, Envigo, Madison, WI, USA) their respective diets at 14 weeks of age; full diet

compositions for these diets have been previously described (PMID: 29401631). After 5 weeks

on the diet, mice were sacrificed following an overnight fast and liver was flash frozen for

analysis as described.

To compare the effect of Met restriction on young and old mice, C57BL/6J. Nia male mice were

obtained from the National Institute on Aging Aged Rodent Colony, and housed in our animal

facility for 3-4 months until reaching 5 and 21 months of age. All animals were then placed on

the Control diet (TD.01084) for 4 weeks, until the mice were 6 and 22 months of age. Mice were

then randomized to either remain on the Control diet, or were placed on the Met-deficient diet

(TD.140119) for three weeks. Liver was collected and flash-frozen in liquid nitrogen harvested

from mice euthanized after an approximately 16-hour fast.

Cell Lines:

HCT116, HEK-293, HEK-293T, HMCF7, Panc1, HepG2, and HepA1 cell lines were cultured at

37_oC with 5.0% CO_2.

<u>Method Details</u>

Metabolite Extraction:

For HCT116 cells, ~1.5x10^6 cells were quickly rinsed 3 times with 1.5 ml ice-cold PBS pH 7.4

before the addition of 1.5 ml of -80$_{\circ}$C 80:20 MeOH:H_2O extraction solvent. Cells were incubated

with extraction solvent at -80$_{\circ}$C for 15 minutes. Next, cells were scraped off their dish and

transferred into a 2 ml microcentrifuge Eppendorf tube which was followed by a 5-minute,

maximum speed centrifugation at 4$_{\circ}$C. The supernatant was transferred to a new 2 ml

microcentrifuge Eppendorf tube. Next, 0.5 ml extraction solvent was added to the remaining

pellet, vortexed, and again centrifugated at maximum speed for 5 minutes at 4$_{\circ}$C. The 2

supernatants from each extraction were pooled and dried completely using a Thermo Fisher

Savant ISS110 SpeedVac. Dried metabolite extracts were resuspended in 80 μl of 85%

acetonitrile following microcentrifugation for 5 minutes at maximum speed at 4$_{\circ}$C to pellet any

remaining insoluble debris. The supernatant was then transferred to a glass vial for LC-MS

analysis.

For *M. musculus* liver tissue, an average tissue weight of 75 mg was powdered in liquid nitrogen

using a mortar and pestle. Powdered tissue was transferred to an individual 1.5 ml

microcentrifuge Eppendorf tube and incubated with 1 ml -80$_{\circ}$C 80:20 MeOH:H_2O extraction

solvent on dry ice for 5 minutes post-vortexing. Tissue homogenate was centrifugated at

maximum speed for 5 minutes at 4$_{\circ}$C. Supernatant was transferred to a 15 ml tube after which

the remaining pellet was resuspended in 0.8 ml -20$_{\circ}$C 40:40:20 ACN:MeOH:H_2O extraction

solvent and incubated on ice for 5 minutes. Tissue homogenate was again centrifugated at

maximum speed for 5 minutes at 4$_{\circ}$C after which the supernatant was pooled with the

previously isolated metabolite fraction. The 40:40:20 ACN:MeOH:H_2O extraction was then

repeated as previously described. Next, 550 μl of pooled metabolite extract for each sample

was transferred to a 1.5 ml microcentrifuge Eppendorf tube and completely dried using a

Thermo Fisher Savant ISS110 SpeedVac. Dried metabolite extracts were resuspended in 450 μl

of 85% ACN following microcentrifugation for 5 minutes at maximum speed at 4$_{\circ}$C to pellet any

remaining insoluble debris. Supernatant was then transferred to a glass vial for LC-MS analysis.

LC-MS Metabolite Analysis:

Each prepared metabolite sample was injected onto a Thermo Fisher Scientific Vanquish

UHPLC with a Waters XBridge BEH Amide column (100 m x 2.1 mm, 3.5 μm) coupled to a

Thermo Fisher Q-Exactive mass spectrometer. Mobile phase (A) consisted of 97% H_2O, 3%

ACN, 20 mM ammonium acetate, and 15 mM ammonium hydroxide pH 9.6. Organic phase (B)

consisted of 100% ACN. Metabolites were resolved using the following linear gradient: 0 min,

85% B, 0.15 ml/min; 1.5 min, 85% B, 0.15 ml/min; 5.5 min, 40% B, 0.15 ml/min; 10 min, 40% B,

0.15 ml/min; 10.5 min, 40% B, 0.3 ml/min; 14.5 min, 40% B, 0.3 ml/min; 15 min, 85% B, 0.15

ml/min; 20 min, 85% B, 0.15 ml/min. The mass spectrometer was operated in positive ionization

mode with a MS1 scan at resolution = 70,000, automatic gain control target = 3×10^6, and scan

range = 60-186 m/z and 187-900 m/z. This protocol was adapted from Mentch et al., 2015.

Individual metabolite data were called using MAVEN (Melamud et al., 2010; Clasquin et al.,

2012) with retention times empirically determined in-house. Peak Area Top values were

analyzed to determine metabolite expression.

1001

1002   *LC-MS DNA Methylation Assay:*

1003

500 ng of cytosine, 5mC, and 5hmC double-stranded DNA standards from Zymo Research

were incubated with Zymo Research DNA Degradase Plus for 3 hours at 37_oC following enzyme

inactivation at 70_oC for 20 minutes. 75 μl of ACN was added to each standard reaction to bring

the final volume to 100 μl. DNA standards were centrifugated at 21,100xg for 5 minutes at 4_oC

after which 75 μl of supernatant was transferred to a glass vial for LC-MS analysis. A range of

0.1 to 100 ng of digested DNA was analyzed via LC-MS to determine each nucleoside's linear

range of detection.

1011

1012   Genomic DNA extracted using a Promega Wizard Genomic DNA Purification Kit after which 1

1013 μg was incubated with Zymo Research DNA Degradase Plus at 37_oC following enzyme

1014   inactivation at 70_oC for 20 minutes. 175 μl of ACN was added to each reaction to bring the final

1015   volume to 200 μl. DNA samples were centrifugated at 21,100xg for 5 minutes at 4_oC after which

1016   150 μl of supernatant was transferred to a glass vial for LC-MS analysis.

1017

Each prepared nucleoside sample was injected onto a Thermo Fisher Scientific Vanquish

UHPLC with a Waters XBridge BEH Amide column (100 m x 2.1 mm, 3.5 μm) coupled to a

Thermo Fisher Q-Exactive mass spectrometer run in positive ionization mode. Mobile phase (B)

consisted of H_2O, 3% ACN, 20 mM ammonium acetate, and 15 mM ammonium hydroxide pH

9.6. Organic phase (B) consisted of 100% ACN. Nucleosides were resolved using the following

linear gradient: 0 min, 85% B, 0.15 ml/min; 1.5 min, 85% B, 0.15 ml/min; 5.5 min, 40% B, 0.15

ml/min; 10 min, 40% B, 0.15 ml/min; 10.5 min, 40% B, 0.3 ml/min; 14.5 min, 40% B, 0.3 ml/min;

15 min, 85% B, 0.15 ml/min; 20 min, 85% B, 0.15 ml/min. The mass spectrometer was operated

in positive ionization mode with a MS1 scan at resolution = 70,000, automatic gain control target

= $3 \times 10_6$, and scan range = 200-300 m/z, followed by a DDA scan with a loop count of 5. DDA

settings were as follows: window size = 2.0 m/z, resolution = 17,500, automatic gain control

target = $1 \times 10_5$, DDA maximum fill time = 50 ms, and normalized collision energy = 30. This

protocol was adapted from Mentch et al., 2015. Individual metabolite data were called using

MAVEN (Melamud et al., 2010; Clasquin et al., 2012) with retention times empirically

determined in-house. Peak Area Top values were analyzed to determine metabolite expression

with dAdenosine being utilized for internal normalization (m/z dCytosine: 228.0978, m/z d5mC:

242.1135, m/z d5hmC: 258.108, m/z dAdenosine: 252.109).

1035

1036 *Histone Isolation and Chemical Derivatization:*

1037

Tissue culture cell pellets were resuspended in 800 μl of ice-cold Buffer A (10 mM Tris-HCl pH

7.4, 10 mM NaCl, and 3 mM $MgCl_2$) supplied with protease and histone deacetylase inhibitors

(10 μg/ml leupeptin, 10 μg/ml aprotinin, 100 μM phenylmethylsulfonyl fluoride, 10 mM

nicotinamide, 1 mM sodium-butyrate, and 4 μM trichostatin A) followed by 80 strokes of light

pestle homogenization in a 1 ml Wheaton dounce homogenizer. Cell homogenate was then

transferred to a 1.5 ml microcentrifuge Eppendorf tube and centrifugated at 800xg for 10

minutes at 4_oC to pellet nuclei. The supernatant was either transferred to a fresh 1.5 ml

Eppendorf tube or discarded. The nuclei pellet was resuspended in 500 μl ice-cold PBS pH 7.4

followed by centrifugation at 800xg for 10 minutes at 4_oC. The supernatant was discarded and

nuclei were again washed with 500 μl ice-cold PBS pH 7.4. Pelleted nuclei were then

resuspended in 500 μl of 0.4N H_2SO_4 and rotated at 4_oC for 4 hours. Samples were

centrifugated at 3,400xg for 5 minutes at 4_oC to pellet nuclear debris and precipitated non-

histone proteins. The supernatant was transferred to a new 1.5 ml Eppendorf tube after which

125 μl of 100% trichloroacetic acid was added and incubated overnight on ice at 4_oC. Samples

were centrifugated at 3,400xg for 5 minutes at 4_oC to pellet precipitated histone proteins. The

supernatant was discarded, after which the precipitant was washed with 1 ml ice-cold acetone

+0.1% HCl. Samples were centrifugated at 3,400xg for 2 minutes at 4_oC and the supernatant

was discarded. This process was repeated except precipitant was washed with 100% ice-cold

acetone. Residual acetone was allowed to evaporate at room temperature for 10 minutes after

which the dried precipitant was dissolved in 125 μl H_2O. Samples were then centrifugated at

21,100xg for 5 minutes at 4$_o$C to pellet any remaining insoluble debris. Supernatant containing

purified histone was then transferred to a new 1.5 ml Eppendorf tube and stored at -20$_o$C until

needed for future analyses.

1061

For *M. musculus* liver tissue, an average tissue weight of 50 mg was homogenized in 800μl ice-

cold Buffer supplemented with protease and HDAC inhibitors (listed above) using a 1 ml

Wheaton dounce homogenizer (20 strokes of the loose pestle followed by 20 strokes of the tight

pestle) and strained through a 100 μM filter before being transferred to a new 1.5 ml Eppendorf

tube. All remaining steps for completing the histone isolation were performed as described

previously in the tissue culture isolation protocol.

1068

To prepare purified histone proteins for LC-MS/MS analysis, 5 μg of histone was diluted with

H$_2$O to a final volume of 9 μl. 1 μl of 1 M triethylammonium bicarbonate was added to each

sample to buffer the solution to a final pH of 7-9. Next, 1 μl of 1:100 propionic anhydride:H$_2$O

was added to each sample and allowed to incubate for 2 minutes at room temperature. The

propionylation reaction was then quenched via the addition of 1 μl 80 mM hydroxylamine which

was allowed to incubate for 20 minutes at room temperature. Next, propionylated histones were

digested with 0.1 μg trypsin for 4 hours at 37$_o$C. Upon completion of trypsin digestion, 5 μl of .02

M NaOH was added to bring the final pH between 9-10. Propionylated, trypsin digested histone

peptides were then N-terminally modified with 1 μl 1:50 phenyl isocyanate:ACN for 1-hour at

37$_o$C. Modified peptides were desalted and eluted-off of Empore™C18 extraction membrane.

Eluted samples were dried completely using a Thermo Fisher Scientific Savant ISS110

SpeedVac, resuspended in 40 μl sample diluent (94.9% H$_2$O, 5% ACN, .1% TFA), and

transferred into glass vials for LC-MS/MS analysis.

1082

1083 *Histone Proteomics Analysis:*

1084

Derivatized histone peptides were injected onto a Dionex Ultimate3000 nanoflow HPLC with a

Waters nanoEase UPLC C18 column (100 m x 150 mm, 3μm) coupled to a Thermo Fisher Q-

Exactive mass spectrometer at 700 nL/ min. Aqueous phase (A) consisted of H$_2$O + 0.1% formic

acid while the mobile phase (B) consisted of acetonitrile + 0.1% formic acid (B). Histone

peptides were resolved using the following linear gradient: 0 min, 2.0% B; 5 min, 2.0% B; 65

min, 35% B; 67 min, 95% B; 77 min, 95% B; 79 min, 2.0% B; 89 min, 2.0% B. Data was

acquired using data-independent acquisition (DIA) mode. The mass spectrometer was operated

with a MS1 scan at resolution = 35,000, automatic gain control target = 1×10^6, and scan range

= 390-910 m/z, followed by a DIA scan with a loop count of 10. DIA settings were as follows:

window size = 10 m/z, resolution = 17,500, automatic gain control target = 1×10^6, DIA

maximum fill time = AUTO, and normalized collision energy = 30.

1096

DIA Thermo .raw files were imported into Skyline open source proteomics software (MacLean et

al., 2010) and matched against a pre-constructed peptide database containing unique peptide

proteoforms from H3 and H4. The identity of each individually called peak was verified by hand

after which total MS1 peak area values were exported for downstream analyses. To control for

differences in ionization efficiencies across histone peptides, all peptides possessing an

identical primary sequence were analyzed as a peptide family which enabled the calculation of

stoichiometric values for each unique peptide proteoform within the family. Stoichiometric values

were compared across conditions to identify altered histone peptide abundances. Furthermore,

combinatorial peptide stoichiometry values were deconvoluted by totaling the stoichiometry

values for all peptides possessing a given PTM independent of any other PTMs that may be

present on other residues of the same primary peptide. A Welch's T-Test with a p-value cutoff of

0.05 was used to determine statistical significance.

1109

1110   Hierarchical clustering was performed in MATLAB R2016a using the 'clustergram' command.

1111   Pearson's Correlation Coefficients were calculated in MATLAB R2016a using the 'corr'

1112   command.

1113

1114   *RNAi Mediated Inhibition of MAT2A and H3K9 HMT Expression:*

1115

1116   For 35 mm tissue culture dishes, 1.5×10^5 HCT116 cells were reverse transfected with 30 pmol

1117   of targeted or control RNAi using 5 μl of Lipofectamine RNAiMAX in addition to 500 μl Opti-

1118   MEM and 2.5 ml RPMI-1640 +10% FBS. MAT2A-RNAi treated HCT116 cells used for LC-MS

1119   metabolite analysis were cultured using dialyzed FBS.

1120

1121

1122   *Methionine Restriction of Tissue Culture Cells:*

1123

All tissue culture cells were initially cultured in RPMI-1640 +10% FBS at $37_{o}C$, 5% CO_2. One

day before Met-restriction, $3.5 \times 10_6$ cells were seeded into 100 mm tissue culture dishes. One-

hour before the restriction began, RPMI-1640 was replaced with fresh, Met replete media. At the

time of Met-restriction, cells were rinsed 2x with 3 ml of sterile PBS pH 7.4 followed by addition

of RPMI-1640 void of Met. Any small molecule inhibitors or vehicle controls would be added at

this time.

1130

1131   For experiments in which Met-restriction was followed by Met repletion, HCT116 cells were

1132   washed with PBS pH 7.4 2x to maximize small molecule inhibitor and DMSO removal before

1133   reintroduction of Met replete RPMI-1640.

1134

1135   HCT116 cells cultured for LC-MS metabolite analysis were cultured in RPMI-1640 containing

1136   10% dialyzed FBS beginning 1-hour prior to beginning Met-restriction.

1137

1138   *SILAC in Tissue Culture:*

1139

197 ml of SILAC RPMI-1640 +10% DFBS void of L-Arg and L-Lys was supplemented with 50mg

isotopically heavy $3C_6 15N_4$ L-Arg HCl and 9.43mg isotopically light L-Lys to obtain final

concentrations of 1.149 mM and 0.219 mM, respectively. HCT116 cells were cultured in the

SILAC media for 7 days (or 9 doubling cycles at a reported doubling time of 21 hours) after

which cells were washed 3x with 5 ml PBS pH 7.4 before replacing the SILAC media with

isotopically light L-Arg and L-Lys RPMI-1640 +10% DFBS void of Met. Half of the cells were

treated with 5 μM UNC0642 or 0.1% DMSO at this time. Furthermore, 0-hour control HCT116

cells which were never exposed to isotopically light L-Arg, Met-restricted RPMI-1640 were

harvested. HCT116 cells cultured in the Met-restricted media were harvested after 24 hours.

1149

1150   *SILAC Shotgun Proteomics Analysis:*

1151

50 μg of cytoplasmic protein extract from the previously described SILAC Met-restriction

experiment were diluted with H_2O into a final volume of 75 μl. Next, dithiothreitol was added to

each sample at a final concentration of 10 mM, followed by a 15-minute incubation at $95_{o}C$. A

sonication water bath was used to resolubilize precipitated proteins post-heat denaturation.

Iodoacetamide was then added to a final concentration of 20 mM, followed by a 20-minute

incubation at room temperature protected from light. Alkylated proteins were precipitated with

337 μl ice-cold acetone for 20 minutes at -20$_o$C. Samples were centrifugated at 21,000xg for 15

minutes at 4$_o$C, the supernatant was discarded, and precipitated protein was resuspended in

400 μl of urea buffer (2 M urea and 100 mM ammonium bicarbonate pH 8). Precipitated proteins

were again solubilized using a sonication water bath. Solubilized proteins were digested into

peptides using 0.8 μg trypsin at 37$_o$C overnight. Upon digest completion, samples were desalted

and eluted-off Empore™C18 extraction membrane. Eluted samples were dried completely using

a Thermo Fisher Scientific Savant ISS110 SpeedVac, after which being resuspended in 50 μl

sample diluent (94.9% H_2O, 5% ACN, .1% TFA) and transferred to glass vials for LC-MS/MS

analysis.

1167

Each prepared cytoplasmic peptide sample was injected onto a Dionex Ultimate3000 nanoflow

HPLC with a Waters nanoEase UPLC C18 column (100 m x 150 mm, 3μm) coupled to a

Thermo Fisher Q-Exactive mass spectrometer at 800 nL/ min. Aqueous phase (A) consisted of

H_2O + 0.1% formic acid while the mobile phase (B) consisted of acetonitrile + 0.1% formic acid

(B). Histone peptides were resolved using the following linear gradient: 0 min, 5.0% B; 5 min,

5.0% B; 120 min, 30% B; 122 min, 95% B; 127 min, 95% B; 129 min, 5.0% B; 145 min, 5.0% B.

Data was acquired using data-dependent acquisition (DDA) mode. The mass spectrometer was

operated with a MS1 scan at resolution = 70,000, automatic gain control target = 1 x 10$_6$, and

scan range = 400-1000 m/z, followed by a DDA scan with a loop count of 20. DDA settings were

as follows: window size = 2.0 m/z, resolution = 17,500, automatic gain control target = 2 x 10$_5$,

DDA maximum fill time = 150 ms, and normalized collision energy = 28.

1179

1180   *MaxQuant Proteomics Analysis:*

1181

1182   Thermo .raw DDA files were analyzed using MaxQuant v1.6.2.6 (Cox et al., 2014) using the

1183   default settings with the following exceptions: Labels: 1, Arg10; Instrument: Orbitrap; LFQ: LFQ

1184   min. ratio count: 2, LFQ min. number of neighbors: 3, LFQ average number of neighbors: 6;

1185   FASTQ File: Proteome: *H. Sapiens* Proteome ID UP000005640.

1186

Identified proteins with a Q-value of 0.01 or above were removed. For the remaining proteins,

LFQ light (L-Arg$_{+0}$) and heavy (L-Arg$_{+10}$) values were summed to generate LFQ total protein

abundance values. LFQ light values were divided by the corresponding LFQ total value to

generate a percent abundance of LFQ light species for each identified protein. Welch's t-Tests

with a p-value cut off of 0.05 were used to determine significant accumulation of LFQ light

species in Met-restricted HCT116 samples relative to 0-hour controls. For proteins determined

to have significant LFQ light species abundance after Met-restriction, LFQ total values were

used to determine the corresponding change in overall protein abundance. Up to 2 null-values

were removed from replicates of an experimental group to account for missed identifications. If

3 of the 4 replicates for a single experimental group contained null-values, that protein was

removed from all experiment groups and therefore excluded from any downstream analyses. A

Welch's t-Test with a p-value cut off of 0.05 was used to determine if an identified protein was

significantly more abundant after Met-restriction relative to 0-hour controls.

1200

1201 *SILAC Histone Proteomics Analysis:*

1202

Histones were isolated, chemically derivatized, and analyzed by LC-MS/MS as described in

"Histone Isolation and Chemical Derivatization" and "Histone Proteomics Analysis." DDA

Thermo .raw files of histone proteins with incorporated isotopically heavy $3C_6$ $15N_4$ L-Arg were

used to construct an H3K9/K14 peptide library using Skyline opensource proteomics software

(MacLean et al., 2010). DIA files were then matched against both isotopically heavy and light L-

Arg peptide libraries after which peptide identities were confirmed by hand. MS1 intensity values

for both isotopically heavy and light L-Arg peptides were summed to generate total MS1 peptide

proteoform intensity values which were then used to calculate the percent of total MS1 peptide

proteoform signal which was contributed by isotopically heavy peptide proteoforms. Welch's T-

Tests with a p-value cutoff of 0.05 were used to determine significant differences in total MS1

peptide proteoform stoichiometric values as well as the percent contribution of isotopically

heavy peptide proteoforms to total peptide proteoform intensity values.

1215

1216 *Chromatin Immunoprecipitation and Sequencing:*

1217

$30 \times 10_6$ HCT116 cells were resuspended in 1 ml MNase digestion buffer (50 mM Tris-HCl pH

7.6, 1 mM $CaCl_2$, and 0.2% Triton X-100) after which 330 μl of the cell suspension was

aliquoted into individual 1.5 ml tubes. Each tube was treated with either 45, 52.5, or 60 units of

Worthington MNase for 5 minutes at 37_oC shaking at 400 rpm to generate mono-nucleosomes.

MNase digestions were stopped via the addition of EDTA to a final concentration of 5mM.

Digested cells were then pooled and lysed at 4_oC using a Covaris S220 Focused-ultrasonicator

with the following settings: Peak Power = 75, Duty Factor = 5, Cycles/Burst = 100, Duration =

30" on/30" off x2. Insoluble debris was centrifugated at 10,000 rpm for 10 minutes at 4_oC after

which the supernatant containing prepared nucleosomes was transferred to a new tube.

Glycerol was added to a final concentration of 5% prior to -80$_o$C storage. For input controls, 43

μl of prepared nucleosomes were aliquoted and saved for further processing at -80$_o$C.

1229

Next, 860 μl of prepared nucleosomes was incubated with 5 μg of the desired antibody, rotating

overnight at 4$_o$C. The nucleosome/antibody solution was added to 50 μl of prepared M-280

Sheep Anti-Rabbit IgG superparamagnetic Dynabeads™ and allowed to rotate at 4$_o$C for 4

hours. Conjugated beads were then washed for 10 minutes, rotating at 4$_o$C under the following

conditions: 3x 1 ml RIPA (10 mM Tris pH 7.6, 1 mM EDTA, 0.1% SDS, 0.1% Na-deoxycholate,

1% Triton X-100) buffer, 2x 1 ml RIPA buffer +0.3 M NaCl, 2x 1 ml LiCl buffer (0.25 M LiCl,

0.5% NP-40, 0.5% Na-deoxycholate), 1x with 1 ml 1xTE +50 mM NaCl. Following the final

wash, beads were centrifugated at 4$_o$C for 3 minutes at 960xg after which the supernatant was

carefully removed and discarded.

1239

To elute immunoprecipitated nucleosome species, beads were resuspended in 210 μl of elution

buffer (50 mM Tris-HCl pH 8.0, 10 mM EDTA, 1% SDS) and incubated at 65$_o$C for 30 minutes

shaking at 400 rpm. Beads were resuspended every 2 minutes via brief vortexing. Next samples

were centrifugated at 16,000xg for 1 minute at room temperature after which 200 μl of

supernatant was transferred to a new 1.5 ml Eppendorf tube. Eluted nucleosome species were

then treated with 2 μl Thermo Fisher Scientific Proteinase K for 2 hours at 55$_o$C. DNA was

recovered from the Proteinase K digestion using the QIAquick PCR Purification Kit.

1247

1248 DNA libraries were prepared by the University of Wisconsin-Madison Biotechnology Center

1249 using the Illumina TruSeq DNA Nano kit and then sequenced across 4 lanes of an Illumina

1250 1x100 HiSeq2500 High Throughput flowcell.

1251

FASTQ files were aligned unpaired using bowtie2-2.1.0 (Langmead and Salzberg, 2012)

against human genome 19. Bowtie2 output SAM files were converted to BAM files, sorted and

indexed using samtools 1.5 (Li et al., 2009). Broad peaks were called from sorted BAM files

using MACS2-2.1.1.20160309 (Zhang et al., 2008) with a Q-value cutoff of 0.1. Input DNA was

used as the control file. All identified peaks possessing a log$_2$ fold-change below 1.75 were

discarded prior to downstream analyses. Bedtools v2.25.0 (Quinlan and Hall, 2010) was used to

identify unique and overlapping across samples and also used to assign raw mapped reads to

pre-specified genomic loci. Genomic loci were annotated using HOMERv4.9 (Heinz et al.,
2010).

1261

1262 *Quantitative PCR:*

1263

1264 RNA was extracted from HCT116 cells using Thermo Fisher Scientific TRIzol and from liver
1265 tissue using Sigma TRI Reagent, both by following the manufacturer's instructions. The
1266 concentration and purity of RNA were determined by absorbance at 260/280 nm.

1267

1268 cDNA was prepared from 1 μg of RNA using RevertAid Reverse Transcriptase for HCT116 RNA
1269 and Superscipt III for liver RNA. Oligo dT primers were used in both cDNA synthesizing
1270 reactions.

1271

1272 HCT116 cDNA was analyzed on a Roche 480 Light Cycler System with PerfeCTa SYBR Green
1273 SuperMix while cDNA generated from liver was analyzed on an Applied Biosystems StepOne
1274 Plus instrument with Sybr Green PCR Master Mix. HCT116 cDNA qPCR reactions were
1275 normalized internally using GAPDH while liver cDNA qPCR reactions were normalized internally
1276 using actin. Primer sequences are available in Table S3.

1277

1278 *RNA sequencing*

1279

RNA libraries were prepared by Novogene using NEBNext adaptors followed by paired end
2x150 sequencing on their Illumina Platform. FASTQ files were trimmed using Trimmomatic
v0.39 (Bolger et al., 2014) and aligned using RSEM v1.2.4 (Li et al., 2011) against the human
genome 18 reference transcriptome. RSEM .genes output files were used for Ebseq
EBMultiTest multiple comparison analysis (Leng et al., 2013). Gene lists associated with
individual EBMultiTest patterns were further analyzed with GSEA 2 (Tamayo et al., 2005;
Mootha et al., 2003)

1287

1288 *MNase Accessibility Assay:*

1289

$6x10^6$ HCT116 cells were resuspended in 1 ml of digestion buffer (50 mM Tris-HCl pH 7.6, 1
mM $CaCl_2$, and 0.2% Triton X-100) in addition to 15 unites of Worthington MNase. MNase
treated cell suspensions were incubated for 5 minutes at 37_oC shaking at 400 rpm. MNase

digestions were stopped via the addition of EDTA to a final concentration of 5 mM. Digested

cells were then pooled and lysed at 4_oC using a Covaris® S220 Focused-ultrasonicator with the

following settings: Peak Power = 75, Duty Factor = 5, Cycles/Burst = 100, Duration = 30" on/30"

off x2. Insoluble debris was pelleted via centrifugation at 10,000 rpm for 10 minutes at 4_oC after

which the supernatant was transferred to a new tube. 150 μl of digested chromatin was treated

with 2 μl of Thermo Fisher Scientific Proteinase K for 90 minutes at 55_oC. DNA was then

purified using the QIAquick PCR Purification Kit. Next, 1 μg of purified DNA was separated on a

2% agarose gel and quantified using GelAnalyzer 2010a.

1301

1302 *Western-Blotting and Image Quantification:*

1303

Protein separated on SDS-PAGE gels were transferred onto 0.2 μm nitrocellulose membrane

and stained with 5 ml of LI-COR REVERT Total Protein Stain for 5 minutes at room temperature

before being blocked in 0.1% fat-free milk PBS pH 7.4 for 1 hour. Primary antibodies were

diluted 1:1000 in 5% fat-free milk PBST pH 7.4 and incubated with the membrane overnight at

4_oC. LI-COR secondary antibodies were used at a 1:15,000 dilution in 5% fat-free milk PBST pH

7.4 and incubated with the membrane for 1 hour at room temperature. Membranes were imaged

using an Odyssey Infrared Imager (model no. 9120). Densitometry was performed using Image

Studio™ Lite software. All data was normalized to REVERT™ Total Protein Stain.

1312

1313 *In vivo Tests:*

1314

1315 For glucose tolerance tests, mice were fasted overnight for 16 hours and then injected with

1316 glucose (1 g/kg) intraperitoneally as previously described (Arriola Apelo et al., 2016). Glucose

1317 measurements were taken using a Bayer Contour blood glucose meter and test strips. Mouse

1318 body composition was determined using an EchoMRI Body Composition Analyzer (EchoMRI$_{TM}$,

1319 Houston, TX, USA) according to the manufacturer's procedures.

1320

1321 *Protein Purification:*

1322

1323 The *H. sapiens* Ehmt1 catalytic subunit *E. coli* expression plasmid used in this study (Addgene

1324 plasmid # 51314) was acquired from addgene, made possible through a gift by Cheryl

1325 Arrowsmith. The *H. sapiens* Ehmt2 catalytic subunit *E. coli* expression plasmid used in this

1326 study was provided by the laboratory of Peter W. Lewis at the University of Wisconsin-Madison.

1327

Transformed Rosetta™ *E. coli* competent cells were cultured in 1L of 2XYT media at 37$_\circ$C to an

O.D. 600 of 0.8. IPTG was then added to each culture at a final concentration of 1mM. Cultures

were allowed to grow for 16 hours at 18$_\circ$C before being harvested and stored at -80$_\circ$C. Pellets

were resuspended in 30ml Buffer A (50 mM NaPi, 250 mM NaCl, 10mM imidazole, pH 7.2) and

lysed using sonicated in the presence of lysozyme. Lysate was centrifugated and the

supernatant was collected. The supernatant was then loaded onto a HisTrap FF nickel column

in line with a GE AKTA FPLC. Protein was eluted off the column using a linear gradient ending

in 100% Buffer B (50 mM NaPi, 250 mM NaCl, 250 mM imidazole, pH 7.2) and collected into

time-dependent fractions. Fractions were analyzed by coomassie staining of SDS-PAGE gels to

assess purity.

1338

1339 *Histone Methyltransferase Activity Assay:*

1340

The methyltransferase activity of recombinantly purified Ehmt1 and Ehmt2 catalytic subunits

was assessed using Promega Corporation's MTase-Glo™ Methyltransferase Assay. Assays

were performed using 12.5 nM enzyme in the presence of saturating concentrations of SAM (30

μM) and unmodified H3K9 peptide (30 μM, WARTKQTARKSTGGKAPR – 3') as well as either

saturating levels of UNC0642 (5 μM) or 0.1% DMSO. Reactions proceeded for 1 hour at 30$_\circ$C

before addition of MTase-Glo™ reagent following the Promega protocol. Luminescence was

detected using a Biotek Synergy H4 Hybrid plate reader. All values detected were determined to

be within the instrument's linear range of detection via the use of a SAH standard curve. To test

for reversibility of enzyme inhibition by small molecule, 1 μM Ehmt1 and Ehmt2 were incubated

with saturating concentrations of UNC0642 (5 μM) or 0.1% DMSO followed overnight 1,000x or

10,000x dialysis with a molecular weight cutoff of 10,000 Da against 1x Reaction Buffer without

BSA (20mM Tris, pH 8.0, 50mM NaCl, 1mM EDTA, 3mM MgCl$_2$, 1mM DTT). Following dialysis,

enzyme and reaction components were diluted using the appropriate dialysis buffers to the

working concentrations described above. The MTase-Glo™ assay was then repeated using

identical conditions as described above.

1356

1357 *Quantification and Statistical Analysis*

1358

1359 Explanations of statistical analyses and parameters are present in the main and supplemental

1360 figure legends as well as in 'Method Details" where appropriate.

1361

1362 *Data and Software Availability*

1363

1364 RNA-sequencing as well as H3K9me1 and H3K9me3 ChIP-sequencing data have been

1365 uploaded to Gene Expression Omnibus (GEO) and will be made publicly available upon

1366 manuscript publication.

1367

1368 **KEY RESOURCES TABLE**

1369

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Rabbit polyclonal anti-H3K9me1	Abcam	Cat#ab176880; RRID#AB_2751009
Rabbit polyclonal anti-H3K9me3	Abcam	Cat#ab8898; RRID#AB_306848
Rabbit polyclonal anti-Histone H3	Abcam	Cat#ab46765; RRID#AB_880439
Rabbit monoclonal anti-alpha Tubulin	Cell Signaling Technologies	Cat#cst2125; RRID#AB_2619646
IRDye® 800CW Goat anti-Rabbit	LI-COR Biosciences	Cat#925-32211; RRID#AB_2651127
Chemicals, Peptides, and Recombinant Proteins		
UNC0642 G9a/GLP Inhibitor	R&D Systems	Cat#5132/10
DNA Degradase Plus	Zymo Research	Cat#E2020
Lipofectamine RNAiMax	Thermo Fisher	Cat#13778030
L-Lysine-2HCl	Thermo Fisher Scientific	Cat#PI89987
L-Arginine-HCl 13C6 15N4	Thermo Fisher	Cat#PI89990
Dimethyl Sulfoxide (DMSO)	Sant Cruz Biotechnology	Cat#sc-358801
Sequencing Grade Modified Trypsin	Promega	Cat#V5113
Propionic Anhydride	Sigma-Aldrich	Cat#240311
Phenyl Isocyanate	Sigma-Aldrich	Cat#78750-25ML
Leupeptin Hemisulfate	Thermo Fisher	Cat#AAJ61188MC
Aprotinin	Gold Biotechnology	Cat#A-655-100
Phenylmethylsulfonyl Fluoride (PMSF)	Gold Biotechnology	Cat#P-470-50
Trichostatin A	Thermo Fisher	Cat#501012404
Sodium Butyrate	Sigma-Aldrich	Cat#303410-100G
Nicotinamide	Sigma-Aldrich	Cat#N3376-500G
Urea	Acros Organics	Cat#197460050
Dithiothreitol (DTT)	Gold Biotechnology	Cat#DTT100
Iodoacetamide	Sigma-Aldrich	Cat#I1149-5G

Ammonium Bicarbonate	Sigma-Aldrich	Cat#09830-500G
Micrococcal Nuclease (MNase)	Worthington Biochemical	Cat#LS004798
Proteinase K	Thermo Fisher	Cat#EO0491
TRIzol™	Thermo Fisher	Cat#15596018
TRI Reagent	Sigma-Aldrich	Cat#T9424-25ML
3M Empore C18 47mm extraction disks	Empore	Cat#2215
Atlantis dC18 NanoEase Column (3 μM, 100 μM, 150 mm)	Waters	Cat#186002209
XBridge BEH Amide Column (3.5 μM, 2.1 mm, 150mm)	Waters	Cat#186004860
Superscript III Reverse Transcriptase	Thermo Fisher	Cat#18080044
RPMI-1640	Thermo Fisher	Cat#11875119
RPMI-1640 for SILAC	Thermo Fisher	Cat#88365
RPMI-1640, no methionine	Thermo Fisher	Cat#A1451701
Fetal Bovine Serum (FBS)	Thermo Fisher	Cat#16000044
Dialyzed Fetal Bovine Serum (DFBS)	Thermo Fisher	Cat#88212
H3K9 unmodified peptide	This paper	N/A
pET28a-LIC-4H4H	Addgene	Plasmid # 51314; RRID#Addgene_51314
pET28a-LIC-EHMT2-913-1193	Peter W. Lewis Laboratory	N/A
Critical Commercial Assays		
PerfeCTa SYBR® Green SuperMix	Quanta Bio	Cat#95054-02K
SYBR Green PCR Master Mix	Thermo Fisher	Cat#4309155
REVERT Total Protein Stain	LI-COR Biosciences	Cat#926-11011
RevertAid First Strand cDNA Synthesis Kit	Thermo Fisher	Cat#K1621
Wizard® Genomic DNA Purification Kit	Promega	Cat#A1120
Dynabeads™ M-280 Sheep Anti Rabbit IgG	Thermo Fisher	Cat#11203D
QIAquick PCR Purification Kit	QIAGEN	Cat#28104
TruSeq DNA Nano	Illumina	Cat#20015964
Axygen™ Light Cycler 480 Compatible PCR Plates	Thermo Fisher	Cat#14-223-212
Promega MTase-Glo™	Promega	Cat#V7602
Deposited Data		

H3K9me1 and H3K9me3 ChIP-sequencing Data	This paper	GEO Accession Number: Available upon manuscript publication
RNA-sequencing Data	This paper	GEO Accession Number: Available upon manuscript publication
Experimental Models: Cell Lines		
H. sapiens: HCT116	ATCC	CCL-247
H. sapiens: HEK-293T	ATCC	CRL-3216
H. sapiens: HEK-293	ATCC	CRL-1573
H. sapiens: HepG2	ATCC	HB-8065
H. sapiens: MCF7	ATCC	HTB-22
H. sapiens: Panc1	ATCC	CRL-1469
M. musculus: Hepa-1c1c7	ATCC	CRL-2026
Experimental Models: Organisms/Strains		
M. musculus: C57BL/6J (9-weeks of age)	The Jackson Laboratory	JAX: 000664
M. musculus: C57BL/6J (2- and 18-months of age)	The National Institute on Aging Rodent Colony	N/A
Oligonucleotides		
5-Methylcytosine and 5-Hydroxymethylcytosine DNA	Zymo Research	Cat#D505
Stealth RNAi Sense Sequence: *H. sapiens* MAT2A: 5′AUCAAGGACAGCAUCACUGAUUUGG-3′	Thermo Fisher	Kera et al., 2013
Stealth RNAi Sense Sequence: Control: 5′-AGGAGGAUAUUACCAUGGACAAGAC-3′	Thermo Fisher	Kera et al., 2013
SUV39H1 RNAi: *H. sapiens*	Novus Biologicals	H00006839-R01-20nmol
SUV39H2 RNAi: *H. sapiens*	Novus Biologicals	H00079723-R01-20nmol
SETDB1 RNAi: *H. sapiens*	Novus Biologicals	H00009869-R01-20nmol
G9a/EHMT2 RNAi: *H. sapiens*	Novus Biologicals	H00010919-R02-20nmol

GLP/EHMT1 RNAi: *H sapiens*	Novus Biologicals	H00079813-R01-20nmol
RT-qPCR Primers	See Table S2	See Table S2
Software and Algorithms		
Bowtie2-2.1.0	Langmead and Salzberg, 2012	http://bowtie-bio.sourceforge.net/bowtie2/index.shtml
Samtools 1.5	Li et al., 2009	http://samtools.sourceforge.net/
MACS2-2.1.1.20160309	Zhang et al., 2008	http://liulab.dfci.harvard.edu/MACS/
Bedtools v2.25.0	Quinlan and Hall, 2010	https://bedtools.readthedocs.io/en/latest/
HOMER v4.9	Heinz et al., 2010	http://homer.ucsd.edu/homer/
GelAnalyzer 2010a	Lazar Software	http://gelanalyzer.com/index.html
MAVEN	Melamud et al., 2010; Clasquin et al., 2012	http://genomics-pubs.princeton.edu/mzroll/index.php
MAXQUANT	Cox et al., 2014	http://www.biochem.mpg.de/5111795/maxquant
Trimmomatic v0.39	Bolger et al., 2014	http://www.usadellab.org/cms/?page=trimmomatic
RSEM v1.2.4	Li et al., 2011	https://deweylab.github.io/RSEM/
GSEA 2	Tamayo et al., 2005 Mootha et al., 2003	http://software.broadinstitute.org/gsea/index.jsp
Skyline	MacLean et al., 2010	https://skyline.ms/project/home/software/Skyline/begin.view
Other		
Control *M. musculus* Diet	Envigo	TD.01084

| Methionine Deficient *M. musculus* Diet | Envigo | TD.140119 |

1370

1371

1372 **Supplemental File Titles and Legends**

1373

1374 Document S1, Related to Figures 3 and S3. Primary Antibody and REVERT Total Protein Stain

1375 Western Blot Membranes.

1376

(*A*) Cytoplasm anti-H3K9me1 (Figure 3E). (*B*) REVERT Total Protein Stain for cytoplasm anti-

H3K9me1 (Figure 3E). (*C*) Cytoplasm anti-H3 (Figure 3E). (*D*) REVERT Total Protein Stain for

cytoplasm anti-H3 (Figure 3E). (*E*) Cytoplasm anti-αTubulin (Figure 3E). (*F*) REVERT Total

Protein Stain for cytoplasm anti-αTubulin (Figure 3E). (*G*) Nucleoplasm anti-αTubulin (Figure

3E). (*H*) REVERT Total Protein Stain for nucleoplasm anti-αTubulin (Figure 3E). (*I*) Purified

histone anti-H3K9me1 (Figure S3F). (*J*) REVERT Total Protein Stain for purified histone anti-

H3K9me1 (Figure S3F).

1384

1385 Table S1, Related to Figures 1-3, 6-7, S1-S3, and S5. LC-MS/MS % Abundance Histone PTM

1386 Data.

1387

1388 Table S2, Related to Figures 2, 5, 7, S3, and S5. RT-qPCR Primers.

1389

1390 Table S3, Related to Figure 3. SILAC Cytoplasmic Proteins.

1391

1392

1393

Figure 1. Methionine Restriction Stimulates Dynamic Histone PTM Response

Figure 2. Decreased SAM Availability Drives Robust Histone Methylation Response

Figure 3. H3K9me1 Abundance under SAM Depletion is Driven by *de novo* Methylation

Figure 4. H3K9me1 is Redistributed Over Repetitive and Transposable Elements During SAM Depletion

Figure 5. *De novo* H3K9 Mono-methylation Preserves Heterochromatin Stability

Figure 6. Adaptive H3K9 Mechanisms Support Epigenetic Persistence Upon Metabolic Recovery

Figure 7. Adaptive and Persistent Epigenetic Responses to SAM Depletion are Robust *in vivo*, Independent of Age

Figure S1. Related to Figure 1.

Figure S2. Related to Figure 2

Figure S3. Related to Figure 3

Figure S4. Related to Figure 6

A

B EHMT1/GLP

C EHMT2/G9a

Figure S5. Related to Figure 6

C

D

A. GO_NUCLEIC_ACID_BINDING_TRANSCRIPTION_FACTOR_ACTIVITY_signal
B. GO_REGULATORY_REGION_NUCLEIC_ACID_BINDING_signal
C. GO_DOUBLE_STRANDED_DNA_BINDING_signal
D. GO_SEQUENCE_SPECIFIC_DNA_BINDING_signal
E. GO_CORE_PROMOTER_PROXIMAL_REGION_DNA_BINDING_signal

Figure S6. Related to Figure 7