Title: Methyl-Metabolite Depletion Elicits Adaptive Responses to Support Heterochromatin

Stability and Epigenetic Persistence

Spencer A. Haws[1,2], Deyang Yu[3,4,5], Cunqi Ye[6], Coral K. Wille[1,6], Long C. Nguyen[8,9], Kimberly A.

Krautkramer[1,2], Jay L. Tomasiewicz[3], Shany E. Yang[3,4], Blake R. Miller[3,4], Wallace H. Liu[1,2],

Kazuhiko Igarashi[8,9], Rupa Sridharan[1,6], Benjamin P. Tu[6], Vincent L. Cryns[3,5,10], Dudley W.

Lamming[3,4,5], and John M. Denu[1,2]*.

[1]Department of Biomolecular Chemistry, SMPH, University of Wisconsin, Madison, WI, 53706,

USA

[2]Wisconsin Institute for Discovery, University of Wisconsin, Madison, WI, 53715, USA

[3]Department of Medicine, SMPH, University of Wisconsin, Madison, WI, 53705, USA

[4]William S. Middleton Memorial Veterans Hospital, Madison, WI, 53705, USA

[5]Molecular & Environmental Toxicology Center, SMPH, University of Wisconsin, Madison, WI,

53705, USA

[6]Department of Biochemistry, University of Texas Southwestern Medical Center, Dallas, TX,

75390, USA

[7]Department of Cell and Regenerative Biology, University of Wisconsin, Madison, WI, 53705,

USA

[8]Department of Biochemistry, Tohoku University Graduate School of Medicine, Sendai, 980-

8575, JPN

[9]Center for Regulatory Epigenome and Diseases, Tohoku University Graduate School of

Medicine, Sendai, 980-8575, JPN

[10]University of Wisconsin Carbone Cancer Center, Madison, WI, 53792, USA

*Correspondence: John M. Denu, john.denu@wisc.edu

Summary
29
S-adenosylmethionine (SAM) is the methyl-donor substrate for DNA and histone
methyltransferases that regulate cellular epigenetic states. This metabolism-epigenome link has
been shown to promote the sensitivity of chromatin methylation patterns to altered SAM
abundance. However, the adaptive mechanisms that allow cells to actively protect epigenetic
information during major fluctuations in SAM availability are unknown. Here, we identify a robust
response to SAM depletion that is highlighted by preferential mono-methylation of H3 Lys 9
(H3K9) at the expense of global losses in histone di- and tri-methylation. These coordinated
pathways drive *de novo* cytoplasmic and nuclear mono-methylation of H3K9 to preserve
heterochromatin stability and support global epigenetic persistence upon metabolic recovery.
Conservation of these responses in both young and old mice further support a role for
epigenetic adaptation to SAM depletion *in vivo*. Together, these studies provide the first
evidence for active epigenetic adaptation and persistence to metabolic stress.
42
43   Keywords: metabolism, methionine, SAM, epigenetics, methylation, chromatin, histone, aging,
44   persistence

Introduction

Metabolism and the epigenome are connected by central metabolites that act as co-substrates for chromatin modifying enzymes, enabling fluctuations in metabolite availability to directly tune a cell's ability to 'write' and 'erase' chromatin post-translational modifications (PTMs) (Fan et al., 2015; Li et al., 2018). Chromatin methylation has been shown to be particularly susceptible to metabolic perturbations as methyltransferases require a single methyl-donor cofactor, S-adenosylmethionine (SAM) (Ducker and Rabinowitz, 2016; Mentch and Locasale, 2017). For example, accumulation of SAM in *S. cerevisiae* results in global increases in histone methylation levels (Ye et al., 2017). On the contrary, reduced SAM availability has been linked with site-specific losses in histone methylation as well as global depression of DNA methylation in several organisms (Shiraki et al., 2014; Ding et al., 2015; Kera et al., 2013; Tang et al., 2017; Shyh-Chang et al., 2013; Mentch et al., 2015; Tang et al., 2015; Hayashi et al., 2018; Towbin et al., 2012; Strekalova et al., 2019; Wang et al., 2019).

Intracellular SAM abundance is dependent on the catalysis of its obligatory precursor metabolite, the essential amino acid methionine (Met). Consumption of popular diets containing low quantities of Met (fish, vegetarian, and vegan) have been shown to significantly correlate with decreased plasma Met concentrations (Farmer, 2014; Schmidt et al., 2016). This link provides an opportunity for diet to directly influence SAM availability. Furthermore, abundance of Met and SAM have both been shown to fluctuate naturally in tune with circadian rhythms (Krishnaiah et al., 2017). Therefore, individuals consuming adequate amounts of dietary Met likely still experience periods of decreased methyl-metabolite availability. While both dietary intake and circadian regulation are capable of influencing Met and SAM abundance, adaptive mechanisms that allow cells to actively respond to – and recover their functional epigenomes from – such metabolic perturbations are unknown. Here, we define the ability of a cell and/or organism to re-establish the epigenome that was present prior to onset of an environmental perturbation or metabolic stress as *epigenetic persistence*.

Adaptive mechanisms that respond to methyl-metabolite depletion likely support critical cellular functions under these conditions. Interestingly, dietary Met-restriction is associated with beneficial metabolic reprogramming and lifespan extension in mammals (Orentreich et al., 1993; Yu et al., 2018; Miller et al., 2005; Brown-Borg et al., 2018; Lees et al., 2014; Green and Lamming, 2018). These phenotypes run counter to negative pathologies associated with dysregulated chromatin methylation (i.e., intellectual disability syndromes, cancers, and premature as well as natural aging) (Greer and Shi, 2012). This suggests cells possess

responsive mechanisms capable of actively maintaining regulation of site specific and/or global

chromatin methylation events during SAM depletion. The goal of this study is to identify and

characterize such mechanisms to better understand how organisms actively protect their

epigenome in response to fluctuations in central metabolite availability.

Here, we identify a robust, conserved chromatin response to metabolically-depleted

SAM levels. This cellular response is characterized by global decreases in higher-state histone

methylation (di- and tri-methyl) and simultaneous coordinated activation of *de novo* H3K9 mono-

methylation involving newly-synthesized and chromatin-bound histone H3. Under extreme

conditions of both reduced SAM levels and inhibited methylation that would yield H3K9me1,

heterochromatin instability and de-repression of constitutively silenced DNA elements is

exacerbated. Acute inhibition of *de* novo H3K9 mono-methylation also leads to the

dysregulation of global histone PTM states upon metabolic recovery, disrupting epigenetic

persistence to decreased SAM availability. Importantly, chronic metabolic depletion of SAM in

both young and old mice recapitulates these *in vitro* findings, revealing a conserved, age-

independent chromatin response. These results implicate *de novo* H3K9 mono-methylation as

an indispensable mechanism to support heterochromatin stability and global epigenetic

persistence in response to SAM depletion.

Results

<u>Methionine Restriction Stimulates Dynamic Histone PTM Response</u>

To comprehensively investigate how disruption of methyl-donor metabolism affects

chromatin methylation states, global histone proteomics and DNA methylation (5mC) analyses

were performed on two distinct *in vitro* and *in vivo* systems. These included Met-restricted

HCT116 human colorectal cancer cells and C57BL/6J mouse liver. Methionine metabolism was

severely repressed in both systems, highlighted by significant reductions in Met and SAM

abundance (Figure 1A and Figure S1A-S1D). Reduced methyl-metabolite availability did not

affect global DNA 5mC abundance in either system (Figure 1B-1C). However, LC-MS/MS

analysis of more than 40 unique histone H3 peptide proteoforms revealed dynamic PTM

responses to Met-restriction in both HCT116 cells and C57BL/6J liver (Figure 1D-1E). In

HCT116 cells, Met-restriction stimulated distinct biphasic changes in global PTM profiles (Figure

1D). Phase I (0 hr-45 min) was rapid and marked by a trending upregulation of H3K4me2/3,

PTMs known to mark transcriptionally active promoters. Phase II (90 min-24 hrs) was

characterized by global decreases in di- and tri- histone methylation. Decreased levels of di-
and tri-methylated peptides were accompanied by increases in acetylated and unmodified
peptide species. Similarly in C57BL/6J mice, histone PTM responses were marked by
significant decreases in histone di- and tri-methylation that essentially matched the patterns
found in HCT116 cells during the prolonged Phase II response (Figure 1E and Figure S1E-S1J).
Decreased higher state (di- and tri-) histone methylation both *in vitro* and *in vivo* highlights a
decreased methylation capacity resulting from prolonged Met and/or SAM depletion. Together,
these observations suggest perturbed methyl-donor metabolism is capable of stimulating
significant changes in histone, but not global 5mC, methylation abundance. Furthermore,
similarities between the metabolic and epigenetic responses to Met-restriction across both
systems support the use of *in vitro* Met-restriction as a model for mechanistic follow-up studies.
124
125 <u>Decreased SAM Availability Drives Robust Histone Methylation Response</u>
126
Dramatic reduction of intracellular Met and SAM correlated with onset of the *in vitro*
Phase I and II histone PTM changes, respectively (Figure 1D and Figure S1A-S1B). This
implies depletion of individual methyl-metabolites may be capable of stimulating distinct histone
modifying pathways. To determine if the global losses in di- and tri- histone methylation *in vitro*
are driven specifically by SAM depletion, two alternative approaches were employed to deplete
cellular SAM independent of Met (Figure 2A). The first approach utilized RNAi knockdown of the
mammalian SAM synthetase MAT2A (methionine adenosyltransferase II alpha). Knockdown via
MAT2A-RNAi treatment decreased *MAT2A* transcript abundance by greater than 97%, resulting
in a near complete deprivation of SAM availability with no effect on Met levels (Figure 2B-2D).
Histone proteomics analysis of MAT2A-RNAi treated cells identified site-specific histone PTM
changes that mirrored those identified in HCT116 cells under Phase II Met-restriction conditions
(r=0.778, p=8.59e^{-10}) (Figure 2E). Using a complementary approach, SAM levels were depleted
by overexpression of a major SAM consumer, PEMT (phosphatidylethanolamine N-
methyltransferase). Ye et al. 2017 have previously shown PEMT overexpression significantly
reduces SAM availability in HEK293T human embryonic kidney cells. Histone proteomics
analysis of *PEMT* overexpressing HEK293T cells identified site-specific PTM changes that
paralleled those of Met-restricted controls (r=0.803, p=3.30e^{-8}) (Figure 2F). Together, two
separate means of reducing SAM levels demonstrate SAM reduction alone is sufficient to inhibit
the methylation capacity of a cell, inducing dynamic changes in histone PTM abundance.

Furthermore, these findings support the use of prolonged Met-restriction as a model of SAM

depletion.

The similar changes in global histone PTM abundance across four SAM depletion

systems were analyzed for the presence of methyl PTMs whose abundances either increased

or were unaffected by this metabolic perturbation. Such PTMs would be candidate mediators of

an adaptive response to SAM depletion. Unexpectedly, all four SAM depletion systems

exhibited identical changes in H3K9 methylation (Figure 2G-2J). SAM depletion stimulated a

decrease in H3K9me2/3 with a corresponding increase in H3K9ac/14ac and unmodified

peptides while net H3K9me1 abundance was unchanged. Preservation of net H3K9me1 levels

in response to SAM depletion was reproducible in Met-restricted MCF7, HEK-293, Panc1, and

HepG2 cell lines, as well as in MAT2A-RNAi treated HepA1 cells (Figure S2A-S2F). The robust

conservation of H3K9me1 abundance in diverse systems marked by reduced methylation

capacity suggest this PTM may be supported by adaptive, active mechanisms.

<u>H3K9me1 Abundance under SAM Depletion is Driven by *de novo* Methylation</u>

Global levels of H3K9me1 could be maintained through two distinct mechanisms during

SAM depletion: de-methylation of H3K9me2/3 and/or *de novo* methylation of unmodified H3K9.

The decrease in H3K9me2/3 and increase in unmodified H3K9 peptide levels across SAM

depletion systems suggest H3K9 can be subjected to facile demethylation while *de novo*

methylation appears enzymatically unfavorable. However, *de novo* H3K9 mono-methylation

would constitute an active, adaptive response to this metabolic stress that is capable of

supporting critical chromatin functions. To investigate the contribution of *de novo* methylation to

global H3K9me1 levels, small molecule (UNC0642) inhibition of H3K9 HMTs Ehmt1 and Ehmt2

was used to block nuclear H3K9 mono-methylation (Liu F et al., 2013). Small molecule

treatment enabled acute inhibition of Ehmt1 and Ehmt2 activity whereas RNAi-mediated

repression of H3K9 HMT expression had significant yet minimal effects on H3K9 PTM

abundance after 24 hours (Figure S3A-S3D). RNAi treatment was limited to 24 hours as

HCT116 cells no longer proliferate beyond 24 hours of Met-restriction (Figure S3E). UNC0642

treatment of Met-replete cells significantly reduced H3K9me1 abundance by 35.6% and

coupling UNC0642 treatment with Met-restriction stimulated a 20.7% decrease in H3K9me1

levels (Figure 3A-3C, Figure S3F). These data suggest Ehmt1/Ehmt2 retain nearly 60% of their

H3K9 mono-methylation activity despite severe *in situ* SAM deprivation and that this activity is

critical for preserving global H3K9me1 abundance. Furthermore, H3K9me2 abundance

decreased significantly in Met-replete cells upon UNC0642 treatment but was unaffected when

UNC0642 was coupled with Met-restriction (Figure 3A-3B). This indicates only the mono-

methylation activity of Ehmt1 and Ehmt2 is preferentially supported during SAM depletion.

In addition to methylation of H3K9 by nuclear enzymes, mono-methylation is known to

be enzymatically catalyzed on cytoplasmic histones prior to nuclear import and deposition onto

chromatin (Loyola A et al., 2006; Pinheiro et al., 2012). Subcellular fractionation of Met-

restricted HCT116 cells coupled with western blot analyses indicated that cytoplasmic

H3K9me1 levels increase 3-fold after 24 hours of Met-restriction, with a similar increase in total

H3 protein levels (Figure 3D-3E). To determine the contribution of cytoplasmic H3K9me1 to the

chromatin-bound nuclear pool after 24 hours of Met-restriction, a pulse-chase, SILAC

experiment was performed (Figure 3F). L-Arg^{+10} isotope incorporation prior to Met-restriction

was determined to be nearly complete with average incorporation for H3K9 peptides greater

than 98% (Figure S3H). Newly synthesized H3 was calculated to comprise 24% of the

chromatin bound histone pool and contributed 18% of the global H3K9me1 signal after 24 hours

of Met-restriction (Figure 3G-3H). UNC0642 treatment did not significantly lessen the

contribution of cytoplasmic H3K9me1 to the total pool, suggesting cytoplasmic H3K9me1 is

largely protected from enzyme catalyzed turnover once deposited onto chromatin under these

conditions (Figure 3H). Modest, insignificant decreases in L-Arg^{+0} H3K9 PTM signal after

UNC0642 treatment can be attributed to reduced incorporation of newly synthesized H3 onto

chromatin. (Figure 3G) Reduced H3 incorporation in UNC0642-treated cells is likely the

consequence of a slower growth rate under these conditions (Figure S3E). Together, these

results suggest ~40% of global H3K9me1 abundance is supported by *de novo* methylation

during Met-restriction with nearly equal contributions from both the cytoplasm and nucleus. The

remaining ~60% of preserved H3K9me1 levels are likely the result of H3K9me2/3 demethylation

(Figure 3I).

The significant upregulation of cytoplasmic H3K9me1 and total H3, coupled with their

incorporation onto chromatin during SAM depletion, were unexpected. These findings suggest

cells utilize scarcely available Met and SAM to support the protein abundance and methylation

of histone H3 under these conditions. Shotgun proteomics was performed on cytoplasmic

protein fractions of the previously described SILAC experiment to determine if histone protein

abundance is uniquely responsive to SAM depletion. Of the more than 2,400 identified proteins,

only 8 possessed significant incorporation of the light L-Arg^{+0} isotope and also increased in

absolute abundance by a $\log_2$ fold-change value greater than 2.0 (Figure 3J). Frequency of Met

residues in the primary amino acid sequence did not affect L-Arg^{+0} isotope incorporation (Figure

S3I). Remarkably, all four core histone subunits, Mat2A, and the methionine importer SLC3A2

(Bröer et al., 2001) were among the 8 proteins (Figure 3J). These results suggest proteins

critical for chromatin stability and methyl-donor metabolism are preferentially generated during

metabolically-induced SAM depletion. To our knowledge, this is also the first example of a

metabolic stress stimulating increased histone protein abundance.

<u>H3K9me1 is Redistributed Over Repetitive and Transposable Elements During SAM Depletion</u>

Multiple, non-redundant mechanisms for H3K9me1 maintenance suggest strong

biological pressure exists to retain this PTM under SAM depleted conditions. One critical

function for H3K9me1 is to serve as a primer for H3K9me2/3 methylation as methyltransferases

that yield these higher methylation states require a mono-methyl substrate (Peters et al., 2001;

Loyola et al., 2006; Loyola et al., 2009). Accordingly, installment of H3K9me2/3 for constitutive

heterochromatin-mediated repression of telomeric, pericentromeric, and centromeric

repetitive/transposable elements requires readily available H3K9me1 (Rea et al., 2000;

Bannister et al., 2001; Lachner et al., 2000). Global H3K9me2/3 losses in response to SAM

depletion suggest these conditions introduce heterochromatin instability, which would create an

increased need for H3K9me1 to restore H3K9me2/3 levels when SAM becomes readily

available.

To determine if H3K9me1 is being maintained globally or redistributed to regions of

susceptible heterochromatin during SAM depletion, ChIP-sequencing was performed. Met-

restriction stimulated a 35% loss in total H3K9me3 peaks abundance (Figure 4A). A higher

percentage of H3K9me3 peaks were lost at LTR, intergenic, and intron annotated loci relative to

the total, while H3K9me3 enrichment at satellite, simple repeat, and SINE annotated loci were

slightly less sensitive to SAM depletion (Figure 4A). Interestingly, H3K9me1 enrichment at

repetitive and transposable loci were either largely unaffected by SAM depletion or increased

(Figure 4B). Only the percent change for intergenic and intron annotated H3K9me1 peaks was

lost to a greater extent than the total peak number.

Maintained or increased levels of H3K9me1 at constitutively repressed regions during

SAM depletion could result from the replacement of H3K9me1 for lost H3K9me3. To investigate

this possibility, we mapped H3K9me1 reads to regions of lost H3K9me3, and determined that

H3K9me1 enrichment increases at 66.4% of nearly $16e^4$ sites (Figure 4C-4D). A higher

percentage of these H3K9me1 enriched sites were annotated for LINE, LTR, simple repeat, and

satellite elements relative to those experiencing decreased H3K9me1 read enrichment (Figure

4E). This indicates H3K9me1 preferentially replaces H3K9me3 at repetitive and transposable

loci during SAM depletion. Figure 4F provides a representative image of H3K9me3 replacement

by H3K9me1 at a pericentric, repetitive loci. Together, these ChIP-sequencing analyses suggest

SAM depletion stimulates the redistribution of H3K9me1 to repetitive and transposable genomic

loci which are susceptible to H3K9 demethylation.

De novo H3K9 Mono-methylation Preserves Heterochromatin Stability

Replacement of H3K9me3 by H3K9me1 may function to preserve heterochromatin

stability and pericentric chromatin repression during SAM depletion by providing the primer for

site-specific H3K9me2/3 methylation. An MNase accessibility assay was used to evaluate global

heterochromatin stability under SAM depleted conditions as MNase can more readily digest

euchromatic DNA compared to heterochromatic DNA. Coupling Met-restriction with UNC0642

treatment resulted in a significantly greater loss and accumulation of di- and mono-nucleosome

species, respectively, compared to Met-restriction alone (Figure 5A-5B). This indicates inhibition

of *de novo* H3K9 mono-methylation during SAM depletion has a more detrimental effect on

global heterochromatin stability than SAM depletion in isolation. To determine if global

decreases in heterochromatin stability resulted in de-repression of transposable DNA elements,

RT-qPCR was employed. Transcript abundance of *LINE 1*, *HERV-K*, and *HERV-R* were all

significantly elevated after SAM depletion. Coupling SAM depletion with UNC0642 treatment

resulted in the further elevation of *LINE1* and *HERV-K* transcript abundance while *HERV-R*

expression was unaffected (Figure 5C-5E). Because SAM depletion-induced losses in

heterochromatin stability and retrotransposon repression are exacerbated with inhibition of

nuclear Ehmt1/2 activity, these data suggest that stimulation of active H3K9 mono-methylation

functions, in part, to preserve heterochromatin stability.

Adaptive H3K9 Mechanisms Support Epigenetic Persistence Upon Metabolic Recovery

Dysregulation of adaptive epigenetic mechanisms during SAM depletion may produce

aberrant changes in histone PTMs that compromise long-term epigenetic persistence upon

metabolic recovery. To determine if *de novo* H3K9me1 is required for epigenetic persistence to

SAM depletion, HCT116 cells were allowed to recover in Met-replete media after being

subjected to Met-restriction coupled with acute vehicle control or UNC0642 treatment (Figure

6A). Histone proteomics analysis confirmed that inhibition of Ehmt1/2 activity during SAM

depletion results in a significant loss of H3K9me1 abundance (Figure 6B-6C). Upon 5 and 24

hours of Met-repletion, vehicle control cells largely regained the original histone PTM state

(Figure 6B). However, histone PTM states of UNC0642 cells acutely treated only during Met-

restriction remained dysregulated after both 5 and 24 hours of Met-repletion (Figure 6B).

Although significant global dysregulation was apparent, the H3K9 peptide proteoforms displayed

the most dynamic differences (Figure 6D-6E). Interestingly, the signature H3K9 PTM response

to SAM depletion described in Figure 3 was largely retained upon Met-repletion in cells acutely

treated with UNC0642 (Figure 6D-6E). H3K9me3 abundance was an exception, recovering to

levels consistent with regained *LINE 1* and *HERV-K/R* repression (Figure 5C-5E).

291   To assess the transcriptional consequences of lost epigenetic persistence, RNA-

sequencing was performed on cells which recovered in Met-replete media after Met-restriction

coupled with acute vehicle control or UNC0642 treatment. Principle Component Analysis (PCA)

of normalized Transcripts Per Million (TPM) values produced 3 distinct clusters, suggesting

altered epigenomes correlate with altered transcript profiles in these cells (Figure 6F). A multiple

condition comparison analysis was used to determine which transcripts distinguished the

clusters from one another (Figure 6G). While 68.4% of identified genes were ubiquitously

expressed across samples (Pattern 1), 27.7% were uniquely expressed in control cells relative

to both recovery groups (Pattern 4) (Figure 6H). This result supports the PCA findings as the

pre-restriction control group is most distinct from both recovery groups. Unique gene expression

profiles for UNC0642 (Pattern 2) and DMSO (Pattern 3) recovery cells contained 1.9% and

0.7%, respectively, of all identified genes. Therefore, these subsets of genes are responsible for

the unique clustering of both Met-restriction recovery groups (Figure 6H). Gene Set Enrichment

Analysis (GSEA) was performed on both Pattern 2 and 3 gene lists to determine functional

differences between each gene set. GSEA of Pattern 2 genes identified 71 significantly enriched

gene ontology (GO) terms possessing an FDR q-value less than 0.25, while an identical

analysis of Pattern 3 genes did not identify any significantly enriched GO terms (Figure 6I-6J).

Of the significantly enriched GO terms in Pattern 2, 66 possessed a positive Normalized

Enrichment Score (NES) when comparing UNC0642 to vehicle control recovery cells (Figure

S4A-S4B). Figure 6K is a representative GSEA running enrichment score figure of a GO term

possessing a positive NES. Only 5 of the significantly enriched GO terms in Pattern 2

possessed a negative NES when comparing UNC0642 to vehicle control recovery cells.

Notably, a majority of the genes assigned to these GO terms were zinc finger binding proteins

(Figure S4C-S4D). Figure 6L is a representative GSEA running enrichment score figure of a GO

term possessing a negative NES.

These data suggest adaptive regulation of H3K9 PTMs in response to SAM depletion is
required for long-term epigenetic persistence upon metabolic recovery. Furthermore, inhibited
epigenetic persistence results in significant transcriptional consequences that impact a diverse
set of cellular functions.
320
321 Adaptive and Persistent Epigenetic Responses to SAM Depletion are Robust *in vivo,*
322 Independent of Age
323
To determine if adaptive and persistent responses to SAM depletion exist *in vivo* and are
age-dependent, 6- and 22-month C57BL/6J mice were subjected to 3 weeks of Met-restriction
followed by 5 weeks of Met-repletion (Figure 7A). Several previous reports have associated
decreased H3K9me2/3 abundance with aging in numerous organisms (Wood et al., 2010;
Larson et al., 2012; Ni et al., 2012; Tvardovskey et al., 2017; Scaffidi and Misteli, 2006). Age-
dependent dysregulation of heterochromatin has been associated with de-repression of
repetitive DNA and transposable elements, as well as DNA damage, factors which may
contribute to disease progression (Wood et al., 2016; De Cecco et al., 2013; De Cecco et al.,
2019; Oberdoerffer and Sinclair, 2007; Booth and Brunet, 2016). Therefore, conservation of
both adaptive and persistent responses to SAM depletion across the lifespan would further
support the functional importance of both mechanisms as an "older" chromatin environment is
thought to be inherently dysregulated.
Remarkably, histone proteomics analysis of liver tissue revealed H3K9 was the only
residue on which PTMs responded similarly to SAM depletion in both 6- and 22-month mice
relative to age matched controls (Figure 7B). This PTM response at H3K9 phenocopied those
identified in the previously described *in vitro* and whole-organism SAM depletion studies, further
highlighting the robust regulation of this residue (Figure 7C). Significant age-dependent
responses to SAM depletion were identified on all remaining histone residues (Figure 7B and
S5A-S5E). Furthermore, significantly elevated *LINE 1*, *SINE B1*, and *SINE B2* transcript
abundance in both 6- and 22-month mice suggest the chromatin instability phenotype
characterized in Figure 5 also occurs during metabolic depletion of SAM *in vivo* (Figure 7D-7E).
After 5 weeks of dietary Met-reintroduction, both young and old mice displayed the ability to re-
establish an epigenome similar to that possessed by age-matched controls (Figure 7B).
Interestingly, 22-month mice appeared to more rapidly adopt the control histone PTM state,
while the 6-month mice still exhibited residual losses in H3K9me3 and increases in H3K9me1
after 5 weeks of recovery (Figure 7F). Consistent with the general recovery of chromatin states,

both age groups largely reestablished repression of *LINE 1*, *SINE B1*, and *SINE B2* expression

(Figure 7G-7H).

Metabolic changes in young and old mice tracked with H3K9 PTMs. Both 6- and 22-

month mice in the Met-restriction group experienced significant losses in total body weight

relative to age matched controls (Figure S5F). Decreases in overall body weight were

comprised of significant losses in both lean and fat mass (Figure S5F). Altered body-weight and

compositions were accompanied by significantly improved glucose tolerance in both young and

old mice (Figure S5G). Upon Met-repletion, mice reacquired their initial metabolic state in an

age-independent manner. All animals experienced significant increases in total body weight,

comprised of elevated lean and fat mass, as well normalized glucose tolerance relative to age-

matched controls (Figure S5H-S5I).

Together, strong association of the H3K9 chromatin response to SAM depletion with

metabolic reprogramming in young and old mice suggest H3K9 regulatory mechanisms are not

only robust across lifespan but may be required to support these metabolic phenotypes.

Furthermore, these data show the ability of epigenetic states to persist upon recovery from a

metabolic challenge is an inherent property of both isolated cells and complex mammals.

Discussion

Here, we provide a comprehensive investigation into the adaptive regulation of

chromatin during a metabolic stress. A highly conserved, robust histone methylation response to

SAM depletion was discovered, characterized by global losses in histone H3 di- and tri-

methylation while H3K9me1 is actively maintained to safeguard against exacerbated

heterochromatin instability and support epigenetic persistence upon metabolic recovery. These

mechanisms involve the enrichment of H3K9me1 in susceptible regions of heterochromatin

through *de novo* methylation on both chromatin-bound and cytoplasmic H3. To our knowledge,

this is the first identified mechanism in which scarce, diet-derived nutrients are funneled toward

the preservation of specific chromatin PTM states.

Zee et al., 2010 have previously determined H3K9me1 has the shortest half-life of 17

measured PTMs, highlighting the importance of *de novo* methylation for supporting global

H3K9me1 abundance in a SAM depleted environment. Here, *de novo* methylation was found to

be critical for more than simply sustaining global H3K9me1 levels under these conditions. ChIP-

sequencing analyses determined H3K9me1 experiences a significant redistribution across

heterochromatin and enrichment at repetitive and transposable DNA elements, particularly in

regions where H3K9me3 is lost. Redistribution of H3K9me1 to these regions is likely critical for

preserving heterochromatin as prevention of *de novo* H3K9 mono-methylation resulted in global

losses in heterochromatin stability and site-specific retrotransposon de-repression. We

hypothesize H3K9me1 mediates repression of retrotransposons during SAM depletion by acting

as a substrate for HMTs which deposit di- and tri-methyl PTMs onto H3K9. (Peters et al., 2001;

Loyola et al., 2006; Loyola et al., 2009; Rea et al., 2000; Bannister et al., 2001; Lachner et al.,

2000). This function is not limited to nuclear derived H3K9me1. Pinheiro et al., 2012 have

previously shown inhibition of cytoplasmic H3K9 mono-methylation through shRNA knockdown

of *PRDM3* and *PRDM16* stimulates pericentric heterochromatin loss and disruption of the

nuclear lamina. Our study demonstrates cytoplasmic derived H3K9me1 comprises a significant

portion (18%) of the total chromatin bound pool after 24 hours of Met-restriction *in vitro*.

Adaptation of H3K9 mono-methylation to SAM depletion is also critical for long-term epigenetic

persistence to this metabolic stress. Prevention of *de novo* H3K9 mono-methylation during SAM

depletion resulted in sustained dysregulation of global histone PTMs upon metabolic recovery,

significantly influencing gene expression patterns. This requirement for an active, adaptive

histone PTM response to support epigenetic persistence upon metabolic recovery is the first

identified mechanism of its kind.

Replication of *in vitro* results *in vivo*, independent of age, strongly supports the discovery

of a universal adaptive and persistent response to SAM depletion. Here, we demonstrate that

the adaptive mechanisms required to support H3K9me1-mediated heterochromatin stability are

inducible in both 6- and 22-month mice after dietary Met-restriction. Furthermore, H3K9me1

maintenance under SAM depleted conditions uniquely correlated with beneficial metabolic

reprogramming (i.e. reduced fat mass and improved glucose tolerance) in both young and old

mice. This suggests the robust regulation of H3K9 PTMs in response to SAM depletion may be

required to support the metabolic reprogramming stimulated by dietary Met-restriction. Global

epigenetic persistence to metabolic SAM depletion was also identified in both 6- and 22-month

mice. Validation of this phenomenon *in vivo*, independent of age, suggests similar mechanisms

critical for maintaining proper regulation of the epigenome during natural fluctuations in the

availability of other metabolic cofactors may exist as well. Such fluctuations could be stimulated

by periods of prolonged fasts, chronic intake of foods lacking essential cofactors (i.e. vegan and

vegetarian dietary patterns), and circadian regulated changes in metabolite availability (Farmer,

2014; Schmidt et al., 2016; Krishnaiah et al., 2017).

Notably, we did not observe an age-associated decrease in H3K9 methylation between

6- and 22-month control diet mice (Table S1). Similar regulation of H3K9 PTMs and mirrored

metabolic responses to SAM depletion across age groups suggest our 22-month mice, the

equivalent to a 60-70 year old human, have not yet experienced a decline in health span

(Flurkey et al., 2007). It will be interesting to determine if the adaptive and persistent epigenetic

mechanisms revealed here are functional in very old animals (i.e. >30 months) where H3K9

methylation is thought to be inherently dysregulated. Disruption of these mechanisms later in an

organism's lifespan could support age-associated de-repression of transposable elements

(Wood et al., 2016; De Cecco et al., 2013; Oberdoerffer and Sinclair, 2007; Booth and Brunet,

2016), contributing to negative phenotypes that accelerate the aging process such as

inflammation (De Cecco et al., 2019; Simon et al., 2019).

Future studies will be needed to determine how H3K9 mono-methyltransferases are able

to maintain their catalytic activity in SAM depleted environments. Favorable steady-state kinetic

constants for H3K9me1 relative to H3K9me2/3 HMTs is one possibility. However, it is difficult to

address this hypothesis by comparing current H3K9 HMT enzymology data due to varied

selections of substrates and assays. Direct shuttling of SAM to H3K9me1 HMTs by Mat2A could

also support H3K9me1 catalytic activity. As SAM is energetically expensive to produce, due to

the 1:1 requirement of Met and ATP molecules for its synthesis, in addition to being highly

unstable (Wu et al., 1983), it would be advantageous for the cell to possess mechanisms which

permit preferential shuttling of SAM during methyl-metabolite depletion. Directed SAM

production and utilization have been proposed in other metabolic contexts (Katoh et al., 2011;

Kera et al., 2013; Li et al., 2015).

Acknowledgements

This research was supported in part by grants from the NIH (S.A.H. –T32 DK007665; K.A.K.

T32 DK007665; W.H.L. T32 DK007665; J.M.D. – R37 GM059785; D.W.L. – AG041765,

AG050135, AG051974, AG056771, AG062328, a New Investigator Program Award (D.W.L.)

and a Collaborative Health Sciences Program Award (V.L.C.) from the Wisconsin Partnership

Program, the V. Foundation for Cancer Research (V.L.C.), and a Glenn Foundation Award for

Research in the Biological Mechanisms of Aging (D.W.L.), as well as startup funds from the

UW-Madison School of Medicine and Public Health and the UW-Madison Department of

Medicine (V.L.C. and D.W.L.). This research was conducted while D.W.L. was an AFAR

Research Grant recipient from the American Federation for Aging Research. D.Y. is supported

in part by a fellowship from the American Heart Association (17PRE33410983). The Lamming

laboratory is supported in part by the U.S. Department of Veterans Affairs (I01-BX004031), and

this work was supported using facilities and resources from the William S. Middleton Memorial

Veterans Hospital. This work does not represent the views of the Department of Veterans
Affairs or the United States Government. We would like to thank Alexis J. Lawton for the
bioinformatics analysis presented in Figure S3I. We would also like to thank the University of
Wisconsin-Madison Biotechnology Center and the University of Wisconsin-Madison Center for
High Throughput Computing for their help in generating and analyzing the ChIP-sequencing
data presented in Figure 5, respectively.

Author Contributions

Conceptualization, S.A.H., D.W.L, V.L.C., and J.M.D.; Methodology, S.A.H., C.Y., C.K.W.,
L.N.C., W.H.L., K.I., B.P.T., D.W.L., and J.M.D.; Validation, S.A.H., D.Y., C.Y., L.N.C., K.A.K.,
K.I., B.P.T., D.W.L., and J.M.D.; Formal Analysis, S.A.H., D.Y., C.K.W., K.A.K., D.W.L., and
J.M.D; Investigation, S.A.H., D.Y., C.Y., L.N.C., K.A.K., J.L.T., S.E.Y., and B.R.M.; Resources,
K.I., B.P.T., V.L.C., D.W.L., and J.M.D.; Writing – Original Draft, S.A.H. and J.M.D.; Writing –
Review & Editing, S.A.H., D.Y., C.Y., C.K.W., W.H.L., K.I., R.S., B.P.T., V.L.C., D.W.L., and
J.M.D.; Visualization, S.A.H., D.Y., and D.W.L.; Supervision and Project Administration, K.I.,
R.S., B.P.T., D.W.L., and J.M.D.; Funding Acquisition, S.A.H., K.I., R.S., B.P.T., V.L.C., D.W.L.,
and J.M.D.

Declaration of Interests

473 J.M.D. is a consultant for FORGE Life Sciences and co-founder of Galilei BioSciences. D.W.L
474 has received funding from, and is a scientific advisory board member of, Aeonian
475 Pharmaceuticals, which seeks to develop novel, selective mTOR inhibitors for the treatment of
476 various diseases. Remaining authors declare no competing interest.

References

1. Arriola Apelo, S.I., Neuman, J.C., Baar, E.L., Syed, F.A., Cummings, N.E., Brar, H.K., Pumper, C.P., Kimple, M.E., and Lamming, D.W. (2016) Alternative rapamycin treatment regimens mitigate the impact of rapamycin on glucose homeostasis and the immune system. Aging Cell. 15, 28-38.

2. Bannister, A.J., Zegerman, P., Pardridge, J.F., Miska, E.A., Thomas, J.O., Allshire, R.C., and Kouzarides, T. (2001) Selective recognition of methylated lysine 9 on histone H3 by the HP1 chromo domain. Nature. 410, 120-124.

3. Bolger, A.M., Lohse, M., and Usadel, B. (2014) Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics. 30, 2114-2120.

4. Booth, L.N. and Brunet, A. (2016) The aging epigenome. Mol. Cell. 62, 728-744.

5. Bröer, A., Friedrich, B., Wagner, C.A., Fillon, S., Ganapathy, V., Lang, F., and Bröer, S. (2001) Association of 4F2hc with light chains LAT1, LAT2 or y+LAT2 requires different domains. Biochem J. 355, 735-731.

6. Brown-Borg, H.M., Rakoczy, S., Wonderlich, J.A., Borg, K.E., and Rojanathammanee, L. (2018) Metabolic adaptation of short-living growth hormone transgenic mice to methionine restriction and supplementation. Ann. N.Y. Acad. Sci. 1418, 118-136.

7. Clasquin, M.F., Melamud, E., and Rabinowitz, J.D. (2012) LC-MS processing with MAVEN: a metabolomic analysis and visualization engine. Curr. Protoc. Bioinformatics. Chapter 14, Unit 14.11. doi:10.1002/0471250953.bi1411s37.

8. Cox, J., Hein, M.Y., Luber, C.A., Paron, I., Nagaraj, N., and Mann, M. (2014) Accurate proteome-wide label-free quantification by delayed normalization and maximal peptide ratio extraction, termed MaxLFQ. Mol. Cell Proteomics. 13, 2513-2526.

9. De Cecco, M., Criscione, S.W., Peckham, E.J., Hillenmeyer, S., Hamm, E.A., Manivannan, J., Peterson, A.L., Kreiling, J.A., Neretti, N., and Sedivy, J.M. (2013) Genomes of replicatively senescent cells undergo global epigenetic changes leading to gene silencing and activation of transposable elements. Aging Cell. 12, 247-256.

10. De Cecco, M., Ito, T., Petrashen, A.P., Elias, A.E., Skivr, N.J., Criscione, S.W., Caligiana, A., Brocculi, G., Adney, E.M., Boeke, J.D., Le, O., Beauséjour, C., Ambati, J., Ambati, K., Simon, M., Seluanov, A., Gorbunova, V., Slagboom, P.E., Helfand, S.L., Neretti, N., and Sedivy, J.M. (2019) L1 drives IFN senescent cells and promotes age-associated inflammation. Nature. 566, 73-78.

11. Ding, W., Smulan, L.J., Hou, N.S., Taubert, S., Watts, J.L., and Walker, A.K. (2015) S-
adenosylmethionine levels govern innate immunity through distinct methylation-
dependent pathways. Cell Metab. 22, 633-645.

12. Ducker, G.S. and Rabinowitz, J.D. (2017) One-carbon metabolism in health and
disease. Cell Metab. 25, 27-42.

13. Fan, J., Krautkramer, K.A., Feldman, J.L., and Denu, J.M. (2015) Metabolic regulation of
histone post-translational modifications. ACS Chem. Biol. 10, 95-108.

14. Farmer, B. (2014) Nutritional adequacy of plant-based diets for weight management:
observations from the NHANES. Am. J. Clin. Nutr. 100 Suppl 1, 365S-368S.

15. Flurkey, K., Currer, J.M., and Harrison, D.E. (2007) The Mouse in Aging Research. In
the Mouse in Biomedical Research 2nd Edition. Fox, J.G., et al., editors. American
College Laboratory Animal Medicine (Elsevier), pp. 637-671.

16. Green, C.L. and Lamming, D.W. (2018) Regulation of metabolic health by essential
dietary amino acids. Mech. Ageing Dev. S0047-6374(18)30079-4. doi:
0.1016/j.mad.2018.07.004.

17. Greer, E.L. and Shi, Y. (2012) Histone methylation: a dynamic mark in health, disease,
and inheritance. Nat. Rev. Genet. 13, 343-357.

18. Hardies, S.C., Wang, L., Zhou, L., Zhao, Y., Casavant, N.C., and Huang, S. (200) LINE-
1 (L1) lineages in the mouse. Mol. Biol. Evol. 17, 616-628.

19. Hayashi, T., Teruya, T., Chaleckis, R., Morigasaki, S., and Yanagida, M. (2018) S-
adenosylmethionine synthetase is required for cell growth, maintenance of G0 phase,
termination of quiescence in fission yeast. iScience 5, 38-51.

20. Heinz, S., Benner, C., Spann, N., Bertolino, E., Lin, Y.C., Laslo, P., Cheng, J.X., Murre,
C., Singh, H., and Glass, C.K. (2010) Simple combinations of lineage-determining
transcription factors prime cis-regulatory elements required for macrophage and B cell
identities. Mol. Cell. 38, 576-589.

21. Kang, Y.K., Park, J.S., Lee, C.S., Yeom, Y.I., Chung, A.S., and Lee, K.K. (1999) Efficient
integration of short interspersed element-flanked foreign DNA via homologous
recombination. J. Biol. Chem. 274, 36585-36591.

22. Katoh, Y., Ikura, T., Hoshikawa, Y., Tashiro, S., Ito, T., Ohta, M., Kera, Y., Noda, T., and
Igarashi, K. (2011) Methionine adenosyltransferase II serves as a transcriptional
corepressor of Maf oncoprotein. Mol. Cell. 41, 554-566.

23. Kera, Y., Katoh, Y., Ohta, M., Matsumoto, M., Takano-Yamamoto, T., and Igarashi, K.
(2013) Methionine adenosyltransferase II-dependent histone H3K9 methylation at the
COX-2 gene locus. J. Biol. Chem. 288, 13592-13601.

24. Klawitter, S., Fuchs, N.V., Upton, K.R., Upton, K.R., Muñoz-Lopez, M., Shukla, R.,
Wang, J., Garcia-Cañadas, M., Lopez-Ruiz, C., Gerhardt, D.J., Sebe, A., Grabundzija, I.,
Merkert, S., Gerdes, P., Pulgarin, J.A., Bock, A., Held, U., Witthuhn, A., Haase, A.,
Sarkadi, B., Löwer, J., Wolvetang, E.J., Martin, U., Ivics, Z., Izsvák, Z., Garcia-Perez,
J.L., Faulkner, G.J., and Schumann, G.G. (2016) Reprogramming triggers endogenous
L1 and Alu retrotransposition in human induced pluripotent stem cells. Nat. Commun.
7:10286. doi: 10.1038/ncomms10286.

25. Konze, K.D., Ma, A., Li, F., Barsyte-Lovejoy, D., Parton, T., Macnevin, C.J., Liu, F., Gao,
C., Huang, X.P., Kuznetsova, E., Rougie, M., Jiang, A., Pattenden, S.G., Norris, J.L.,
James, L.I., Roth, B.L., Brown, P.J., Frye, S.V., Arrowsmith, C.H., Hahn, K.M., Wang,
G.G., Vedadi, M., and Jin, J. (2013) An orally bioavailable chemical probe of lysine
methyltransferases EZH2 and EZH1. ACS Chem. Biol. 8, 1324-1334.

26. Krishnaiah, S.Y., Wu, G., Altman, B.J., Growe, J., Rhoades, S.D., Coldren, F.,
Venkataraman, A., Olarerin-George, A.O., Francey, L.J., Mukherjee, S., Girish, S.,
Selby, C.P., Cal, S., Er, U., Sianati, B., Sengupta, A., Anafi, R.C., Kavakli, I.H., Sancar,
A., Baur, J.A., Dang, C.V., Hogenesch, J.B., and Weljie, A.M. (2017) Clock regulation of
metabolites reveals coupling between transcription and metabolism. Cell Metab. 35,
961-974.

27. Lachner, M., O'Carroll, D., Rea, S., Mechtler, K., and Jenuwein, T. (2001) Methylation of
histone H3 lysine 9 creates a binding site for HP1 proteins. Nature. 410, 116-120.

28. Larson, K., Yan, S.J., Tsurumi, A., Liu, J., Zhou, J., Gaur, K., Guo, D., Eickbush, T.H.,
and Li, W.X. (2012) Heterochromatin formation promotes longevity and represses
ribosomal RNA synthesis. PLoS Genet. 8, e1002473. doi:
10.1371/journal.pgen.1002473.

29. Lees, E.K., Król, E., Grant, L., Shearer, K., Wyse, C., Moncur, E., Bykowska, A.S.,
Mody, N., Gettys, T.W., and Delibegovic, M. (2014) Methionine restriction restores a
younger metabolic phenotype in adult mice with alterations in fibroblast growth factor 21.
Aging Cell. 13, 817-827.

30. Leng, N., Dawson, J.A., Thomson, J.A., Ruotti, V., Rissman, A.I., Smits, B.M., Haag,
J.D., Gould, M.N., Stewart, R.M., and Kendziorski, C. (2013) EBSeq: an empirical Bayes
hierarchical model for inference in RNA-seq experiments. Bioinformatics. 29, 1035-1043.

31. Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G.,
578   Abecasis, G., Durbin, R., and 1000 Genome Project Data Processing Subgroup. (2009)
579   The sequence alignment/map format and SAMtools. Bioinformatics. 25, 2078-2079.

580   32. Li, B. and Dewey, C.N. (2011) RSEM: accurate transcript quantification from RNA-Seq
581   data with or without a reference genome. BMC Bioinform. doi: 10.1186/1471-2105-12-
582   323.

583   33. Li, S., Swanson, S.K., Gogol, M., Florens, L., Washburn, M.P., Workman, J.L, and
584   Suganuma, T. (2015) Serine and SAM responsive complex SESAME regulates histone
585   modification crosstalk by sensing cellular metabolism. Mol. Cell. 60, 408-421.

586   34. Li, X., Egervari, G., Wang, Y., Berger, S.L., and Lu, Z. (2018) Regulation of chromatin
587   and gene expression by metabolic enzymes and metabolites. Nat. Rev. Mol. Cell Biol.
588   19, 563-578.

589   35. Liu, F., Barsyte-Lovejoy, D., Li, F., Xiong, Y., Korboukh, V., Huang, X.P., Allali-Hassani,
590   A., Janzen, W.P., Roth, B.L., Frye, S.V., Arrowsmith, C.H., Brown, P.J., Vedadi, M., and
591   Jin, J. (2013) Discovery of an in vivo chemical probe of the lysine methyltransferases
592   G9a and GLP. J. Med. Chem. 56, 8931-8942.

593   36. Loyola, A., Bonaldi, T., Roche, D., Imhof, A., and Almouzni, G. (2006) PTMs on H3
594   variants before chromatin assembly potentiate their final epigenetic state. Mol. Cell. 24,
595   309-316.

596   37. Loyola, A., Tagami, H., Bonaldi, T., Roche, D., Quivy, J.P., Imhof, A., Nakatani, Y., Dent,
597   S.Y. and Almouzni, G. (2009) The HP1α-CAF1-SETDB1-containing complex provides
598   H3K9me1 for Suv39-mediated K9me3 in pericentric heterochromatin. Embo Rep. 10,
599   769-775.

600   38. MacLean, B., Tomazela, D.M., Shulman, N., Chamber, M., Finney, G.L., Frewen, B.,
601   Kern, R., Tabb, D.L., Liebler, D.C., and MacCoss, M.J. (2010) Skyline: an open source
602   document editor for creating and analyzing targeted proteomics experiments.
603   Bioinformatics. 26, 966-968.

604   39. Mauger, O., Klinck, R., Chabot, B., Muchardt, C., Allemand, E., and Batsché, E. (2015)
605   Alternative splicing regulates the expression of G9A and SUV39H2 methyltransferases,
606   and dramatically changes SUV39H2 functions. Nucleic Acids Res. 43, 1869-1882.

607   40. Melamud, E., Vastag, L, and Rabinowitz, J.D. (2010) Metabolomic analysis and
608   visualization engine for LC-MS data. Anal. Chem. 82, 9818-9826.

609   41. Mentch S.J. and Locasale J.W. (2016) One-carbon metabolism and epigenetics:
610   understanding the specificity. Ann N. Y. Acad. Sci. 1363, 91-98.

42. Mentch, S.J., Mehrmohamadi, M., Huang, L., Liu, X., Gupta, D., Mattocks, D., Gómez

Padilla, P., Ables, G., Bamman, M.M., Thalacker-Mercer, A.E. Nichenametla, S.N., and

Locasale, J.W. (2015) Histone methylation dynamics and gene regulation occur through

the sensing of one-carbon metabolism. Cell Metab. 22,861-873.

43. Miller, R.A., Buehner, G., Chang, Y., Harper, J.M., Sigler, R., and Smith-Wheelock, M.

(2005) Methionine-deficient diet extends mouse lifespan, slows immune and lens aging,

alters glucose, T4, IGF-I and insulin levels, increases hepatocyte MIF levels and stress

resistance. Aging Cell. 4, 119-125.

44. Mootha, V.K., Lindgren, C.M., Eriksson, K.F., Subramanian, A., Sihag, S., Lehar, J.,

Puigserver, P., Carlsson, E., Ridderstråle, M., Laurila, E., Houstis, N., Daly, M.J.,

Patterson, N., Mesirov, J.P., Golub, T.R., Tamayo, P., Spiegelman, B., Lander, E.S.,

Hirschhorn, J.N., Altshuler, D., and Groop, L.C. (2003) PGC-1alpha responsive genes

involved in oxidative phosphorylation are coordinately downregulated in human diabetes.

Nat. Genet. 34, 267-273.

45. Ni, Z., Ebata, A., Alipanahiramandi E., and Lee, S.S. (2012) Two SET domain containing

genes link epigenetic changes and aging in Caenorhabditis elegans. Aging Cell. 11, 315-

325.

46. Northcott, P.A., Nakahara, Y., Wu, X., Feuk, L., Ellison, D.W., Croul, S., Mack, S.,

Kongkham, P.N., Peacock, J., Dubuc, A., Ra, Y.S., Zilberberg, K., Mcleod, J., Scherer,

S.W., Rao, J.S., Eberhart, C.G., Grajkowska, W., Gillespie, Y., Lach, B., Grundy, R.,

Pollack, I.F., Hamilton, R.L., Van Meter, T., Carlotti, C.G., Boop, F., Bigner, D.,

Gilbertson, R.J., Rutka, J.T., and Taylor, M.D. (2009) Multiple recurrent genetic events

converge on control of histone lysine methylation in medulloblastoma. Nat. Genet. 41,

465-472.

47. Oberdoerffer, P. and Sinclair, D.A. (2007) The role of nuclear architecture in genome

instability and ageing. Nat. Rev. Mol. Cell Biol. 8, 692-702.

48. Orentreich, N., Matias, J.R., DeFelice, A., and Zimmerman, J.A. (1993) Low methionine

ingested by rats extends life span. J. Nutr. 123, 269-274.

49. Peters, A.H., O'Carroll, D., Scherthan, H., Mechtler, K., Sauer, S., Schöfer, C.,

Weipoltshammer, K., Pagani, M., Lachner, M., Kohlmaier, A., Opravil, S., Doyle, M.,

Sibilia, M., and Jenuwein, T. (2001) Loss of the Suv39h histone methyltransferase

impairs mammalian heterochromatin and genome stability. Cell. 107, 323-327.

50. Pinheiro, I., Margueron, R., Shukeir, N., Eisold, M., Fritzsch, C., Richter, F.M., Mittler, G.,

Genoud, C., Goyama, S., Kurokawa, M., Son, J., Reinberg, D., Lachner, M., and

Jenuwein, T. (2012) Prdm3 and Prdm16 are H3K9me1 methyltransferases required for
mammalian heterochromatin integrity. Cell. 150, 948-960.

51. Qi, F., Shang, K., Zhang, J., Chuai, S., Kong, D., Zhou, T., Fu, S., Liang, Y., Li, C.,
Chen, Z., Zhao, Y., Yu, Z., Huang, Z., Hu, M., Ying, H., Chen, Z., Zhang, Y., Xing, F.,
Zhu, J., Xu, H., Zhao, K., Lu, C., Atadja, P., Xia, Z.X., Li, E., and Shou, J. (2015) Histone
methyltransferase SETDB1 regulates liver cancer cell growth through methylation of
p53. Nat. Commun. 6, 8651.

52. Quinlan, A.R. and Hall, I.M. (2010) BEDTools: a flexible suite of utilities for comparing
genomic features. Bioinformatics. 26, 841-842.

53. Rea S., Eisenhaber, F., O'Carroll, D., Strahl, B.D., Sun, Z.W., Schmid, M., Opravil, S.,
Mechtler, K., Pontin, C.P., Allis, C.D., and Jenuwein, T. (2000) Regulation of chromatin
structure by site-specific histone H3 methyltransferases. Nature. 406, 593-599.

54. Rhyu, D.W., Kang, Y.J., Ock, M.S., Eo, J.W., Choi, Y.H., Kim, W.J., Leem, S.H., Yi, J.M.,
Kim, H.S., and Cha, H.J. (2014) Expression of human endogenous retrovirus env genes
in the blood of breast cancer patients. Int. J. Mol. Sci. 15, 9173-9183.

55. Scaffidi, P. and Misteli, T. (2006) Lamin A-dependent nuclear defects in human aging.
Science. 312, 1059-1063.

56. Schmidt, J.A., Rinaldi, S., Scalber, A., Ferrari, P., Achaintre, D., Gunter, M.J., Appleby,
P.N., Key, T.J., and Travis, R.C. (2016) Plasma concentrations and intakes of amino
acids in male meat-eaters, fish-eaters, vegetarians and vegans: a cross-sectional
analysis in EPIC-Oxford cohort. Eur. J. Clin. Nutr. 70, 306-312.

57. Shiraki, N., Shiraki, Y., Tsuyama, T., Obata, F., Miura, M., Nagae, G., Aburatani, H.,
Kume, K., Endo, F., and Kume, S. (2014) Methionine metabolism regulates maintenance
and differentiation of human pluripotent stem cells. Cell Metab. 19, 780-794.

58. Shyh-Chang, N., Locasale, J.W., Lyssiotis, C.A., Zheng, Y., Teo, R.Y.,
Ratanasirintrawoot, S., Zhang, J., Onder, T., Unternaehrer, J.J., Zhu, H., Asara, J.M.,
Daley, G.Q., and Cantley, L.C. (2013) Influence of threonine metabolism on S-
adonsylmethionine and histone methylation. Science. 339, 222-226.

59. Simon, M., Van Meter, M., Ablaeva, J., Ke, Z., Gonzalez, R.S., Taguchi, T., De Cecco,
M., Leonova, K.I., Kogan, V., Helfand, S.L., Neretti, N., Roichman, A., Cohen, H.Y.,
Meer, M.V., Gladyshev, V.N., Antoch, M.P., Gudkov, A.V., Sedivy, J.M., Seluanov, A.,
and Gorbunova, V. (2019) LINE1 derepression in aged wild-type and SIRT6-deficient
mice drives inflammation. Cell Metab. 29, 871-885.

60. Strekalova, E., Malin, D., Weisenhorn, E.M.M., Russell, J.D., Hoelper, D., Jain, A., Coon,
J.J., Lewis, P.W., and Cryns, V.L. (2019) S-adenosylmethionine biosynthesis is a
targetable metabolic vulnerability of cancer stem cells. Breast Cancer Res. Treat. 175,
39-50.

61. Subramanian, A., Tamayo, P., Mootha, V.K., Mukherjee, S., Ebert, B.L., Gillette, M.A.,
Paulovich, A., Pomeroy, S.L., Golub, T.R., Lander, E.S., and Mesirov, J.P. (2005) Gene
set enrichment analysis: a knowledge-based approach for interpreting genome-wide
expression profiles. Proc. Natl. Acad. Sci.102, 15545-15550.

62. Tang, S., Fang, Y., Huang, G., Xu, X., Padilla-Banks, E., Fan, W., Xu, Q., Sanderson,
S.M., Foley, J.F., Dowdy, S., McBurney, M.W., Fargo, D.C., Williams, C.J., Locasale,
J.W., Guan, Z., and Li, X. (2017) Methionine metabolism is essential for SIRT1-regulated
mouse embryonic stem cell maintenance and embryonic development. EMBO J. 36,
3175-3193.

63. Tang, X., Keenan, M.M., Wu, J., Lin, C.A., Dubois, L., Thompson, J.W., Freedland, S.J.,
Murphy, S.K., and Chi, J.T. (2015) Comprehensive profiling of amino acid response
uncovers unique methionine-deprived response dependent on intact creatine
biosynthesis. PLoS Genet. 11, e1005158. doi: 10.1371/journal.pgen.1005158.

64. Towbin, B.D., González-Aguilera, C., Sack, R., Gaidatzis, D., Kalck, V., Meister, P.,
Askjaer, P., and Gasser, S.M. (2012) Step-wise methylation of histone H3K9 positions
heterochromatin at the nuclear periphery. Cell. 150, 934-937.

65. Tvardovskiy, A., Schwämmle, V., Kempf, S.J., Rogowska-Wrzesinska, A., and Jensen,
O.N. (2017) Accumulation of histone variant H3.3 with age is associated with profound
changes in histone methylation landscape. Nucleic Acids Res. 45, 9272-9289.

66. Wang, Z., Yip, L.Y., Lee, J.H.J., Wu, Z., Chew, H.Y., Chong, P.K.W., Teo, C.C., Ang,
H.Y.K., Peh, K.L.E., Yuan, J., Ma, S., Choo, L.S.K., Basri, N., Jiang, X., Yu, Q., Hillmer,
A.M., Lim, W.T., Lim, T.K.H., Takano, A., Tan, E.H., Tan, D.S.W., Ho, Y.S., Lim, B., and
Tam, W.L. (2019) Methionine is a metabolic dependency of tumor-initiating cells. Nat.
Med. 25, 825-837.

67. Watson, G.W., Wickramasekara, S., Palomera-Sanchez, Z., Black, C., Maier, C.S.,
Williams, D.E., Dashwood, R.H., and Ho, E. (2014) SUV39H1/H3K9me3 attenuates
sulforaphane-induced apoptotic signaling in PC3 prostate cancer cells. Oncogenesis. 3,
e131.

68. Wood, J.G., Hillenmeyer, S., Lawrence, C., Chang, C., Hosier, S., Lightfoot, W.,
Mukherjee, E., Jiang, N., Schorl, C., Brodsky, A.S., Neretti, N., and Helfand, S.L. (2010)
Chromatin remodeling in the aging genome of Drosophila. Aging Cell. 9, 971-978.
69. Wood, J.G., Jones, B.C., Jiang, N., Chang, C., Hosier, S., Wickremesinghe, P., Garcia,
M., Harnett, D.A., Burhenn, L., Neretti, N., and Helfand, S.L. (2016) Chromatin-modifying
genetic interventions surpress age-associated transposable element activation and
extend life span in *Drosophila*. Proc. Natl. Acad. Sci. 113, 11277-11282.
70. Wu, S.E., Huskey, W.P., Borchardt, R.T., and Schowen, R.L. (1983) Chiral instability at
the sulfur of S-adenosylmethionine. Biochemistry. 22, 2828-2832.
71. Ye, C., Sutter, B.M., Wang, Y., Kuang, Z., and Tu, B.P. (2017) A metabolic function for
phospholipid and histone methylation. Mol. Cell 66, 180-193.
72. Yu, D., Yang, S.E., Miller, B.R., Wisinski, J.A., Sherman, D.S., Brinkman, J.A.,
Tomasiewicz, J.L., Cummings, N.E., Kimple, M.E., Cryns, V.L., and Lamming, D.W.
(2018) Short-term methionine deprivation improves metabolic health via sexually
dimorphic, mTORC-1 independent mechanisms. FASEB J. 32, 3471-3482.
73. Zhang, Y., Liu, T., Meyer, C.A., Eeckhoute, J., Johnson, D.S., Bernstein, B.E.,
Nusbaum, C., Myers, R.M., Brown, M., Li, W., and Liu, X.S. (2008) Model-based
analysis of ChIP-Seq (MACS). Genome Biol. 9, R137. doi: 10.1186/gb-2008-9-9-r137.
74. Zee, B.M., Levin, R.S., Xu, B., Leroy, G., Wingreen, N.S., and Garcia, B.A. (2010) In
vivo residue-specific histone methylation dynamics. J. Biol. Chem. 285, 3341-3350.
730

Main Figure Titles and Legends

Figure 1. Methionine Restriction Stimulates Dynamic Histone PTM Response

(*A*) Metabolic pathway diagram illustrating changes in Met-cycle metabolite abundance after 24 hours and 5 weeks of Met-restriction in HCT116 cells and C57BL/6J liver, respectively. n=5, error bars represent SD, *p<0.05 (Welch's t-Test). (*B-C*) Plots illustrating global, relative 5mC DNA methylation levels during Met-restriction. n≥3, error bars represent SD, *p<0.05 (Welch's t-Test) (*D-E*) Hierarchichal clustered heatmaps of LC-MS/MS generated $\log_2$ fold-change stoichiometric histone H3 peptide proteoform values relative to 0-hour or chow diet controls in HCT116 cells and C57BL/6J liver, respectively. n≥3, *p<0.05, **p<.01 (Welch's t-Test). See also Table S1 and Figure S1.

Figure 2. Decreased SAM Availability Drives Robust Histone Methylation Response

(*A*) Pathway diagram illustrating experimental approaches to depleting intracellular SAM availability. (*B*) Bar graph of relative *MAT2A* mRNA abundance in HCT116 cells as measured by RT-qPCR. n≥2, error bars represent SD, *p<0.05 (Welch's t-Test). (*C-D*) Dot plots of relative Met and SAM levels in HCT116 cells. n≥4, *p<0.05 (Welch's t-test). (*E-F*) Correlation plot of LC-MS/MS generated $\log_2$ fold-change stoichiometric values for individual histone H3 peptide proteoforms. n=3. (*G*-J) Bar graphs illustrating LC-MS/MS generated $\log_2$ fold-changes for individual H3K9 PTMs. n=3 *p<0.05 (Welch's t-test) See also Table S1, Table S2, and Figure S2.

Figure 3. H3K9me1 Abundance under SAM Depletion is Driven by *de novo* Methylation

(*A-B*) Bar graphs illustrating LC-MS/MS generated $\log_2$ fold-changes for individual H3K9 PTMs in HCT116 cells. n≥2, error bars represent SD, *p<0.05 (Welch's t-Test). (*C*) Bar graph illustrating percent loss in global H3K9me1, as measured by LC-MS/MS, in response to 5μm UNC0642 treatment. n≥2, error bars represent SD, *p<0.05 (Welch's t-Test) (*D*) Scatter plot of relative, total H3 and H3K9me1 cytoplasmic protein abundance in HCT116 cells as measured by western blot. Error bars represent SD, n=3, *p<0.05 (Welch's t-Test). (*E*) Representative western blot images of those used for the quantification presented in Figure 3E. Alpha-Tubulin is used as a cytoplasm marker to confirm efficient sub-cellular fractionation. See Document S1 for REVERT Total Protein Stain images used for normalization. (*F*) Diagram illustrating the

SILAC experimental design. (*G*) Bar graph illustrating contribution of newly synthesized histone

H3 to the total chromatin bound pool after 24 hours of Met-restriction in HCT116 cells as

measured by LC-MS/MS. n≥2, error bars represent SD, *p<0.05 (Welch's t-test). (*H*) Bar graph

illustrating ratio of light:heavy L-Arg isotope presence in H3K9 proteoforms after 24 hours of

Met-restriction in HCT116 cells as measured by LC-MS/MS. n≥2, Error bars represent SD,

*p<0.05 (Welch's t-Test). (*I*) Pie chart depicting sources of H3K9me1 during SAM depletion. (*J*)

Scatter plot of 373 cytoplasmic proteins identified by LC-MS/MS with significant incorporation of

L-Arg $^{+0}$ that were also significantly upregulated after 24 hours of Met-restriction in 0.1% DMSO

treated HCT116 cells. n=4, Statistical significance of p<0.05 (Welch's t-Test). See also Table

S1, Table S3, and Figure S3.

Figure 4. H3K9me1 is Redistributed Over Repetitive and Transposable Elements During SAM

Depletion

(*A-B*) Bar graphs illustrating percent changes in H3K9 PTM peak number at various annotated

loci after 24 hours of Met-restriction in HCT116 cells. (*C-D*) Plots depicting changes in

normalized H3K9 PTM ChIP-sequencing reads to loci of lost H3K9me3 after 24 hours of Met-

restriction in HCT116 cells. (*E*) Bar graphs illustrating percent changes in loci annotation for

regions of lost H3K9me3 experiencing increased relative to decreased H3K9me1 ChIP read

enrichment after 24 hours of Met-restriction in HCT116 cells. (*F*) Representative ChIP-

sequencing track image.

Figure 5. *De novo* H3K9 Mono-methylation Preserves Heterochromatin Stability

(*A*) Bar graph illustrating the percent abundances of 5 distinct nucleosome species following

MNase digestion of HCT116 cells. n≥3, error bars represent SD, *p<0.05 (Welch's t-Test). (*B*)

Representative DNA agarose gel image of those used for the quantification presented in Figure

5A. (*C-E*) Bar graphs illustrating relative mRNA abundances of *LINE* 1, *HERV-K,* and *HERV-R*

in HCT116 cells as measured by RT-qPCR. n≥4, error bars represent SD, *p<0.05 (Welch's t-

Test). See also Table S2.

Figure 6. Adaptive H3K9 Mechanisms Support Epigenetic Persistence Upon Metabolic

Recovery

(*A*) Diagram illustrating the experimental design for replenishing Met and SAM availability after

24 hours of Met-restriction coupled with acute 5 μM UNC0642 or 0.1% DMSO treatment in

HCT116 cells. (*B*) Heatmap of $\log_2$ fold-change stoichiometric values for histone H3 and H4

peptides as measured by LC-MS/MS. n=4, *p<0.05, **p<0.01 (Welch's t-Test). (*C-E*) Bar graphs

illustrating $\log_2$ fold-changes for individual H3K9 PTMs in HCT116 cells as measured by LC-

MS/MS. n=4, error bars represent SD, *p<0.05 (Welch's t-Test). (*F*) PCA diagram of TPM

values. (*G*) EBseq multiple condition comparison diagram. (*H*) Pie chart illustrating percent

allocation of genes identified via RNA-sequencing to all EBseq patterns. (*I-J*) Scatter plots

illustrating NES and FDR q-value for GSEA identified GO terms. (*K-L*) Representative GSEA

running enrichment score figures. See also Table S1 and Figure S4.

Figure 7. Adaptive and Persistent Epigenetic Responses to SAM Depletion are Robust *in vivo*,

Independent of Age

(*A*) Diagram depicting the experimental design for *in vivo* Met-restriction of 6- and 22-month

C57BL/6J mice followed by dietary Met-reintroduction. (*B*) Heatmap of $\log_2$ fold-change

stoichiometric values for histone H3 and H4 peptides in C57BL/6J liver relative to age-matched

controls as measured by LC-MS/MS. n≥5, *p<0.05, **p<0.01 (Welch's t-Test). (*C*) Bar graph

illustrating $\log_2$ fold-changes for individual H3K9 PTMs in liver from Met-restricted C57BL/6J

mice relative to age-matched controls as measured by LC-MS/MS. n≥5, error bars represent

SD, *p<0.05 (Welch's t-Test). (*D-E*) Bar graphs illustrating relative mRNA abundances of *LINE*

1, *SINE B1*, and *SINE B2* in C57BL/6J liver as measured by RT-qPCR. n≥5, error bars

represent SD, *p<0.05 (Student's t-Test). (*F*) Bar graph illustrating $\log_2$ fold-changes for

individual H3K9 PTMs in liver from C57BL/6J relative to age-matched controls as measured by

LC-MS/MS. n≥5, error bars represent SD, *p<0.05, (Welch's t-Test). (*G-H*) Bar graphs

illustrating relative mRNA abundances of *LINE 1*, *SINE B1*, and *SINE B2* in C57BL/6J liver as

measured by RT-qPCR. n≥5, error bars represent SD, *p<0.05 (Student's t-Test). See also

Table S1, Table S2, and Figure S5.

Supplemental Figure Titles and Legends

<u>Figure S1, Related to Figure 1</u>

(*A-D*) Scatter plots of relative abundance values for key Met-cycle metabolites in HCT116 cells

as measured by LC-MS. n=5, error bars represent SD, *p<0.05 (Welch's t-Test). (*E-J*)

Correlation plots of LC-MS/MS generated $\log_2$ fold-change stoichiometric values for individual

histone H3 peptide proteoforms. n≥3. See also Table S1.

<u>Figure S2, Related to Figure 2</u>

(*A*) Diagram illustrating the experimental design for determining the conservation of H3K9

regulation in response to SAM depletion across various biological systems. (*B-F*) Bar graphs

illustrating $\log_2$ fold-changes for individual H3K9 in response to *in vitro* SAM depletion. n=3, error

bars represent SD, n=3, *p<0.05 (Welch's t-Test). See also Table S1.

<u>Figure S3, Related to Figure 3</u>

(*A-C*) Scatter plots of relative HMT mRNA abundance in HCT116 cells as measured by RT-

qPCR. n=4, error bars represent SD, *p<0.05 (Welch's t-Test). (*D*) Heatmap of $\log_2$ fold-change

stoichiometric values for unique histone H3K9 peptide proteoforms in HCT116 cells. n=4,

*p<0.05 (Welch's t-Test). (*E*) Scatter plot of relative cell counts for attached HCT116 cells. n=4,

error bars represent SD, *p<0.05 (Welch's t-Test). (F) Bar graph of relative H3K9me1

abundance in HCT116 cells as measured by western blot with accompanied representative blot

image. See Document S1 for REVERT Total Protein Stain image used for normalization. Error

bars represent SD, n≥3, *p<0.05 (Welch's t-Test). (*G*) Bar graphs illustrating $\log_2$ fold-changes

for individual H3K9 PTMs. n≥3, *p<0.05 (Welch's t-Test). (*H*) Bar graph showing average

incorporation of L-Arg^{+10} in H3K9 peptide proteoforms. n=4, error bars represent SD. (*I*) Density

plot illustrating the percentage of Met residue frequency in the primary amino acid sequence of

all identified cytoplasmic proteins as well as those determined to be preferentially translated.

See also Table S1, Table S2, and Table S3.

<u>Figure S4, Related to Figure 6</u>

(*A*) GSEA heatmap of relative gene expression across positive NES GO terms for Pattern 2. (B)

GSEA set-to-set heatmap illustrating gene enrichment overlap across positive NES GO terms

for Pattern 2. (*C*) GSEA heatmap of relative gene expression across negative NES GO terms

for Pattern 2. (D) GSEA set-to-set heatmap illustrating gene enrichment overlap across negative

NES GO terms for Pattern 2.

<u>Figure S5, Related to Figure 7</u>

(*A-E*) Bar graphs illustrating $\log_2$ fold-changes for individual H3K4, H3K18/K23, H3K27, H3K36,

and H4 PTMs in liver from Met-restricted C57BL/6J mice relative to age-matched controls as

measured by LC-MS/MS. $n \geq 5$, error bars represent SD, $*p < 0.05$ (Welch's t-Test). (F) Bar graph

illustrating changes in fat, lean mass, and total body mass (BW) after 3 weeks of control or Met-

restricted diet feeding. n=14, error bars represent SEM, $*p < 0.05$ (two-way ANOVA). (*G*)

Glucose tolerance test in 6- and 22-month C57BL/6J mice after 2 weeks of control or Met-

restricted diet feeding. n=14, error bars represent SEM, $*p < 0.05$ (two-way ANOVA). (*H*) Bar

graph illustrating changes in fat, lean mass, and total body mass (BW) after 5 weeks of control

diet feeding following either 3 weeks of control or Met-restricted diet feeding. $n \geq 6$, error bars

represent SEM, $*p < 0.05$ (two-way ANOVA). (*I*) Glucose tolerance test in 6- and 22-month

C57BL/6J mice after 4 weeks of control diet feeding following either 3 weeks of control or Met-

restricted diet feeding. $n \geq 7$, error bars represent SEM, $*p < 0.05$ (two-way ANOVA). See also

Table S1 and Table S2.

Star Methods Text

<u>Contact for Reagent and Resource Sharing:</u>

Further information and requests for resources and reagents should be directed to and will be

fulfilled by the Lead Contact, John M. Denu (john.denu@wisc.edu).

<u>Experimental Model and Subject Details</u>

Animals and Diets:

All experiments involving animals were approved by the Institutional Animal Care and Use
Committee of the William S. Middleton Memorial Veterans Hospital, Madison WI. To test the
effect of Met deprivation in the context of young mice, 9-week-old male C57BL/6J mice were
purchased from The Jackson Laboratory (Bar Harbor, ME, USA) and placed on either an amino
acid defined Control diet (TD.01084, Envigo, Madison, WI, USA) or a Met-deficient diet
(TD.140119, Envigo, Madison, WI, USA) their respective diets at 14 weeks of age; full diet
compositions for these diets have been previously described (PMID: 29401631). After 5 weeks
on the diet, mice were sacrificed following an overnight fast and liver was flash frozen for
analysis as described.
904
To compare the effect of Met restriction on young and old mice, C57BL/6J. Nia male mice were
obtained from the National Institute on Aging Aged Rodent Colony, and housed in our animal
facility for 3-4 months until reaching 5 and 21 months of age. All animals were then placed on
the Control diet (TD.01084) for 4 weeks, until the mice were 6 and 22 months of age. Mice were
then randomized to either remain on the Control diet, or were placed on the Met-deficient diet
(TD.140119) for three weeks. Liver was collected and flash-frozen in liquid nitrogen harvested
from mice euthanized after an approximately 16-hour fast.
912
913 *Cell Lines:*
914
915 HCT116, HEK-293, HEK-293T, HMCF7, Panc1, HepG2, and HepA1 cell lines were cultured at
916 37°C with 5.0% CO_2.
917
918 <u>Method Details</u>
919
920 *Metabolite Extraction:*
921
For HCT116 cells, ~1.5x10^6 cells were quickly rinsed 3 times with 1.5 ml ice-cold PBS pH 7.4
before the addition of 1.5 ml of -80°C 80:20 MeOH:H_2O extraction solvent. Cells were incubated
with extraction solvent at -80°C for 15 minutes. Next, cells were scraped off their dish and
transferred into a 2 ml microcentrifuge Eppendorf tube which was followed by a 5-minute,
maximum speed centrifugation at 4°C. The supernatant was transferred to a new 2 ml
microcentrifuge Eppendorf tube. Next, 0.5 ml extraction solvent was added to the remaining
pellet, vortexed, and again centrifugated at maximum speed for 5 minutes at 4°C. The 2

supernatants from each extraction were pooled and dried completely using a Thermo Fisher
Savant ISS110 SpeedVac. Dried metabolite extracts were resuspended in 80 μl of 85%
acetonitrile following microcentrifugation for 5 minutes at maximum speed at 4°C to pellet any
remaining insoluble debris. The supernatant was then transferred to a glass vial for LC-MS
analysis.
934
For *M. musculus* liver tissue, an average tissue weight of 75 mg was powdered in liquid nitrogen
using a mortar and pestle. Powdered tissue was transferred to an individual 1.5 ml
microcentrifuge Eppendorf tube and incubated with 1 ml -80°C 80:20 MeOH:H_2O extraction
solvent on dry ice for 5 minutes post-vortexing. Tissue homogenate was centrifuged at
maximum speed for 5 minutes at 4°C. Supernatant was transferred to a 15 ml tube after which
the remaining pellet was resuspended in 0.8 ml -20°C 40:40:20 ACN:MeOH:H_2O extraction
solvent and incubated on ice for 5 minutes. Tissue homogenate was again centrifuged at
maximum speed for 5 minutes at 4°C after which the supernatant was pooled with the
previously isolated metabolite fraction. The 40:40:20 ACN:MeOH:H_2O extraction was then
repeated as previously described. Next, 550 μl of pooled metabolite extract for each sample
was transferred to a 1.5 ml microcentrifuge Eppendorf tube and completely dried using a
Thermo Fisher Savant ISS110 SpeedVac. Dried metabolite extracts were resuspended in 450 μl
of 85% ACN following microcentrifugation for 5 minutes at maximum speed at 4°C to pellet any
remaining insoluble debris. Supernatant was then transferred to a glass vial for LC-MS analysis.
949
950 *LC-MS Metabolite Analysis:*
951
Each prepared metabolite sample was injected onto a Thermo Fisher Scientific Vanquish
UHPLC with a Waters XBridge BEH Amide column (100 m x 2.1 mm, 3.5 μm) coupled to a
Thermo Fisher Q-Exactive mass spectrometer. Mobile phase (A) consisted of 97% H_2O, 3%
ACN, 20 mM ammonium acetate, and 15 mM ammonium hydroxide pH 9.6. Organic phase (B)
consisted of 100% ACN. Metabolites were resolved using the following linear gradient: 0 min,
85% B, 0.15 ml/min; 1.5 min, 85% B, 0.15 ml/min; 5.5 min, 40% B, 0.15 ml/min; 10 min, 40% B,
0.15 ml/min; 10.5 min, 40% B, 0.3 ml/min; 14.5 min, 40% B, 0.3 ml/min; 15 min, 85% B, 0.15
ml/min; 20 min, 85% B, 0.15 ml/min. The mass spectrometer was operated in positive ionization
mode with a MS1 scan at resolution = 70,000, automatic gain control target = 3 x 10^6, and scan
range = 60-186 m/z and 187-900 m/z. This protocol was adapted from Mentch et al., 2015.
Individual metabolite data were called using MAVEN (Melamud et al., 2010; Clasquin et al.,

2012) with retention times empirically determined in-house. Peak Area Top values were

analyzed to determine metabolite expression.

LC-MS DNA Methylation Assay:

500 ng of cytosine, 5mC, and 5hmC double-stranded DNA standards from Zymo Research

were incubated with Zymo Research DNA Degradase Plus for 3 hours at 37°C following enzyme

inactivation at 70°C for 20 minutes. 75 µl of ACN was added to each standard reaction to bring

the final volume to 100 µl. DNA standards were centrifugated at 21,100xg for 5 minutes at 4°C

after which 75 µl of supernatant was transferred to a glass vial for LC-MS analysis. A range of

0.1 to 100 ng of digested DNA was analyzed via LC-MS to determine each nucleoside's linear

range of detection.

Genomic DNA extracted using a Promega Wizard Genomic DNA Purification Kit after which 1

µg was incubated with Zymo Research DNA Degradase Plus at 37°C following enzyme

inactivation at 70°C for 20 minutes. 175 µl of ACN was added to each reaction to bring the final

volume to 200 µl. DNA samples were centrifugated at 21,100xg for 5 minutes at 4°C after which

150 µl of supernatant was transferred to a glass vial for LC-MS analysis.

Each prepared nucleoside sample was injected onto a Thermo Fisher Scientific Vanquish

UHPLC with a Waters XBridge BEH Amide column (100 m x 2.1 mm, 3.5 µm) coupled to a

Thermo Fisher Q-Exactive mass spectrometer run in positive ionization mode. Mobile phase (B)

consisted of H_2O, 3% ACN, 20 mM ammonium acetate, and 15 mM ammonium hydroxide pH

9.6. Organic phase (B) consisted of 100% ACN. Nucleosides were resolved using the following

linear gradient: 0 min, 85% B, 0.15 ml/min; 1.5 min, 85% B, 0.15 ml/min; 5.5 min, 40% B, 0.15

ml/min; 10 min, 40% B, 0.15 ml/min; 10.5 min, 40% B, 0.3 ml/min; 14.5 min, 40% B, 0.3 ml/min;

15 min, 85% B, 0.15 ml/min; 20 min, 85% B, 0.15 ml/min. The mass spectrometer was operated

in positive ionization mode with a MS1 scan at resolution = 70,000, automatic gain control target

= 3×10^6, and scan range = 200-300 m/z, followed by a DDA scan with a loop count of 5. DDA

settings were as follows: window size = 2.0 m/z, resolution = 17,500, automatic gain control

target = 1×10^5, DDA maximum fill time = 50 ms, and normalized collision energy = 30. This

protocol was adapted from Mentch et al., 2015. Individual metabolite data were called using

MAVEN (Melamud et al., 2010; Clasquin et al., 2012) with retention times empirically

determined in-house. Peak Area Top values were analyzed to determine metabolite expression
with dAdenosine being utilized for internal normalization (m/z dCytosine: 228.0978, m/z d5mC:
242.1135, m/z d5hmC: 258.108, m/z dAdenosine: 252.109).
999
1000 *Histone Isolation and Chemical Derivatization:*
1001
Tissue culture cell pellets were resuspended in 800 μl of ice-cold Buffer A (10 mM Tris-HCl pH
7.4, 10 mM NaCl, and 3 mM $MgCl_2$) supplied with protease and histone deacetylase inhibitors
(10 μg/ml leupeptin, 10 μg/ml aprotinin, 100 μM phenylmethylsulfonyl fluoride, 10 mM
nicotinamide, 1 mM sodium-butyrate, and 4 μM trichostatin A) followed by 80 strokes of light
pestle homogenization in a 1 ml Wheaton dounce homogenizer. Cell homogenate was then
transferred to a 1.5 ml microcentrifuge Eppendorf tube and centrifugated at 800xg for 10
minutes at 4°C to pellet nuclei. The supernatant was either transferred to a fresh 1.5 ml
Eppendorf tube or discarded. The nuclei pellet was resuspended in 500 μl ice-cold PBS pH 7.4
followed by centrifugation at 800xg for 10 minutes at 4°C. The supernatant was discarded and
nuclei were again washed with 500 μl ice-cold PBS pH 7.4. Pelleted nuclei were then
resuspended in 500 μl of 0.4N H_2SO_4 and rotated at 4°C for 4 hours. Samples were
centrifugated at 3,400xg for 5 minutes at 4°C to pellet nuclear debris and precipitated non-
histone proteins. The supernatant was transferred to a new 1.5 ml Eppendorf tube after which
125 μl of 100% trichloroacetic acid was added and incubated overnight on ice at 4°C. Samples
were centrifugated at 3,400xg for 5 minutes at 4°C to pellet precipitated histone proteins. The
supernatant was discarded, after which the precipitant was washed with 1 ml ice-cold acetone
+0.1% HCl. Samples were centrifugated at 3,400xg for 2 minutes at 4°C and the supernatant
was discarded. This process was repeated except precipitant was washed with 100% ice-cold
acetone. Residual acetone was allowed to evaporate at room temperature for 10 minutes after
which the dried precipitant was dissolved in 125 μl H_2O. Samples were then centrifugated at
21,100xg for 5 minutes at 4°C to pellet any remaining insoluble debris. Supernatant containing
purified histone was then transferred to a new 1.5 ml Eppendorf tube and stored at -20°C until
needed for future analyses.
1025
For *M. musculus* liver tissue, an average tissue weight of 50 mg was homogenized in 800μl ice-
cold Buffer supplemented with protease and HDAC inhibitors (listed above) using a 1 ml
Wheaton dounce homogenizer (20 strokes of the loose pestle followed by 20 strokes of the tight

pestle) and strained through a 100 μM filter before being transferred to a new 1.5 ml Eppendorf

tube. All remaining steps for completing the histone isolation were performed as described

previously in the tissue culture isolation protocol.

1032

To prepare purified histone proteins for LC-MS/MS analysis, 5 μg of histone was diluted with

H_2O to a final volume of 9 μl. 1 μl of 1 M triethylammonium bicarbonate was added to each

sample to buffer the solution to a final pH of 7-9. Next, 1 μl of 1:100 propionic anhydride:H_2O

was added to each sample and allowed to incubate for 2 minutes at room temperature. The

propionylation reaction was then quenched via the addition of 1 μl 80 mM hydroxylamine which

was allowed to incubate for 20 minutes at room temperature. Next, propionylated histones were

digested with 0.1 μg trypsin for 4 hours at 37^{o}C. Upon completion of trypsin digestion, 5 μl of .02

M NaOH was added to bring the final pH between 9-10. Propionylated, trypsin digested histone

peptides were then N-terminally modified with 1 μl 1:50 phenyl isocyanate:ACN for 1-hour at

37^{o}C. Modified peptides were desalted and eluted-off of Empore™C18 extraction membrane.

Eluted samples were dried completely using a Thermo Fisher Scientific Savant ISS110

SpeedVac, resuspended in 40 μl sample diluent (94.9% H_2O, 5% ACN, .1% TFA), and

transferred into glass vials for LC-MS/MS analysis.

1046

1047   *Histone Proteomics Analysis:*

1048

Derivatized histone peptides were injected onto a Dionex Ultimate3000 nanoflow HPLC with a

Waters nanoEase UPLC C18 column (100 m x 150 mm, 3μm) coupled to a Thermo Fisher Q-

Exactive mass spectrometer at 700 nL/ min. Aqueous phase (A) consisted of H_2O + 0.1% formic

acid while the mobile phase (B) consisted of acetonitrile + 0.1% formic acid (B). Histone

peptides were resolved using the following linear gradient: 0 min, 2.0% B; 5 min, 2.0% B; 65

min, 35% B; 67 min, 95% B; 77 min, 95% B; 79 min, 2.0% B; 89 min, 2.0% B. Data was

acquired using data-independent acquisition (DIA) mode. The mass spectrometer was operated

with a MS1 scan at resolution = 35,000, automatic gain control target = 1 x 10^6, and scan range

= 390-910 m/z, followed by a DIA scan with a loop count of 10. DIA settings were as follows:

window size = 10 m/z, resolution = 17,500, automatic gain control target = 1 x 10^6, DIA

maximum fill time = AUTO, and normalized collision energy = 30.

1060

DIA Thermo .raw files were imported into Skyline open source proteomics software (MacLean et
al., 2010) and matched against a pre-constructed peptide database containing unique peptide
proteoforms from H3 and H4. The identity of each individually called peak was verified by hand
after which total MS1 peak area values were exported for downstream analyses. To control for
differences in ionization efficiencies across histone peptides, all peptides possessing an
identical primary sequence were analyzed as a peptide family which enabled the calculation of
stoichiometric values for each unique peptide proteoform within the family. Stoichiometric values
were compared across conditions to identify altered histone peptide abundances. Furthermore,
combinatorial peptide stoichiometry values were deconvoluted by totaling the stoichiometry
values for all peptides possessing a given PTM independent of any other PTMs that may be
present on other residues of the same primary peptide. A Welch's T-Test with a p-value cutoff of
0.05 was used to determine statistical significance.
1073
1074 Hierarchical clustering was performed in MATLAB R2016a using the 'clustergram' command.
1075 Pearson's Correlation Coefficients were calculated in MATLAB R2016a using the 'corr'
1076 command.
1077
1078 *RNAi Mediated Inhibition of MAT2A and H3K9 HMT Expression:*
1079
1080 For 35 mm tissue culture dishes, 1.5×10^5 HCT116 cells were reverse transfected with 30 pmol
1081 of targeted or control RNAi using 5 μl of Lipofectamine RNAiMAX in addition to 500 μl Opti-
1082 MEM and 2.5 ml RPMI-1640 +10% FBS. MAT2A-RNAi treated HCT116 cells used for LC-MS
1083 metabolite analysis were cultured using dialyzed FBS.
1084
1085
1086 *Methionine Restriction of Tissue Culture Cells:*
1087
All tissue culture cells were initially cultured in RPMI-1640 +10% FBS at 37°C, 5% CO_2. One
day before Met-restriction, 3.5×10^6 cells were seeded into 100 mm tissue culture dishes. One-
hour before the restriction began, RPMI-1640 was replaced with fresh, Met replete media. At the
time of Met-restriction, cells were rinsed 2x with 3 ml of sterile PBS pH 7.4 followed by addition
of RPMI-1640 void of Met. Any small molecule inhibitors or vehicle controls would be added at
this time.
1094

1095 For experiments in which Met-restriction was followed by Met repletion, HCT116 cells were

1096 washed with PBS pH 7.4 2x to maximize small molecule inhibitor and DMSO removal before

1097 reintroduction of Met replete RPMI-1640.

1098

1099 HCT116 cells cultured for LC-MS metabolite analysis were cultured in RPMI-1640 containing

1100 10% dialyzed FBS beginning 1-hour prior to beginning Met-restriction.

1101

1102 *SILAC in Tissue Culture:*

1103

197 ml of SILAC RPMI-1640 +10% DFBS void of L-Arg and L-Lys was supplemented with 50mg

isotopically heavy $3C_6$ $15N_4$ L-Arg HCl and 9.43mg isotopically light L-Lys to obtain final

concentrations of 1.149 mM and 0.219 mM, respectively. HCT116 cells were cultured in the

SILAC media for 7 days (or 9 doubling cycles at a reported doubling time of 21 hours) after

which cells were washed 3x with 5 ml PBS pH 7.4 before replacing the SILAC media with

isotopically light L-Arg and L-Lys RPMI-1640 +10% DFBS void of Met. Half of the cells were

treated with 5 μM UNC0642 or 0.1% DMSO at this time. Furthermore, 0-hour control HCT116

cells which were never exposed to isotopically light L-Arg, Met-restricted RPMI-1640 were

harvested. HCT116 cells cultured in the Met-restricted media were harvested after 24 hours.

1113

1114 *SILAC Shotgun Proteomics Analysis:*

1115

50 μg of cytoplasmic protein extract from the previously described SILAC Met-restriction

experiment were diluted with H_2O into a final volume of 75 μl. Next, dithiothreitol was added to

each sample at a final concentration of 10 mM, followed by a 15-minute incubation at $95^{\circ}C$. A

sonication water bath was used to resolubilize precipitated proteins post-heat denaturation.

Iodoacetamide was then added to a final concentration of 20 mM, followed by a 20-minute

incubation at room temperature protected from light. Alkylated proteins were precipitated with

337 μl ice-cold acetone for 20 minutes at $-20^{\circ}C$. Samples were centrifuged at 21,000xg for 15

minutes at $4^{\circ}C$, the supernatant was discarded, and precipitated protein was resuspended in

400 μl of urea buffer (2 M urea and 100 mM ammonium bicarbonate pH 8). Precipitated proteins

were again solubilized using a sonication water bath. Solubilized proteins were digested into

peptides using 0.8 μg trypsin at $37^{\circ}C$ overnight. Upon digest completion, samples were desalted

and eluted-off Empore™C18 extraction membrane. Eluted samples were dried completely using

a Thermo Fisher Scientific Savant ISS110 SpeedVac, after which being resuspended in 50 μl

sample diluent (94.9% H_2O, 5% ACN, .1% TFA) and transferred to glass vials for LC-MS/MS

analysis.

1131

Each prepared cytoplasmic peptide sample was injected onto a Dionex Ultimate3000 nanoflow

HPLC with a Waters nanoEase UPLC C18 column (100 m x 150 mm, $3\mu m$) coupled to a

Thermo Fisher Q-Exactive mass spectrometer at 800 nL/ min. Aqueous phase (A) consisted of

H_2O + 0.1% formic acid while the mobile phase (B) consisted of acetonitrile + 0.1% formic acid

(B). Histone peptides were resolved using the following linear gradient: 0 min, 5.0% B; 5 min,

5.0% B; 120 min, 30% B; 122 min, 95% B; 127 min, 95% B; 129 min, 5.0% B; 145 min, 5.0% B.

Data was acquired using data-dependent acquisition (DDA) mode. The mass spectrometer was

operated with a MS1 scan at resolution = 70,000, automatic gain control target = 1×10^6, and

scan range = 400-1000 m/z, followed by a DDA scan with a loop count of 20. DDA settings were

as follows: window size = 2.0 m/z, resolution = 17,500, automatic gain control target = 2×10^5,

DDA maximum fill time = 150 ms, and normalized collision energy = 28.

1143

1144   *MaxQuant Proteomics Analysis:*

1145

1146   Thermo .raw DDA files were analyzed using MaxQuant v1.6.2.6 (Cox et al., 2014) using the

1147   default settings with the following exceptions: Labels: 1, Arg10; Instrument: Orbitrap; LFQ: LFQ

1148   min. ratio count: 2, LFQ min. number of neighbors: 3, LFQ average number of neighbors: 6;

1149   FASTQ File: Proteome: *H. Sapiens* Proteome ID UP000005640.

1150

Identified proteins with a Q-value of 0.01 or above were removed. For the remaining proteins,

LFQ light (L-Arg^{+0}) and heavy (L-Arg^{+10}) values were summed to generate LFQ total protein

abundance values. LFQ light values were divided by the corresponding LFQ total value to

generate a percent abundance of LFQ light species for each identified protein. Welch's t-Tests

with a p-value cut off of 0.05 were used to determine significant accumulation of LFQ light

species in Met-restricted HCT116 samples relative to 0-hour controls. For proteins determined

to have significant LFQ light species abundance after Met-restriction, LFQ total values were

used to determine the corresponding change in overall protein abundance. Up to 2 null-values

were removed from replicates of an experimental group to account for missed identifications. If

3 of the 4 replicates for a single experimental group contained null-values, that protein was

removed from all experiment groups and therefore excluded from any downstream analyses. A

Welch's t-Test with a p-value cut off of 0.05 was used to determine if an identified protein was
significantly more abundant after Met-restriction relative to 0-hour controls.

1164

1165 *SILAC Histone Proteomics Analysis:*

1166

Histones were isolated, chemically derivatized, and analyzed by LC-MS/MS as described in
"Histone Isolation and Chemical Derivatization" and "Histone Proteomics Analysis." DDA
Thermo .raw files of histone proteins with incorporated isotopically heavy $3C_6$ $15N_4$ L-Arg were
used to construct an H3K9/K14 peptide library using Skyline opensource proteomics software
(MacLean et al., 2010). DIA files were then matched against both isotopically heavy and light L-
Arg peptide libraries after which peptide identities were confirmed by hand. MS1 intensity values
for both isotopically heavy and light L-Arg peptides were summed to generate total MS1 peptide
proteoform intensity values which were then used to calculate the percent of total MS1 peptide
proteoform signal which was contributed by isotopically heavy peptide proteoforms. Welch's T-
Tests with a p-value cutoff of 0.05 were used to determine significant differences in total MS1
peptide proteoform stoichiometric values as well as the percent contribution of isotopically
heavy peptide proteoforms to total peptide proteoform intensity values.

1179

1180 *Chromatin Immunoprecipitation and Sequencing:*

1181

30×10^6 HCT116 cells were resuspended in 1 ml MNase digestion buffer (50 mM Tris-HCl pH
7.6, 1 mM $CaCl_2$, and 0.2% Triton X-100) after which 330 μl of the cell suspension was
aliquoted into individual 1.5 ml tubes. Each tube was treated with either 45, 52.5, or 60 units of
Worthington MNase for 5 minutes at 37°C shaking at 400 rpm to generate mono-nucleosomes.
MNase digestions were stopped via the addition of EDTA to a final concentration of 5mM.
Digested cells were then pooled and lysed at 4°C using a Covaris S220 Focused-ultrasonicator
with the following settings: Peak Power = 75, Duty Factor = 5, Cycles/Burst = 100, Duration =
30" on/30" off x2. Insoluble debris was centrifuged at 10,000 rpm for 10 minutes at 4°C after
which the supernatant containing prepared nucleosomes was transferred to a new tube.
Glycerol was added to a final concentration of 5% prior to -80°C storage. For input controls, 43
μl of prepared nucleosomes were aliquoted and saved for further processing at -80°C.

1193

Next, 860 μl of prepared nucleosomes was incubated with 5 μg of the desired antibody, rotating
overnight at 4°C. The nucleosome/antibody solution was added to 50 μl of prepared M-280

Sheep Anti-Rabbit IgG superparamagnetic Dynabeads™ and allowed to rotate at 4°C for 4

hours. Conjugated beads were then washed for 10 minutes, rotating at 4°C under the following

conditions: 3x 1 ml RIPA (10 mM Tris pH 7.6, 1 mM EDTA, 0.1% SDS, 0.1% Na-deoxycholate,

1% Triton X-100) buffer, 2x 1 ml RIPA buffer +0.3 M NaCl, 2x 1 ml LiCl buffer (0.25 M LiCl,

0.5% NP-40, 0.5% Na-deoxycholate), 1x with 1 ml 1xTE +50 mM NaCl. Following the final

wash, beads were centrifugated at 4°C for 3 minutes at 960xg after which the supernatant was

carefully removed and discarded.

1203

To elute immunoprecipitated nucleosome species, beads were resuspended in 210 μl of elution

buffer (50 mM Tris-HCl pH 8.0, 10 mM EDTA, 1% SDS) and incubated at 65°C for 30 minutes

shaking at 400 rpm. Beads were resuspended every 2 minutes via brief vortexing. Next samples

were centrifugated at 16,000xg for 1 minute at room temperature after which 200 μl of

supernatant was transferred to a new 1.5 ml Eppendorf tube. Eluted nucleosome species were

then treated with 2 μl Thermo Fisher Scientific Proteinase K for 2 hours at 55°C. DNA was

recovered from the Proteinase K digestion using the QIAquick PCR Purification Kit.

1211

1212   DNA libraries were prepared by the University of Wisconsin-Madison Biotechnology Center

1213   using the Illumina TruSeq DNA Nano kit and then sequenced across 4 lanes of an Illumina

1214   1x100 HiSeq2500 High Throughput flowcell.

1215

FASTQ files were aligned unpaired using bowtie2-2.1.0 (Langmead and Salzberg, 2012)

against human genome 19. Bowtie2 output SAM files were converted to BAM files, sorted and

indexed using samtools 1.5 (Li et al., 2009). Broad peaks were called from sorted BAM files

using MACS2-2.1.1.20160309 (Zhang et al., 2008) with a Q-value cutoff of 0.1. Input DNA was

used as the control file. All identified peaks possessing a $\log_2$ fold-change below 1.75 were

discarded prior to downstream analyses. Bedtools v2.25.0 (Quinlan and Hall, 2010) was used to

identify unique and overlapping across samples and also used to assign raw mapped reads to

pre-specified genomic loci. Genomic loci were annotated using HOMERv4.9 (Heinz et al.,

2010).

1225

1226   *Quantitative PCR:*

1227

1228 RNA was extracted from HCT116 cells using Thermo Fisher Scientific TRIzol and from liver

1229 tissue using Sigma TRI Reagent, both by following the manufacturer's instructions. The

1230 concentration and purity of RNA were determined by absorbance at 260/280 nm.

1231

1232 cDNA was prepared from 1 μg of RNA using RevertAid Reverse Transcriptase for HCT116 RNA

1233 and Superscipt III for liver RNA. Oligo dT primers were used in both cDNA synthesizing

1234 reactions.

1235

1236 HCT116 cDNA was analyzed on a Roche 480 Light Cycler System with PerfeCTa SYBR Green

1237 SuperMix while cDNA generated from liver was analyzed on an Applied Biosystems StepOne

1238 Plus instrument with Sybr Green PCR Master Mix. HCT116 cDNA qPCR reactions were

1239 normalized internally using GAPDH while liver cDNA qPCR reactions were normalized internally

1240 using actin. Primer sequences are available in Table S3.

1241

1242 *RNA sequencing*

1243

RNA libraries were prepared by Novogene using NEBNext adaptors followed by paired end

2x150 sequencing on their Illumina Platform. FASTQ files were trimmed using Trimmomatic

v0.39 (Bolger et al., 2014) and aligned using RSEM v1.2.4 (Li et al., 2011) against the human

genome 18 reference transcriptome. RSEM .genes output files were used for Ebseq

EBMultiTest multiple comparison analysis (Leng et al., 2013). Gene lists associated with

individual EBMultiTest patterns were further analyzed with GSEA 2 (Tamayo et al., 2005;

Mootha et al., 2003)

1251

1252 *MNase Accessibility Assay:*

1253

6×10^6 HCT116 cells were resuspended in 1 ml of digestion buffer (50 mM Tris-HCl pH 7.6, 1

mM $CaCl_2$, and 0.2% Triton X-100) in addition to 15 unites of Worthington MNase. MNase

treated cell suspensions were incubated for 5 minutes at 37°C shaking at 400 rpm. MNase

digestions were stopped via the addition of EDTA to a final concentration of 5 mM. Digested

cells were then pooled and lysed at 4°C using a Covaris® S220 Focused-ultrasonicator with the

following settings: Peak Power = 75, Duty Factor = 5, Cycles/Burst = 100, Duration = 30" on/30"

off x2. Insoluble debris was pelleted via centrifugation at 10,000 rpm for 10 minutes at 4°C after

which the supernatant was transferred to a new tube. 150 μl of digested chromatin was treated

with 2 μl of Thermo Fisher Scientific Proteinase K for 90 minutes at 55^{o}C. DNA was then

purified using the QIAquick PCR Purification Kit. Next, 1 μg of purified DNA was separated on a

2% agarose gel and quantified using GelAnalyzer 2010a.

1265

1266   *Western-Blotting and Image Quantification:*

1267

Protein separated on SDS-PAGE gels were transferred onto 0.2 μm nitrocellulose membrane

and stained with 5 ml of LI-COR REVERT Total Protein Stain for 5 minutes at room temperature

before being blocked in 0.1% fat-free milk PBS pH 7.4 for 1 hour. Primary antibodies were

diluted 1:1000 in 5% fat-free milk PBST pH 7.4 and incubated with the membrane overnight at

4^{o}C. LI-COR secondary antibodies were used at a 1:15,000 dilution in 5% fat-free milk PBST pH

7.4 and incubated with the membrane for 1 hour at room temperature. Membranes were imaged

using an Odyssey Infrared Imager (model no. 9120). Densitometry was performed using Image

Studio™ Lite software. All data was normalized to REVERT™ Total Protein Stain.

1276

1277   *In vivo Tests:*

1278

1279   For glucose tolerance tests, mice were fasted overnight for 16 hours and then injected with

1280   glucose (1 g/kg) intraperitoneally as previously described (Arriola Apelo et al., 2016). Glucose

1281   measurements were taken using a Bayer Contour blood glucose meter and test strips. Mouse

1282   body composition was determined using an EchoMRI Body Composition Analyzer (EchoMRI™,

1283   Houston, TX, USA) according to the manufacturer's procedures.

1284

1285   <u>Quantification and Statistical Analysis</u>

1286

1287   Explanations of statistical analyses and parameters are present in the main and supplemental

1288   figure legends as well as in 'Method Details" where appropriate.

1289

1290   <u>Data and Software Availability</u>

1291

1292   RNA-sequencing as well as H3K9me1 and H3K9me3 ChIP-sequencing data have been

1293   uploaded to Gene Expression Omnibus (GEO) and will be made publicly available upon

1294   manuscript publication.

1295

1296 **KEY RESOURCES TABLE**

1297

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Rabbit polyclonal anti-H3K9me1	Abcam	Cat#ab176880; RRID#AB_2751009
Rabbit polyclonal anti-H3K9me3	Abcam	Cat#ab8898; RRID#AB_306848
Rabbit polyclonal anti-Histone H3	Abcam	Cat#ab46765; RRID#AB_880439
Rabbit monoclonal anti-alpha Tubulin	Cell Signaling Technologies	Cat#cst2125; RRID#AB_2619646
IRDye® 800CW Goat anti-Rabbit	LI-COR Biosciences	Cat#925-32211; RRID#AB_2651127
Chemicals, Peptides, and Recombinant Proteins		
UNC0642 G9a/GLP Inhibitor	R&D Systems	Cat#5132/10
DNA Degradase Plus	Zymo Research	Cat#E2020
Lipofectamine RNAiMax	Thermo Fisher	Cat#13778030
L-Lysine-2HCl	Thermo Fisher Scientific	Cat#PI89987
L-Arginine-HCl 13C6 15N4	Thermo Fisher	Cat#PI89990
Dimethyl Sulfoxide (DMSO)	Sant Cruz Biotechnology	Cat#sc-358801
Sequencing Grade Modified Trypsin	Promega	Cat#V5113
Propionic Anhydride	Sigma-Aldrich	Cat#240311
Phenyl Isocyanate	Sigma-Aldrich	Cat#78750-25ML
Leupeptin Hemisulfate	Thermo Fisher	Cat#AAJ61188MC
Aprotinin	Gold Biotechnology	Cat#A-655-100
Phenylmethylsulfonyl Fluoride (PMSF)	Gold Biotechnology	Cat#P-470-50
Trichostatin A	Thermo Fisher	Cat#501012404
Sodium Butyrate	Sigma-Aldrich	Cat#303410-100G
Nicotinamide	Sigma-Aldrich	Cat#N3376-500G
Urea	Acros Organics	Cat#197460050
Dithiothreitol (DTT)	Gold Biotechnology	Cat#DTT100
Iodoacetamide	Sigma-Aldrich	Cat#I1149-5G

Ammonium Bicarbonate	Sigma-Aldrich	Cat#09830-500G
Micrococcal Nuclease (MNase)	Worthington Biochemical	Cat#LS004798
Proteinase K	Thermo Fisher	Cat#EO0491
TRIzol™	Thermo Fisher	Cat#15596018
TRI Reagent	Sigma-Aldrich	Cat#T9424-25ML
3M Empore C18 47mm extraction disks	Empore	Cat#2215
Atlantis dC18 NanoEase Column (3 μM, 100 μM, 150 mm)	Waters	Cat#186002209
XBridge BEH Amide Column (3.5 μM, 2.1 mm, 150mm)	Waters	Cat#186004860
Superscript III Reverse Transcriptase	Thermo Fisher	Cat#18080044
RPMI-1640	Thermo Fisher	Cat#11875119
RPMI-1640 for SILAC	Thermo Fisher	Cat#88365
RPMI-1640, no methionine	Thermo Fisher	Cat#A1451701
Fetal Bovine Serum (FBS)	Thermo Fisher	Cat#16000044
Dialyzed Fetal Bovine Serum (DFBS)	Thermo Fisher	Cat#88212
Critical Commercial Assays		
PerfeCTa SYBR® Green SuperMix	Quanta Bio	Cat#95054-02K
SYBR Green PCR Master Mix	Thermo Fisher	Cat#4309155
REVERT Total Protein Stain	LI-COR Biosciences	Cat#926-11011
RevertAid First Strand cDNA Synthesis Kit	Thermo Fisher	Cat#K1621
Wizard® Genomic DNA Purification Kit	Promega	Cat#A1120
Dynabeads™ M-280 Sheep Anti Rabbit IgG	Thermo Fisher	Cat#11203D
QIAquick PCR Purification Kit	QIAGEN	Cat#28104
TruSeq DNA Nano	Illumina	Cat#20015964
Axygen™ Light Cycler 480 Compatible PCR Plates	Thermo Fisher	Cat#14-223-212
Deposited Data		
H3K9me1 and H3K9me3 ChIP-sequencing Data	This paper	GEO Accession Number: Available upon manuscript publication
RNA-sequencing Data	This paper	GEO Accession Number: Available upon manuscript publication

Experimental Models: Cell Lines		
H. sapiens: HCT116	ATCC	CCL-247
H. sapiens: HEK-293T	ATCC	CRL-3216
H. sapiens: HEK-293	ATCC	CRL-1573
H. sapiens: HepG2	ATCC	HB-8065
H. sapiens: MCF7	ATCC	HTB-22
H. sapiens: Panc1	ATCC	CRL-1469
M. musculus: Hepa-1c1c7	ATCC	CRL-2026
Experimental Models: Organisms/Strains		
M. musculus: C57BL/6J (9-weeks of age)	The Jackson Laboratory	JAX: 000664
M. musculus: C57BL/6J (2- and 18-months of age)	The National Institute on Aging Rodent Colony	N/A
Oligonucleotides		
5-Methylcytosine and 5-Hydroxymethylcytosine DNA	Zymo Research	Cat#D505
Stealth RNAi Sense Sequence: *H. sapiens* MAT2A: 5′AUCAAGGACAGCAUCACUGAUUUGG-3′	Thermo Fisher	Kera et al., 2013
Stealth RNAi Sense Sequence: Control: 5′-AGGAGGAUAUUACCAUGGACAAGAC-3′	Thermo Fisher	Kera et al., 2013
SUV39H1 RNAi: *H. sapiens*	Novus Biologicals	H00006839-R01-20nmol
SUV39H2 RNAi: *H. sapiens*	Novus Biologicals	H00079723-R01-20nmol
SETDB1 RNAi: *H. sapiens*	Novus Biologicals	H00009869-R01-20nmol
G9a/EHMT2 RNAi: *H. sapiens*	Novus Biologicals	H00010919-R02-20nmol
GLP/EHMT1 RNAi: *H sapiens*	Novus Biologicals	H00079813-R01-20nmol
RT-qPCR Primers	See Table S2	See Table S2
Software and Algorithms		
Bowtie2-2.1.0	Langmead and Salzberg, 2012	http://bowtie-bio.sourceforge.net/bowtie2/index.shtml

Samtools 1.5	Li et al., 2009	http://samtools.sourceforge.net/
MACS2-2.1.1.20160309	Zhang et al., 2008	http://liulab.dfci.harvard.edu/MACS/
Bedtools v2.25.0	Quinlan and Hall, 2010	https://bedtools.readthedocs.io/en/latest/
HOMER v4.9	Heinz et al., 2010	http://homer.ucsd.edu/homer/
GelAnalyzer 2010a	Lazar Software	http://gelanalyzer.com/index.html
MAVEN	Melamud et al., 2010; Clasquin et al., 2012	http://genomics-pubs.princeton.edu/mzroll/index.php
MAXQUANT	Cox et al., 2014	http://www.biochem.mpg.de/5111795/maxquant
Trimmomatic v0.39	Bolger et al., 2014	http://www.usadellab.org/cms/?page=trimmomatic
RSEM v1.2.4	Li et al., 2011	https://deweylab.github.io/RSEM/
GSEA 2	Tamayo et al., 2005 Mootha et al., 2003	http://software.broadinstitute.org/gsea/index.jsp
Skyline	MacLean et al., 2010	https://skyline.ms/project/home/software/Skyline/begin.view
Other		
Control *M. musculus* Diet	Envigo	TD.01084
Methionine Deficient *M. musculus* Diet	Envigo	TD.140119

1298

1299

Supplemental File Titles and Legends

Document S1, Related to Figures 3 and S3. Primary Antibody and REVERT Total Protein Stain Western Blot Membranes.

(*A*) Cytoplasm anti-H3K9me1 (Figure 3E). (*B*) REVERT Total Protein Stain for cytoplasm anti-H3K9me1 (Figure 3E). (*C*) Cytoplasm anti-H3 (Figure 3E). (*D*) REVERT Total Protein Stain for cytoplasm anti-H3 (Figure 3E). (*E*) Cytoplasm anti-αTubulin (Figure 3E). (*F*) REVERT Total Protein Stain for cytoplasm anti-αTubulin (Figure 3E). (*G*) Nucleoplasm anti-αTubulin (Figure 3E). (*H*) REVERT Total Protein Stain for nucleoplasm anti-αTubulin (Figure 3E). (*I*) Purified histone anti-H3K9me1 (Figure S3F). (*J*) REVERT Total Protein Stain for purified histone anti-H3K9me1 (Figure S3F).

Table S1, Related to Figures 1-3, 6-7, S1-S3, and S5. LC-MS/MS % Abundance Histone PTM Data.

Table S2, Related to Figures 2, 5, 7, S3, and S5. RT-qPCR Primers.

Table S3, Related to Figure 3. SILAC Cytoplasmic Proteins.

Figure 1. Methionine Restriction Stimulates Dynamic Histone PTM Response

Figure 2. Decreased SAM Availability Drives Robust Histone Methylation Response

Figure 3. H3K9me1 Abundance under SAM Depletion is Driven by *de novo* Methylation

Figure 4. H3K9me1 is Redistributed Over Repetitive and Transposable Elements During SAM Depletion

A
Met-R H3K9me3 Δ % Peak Number

B
Met-R H3K9me1 Δ % Peak Number

C H3K9me3
log₂(Met-R/0hr) ChIP Reads
7.66%
92.34%
Lost H3K9me3 Loci after Met-R (10⁴)

D H3K9me1
log₂(Met-R/0hr) ChIP Reads
66.44%
33.56%
Lost H3K9me3 Loci after Met-R (10⁴)

E
Δ % Annotation (H3K9me1 Enriched/Depleted)

F

Chr4 - Simple Repeat
3,752 bp
Control K9me3 [0 - 15]
Control K9me1 [0 - 15]
Met-R K9me3 [0 - 15]
Met-R K9me1 [0 - 15]

Figure 5. *De novo* H3K9 Mono-methylation Preserves Heterochromatin Stability

Figure 6. Adaptive H3K9 Mechanisms Support Epigenetic Persistence Upon Metabolic Recovery

Figure 7. Adaptive and Persistent Epigenetic Responses to SAM Depletion are Robust *in vivo*, Independent of Age

Figure S1. Related to Figure 1.

Figure S2. Related to Figure 2

Figure S3. Related to Figure 3

Figure S4. Related to Figure 6

C

GO_NUCLEIC_ACID_BINDING_TRANSCRIPTION_FACTOR_ACTIVITY_signal
GO_REGULATORY_REGION_NUCLEIC_ACID_BINDING_signal
GO_DOUBLE_STRANDED_DNA_BINDING_signal
GO_SEQUENCE_SPECIFIC_DNA_BINDING_signal
GO_CORE_PROMOTER_PROXIMAL_REGION_DNA_BINDING_signal

D

A. GO_NUCLEIC_ACID_BINDING_TRANSCRIPTION_FACTOR_ACTIVITY_signal
B. GO_REGULATORY_REGION_NUCLEIC_ACID_BINDING_signal
C. GO_DOUBLE_STRANDED_DNA_BINDING_signal
D. GO_SEQUENCE_SPECIFIC_DNA_BINDING_signal
E. GO_CORE_PROMOTER_PROXIMAL_REGION_DNA_BINDING_signal

Figure S5. Related to Figure 7