Darwinian properties and their trade-offs in autocatalytic RNA networks

Sandeep Ameta[1,‡], Simon Arsène[1,‡], Sophie Foulon[1], Baptiste Saudemont[1], Bryce E. Clifton[2], Andrew D. Griffiths[1,], Philippe Nghe[1,*]*

Affiliations

[1]Laboratoire de Biochimie, CNRS UMR8231, Chimie Biologie Innovation, ESPCI Paris, 10 Rue Vauquelin, 75005, Paris, France

[2]School of Chemistry and Biochemistry, Georgia Institute of Technology, Atlanta, Georgia 30332-0400

‡Sandeep Ameta and Simon Arsène contributed equally to the work.

Correspondence to: andrew.griffiths@espci.fr, philippe.nghe@espci.fr

Abstract

Evolution *via* template-based replication[1-4] was probably preceded by a more rudimentary form of evolution based on networks of autocatalytic reactions[5-8]. However, reaction networks possessing the Darwinian properties of variation (in composition, the catalysts present and their relative amounts), differential reproduction (accumulation of products), and heredity (persistence of composition), have so far not been identified. Here we show that networks of catalytic RNAs can possess certain properties of Darwinian systems, that these properties are controlled by network topology, and characterize important trade-offs between them. By combining barcoded sequencing with droplet microfluidics, we screened ~20,000 reactions corresponding to more than 1,800 distinct networks of ribozymes that catalyse their own formation from RNA fragments. We found that more highly connected networks tend to reproduce more quickly (accumulate more ribozymes) and be more robust to perturbations, indicating a trade-off between variation and reproduction. Variations are strongest when adding upstream ribozymes with novel reaction specificities (innovations) which target weakly connected networks. In turn, innovations increase connectivity, thus buffer against further perturbations, highlighting a second trade-off between robustness and variation. Despite this, phases of compositional robustness can alternate with sudden variations of the composition across trajectories of network growth by accretion of novel species. Finally, heredity is found to be limited by the activity of self-assembled, non-covalent ribozymes. Our findings show that connectivity in reaction networks not only determines the probability of autocatalytic sets in chemistries[8], but also their potential for evolution. They are directly relevant to scenarios where early evolution is driven by environmental heterogeneity[9,10], as it depends both on robustness (persistence being necessary for selection to act) and susceptibility to changes (in order to explore novel states). More broadly, they provide guiding rules for chemistries capable of Darwinian evolution and constrain scenarios of the origins of life.

Main

In the prebiotic world, the spontaneous appearance of an RNA polymerase ribozyme (a replicase) with sufficient processivity to allow self-replication and enough fidelity to avoid an error catastrophe[5,6] seems unlikely, given that known replicases are long (>165 nt) and structurally complex[1-4]. However, theoretical studies suggest that earlier modes of evolution could have been supported by autocatalytic sets, where reproduction results from networks of more rudimentary catalysts[11-13]. Consistently, recent experiments indicate that RNA polymerase ribozymes can assemble from catalytic networks of RNA oligomers[14], suggesting that replicases may have emerged as components of such networks. Nevertheless, sustaining evolution in reaction networks is by no means trivial, as the Darwinian properties of variation, heredity and selection are mediated by chemical compositions (the proportion of different chemical species) rather than the copying of a sequence. Furthermore, the possibility to evolve chemistries may be constained by trade-offs between these properties. For instance, robustness to environmental perturbations and persistence of compositions are necessary for selection to act, but must be balanced with variation to explore novel states. So far, none of these Darwinian properties, nor their interplay, have been experimentally studied at a large scale in a prebiotically relevant system.

We studied an experimental model of autocatalytic RNAs derived from the group I intron of the *Azoarcus* bacterium[15]. Fragments (denoted WXY and Z) assemble into non-covalent complexes that catalyze the formation of more efficient covalent ribozymes[16,17] (denoted WXYZ, Fig. 1a), which in turn catalyze with higher efficiency the formation of further covalent ribozymes, allowing the formation of autocatalytic networks[16]. Three nucleotide long sequences, called internal guide (IGS) and target (tag), located at extremities of the WXY fragments[16] (Fig. 1a, top left), determine catalytic specificity by base-pairing between a ribozyme IGS and a fragment tag. Combinations of such fragments yield networks of diverse connectivity (Fig 1a, bottom center)[18]. These networks can be represented in a coarse-grained manner by directed graphs (Fig. 1a, top right): a node representing both non-covalent and covalent ribozyme species with the same IGS and tag, and a directed edge points from an upstream ribozyme species to the downstream ribozyme species whose formation it catalyses. We developed a method to generate and measure a high diversity of such RNA reaction networks, using droplet microfluidics coupled to barcoded Next-Generation Sequencing (Fig. 1b-c, Extended Data Fig. 1). It consists of first producing a library of 5 pL droplets containing 24 different combinations of the 16 WXY fragments (Extended Data Table 1), together with

hairpin RNA reporters that do not react but enable later identification of the mixture of WXY fragments by sequencing. The 24 initial combinations were selected among a random sample by minimizing their mutual overlap while covering as evenly as possible network sizes from 1 to 16 species after combinatorial droplet fusion (see below and Methods). Random sets comprising 1 to 5 droplets of this initial library were then electrocoalesced[19] with a 50 pL droplet containing the reaction buffer and Z fragments. After incubation at 48°C for 1 hour, the composition of each droplet was analyzed by droplet-level RNA sequencing adapted from single-cell transcriptomics[20]. Hydrogel beads carrying cDNA primers with a barcode specific to each bead (Extended Data Fig. 2) were encapsulated one-by-one in droplets then fused with the RNA containing droplets, followed by primer release and cDNA synthesis (Fig. 1c, Extended Data Fig. 1, and Methods). Barcoded cDNAs were recovered and sequenced. All the reads from a same droplet carry the same barcode (Extended Data Fig. 3a, b).

The initial combinations of fragments (encoded by hairpin RNA reporters, Extended Data Fig. 3, 4), and the final fraction of ribozymes produced during incubation were determined in 20,038 droplets, comprising 1,837 unique networks with on average 11 replicates each (Fig. 1d, Extended Data Fig. 5a), indicating a mean precision of ~6% in species fractions (Extended Data Fig. 5b). Repeatability was tested with a full experimental replicate (r=0.84 between species fractions), and droplet cross-talk was quantified to be ~13% (Extended Data Fig. 5c-e).

We observed highly heterogeneous patterns of covalent ribozyme accumulation, demonstrating the existence of interdependences between reaction network components (Fig. 1d). Indeed, if every ribozyme species were reacting independently, the relative rank between any pair of ribozyme species would be conserved across all networks. On the contrary, we observed that for 63% of ribozyme pairs, their relative ranking differed in at least 10% of the networks (Extended Data Fig. 6a), extending previous findings of non-conserved ranking in a small set of such networks[21]. This diversity in species fractions in networks was well predicted by a kinetic model without fitting parameters (Methods) by adding the first order effective catalytic rates of non-covalent and covalent ribozymes measured independently for individual IGS-tag pairs (r=0.83, p<10^{-3}, Fig. 1 e-f, Extended Data Fig. 6 b-d).

Comparing networks generated from distinct substrate sets also revealed differences in growth, quantified as the concentration of covalent ribozyme accumulated during the reaction relatively to hairpin reporters of known concentration (Fig. 2a): yield varied 10-fold across networks comprising between 2 and 15 species (see Extended Data Table 2), and up to 6-fold within networks of the same size but with different topologies (Fig. 2a).

We then quantified variations in response to the addition of WXY fragments allowing the
formation of a novel ribozyme. This corresponds to comparing networks G and G' differing by
a single node a (networks connected by a curved line in Fig. 1d). Denoting V the set of nodes
common to G and G', the total perturbation is defined as $p_{G \to G'} = \sum_{v \in V} |y'_v - y_v|$ where y_v and
y'_v are the respective fractions of species v in G and G', both normalized within V (Fig. 2b). By
definition, the maximum is $p_{G \to G'} = 2$ and is reached for a full switch in species composition,
for example going from $(y_u, y_v) = (0, 1)$ to $(y'_u, y'_v) = (1, 0)$ in $V = \{u, v\}$.
Averaging for each network the effect of all perturbations (green line, Fig. 2c,) revealed large
differences from marginally to strongly perturbable networks (mean $\bar{p}_{G \to G'}$ from 0.1 to 1).
Furthermore, we observed a trade-off between growth and variation (Fig. 2d), implying that
high yield networks tend to be poorly perturbable.
We aimed to interpret the diversity of observed responses based on network connectivity. The
kinetic model does not provide such a direct interpretation as it consists of a mixture between
two regimes, where either only non-covalent or only covalent ribozymes are active. For non-
covalent ribozymes only, species fractions should be predicted by in-degree centrality, a
network-theoretic measure which accounts only for catalysis by directly upstream ribozymes[22].
In contrast, for covalent ribozymes only, species fractions should be predicted by eigenvector
centrality, which accounts for longer catalytic chains[22]. Despite these differences, the in-degree
centrality was found to be a good approximation of the eigenvector centrality (for our dataset,
R=0.83, p-value<10[-5], Extended Data Fig. 7a), as is already known in general[23], except for
strongly selfish ribozymes which fraction is underestimated (13% of the dataset, Extended Data
Fig. 7b).
The in-degree centrality approximation allowed an analytical derivation of the perturbations
(Supplementary File 1),

$$149 \qquad p_{G \to G'} = 2n \frac{1 - n \frac{m}{\sigma_G/e}}{\sigma_G/e + n} \ (1),$$

with 4 control parameters (Fig. 3a): n is the *perturbation breadth*, the number of targets of the
new node a introduced in the network; m is the *catalytic novelty*, the number of catalysts
already present in G with the same IGS as a (novelty being higher for lower m); σ_G is the
background strength, the sum of all edge weights in the network, and; e is the catalytic strength
e of edges outgoing from a as determined by its IGS.
The influence of parameters $\sigma_G, e, m,$ and n on $p_{G \to G'}$ predicted by the in-degree approximation
(Equation 1, continuous curves in Fig. 3b) was similar to those predicted by eigenvector

centrality (Fig. 3b, dots), and, more importantly, were well verified experimentally (Fig. 3b,

compare left and right panels). First, perturbation strength was impacted much more strongly

by catalytic novelty m (Fig. 3b, bottom) than by perturbation breadth n for the values tested

here (n = 1 or 2, Fig. 3b, top). Second, Equation (1) poses the condition $\sigma_G/e \leq n$ for

significant variations, consistent with high values $p_{G \to G'}$ being observed only when $\sigma_G/e < 2$

(given that $n \leq 2$, Fig. 3b, top). Overall, the highest perturbations $p_{G \to G'}$ required a catalytic

innovation (m = 0, i.e. a new IGS) to be combined with a low normalized background strength

($\sigma_G/e \leq 1$). Perturbations were indeed up to 10 times higher in such cases, compared to cases

where either $m > 0$ or $\sigma_G/e > 2$ (Fig. 3b, bottom). Interestingly, the model indicated that

perturbations should peak at intermediate values of σ_G/e when $m > 0$ (purple line, Fig 3b, right),

which was consistently observed in the data but not significant (purple line, Fig 3b, left). The

predictions of Equation 1 were also verified for other values of the parameters (Extended Data

Fig. 8) and at the level of single nodes (Extended Data Fig. 9). In addition, the attenuation of

perturbations with large σ_G/e combined with the weak but significant correlation between

growth and σ_G (R=0.5, p<10^{-5}, Fig. 3c) explains part of the trade-off between growth and

perturbability reported in Fig. 2d.

To test the interplay between robustness and variation in scenarios where novel species would

appear either spontaneously[13] or due to changes in substrates provided from the prebiotic

milieu[9,10], we analyzed cumulative perturbations across trajectories of network growth, starting

from networks with three nodes, randomly adding one node at a time (Fig. 4a). We have seen

that strong perturbations require catalytic innovations (m = 0). In such case, Equation 1

reduces to $p_{G \to G'} = 2/(1 + \sigma_G/ne)$. The ratio σ_G/ne expresses a second trade-off, between

the robustness induced by the background strength of the network σ_G, and the variation induced

by the novel catalyst as characterized by n and e.

As for a large perturbation to occur, ne must be comparable to the σ_G of the perturbed network,

σ_G would roughly double after a strong perturbation, enhancing robustness to further variation.

Consequently, strong variations should be followed by small ones, and result in inflexions

(change in curvature) in cumulative perturbation trajectories. We quantified the corresponding

inflexions by their sharpness (third derivative at inflexion, Extended Data Fig. 10a), and

categorized them as strong when in the top 25% of sharpness (Fig. 4b). Comparing the

distributions of sharpness for all inflexion points showed that they are, as predicted,

significantly sharper for catalytic innovations (Fig. 4c). Consistently, the number of strong

inflexions correlates with the number of catalytic innovations per trajectory (Fig. 4d).

Although introducing a node causing a strong perturbation buffers the resulting network against
further perturbations, subsequent variations along trajectories are still possible, as exemplified
in Fig. 4a, b. By computationally analysing trajectories for an extended repertoire of specific
interactions, we found that sustained variation requires perturbing species of increasing
targeting breadth n (Fig. 4e) and increased waiting time between variation events (until the
diversity of IGS/tag pairs saturates) Fig. 4f. The former allows the ever larger σ_G values to be
overcome, while the latter corresponds to the build-up of weakly connected nodes that can
become targets. Interestingly, the trade-off between σ_G and ne translates at the level of
trajectories: chemistries with a high density of catalytic interactions lead to many trajectories
with a few inflexions, while sparse chemistries lead to fewer trajectories but with many
inflexions (Fig. 4g).

We have shown experimentally that a diversity network behaviour can be generated from a
small set of interactions, and that these networks can possess Darwinian properties. What is
missing, therefore, for such a system to evolve? First, we found that network topology imposes
trade-offs between growth and variation, and between variation and robustness. In particular,
the density of interactions in a chemistry should neither too sparse, nor too dense to potentially
allow evolution. Second, heredity, which in the context of autocatalytic networks depends on
the persistence of the chemical composition over time is weakened due to catalysis by self-
assembled, non-covalent, complexes. This leads to relaxation of chemical compositions toward
states determined by the substrates rather than by the covalent catalysts transmitted from pre-
existing networks (Extended Data Fig. 11). Reducing catalysis by self- non-covalent complexes
may sufficiently enhance heritability for the system to evolve, if combined with selection and
variation. On the other hand, self-assembled catalysts could facilitate the emergence of novel
catalysts from variation in the substrates provided from environment.

214
215
216

References

1. Horning, D. P. & Joyce, G. F. Amplification of RNA by an RNA polymerase ribozyme. *Proc. Natl. Acad. Sci. U.S.A.* **113**, 9786-9791 (2016).
2. Johnston, W. K., Unrau, P. J., Lawrence, M. S., Glasner, M. E. & Bartel, D. P. RNA-catalyzed RNA polymerization: accurate and general RNA-templated primer extension. *Science* **292**, 1319-1325 (2001).
3. Kim, D. E. & Joyce, G. F. Cross-catalytic replication of an RNA ligase ribozyme. *Chem. Biol.* **11**, 1505-1512 (2004).
4. Wochner, A., Attwater, J., Coulson, A. & Holliger, P. Ribozyme-catalyzed transcription of an active ribozyme. *Science* **332**, 209-212 (2011).
5. Eigen, M. Selforganization of matter and the evolution of biological macromolecules. *Naturwissenschaften* **58**, 465-523 (1971).
6. Eigen, M. & Schuster, P. The hypercycle. A principle of natural self-organization. Part A: Emergence of the hypercycle. *Naturwissenschaften* **64**, 541-565 (1977).
7. Hordijk, W., Steel, M. & Kauffman, S. The structure of autocatalytic sets: evolvability, enablement, and emergence. *Acta. Biotheor.* **60**, 379-392 (2012).
8. Kauffman, S. A. Autocatalytic sets of proteins. *J. Theor. Biol.* **119**, 1-24 (1986).
9. Abbott, S. S., Harrison, T. M., Schmitt, A. K. & Mojzsis, S. J. A search for thermal excursions from ancient extraterrestrial impacts using Hadean zircon Ti-U-Th-Pb depth profiles. *Proc. Natl. Acad. Sci. U.S.A.* **109**, 13486-13492 (2012).
10. Stueken, E. E. *et al.* Did life originate from a global chemical reactor? *Geobiology* **11**, 101-126 (2013).
11. Jain, S. & Krishna, S. A model for the emergence of cooperation, interdependence, and structure in evolving networks. *Proc. Natl. Acad. Sci. U.S.A.* **98**, 543-547 (2001).
12. Nghe, P. *et al.* Prebiotic network evolution: six key parameters. *Mol. Biosyst.* **11**, 3206-3217 (2015).
13. Vasas, V., Fernando, C., Santos, M., Kauffman, S. & Szathmáry, E. Evolution before genes. *Biol. Direct* **7**, 1; discussion 1 (2012).
14. Mutschler, H., Wochner, A. & Holliger, P. Freeze-thaw cycles as drivers of complex ribozyme assembly. *Nat. Chem.* **7**, 502-508 (2015).
15. Reinhold-Hurek, B. & Shub, D. A. Self-splicing introns in tRNA genes of widely divergent bacteria. *Nature* **357**, 173-176 (1992).
16. Draper, W. E., Hayden, E. J. & Lehman, N. Mechanisms of covalent self-assembly of the Azoarcus ribozyme from four fragment oligonucleotides. *Nucleic Acids Res.* **36**, 520-531 (2008).
17. Hayden, E. J., von Kiedrowski, G. & Lehman, N. Systems chemistry on ribozyme self-construction: evidence for anabolic autocatalysis in a recombination network. *Angew. Chem. Int. Ed. Engl.* **47**, 8424-8428 (2008).
18. Vaidya, N. *et al.* Spontaneous network formation among cooperative RNA replicators. *Nature* **491**, 72-77 (2012).
19. Chabert, M., Dorfman, K. D. & Viovy, J. L. Droplet fusion by alternating current (AC) field electrocoalescence in microchannels. *Electrophoresis* **26**, 3706-3715 (2005).
20. Klein, A. M. *et al.* Droplet barcoding for single-cell transcriptomics applied to embryonic stem cells. *Cell* **161**, 1187-1201 (2015).
21. Yeates, J. A., Hilbe, C., Zwick, M., Nowak, M. A. & Lehman, N. Dynamics of prebiotic RNA reproduction illuminated by chemical game theory. *Proc. Natl. Acad. Sci. U.S.A.* **113**, 5030-5035 (2016).
22. Yeates, J. A. M., Nghe, P. & Lehman, N. Topological and thermodynamic factors that influence the evolution of small networks of catalytic RNA species. *RNA* **23**, 1088-1096 (2017).
23. Valente, T. W., Coronges, K., Lakon, C. & Costenbader, E. How Correlated Are Network Centrality Measures? *Connect (Tor. Ont.)* **28**, 16-26 (2008).

268 24. Krzywinski, M. *et al.* Circos: an information aesthetic for comparative genomics. *Genome Res.*
269   **19**, 1639-1645 (2009).
270

Figures

Fig. 1. Experimental set-up and RNA network compositional landscape. a, Top left: RNA fragments, named WYX and Z, can be assembled into full length ribozymes WXYZ (both non-covalent and covalent catalysts). WYX fragments each comprise 3 nucleotide long IGS and tag sequences, respectively denoted g**M**g and c**N**u, where **M**, **N** can be varied, resulting in 16 different combinations[18]. Bottom center panel: The base-pairing interactions between **M** and **N** determine the specificity of full length catalysts for their substrate fragments. In this example, each of the 3 different MN ribozyme species catalyzes the formation of another species from fragments provided as substrates, forming an autocatalytic cycle. Dotted and bold arrows show catalysis carried out by non-covalent and covalent ribozymes, respectively. Note that for clarity, only Watson-Crick IGS-tag

interactions are shown here. Any other non-Watson pair will have a weaker but non-zero activity (Extended Data Fig. 6c), therefore all networks possess some degree of autocatalysis. Top right panel: a simplified network representation shows catalytic species as nodes with their MN identity, and catalytic interactions as directed edges (corresponding to the dotted and bold black arrows of the bottom center panel). **b**, Random sets of ~3 droplets (small black disks), each containing different mixtures of WXY fragments, are fused by electrocoalescence in a microfluidic device with a droplet containing Z fragments and the reaction buffer (larger gray disks). This combinatorial process leads to a large diversity of catalytic networks, each contained in a different droplet (Methods). Substrate fragments are initially co-diluted with non-reactive hairpin RNA reporters that later allow the initial composition to be retrieved by sequencing. **c**, Droplet-level barcoded sequencing: each RNA-containing droplet is analyzed after incubation by fusing a sample of it to another droplet containing a hydrogel bead carrying DNA barcodes. These barcodes, specific to each bead, serve as primers for the reverse transcription of the RNAs, including the ribozymes formed during incubation, and the hairpin RNA reporters to identify initial condition. **d**, Compositional landscape of the 1837 unique measured networks[24]: Each ray is made of 16 boxes corresponding to the 16 possible MN species. A blank box means that the MN ribozyme is absent in this network, as determined from hairpin sequencing. Otherwise the box color indicates the measured fraction of the MN species. Grey arcs connect networks that differ from each other by a single node (addition or removal). **e**, Left: schematic of an *Azoarcus* network. Right: top, experimentally determined node fractions for the same network (error bars are ±1 s.d. over n=425 droplets); bottom, theoretical results with the kinetic model. **f**, Fraction obtained with the kinetic model versus measured fraction for every node that is a member of a 4 node-network. Bins with less than 10 points are discarded. Dark grey dots are mean values, quartile boxplots have 5^{th} - 95^{th} percentiles whiskers with flier points. The dotted grey line is the identity line. P-value < 0.001.

Fig. 2. Network growth and perturbation. a, Network total yield (concentration of WXYZ catalysts in μM) distribution of networks with four nodes and at least 10 replicates (for data on all sizes see Extended Data Table 2). Networks are ranked (x-axis) based on the mean across network replicates of the total WXYZ concentration (y-axis). The error bars show the Standard Error of the Mean. **b**, Example showing the addition of node a (GG, orange) to network G resulting in network G'. Top: schematic. Bottom: experimentally determined node fractions for the same network (error bars are ±1 s.d. over n=109 droplets (G) and n=15 droplets (G') **c**, Perturbation distribution of networks with four nodes. Networks are ranked (x-axis) based on the mean perturbation (y-axis). Each position on the x-axis corresponds to a certain network G, for which the perturbation has been computed for all possible single node additions present in the dataset, leading to several y-axis values corresponding to different G' networks. The green line is the average perturbation taken over all G' networks for each network G. **d**, Perturbation $p_{G \rightarrow G'}$ for networks with 3, 4 and 5 nodes plotted against network yield in μM for perturbations involving the addition of a novel catalysts (with G/C as IGS) with at least one target. The number of strongly perturbable ($p_{G \rightarrow G'}$>0.8, above the green dotted line) and high yield networks (>0.08 μM, right hand side of red dotted line) is very low compared to the null hypothesis of independence of perturbability and yield (one-sided Fisher exact test, N=162, odds ratio < 0.1, 95% confidence interval 0.01-0.66, p-value=1.6.10^{-3}).

Fig. 3. Network perturbation analysis. a, Schematic of three situations where the parameters n, the number of targets of added node a, and m, the number of nodes sharing the same IGS as a, have different values. **b**, The perturbation $p_{G \to G'}$ is plotted against σ_G/e for different values of the parameters m and n for both the experimental data (left, mean and standard deviation per bin, bins with less than 3 points are discarded) and results predicted by in-degree centrality using the analytical expression (right, solid line) and with fractions predicted by eigenvector centrality (right, solid circles) for the networks with a number of nodes between 3 and 6 before addition. Results for other parameter values are shown in Extended Data Fig. 8. **c**, Network yield (μM) is plotted against σ_G for network with four nodes (see Methods for further details on how network yield is determined).

a Network growth trajectory by sequential addition of nodes

328
329
Fig. 4. Perturbation dynamics across network growth trajectories. a, Example of a network growth trajectory
where at each step, a new node is added. Nodes resulting in strong perturbations (see panel b) are in green, and
correspond to the IGS/tag pairs C-G and G-C. **b**, Examples of measured cumulative perturbation over trajectories,
plotted against the number of node additions, by number of strong inflexions points (colored). The latter are
determined as the top 25% in sharpness (absolute value of the third derivative, Extended Data Fig. 10a) measured
over all trajectories. The asterisk (*) denotes the perturbation trajectory of the example shown in panel a. **c**,
Distribution of sharpness for inflexion points associated with a catalytic innovation (orange, n=4,462) or not (grey,
n=934). Catalytic innovations are defined as the introduction of strong IGS/tag interactions (CG or GC, Extended
Data Fig. 6 b, c). that were not present until node addition. The dotted line is the mean of each distribution and
the significance of the difference between the two distributions is reported (Mann–Whitney U test, p < 0.001). **d**,
Distribution of the number of inflexion points within the 75th percentile sharpness per trajectory, depending on the
number catalytic innovations per trajectory. **e-g**, Computational study of network growth trajectories for
chemistries with varying number of IGS/tag pairs and with varying degrees of catalytic density points are

determined, as before, based on their sharpness (within the 75th percentile) here computed along a random sample
of 1000 trajectories growing from 2 to 100 nodes. Catalytic densities are varied by random removal among the
pool of all possible specific IGS/tag interactions. **e**, Targeting breath (number of targets) of catalysts causing strong
perturbations, as the function of the inflexion rank. **f**, Waiting time (number of species additions) between two
strong inflexions, as a function of the inflexion rank (e.g.: rank 5 is the fifth inflexion observed along a growth
trajectory) **g**, Proportions of trajectories with a given number of strong inflexion points plotted against catalytic
density for a chemistry comprising up to 24 different IGS/tag pairs.
350

Methods

General material & methods. All experiments used DNAse/RNase free water (Thermo Fisher Scientific Product No.: 10977035). Chemicals were purchased from Sigma-Aldrich unless specified otherwise. 4-(2-Hydroxyethyl)-1-piperazinepropanesulfonic acid (EPPS) was purchased from Alfa Aesar (Product no.: J60511, CAS no.: 16052-06-5). Denaturing polyacrylamide gels (12%) were prepared using gel stock solution (Roth, Product no.: A431.1) and run in 1X Tris-Borate EDTA buffer. DNA/RNA oligonucleotides (Supplementary File 2) were obtained from IDT DNA technologies.

Transcription of RNAs. RNAs were *in vitro* transcribed as described in[25], using DNA templates produced as described in the Supplementary File 1. In the template for the WXY fragments the IGS is duplicated at position 25 to allow identification by sequencing. The RNAs were extracted with Phenol-Chloroform-Isoamyl alcohol, treated with DNase I (Thermo Fisher Scientific, Product No.: EN0521) and purified on polyacrylamide gels. RNA concentrations were measured using Qubit® RNA HS Assay Kit (Thermo Scientific, Product No.: Q32852).

Microfluidic devices and set-up. Droplet microfluidic experiments used HFE 7500 fluorinated oil (3M™ Novec™, Product No.:98-0212-2928-5) and a fluorosurfactant (RAN Biotechnologies, Product No.: 008-FluoroSurfactant). Devices were designed with AutoCAD software (Supplementary File 4), printed at Selba SA. PDMS (polydimethylsiloxane) devices were fabricated by soft-lithography as described in[26,27]. For electrocoalescence, indium tin oxide coated glass slides were used (Delta Technologies, Product No.: CG-90IN-S215), with channels containing 3 M NaCl liquid electrodes[28]. After plasma bonding, channels were made hydrophobic with 2% 1H,1H,2H,2H-perfluorodecyltrichlorosilane (ABCR, Product No.: AB111155) in HFE 7500. Droplet-based microfluidics experiments were performed using an inverted microscope (Nikon Eclipse *Ti*) as described earlier[27]. Flows were controlled by either syringe pumps (Harvard Apparatus Inc.) or by air-pressure control pumps (MFCS™-EZ, Fluigent SA).

Droplet microfluidic experiments. The complete experimental system consists of the steps **A** to **E** (Extended Data Fig. 1): (**A**) *Initial emulsions production:* Using all possible 16 WXY RNA fragments ($_{gMg}WXY_{cNu}$)[18], 24 unique different WXY fragments combinations were prepared in separate tubes (Extended Data Table 1). To determine those combinations, tube contents were computationally drawn randomly among the 16 species. The total number of fragment types to distribute over all tubes (with possible redundancies) was set to 50, which was computationally

found to maximize the number of distinct networks formed after random droplet fusions (10,000

iterations). Once this was fixed, network diversity was further maximized based on spectral

graph criteria, by selecting the fragment assignment (among 100,000 random realizations)

maximizing the Shannon entropy of the first eigenvalue and spectral gap distributions, across

networks resulting again from the simulation of 10,000 random droplet fusions.

Experimentally, for each combination, 1 µM of respective WXY fragment(s) was heated at

48°C and cooled down to room temperature over 10 min (0.1°C/sec) and mixed with hairpin

RNA reporter at a final concentration of 0.03 µM. All 24 mixes were encapsulated individually

in 5 pL droplets using flow focusing[29] on a dropmaker device (Supplementary File 4) using

flow rates of 100-150 µL/h for the aqueous (RNA) and 150-170 µL/h for the oil (HFE 7500

with 2% surfactant) phases, each emulsion being collected for 5 min. All 24 emulsions were

collected together in a 1.5 mL collection tube (with a PDMS cap) containing oil with 2%

surfactant and mixed thoroughly. A 1.6 µM solution of Z RNA fragment in 1X *Azoarcus*

reaction buffer (30 mM EPPS pH 7.4, 20 mM $MgCl_2$), was folded in the same manner as WXY

RNA, and encapsulated in 50 pL droplets using a 50 pL drop-making device (Supplementary

File 4), at flow rates of 350 µL/h for the aqueous (RNA) and 250 µL/h for the oil (HFE 7500

with 2% surfactant) phases, droplets being collected in a 0.5 mL collection tube (with a PDMS

cap). All flows were driven by syringe pumps. (**B**) *Electrocoalescence of initial emulsions*: The

5 pL droplets (containing WXY fragments) and 50 pL droplets (containing Z fragment and

reaction buffer) were re-injected into separate channels of a droplet fusion device

(Supplementary File 4). Both 5 pL and 50 pL droplets were spaced using HFE 7500 oil

containing 2% surfactant and brought together into the same channel, where they were

electrocoalesced[19,28] using an AC electrical field (Agilent 33522A waveform generator, sine

function, 4 kHz, 400 mV, 50 Ω) amplified 10^3 times (TREK high-voltage amplifier). Flows

were regulated such that, on average, between one and five 5 pL droplets coalesce with one 50

pL droplet (Supplementary File 5). This resulted in ~1/10th dilution of WXY RNA concentration

in the fused droplets (the Z fragment is in stoichiometric excess). All flows were driven by a

pressure control system (MFCS™-EZ, Fluigent SA), and the desired fusion frequency achieved

by carefully adjusting and coupling the flows with pressure values of 380-400, 350-380, 470-

520 mbar for 50 pL droplet emulsion, 5 pL droplet emulsion and spacer oil channel,

respectively. Electrocoalescence and collection (in a 0.2 mL tube with a PDMS cap) was

performed for ~ 3.5 h and the emulsions were stored on ice during the entire process. To monitor

the fusion frequency during the collection, fusion events were counted from videos recorded

every 20 min, showing very stable fusion frequency, closely comparable to what was obtained

by sequencing hairpin RNA reporters (Extended Data Fig. 4). (**C**) *Incubation and splitting of droplets:* After collection, fused droplets were incubated at 48°C in a thermo-block for 1 h to allow the accumulation of ribozymes. In order to dilute the MgCl$_2$ during the reverse transcription (RT) step, these droplets (~60-65 pL) were re-injected (spaced with HFE 7500 oil containing 2% surfactant) and split by a T-junction device[30] (Supplementary File 4) into smaller droplets (~5 pL) prior to the RT step. Flow rates were maintained using pressures of 800-1000 mbar and 150-250 mbar, respectively for the oil separator channel and droplet inlet. (**D**) *Droplet barcoding and RT in droplet*: To sequence the RNA in each split droplet we developed a strategy combining droplet barcoding and RT in droplets, inspired by single-cell transcriptomics methods described earlier[20,31]. Here barcoded hydrogel beads are individually encapsulated in droplets with all the necessary reagents for RT and fused with RNA network containing droplets. The details of each step are as follows: (*a) Barcoded hydrogel bead synthesis*: Hydrogel beads were produced by co-encapsulating 10% (w/w) polyethylene glycol diacrylate (PEG-DA) 6000 (Sigma, Product No.: 701963), 1% (w/w) PEG-DA-700 (Sigma, Product No.: 455008), 400 µM of acrydite-modified dsDNA with a 4 base sticky-end (Oligo 10, top, carrying a 5'-acrydite modification, and Oligo 11, bottom, Supplementary File 2), 10 µM of FITC-Na (Fluorescein isothiocynate), 1% (v/v) photo-initiator (2-hydroxy-2-methylpropiophenone, Sigma, Product No.: 405655) in buffer (75 mM Trizma HCl pH 7.4, 50 mM NaCl) using flow focusing[32] on a dropmaker device (Supplementary File 4). 9 pL droplets were produced using flows of 150 µL/h for aqueous solution and 500 µL/h for oil (HFE 7500 with 2% surfactant). Droplets were polymerized by UV irradiation (UV omnicure AC475-365, Lumen Dynamics, ~360 mW exposure, λ_{nm} = 365 nm) through a PTFE tubing and collected in 5 mL tube (tubing internal diameter = 0.3 mm, distance from lamp 3 cm, tubing length = 12 cm; approximate exposure time = 47 s). Prior to UV irradiation, the complete experiment set-up was maintained in the dark to avoid any spontaneous polymerization. The collected droplets were washed once with 4 mL hexane to break the emulsion and then the beads washed 3 times with 4 mL binding-wash buffer (20 mM Trizma-HCl pH 7.4, 50 mM NaCl, 0.1% of Tween 20) by centrifuging at 3000 g for 2 min. Beads were then filtered using Steriflip 20 µm Nylon filters (Millipore, Product No.: SCNY00020) and stored in binding-wash buffer supplemented with 1 mM EDTA. In contrast to earlier methods[20,31], barcodes were built on the beads using a split-and-pool ligation strategy[33] (Extended Data Fig. 2). For this, 250 µL (~10 million) of pelleted beads were washed 3 times with 4 mL of binding-wash buffer by centrifuging at 3000g for 2 min, and were subjected to a first ligation step. Ligation was performed on a 2 mL reaction scale with 1X T7 DNA Ligase buffer (New England Biolabs), 4 µM dsDNA adaptor with two

different 4 base sticky-ends, the first being complementary to the 4 base sticky-end on the

beads, and containing a BclI restriction site and the Read 2 illumina adaptor sequence (Oligo

12, top and Oligo 13, bottom, Supplementary File 2), 30 U/µL of T7 DNA Ligase (New England

Biolabs, Product No.: M0318L), at room temperature for 25 min with agitation at 600 rpm.

Beads were then washed 5 times with 4 mL of binding-wash buffer and pelleted after the last

wash. For the next ligation step, a master-mix (1.6 mL) was prepared to containing 1X T7 DNA

Ligase buffer, 30 U/µL of T7 DNA Ligase and divided into the wells of a 96-well deep well

plate (16 µL per well), pre-filled with 4 µL of 20 µM 1st barcode index (Index A)

(Supplementary File 2), with a different index in each well. These indexes had two different 4

base sticky-ends, the first being complementary to the free sticky-end of oligos ligated in

previous step (dsDNA, Oligo 12 and 13). The plate was sealed and incubated at 25°C with 600

rpm for 25 min. Then the ligase was heat inactivated at 65°C for 10 min and the content of all

the wells was pooled after adding 200 µL of binding-wash buffer containing 1 mM EDTA.

After pooling, the beads were washed 7 times with 4 mL of binding-wash buffer. The complete

process of split-and-pool was repeated to ligate the 2nd (Index B) and 3rd (Index C) barcode

indexes (Supplementary File 2) generating a total diversity of 8.8 x 10^5 (96^3) barcodes. After

the final pooling step, a partially double-stranded sequence was ligated to all the beads using

the same ligation protocol (Oligo 14, top and Oligo 15, bottom, Supplementary File 2). This

sequence comprises a sticky-end complementary to the free sticky-end on the 3rd index, and a

single-stranded 33 base long 3'-overhang at the other end (Extended Data Fig. 2b). Out of these

33 nucleotides, the first 8 are random nucleotides used as unique molecular identifiers (UMIs)[34]

and remaining 25 nucleotides, which function as a primer for RT, are complementary to the 3'

end of the WXYZ ribozyme and hairpin RNA reporter (both contain the same primer binding

region). The final barcoded hydrogel bead library was washed 5 times with 4 mL of binding-

wash buffer supplemented with 1 mM EDTA and re-suspended in the same buffer. (b)

Encapsulation and fusion: The barcoded beads (50 µL) were washed 5 times with 500 µL of

binding-wash buffer by centrifuging at 11866 g for 1 min. Washed beads were then mixed with

2.5 µL of 10 mM dNTPs, 10 µL of 5X SuperScript III reaction buffer (Invitrogen), 2.5 µL of

100 mM dithiothreitol (DTT), 0.4% of Tween20 (10%), 100 units of SUPERase·In^{TM} inhibitor

(Thermo Fisher Scientific, Product No.: AM2694) and incubated at 37°C for 30 min to ensure

diffusion of all reactants within the beads. After this, the beads were centrifuged as above and

excess of liquid over the beads was removed. Then, at 4°C, 500 units of reverse transcriptase

(SuperScript III, Thermo Fisher Scientific, Product No.: 18080044) and 25 units of BclI

restriction enzyme (New England Biolabs, Product No.: R0160L) were added to the beads and

mixed thoroughly. These hydrogel beads together with the other reagents were encapsulated

individually in ~50 pL droplets (Supplementary File 6), fused with RNA droplets on the same

microfluidic device (Supplementary File 4). The injection of close-packed deformable beads

resulting in >99% of droplets containing a single barcoded hydrogel bead. The 5 pL RNA

containing droplets and the 50 pL bead containing droplets were fused at a 1:10 ratio to ensure

that, in the majority of cases, no more than one of the formers is fused with one of the latter.

For this, flows were pressure controlled using, respectively, 550, 225, 650, 450 mbar for

hydrogel beads, 5 pL droplets, oil to control droplet spacing (HFE 7500 with 2% surfactant)

and oil for beads encapsulation (HFE 7500 with 2% surfactant). Electrocoalescence was

performed as described above. Fused droplets were collected in 0.2 mL collector tube (with

PDMS cap) containing oil with 2% surfactant. (c) *cDNA synthesis and extraction*: The fused

droplets were then incubated at 60°C in a thermo-block for 1 h to release the DNA barcodes

using the BclI restriction enzyme and perform cDNA synthesis in droplets. The emulsion was

broken post incubation by adding 2 volumes of 1*H*, 1*H*, 2*H*, 2*H*-Perfluoro-1-octanol and 100

µL water. The pooled barcoded cDNAs are then isopropanol precipitated and re-suspended in

40 µL of water. **(E)** _Sequencing sample preparation_: In order to block extension of the non-

elongated primers, 20 µL of cDNA was mixed with 1X TdT reaction buffer (New England

Biolabs), 0.25 mM $CoCl_2$, 0.4 mM ddCTP (Roche CustomBiotech, Product No.: 12158183103)

and 0.4 U/µL of Terminal Transferase (TdT, New England Biolabs, Product No.: M0315L).

The reaction was carried out in 50 µL volume and incubated for 30 min at 37°C before heat

inactivation at 70°C for 10 min. The TdT treated cDNAs were purified using magnetic beads

(AMPure XP, Beckman Coulter, Product No.: A63881) following the manufacturer's protocol.

Sequencing adaptors were then appended using two sequential PCR steps (Extended Data Fig.

3a). The first PCR was performed in multiples of 20 µL volume where 2 µL of TdT treated

reverse transcription reaction was mixed with 0.5 µM of Oligo 16 (also containing a 6

nucleotide stretch used as sample barcode to multiplex different samples for the sequencing

run) and 0.5 µM of Oligo 17 (Supplementary File 2) as forward and reverse primers in 1x PCR

buffer (Thermo Scientific), 0.2 mM dNTPs, 0.01 U/µL of polymerase (Thermo Scientific

Phusion Hot Start II, Product No.: F459) using the following protocol: initial denaturation

98°C/30sec, then 18 cycles of denaturing 98°C/10 s, annealing and extension 72°C/1 min, and

a final extension of 72°C/5 min. PCR products were purified using AMPure XP magnetic beads

and eluted in 34 µL of water. The second PCR was performed as above, but in 40 µL volume

with 4 µL as PCR I purified product as template, using Oligo 18 and Oligo 19 (Supplementary

File 2) as forward and reverse primers, and using the following protocol: initial denaturation

98°C/30sec, then 10 cycles of denaturing 98°C/10 s, annealing 56°C/30 s, extension 72°C/30

s, and a final extension of 72°C/5 min. The final sample was purified using AMPure XP

magnetic beads, resuspended in 34 µL of water, analysed on TapeStation (Agilent 2200

TapeStation, using high sensitivity D1000 ScreenTape®, Product No.: 5067-5584) and

quantified by Qubit® dsDNA HS Assay Kit (Thermo Scientific, Product No.: Q32854).

Libraries were sequenced using the Illumina NextSeq 550 system in 2*150 High Output mode

at the Genotyping and Sequencing Core Facility, ICM Paris (iGenSeq, Institut du cerveau et de

la moelle épinière).

Sequencing data processing. Custom software was written to process sequencing data and

further data analysis steps. The libraries were sequenced in paired-end mode with read 1 and

read 2 of 180 bp and 120 bp, respectively. Read 1 was used to determine the sample barcode

(used for multiplexing samples for sequencing) and the RNA molecule identity (either an

Azoarcus ribozyme with specific IGS-tag pair or a hairpin RNA reporter). Read 2 was used to

determine the UMIs[34] and the droplet barcode. The structure of read 1 and 2 is summarized in

Extended Data Fig. 3b. *UMI normalization*: Four meta fields were associated for each pair of

reads: sample barcode, droplet barcode, UMI and RNA molecule identity (details in

Supplementary File 1). Reads missing any of these were discarded (~30%). The reads with the

same meta information were merged and the read count was added as an extra meta field. This

allows to filter out noise due to sequencing errors, given that read counts resulting from such

errors are significantly lower (threshold indicated on Extended Data Fig. 3e). The filtering

thresholds for the minimum number of reads per UMI were determined separately for the

Azoarcus ribozymes and hairpin RNA reporters based on the shape of the distribution of number

of reads per UMI (Extended Data Fig. 3e-f). *Final data processing*: Filtered UMI-normalized

reads were then clustered by droplet barcode to count the number of UMIs per type of RNA

molecule per droplet barcode (Extended Data Fig. 3f). For each droplet barcode, only hairpin

RNA reporters where UMIs made up ≥7.5% of the total of hairpin UMIs were retained. This

threshold value was set to optimally match the fusion distribution measured by video (Extended

Data Fig. 3g). Within each droplet, ribozyme sequences that did not corresponding to a correct

hairpin reporter were discarded. Then, only the droplet barcodes with more than 10 UMIs

associated with hairpin RNA reporters and 20 with ribozymes were retained and the fraction of

each node in the network was computed. As before, these thresholds were chosen to closely

match the observed distribution of drop fusion determined from videos acquired during the

experiments (Extended Data Fig. 3h, i). Processed network compositions and mean node

fractions are publicly available for all unique networks (Supplementary File 3). To determine
network yield per droplet, first we computed a droplet-specific UMI to concentration
conversion rate by considering total hairpin reporter UMIs as well as number of different
hairpin reporters identified. This was then used to convert total number of ribozyme UMIs into
a yield value.

Controls for droplet-level sequencing biases. Four 5 pL emulsions (A, B, C and D) were
prepared, each with: a distinct pair of hairpin RNA reporters at a final concentration of 5 nM, a
set of 4 WXYZ ribozymes with the same IGS but in different proportions (total concentrations
summing to 100 nM) and 1.6 µM of Z fragment. The four emulsions were mixed and then
singly fused to droplets containing hydrogel beads as described above to perform droplet level
sequencing. The data was processed as above using the same thresholding strategy. We
quantified the bias as the proportion of droplets containing pure (A, B, C or D) and compared
to mixed populations (AB, AC, AD, BC, BD, CD, ABC, ABD, BCD and ABCD, Extended
Data Fig. 5d).

Experimental measurement of catalysis by non-covalent and covalent ribozymes. Self-
assembly of *Azoarcus* ribozymes with different IGS/tag pairs were re-measured in 20 mM
MgCl$_2$, using the strategy developed by von Kiedrowski[35], as described[21]. For this, the initial
rate of formation of WXYZ from WXY and Z fragments was measured as a function of the
concentration of doped covalent WXYZ ribozymes (with same IGS and tag as the WXY
fragment). The concentration of RNA fragments were identical to the droplet experiments (0.1
µM of WXY fragment and 1.6 µM of Z fragment) with additional but minor amounts (0.001
µM) of WXY fragment, radiolabeled using T4 Polynucleotide Kinase (New England Biolabs)
and γ-32P ATP (Perkin-Elmer). In accordance to previous measurements[21], the initial rate of
formation of WXYZ (v_0) was found to depend linearly on the concentration of the doped
WXYZ. This linear relationship can be described with the following linear equation $v_0 = \alpha x +$
β (Extended Data Fig. 6 b,c) where, x is the concentration of the covalent ribozyme, α is the
slope for an IGS/tag pair, which quantifies the WXYZ synthesis by covalent ribozymes, and β
is the y-intercept, which quantifies the WXYZ synthesis by non-covalent ribozymes (trans-
catalysis by non-covalent complex of WXY and Z fragments)[18]. Wobble pairs (GU and UG)
and one of the mismatched IGS/tag pairs (AG) were also measured and found to have negligible
values for α and β.

Kinetic model. In a network, covalent catalysts from species j, whose concentration is x_j,
contributes to the rate of formation of covalent catalysts from species i with the term $\alpha_{ij} x_j$

while non-covalent catalysts contribute with the constant term β_{ij}. α_{ij} and β_{ij} terms were

measured experimentally and independently from the network experiments, for single IGS-tag

pairs (Extended Data Fig. 6b-c). The production rate of each species is $\dot{x}_i = \sum_j \alpha_{ij} \cdot x_j + \beta_{ij}$,

resulting in a linear system of ordinary differential equations for each network depending on

the IGS-tag combinations provided from the RNA fragments.

Additional references

25.   Arsene, S., Ameta, S., Lehman, N., Griffiths, A. D. & Nghe, P. Coupled catabolism and anabolism
594   in autocatalytic RNA sets. *Nucleic Acids Res*. **46**, 9660-9666 (2018).
595   26.   Duffy, D. C., McDonald, J. C., Schueller, O. J. & Whitesides, G. M. Rapid Prototyping of
596   Microfluidic Systems in Poly(dimethylsiloxane). *Anal. Chem*. **70**, 4974-4984 (1998).
597   27.   Mazutis, L. et al. Single-cell analysis and sorting using droplet-based microfluidics. *Nat. Protoc*.
598   **8**, 870-891 (2013).
599   28.   Sciambi, A. & Abate, A. R. Generating electric fields in PDMS microfluidic devices with salt
600   water electrodes. *Lab Chip* **14**, 2605-2609 (2014).
601   29.   Anna, S. L., Bontoux, N. & Stone, H. A. Formation of dispersions using "flow focusing" in
602   microchannels. *Appl. Phys. Lett*. **82**, 364-366 (2003).
603   30.   Link, D. R., Anna, S. L., Weitz, D. A. & Stone, H. A. Geometrically mediated breakup of drops in
604   microfluidic devices. *Phys. Rev. Lett*. **92**, 054503 (2004).
605   31.   Macosko, E. Z. et al. Highly Parallel Genome-wide Expression Profiling of Individual Cells Using
606   Nanoliter Droplets. *Cell* **161**, 1202-1214 (2015).
607   32.   Anna, S. L., Bontoux, N. & Stone, H. A. Formation of dispersions using "flow focusing" in
608   microchannels. *Appl. Phys. Lett*. **82**, 364-366 (2003).
609   33.   Grosselin, K. et al. High-throughput single-cell ChIP-seq identifies heterogeneity of chromatin
610   states in breast cancer. *Nat. Genet*. **51**, 1060-1066 (2019).
611   34.   Kivioja, T. et al. Counting absolute numbers of molecules using unique molecular identifiers.
612   *Nat. Methods* **9**, 72-74 (2011).
613   35.   von Kiedrowski, G. A Self-Replicating Hexadeoxynucleotide. *Angew. Chem. Int. Ed. Engl*. **25**,
614   932 (1986).

Extended Data

618
Extended Data Fig. 1. Schematic of the complete experimental set-up. The different steps involved in the
experiment. See Methods for details. On a first microfluidic device, 24 different initial emulsions, comprising 5
pL droplets, were prepared, each containing a unique combination of WXY fragments (Extended Data Table 1),
and a unique molecular hairpin RNA reporter. These emulsions were collected in a single tube (1.5 mL) and mixed
thoroughly. These 5 pL droplets were then re-injected into a second microfluidic device, where they were fused
by electrocoalescence with 50 pL droplets, produced on a separate chip, containing Z fragment and the reaction
buffer. The pairing of 5 pL and 50 pL droplets was controlled such that one to five 5 pL droplets fuse with one 50
pL droplet. After fusion, the resulting emulsion, comprising 65 pL droplets, was incubated at 48°C for 1 h, and
then split into 5 pL droplets on a third microfluidic device to reduce the volume. These droplets were re-injected
into a fourth microfluidic device, where they were fused with 50 pL droplets, produced on the same chip,
containing single hydrogel beads carrying barcoded cDNA primers, and reverse transcription reaction reagents.
The ratio of 5 pL to 50 pL droplets was kept low (1:10) such that only one 5 pL RNA droplet was fused with one
50 pL droplet. The resulting emulsion was incubated at 60°C for 1 h to allow release of barcoded primers from the
beads and cDNA synthesis. Then, the emulsion was broken, the barcoded cDNA was amplified by PCR to append
sequencing adaptors, and sequenced.

634
Extended Data Fig. 2. Barcoded hydrogel beads. a, Synthesis of barcoded hydrogel beads using a split and pool
ligation strategy (details in Methods). On each bead, the clonal population of barcodes is built by successive
ligation of indexes in one of the wells of 3 successive 96-well plates, in which each well contains a different index.
Between each ligation step, beads are pooled before being distributed randomly in the 96 wells of the next plate.
After the ligation of three indexes (A, B and C), a common oligo containing a linker (double stranded), UMIs (8
nt, single stranded), and the RNA binding region for reverse transcription (25 nt, single stranded) is ligated. **b**,
Final structure of a barcode: linker, restriction site (RE), Read 2 Illumina sequence (Rd2), 3 barcode indexes (A,
B and C) separated by linker regions, UMI and cDNA primer. The sequences of the ligated 4 base sticky-ends in
the linker regions are indicated. **c**, Capillary electrophoresis (TapeStation, Agilent) data of the barcoded primers
released from beads by restriction digestion. The length at the main peak (~70%) corresponds to fully ligated

barcodes with three indexes. Note that the actual length of full barcodes is 152 bp compared to the length indicated by TapeStation (210 bp) due to barcodes being partially single-stranded. See Supplementary File 1 for details. **d,** Distribution of percentage of the most abundant barcode sequence per bead is plotted for n=71 beads obtained by sequencing of barcodes released from single beads in microplate wells. See Supplementary File 1 for details.

652 **Extended Data Fig. 3. Sequencing and analysis. a**, Schematic of PCR steps to append sequencing adaptors (see
653 Methods). In the first step, barcoded cDNAs are amplified with a gene-specific forward primer, which also contains
654 a barcode for multiplexing different samples for sequencing, and a common read 2 reverse primer. The PCR
655 products from first PCR are then subjected to a second PCR to add adaptors compatible with Illumina sequencing
656 platforms. **b-c**, Sequencing read structures **b**, Generic paired-end read structure. **c**, Detailed structure of the reads

for *Azoarcus* ribozymes and for hairpin RNA reporters. **d**, Structure of molecular hairpin RNA reporters, which share the same 5' and 3' regions as WXYZ to allow amplification. These hairpins have a variable loop region of 4 nucleotide serving as barcode. **e**, Distribution of number of UMIs as a function of the number of reads per UMI for reads associated with *Azoarcus* ribozymes or hairpin RNA reporters. The threshold, indicated by the gray line, is determined by the intersection of the the "noise" distribution with very low number of reads per UMI and the "signal" distribution with higher mean number of reads per UMI. Any UMIs with less reads than this threshold were discarded. Note that x-axis scale for the plots is different because of the more efficient amplification of hairpin reporters during PCR due to their smaller size. **f**, Distribution of number of UMIs (ribozyme and hairpin) per droplet barcode. **g-i**, Choosing the thresholds. The distance to expectation is the Euclidian distance between the distribution of the number of hairpin RNA reporters per droplet barcode once the three thresholds (see below) have been applied, and the distribution of droplets at the fusion step measured by video acquisition. The three thresholds are: **g**, the minimum percentage of total hairpin RNA reporter UMIs used to declare the corresponding hairpin reporter to be part of the coding set for a given droplet barcode (chosen value = 7.5%), **h**, the minimum number of hairpin RNA reporter UMIs for a droplet barcode to be included in the final dataset (chosen value = 10), and **i**, the minimum number *Azoarcus* ribozyme UMIs for a droplet barcode to be included in the final dataset (chosen value = 20).

Extended Data Fig. 4. Identification of the network structure at the droplet level using hairpin RNA
reporters. a, Schematic illustrating identification of a network structure using hairpin RNA reporters. The hairpin
RNAs where the UMI counts make up ≥7.5% of the total number of UMI counts for hairpin RNAs are considered
as part of the coding set. This can be used to derive network structure in addition to node identification using an
internal mutation in WXYZ ribozyme (Extended Data Fig. 3c, Supplementary File 1). The number of UMIs for
each node of the identified network structure gives the node fraction. **b**, In orange, measured distribution of the
number of hairpin RNA reporters per droplet barcode in the final dataset. In dark grey, distribution of number of
5 pL droplets (containing initial WXY emulsions) fused with a 50 pL droplet (containing reaction buffer and Z)
measured by video acquisition (Methods). Pearson's correlation coefficient is reported (p<0.001, n=13). **c**, In
green, distribution of number of nodes per network per droplet barcode in the final dataset. In dark grey, expected
distribution from simulating the library fusion step with the same number of droplets as in the data and with the
fusion frequencies measured by video acquisition (Methods). Pearson's correlation coefficient is reported
(p<0.001, n=17). **d**, Histogram of distribution of the proportion of hairpin RNA reporter-coded ribozyme UMIs,
defined as the ratio between the total numbers of UMIs associated with nodes identified by the hairpin reporters
and the total number of UMIs for *Azoarcus* ribozyme identified by the internal mutation (Supplementary File 1)
for the same droplet barcode. A value of 1 is reached when all ribozyme UMIs predicted by the set of hairpin
reporters are identified. **e**, Correlation between the measured and expected network proportions. Network
proportion is calculated as the ratio between the number of replicates of a given network and the total number of
replicates for all networks. Expected network proportion is obtained by simulating the library fusion step with
50,000 droplets mimicking the experimental set-up using the fusion frequencies measured by video acquisition.
About 1/3 of the networks in the data do not have any replicate in the simulation and are excluded from the analysis.
Bins with less than 10 points were discarded. The box extends from the lower to upper quartile values of the data,
with a dot at the mean. The whiskers extend from the 5th percentile to the 95th. Flier points are those past the end

of the whiskers. The dotted orange line is the identity line. Pearson's correlation coefficient is reported (p<0.001,
n=2160).

700
701
Extended Data Fig. 5. Measurement precision. a, Proportion of unique networks having replicates. 56.0% of
the networks in the final dataset have at least duplicates and 18.3% of them have at least 10 replicates. **b**,
Distribution of node fraction standard deviation where standard deviation is measured from the networks with at
least 5 replicates. The dotted grey line is the mean of the distribution of standard deviation. The result indicates a
mean precision of ~6% in species fractions. **c**, The complete experiment (Methods, Extended Data Fig. 1) was
replicated and node fractions in both replicates were measured. Node fractions measured in one experimental
replicate are plotted against the other. Data is restricted to the common set of networks present in the two replicates.
Data points are binned in 20 linearly spaced bins according to their x-axis component. Bins with less than 10 points
are discarded. The box extends from the lower to upper quartile values of the data, with a dot at the mean. The
whiskers extend from the 5[th] percentile to the 95[th]. Flier points are those past the end of the whiskers. The dotted
orange line is the identity line. Pearson's correlation coefficient is reported (p<0.001, n=2047). These technical
replicates demonstrated high repeatability (r=0.84 between species fractions) **d-e**, Control for cross-talk and
quantification bias during droplet barcoding and sequencing. Barcoded sequencing was performed on a mix of
four emulsions (A, B, C and D) each containing a mixture of ribozymes in known proportions as well as a pair of

specific hairpin RNA reporters for identification (Methods). **d**, Measured proportions of each population of
droplets (orange) and expectations (green). The result indicates 87% of sequenced droplet contained a single set
of ribozymes which demonstrate a low cross-talk between droplets. **e**, Measured against expected ribozymes
concentrations (normalized) aggregating data from the four emulsions show high correlation (r = 0.91) and rule
out biases of the droplet level sequencing. Each dot corresponds to a ribozyme in one of the emulsions (4 ribozymes
in each of the 4 emulsions). Error bars are standard deviations. Grey dotted line is the identity line. Pearson
correlation coefficient p<0.001 in all cases.

Extended Data Figure 6. Fitting of edge weights in *Azoarcus* networks across various models of node centrality and comparison of the best model with experimental measurements. a, For each pair of nodes, fractions are extracted in all networks that contain the two nodes. The color code indicates the proportion of cases where one node (row) has higher fraction compared to the other node (column). **b-c**, Experimental measurements to extract catalytic parameters used to describe the *Azoarcus* network system[21,35] (Methods). **b**, Plot showing

reaction kinetics for all four Watson-Crick IGS/tag interactions. Initial rates (v_0) of WXYZ formation from WXY
and Z RNA fragments were measured by doping the reaction with covalent WXYZ ribozyme with the same IGS
and tag as the WXY fragment, as reported by Yeates *et al.*[21]. **c**, Table showing the α and β parameters derived
from fitting a linear relationship (top) between the initial rate of formation of WXYZ ribozymes (v_0) as a function
of doped covalent WXYZ ribozyme concentration (x)[21,35]. Here α quantifies the synthesis of WXYZ ribozymes
catalyzed by covalent ribozymes and, is derived from the slope of this relationship (plot in **b**). Whereas, β quantifies
the synthesis of WXYZ ribozymes by non-covalent ribozymes and is derived from the intercept of this relationship
(plot in **b**). **d**, Fraction obtained with the kinetic model versus measured fraction for networks with same number
of nodes. Bins with less than 10 points are discarded. Dark grey dots are averages; quartile boxplots have 5th - 95th
percentiles whiskers with flier points. The dotted orange line is the identity line.
740

741
Extended Data Fig. 7. Network composition predictions using eigenvector centrality and in-degree
centrality a, Covalent ribozyme fraction predicted by eigenvector centrality versus in-degree centrality for all
networks of the experimental dataset (N=1837). In grey are species participating to networks containing outlier
ribozymes. The latter are ribozyme species which fraction is underestimated by more than 20% by the in-degree
centrality compared to the eigenvector centrality. **b**, Networks containing outliers tend to have selfish nodes
(bottom) compared to others (top). Selfishness is quantified here as the in-degree of nodes with self-loops and no
out-going edges.
749

750
751 **Extended Data Fig. 8. Analytical predictions of Equation 1 with all possible values of the parameters n and**
752 **m.** Network perturbation $p_{G \to G'}$ upon addition of a new node is plotted against normalized background strength
753 σ_G/e for all possible values of the catalytic novelty m (panels) and of perturbation breadth n (color coded).

Extended Data Fig. 9. Node perturbation $p_{G \to G'}^{v}$ upon addition of a new node a to network G, comparing experimental data with analytical predictions. a, $p_{G \to G'}^{v}$ plotted against normalized background strength σ_G/e for different values of catalytic novelty m and perturbation breadth n in the case where v is a target for a and for both the experimental data (left) and the analytical results (right). For details of how analytical predictions were derived see Supplementary File 1. Data points were distributed along the x-axis in 20 log-spaced bins between $10^{-0.5}$ and $10^{2.5}$. Bins with less than 3 points were discarded and the mean and standard deviation for each bin are plotted. Networks with less than 3 nodes were discarded for the analysis. **b**, Same as **a** but the case where v is not target for a.

Extended Data Fig. 10. Determination of strong inflexion points and influence of the number of different IGS/tag interactions on the number of inflexion points per trajectory. a, Cumulative perturbation for four representative trajectories from the experimental data are plotted (first row) as well as the first derivative (second row), the second derivative (third row) and the net third derivative (fourth row). The derivative is the difference between two successive points divided by the width of the interval (here 1). Inflexion points are first detected with the second derivative when it switches from positive to negative and the sharpness of these is measured with the net third derivative. Derivatives were computed as the difference of values between n and n+1 nodes. Strong inflexion points are defined as being in the top 25% in sharpness and are indicated with dark grey rectangles. An extra-step with no perturbation is added at the beginning of the trajectory to detect starting strong inflexion points. **b**, Proportion of trajectories with a given number of inflexions plotted against catalytic density for different numbers of IGS/tag combinations. Note that panels corresponding to 6 and 7 inflexions show no trajectory with inflexions when the number of IGS/tag pairs equals 4. Hence analysis of trends holds only when the number of IGS/tag pairs exceeds the number of inflexions, as mentioned in the main text. These results were obtained by simulating 20,000 trajectories as a function of catalytic density values (defined as the proportion of the non-null specific interaction between an IGS and a tag over the complete set of possible interactions) for various number of IGS/tag. See Fig. 4g for the details.

782
783
Extended Data Figure 11. Relaxation of chemical compositions to states due to non-covalent catalysts. a,
Schematic of two serial transfer experiments with different initial conditions and two generations, G_0 and G_1,
incubated for an hour at 48°C as in the droplet experiment. Top: control experiment (unperturbed network), where
only the fragment for the AC, AU, UA and UG ribozymes are provided at each round, with concentrations 0.1 μM
for WXY fragments and 1.6 μM for Z. Bottom: same as top, but the network is perturbed at G_0 by addition of CA
fragments at the same concentration. **b,** Relative fraction of covalent ribozymes AC, AU, UA and UG produced
during each generation of the serial transfer. Although introduction of fragments at G_0 has a significant impact on
the final covalent ribozyme composition, these differences are not preserved at G_1 where the composition relaxes
toward the one of the unperturbed network.
793
794

795 **Extended Data Table 1. Different combination of WXY fragments in initial 5 pL emulsions.**

Combination no.	$_gM_g$WXY$_cN_u$ fragments (MN)[#]	Hairpin RNA reporter variable region[¶]
1	UA	UUUU
2	CA	ACUG
3	GA	AAUU
4	AG	GAUA
5	CU	GUGG
6	GC	UAUG
7	UU	CCCA
8	CU, UG	AAGG
9	AC	GGAA
10	CA, GA, CG, UA, AU	AUAG
11	GG, GC	UAAC
12	AU	UGUA
13	CA, UC, GA, CC	UACA
14	AC, AG	CGUG
15	UA, GU, GG, CG, UG, AG, UU, UC	GCCG
16	GG	GAAU
17	AU, CA, AA, AG	GGGU
18	GC	CGCC
19	GU	UUGG
20	CC, GU, CA	GUCC
21	GG, GU, CC	UAGC
22	UU, CU	AAAA
23	CU, CA	CUAA
24	UC	AUGC

796 [#]All 16 possible $_gM_g$WXY$_cN_u$ RNA fragments were used where 'M' is the middle nucleotide of IGS and 'N' is the
797 middle nucleotide of tag[18]. [¶]4 nucleotide variable region of the hairpin RNA reporter added to barcode the
798 composition.
799

800   **Extended Data Table 2. Variation in growth for different sizes of networks.**

Number of nodes	Number of networks	Mean yield (nM)	SD yield (nM)	Minimum growth (nM)	Maximum growth (nM)	Fold difference max/min growth
2	22	37.3	19.1	10.4	95.1	9.1
3	48	49.6	29.5	19.3	151.2	7.8
4	60	56.0	26.4	25.6	146.9	5.8
5	47	84.0	37.5	26.4	162.1	6.1
6	47	113.9	43.8	52.4	233.3	4.5
7	27	142.7	38.6	91.9	222.6	2.4
8	17	181.7	67.9	92.6	373.1	4.0
9	9	280.6	45.4	184.4	328.3	1.8
10	16	303.4	60.3	203.4	405.5	2.0
11	10	371.9	86.9	236.8	481.3	2.0
12	15	371.3	95.7	240.8	654.6	2.7
13	10	367.8	70.7	260.2	452.6	1.7
14	5	419.1	36.2	381.0	459.5	1.2
15	3	394.9	40.7	359.8	439.5	1.2