TITLE

Interactions between the N- and C- termini of mechanosensitive ion channel AtMSL10 support a

three-step mechanism for activating its signaling function

AUTHORS/AFFILIATIONS

Debarati Basu[1], Jennette M. Shoots[1], Alex Harkess[2], Kira M. Veley[2] and Elizabeth S. Haswell[1]

[1]NSF Center for Engineering Mechanobiology, Dept. of Biology, Washington University in St.

Louis, St. Louis MO 63130, [2]Donald Danforth Plant Science Center, St. Louis, Missouri 63132.

CONTACT INFORMATION

ehaswell@wustl.edu

RUNNING TITLE

Interaction between the N- and C-termini of MSL10

ABSTRACT

Although a growing number of mechanosensitive ion channels are being identified in plant systems, the molecular mechanisms by which they function remain to be established. Overexpression of the mechanosensitive ion channel MSL (MscS-Like)10 fused to GFP triggers a number of developmental and cellular phenotypes including the induction of cell death, and this function is regulated by seven phosphorylation sites in its soluble N-terminus. Here, we show that these and other phenotypes required neither overexpression nor a tag and could be also induced by a point mutation in the soluble C-terminus (S640L). In all cases, these gain-of-function phenotypes were suppressed by growth at elevated temperatures. The promotion of cell death and hyperaccumulation of H_2O_2 in *35S:MSL10^{S640L}-GFP* overexpression lines was suppressed by N-terminal phosphomimetic substitutions, and the soluble N- and C-terminal domains of MSL10 physically interacted. Finally, transcriptomic analysis revealed that the complex network of genes induced by these gain-of-function *MSL10* alleles overlap with established biotic and abiotic stress-responsive pathways. We propose a three-step model by which tension-activated conformational changes in the C-terminus activate the N-terminus, leading to its dephosphorylation and the induction of adaptive responses. Taken together, this work expands our understanding of the molecular mechanisms of mechanotransduction in plants.

KEYWORDS

MscS-Like, MSL10, mechanosensitive ion channel, cell death, reactive-oxygen species

INTRODUCTION

Mechanical forces, whether endogenous or exogenous in origin, are important stimuli that help direct plant growth, development, and immune responses. External mechanical perturbations include touch, wounding, wind, vibration, gravity, osmotic stress, and pathogen invasion (Monshausen and Haswell, 2013; Jayaraman et al., 2014). Endogenous mechanical forces include turgor pressure, a fundamental cue influencing cell expansion (Hamant and Haswell, 2017; Kierzkowski, 2019). Before plants can coordinate a cellular response to these varied stimuli, they must be able to precisely sense and transduce mechanical cues.

Mechanosensitive (MS) ion channels provide a fast and efficient molecular mechanism for transducing mechanical stimuli into intracellular signals. They are found in all domains of life, and are highly diverse in terms of their structure, ion selectivity, regulation, and physiological roles (Martinac et al., 2013; Ranade et al., 2015; Basu and Haswell, 2017). However, all MS ion channels share two common attributes: opening in response to mechanical force (lateral membrane tension, forces relayed from the cytoskeleton, or a combination of both) and facilitating the passive flow of ions across membrane (Bavi et al., 2017).

In plants, there is a growing list of predicted and established MS ion channel families, which includes the MscS-Like (MSL, Basu and Haswell, 2017), Two Pore Potassium (TPK, Maathuis, 2011), Mid1-Complementing Activity (MCA, Nakagawa et al., 2007) and Reduced Hyperosmolality-induced [Ca^{2+}] Increase (OSCA, (Hou et al., 2014; Yuan et al., 2014; Murthy et al., 2018) families, as well as homologs of the animal MS ion channel Piezo (Zhang et al., 2019). Plant MS channels have been implicated in a wide range of physiological functions, including organellar osmoregulation (Veley et al., 2012), stomatal closure (Gobert et al., 2007; Yuan et al., 2014), cold tolerance (Mori et al., 2018), hyperosmolarity-evoked Ca^{2+} influx (Yuan et al., 2014), penetration of hard substrates (Nakagawa et al., 2007), pathogen-triggered immunity (Zhang et al., 2017) and viral resistance (Zhang et al., 2019). However, the detailed molecular mechanisms by which MS channels participate in these events need further investigation.

The MSL family of MS ion channels was first identified based on homology to the canonical MS ion channel from *Escherichia coli,* Mechanosensitive ion channel of Small conductance (*Ec*MscS)

(Pivetti et al., 2003; Haswell and Meyerowitz, 2006; Haswell et al., 2008). There are ten MSLs

encoded in the *Arabidopsis thaliana* genome, and they are localized to mitochondria (MSL1, Lee

et al., 2016), to chloroplasts (MSL2/3, Haswell and Meyerowitz, 2006) or to the plasma membrane

(MSL8, MSL9 and MSL10, Haswell et al., 2008; Hamilton et al., 2015). The tension-gated ion

flux of MSL1, MSL8, and MSL10 have been examined by single channel patch-clamp

electrophysiology (Maksaev and Haswell, 2012; Hamilton et al., 2015; Lee et al., 2016).

MSL2/MSL3 and MSL8 are thought to serve as osmotic safety valves in plastids and in pollen,

respectively (Veley et al., 2012; Hamilton et al., 2015). MSL1 plays a poorly understood role

maintaining redox homeostasis in mitochondria (Lee et al., 2016).

While MSL10 was the first in the family to be characterized by electrophysiology, its physiological

function is still under study. To date, no visible loss-of function phenotypes have been reported in

msl10-1 null mutants (nor in double *msl9-1 msl10-1* mutants, where MSL10's closest homolog

MSL9 is also ablated). Overexpression of *MSL10-GFP* results in growth retardation, ectopic cell

death, constitutive production of H_2O_2, and induction of genes involved in ROS accumulation,

senescence, and abiotic and biotic stress responses (Veley et al., 2014). These gain-of-function

phenotypes can be attributed to a single domain of MSL10, the soluble N-terminus; in a transient

Nicotiana benthamiana expression system, this domain can induce cell death on its own (Veley et

al., 2014). In another study, we showed that variants of MSL10-GFP harboring pore-blocking

lesions also induce cell death in this system (Maksaev et al., 2018). None of the phenotypes

associated with *MSL10-GFP* overexpression are observed in plants overexpressing a version of

MSL10-GFP harboring four phospho-mimetic substitutions in the N-terminus (*MSL10 $^{S57D, S128D,}$*

$^{S131E, T136D}$*-GFP*, or *MSL10^{4D}-GFP*), though this variant has MS ion channel activity, protein levels,

and subcellular localization indistinguishable from the wild type.

Thus, dephosphorylation of the MSL10 N-terminus activates *MSL10-GFP* to trigger cell death

when overexpressed in Arabidopsis or in *N. benthamiana*. Ion flux through the channel pore,

however, is not required. We infer that under normal conditions, MSL10 remains in its inactive

state (i.e. with a phosphorylated N-terminus), and that overexpression overwhelms the kinase that

normally maintains it in its inactive form, thereby triggering a cell death signaling cascade. It has

been shown that the opening of *Ec*MscS is accompanied by structural rearrangement of the soluble

C-terminal domain (Bass et al., 2002; Wang et al., 2008; Machiyama et al., 2009; Rowe et al.,

2014). If similar rearrangements occur in MSL10, the N-terminus could also undergo tension-

induced conformational changes that lead to activation, either directly from the channel pore or

indirectly from the C-terminal domain.

Indeed, a functional link between the N- and the C-termini of MSL10 was suggested by a recent

report by the Zhou group (Zou et al., 2015). The *rea1 (RAP2.6 expresser in shoot apex)* mutant

harbors an EMS-induced point mutation (S640L) located in the soluble C-terminus of *MSL10* that

leads to increased expression of a wound-responsive luciferase reporter gene. Mutant *rea1* plants

exhibit growth retardation and ectopic cell death, strikingly reminiscent of *MSL10-GFP*

overexpression lines. The *rea1* mutants also exhibited shorter petioles, accumulation of

anthocyanin pigments, lack of apical dominance, wound-induced hyperaccumulation of JA, and

altered expression of genes involved in JA biosynthesis and response (*LIPOXYGENASE2* or

LOX2, PLANT DEFENSIN or *PDF1.2,* and *ALLENE OXIDASE* or *AOS)*.

The *rea1* mutant thus provides the opportunity to probe MSL10 function in an endogenous context.

Its phenotypic similarity to *MSL10-GFP* overexpression lines suggested that growth retardation,

ROS accumulation, and cell death might reflect physiological functions of the MSL10 protein,

though it was also possible that these similarities might be only superficial. Here we present an in-

depth comparison of the developmental, cellular and gene expression phenotypes resulting from

MSL10-GFP overexpression, the *rea1* lesion, and genomic phospho-variants of *MSL10*. The

results from these experiments prompted us to assess genetic and physical interactions between

the soluble N- and C- termini of MSL10 and to characterize the gene expression pattern connecting

MSL10 activation to downstream phenotypic effects. Our results suggest a three-step process of

MSL10 activation that transduces the effect of increased membrane tension from soluble C-

terminus to the N-terminus of the channel, leading to phosphorylation and subsequently triggering

downstream gene expression networks.

RESULTS

Conservation of functionally significant residues in the soluble N- and C-terminal domains

of putative MSL10 orthologs

We investigated the conservation of eight residues previously identified as important for MSL10

function: seven phosphorylation sites in the soluble N-terminus (S29, S46, S48, S57, S128, S131,

and T136; and a C-terminal amino acid that is mutated in the *rea1* mutant (S640, Zou et al., 2015)

(**Figure 1A**). The predicted N-terminal domains of putative MSL10 homologs from other

flowering plants showed surprisingly low sequence conservation. However, when analysis was

restricted to the *Brassicacae* family, the 33-amino-acid region containing S29, S46, S48 and S57,

and the 15-amino-acid region containing S128, S131, and T136 showed 73% amino acid identity

with the *Brassica napus* homolog (**Figure 1B**). The sequence of the MSL10 C-terminus

surrounding S640 was highly conserved among angiosperms (90% amino acid identity with the

putative homolog from *B. napus,* 60% identity with that from *Oryza sativa*), and S640 itself was

conserved in 11 of the 15 sequences analyzed (**Figure 1C**). To summarize, phosphosites in the

soluble N-terminus known to modulate the *MSL10-GFP* overexpression phenotype were not well-

conserved, while S640, a residue in the soluble C-terminus known to affect MSL10 function, was

well-conserved among the sequences analyzed.

rea1* is a recessive gain-of-function *MSL10* allele, renamed *msl10-3G

To further characterize the *rea1* allele, we backcrossed *rea1* plants to wild type *Col-0* plants. As

shown in **Figure S1A**, the resulting F1 hybrid plants resembled the wild type parent, indicating

that the mutation responsible for the phenotypes observed in the *rea1* mutant is recessive,

consistent with previous findings (Zou et al., 2015). After self-pollination, the resulting F2

population segregated into wild-type and *rea1* phenotypes at a ratio of 3:1, as expected for a

recessive allele, and these results were confirmed by PCR genotyping (**Figure S1C**). In addition,

rea1 plants were outcrossed to *msl10-1* null mutant plants. The F1 progenies resembled *rea1* in

terms of leaf morphology but attained intermediate height after 8 weeks of growth (**Figure S1B**).

After self-pollination, resulting F2 populations similarly segregated approximately 1:2:1 with

respect to height at late stages of development (wild type like: intermediate: *rea1* phenotype);

phenotypes were confirmed by PCR genotyping (**Figure S1C**). These data show that the

phenotypic effects of the *rea1* lesion depend on the presence or absence of the WT *MSL10* allele,

suggest that they may be dosage-dependent (as it requires two copies of the *rea1* mutant allele to

generate the full *rea1* phenotype), and confirm that the phenotypes associated with the *rea1* mutant

are due to a recessive gain-of-function point mutation in the *MSL10* gene. This allele is hereafter

called *msl10-3G*.

MSL10-GFP overexpression, the *msl10-3G* allele, and the $MSL10g^{7A}$ transgene produce

similar phenotypes and gene expression patterns

Strong constitutive overexpression of a gene often results in pleiotropic phenotypes that may not

reflect its normal function (Zhang et al., 2003a; Kovalchuk et al., 2013), and the addition of tags

can alter protein localization, regulation, stability or function (Spartz et al., 2012). Zou et al. (2015)

reported that neither estradiol-induced expression of *MSL10-FLAG* nor constitutive expression of

untagged *MSL10* resulted in the phenotypes seen with constitutive expression of *MSL10-GFP*,

suggesting that large tags may perturb the function of MSL10. Furthermore, we were previously

unable to retrieve plants constitutively expressing $MSL10^{4A}–GFP$ (Veley et al., 2014), and while

we proposed that the over-expression of phospho-dead *MSL10* was lethal, it remained possible that

an unrelated defect in the transgene prevented us from isolating any transgenic plants.

To determine if the phenotypes associated with *MSL10-GFP* overexpression could be replicated

at endogenous expression levels and without a GFP tag, we generated constructs to drive

expression of untagged wild-type *MSL10 (MSL10g)*, phospho-dead *MSL10* S29A, S46A, S48A,

S57A, S128A, S131A, and T136A ($MSL10g^{7A}$) or phospho-mimetic *MSL10* S29D, S46D, S48D,

S57D, S128D, S131E, and T136D ($MSL10g^{7D}$), within the native *MSL10* genomic context. These

constructs were introduced into the *msl10-1* null mutant background and three independent

homozygous transgenic lines from each transformation that accumulated *MSL10* transcripts at

levels similar or slightly higher than those of wild-type plants (**Figure S2A**) were selected for

further analysis.

Figure 1. *MSL10-GFP* overexpression lines, *msl10-3G* mutants, and *MSL10g^{7A}* lines have similar whole-plant phenotypes. (A) Predicted topology of an AtMSL10 monomer. Teal, cytosolic N-terminus; yellow, transmembrane helices; green, cytosolic C-terminus. Black triangles denote the positions of the residues mutated in this study. The MscS domain is as defined in (Veley et al. 2014). (B) Multiple alignment of protein sequences corresponding to the N-terminus of putative AtMSL10 orthologs in the Brassicaceae family. Darker teal shading indicates higher conservation score and identical amino acids with an asterisk. (C) Multiple alignment of protein sequences corresponding to the C-terminus of putative AtMSL10 orthologs in selected flowering plants. Conservation of residues is indicated as in (B). (D) and (E) T4 homozygous lines overexpressing *MSL10-GFP* in the Col-0 background (line 15-2), T3 homozygous lines expressing *MSL10g*, or *MSL10g^{7D}* in the *msl10-1* background, and T3 segregating or T3 homozygous lines expressing *MSL10g^{7A}* in the *msl10-1* background were selected for comparison. (D) Whole-plant images of 25-day-old plants of the indicated genotypes grown side-by-side at 21°C under a 24-h light regime. Bar = 0.5 cm. (E) Petiole length of the fourth or fifth leaf from 28-day-old plants. Different letters indicate significant differences as determined by one-way ANOVA followed by Tukey's post-hoc test (P < 0.05). Error bars indicate ± SD of three replicates (n = 18 per replicate).

Like the *MSL10-GFP* overexpression lines, the *msl10-3G* mutant and three *MSL10g^{7A}* lines all

exhibited reduced rosette size, fresh weight and plant height compared to wild-type (**Figure 1D**,

S2B-D). Four- to five-week-old plants from these same lines also exhibited shorter petioles and

broader leaf blades compared to wild-type plants (**Figure 1E, S2E-F**). Unlike *MSL10-GFP*

overexpression lines, the *msl10-3G* mutant and *MSL10g^{7A}* lines lacked apical dominance (**Figure**

S2G). The *msl10-1* null mutant and *msl10-1* mutants expressing *MSL10g* or *MSL10g^{7D}* were

phenotypically indistinguishable from wild-type plants.

We next tested if *msl10-3G* mutants, *MSL10^{7A}g* lines, and *MSL10-GFP* overexpression lines have

similar gene expression patterns. As shown in **Figure 2A**, four genes previously shown to be

upregulated in *MSL10-GFP* overexpression lines (*SENESCENCE ASSOCIATED GENE 12*

(*SAG12*), *α-DIOXYGENASE (DOX1), PEROXIDASE-34 (PERX34),* and *OSMOTIN-LIKE*

PROTEIN-34 (OSM34) (Veley et al., 2014) were also upregulated in *msl10-3G* (4- to 5-fold) and

MSL10g^{7A} lines (3- to 5-fold) compared to wild-type plants, although not to the same levels as in

MSL10-GFP overexpression lines (7- to 20-fold). Similarly, four genes previously shown to be

upregulated in *rea1* mutants, *LIPOXYGENASE2* (*LOX2,*) *PLANT DEFENSIN* (*PDF1.2), ALLENE*

OXIDASE (*AOS)* and *RAP2.6* (Zou et al., 2015) were also upregulated in *MSL10-GFP*

overexpression lines (14- to 35-fold), and in *MSL10g^{7A}* lines (5- to 12-fold) (**Figure 2B**). Mutant

msl10-1, *MSL10g*, and *MSL10g^{7D}* lines did not exhibit statistically significant differences in

expression of any of these genes compared to wild-type plants. We also observed upregulation of

SAG12, DOX1, OSM34 and *PERX34* and ectopic cell death in response to inducible expression of

wild-type and phospho-dead (but not phospho-mimetic) MSL10 (**Figure S3).**

Thus, overexpression of *MSL10-GFP*, expression of untagged *MSL10^{7A}* at endogenous levels in

the *msl10-1* background, and the C-terminal point mutation in the *msl10-3G* allele all lead to

growth retardation, defects in petiole length, and the upregulation of eight hallmark genes. While

the *MSL10-GFP* overexpression line *15-2* was the most severely affected in all cases, and the only

line to display yellowish-brown patches on rosette leaves (**Figure 1D**), all of these phenotypes

could be generated to various degrees of severity without overexpression and/or a C-terminal tag,

and thus are likely to be related to the normal function of MSL10. Furthermore, these phenotypes

are a direct, rather than developmental, effect of ectopic MSL10 activation.

We interpret these data to mean that high levels of wild-type MSL10, basal levels of MSL10 that

is dephosphorylated at the N-terminus, or basal levels of MSL10 harboring the S640L lesion all

lead to a set of pleiotropic phenotypes through the same, as yet unknown, molecular mechanism.

We note that a lack of MSL10-specific antibodies prevented us from measuring protein levels in

untagged lines, so it is formally possible that MSL10 phospho-dead lesions or the S640L lesion

alter protein stability in the absence of a GFP tag. However, these lesions do not alter stability

when fused to a GFP tag (see Veley et al., 2014 and **Figure 4B** below).

Figure 2. *MSL10-GFP* **overexpression lines,** *msl10-3G* **mutants, and** *MSL10g⁷ᴬ* **lines display similar expression profiles.** Quantitative RT-PCR analysis of four genes previously shown to be upregulated in *35S:MSL10-GFP* overexpression lines (A) and four genes previously shown to be upregulated in the *msl10-3G* allele (B). cDNA was synthesized from total RNA extracted from rosette leaves of three-week-old plants grown at 21°C under 24 hours of light. Expression levels of respective genes were normalized to both *EF1α* and *UBQ5*. Mean fold-change values relative to the wild type are plotted, with error bars showing ± SE of the mean of three biological replicates Different letters indicate significant difference as determined by one-way ANOVA followed by Tukey's post-hoc test (P < 0.05). For transgenics, two independent T3 or T4 homozygous lines were selected for comparison.

Growth at high temperatures suppresses phenotypes associated with overexpression of

***MSL10-GFP,* the *MSL10g⁷ᴬ* lines, and the *msl10-3G* allele**

Growth at elevated temperature can ameliorate growth defects and suppress ectopic cell death in

a number of mutant backgrounds (Gray et al., 1998; Ichimura et al., 2006; Koini et al., 2009; Quint

et al., 2009; Wierzba and Tax, 2016), suggesting that the same might be true for *MSL10-GFP*

overexpression. When grown at 30°C, *MSL10-GFP* overexpressing lines were nearly

indistinguishable from wild-type, *msl10-1*, or *msl9-1; msl10-1* plants in terms of rosette size and

fresh weight (**Figure 3A** and **S4A**). In these lines, *MSL10* transcript levels were unchanged or only

slightly reduced (**Figure 3B**), and neither MSL10-GFP protein abundance nor localization at the

periphery of the cell was altered (**Figure 3C** and **S4B**). Thus, growth at high temperatures did not

relieve *MSL10-GFP* overexpression-associated phenotypes by altering protein abundance,

stability or subcellular localization.

Figure 3. Growth at high temperatures suppresses phenotypes associated with _MSL10-GFP_ overexpression. Wild type, _msl10-1_, and three independent T4 homozygous lines overexpressing _MSL10-GFP_ in the Col-0 background were grown side-by-side on soil under 24 h light regime and at either 21°C or 28°-30°C. (A) Images of whole three-week-old plants. Bar = 0.5 cm. (B) Relative transcript abundance of _MSL10_ analyzed by with qRT-PCR from wild type, _msl10-1_, and _35S:MSL10-GFP_ line 15-2. Inset compares _MSL10_ expression levels in wild-type and _msl10-1_ lines only. (C) Immunoblot analysis of MSL10-GFP levels in rosette leaves of two-week old plants. MSL10-GFP was detected with an anti-GFP primary antibody (top), and then the blot was stripped and re-probed with an anti-α-tubulin primary antibody (bottom). Expected protein sizes are indicated at the left according to a commercially available standard. (D) Colorimetric quantification of NBT-formazan deposition in rosette leaves (see Methods). FW, fresh weight. Data are means ± SD of three replicates (n=18). (E) Hydrogen peroxide (H_2O_2) content, quantified as mmol H_2O_2 per gram dry weight (DW) of rosette leaves using the Amplex Red-coupled fluorescence quantitative assay. Data are means ± SD of three replicates (n=14). (F) Ectopic cell death detected by Trypan blue staining (see Methods). Data are means ± SD of three replicates (n=7-10). (G) Elevation of transcripts in MSL10-GFP overexpression lines is suppressed by high temperature. RNA extraction, cDNA synthesis, and qRT-PCR analyses in (B) and (G) were performed as in Figure 2.

High temperatures also suppressed the cellular phenotypes of *MSL10-GFP* overexpression lines.
Strikingly, levels of O^{2-} (assayed by Nitroblue tetrazolium (NBT); **Figure 3D)**, H_2O_2 (assayed by
DAB staining and Amplex Red-coupled fluorescence assay; **Figure 3E**,) and ectopic cell death
(assessed by Trypan blue staining; **Figure 3F, S5B**) were strongly suppressed in *MSL10-GFP*
overexpression lines grown at high temperatures. In addition, *PERX34, DOX1, OSM34,* and
SAG12 transcript levels in *MSL10-GFP* overexpression lines were reduced to wild-type levels
when grown at 28°C-30°C (**Figure 3G)**. We also observed that the growth defects, ROS
accumulation and ectopic cell death of lines expressing the *MSL10g^{7A}* transgene and *msl10-3G*
mutants was suppressed when plants were grown at high temperatures (**Figures S5 and S6A-F**).
No apparent changes in *MSL10* transcript levels were detected in *MSL10g^{7A}* or *msl10-3G* lines
when grown at high temperature (**Figure S6G)**. Taken together, our data show that growth at 28-
30°C suppresses all phenotypes associated with the presence of over-active MSL10.
251

Phospho-mimetic lesions in the MSL10 N-terminus suppress *msl10-3G* phenotypes
Given that mutations in the N-terminus (*MSL10g^{7A}*) and the C-terminus (*msl10-3G*) produce
similar phenotypes, we tested for a genetic interaction between these two soluble domains. The
msl10-3G (S640L) mutation was introduced into the *35S:MSL10-GFP* and *35S:MSL10^{7D}-GFP*
transgenes to make *MSL10^{S640L}-GFP* and *MSL10^{7D,S640L}-GFP*. As expected, overexpression of
MSL10-GFP and *MSL10^{S640L}-GFP* lead to growth retardation, ectopic cell death (as assessed by
the occurrence of yellowish-brown lesions on rosette leaves and Trypan blue staining), and
enhanced H_2O_2 accumulation in rosette leaves, while overexpression of phospho-mimetic
MSL10^{7D}-GFP did not (**Figure 4A)**. *MSL10^{7D,S640L}-GFP* plants were indistinguishable from wild
type or *MSL10^{7D}-GFP* plants. Immunoblotting showed that these phenotypic differences cannot
be attributed to differences in protein abundance (**Figure 4B),** and likely reflect the inability of
MSL10^{7D}-GFP and MSL10^{7D,S640L}-GFP to activate downstream signaling. Thus, N-terminal
phospho-mimetic substitutions prevent or block the phenotypes produced by the C-terminal lesion
S640L found in the *msl10-3G* mutant.
266

Figure 4. Phospho-mimetic substitutions in the MSL10 N-terminus suppress phenotypes associated with overexpression of MSL10^{S640L}. (A) Top row: Images of three-week-old plants grown at 21°C overexpressing wild-type *MSL10-GFP*, phospho-mimetic *MSL10^{7D}-GFP*, *MSL10^{S640L}-GFP*, or *MSL10^{7D,S640L}-GFP*. Overexpression lines were generated in the *msl9-1;msl10-1* background, phenotypically indistinguishable from *msl10-1*. Two independent homozygous T3 lines for each transgene are shown. Bar = 0.5 cm. Middle row: Trypan blue staining of four-week old leaves from the above T3 plants to visualize ectopic cell death. Bottom row: DAB staining of five-week-old leaves to detect the accumulation of H$_2$O$_2$. For leaf images, bar = 0.2 mm. (B) Immunoblot showing that plants harboring various *MSL10-GFP* transgenes accumulate MSL10-GFP to similar levels. Western blot performed as in Figure 3C. The two forms of MSL10-GFP that migrate slower on SDS-PAGE may result from post-translational modifications.

The soluble N- and C- termini of MSL10 interact directly in two protein-protein interaction assays

To assess whether this genetic interaction between the N- and C-termini of MSL10 might be mediated through physical interactions, we first employed the mating-based split-ubiquitin yeast two-hybrid assay (mbSUS) (Reinders et al., 2002). We previously showed that MSL10 can interact with itself in this assay, but not with close relative MSL9—as expected for a homomeric channel (Veley et al., 2014). We repeated these results and further observed that MSL10 does not interact with an unrelated membrane protein, KAT1 (Obrdlik et al., 2004) (**Figure 5A)**.

To test for interactions specific to the N- and C-termini, we first split MSL10 into two halves, the first comprising the N-terminal domain and the first four TM helices (MSL10$_{1-460}$), and the second comprising the fifth and sixth TM helix as well as the C-terminal domain (MSL10$_{461-734}$). These two halves of MSL10 displayed a strong interaction, providing support for intra- or inter-molecular interactions between different domains of MSL10 (**Figure 5A**). Furthermore, the soluble N-terminus (MSL10$_{1-164}$) and the C-terminal half of MSL10 (MSL10$_{461-734}$) interacted, as did the N-terminal half of MSL10 (MSL10$_{1-460}$) and its soluble C-terminus (MSL10$_{553-734}$). This result suggested that the middle part of the protein—which contains all the TM helices—is not required for self-association. Almost no interaction was detected between the N-terminal half and itself or between the C-terminal half and itself. None of these interactions were affected by the presence of the S640L lesion (indicated as MSL10*).

To validate these mbSUS interactions, to investigate whether they can occur *in planta*, and to assess interactions without requiring one partner to be tethered to the membrane, we employed a bimolecular fluorescence complementation (BiFC) assay as described in Gehl et al. (2009). The N- and C-terminal halves of YFP (YN- and YC-) were fused to MSL10 variants and transiently expressed in leaf epidermal cells of *Nicotiana benthamiana*. As expected, co-infiltration of MSL10-YN with MSL10-YC resulted in strong YFP signal at the periphery of the cell **(Figure 5B)**. Strong YFP fluorescence was also detected when MSL10$_{553-734}$-YN was co-infiltrated with MSL10$_{1-164}$-YC (**Figure 5B**) or with MSL10$_{1-460}$-YC (**Figure S7B**). The C-terminal half of MSL10 formed aggregates **(Figure S7A)** so we only used the soluble C-terminus (MSL10 $_{553-734}$) in this study. Neither the N-terminus nor the C- terminus of MSL10 was able to self-associate in this assay, nor to interact with unfused YN or YC. Furthermore, neither of these domains were able to interact with MSL9 or KAT1 (**Figure 5B, Figure S7B**). Consistent with the mbSUS assay results, the S640L mutation (again indicated as MSL10*) did not affect any interactions; nor did the introduction of phospho-mimetic or phospho-dead lesions into the N-terminal sequences (**Figure 5B, Figure S7B**). These results demonstrate a direct physical interaction between the soluble N- and C- termini of MSL10 and show that it does not require tethering to the membrane and is unaffected by the S640L lesion or by the phosphorylation status of the N-terminus.

Figure 5. The soluble N- and C- termini of MSL10 interact. (A) Split ubiquitin yeast two-hybrid assay. The strength of interaction between MSL10 truncations was determined in liquid assays (see Methods). The NubWT moiety is used as a positive control. Data presented are means ± SD of three replicates. Different letters indicate significant differences as determined by one-way ANOVA followed by Tukey's post-hoc test (P < 0.05). (B) Confocal laser scanning micrographs of the abaxial surface of N. benthamiana leaves 3 to 5 days after infiltration with Agrobacterium harboring the indicated BiFC construct pairs. Scale bar, 50 µm. Identical acquisition settings were used in all images. All experiments were repeated at least three times. MSL10* indicates MSL10^{S640L}.

A temperature-sensitive gene network induced by *MSL10-GFP* overexpression is associated with multiple abiotic and biotic stress responses

To begin to understand the pathways downstream of MSL10 activation, we performed a transcriptome analysis, exploiting the temperature sensitivity of the *35S:MSL10-GFP* phenotype. The workflow is diagrammed in **Figure 6A**. Briefly, wild-type plants and *35S:MSL10-GFP* line *12-3* were grown for three weeks at 21°C, then half of the plants were transferred to 30°C. After 24 hours, rosette tissue was collected from duplicate pools, from both genotypes grown under both temperature regimes, resulting in a total of 8 cDNA libraries. An average of 21,910,736 reads were sequenced across all libraries with an average pseudoalignment rate of 95.40%. An assessment of sample clustering using the plotMDS function of edgeR v3.18.1; (Robinson et al., 2010)is shown in **Figure S8A**. A *k*-means clustering analysis was performed across all genotypes and temperature regimes yielded eight distinct clusters with unique response patterns. 'Cluster 2' comprised genes upregulated in the *MSL10-GFP* overexpression line at 21°C, but not at 30°C, and not in wild-type plants at either temperature (**Figure 6B, Table S2**). All eight marker genes upregulated in *MSL10* gain-of-function mutants shown in **Figure 3** were present in Cluster 2 (**Figure S8B**), providing some validation for our procedure.

Next, the 5828 unique differentially expressed genes in Cluster 2 were subjected to a gene ontology (GO) search using an overrepresentation test (PANTHER 14.1; Mi et al., 2017) and a total of 139 GO terms were found (**Table S3**). The top seven significantly overrepresented GO terms with a p-value cut-off of p=E-25, ranked by their *p*-values, are shown in **Figure 6C.** All seven categories concern the response to environmental stresses of various types, including chemicals and other organisms. Other biological processes such as 'response to acid chemical' 'cellular catabolic process' and 'multi-organism process' were also enriched.

To identify candidate genes for further validation, and to see if a similar expression pattern was observed in transgenic *MSL10^{7A}g* plants and *msl10-3G* mutants, we selected five candidate genes which satisfied the following three criteria: 1) is one of the top 100 genes identified in **Table S2** (highlighted in yellow) that was differentially expressed in *MSL10-GFP* overexpression lines at low temperatures with a threshold filter value >10, 2) belongs to at least two of the overrepresented GO terms shown in **Figure 6C** when these top 100 genes were subject to GO analysis, and 3) there

was minimal difference in the threshold value between the two biological replicates (**Table S4**).

We measured expression of these five genes (*SENESCENCE ASSOCIATED GENE* (*SAG13*),

LIPOXYGENASE3 (*LOX3*), *TRICHOME BIREFRINGENCE-LIKE 8* (*TBL8*), *ABA-INDUCED*

PP2C GENE-1 (*PP2C-1*) and *ELICITOR-ACTIVATED GENE 3-2* (*ELI3-2*)) in all *MSL10* gain-

of-function lines. All five genes were expressed at levels 75- to 150-fold higher in the *MSL10-*

GFP overexpression line than in the wild type, and a similar upregulation of these genes was

displayed in *msl10-3G* (30- to 72-fold) and *MSL10g^{7A}* lines (30- to 62-fold) compared to wild-type

plants (**Figure 6D**). In contrast, *msl10-1*, *MSL10g*, and *MSL10g^{7D}* lines did not exhibit detectable

changes in expression of any of these genes. These data validate the RNA-Seq analysis and provide

both a list of candidate marker and a starting point for a better understanding of the pathways that

lead to the phenotypic defects associated with *MSL10* activation. Finally, these results provide

further evidence that all the three *MSL10* gain-of-function alleles have similar signaling outputs.

Figure 6. Transcriptome profiling identifies downstream genes associated with *MSL10* gain-of-function lines. (A) Overview of our experimental approach. (B) Averaged expression profiles of gene clusters were identified by centroid based k-means clustering using a K-means function in R. The $\log_2$-transformed expression levels are plotted on the y-axis and samples with biological replicates on the x-axis. (C) The top seven Gene Ontology classes of genes belonging to Cluster 2 as determined through the PANTHER statistical overrepresentation test. The number of genes corresponding to each overrepresented GO term and its associated p values are shown. (D) Five genes selected from the *MSL10-GFP* overexpression RNA-Seq Cluster 2 were upregulated in *MSL10-GFP* overexpression lines, the *msl10-3G* mutant, and lines expressing the *MSL10g^{7A}* transgene. RNA extraction, cDNA synthesis, and qRT-PCR analyses were performed as in Figure 3. Plants were grown at 21°C under 24-h of light.

DISCUSSION

Here we present several lines of evidence in support of a direct interaction between the soluble N-terminus and the soluble C-terminus of the plant mechanosensitive ion channel AtMSL10. First, a suite of temperature-sensitive phenotypes was produced by overexpression of *MSL10-GFP*, by native expression of phospho-dead substitutions in MSL10's N-terminus, or by the S640L point mutation in its C-terminus. Second, phospho-mimetic substitutions suppressed the effect of S640L when both types of lesions were present in the same monomer. Third, the soluble N- and C-termini of MSL10 interacted in two protein-protein interaction assays in a manner independent of phospho-variant mutations or the S640L lesion. In addition, we showed that MSL10 activation resulted in the upregulation of genes involved in a range of pathways that can be generally characterized as responses to biotic and abiotic stress.

To date, functional characterization of MSL10 has been based on overexpression of *MSL10-GFP* in *N. benthamiana* and Arabidopsis (Veley et al., 2014). Here, we show that two additional types of *MSL10* gain-of-function lines (plants expressing endogenous levels of untagged $MSL10g^{7A}$ or harboring the *msl10-3G* allele) phenocopy *MSL10-GFP* overexpression lines, as does inducible overexpression of MSL10^{7A} (**Figure 1**, **2**, **S2**, **S5**). Thus, the ability of MSL10 to induce growth retardation, ectopic cell death, hyperaccumulation of H_2O_2, and the induction of genes related to ROS, and cell death does not require overexpression or the presence of a tag, and therefore is likely to be related to the normal function of MSL10. We note that plants overexpressing untagged wild type *MSL10* do not show these phenotypes (Zou et al., 2015). Perhaps, in this context, a large tag improves MSL10 stability.

Furthermore, the phenotypes associated with these *MSL10* gain-of-function lines was alleviated by growth at elevated temperature, while protein abundance and localization at the cell periphery remained unchanged (**Figure 3, S4** and **S6**). There is a growing list of plant defense response mutants which display partial or complete suppression of similar growth defects and ectopic cell death when grown at high temperatures (Yang et al., 2007; Gao et al., 2009; Wang et al., 2011; Wierzba and Tax, 2016), suggesting that a shared pathway leading to these phenotypes is

temperature sensitive. Alternatively, growth at high temperature could affect MSL10 activity due

to alterations in membrane fluidity (Fan and Evans, 2015).

MSL10 likely functions as a homo-oligomer similar to *Ec*MscS, a homoheptameric (Bass et al.,

2002; Sukharev, 2002). Thus, the interactions we observe between the N- and the C-terminus could

be between domains on the same monomer, or between domains on different monomers once they

are assembled into a channel. To distinguish between these two possibilities will require further

study. These interactions occur in the absence of the membrane spanning domains, at least in the

transient expression system used for BiFC (**Figure 5B**). Neither N-terminal phosphorylation status

nor the C-terminal lesion S640L had any impact on the self-association of MSL10 in either BiFC

or mbSUS assays, suggesting that this is a stable interaction and is not a regulated step in the

activation of MSL10 (**Figure 5**).

Based on these results, we propose a three-step model for MSL10 activation, illustrated in **Figure**

7. According to this model, phosphorylation of its N-terminal domain maintains MSL10 in its

inactive form (**Figure 7A**). When membrane tension is increased, the channel opens, first resulting

in a structural rearrangement at the C-terminus. Second, this conformational change is transduced

to the N-terminal domain. Third, the subsequent conformational change of the N-terminus results

in its dephosphorylation. The dephosphorylated N-terminal domain (depicted in red) is then

capable of triggering a cell death signaling cascade through an unknown mechanism.

Dephosphorylation could arise through conformational changes that directly (or indirectly through

accessory proteins like 14-3-3 proteins (Moeller et al., 2016) make the N-terminus more accessible

to cytosolic phosphatases or less accessible to kinases. Conformational changes that lead to altered

phosphorylation status has been previously shown to activate hormone-receptor complexes in

plants (Zhang et al., 2014; Farrell and Breeze, 2018; Wang et al., 2019).

In the case of *MSL10-GFP* overexpression (**Figure 7B**), we propose that the appropriate kinase is

not sufficient to maintain the N-terminus in its dephosphorylated and deactivated, form, leading to

constitutive activation of the cell death signaling pathway. Phospho-dead versions of MSL10

(MSL10g^{7A}) are constitutively active and do not require added tension; conversely, phospho-

mimetic version of MSL10 (MSL10g^{7D}) are maintained in an inactive state (**Figure 7C**). As for

plants harboring the *MSL10^{S640L}-GFP* transgene or the *msl10-3G* mutant (**Figure 7D**), the S640L

mutation may mimic the conformational change that occurs during opening of the channel,

bypassing the normal activation of signaling by membrane tension and leading to constitutive

activation. Alternatively, S640L may alter the tension sensitivity or conductivity of MSL10. This

effect, however, is blocked by the presence of N-terminal phospho-mimetic substitutions, as shown

in plants constitutively expressing *MSL10$^{7D\,,S640L}$-GFP*. Thus, this three-step model is sufficient to

explain the phenotypic and genetic interaction data documented above.

The results presented here also begin to address how the activation of MSL10 leads to phenotypic

consequences. Our transcriptomic analyses show that MSL10 activation promotes the expression

of genes previously seen to be involved in the response to osmotic stress, biotic stress and

wounding (**Figure 6D**). In fact, signaling networks originating from diverse abiotic, biotic, and

mechanical stresses are highly integrated and often culminate in a common set of responses

(Walley et al., 2007; Macqueen and Bergelson, 2016; Zhang et al., 2016). *MSL10-GFP*

overexpressing lines resemble a number of other mutant or transgenic lines associated with

constitutive induction of biotic and abiotic stress responses, including the gain-of-function mutant

SUPPRESSOR OF npr1-1, CONSTITUTIVE 1 (snc1), CHILLING SENSITIVE 2 (chs2), LESION

SIMULATING DISEASE 1 (lsd1) and plants overexpressing a *TIR-X* gene (Zhang et al., 2003b;

Huang et al., 2010a; Huang et al., 2010b; Kato et al., 2014). Both abiotic and biotic stressors such

as dehydration, flooding, or fungal penetration could induce physical alterations in the plasma

membrane that are subsequently perceived as lateral membrane tension.

In summary, our three-step model for MSL10 activation, which involves an intra- and/or

intermolecular interaction between the soluble N- and C- termini of MSL10, explains the

phenotypic similarity shared by plants overexpressing MSL10-GFP, expressing the *MSL10^{7A}*

transgene, or harboring the *msl10-3G* allele. Gene expression analyses provide evidence that

MSL10 may serves as a broad sensor of membrane perturbation that, when over-activated, leads

to cell death. These results begin to build the groundwork for future investigations into the

molecular mechanism by which the mechanosensitive ion channel MSL10 functions.

Figure 7. A three-step model for the regulation of MSL10's cell death signaling function

For simplicity, only the soluble N- and C-terminal domains of MSL10 are shown. They could be from the same or from different monomers. The inactive N-terminus, active N-terminus, and C-terminus are represented by rectangles colored blue, red, and green, respectively. Proposed conformational changes are indicated by changes in the shape of each rectangle. This model explains the signaling output (red arrow) observed in wild type plants (A), in response to MSL10-GFP overexpression (B), in the presence of MSL10 phospho-variants (C), in the presence of alleles or transgenes harboring the S640L mutation in the C-terminus (D, top), or in the presence of a transgene that combines S640L with phospho-mimetic substitutions in the N-terminus (D, bottom).

METHODS

Sequence alignments

The full-length amino acid sequence of *Arabidopsis thaliana* MSL10 was used as a BLASTp query to search for homologs in selected plant species. The protein with the highest identity to *At*MSL10 in each species was chosen for the final alignment, using the PRALINE multiple sequence alignment server (Bawono et al., 2017).

Plant lines and growth conditions

The plants used in this study are all in the Arabidopsis *Col-0* ecotype background. The T-DNA insertion lines, including *msl10-1* (SALK_114626) and *msl9-1* (SALK_114361) was obtained from the Arabidopsis Biological Resource Center (Haswell et al., 2008). The *rea1/msl10-3G* mutant was obtained from the J.-M. Zhou lab (ShanghaiTech University, Shanghai). In most experiments, plants were grown on soil at 21°C under a 24-h light regime ($\sim$120 μmol m^{-2}s^{-1}). For experiments done at elevated temperature, plants were grown on soil at high temperature that ranges from 28°C to 30°C under a 24-h light regime ($\sim$120 μmol m^{-2}s^{-1}). Backcrosses and outcrosses were made through standard techniques and genotyped with PCR-based markers. The *MSL10-GFP* overexpression lines are the same as those described in an earlier study (Veley et al., 2014).

Genotyping

DNA was extracted by grinding tissue in extraction buffer (200 mM Tris-HCl pH 7.5, 250 mM NaCl, 250 mM EDTA, 0.5% SDS) and precipitating with an equal volume of isopropanol. The *msl10-3G/rea1* allele was genotyped by amplifying the genomic region surrounding the point mutation using gene specific oligos listed in **Supplemental Table 5** followed by digestion with *TaqIα* restriction enzyme (NEB), which only digests the wild-type *MSL10* gene. PCR genotyping of *msl10-1* and *msl9-1* alleles was performed as described (Haswell et al., 2008).

Cloning and generation of transgenic plants

MSL10^{S640L} cDNA constructs were generated through site-directed mutagenesis as described (Jensen and Haswell, 2012) using pENTRY clones (pENTR+*MSL10* and pENTR+*MSL10^{7D}*) as

template (Veley et al., 2014). They were cloned into pEarleyGate103 destination vectors (Earley

et al., 2006) using LR recombination. To construct *gMSL10* plasmids, the *MSL10* gene was

amplified from genomic wild-type *Col-0* DNA using gene-specific primers listed in **Table S5** and

cloned into pENTR/D-TOPO entry vector (Thermo Fisher Scientific), then recombined into

pBGW destination vectors (Karimi et al., 2002) by LR recombination. The *MSL10g^{7A}* and

MSL10g^{7D} constructs were generated in a single reaction using a two-fragment Gibson Assembly,

using Gibson Assembly NEB Mix and overlapping primers following manufacturer's

recommendation (Thermo Fisher Scientific). The assembled plasmids were then recombined into

pBGW destination vectors by LR recombination. All primers used for Gibson cloning are listed in

Table S5. For generating constructs for conditional expression of *MSL10* and its phospho-variants

(*MSL10^{4A}* and *MSL10^{4D}*) under the dexamethasone (DEX)- inducible promoter, the Gateway

cassette-containing region from pEarleyGate100 (Earley et al., 2006), was amplified from the

plasmid introduced into the binary expression vector pTA7002 (Aoyama and Chua, 1997) using

XhoI and SpeI restriction sites. pENTR constructs containing coding region of *MSL10*, *MSL10^{4A}*

and *MSL10^{4D}* (Veley et al., 2014) were used in recombination reactions with the pTA7002 vector

using LR clonase.

Plant transformation

495 All binary constructs were transformed into wild-type *Col-0, msl10-1* or *msl9-1; msl10-1* plants

496 with *Agrobacterium tumefaciens* GV3101 by floral dip (Clough and Bent, 1998). Homozygous T3

497 or T4 lines with a single transgene insertion were identified using selectable markers, PCR

498 genotyping, RT-PCR, and/or fluorescent GFP expression and immunodetection.

Gene expression analysis

Quantitative reverse-transcription polymerase chain reaction (qRT-PCR) was performed as

described (Hamilton et al., 2015) with minor modifications. Total RNA was isolated from the

rosette leaves of healthy 3-week-old plants (before yellow necrotic lesions develop) using the

RNeasy Plant Mini Kit (Qiagen). All results shown include data from three biological replicates;

for each biological replicate, three technical replicates were performed for each of three samples.

Transcript levels for each gene was normalized against the geometric mean of the threshold cycle

(ct) values of the two reference genes (Biazzi et al., 2015), namely *UBQ5* and *EF1α.* Finally,

relative abundance of transcripts was calculated using the $2^{-\Delta\Delta Ct}$ method (Livak and Schmittgen

2001). Semi-quantitative RT-PCR was performed as described (Veley et al., 2014). For **Figure**

S3A, 10 µM DEX (Sigma-Aldrich), dissolved in 0.016% ethanol, was infiltrated into 5-week-old

leaves. Tissue was harvested for RNA isolation 12 h post-infiltration. The primers used are listed

in **Table S5.**

Immunodetection

Total proteins were extracted from non-chlorotic leaves of three-week-old plants as described

(Veley et al. 2014). Proteins samples were resolved by 10% SDS-PAGE and transferred to

polyvinylidene difluoride membranes (Millipore) for 16 h at 100 mA. Transferred proteins were

probed with anti-tubulin (Sigma, 1:10,000 dilution) and anti-GFP (Takara Bio, 1:5,000 dilution)

antibodies with incubation for 1 h and 16 h, respectively. After incubation for 1 h with horseradish

peroxidase-conjugated anti-mouse secondary antibodies (1:10,000 dilution; Millipore). Detection

was performed using the SuperSignal West Dura Detection Kit (Thermo Fisher Scientific).

Statistical analyses

Statistical evaluations were conducted using R Studio software (RStudio). Statistical differences

were analyzed as indicated in the figure legends.

Trypan blue staining and quantification

Cell death was visualized in leaves of three to four-week-old soil-grown plants using Trypan blue

staining as described (Veley et al., 2014). Images were obtained with an Olympus DP80 equipped

with a cooled color digital camera. The size of Trypan blue stained regions was quantified using

ImageJ software as described (Fernández-Bautista et al., 2016).

Detection and quantification of reactive oxygen species

Superoxide anion radical accumulation was detected by NitroBlue Tetrazolium chloride (NBT;

Sigma) as described previously (Wilson et al., 2016). The amount of formazan was determined by

measuring lysates at an absorbance of 700 nm using a 96-well microplate reader (Infinite 200 PRO;

Tecan). Absorbance reads were normalized to the fresh weight of the leaves. H_2O_2 levels were

measured by 3,3′-diaminobenzidine (DAB, Sigma) staining (Veley et al., 2014). For the

quantitative measurement of hydrogen peroxide concentrations, the Amplex Red

Hydrogen/Peroxidase Assay Kit (Molecular Probes, Invitrogen) was used, following the

manufacturer's instructions and as described in (Wilson et al., 2016).

Transcriptomic analysis by RNA sequencing

Wild type and the *MSL10-GFP* overexpressing line, *12-3* were grown for three-weeks at 21°C

under 24h light regime (∼120 μmol $m^{-2}s^{-1}$), then half of the plants were transferred to 30°C for

24 hours. RNA was extracted from the entire rosette tissue of three-week-old plants using the Plant

RNeasy Kit following manufacturer's instructions. The RNA samples were outsourced to Cofactor

Genomics (St. Louis, MO) for poly-A purification, library construction, sequencing and

preliminary analysis. Illumina TruSeq unstranded mRNA libraries were prepared and sequenced

on an Illumina HiSeq2500 with single end 50nt reads. Raw data for each library was assessed

using FASTQC, which revealed no adaptor contamination. The raw data were aligned directly to

the Araport11 cDNA annotations using kallisto v0.44.0 (Bray et al., 2016). Raw counts were used

as input for edgeR v3.18.1 (Robinson et al., 2010), normalized to library size, and initially filtered

to retain all cDNAs with at least 1 count per million mapped reads (cpm) in at least two libraries.

A multidimensional scaling (MDS) plot was constructed with the plotMDS() function (Smyth,

2005). Library cpm counts were scaled, k-means clustered with the kmeans function in R and

centroids were visually inspected for upregulation in the "MSL10 O/E 21°C" condition. To

identify genes with increased pseudocounts in the "MSL10 O/E 21°C" condition relative to all

other libraries, cDNAs were annotated for function using ThaleMine v1.10.4 (Krishnakumar et al.,

2017). For GO enrichment analysis, differentially expressed nonredundant genes (N = 5828) were

assessed using PANTHER version 14.0 (Mi et al., 2017). A statistical overrepresentation test using

default settings (P < 0.0001, Bonferroni correction) was used to determine overrepresented GO

terms for the PANTHER biological process data set.

Split-Ubiquitin Yeast two hybrid assay (mbSUS)

Intramolecular interaction between MSL10 N- and C-termini was determined using the mating-

based split-ubiquitin system as described (Obrdlik et al., 2004; Lee et al., 2019). Strength of

interaction is quantified by CPRG, a chlorophenol red-β-D-galactopyranoside colorimetric

reporter (Bhaskara et al., 2012). The PCR primers used for creating constructs for mbSUS assay

are listed in **Table S5.** All the yeast vectors were obtained from Arabidopsis Biological Resource

Center.

Bimolecular Fluorescence Complementation (BiFC) assays

Entry vectors containing various truncated versions of MSL10 coding region were recombined

into the binary vector pDEST-VYCE(R)GW or pDEST-VYNE(R)GW (Gehl et al., 2009), which

carry the C-terminal or N-terminal fragment of Venus YFP, respectively, using LR Clonase. The

PCR primers used for BiFC assay are listed in **Table S5**. All the BiFC fusion proteins were tagged

at the C-terminus. The plasmids were introduced into *Agrobacterium strain GV3101* and pairwise

combinations were co-infiltrated into 4- to 6-week-old *N. benthamiana* leaves as described (Waadt

and Kudla, 2008). To suppress posttranscriptional gene silencing, each construct pair was co-

infiltrated with *Agrobacterium* strain *AGL-1* harboring p19 (Voinnet et al., 2002). Infiltrated

abaxial leaf areas were examined for YFP signal using a confocal microscope (Olympus Fluoview

FV 3000) at 3 to 5 d post-inoculation. The experiments were performed at least three times using

different batches of plants; for each biological replicate, three independent *N. benthamiana* plants

were infiltrated. The microscope and camera settings used were as described (Veley et al., 2014).

Accession numbers

GenBank accession numbers used in **Figure 1** are: *Arabidopsis thaliana* (NP_196769.1),

Arabidopsis lyrata (XP_002873549.1), *Brassica rapa* (XP_009121883.1), *Brassica napus*

(XP_013676093.1), *Camelina sativa* (XP_010453270.1), *Populus trichocarpa*

(POPTR_0006s14640g), *Medicago truncatula* (XP_003603202.2), *Vitis vinefera*

(XP_002279755.1), *Solanum lycopersicum* (XP_004245056.1), *Solanum tuberosum*

(XP_006350354.1), *Zea mays* (XP_008649202.1), *Oryza sativa* (XP_015641284.1), *Sorghum*

bicolor (XP_002438025.1), *Brachypodium distachyon* (XP_003560953.1), *Setaria italica*

(XP_004964936.1).

The rest of the genes referred to in this study correspond to the following Arabidopsis Genome

Initiative locus identifiers: *MSL9 (At5G19520), MSL10 (At5G12080), RAP2.6 (At1g43160), AOS*

(At5g42650), LOX2 (At3g45140), PDF1.2 (At5G44420), SAG12 (At3g20770), PERX34

(At3g49120), DOX1 (At3g01420), OSM34 (At4G11650), UBQ5 (At3g62250), EF1α (At5g60390),

SAG13 (At2g29350), LOX3 (At1g17420), ELI3-2 (At4g37990). PP2C-1 (At5g59220), TBL8
(At3g11480).
603

604 **AUTHOR CONTRIBUTIONS**
605

612 **ACKNOWLEDGMENTS**
613

We thank J.-M. Zhou (ShanghaiTech University, Shanghai) for providing *rea1* mutant seeds and
the Arabidopsis Biological Resource Center for seeds and for vectors used in BiFC and mbSUS
assays. We also are grateful to Ivan Radin, Yanbing Wang, Kari Miller and Angela Schlegel for
critical reading of the manuscript. This work was supported by NSF MCB-1253103 and the NSF
Center for Engineering Mechanobiology Award 1548571. J.M.S. was supported by NSF Graduate
Research Fellowship DGE-1745038. A.H was supported by an NSF National Plant Genome
Initiative Postdoctoral Research Fellowship in Biology Award 1611853. No conflicts of interest
declared.

REFERENCES

Aoyama, T., and Chua, N. (1997). A glucocorticoid-mediated transcriptional induction system
in transgenic plants. *Plant Journal* **11**:605–612.
Bass, R. B., Strop, P., Barclay, M., and Rees, D. C. (2002). Crystal structure of *Escherichia*
coli MscS, a voltage-modulated and mechanosensitive channel. *Science* **298**:1582–1587.
Basu, D., and Haswell, E. S. (2017). Plant mechanosensitive ion channels: an ocean of
possibilities. *Current Opinion in Plant Biology* **40**:43–48.
Bavi, N., Nikolaev, Y. A., Bavi, O., Ridone, P., Martinac, A. D., Nakayama, Y., Cox, C. D.,
and Martinac, B. (2017). Principles of mechanosensing at the membrane interface. In *The*

Biophysics of Cell Membranes: Biological Consequences (ed. Epand, R. M. and

Ruysschaert, J.-M.), pp. 85–119. Singapore: Springer Singapore.

Bawono, P., Dijkstra, M., Pirovano, W., Feenstra, A., Abeln, S., and Heringa, J. (2017).

Multiple sequence alignment. In *Bioinformatics: Volume I: Data, Sequence Analysis, and*

Evolution (ed. Keith, J. M.), pp. 167–189. New York, NY: Springer New York.

Bhaskara, G. B., Nguyen, T. T., and Verslues, P. E. (2012). Unique drought resistance

functions of the highly ABA-induced clade A protein phosphatase 2Cs. *Plant Physiology*

160:379–395.

Biazzi, E., Carelli, M., Tava, A., Abbruscato, P., Losini, I., Avato, P., Scotti, C., and

Calderini, O. (2015). CYP72A67 catalyzes a key oxidative step in *Medicago truncatula*

hemolytic saponin biosynthesis. *Molecular Plant* **8**:1493–1506.

Bray, N. L., Pimentel, H., Melsted, P., and Pachter, L. (2016). Near-optimal probabilistic

RNA-seq quantification. *Nature Biotechnology* **34**:525–527.

Clough, S. J., and Bent, A. F. (1998). Floral dip: a simplified method for Agrobacterium -

mediated transformation of *Arabidopsis thaliana*. *The Plant Journal* **16**:735–743.

Earley, K. W., Haag, J. R., Pontes, O., Opper, K., Juehne, T., Song, K., and Pikaard, C. S.

(2006). Gateway-compatible vectors for plant functional genomics and proteomics. *Plant*

Journal **45**:616–629.

Fan, W., and Evans, R. M. (2015). Turning up the heat on membrane fluidity. *Cell* **161**:962–

963.

Farrell, B., and Breeze, A. L. (2018). Structure, activation and dysregulation of fibroblast

growth factor receptor kinases: perspectives for clinical targeting. *Biochemical Society*

Transactions **46**:1753–1770.

Fernández-Bautista, N., Domínguez-Núñez, J. A., Moreno, M. M. C., and Berrocal-Lobo,

M. (2016). Plant tissue trypan blue staining during phytopathogen infection. *Bio-protocol*

6:e2078.

Gao, M., Wang, X., Wang, D., Xu, F., Ding, X., Zhang, Z., Bi, D., Cheng, Y. T., Chen, S.,

Li, X., et al. (2009). Regulation of cell death and innate immunity by two receptor-like

kinases in Arabidopsis. *Cell Host and Microbe* **6**:34–44.

Gehl, C., Waadt, R., Mendel, R., and Ha, R. (2009). New GATEWAY vectors for high-

throughput analyses of protein – protein interactions by bimolecular fluorescence

complementation. *Molecular Plant* **2**:1051–1058.

Gobert, A., Isayenkov, S., Voelker, C., Czempinski, K., and Maathuis, F. J. M. (2007). The

two-pore channel *TPK1* gene encodes the vacuolar K$^+$ conductance and plays a role in K$^+$

homeostasis. *Proceedings of the National Academy of Sciences of the United States of*

America **104**:1–6.

Gray, W. M., Ostin, A., Sandberg, G., Romano, C. P., and Estelle, M. (1998). High

temperature promotes auxin-mediated hypocotyl elongation in Arabidopsis. *Proceedings of*

the National Academy of Sciences of the United States of America **95**:7197–7202.

Hamant, O., and Haswell, E. S. (2017). Life behind the wall: sensing mechanical cues in

plants. *BMC Biology* **15**:59.

Hamilton, E. S., Jensen, G. S., Maksaev, G., Katims, A., Sherp, A. M., and Haswell, E. S.

(2015). Mechanosensitive channel MSL8 regulates osmotic forces during pollen hydration

and germination. *Science* **350**:438–441.

Haswell, E. S., and Meyerowitz, E. M. (2006). MscS-like proteins control plastid size and

shape in *Arabidopsis thaliana*. *Current Biology* **16**:1–11.

Haswell, E. S., Peyronnet, R., Barbier-Brygoo, H., Meyerowitz, E. M., and Frachisse, J. M.

(2008). Two MscS homologs provide mechanosensitive channel activities in the

Arabidopsis root. *Current Biology* **18**:730–734.

Hou, C., Tian, W., Kleist, T., He, K., Garcia, V., Bai, F., Hao, Y., Luan, S., and Li, L.

(2014). Osmosensitive calcium-permeable cation channels 632 DUF221 proteins are a

family of osmosensitive calcium- permeable cation channels conserved across eukaryotes.

Cell Research **24**:632–635.

Huang, X., Li, Y., Zhang, X., Zuo, J., and Yang, S. (2010a). The Arabidopsis LSD1 gene

plays an important role in the regulation of low. *New Phytologist* **187**:301–12.

Huang, X., Li, J., Bao, F., Zhang, X., and Yang, S. (2010b). A gain-of-function mutation in

the Arabidopsis disease resistance gene *RPP4* confers sensitivity to low temperature. *Plant*

Physiology **154**:796–809.

Ichimura, K., Casais, C., Peck, S. C., Shinozaki, K., and Shirasu, K. (2006). MEKK1 is

required for MPK4 activation and regulates tissue-specific and temperature-dependent cell

death in Arabidopsis. *Journal of Biological Chemistry* **281**:36969–36976.

Jayaraman, D., Gilroy, S., and Ané, J. M. (2014). Staying in touch: Mechanical signals in

plant-microbe interactions. *Current Opinion in Plant Biology* **20**:104–109.

Jensen, G. S., and Haswell, E. S. (2012). Functional analysis of conserved motifs in the

mechanosensitive channel homolog MscS-Like2 from *Arabidopsis thaliana*. *PLoS ONE* **7**.

Karimi, M., Inzé, D., and Depicker, A. (2002). GATEWAY™ vectors for Agrobacterium-

mediated plant transformation. *Trends in Plant Science* **7**:193–195.

Kato, H., Saito, T., Ito, H., Komeda, Y., and Kato, A. (2014). Overexpression of the *TIR-X*

gene results in a dwarf phenotype and activation of defense-related gene expression in

Arabidopsis thaliana. *Journal of Plant Physiology* **171**:382–388.

Kierzkowski, D. (2019). Cellular basis of growth in plants : geometry matters. *Current Opinion*

in Plant Biology **47**:56–63.

Koini, M. A., Alvey, L., Allen, T., Tilley, C. A., Harberd, N. P., Whitelam, G. C., and

Franklin, K. A. (2009). High temperature-mediated adaptations in plant architecture

require the bHLH transcription factor PIF4. *Current Biology* **19**:408–413.

Kovalchuk, N., Jia, W., Eini, O., Morran, S., Pyvovarenko, T., Fletcher, S., Bazanova, N.,

Harris, J., Beck-Oldach, K., Shavrukov, Y., et al. (2013). Optimization of TaDREB3

gene expression in transgenic barley using cold-inducible promoters. *Plant Biotechnology*

Journal **11**:659–670.

Krishnakumar, V., Contrino, S., Cheng, C. Y., Belyaeva, I., Ferlanti, E. S., Miller, J. R.,

Vaughn, M. W., Micklem, G., Town, C. D., and Chan, A. P. (2017). Thalemine: A

warehouse for Arabidopsis data integration and discovery. *Plant and Cell Physiology* **58**:e4.

Lee, C. P., Maksaev, G., Jensen, G. S., Murcha, M. W., Wilson, M. E., Fricker, M., Hell, R.,

Haswell, E. S., Millar, A. H., and Sweetlove, L. J. (2016). MSL1 is a mechanosensitive

ion channel that dissipates mitochondrial membrane potential and maintains redox

homeostasis in mitochondria during abiotic stress. *Plant Journal* **88**:809–825.

Lee, J. S., Wilson, M. E., Richardson, R. A., and Haswell, E. S. (2019). Genetic and physical

interactions between the organellar mechanosensitive ion channel homologs MSL1, MSL2,

and MSL3 reveal a role for inter-organellar communication in plant development. *Plant*

Direct **3**:e00124.

Livak, K. J., and Schmittgen, T. D. (2001). Analysis of relative gene expression data using

real-time quantitative PCR and the 2-$\Delta\Delta$CT method. *Methods* **25**:402–408.

Maathuis, F. J. M. (2011). Vacuolar two-pore K^+ channels act as vacuolar osmosensors. *New*

Phytologist **191**:84–91.

Machiyama, H., Tatsumi, H., and Sokabe, M. (2009). Structural changes in thecytoplasmic

domain of the mechanosensitive channel MscS during opening. *Biophysical Journal*

97:1048–1057.

Macqueen, A., and Bergelson, J. (2016). Modulation of R-gene expression across

environments. *Journal of Experimental Botany* **67**:2093–2105.

Maksaev, G., and Haswell, E. S. (2012). MscS-Like10 is a stretch-activated ion channel from

Arabidopsis thaliana with a preference for anions. *Proceedings of the National Academy of*

Sciences of the United States of America **109**:19015–20.

Maksaev, G., Shoots, J. M., Ohri, S., and Haswell, E. S. (2018). Nonpolar residues in the

presumptive pore-lining helix of mechanosensitive channel MSL10 influence channel

behavior and establish a nonconducting function. *Plant Direct* **2**:e00059.

Martinac, B., Nomura, T., Chi, G., Petrov, E., Rohde, P. R., Battle, A. R., Foo, A.,

Constantine, M., Rothnagel, R., Carne, S., et al. (2013). Bacterial mechanosensitive

channels: Models for studying mechanosensory transduction. *Antioxidants & Redox*

Signaling **20**:952–969.

Mi, H., Huang, X., Muruganujan, A., Tang, H., Mills, C., Kang, D., and Thomas, P. D.

(2017). PANTHER version 11: Expanded annotation data from Gene Ontology and

Reactome pathways, and data analysis tool enhancements. *Nucleic Acids Research*

45:D183–D189.

Moeller, H. B., Slengerik-Hansen, J., Aroankins, T., Assentoft, M., MacAulay, N.,

Moestrup, S. K., Bhalla, V., and Fenton, R. A. (2016). Regulation of the water channel

aquaporin-2 via 14-3-3θ and -ζ. *Journal of Biological Chemistry* **291**:2469–2484.

Monshausen, G. B., and Haswell, E. S. (2013). A force of nature: Molecular mechanisms of

mechanoperception in plants. *Journal of Experimental Botany* **64**:4663–4680.

Mori, K., Renhu, N., Naito, M., Nakamura, A., Shiba, H., Yamamoto, T., Suzaki, T., Iida,

H., and Miura, K. (2018). Ca^{2+}-permeable mechanosensitive channels MCA1 and MCA2

mediate cold-induced cytosolic Ca^{2+} increase and cold tolerance in Arabidopsis. *Scientific*

reports **8**:550.

Murthy, S. E., Dubin, A. E., Whitwam, T., Jojoa-Cruz, S., Cahalan, S. M., Mousavi, S. A.

R., Ward, A. B., and Patapoutian, A. (2018). OSCA/TMEM63 are an evolutionarily

conserved family of mechanically activated ion channels. *eLife* **7**:e41844.

Nakagawa, Y., Katagiri, T., Shinozaki, K., Qi, Z., Tatsumi, H., Furuichi, T., Kishigami, A., Sokabe, M., Kojima, I., Sato, S., et al. (2007). Arabidopsis plasma membrane protein crucial for Ca^{2+} influx and touch sensing in roots. *Proceedings of the National Academy of Sciences of the United States of America* **104**:3639–3644.

Obrdlik, P., El-Bakkoury, M., Hamacher, T., Cappellaro, C., Vilarino, C., Fleischer, C., Ellerbrok, H., Kamuzinzi, R., Ledent, V., Blaudez, D., et al. (2004). K^+ channel interactions detected by a genetic system optimized for systematic studies of membrane protein interactions. *Proceedings of the National Academy of Sciences of the United States of America* **101**:12242–12247.

Pivetti, C. D., Yen, M.-R., Miller, S., Busch, W., Tseng, Y.-H., Booth, I. R., and Saier Jr, M. H. (2003). Two families of mechanosensitive channel proteins. *Microbiology and Molecular Biology Reviews* **67**:66–85.

Quint, M., Barkawi, L. S., Fan, K.-T., Cohen, J. D., and Gray, W. M. (2009). Arabidopsis IAR4 modulates auxin response by regulating auxin homeostasis. *Plant Physiology* **150**:748–758.

Ranade, S. S., Syeda, R., and Patapoutian, A. (2015). Mechanically activated ion channels. *Neuron* **87**:1162–1179.

Reinders, A., Schulze, W., Thaminy, S., Stagljar, I., Frommer, W. B., and Ward, J. M. (2002). Intra- and intermolecular interactions in sucrose transporters at the plasma membrane detected by the split-ubiquitin system and functional assays. *Structure* **10**:763–772.

Robinson, M. D., McCarthy, D. J., and Smyth, G. K. (2010). edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics (Oxford, England)* **26**:139–140.

Rowe, I., Anishkin, A., Kamaraju, K., Yoshimura, K., and Sukharev, S. (2014). The cytoplasmic cage domain of the mechanosensitive channel MscS is a sensor of macromolecular crowding. *The Journal of General Physiology* **143**:543–557.

Smyth, G. K. (2005). Linear models for microarray data. In *Bioinformatics and Computational Biology Solutions Using R and Bioconductor* (ed. Gentleman, R.), Carey, V. J., Huber, W., Irizarry, R. A., and Dudoit, S., pp. 397–420. New York, NY: Springer New York.

Spartz, A. K., Lee, S. H., Wenger, J. P., Gonzalez, N., Itoh, H., Inzé, D., Peer, W. A.,

Murphy, A. S., Overvoorde, P. J., and Gray, W. M. (2012). The SAUR19 subfamily of

SMALL AUXIN UP RNA genes promote cell expansion. *The Plant Journal* **70**:978–990.

Sukharev, S. (2002). Purification of the small mechanosensitive channel of *Escherichia coli*

(MscS): the subunit structure, conduction, and gating characteristics in liposomes.

Biophysical Journal **83**:290–298.

Veley, K. M., Marshburn, S., Clure, C. E., and Haswell, E. S. (2012). Mechanosensitive

channels protect plastids from hypoosmotic stress during normal plant growth. *Current*

Biology **22**:408–413.

Veley, K. M., Maksaev, G., Frick, E. M., January, E., Kloepper, S. C., and Haswell, E. S.

(2014). *Arabidopsis* MSL10 has a regulated cell death signaling activity that is separable

from its mechanosensitive ion channel activity. *The Plant Cell* **26**:3115–3131.

Voinnet, O., Pinto, Y. M., and Baulcombe, D. C. (2002). Suppression of gene silencing: A

general strategy used by diverse DNA and RNA viruses of plants. *Proceedings of the*

National Academy of Sciences **96**:14147–14152.

Waadt, R., and Kudla, J. (2008). In plant visualization of protein interactions using

bimolecular fluorescence complementation (BiFC). *Cold Spring Harbor Protocols* **3**:1–8.

Walley, J. W., Coughlan, S., Hudson, M. E., Covington, M. F., Kaspi, R., Banu, G.,

Harmer, S. L., and Dehesh, K. (2007). Mechanical stress induces biotic and abiotic stress

responses via a novel cis-element. *PLoS Genetics* **3**:1800–1812.

Wang, W., Dong, C., Naismith, J. H., Black, S. S., Edwards, M. D., Miller, S., Morrison, E.

L., Bartlett, W., and Booth, I. R. (2008). The structure of an open form of an *E. coli*

mechanosensitive channel at 3.45 å resolution. *Science* **321**:1179–1183.

Wang, Z., Meng, P., Zhang, X., Ren, D., and Yang, S. (2011). BON1 interacts with the protein

kinases BIR1 and BAK1 in modulation of temperature-dependent plant growth and cell

death in Arabidopsis. *Plant Journal* **67**:1081–1093.

Wang, J., Wang, J., Hu, M., Wu, S., Qi, J., Wang, G., Han, Z., Qi, Y., Gao, N., Wang, H.-

W., et al. (2019). Ligand-triggered allosteric ADP release primes a plant NLR complex.

Science **364**:eaav5868.

Wierzba, M. P., and Tax, F. E. (2016). An allelic series of *bak1* mutations differentially alter

bir1 cell death, immune response, growth, and root development phenotypes in *Arabidopsis*

thaliana. *Genetics* **202**:689–702.

Wilson, M. E., Mixdorf, M., Berg, R. H., and Haswell, E. S. (2016). Plastid osmotic stress
influences cell differentiation at the plant shoot apex. *Development* **143**:3382–3393.

Yang, H., Yang, S., Li, Y., and Hua, J. (2007). The Arabidopsis BAP1 and BAP2 genes are
general inhibitors of programmed cell death. *Plant Physiology* **145**:135–146.

Yuan, F., Yang, H., Xue, Y., Kong, D., Ye, R., Li, C., Zhang, J., Theprungsirikul, L., Shrift,
T., Krichilsky, B., et al. (2014). OSCA1 mediates osmotic-stress-evoked Ca^{2+} increases
vital for osmosensing in Arabidopsis. *Nature* **514**:367–71.

Zhang, X., Liang, R., Chen, X., Yang, F., and Zhang, L. (2003a). Transgene inheritance and
quality improvement by expressing novel HMW glutenin subunit (HMW-GS) genes in
winter wheat. *Chinese Science Bulletin* **48**:771–776.

Zhang, Y., Goritschnig, S., Dong, X., and Li, X. (2003b). A gain-of-function mutation in a
plant disease resistance gene leads to constitutive activation of downstream signal
transduction pathways in *suppressor of npr1-1, constitutive 1*. *The Plant Cell* **15**:2636–
2646.

Zhang, J., Yu, J., and Wen, C.-K. (2014). An alternate route of ethylene receptor signaling.
Frontiers in Plant Science **5**:1–6.

Zhang, M., Koh, J., Liu, L., Shao, Z., Liu, H., Hu, S., Zhu, N., Dufresne, C. P., Chen, S.,
and Wang, Q. (2016). Critical role of COI1-dependent jasmonate pathway in AAL toxin-
induced PCD in tomato revealed by comparative proteomics. *Scientific Reports* **6**:1–14.

Zhang, Z., Tateda, C., Jiang, S., Shrestha, J., Jelenska, J., Speed, D. J., and Greenberg, J.
T. (2017). A suite of receptor-like kinases and a putative mechano-sensitive channel are
involved in autoimmunity and plasma membrane – based defenses in Arabidopsis.
Molecular Plant-Microbe Interactions **30**:150–160.

Zhang, Z., Tong, X., Liu, S.-Y., Chai, L.-X., Zhu, F.-F., Zhang, X.-P., Zou, J.-Z., and Wang,
X.-B. (2019). Genetic analysis of a Piezo-like protein suppressing systemic movement of
plant viruses in *Arabidopsis thaliana*. *Scientific Reports* **9**:3187.

Zou, Y., Chintamanani, S., He, P., Fukushige, H., Yu, L., Shao, M., Zhu, L., Hildebrand, D.
F., Tang, X., and Zhou, J.-M. (2015). A gain-of-function mutation in Msl10 triggers cell
death and wound-induced hyperaccumulation of JA in Arabidopsis. *Journal of Integrative*
Plant Biology **58**:600–609.

Figure 1. *MSL10-GFP* overexpression lines, *msl10-3G* mutants, and *MSL10g^{7A}* lines have similar whole-plant phenotypes. **(A)** Predicted topology of an *At*MSL10 monomer. Teal, cytosolic N-terminus; yellow, transmembrane helices; green, cytosolic C-terminus. Black triangles denote the positions of the residues mutated in this study. The MscS domain is as defined in (Veley et al. 2014). **(B)** Multiple alignment of protein sequences corresponding to the N-terminus of putative *At*MSL10 orthologs in the *Brassicaceae* family. Darker teal shading indicates higher conservation score and identical amino acids with an asterisk. **(C)** Multiple alignment of protein sequences corresponding to the C-terminus of putative *At*MSL10 orthologs in selected flowering plants. Conservation of residues is indicated as in **(B). (D)** and **(E)** T4 homozygous lines overexpressing *MSL10-GFP* in the *Col-0* background (line *15-2*), T3 homozygous lines expressing *MSL10g*, or *MSL10g^{7D}* in the *msl10-1* background, and T3 segregating or T3 homozygous lines expressing *MSL10g^{7A}* in the *msl10-1* background were selected for comparison. **(D)** Whole-plant images of 25-day-old plants of the indicated genotypes grown side-by-side at 21°C under a 24-h light regime. Bar = 0.5 cm. **(E)** Petiole length of the fourth or fifth leaf from 28-day-old plants. Different letters indicate significant differences as determined by one-way ANOVA followed by Tukey's post-hoc test ($P < 0.05$). Error bars indicate ± SD of three replicates (n = 18 per replicate).

Figure 2. *MSL10-GFP* overexpression lines, *msl10-3G* mutants, and *MSL10g^{7A}* lines display similar expression profiles. Quantitative RT-PCR analysis of four genes previously shown to be upregulated in *35S:MSL10-GFP* overexpression lines **(A)** and four genes previously shown to be upregulated in the *msl10-3G* allele **(B)**. cDNA was synthesized from total RNA extracted from rosette leaves of three-week-old plants grown at 21°C under 24 hours of light. Expression levels of respective genes were normalized to both *EF1α* and *UBQ5*. Mean fold-change values relative to the wild type are plotted, with error bars showing ± SE of the mean of three biological replicates Different letters indicate significant difference as determined by one-way ANOVA followed by Tukey's post-hoc test ($P < 0.05$). For transgenics, two independent T3 or T4 homozygous lines were selected for comparison.

Figure 3. Growth at high temperatures suppresses phenotypes associated with *MSL10-GFP* overexpression. Wild type, *msl10-1*, and three independent T4 homozygous lines expressing

35S:MSL10-GFP in the Col-0 background were grown side-by-side on soil under 24 h light regime

and at either 21°C or 28°-30°C. **(A)** Images of whole three-week-old plants. Bar = 0.5 cm. **(B)**

Relative transcript abundance of *MSL10* analyzed by with qRT-PCR from wild type, *msl10-1*, and

35S:MSL10-GFP line *15-2*. Inset compares *MSL10* expression levels in wild-type and *msl10-1*

lines only. **(C)** Immunoblot analysis of MSL10-GFP levels in rosette leaves of two-week old

plants. MSL10-GFP was detected with an anti-GFP primary antibody (top), and then the blot was

stripped and re-probed with an anti-α-tubulin primary antibody (bottom). Expected protein sizes

are indicated at the left according to a commercially available standard. **(D)** Colorimetric

quantification of NBT-formazan deposition in rosette leaves (see Methods). FW, fresh weight.

Data are means ± SD of three replicates (n=18). **(E)** Hydrogen peroxide (H_2O_2) content, quantified

as mmol H_2O_2 per gram dry weight (DW) of rosette leaves using the Amplex Red-coupled

fluorescence quantitative assay. Data are means ± SD of three replicates (n=14). **(F)** Ectopic cell

death detected by Trypan blue staining (see Methods). Data are means ± SD of three replicates

(n=7-10). **(G)** Elevation of transcripts in *MSL10-GFP* overexpression lines is suppressed by high

temperature. RNA extraction, cDNA synthesis, and qRT-PCR analyses in **(B)** and **(G)** were

performed as in Figure 2.

Figure 4. Phospho-mimetic substitutions in the MSL10 N-terminus suppress phenotypes

associated with overexpression of MSL10^{S640L}. (A) Top row: Images of three-week-old plants

grown at 21°C overexpressing wild-type *MSL10-GFP,* phospho-mimetic *MSL10^{7D}-GFP,*

MSL10^{S640L}-GFP, or *MSL10^{7D,S640L}-GFP*. Overexpression lines were generated in the *msl9-*

1;msl10-1 background, phenotypically indistinguishable from *msl10-1*. Two independent

homozygous T3 lines for each transgene are shown. Bar = 0.5 cm. **Middle row:** Trypan blue

staining of four-week old leaves from the above T3 plants to visualize ectopic cell death. **Bottom**

row: DAB staining of five-week-old leaves to detect the accumulation of H_2O_2. For leaf images,

bar = 0.2 mm. **(B)** Immunoblot showing that plants harboring various *MSL10-GFP* transgenes

accumulate MSL10-GFP to similar levels. Western blot performed as in Figure 3C. The two forms

of MSL10-GFP that migrate slower on SDS-PAGE may result from posttranslational

modifications.

Figure 5. The soluble N- and C- termini of MSL10 interact. (A) Split ubiquitin yeast two-hybrid assay. The strength of interaction between MSL10 truncations was determined in liquid assays (see Methods). The NubWT moiety is used as a positive control. Data presented are means ± SD of three replicates. Different letters indicate significant differences as determined by one-way ANOVA followed by Tukey's post-hoc test ($P < 0.05$). **(B)** Confocal laser scanning micrographs of the abaxial surface of *N. benthamiana* leaves 3 to 5 days after infiltration with *Agrobacterium* harboring the indicated BiFC construct pairs. Scale bar, 50 μm. Identical acquisition settings were used in all images. All experiments were repeated at least three times. MSL10* indicates MSL10^{S640L}.

Figure 6. Transcriptome profiling identifies downstream genes associated with *MSL10* gain-of-function lines. **(A)** Overview of our experimental approach. (**B)** Averaged expression profiles of gene clusters were identified by centroid based *k*-means clustering using a *K*-means function in R. The log$_2$-transformed expression levels are plotted on the y-axis and samples with biological replicates on the x-axis. **(C)** The top seven Gene Ontology classes of genes belonging to Cluster 2 as determined through the PANTHER statistical overrepresentation test. The number of genes corresponding to each overrepresented GO term and its associated *p* values are shown. **(D)** Five genes selected from the *MSL10-GFP* overexpression RNA-Seq Cluster 2 were upregulated in *MSL10-GFP* overexpression lines, the *msl10-3G* mutant, and lines expressing the *MSL10g^{7A}* transgene. RNA extraction, cDNA synthesis, and qRT-PCR analyses were performed as in **Figure 3.** Plants were grown at 21°C under 24-h of light.

Figure 7. A three-step model for the regulation of MSL10's cell death signaling function
For simplicity, only the soluble N- and C-terminal domains of MSL10 are shown. They could be from the same or from different monomers. The inactive N-terminus, active N-terminus, and C-terminus are represented by rectangles colored blue, red, and green, respectively. Proposed conformational changes are indicated by changes in the shape of each rectangle. This model explains the signaling output (red arrow) observed in wild type plants **(A)**, in response to MSL10-GFP overexpression **(B)**, in the presence of MSL10 phospho-variants **(C)**, in the presence of alleles or transgenes harboring the S640L mutation in the C-terminus (**D**, top), or in the presence of a

transgene that combines S640L with phospho-mimetic substitutions in the N-terminus (**D**,
bottom).

Supplemental Figure 1. *msl10-3G* is a recessive gain-of-function allele.

(A) Representative images of parental and F1 offspring plants from a *msl10-3G* backcross to wild
type *Col-0*. Top, four-week-old; bottom, eight-week-old plants. Bar = 0.5 cm. **(B)** Representative
images of parental and F1 offspring from a *msl10-3G* outcross to *msl10-1*. Top, four-week-old
plants; bottom, twelve-week-old plants. Bar = 0.5 cm. **(C)** Table showing segregation of *msl10-
3G* inheritance data compared to the expected ratios using a chi-squared analysis. Plants were
randomly chosen for genotyping.

Supplemental Figure 2. *msl10-3G* allele and *MSL10g^{7A}* transgene exhibit similar reduction
in fresh weight, petiole length, plant height and apical dominance to *35S:MSL10-GFP*

(A) Relative transcript abundance of *MSL10* in the indicated lines. Left panel displays
accumulation of *MSL10* transcripts in plants harboring the *35S:MSL10-GFP* transgene, genomic
MSL10g phospho-variants, the EMS-induced point mutant allele *msl10-3G* or the null mutant
allele *msl10-1* compared to wild-type *Col-0*. Right panel depicts *MSL10* expression only in wild
type, null mutant allele and genomic *MSL10g* phospho-variants. cDNA was synthesized from total
RNA extracted from rosette leaves of three-week-old plants. Expression levels of respective genes
were analyzed with qRT-PCR and normalized to both *EF1α* and *UBQ5*. Mean fold-change values
relative to the wild type are plotted, with error bars showing ±SE of the mean of three biological
replicates. **(B)** Average fresh weight of rosettes clipped from three-week-old plants of the indicated
genotypes. Error bars indicate ±SD of three replicates (n=10 per replicate). **(C)** Plant height
measurement from eight-week-old plants from indicated genotypes. Error bars indicate ±SD of
three replicates (n=20-30 per replicate). Different letters indicate significant difference as
determined by one-way ANOVA followed by Scheffe's post-hoc test ($P < 0.05$) for unbalanced
samples. **(D)** Representative images of eight-week-old adult plants from indicated genotypes. **(E)**
Leaf blade length of seventh leaf from 35-day-old plants from the indicated genotypes. Error bars
indicate ±SD of three replicates (n=18 per replicate). **(F)** Leaf blade width of fourth or fifth leaf
from 35-day-old plants from the indicated genotypes. Error bars indicate ±SD of three replicates
(n=18 per replicate). **(G)** Average number of floral apices were counted from six to seven-week-

old plants of the indicated genotypes. Error bars indicate ±SD of three replicates (n=20 per

replicate). The T4 homozygous transgenic lines expressing either *35S:MSL10-GFP* in *Col-0*

background (*15-2*) and T3 homozygous transgenic lines expressing *MSL10g*, or *MSL10g^{7D}* in the

msl10-1 background, and T3 segregating or T3 homozygous lines for *MSL10g^{7A}* were selected for

comparison. Plants of indicated genotypes were grown side-by-side in soil at 21°C under a 24- h

light regime. Different letters indicate significant differences, as determined by one-way ANOVA

followed by Tukey's post-hoc test (P < 0.05).

Supplemental Figure 3. DEX-inducible overexpression of *MSL10* and *MSL10^{4A}* promotes the

upregulation of MSL10-associated marker genes and ectopic cell death. Representative

images of five-week old plants grown under a short-day photoperiod (10 h light/14 h dark) sprayed

with 0.016% ethanol (mock; top row) or 30 μM DEX (bottom row) for 5 days. Expression profiles

of five marker genes before and after DEX treatment of transgenic plants expressing *DEX:MSL10*

and its phospho-variants using semi-quantitative RT-PCR. Rosette leaves from five-week-old

plants were infiltrated with 10 μM DEX or mock (0.016% ethanol), and tissue was harvested 12 h

post infiltration. Total RNA was isolated from *Col-0*, and from two independent homozygous T3

lines expressing either *DEX:MSL10*, *DEX:MSL10^{4A}* or *DEX:MSL10^{4D}*. Expression of *UBQ5* was

used as a control.

Supplemental Figure 4. Growth at elevated temperature rescues the whole plant phenotypes

associated with overexpression of *MSL10-GFP*. (A) Comparison of average fresh weight of

rosettes clipped from three-week-old plants of the indicated genotypes grown side-by-side at 21°C

or 28-30°C under a 24-h light regime. Error bars indicate SD of three replicates (n=10 per

replicate). FW= fresh weight. **(B)** Confocal images of epidermal cells showing GFP signal from

three-week-old plants of indicated genotypes. WT was used as a background control. T4

homozygous transgenic lines overexpressing *MSL10-GFP* were used.

Supplemental Figure 5. High temperature suppresses the accumulation of H_2O_2 and the

incidence of ectopic cell death associated with *MSL10* gain-of function lines. Leaves from

three-week-old plants from each line grown either at 21°C (top row) or at an elevated temperature

of 28-30°C (bottom row) under 24-h light regime were used for the experiment. Three independent

T3 homozygous transgenic lines expressing *MSL10g, MSL10g^{7A}* or *MSL10g^{7D}* in the *msl10-1*

background were selected for comparison. **(A)** Elevated levels of H_2O_2 detected by DAB staining.

Representative bright-field images of rosette leaves are shown. Bar = 200 μm. **(B)** Visualization

of ectopic cell death incidence by Trypan blue staining. Bar = 200 μm.

1006

Supplemental Figure 6. Growth retardation, accumulation of ROS and incidence of ectopic

cell death associated with expression of the *MSL10g^{7A}* transgene and the *msl10-3G* allele were

suppressed by growth at high temperature. Three independent T3 homozygous transgenic lines

expressing *MSL10g, MSL10g^{7A}*, or *MSL10g^{7D}* in the *msl10-1* background were selected for

comparison. Plants of indicated genotypes were grown side-by-side on soil under a 24 h light

regime at either 21°C or 28°-30°C. Different letters indicate significant differences, as determined

by two-way ANOVA followed by Tukey's post-hoc test ($P < 0.05$). **(A)** Morphological phenotypes

of four-week-old plants of the indicated genotypes. Bar = 0.5 cm. **(B)** Plant height of the indicated

genotypes. Data are means ± SD of three replicates (n=15-18). **(C)** Whole-plant images of eight-

week-old plants of the indicated genotypes. Bar = 0.5 cm. **(D)** Colorimetric quantification of NBT-

formazan deposition on rosette leaves. FW, fresh weight. Data are means ± SD of three replicates

(n=18). **(E)** Hydrogen peroxide (H_2O_2) content, quantified as the H_2O_2 (mmol) per gram dry

weight (DW) of rosette leaves using Amplex Red-coupled fluorescence quantitative assay. Values

are means ± SD of three replicates (n=14). **(F)** Quantification of ectopic cell death detected by

Trypan blue staining. Data are means ± SD of three replicates each one consisting of 7 to 10 leaves.

(G) Semi-quantitative RT-PCR analysis of *MSL10* from lines expressing *MSL10g* transgene or its

phospho-variants and *msl10-3G* lesion. cDNA was synthesized from total RNA extracted from

three-week-old plants of the indicated genotypes. *EF1α* was used as a loading control.

1025

Supplemental Figure 7. Additional controls for the BiFC assay. (A) The C-terminal half of

MSL10 forms aggregates when fused to GFP. Representative confocal images of abaxial *N.*

benthamiana leaf epidermis from plants transiently expressing truncated MSL10$_{461-734}$-GFP and

MSL10 $^{S640L}_{461-734}$-GFP. Images were taken 5 d post-infiltration. Scale bar, 50 μm. **(B)** *N.*

benthamiana leaves co-infiltrated with constructs fused to the split YFP were observed under the

confocal microscope 3-5 days after infiltration. Scale bar, 50 μm. (**Top row**) Neither domain

interacts with KAT1. (**Middle row**) Neither phosphorylation status of the N-terminus nor the

S640L lesion in the C-terminus affect the interaction. either reconstituted YFP fluorescence signal

from positive interaction or marked reduction in YFP signal in case of negative interaction from

indicated combinations. (Bottom row) Negative controls for the N-terminus with the S640L lesion.

Scale bar, 50 μm.

1037

Supplemental Figure 8. Analysis of reliability of RNA-seq samples using multi-dimensional

scaling plot (MDS) and correlation between RNA-seq and qRT-PCR derived expression

data. **(A)** MDS plot indicates biological variation across treatments and genotypes. MDS plot was

generated using edgeR (Robinson et al., 2010). FC, Fold change; dim, dimension. **(B)** Table

showing the list of hallmark genes associated with *MSL10* gain-of-function lines that are also

identified in the RNA-seq analysis.

1044

1045   **Supplemental Table 1.** mRNA sequencing and alignment statistics

1046

1047   **Supplemental Table 2. List of differentially expressed genes identified in the transcriptomic**

1048   **analysis belonging to Cluster 2.** Marked in yellow are the two columns corresponding to the two

1049   replicates of *35S:MSL10-GFP* overexpression lines grown at 21°C.

1050

1051   **Supplemental Table 3. Tabulated list of overrepresented GO terms (biological process) and**

1052   **corresponding P values from the enrichment analysis performed using the PANTHER**

1053   **classification tool**

1054

1055   **Supplemental Table 4. List of selected genes with threshold value >10 and significantly**

1056   **enriched in at least two overrepresented GO categories.** Marked in yellow are the two columns

1057   corresponding to the two replicates of *35S:MSL10-GFP* overexpression lines grown at 21°C.

1058   Marked in green are the genes selected for qRT-PCR in this study.

1059

1060   **Supplemental Table 5**. List of primers used in this study.