The influence of substrate concentration on the culturability of heterotrophic soil microbes

isolated by high-throughput dilution-to-extinction cultivation

Authors: Ryan P. Bartelme[1], Joy M. Custer[1], Christopher L. Dupont[2], Josh L. Espinoza[2],

Manolito Torralba[2], Banafshe Khalili[3], Paul Carini[1]*

Affiliations:

[1]Department of Environmental Science, University of Arizona, Tucson, AZ 85721

[2] Department of Environment and Sustainability, J. Craig Venter Institute, La Jolla, CA 92037

[3]Department of Ecology and Evolutionary Biology, University of California, Irvine, CA 92697

*Corresponding author: paulcarini@email.arizona.edu

Abstract

The vast majority of microbes inhabiting oligotrophic shallow subsurface soil environments have

not been isolated or studied under controlled laboratory conditions. In part, the challenges

associated with isolating shallow subsurface microbes may persist because microbes in deeper

soils are adapted to low nutrient availability or quality. Here we use high-throughput dilution-to-

extinction culturing to isolate shallow subsurface microbes from a conifer forest in Arizona,

USA. We hypothesized that the concentration of heterotrophic substrates in microbiological

growth medium would affect which microbial taxa were culturable from these soils. To test this,

we diluted extracted cells into one of two custom-designed defined growth media that differed

only by a 100-fold difference in the concentration of amino acids and organic carbon. Across

both media, we isolated a total of 133 pure cultures, all of which were classified as

Actinobacteria or Alphaproteobacteria. The substrate availability dictated which actinobacterial

phylotypes were culturable but had no significant effect on the culturability of

Alphaproteobacteria. We isolated cultures that were representative of the most abundant

phylotype in the soil microbial community (*Bradyrhizobium* spp.) and representatives of five of

the top 10 most abundant *Actinobacteria* phylotypes, including *Nocardioides* spp.,

Mycobacterium spp., and several other phylogenetically-divergent lineages. Flow cytometry of

nucleic acid-stained cells showed that cultures isolated on low-substrate medium had

significantly lower nucleic-acid fluorescence than those isolated on high-substrate medium.

These results show that dilution-to-extinction is an effective method to isolate abundant soil

microbes and the concentration of substrates in culture medium influences the culturability of

specific microbial lineages.

Importance

Isolating environmental microbes and studying their physiology under controlled conditions is an

essential aspect of understanding their ecology. Subsurface ecosystems are typically nutrient-

poor environments that harbor diverse microbial communities—the majority of which are thus

far uncultured. In this study, we use modified high-throughput cultivation methods to isolate

subsurface soil microbes. We show that a component of whether a microbe is culturable from

subsurface soils is the concentration of growth substrates in the culture medium. Our results offer

new insight into technical approaches and growth medium design that can be used to access the

uncultured diversity of soil microbes.

Introduction:

Soil microbial communities are tremendously diverse and mediate crucial aspects of plant fertility, biogeochemistry, pollutant mitigation, and carbon sequestration (1–4). While the diversity and community composition of surface soils have been relatively well described, we know far less about the microbes inhabiting deeper soils (defined here as >10 cm below the surface), despite their key role in soil formation and mineralization of key plant nutrients. In contrast to surface soils that are typically rich in plant-derived compounds, subsurface soils are often characterized by lower amounts of mineralizable nitrogen, phosphorus and organic carbon—much of which has a long residence time and is relatively recalcitrant to microbial degradation (5–9). The temperature and soil moisture of subsurface soils are also less variable than shallower soils that are exposed to seasonal changes in temperature and precipitation (10). These relatively stable and low-nutrient conditions found at depth constrain both the amount of microbial biomass present in the subsurface and the structure of these microbial communities (11–14). Many of the microbial taxa that are abundant in these subsurface environments are underrepresented in microbial culture and genome databases (11). Thus, there are large knowledge gaps in our understanding of the biology of a major fraction of subsurface soil microbes.

Because subsurface soils are low-nutrient habitats, part of the challenge associated with culturing and studying the microbes that live belowground may be that they require low nutrient concentrations in order to be isolated or propagated in the laboratory (15). These microbes— often referred to as 'oligotrophs'—are capable of growing in conditions where the supply or quality of nutrition is poor. Although oligotrophs dominate most free-living microbial

ecosystems (16), the concept of oligotrophy itself is enigmatic. There is no coherent definition of

what constitutes oligotrophic metabolism aside from their ability to grow at 'low' nutrient

concentrations—a definition that itself is arbitrary (17, 18). Kuznetsov *et al*. (19), identified three

groups of cultivatable oligotrophs: 1) microbes that can be isolated on nutrient-poor medium, but

cannot be subsequently propagated; 2) microbes that can be isolated on nutrient-poor medium

but can be subsequently propagated on nutrient-rich medium; and 3) microbes that require

special nutrient-poor medium for both isolation and propagation. Although the molecular and

genetic mechanisms that distinguish these three categories are poorly understood, several traits

of oligotrophs have emerged from the study of microbes that numerically dominate oligotrophic

ecosystems. For example, oligotrophs are typically small, slowly growing cells (20–23). The

genome sizes of numerous lineages of microbes that dominate oligotrophic marine ecosystems

tend to be highly reduced—an indication that microbial oligotrophy may be tied to reduction of

genome size (24–26). These 'streamlined' genomes often code for fewer copies of the rRNA

gene operon and transcriptional regulator genes than microbes with larger genomes, suggesting

oligotrophs lack the ability to sense and rapidly respond to varying environmental conditions (12,

16). Instead, genomic inventories of marine oligotrophs suggest a reliance on broad-specificity,

high-affinity transporters that are constitutively expressed (22, 26, 28, 29).

While the activities of abundant and ubiquitous microbes that inhabit oligotrophic marine

environments have been extensively investigated in recent years (24, 30, 31), far fewer studies

have focused on the activities of microbes that dominate oligotrophic soil environments. Several

soil studies used low-throughput techniques to show that reduced-nutrient solid media facilitated

the isolation important soil microbes that were previously uncultured (32–35). While several

agar-based high-throughput approaches have been used to isolate diverse microbes (36, 37),

these approaches may not be appropriate to isolate microbes that thrive at micromolar amounts

of growth substrate and do not form detectable colonies on solid media. Here, we adapt existing

high-throughput dilution-to extinction protocols, originally developed for isolating abundant

aquatic oligotrophic bacteria, to facilitate the isolation of soil microbes. We hypothesized that the

concentration of heterotrophic growth substrates in a growth medium would constrain which taxa

were able to be isolated on a custom-designed defined artificial medium. We tested this by

extracting cells from oligotrophic subsurface soils using Nycodenz buoyant density

centrifugation (38), and inoculating high-throughput dilution-to-extinction experiments in two

defined media that contained a 100-fold difference in the amounts of heterotrophic growth

substrates. We isolated several bacteria that were representative of abundant phylotypes found in

the original soil microbial community and two lineages representative of uncultured groups of

microbes. In these experiments, the substrate concentration significantly influenced which

actinobacterial genera were culturable in the laboratory but no effect on which

alphaproteobacterial lineages were culturable. Moreover, we show that cells isolated on low

nutrient medium had significantly lower SYBR green I nucleic acid fluorescence, suggesting

microbes isolated on low nutrient medium may contain reduced nucleic acid content relative to

those isolated on higher nutrient medium.

Results:

We collected shallow subsurface soil (55 cm) from the Oracle Ridge field site in a mid-

elevation conifer forest that is part of the Santa Catalina Mountains Critical Zone Observatory in

Arizona, USA. These soils contained very low amounts of total organic carbon (0.095%) and N-

NO_3 (0.3 ppm) indicating they were highly oligotrophic (Supplementary Fig. 1). We adapted

existing high-throughput dilution-to-extinction approaches designed for aquatic microbes (39,

40) to culture soil microbes from these samples (Fig. 1). The primary modification to existing

protocols was to add a buoyant density centrifugation cell-separation step to detach inoculum

cells from mineral soils prior to diluting cells into growth medium. To do this, we vortexed soil

in a cell extraction buffer containing a nonionic surfactant and a dispersing agent. We layered

this soil-buffer slurry over 80% Nycodenz and centrifuged it. During centrifugation, the mineral

components of soil migrated through the Nycodenz, while cells 'floated' on the surface of the

Nycodenz. We extracted cells located at the Nycodenz interface, stained them with SYBR Green

I, and counted them on a flow cytometer. This extraction yielded $1.28 \times 10_5$ cells ml$_{-1}$ from 0.5 g

wet soil. We diluted the extracted cells to an average of 5 cells well$_{-1}$ in deep-well

polytetrafluoroethylene 96-well plates containing a custom-designed and defined growth

medium that we named 'Artificial Subterranean Medium' (ASM), with low or high

concentrations of heterotrophic growth substrates (ASM-low and ASM-high, respectively) (Fig.

1). The ASM-low and ASM-high media contained identical inorganic mineral and vitamin

amendments but a 100-fold difference in the concentration of organic carbon and amino acids

(Supplementary Table 1). We designed these media to facilitate the growth of diverse

chemoheterotrophic microbes by including an array of simple carbon compounds, polymeric

carbon substrates, and individual amino acids (Supplementary Table 1). We prepared triplicate

96-well plates for each growth medium formulation. These dilution-to-extinction experiments

were screened for growth with flow cytometry after 4 weeks of incubation and again after 11

weeks of incubation (Fig. 1). Wells displaying growth (defined as those wells displaying $1.0 \times$

10_4 cells ml$_{-1}$) were sub-cultured into larger volumes and subsequently cryo-preserved and

identified by 16S rRNA gene sequencing (Fig. 1).

Across both medium types, a total of 214 wells (119 for ASM-low and 95 for ASM-high)

displayed growth after 11 weeks of incubation. We successfully propagated 182 (85%) of the

cultures from microtiter plates to polycarbonate flasks containing fresh medium. Of the cultures

that successfully propagated, we confirmed that 73% (133 cultures) were pure cultures by

amplifying and sequencing full-length 16S rRNA gene sequences from genomic DNA

extractions. The remaining 49 cultures were mixed (Foreword and Reverse 16S rRNA sequence

reads did not assemble due to base ambiguities) or, in rare instances, did not amplify under

several amplification conditions. We defined microbial culturability using Button's definition of

microbial 'viability' as determined in dilution-to-extinction experiments (41). Here,

'culturability' is defined as the ratio of cells that grew into detectable cultures to the total number

of cells initially diluted into a cultivation chamber (41). By the end of the experiment, we

approached ~20% culturability across both medium formulations (Fig. 2). In general, microbial

culturability was higher for ASM-low than for ASM-high, but this effect was only significant

after 4 weeks of incubation (Fig. 2; Wilcoxon rank-sum test $P \leq 0.05$ at 4 weeks).

We assigned taxonomy to each 16S rRNA gene sequence using the SILVA database. All

pure cultures isolated on ASM-low and ASM-high belonged to one of two bacterial phyla:

Actinobacteria (110 cultures; 83% of the pure cultures) or Proteobacteria (23 cultures, all

Alphaproteobacteria; 17% of the pure cultures) (Fig. 3). Across all experiments, the genera

assigned to bacteria isolated on ASM-low were significantly distinct from those isolated on

ASM-high (Kruskal-Wallis rank sum, χ_2=19.05, P=1.28 × 10$_{-5}$). However, these differences were

largely driven by significant differences in culturability across medium types for the

Actinobacteria, but not for the Alphaproteobacteria (Dunn test, $P \leq 0.000$ for Actinobacteria and

P=0.128 for Alphaproteobacteria).

Just over half of the Alphaproteobacteria (57%) were isolated on ASM-low medium and

the remaining 43% on ASM-high (Fig. 3 and Supplementary Fig. 2). Cultures that classified as

Bradyrhizobium spp. were the most frequent alphaproteobacterial isolates (13 isolates), seven of

which were isolated on ASM-low medium. Cultures classified as *Reyranella* spp. and *Nordella*

spp. were also isolated on both ASM-high and ASM-low medium. Of the remaining five

Proteobacteria cultures, three were isolated on ASM-low (*Afipia* (1 culture), *Rhizobiales* (1

culture), *Bauldia* (1 culture)), and two were isolated on ASM-high (*Pseudolabrys* (1 culture) and

a *Xanthobacteraceae* sp. (1 culture)). The actinobacterial cultures belonged to three classes:

Actinobacteria (107 cultures), Thermoleophilia (2 cultures), and Acidimicrobiia (1 culture). Of

these Actinobacteria, 65 (59%) were isolated on ASM-low, and 45 (41%) on ASM-high. The

cultures were numerically dominated by two genera that were differentially isolated on ASM-

low and ASM-high: *Nocardioides* and *Mycobacterium*. *Nocardioides* spp. (46 cultures) were

exclusively isolated on ASM-low medium (Fig. 3 and Supplementary Fig. 3). Other cultures that

were isolated on ASM-low included those classified as *Arthrobacter* (3 cultures), *Marmoricola*

(2 cultures), *Nakamurella* (2 cultures), *Aeromicrobium* (1 culture), *Blastococcus* (1 culture), and

Patulibacter (1 culture) (Fig. 3 and Supplementary Fig 3). While the majority of cultures

classified as *Mycobacterium* sp. were isolated on ASM-high (38 cultures), we isolated seven

mycobacterial cultures on ASM-low medium—five of which form a phylogenetically-distinct

cluster from those isolated on ASM-high (Fig. 3 and Supplementary Fig. 3). Other

actinobacterial cultures isolated on ASM-high included *Jatrophihabitans* (4 cultures),

Conexibacter (1 culture), and *Amycolatopsis* (1 culture).

Interestingly, we isolated what are likely the first members of two novel actinobacterial

lineages on ASM-low. The first such culture—*Microtrichales* sp. str. AZCC_0197—belongs to

the *Microtrichales* order of the Acidimicrobiia class. The best 16S rRNA gene sequence match

to an existing isolate is 93.4% identity to *Aquihabitans daechunggensis* str. G128. However,

strain AZCC_0197 more closely matched numerous 16S rRNA gene sequences from

environmental clones of uncultured Acidimicrobiia. The second lineage—*Frankiales* sp. strains

AZCC_0102 and AZCC_0072—classified as members of the *Frankiales* order of the

Actinobacteria class with best matches of <97% nucleotide identity to existing *Frankiales*

isolates (42).

Several of the microbes we isolated were representative of abundant members of the

subsurface soil microbial community at the Oracle Ridge site. We matched the 16S rRNA gene

sequences from our cultures to the phylotypes derived from the 55 cm Oracle Ridge soil sample.

The 16S rRNA gene sequences from our cultures matched 13 phylotypes (at 97% identity; Fig.

4) that account for $11.0 \pm 1.6\%$ (mean $\pm$ SD, n=3) of the total amplifiable microbial community.

For example, the 16S rRNA gene sequences from our *Bradyrhizobium* isolates match a single

Bradyrhizobium phylotype that was the most abundant phylotype at 55 cm (relative abundance

of $5.7 \pm 0.3\%$ (mean $\pm$ SD, n=3); Fig. 4). Additionally, we isolated representatives of abundant

Actinobacteria (Fig. 4) including: two *Mycobacterium* phylotypes (the 11th and 17th most

abundant phylotypes overall); *Nocardioides* (the 13th most abundant phylotype overall); and two

Arthrobacter phylotypes (16th and 1,271st most abundant phylotypes overall). The other

Actinobacteria cultured in these experiments represent rarer phylotypes in bulk soils. The 16S

rRNA gene sequences from several of our pure cultures did not match any of the phylotypes

derived from these soils at ≥97% identity, including *Nakamurella* (2 cultures)*, Nocardioides* (5

cultures), *Mycobacterium* (1 culture), *Jatrophihabitans* (1 culture), *Patulibacter* (1 culture),

Conexibacter (1 culture), *Rhizobiales* sp. (1 culture), *Reyranella* (1 culture), and *Microtrichales*

sp. str. AZCC_0197.

In many environments, relative nucleic acid content can be estimated with flow

cytometry analysis of cells stained with nucleic acid staining dyes (43–45). Given that many

microbes inhabiting low-nutrient environments exhibit reduced genome sizes, we sought to

determine whether the nucleic acid fluorescence measured for our cultures partitioned by the

growth medium they were isolated on. For each culture, we identified the closest match to an

available genome sequence and found that the genome length of these 'best hit' matches was

significantly correlated with the average fluorescence of SYBR Green I-stained cells

(Spearman's rho=0.3 P=0.0012), indicating that the nucleic acid fluorescence we quantified by

flow cytometry was indeed indicative of genome size differences. We found that cultures

isolated on ASM-low exhibited significantly lower mean nucleic acid fluorescence than those

isolated on ASM-high (Fig. 5; Kruskal-Wallis Rank sum χ_2=24.8 P=6.27 $\times$ 10$_{-7}$). The overall

mean fluorescence was not significantly different across the phyla assigned to each isolate

(Kruskal-Wallis Rank sum χ_2=0.210, P=0.647), but was significant across individual genera

assignments (Kruskal-Wallis Rank sum χ_2=62.4 P=2.98 $\times$10$_{-6}$). Moreover, the mean nucleic acid

fluorescence values within a given genera were similar (Supplementary Fig. 4). For example,

Mycobacterium isolates had relatively high nucleic acid fluorescence, regardless of which

medium they were isolated on (Supplementary Fig. 4). In contrast, *Nocardioides* (ASM-low) and

Jatrophihabitans (ASM-high) displayed relatively low nucleic acid fluorescence. Interestingly,

when cultured on ASM-high, we observed a clear nucleic acid fluorescence dichotomy across the

Bradyrhizobium isolates (Supplementary Fig. 4).

Discussion:

We designed a proof-of-concept workflow to determine the feasibility of high-throughput

dilution-to-extinction cultivation for the isolation of soil microbes. The method was based on a

workflow for isolating microbes from oligotrophic marine environments (39). However, unlike

aquatic samples, microbial cells in soils are heterogeneously dispersed within, or attached to, a

complex matrix comprised of non-cellular organic matter and minerals. The complexity of this

soil matrix complicates accurate enumeration of viable cells because mineral and organic matter

can interfere with flow cytometry. To circumvent these issues, we separated cells by gently

shaking soils in a cell extraction buffer containing a dispersing agent and a nonionic surfactant.

Cells were separated from this slurry by buoyant density centrifugation (Fig. 1). This procedure

allowed cells to be floated on top of a dense solution of Nycodenz while allowing minerals to

migrate through the Nycodenz solution (46).

We estimated the expected culturability and calculated the actual culturability using the

statistical framework of dilution culture growth outcomes described by Button, *et al.* (41). The

expected number of pure cultures ($\hat{u}$) was estimated across all experiments using the formula $\hat{u} =$

$-n(1-p) \times \ln(1-p)$. Where p is the proportion of wells displaying growth (214 growth chambers

displaying growth ÷ 576 chambers inoculated = 0.37) and n is the number of inoculated

chambers (576 chambers in total). Based on this equation, we expected $\hat{u} = 168$ pure cultures

across all experiments. The number of pure cultures we obtained (133 cultures) was within 21%

of this value. However, this result is conservative because it does not account for the cultures that

were initially scored as positive for growth but did not successfully subculture. Some of these

cultures may be oligotrophic taxa that were initially cultivatable but failed to successfully

propagate, as described by Kuznetsov et al. (19). Alternatively, cultures that failed to propagate

from microtiter plates to larger volumes might have been false positives, where flow cytometer

instrument noise or well-to-well carryover was mistaken for a low-density culture. The mean

culturability we observed for a given experiment (1.4%-11%; Fig. 2) was comparable to dilution-

to-extinction cultivation studies of marine microbes, which report 0.5%-14.3% culturability (40).

Similar to previous observations for soil microbes (34), we observed increased culturability with

longer incubation times (Fig. 2). We speculate that the culturability was higher on ASM-low than

on ASM-high because the cells were extracted from an extremely oligotrophic soil sample

(Supplementary Fig. 1) and perhaps the results would have been different if the original

inoculum originated from more productive soils.

The concentration of heterotrophic growth substrates in the isolation medium significantly

influenced our ability to isolate certain actinobacterial lineages (Fig. 3). In particular,

Nocardioides were exclusively isolated on ASM-low and most *Mycobacterium* isolates were

isolated on ASM-high. We designed our media to include a defined but diverse range of carbon

substrates that have been successfully used to isolate chemoheterotrophic microbes from soil or

oligotrophic taxa from other environments (34, 50). Carbon type and availability are crucial for

heterotrophic soil microbes because carbon acts as both a source of electrons for respiration and

carbon for biomass. To accommodate this requirement, many common microbial growth

medium formulations for heterotrophic microbes supply diverse growth substrates (yeast extract

or casein digests, for example), usually at concentrations much higher than are normally

available *in situ*. Two key assumptions made with these common media formulations are that: 1)

microbes will use only the relevant constituents and any remaining compounds will have minor

or no effect on microbial growth; and 2) microbes grow optimally in the laboratory when

nutrient availability is much greater than their half saturation constant (47). While many

commonly studied microbes have the capacity to grow on complex, high-nutrient formulations,

environmental nucleic acid data informs us that the vast majority of Earth's microbes remain

uncultured (48, 49). Our results indicate that the concentration of nutrients in a growth medium

may be as important as the constituents of the growth medium for cultivation of uncultured

environmental microbes.

Numerous studies have demonstrated that dilute growth media is superior to substrate

rich growth media for the isolation of novel soil microbes (34, 35, 51, 52). However, the

physiological explanation as to why low-nutrient media facilitates the growth of diverse

microbes, or high nutrient concentrations inhibit the growth of some taxa, remain unclear. One

possible explanation for these concentration-dependent effects may be that growth medium

formulations applied at high concentrations contain high amounts of inhibitory substances—

substances that are reduced to non-inhibitory levels in dilute medium formulations. For example,

a key amino acid transporter in *Chlamydia trachomatis* can be blocked by non-essential amino

acids, preventing the transport of required amino acids, resulting in growth inhibition (53). A

similar phenomenon was demonstrated in the extreme marine oligotroph '*Candidatus*

Pelagibacter ubique', where alanine was conditionally required for cell division but abolished

growth at higher concentrations (54). Furthermore, reactive oxygen species can be produced

during the autoclaving of nutrient-rich medium that either directly inhibit growth or combine

with organics in the medium to form inhibitory compounds (55, 56). Finally, growth inhibition

may be the result of misbalanced regulation of growth or accumulation of nutrient storage

structures (poly-β-hydroxybutyrate, for example) ultimately leading to cell lysis (57). A better

understanding of the mechanism(s) that enable growth on low nutrient medium—or prevent

growth on high-nutrient medium—may help us design better strategies for isolating uncultivated

lineages. Critically, the collection of cultures we describe here, that were isolated on medium

with identical constituents applied at different concentrations, is a first step towards an

experimental method capable of addressing these questions.

While several of the taxa we isolated were abundant microbial members of the shallow

subsurface microbial community (Fig. 4), other isolates were rare or not identified in the

cultivation-independent soil microbial community. The cultivation of additional microbial

phylotypes that were not observed in molecular analyses of the same samples has been observed

(58, 59). The dilution-to-extinction approach we used here favors the cultivation of abundant

microbes in a given sample (41), such that the isolation of rarer taxa or taxa that were not

observed in the original sample was unexpected. There are several possible explanations for this

observation. First, the buoyant density separation protocol that we used to separate cells may

have introduced biases. Nycodenz cell separation approaches do not recover all microbial cells

from soil and the separable fraction can be compositionally distinct from the non-separated soils

(61). These effects could potentially skew the proportions of microbes that were diluted into

microtiter plate wells. Alternatively, the absence of a particular taxon in a soil microbiome

analysis may be the result of insufficient sequencing depth (58). Moreover, the 'universal'

primers used in the soil microbiome analysis (60) may not have primed DNA from some of the

divergent lineages we cultured as efficiently as other phylotypes in the soil microbiome, resulting

in either underrepresentation of these phylotypes in the original community or no amplification

at all. We do not have sufficient evidence indicating which of these scenarios may explain our

ability to culture cells that were not apparent in the microbiome analysis. Finally, as is true in

any microbial cultivation experiment, there were many taxa that we did not isolate. In particular,

these soils contained high relative abundances of Verrucomicrobia related to '*Candidatus*

Udaeobacter copiosus' (62), and Acidobacteria (Subgroup 6), which belong to highly sought-

after lineages of uncultured microbes (63). The reasons for not isolating these (and other)

lineages are numerous but may be the result of inappropriate medium composition (64, 65), toxic

compounds in the cell separation constituents, long doubling times (> ~6 days), or dormancy

(reviewed in (66)).

We provide evidence that microbes cultured from oligotrophic soils on low-nutrient medium

may have reduced nucleic acid content relative to microbes isolated on richer medium (Fig. 5).

Depending on the taxa in question, and their effective population size, microbial genome

reduction can be driven by either genetic drift or 'streamlining' selection (reviewed in (67)).

Genome streamlining is strongly linked with microbial oligotrophy in free-living aquatic

microbes as a mechanism to reduce the overhead cost of replication in periodically nutrient-

limited environments (reviewed in (21)). However, direct evidence for genome streamlining in

terrestrial microbes has been elusive. For example, metagenome-assembled genomes of abundant

and ubiquitous uncultured Verrucomicrobia suggest that some lineages may contain reduced

genome sizes (62). A more recent study showed that fire-affected warm soils selected for groups

of microbes with significantly smaller genomes than cooler soils (68). Yet, there are few

definitive ways to identify the growth preferences of taxa with reduced genomes short of

culturing them and studying their growth dynamics under controlled conditions. The appearance

of reduced nucleic acid content in cultures isolated on ASM-low may be an indication that

genome reduction may be a successful life strategy for soil oligotrophs. Alternative explanations

for the apparent differences in nucleic acid content in microbes cultured in ASM-low may be

that: 1) the ploidy of stationary-phase cells grown in ASM-low may be lower than those isolated

in ASM-high; 2) unknown cellular constituents of cells grown in ASM-low may quench SYBR

Green I fluorescence in the assay conditions we used; or 3) cells isolated in ASM-high may form

small microaggregates that are not completely dispersed prior to flow cytometry.

The development of cultivation techniques emphasizing the high-throughput and sensitive

detection of microbial growth on low-nutrient medium revolutionized the field of aquatic

microbial ecology by culturing microbes that were previously 'unculturable' using standard

techniques (39, 40, 50, 69, 70). Here we show that similar cultivation principles facilitate the

cultivation of abundant soil microbes. We demonstrate that, in addition to scrutinizing the

nutritional composition of a given growth medium, the concentration of growth substrates in the

growth medium must also be considered. Although we do not yet understand the mechanism of

substrate-induced growth inhibition, there is evidence that this phenomenon is widespread and

may impede laboratory cultivation efforts. Future studies to deduce the molecular mechanisms of

substrate-induced growth inhibition will likely lead to new cultivation approaches that will allow

us to isolate abundant free-living oligotrophic microbes.

Materials and Methods:

Soil source and nutrient analysis: Fresh soil samples were collected from a soil pit on

August 16, 2017 at the Oracle Ridge site in the Catalina Jemez Critical Zone Observatory

(coordinates: 32.45 N, -110.74 W, elevation 2,103 m). The mean annual precipitation at Oracle

Ridge is 87 cm year-1 and the mean annual surface temperature is 12ºC (71). The soils were

Typic Ustorthent (71). The dominant vegetation at the site is ponderosa pine (*Pinus ponderosa*),

with sparse Douglas-fir (*Pseudotsuga menziesii*). We collected ~300 g subsamples from 0-5 cm,

10 cm, 20 cm, 30 cm, 40 cm and 55 cm depths. Soils were kept cool with ice packs for <4 h

while in transit to the laboratory. At the laboratory, the soils were sieved to 2 mm and kept at 4ºC

for 50 days at which point cells were separated from mineral soil. Standard soil chemical

analyses were performed at the Colorado State University Soil Water and Plant Testing

Laboratory using their standard protocols. We analyzed the microbial community composition at

each depth (see *'Soil microbial community analysis,'* below) and conducted cultivation

experiments from the 55 cm soil sample.

Soil microbial community analysis: We extracted DNA from 1.0 g subsamples (n=3) using

MoBio PowerSoil DNA extraction kits using the manufacturer's instructions. We amplified 16S

rRNA gene fragments using 515F-Y (5'-TATGGTAATTGTGTGYCAGCMGCCGCGGTAA-

3') and 926R (5'-AGTCAGTCAGGGCCGYCAATTCMTTTRAGT-3') (72). PCR products

were purified using the QIAquick PCR purification kit (Qiagen, Germantown, MD) per

manufacturer's specification. Cleaned products were quantified using Tecan fluorometric

methods (Tecan Group, Mannedorf, Switzerland), normalized, and pooled for Illumina MiSeq

sequencing using custom sequencing primers and the MiSeq Reagent v2 500 cycle Kit (Roche,

Branford, CT) following the manufacturer's protocols. We identified phylotypes based on the

generation of *de novo* operational taxonomic units (OTUs) from raw Illumina sequence reads

using the UPARSE pipeline at a stringency of 97% identity (73). Paired-end reads were trimmed

of adapter sequences, barcodes and primers prior to assembly. We discarded low-quality and

singleton sequences and dereplicated the remaining sequences before calculating relative

abundances. Chimera filtering of the sequences was completed during clustering while taxonomy

was assigned to the OTUs with mothur (74) using version 123 of the SILVA 16S ribosomal

RNA database (75) as the reference. We generated OTU and taxonomy assignment tables for

subsequent analyses.

Cell separation: Cells were separated from sieved soils using buoyant density centrifugation

with Nycodenz modified from (38) to isolate viable cells. Briefly, we added 0.5 g wet soil to

44.8 ml of cell extraction buffer (137.5 mM NaCl, 26.78 mM tetrasodium pyrophosphate, and

0.27% (v/v) Tween 80). The soil-buffer slurry was vortexed for 30 seconds and shaken

horizontally on a platform shaker for 2 hours at 4°C. We layered 15 ml aliquots of this soil-

buffer slurry over 10.0 ml of 80% (w/v) Nycodenz solution in 50 mM tetrasodium

pyrophosphate. We used 50 ml Nalgene Oak Ridge high-speed polycarbonate centrifuge tubes

for buoyant density centrifugations. Tubes containing the soil-buffer solution with Nycodenz

were centrifuged at $17,000 \times g$ for 30 minutes at 16°C. We extracted 3×0.5 mL aliquots from

the resulting buoyant density preparation at a location of ~25 mm above the bottom of the tube

(coincident with the approximate level of the top of the Nycodenz solution) to sterile

microcentrifuge tubes containing 1.0 ml 137.5 mM NaCl. The microcentrifuge tubes containing

Nycodenz/NaCl were vortexed and centrifuged for 20 min at $17,000 \times g$. The resulting cell

pellets were resuspended in 137.5 mM NaCl, pooled and stored at 4°C.

Medium design rationale: The ASM media were custom-designed to facilitate the growth of

a broad range of soil chemoheterotrophic microbes (Supplementary Table 1). Both ASM-high

and ASM-low were buffered with phosphate. To this, we added minerals at concentrations

derived from an 'artificial rainwater' recipe (76), trace elements as described in trace element

solution SL-10 with the addition of $LaCl_3$, and vitamins as described elsewhere (54). We added

heterotrophic growth substrates that included 21 amino acids and a diverse range of simple

carbon substrates including 2-C substrates (glycerol, acetate), 3-C substrates (pyruvate), 4-C

substrates (succinate, butyrate, isobutyrate), 5-C substrates (ribose, valerate), a 6-C substrate

(glucose), an 8-C substrate (NAG) and a 10-C substrate (decanoic acid) (Supplementary Table

1). We also added several polymeric growth substrates including pectin, methylcellulose,

alginate, starch, and xylan (Supplementary Table 1). We calculated the added carbon amount to

be ~200 mg C L_{-1} for ASM-high and ~2 mg C L_{-1} for ASM-low.

Dilution-to-extinction: An aliquot of cells extracted from the buoyant density separation were

fixed with 1.75% (final v/v) formaldehyde and stained with SYBR Green I (final stain

concentration was a 1:4,000 dilution of commercial stock) for 3.5 h at room temperature in the

dark. Cells were enumerated using a Millipore Guava flow cytometer, as described elsewhere

(54). We diluted cells into Artificial Subterranean Medium (ASM)-high or ASM-low nutrient

medium (Supplementary Table 1) to a density of 5 cells mL_{-1} and aliquoted 1.0 ml of the dilute

cell suspension into the wells of 2 ml polytetrafluoroethylene 96-well microtiter plates (Cowie

Technology, New Castle, DE) so that on average each well contained 5 cells. Plates were

covered with plastic lids that allow air circulation and incubated at 16°C in the dark in aerobic

conditions. We screened the dilution-to-extinction plates for growth by fixing (1.75%

formaldehyde) and staining (1:4,000 dilution of commercial SYBR Green I stock) aliquots for

18 h in the dark at room temperature and counting by flow cytometry (EMD-Millipore Guava

EasyCyte), as described previously (54). We screened plates for growth at 4 and 11 weeks after

inoculation. Positive cultures were defined as cultures that exceeded $1.0 \times 10_4$ cells ml_{-1}.

Actual and theoretical culturability estimates: Culturability estimates were determined by the

equation: $V = -\ln(1-p)/X$, where *V* is the estimated culturability, *p* is the proportion of inoculated

cultivation chambers that displayed measurable growth (number of chambers positive for growth

÷ total number of chambers inoculated), and *X* is the number of cells added to each cultivation

chamber as estimated from dilutions (41). The number of pure cultures (*û*) was estimated as

follows: $\hat{u} = -n(1-p) \times ln(1-p)$, where n is the number of inoculated growth chambers and p is the

proportion of inoculated wells displaying growth (41).

Culture transfer and storage: We subcultured positive growth chambers into 25 ml of the

respective growth medium (ASM-high or ASM-low) in acid-washed, sterile polycarbonate flasks

and incubated them at 16ºC. At the time of transfer, we assigned cultures a unique Arizona

Culture Collection (AZCC) number. Flasks were monitored for growth every other week for 2

months. Flasks displaying growth within two months were cryopreserved in 10% glycerol and

stored at -80ºC. If no growth appeared within two months, the cultures were discarded and the

assigned AZCC number was retired.

Mean fluorescence calculations: We calculated the mean fluorescence of each culture from

the subcultures grown in 25 ml volumes at 12-15 weeks after inoculating. Culture aliquots were

fixed and stained for 15-18 h as described above in 'Dilution-to-extinction.' We manually gated

histograms of the intensity of SYBR Green I fluorescence (in arbitrary units) and extracted the

mean fluorescence of the gated peak for each culture using the GuavaSoft software package.

'Best hit' genomes were determined by blasting the full length 16S rRNA gene sequence of our

isolates against the NCBI Microbial Genomes database using web-blast (77). We extracted the

total genome length from each best-hit genome.

Culture identification: Cultures were identified by full length 16S rRNA gene sequencing.

Briefly, we filtered 5-10 ml of cell biomass from 25 mL cultures on to 0.2 µm-pore size Supor

filters and extracted DNA using a Qiagen PowerSoil DNA extraction kit following the

manufacturer's instructions. We amplified full-length 16S rRNA genes from the resulting DNA

using the 27F-1492R primer set (27F: 5'-AGAGTTTGATCMTGGCTCAG-3'; 1492R: 5'-

ACCTTGTTACGACTT-3' (78)). The reaction mix consisted of Promega's GoTaq HotStart 2x

PCR master mix with final concentrations of 0.4 µM 27F and 0.4µM 1492R primers, and 1-11.5

µl of template DNA, in a total reaction volume of 25 µL. The thermocycling profile was $1 \times$

94°C for 10 min followed by 36 cycles of: 94°C for 45 s, 50°C for 90 s, 72°C for 90 s; and a

single 72°C extension for 10 min. The resulting amplicons were cleaned and Sanger sequenced

from both the 27F and 1492R primer by Eurofins Genomics (Louisville, KY, USA) using their

standard techniques. Sequences were curated using 4Peaks (79) and Geneious Prime v2019.0.1

(80). Reads were trimmed and assembled using the moderate setting in Geneious. Forward and

reverse Sanger PCR reads that failed to build a full length 16S rRNA gene with these metrics

were considered "mixed" cultures and not analyzed further.

Culture taxonomy and determination of taxonomic differences across growth medium

formulations: High-quality full length 16S rRNA gene sequences from the cultures were used to

assign taxonomy and reconstruct phylogenetic relationships. We assigned taxonomy to all

assembled 16S rRNA gene sequences using the SILVA database SINA aligner v128 (81). A

Shapiro-Wilk test of normality was conducted in base R (82) on the distribution of SILVA

genera assignments from both media types. After concluding the data were non-parametric, we

performed a Kruskal-Wallis test in R (Assigned Genus ~ Media Type). We performed a *post hoc*

analysis (Dunn test, in R) to determine whether culturability within a Phylum varied by growth

medium type.

Taxonomic selection for phylogenetic reconstruction: To reconstruct a phylogeny of full

length 16S rRNA genes, our culture sequences were compared to NCBI's Microbial Genomes

and environmental sequence databases using web-blast (77). The top five hits for each sequence

from each NCBI database were chosen based on the highest percent coverage and lowest e-value

score and included in the reconstruction. *E. coli* K-12 was used as the outgroup of the

Alphaproteobacterial phylogeny, and *Bacillus subtilis* was used as the outgroup for the

Actinobacterial tree. These sequences aligned with MAFFT (83) with turn checking enabled. The

alignment was then trimmed using TrimAl (84) with the 'automated1' setting to optimize

sequence trimming for maximum likelihood phylogenetic analyses. We reconstructed

phylogenetic relationships from this trimmed alignment in the CIPRES Gateway (85).

Maximum-Likelihood (ML) trees were constructed using IQ-TREE with 10,000 ultrafast

bootstrap trees and Bayesian Information Criterion to select the best fit nucleic acid substitution

model (86, 87). For Actinobacteria, we used the SYM+R10 model and for Alphaproteobacteria

we used the GTR+F+I+G4 model. After an initial round of ML trees, sequence alignments were

heuristically curated with IQ-TREE to eliminate sequences that appeared in the tree more than

once. Finalized ML trees were then imported into the ARB environment (88), where any

duplicate sequences from our AZCC cultures were added to the ML trees through ARB's quick

add parsimony function. Final trees were visualized with FigTree (89).

Environmental contextualization of AZCC isolates: We matched the AZCC isolate full length

16S rRNA gene sequences against a database of the clustered OTUs from the shallow soil depth

profile samples (see *'soil microbial community analysis,'* above) using the usearch_global

command (90) at a stringency of ≥97% identity, in both strand orientations, with maxaccepts=1

and maxrejects=0.

Data Availability: Full-length Sanger Sequenced 16S rRNA gene sequences are available

on NCBI GenBank, accession numbers: MK875836 - MK875967. Illumina data from the 55 cm

Oracle Ridge community are available on the NCBI SRA, accession numbers: SRR9172130-

SRR9172198.

Acknowledgements: The authors would like to thank Nathan Abramson, Jasper Bloodsworth,

Brenna Bourque, Amanda Howe, Bridget Taylor, and H. James Tripp for assisting with sample

collection, culture maintenance, and DNA extractions relevant to this work. Funding from this

work came from startup funds provided to PC from the University of Arizona's Technology and

Research Initiative Fund (the Water, Environmental, and Energy Solutions initiative), and seed

grants from the Center for Environmentally Sustainable Mining and The University of Arizona

College of Agriculture and Life Sciences. Work at JCVI was supported by P01AI118687.

References:

1. Chaparro JM, Sheflin AM, Manter DK, Vivanco JM. 2012. Manipulating the soil

microbiome to increase soil health and plant fertility. Biol Fertil Soils 48:489–499.

2. Long PE, Williams KH, Hubbard SS, Banfield JF. 2016. Microbial Metagenomics

Reveals Climate-Relevant Subsurface Biogeochemical Processes. Trends Microbiol

24:600–610.

3. Fernández-Luqueño F, Valenzuela-Encinas C, Marsch R, Martínez-Suárez C, Vázquez-

Núñez E, Dendooven L. 2011. Microbial communities to mitigate contamination of PAHs

in soil-possibilities and challenges: A review. Environ Sci Pollut Res 18:12–30.

4. Tiedje JM, Asuming-Brempong S, Nüsslein K. 1999. Opening the black box of soil

microbial diversity. Appl Soil Ecol 13:109–122.

5. Jobbagy EG, Jackson RB. 2000. The Vertical Distribution of Soil Organic Carbon and Its

Relation to Climate and Vegetation. Ecol Appl 10:423–436.

6. Fierer N, Schimel JP, Holden PA. 2003. Variations in microbial community composition

through two soil depth profiles. Soil Biol Biochem 35:167–176.

7.   Ajwa HA, Rice CW, Sotomayor D. 1998. Carbon and Nitrogen Mineralization in

Tallgrass Prairie and Agricultural Soil Profiles. Soil Sci Soc Am J 62:942–951.

8.   Jobbágy EG, Jackson RB. 2001. The Distribution of Soil Nutrients with Depth: Global

Patterns and the Imprint of Plants. Biogeochemistry 53:51–77.

9.   Balesdent J, Basile-Doelsch I, Chadoeuf J, Cornu S, Derrien D, Fekiacova Z, Hatté C.

2018. Atmosphere–soil carbon transfer as a function of soil depth. Nature 559:599–602.

10.   Hillel D. 2013. Fundamentals of Soil PhysicsFundamentals of Soil Physics.

11.   Brewer TE, Aronson EL, Arogyaswamy K, Billings SA, Botthoff JK, Campbell AN, Dove

NC, Fairbanks D, Gallery RE, Hart SC, Kaye J, King G, Logan G, Lohse KA, Maltz MR,

Mayorga E, O'Neill C, Owens SM, Packman A, Pett-Ridge J, Plante AF, Richter DD,

Silver WL, Yang WH, Fierer N. 2019. Ecological and genomic attributes of novel

bacterial taxa that thrive in subsurface soil horizons. MBio 10:e01318-19.

12.   Eilers KG, Debenport S, Anderson S, Fierer N. 2012. Digging deeper to find unique

microbial communities: The strong effect of depth on the structure of bacterial and

archaeal communities in soil. Soil Biol Biochem 50:58–65.

13.   Blume E, Bischoff M, Reichert JM, Moorman T, Konopka A, Turco RF. 2002. Surface

and subsurface microbial biomass, community structure and metabolic activity as a

function of soil depth and season. Appl Soil Ecol 20:171–181.

14.   Spohn M, Klaus K, Wanek W, Richter A. 2016. Microbial carbon use efficiency and

biomass turnover times depending on soil depth - Implications for carbon cycling. Soil

Biol Biochem 96:74–81.

15.   Overmann J, Abt B, Sikorski J. 2017. Present and Future of Culturing Bacteria. Annu Rev

Microbiol 71:711–730.

16. Gao Y, Wu M. 2018. Free-living Bacterial Communities Are Mostly Dominated by

Oligotrophs. bioRxiv.

17. Morita RY. 1997. Bacteria in Oligotrophic Environments: Starvation-Survival Life Styles,

1st ed. Springer.

18. Schut F, Prins RA, Gottschal JC. 1997. Oligotrophy and pelagic marine bacteria: facts and

fiction. Aquat Microb Ecol 12:177–202.

19. Kuznetsov SI, Dubinina GA, Lapteva NA. 1979. Biology of Oligotrophic Bacteria. Annu

Rev Microbiol 33:377–387.

20. Cho J-C, Giovannoni SJ. 2004. Cultivation and Growth Characteristics of a Diverse

Group of Oligotrophic Marine Gammaproteobacteria. Appl Environ Microbiol 70:432–

440.

21. Giovannoni SJ, Cameron Thrash J, Temperton B. 2014. Implications of streamlining

theory for microbial ecology. ISME J 8:1553–1565.

22. Lauro FM, McDougald D, Thomas T, Williams TJ, Egan S, Rice S, DeMaere MZ, Ting L,

Ertan H, Johnson J, Ferriera S, Lapidus A, Anderson I, Kyrpides N, Munk AC, Detter C,

Han CS, Brown M V., Robb FT, Kjelleberg S, Cavicchioli R. 2009. The genomic basis of

trophic strategy in marine bacteria. Proc Natl Acad Sci 37:15527–15533.

23. Poindexter JS. 1981. Chapter 2, Oligotrophy: Fast and Famine Existence, p. 63–90. *In*

Advances in Microbial Ecology Volume 5.

24. Giovannoni SJ. 2017. SAR11 Bacteria: The Most Abundant Plankton in the Oceans. Ann

Rev Mar Sci 9:231–255.

25. Swan BK, Tupper B, Sczyrba A, Lauro FM, Martinez-Garcia M, Gonzalez JM, Luo H,

Wright JJ, Landry ZC, Hanson NW, Thompson BP, Poulton NJ, Schwientek P, Acinas

SG, Giovannoni SJ, Moran MA, Hallam SJ, Cavicchioli R, Woyke T, Stepanauskas R.

2013. Prevalent genome streamlining and latitudinal divergence of planktonic bacteria in

the surface ocean. Proc Natl Acad Sci 110:11463–11468.

26.   Santoro AE, Dupont CL, Richter RA, Craig MT, Carini P, McIlvin MR, Yang Y, Orsi

WD, Moran DM, Saito MA. 2015. Genomic and proteomic characterization of "

Candidatus Nitrosopelagicus brevis": An ammonia-oxidizing archaeon from the open

ocean. Proc Natl Acad Sci 112:1173–1178.

27.   Eichorst SA, Kuske CR, Schmidt TM. 2011. Influence of Plant Polymers on the

Distribution and Cultivation of Bacteria in the Phylum Acidobacteria . Appl Environ

Microbiol 77:586–596.

28.   Noell SE, Giovannoni SJ. 2019. SAR11 Bacteria Have a High Affinity and

Multifunctional Glycine Betaine Transporter. Environ Microbiol 00:00–00.

29.   Carini P, Dupont CL, Santoro AE. 2018. Patterns of thaumarchaeal gene expression in

culture and diverse marine environments. Environ Microbiol 20:2112–2124.

30.   Könneke M, Bernhard AE, de la Torre JR, Walker CB, Waterbury JB, Stahl D a. 2005.

Isolation of an autotrophic ammonia-oxidizing marine archaeon. Nature 437:543–6.

31.   Santoro AE, Casciotti KL. 2011. Enrichment and characterization of ammonia-oxidizing

archaea from the open ocean: Phylogeny, physiology and stable isotope fractionation.

ISME J 5:1796–1808.

32.   Hu SJ, Van Bruggen AHC, Grünwald NJ. 1999. Dynamics of bacterial populations in

relation to carbon availability in a residue-amended soil. Appl Soil Ecol 13:21–30.

33.   Saito A, Mitsui H, Hattori R, Minamisawa K, Hattori T. 1998. Slow-growing and

oligotrophic soil bacteria phylogenetically close to Bradyrhizobium japonicum. FEMS

Microbiol Ecol 25:277–286.

34.   Davis KER, Joseph SJ, Janssen PH. 2005. Effects of growth medium, inoculum size, and

incubation time on culturability and isolation of soil bacteria. Appl Environ Microbiol

71:826–834.

35.   Joseph SJ, Hugenholtz P, Sangwan P, Osborne CA, Janssen PH. 2003. Laboratory

cultivation of widespread and previously uncultured soil bacteria. Appl Environ Microbiol

69:7210–5.

36.   Zengler K. 2017. Protocols for High-Throughput Isolation and Cultivation, p. 27–35. *In*

McGenity, TJ, Timmis, KN, Balbina, N (eds.), Hydrocarbon and Lipid Microbiology

Protocols: Isolation and Cultivation. Springer Berlin Heidelberg, Berlin, Heidelberg.

37.   Jiang CY, Dong L, Zhao JK, Hu X, Shen C, Qiao Y, Zhang X, Wang Y, Ismagilov RF,

Liu SJ, Du W. 2016. High-throughput single-cell cultivation on microfluidic streak plates.

Appl Environ Microbiol 82:2210–2218.

38.   Khalili B, Weihe C, Kimball S, Schmidt KT, Martiny JBH. 2019. Optimization of a

Method To Quantify Soil Bacterial Abundance by Flow Cytometry. mSphere 4:1–10.

39.   Stingl U, Tripp HJ, Giovannoni SJ. 2007. Improvements of high-throughput culturing

yielded novel SAR11 strains and other abundant marine bacteria from the Oregon coast

and the Bermuda Atlantic Time Series study site. ISME J 1:361–371.

40.   Connon SA, Givannoni SJ. 2002. High-throughput methods for culturing microorganisms

in very-low-nutrient media. Appl Env Microbiol 68:3878–3885.

41.   Button DK, Schut F, Quang P, Martin R, Robertson BR. 1993. Viability and isolation of

marine bacteria by dilution culture: theory, procedures, and initial results. Appl Environ

Microbiol.

42.   Tahon G, Willems A. 2017. Isolation and characterization of aerobic anoxygenic

phototrophs from exposed soils from the Sør Rondane Mountains, East Antarctica. Syst

Appl Microbiol 40:357–369.

43.   Wang Y, Hammes F, Boon N, Chami M, Egli T. 2009. Isolation and characterization of

low nucleic acid (LNA)-content bacteria. ISME J 3:889–902.

44.   Proctor CR, Besmer MD, Langenegger T, Beck K, Walser JC, Ackermann M, Bürgmann

H, Hammes F. 2018. Phylogenetic clustering of small low nucleic acid-content bacteria

across diverse freshwater ecosystems. ISME J 12:1344–1359.

45.   Liu J, Hao Z, Ma L, Ji Y, Bartlam M, Wang Y. 2016. Spatio-temporal variations of high

and low nucleic acid content bacteria in an exorheic river. PLoS One 11.

46.   Liu J, Li JQ, Feng L, Cao H, Cui Z. 2010. An improved method for extracting bacteria

from soil for high molecular weight DNA recovery and BAC library construction. J

Microbiol 48:728–733.

47.   Monod J. 1949. The Growth of Bacterial Cultures. Annu Rev Microbiol 3:371–394.

48.   Lloyd KG, Steen AD, Ladau J, Yin J, Crosby L. 2018. Phylogenetically Novel Uncultured

Microbial Cells Dominate Earth Microbiomes. mSystems 3:e00055-18.

49.   Steen AD, Crits-Christoph A, Carini P, DeAngelis KM, Fierer N, Lloyd KG, Cameron

Thrash J. 2019. High proportions of bacteria and archaea across most biomes remain

uncultured. ISME J.

50.   Rappe MS, Connon SA, Vergin KL, Giovannoni SJ. 2002. Cultivation of the ubiquitous

SAR11 marine bacterioplankton clade. Nature 418:630–633.

51.   Sait M, Hugenholtz P, Janssen PH. 2002. Cultivation of globally distributed soil bacteria

from phylogenetic lineages previously only detected in cultivation-independent surveys.

Environ Microbiol 4:654–666.

52. Janssen PH, Yates PS, Grinton BE, Taylor PM, Sait M. 2002. Improved culturability of

soil bacteria and isolation in pure culture of novel members of the divisions Acidobacteria,

Actinobacteria, Proteobacteria, and Verrucomicrobia. Appl Environ Microbiol 68:2391–6.

53. Braun PR, Al-Younes H, Gussmann J, Klein J, Schneider E, Meyer TF. 2008. Competitive

inhibition of amino acid uptake suppresses chlamydial growth: Involvement of the

chlamydial amino acid transporter BrnQ. J Bacteriol 190:1822–1830.

54. Carini P, Steindler L, Beszteri S, Giovannoni SJ. 2013. Nutrient requirements for growth

of the extreme oligotroph "Candidatus Pelagibacter ubique" HTCC1062 on a defined

medium. ISME J 7:592–602.

55. Tanaka T, Kawasaki K, Daimon S, Kitagawa W, Yamamoto K, Tamaki H, Tanaka M,

Nakatsu CH, Kamagata Y. 2014. A Hidden Pitfall in the Preparation of Agar Media

Undermines Microorganism Cultivability. Appl Environ Microbiol 80:7659–7666.

56. Carlsson J, Nyberg G, Wrethén J. 1978. Hydrogen peroxide and superoxide radical

formation in anaerobic broth media exposed to atmospheric oxygen. Appl Environ

Microbiol 36:223–229.

57. Baxter M, Sieburth JM. 1984. Metabolic and ultrastructural response to glucose of two

eurytrophic bacteria isolated from seawater at different enriching concentrations. Appl

Environ Microbiol 47:31–38.

58. Lau JT, Whelan FJ, Herath I, Lee CH, Collins SM, Bercik P, Surette MG. 2016. Capturing

the diversity of the human gut microbiota through culture-enriched molecular profiling.

Genome Med 8:1–10.

59.   Diakite A, Dubourg G, Dione N, Afouda P, Bellali S, Ngom II, Valles C, Million M,

Levasseur A, Cadoret F, Lagier JC, Raoult D. 2019. Extensive culturomics of 8 healthy

samples enhances metagenomics efficiency. PLoS One 14.

60.   Walters W, Hyde ER, Berg-lyons D, Ackermann G, Humphrey G, Parada A, Gilbert JA,

Jansson JK, Caporaso JG, Fuhrman JA, Apprill A, Knight R. 2015. Improved Bacterial

16S rRNA Gene (V4 and V4-5) and Fungal Internal Transcribed Spacer Marker Gene

Primers. mSystems 1:1–10.

61.   Portillo MC, Leff JW, Lauber CL, Fierer N. 2013. Cell size distributions of soil bacterial

and archaeal taxa. Appl Environ Microbiol 79:7610–7617.

62.   Brewer TE, Handley KM, Carini P, Gilbert JA, Fierer N. 2016. Genome reduction in an

abundant and ubiquitous soil bacterium "Candidatus Udaeobacter copiosus." Nat

Microbiol 2.

63.   Carini P. 2019. A "Cultural" Renaissance: Genomics Breathes New Life into an Old Craft

Paul. mSystems 4:e00092-19.

64.   Olsen RA, Bakken LR. 1987. Viability of soil bacteria: Optimization of plate-counting

technique and comparison between total counts and plate counts within different size

groups. Microb Ecol 13:59–74.

65.   Aagot N, Nybroe O, Nielsen P, Johnsen K. 2001. An Altered Pseudomonas Diversity Is

Recovered from Soil by Using Nutrient-Poor Pseudomonas-Selective Soil Extract Media.

Appl Environ Microbiol 67:5233–5239.

66.   Lennon JT, Jones SE. 2011. Microbial seed banks: The ecological and evolutionary

implications of dormancy. Nat Rev Microbiol 9:119–130.

67.   Batut B, Knibbe C, Marais G, Daubin V. 2014. Reductive genome evolution at both ends

688   of the bacterial population size spectrum. Nat Rev Microbiol 12:841–850.

689 68. Sorensen JW, Dunivin TK, Tobin TC, Shade A. 2019. Ecological selection for small

690   microbial genomes along a temperate-to-thermal soil gradient. Nat Microbiol 4:55–61.

691 69. Kim S, Kang I, Seo JH, Cho JC. 2019. Culturing the ubiquitous freshwater actinobacterial

692   acI lineage by supplying a biochemical 'helper' catalase. ISME J.

693 70. Salcher MM, Schaefle D, Kaspar M, Neuenschwander SM, Ghai R. 2019. Evolution in

694   action: Habitat-transition leads to genome-streamlining in Methylophilaceae. ISME J

695   651331.

696 71. Lybrand RA, Rasmussen C. 2015. Quantifying climate and landscape position controls on

697   soil development in semiarid ecosystems. Soil Sci Soc Am J 79:104–116.

698 72. Parada AE, Needham DM, Fuhrman JA. 2016. Every base matters: Assessing small

699   subunit rRNA primers for marine microbiomes with mock communities, time series and

700   global field samples. Environ Microbiol 18:1403–1414.

701 73. Edgar RC. 2013. UPARSE: Highly accurate OTU sequences from microbial amplicon

702   reads. Nat Methods 10:996–998.

703 74. Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, Lesniewski RA,

704   Oakley BB, Parks DH, Robinson CJ, Sahl JW, Stres B, Thallinger GG, Horn DJ Van,

705   Weber CF. 2009. Introducing mothur : Open-Source , Platform-Independent , Community-

706   Supported Software for Describing and Comparing Microbial Communities □. Appl

707   Environ Microbiol 75:7537–7541.

708 75. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glöckner FO.

709   2013. The SILVA ribosomal RNA gene database project: improved data processing and

710   web-based tools. Nucleic Acids Res 41:D590-6.

76.   Kox MAR, Lüke C, Fritz C, van den Elzen E, van Alen T, Op den Camp HJM, Lamers

LPM, Jetten MSM, Ettwig KF. 2016. Effects of nitrogen fertilization on diazotrophic

activity of microorganisms associated with Sphagnum magellanicum. Plant Soil 406:83–

100.

77.   Johnson M, Zaretskaya I, Raytselis Y, Merezhuk Y, McGinnis S, Madden TL. 2008.

NCBI BLAST: a better web interface. Nucleic Acids Res.

78.   Lane DJ. 1991. Nucleic acid techniques in bacterial systematicsDevelopment and

Application of Nucleic Acid Probes.

79.   Griekspoor A, Groothuis T. 4Peaks. 1.8. Nucleobytes, Amsterdam, Netherlands.

80.   2019. Geneious Prime. v2019.0.1. Biomatters, Ltd., Auckland, New Zealand.

81.   Pruesse E, Peplies J, Glöckner FO. 2012. SINA : Accurate high-throughput multiple

sequence alignment of ribosomal RNA genes. Bioinformatics 28:1823–1829.

82.   R Core Team. 2014. R: A Language and Environment for Statistical Computing. 3.1.1. R

Foundation for Statistical Computing, Vienna, Austria.

83.   Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7:

improvements in performance and usability. Mol Biol Evol.

84.   Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. 2009. trimAl: A tool for automated

alignment trimming in large-scale phylogenetic analyses. Bioinformatics 25:1972–1973.

85.   Miller MA, Pfeiffer W, Schwartz T. 2010. Creating the CIPRES Science Gateway for

inference of large phylogenetic treesGateway Computing Environments Workshop.

86.   Minh BQ, Nguyen MAT, Von Haeseler A. 2013. Ultrafast approximation for phylogenetic

bootstrap. Mol Biol Evol 30:1188–1195.

87.   Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. 2014. IQ-TREE: A Fast and

Effective Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. Mol

Biol Evol 32:268–274.

88.	Ludwig W, Strunk O, Westram R, Richter L, Meier H, Yadhukumar, Buchner A, Lai T,

Steppi S, Jobb G, Förster W, Brettske I, Gerber S, Ginhart AW, Gross O, Grumann S,

Hermann S, Jost R, König A, Liss T, Lüßmann R, May M, Nonhoff B, Reichel B,

Strehlow R, Stamatakis A, Stuckmann N, Vilbig A, Lenke M, Ludwig T, Bode A,

Schleifer K. 2004. ARB: a software environment for sequence data. Nucleic Acids Res

32:1363–1371.

89.	Rambaut A, Drummond A. 2016. FigTree. 1.4.3.

90.	Edgar RC. 2010. Search and clustering orders of magnitude faster than BLAST.

Bioinformatics 26:2460–2461.

91.	Carini P, Marsden PJ, Leff JW, Morgan EE, Strickland MS, Fierer N. 2016. Relic DNA is

abundant in soil and obscures estimates of soil microbial diversity. Nat Microbiol

2:16242.

92.	Delgado-Baquerizo M, Oliverio AM, Brewer TE, Benavent-González A, Eldridge DJ,

Bardgett RD, Maestre FT, Singh BK, Fierer N. 2018. A global atlas of the dominant

bacteria found in soil. Science (80-) 359:320–325.

Figure 1: Dilution-to-extinction workflow. Soils were collected and brought to the lab where

they were homogenized in cell extraction buffer, layered over a Nycodenz solution, and

centrifuged (A). The cell layer was extracted from the Nycodenz solution and counted with flow

cytometry (B). Counted cells were diluted into growth medium in 96-well microtiter plates to an

average density of 5 cells well$_{-1}$ (C). After incubation, the 96-well microtiter plates were

screened for growth with flow cytometry, and wells displaying growth were subcultured into

larger volumes (D). After incubating the subcultures, flasks displaying growth were identified by

16S rRNA gene sequencing and molecular phylogeny (E). Aliquots of these identified

subcultures were cryopreserved at -80℃ (F).

Figure 2: Microbial culturability (v) was greater on ASM-low than on ASM-high. Bar

heights are the mean percent culturability ± standard deviation in 96 well microtiter plates (n=3)

as calculated from the initial cell inoculum and the proportion of wells positive for growth (41).

Double asterisks indicate Wilcoxon rank sum test P values ≤ 0.05.

Figure 3: ASM-low and ASM-high cultured distinct Alphaproteobacteria (a) and

Actinobacteria (b). Bar heights are the number of cultures obtained for each taxon and are

colored by the medium type on which they were isolated. The genera assigned to bacteria

isolated on ASM-low were distinct from those isolated on ASM-high (Kruskal-Wallis rank sum,

χ_2=19.05, P=1.28 × 10$_{-5}$). These differences were driven by differences in culturability across

medium types for actinobacterial genera, but not for alphaproteobacterial genera (Dunn test,

P≤0.000 for Actinobacteria and P=0.128 for Alphaproteobacteria).

Figure 4: The cultures isolated in this study were representative of several abundant soil lineages that show dynamic depth distributions in Oracle Ridge soils. Points are the mean relative abundances ± standard deviation (n=3) of 16S rRNA gene sequence phylotypes that matched the 16S rRNA gene sequences obtained from cultured isolates at ≥97% identity. Error bars that are not visible are located behind the symbol. Assigned genera names and the average (n=3) relative rank of each phylotype at 55 cm are shown. Cultures classified at the genera level as *Mycobacterium* and *Arthrobacter* cultures matched more than one phylotype in the

cultivation-independent surveys. The best-matching OTU number is designated in parentheses.

Triangles are Alphaproteobacteria. Circles are Actinobacteria.

Figure 5: The mean nucleic acid fluorescence of taxa isolated on ASM-low was significantly

lower than for those microbes isolated on ASM-high. Points are the mean natural logarithm

(ln) of the quantified nucleic acid fluorescence (in arbitrary units [AU]) of fixed and SYBR

green I-stained stationary phase cultures. The mean fluorescence value was obtained from

manually-gated histogram plots of fluorescence within the Guava EasyCyte software. Only those

cultures that were defined as pure cultures are plotted. The horizontal line in each plot is the

mean fluorescence value and the box surrounding the mean is a 95% confidence interval.

Shading illustrates the relative distribution of fluorescence values within each medium type.

Supplementary Table and Figure Captions.

Supplementary Table 1. Artificial Subterranean Growth Medium (ASM) -low and -high

formulations including concentrations of carbon sources, amino acids, vitamins, trace metals,

amino acids, and inorganic nutrients.

Supplementary Figure 1: The soils used for cultivation were oligotrophic. (A) points are the

measured soil organic carbon content of 184 soil samples collected from shallow subsurface pits

located across the United States (from ref. (11)). (B) points are the measured NO_3-N content of

121 soils (113 surface soils) collected across the United States (from refs (91, 92)). The organic

carbon percent and NO_3-N measured in our 55 cm sample from Oracle Ridge is illustrated with a

red dashed line in both panels. Our sample falls on the low end of both organic carbon content

and NO_3-N content across this broad selection of soils.

Supplementary Figure 2: Alphaproteobacterial IQ-TREE Maximum Likelihood Full

Length 16S rRNA gene sequence phylogeny. The tree was constructed using a GTR+F+I+G4

nucleotide substitution model iterated over 10,000 ultrafast bootstraps. The scale bar indicates

the FigTree proportionally transformed branch length. The 16S rRNA gene sequence from *E.*

coli K-12 was used as an outgroup. Clade bootstrap support is indicated by colored nodes as

indicated in the legend. This tree includes 16S rRNA gene sequences that amplified from all pure

alphaproteobacterial cultures isolated in this study (bolded black leaves denoted with 'AZCC'),

as well as their best sequence matches to NCBI microbial genomes (dark blue leaves), NCBI 16S

rRNA gene sequences from cultured isolates (orange leaves), and NCBI 16S rRNA gene

sequences from environmental clones (green leaves). The growth medium concentration on

which each AZCC culture was isolated are indicated by colored bars on the right side of the

phylogenetic tree (red for ASM-high and blue for ASM-low).

Supplementary Figure 3: Actinobacterial IQ-TREE Maximum Likelihood Full Length 16S

rRNA gene sequence phylogeny. The tree was constructed using a SYM+R10 nucleotide

substitution model iterated over 10,000 ultrafast bootstraps. The scale bar indicates the FigTree

proportionally transformed branch length. The 16S rRNA gene sequence from *B. subtilis* DSM-

10 was used as an outgroup. Clade bootstrap support is indicated by colored nodes as indicated

in the legend. This tree includes 16S rRNA gene sequences that amplified from all pure

actinobacterial cultures isolated in this study (bolded black leaves denoted with 'AZCC'), as well

as their best sequence matches to NCBI microbial genomes (dark blue leaves), NCBI 16S rRNA

gene sequences from cultured isolates (orange leaves), and NCBI 16S rRNA gene sequences

from environmental clones (green leaves). The growth medium concentration on which each

AZCC culture was isolated are indicated by colored bars on the right side of the phylogenetic

tree (red for ASM-high and blue for ASM-low). Actinobacterial classes are indicated by colored

vertical bars to the right of the tree figure: Thermoleophilia (purple), Acidimicrobiia (teal), and

Actinobacteria (Orange-Red).

Supplementary Figure 4: Distribution of nucleic acid fluorescence values by Phylum (a)

and Genus (b). Data are natural logarithm-transformed nucleic acid fluorescence values of fixed

and SYBR Green I-stained stationary phase cultures. Plots show all genera with n ≥ 3 cultures in

one or both culture media types. Shaded regions illustrate the distribution of fluorescence values

within culture media groups.