The concentration of heterotrophic growth substrates in culture medium is a crucial determinant of the culturability of subsurface soil microbes

Authors: Ryan P. Bartelme[1], Joy M. Custer[1], Christopher L. Dupont[2], Josh L. Espinoza[2], Manolito Torralba[2], Banafshe Khalili[3], Paul Carini[1*]

Affiliations:

[1]Department of Soil, Water and Environmental Science, University of Arizona, Tucson, AZ 85721

[2]J. Craig Venter Institute, La Jolla, CA 92037

[3]Department of Ecology and Evolutionary Biology, University of California, Irvine, CA 92697

*Corresponding author: paulcarini@email.arizona.edu

Abstract

Soil microbial communities have crucial roles in soil fertility and biogeochemistry. While surface soils have been relatively well characterized, we know far less about the microbiology of oligotrophic shallow subsurface soils that often contain microbial taxa that are distinct from surface soils. We hypothesized that the concentration of heterotrophic growth substrates in microbiological growth medium is a critical determinant of the ability to culture subsurface soil microbes. To test this, we conducted dilution-to-extinction experiments with microbial cells separated from subsurface soils collected in a mid-elevation conifer forest in Arizona, USA. We diluted cells into two defined growth media with identical concentrations of inorganic nutrients and vitamins but contained a 100-fold difference in the availability of an amino acid and organic carbon mixture. Across both media formulations, we isolated a total of 133 pure cultures. The concentration of heterotrophic growth substrate in the medium facilitated the isolation of taxonomically distinct microbes. We isolated representative cultures of the most abundant member in the soil community (*Bradyrhizobium* spp.) and representatives of five of the top 10 most abundant *Actinobacteria* phylotypes, including *Nocardioides* spp., *Mycobacterium* spp., and several other phylogenetically divergent lineages. Flow cytometric analysis of the cultures showed that SYBR green-stained cells isolated on medium with reduced heterotrophic substrate concentrations had significantly lower nucleic-acid fluorescence than those isolated on medium with higher substrate concentrations. These results show that medium substrate concentration can facilitate the isolation of certain microbial lineages and that dilution-to-extinction is an effective method to isolate abundant soil microbes.

Importance

Isolating environmental microbes and studying their physiology under controlled conditions is an essential aspect of understanding their ecology. In this study, we use modified high-throughput cultivation methods to isolate subsurface soil microbes. Subsurface ecosystems are typically nutrient-poor environments that harbor diverse microbial communities—the vast majority of which are uncultured. We show that a critical component of whether a microbe is

45 culturable from subsurface soils is the concentration of growth substrates in the culture
46 medium. Our results offer new insight into technical approaches and growth medium design
47 that can be used to access the vast uncultured diversity of soil microbes.
48
Introduction:
Soil microbial communities are tremendously diverse and mediate crucial aspects of plant
fertility, biogeochemistry, pollutant mitigation, and carbon sequestration (1–4). While the
diversity and community composition of surface soils have been relatively well described, we
know far less about the microbes inhabiting deeper soils, despite their key role in soil
formation and mineralization of key plant nutrients. In contrast to surface soils that are
typically rich in plant-derived compounds, subsurface soils are often oligotrophic and
characterized by lower amounts of organic carbon, mineralizable nitrogen, and phosphorus (5–
8). The temperature and soil moisture of subsurface soils also vary less over time than surface
environments (9). As depth from the surface increases, the diversity of subsurface microbial
communities decreases and the dissimilarity between the subsurface microbial community
composition increases relative to surface microbial communities (10, 11). Moreover, the
microbial taxa that are abundant in subsurface environments are generally absent from
microbial culture and genome databases (10). Thus, we do not understand fundamental
biological aspects of a major fraction of subsurface soil microbes (10).
Part of the challenge in culturing and studying subsurface microbes is that many of
these uncultured lineages likely require low nutrient concentrations for optimal growth. These
microbes are termed 'oligotrophs' and are paradoxical to what we understand about most
cultivated microbes that tend to grow best when nutrient availability is high. Although
oligotrophs dominate most free-living microbial ecosystems (12), the concept of oligotrophy is
enigmatic: there is no consensus on what defines oligotrophic metabolism. Several general
traits of oligotrophic microbes have emerged from studies of aquatic oligotrophs. For example,
oligotrophs are small, slowly growing cells (13–16). The genomes of marine oligotrophs tend to
be streamlined often coding for fewer copies of the rRNA gene operon and transcriptional
regulators than microbes capable of robust growth at high nutrient availability, suggesting
oligotrophs lack the ability to sense and rapidly respond to varying environmental conditions
(12, 14, 17). Instead, genomic inventories in oligotrophs point toward a reliance on broad-
specificity, high-affinity transport functions and relatively constitutive gene expression (15, 18–
20). Because oligotrophs do not attain high yields on low nutrient media, traditional methods
of quantifying microbial growth are ineffective (for example, optical density measurements
and plating on solid media) (16, 21, 22).
While the physiology of aquatic oligotrophs have been fairly well studied, far fewer studies
have focused on the physiology of oligotrophic soil microbes. Although many soil cultivation
approaches rely on the use of low nutrient media (23–26), we lack a fundamental
understanding about why these media facilitate the isolation of distinct taxa and often
produce higher culturable counts than nutrient-rich media. We hypothesized that the
concentration of heterotrophic growth substrates in a growth medium would constrain the
taxa that grow on a defined artificial medium. We tested this by extracting cells from
oligotrophic subsurface soils and inoculating high-throughput dilution-to-extinction
experiments on a defined medium with differing amounts of heterotrophic substrates. We

isolated several bacteria representing abundant microbial community members. In these
experiments, the substrate content significantly influenced which taxa grew in the laboratory.
Moreover, we provide evidence that cells isolated on low nutrient medium may contain
reduced nucleic acid content relative to those isolated on higher nutrient medium.
93
Results:
We collected shallow subsurface soil (55 cm) from the Oracle Ridge field site in a mid-
elevation conifer forest that is part of the Santa Catalina Mountains Critical Zone Observatory
in Arizona, USA. We modified high-throughput dilution-to-extinction approaches designed for
aquatic microbes (21, 27) to culture shallow subsurface soil microbes from these samples (Fig.
1). The primary modification to existing protocols was to add a cell-separation step to detach
inoculum cells from mineral soils prior to diluting them to extinction. We diluted the inoculum
to an average of 5 cells well^{-1} into deep-well polytetrafluoroethylene 96-well plates containing
a defined Artificial Subterranean Medium (ASM), having either low or high concentrations of
heterotrophic growth substrates (ASM-low and ASM-high, respectively). The ASM-low and
ASM-high media contained identical inorganic mineral and vitamin amendments but a 100-
fold difference in the concentration of organic carbon and amino acids (Supplementary Table
1). We prepared triplicate 96-well plates for each growth medium formulation. These dilution-
to-extinction experiments were screened for growth with flow cytometry after ~4 weeks of
incubation and again after ~11 weeks of incubation. Wells displaying growth were sub-cultured
into larger volumes and subsequently identified by 16S rRNA gene sequencing.
Across both medium types, a total of 214 wells (119 for ASM-low and 95 for ASM-high)
exceeded our growth threshold (1.0 × 10^4 cells ml^{-1}) after 11 weeks of incubation. We
successfully propagated 182 (85%) of the cultures from microtiter plates to polycarbonate
flasks containing fresh medium. Of the cultures that successfully propagated, we confirmed
that 73% (133 cultures) were pure cultures by amplifying and sequencing full-length 16S rRNA
gene sequences. The remaining 49 cultures were either mixed (16S rRNA genes did not
assemble due to base ambiguities) or did not amplify under several amplification conditions.
We defined microbial culturability as the ratio of cells capable of being isolated in a given
condition to the total number of cells initially diluted into a cultivation chamber (28). By the
end of the experiment, we approached ~20% culturability across both medium formulations
(Fig. 2). In general, microbial culturability was higher for ASM-low than for ASM-high, but this
effect was only significant after 4 weeks of incubation (Fig. 2; Wilcoxon rank-sum test $P \leq 0.05$
at 4 weeks).
Cultures isolated on ASM-high were taxonomically distinct (at a genus level) from those
isolated on ASM-low (Kruskall-Wallis rank sum, $P=1.28 \times 10^{-5}$). All pure cultures grown on both
ASM-low and ASM-high belonged to one of two bacterial phyla: Actinobacteria (110 cultures;
83% of the pure cultures) or Proteobacteria (23 cultures; 17% of the pure cultures) (Fig. 3). The
Proteobacterial cultures were exclusively Alphaproteobacteria, 57% of which were isolated on
ASM-low medium and the remainder (43%) were isolated on ASM-high (Fig. 3 and
Supplementary Fig. 1). Cultures that classified as *Bradyrhizobium* spp. were the most frequent
alphaproteobacterial isolates (13 isolates), seven of which were isolated on ASM-low medium.
Cultures classified as *Reyranella* spp. and *Nordella* spp. were also isolated on both ASM-high

and ASM-low medium. The remaining five alphaproteobacterial cultures were taxonomically
distinct—three were isolated on ASM-low and two on ASM-high.
The taxonomy of the actinobacterial cultures was split among three classes:
Actinobacteria (107 cultures), Thermoleophilia (2 cultures), and Acidimicrobiia (1 culture). Of
these Actinobacteria, 65 (59%) were isolated on ASM-low, and 45 (41%) on ASM-high. The
actinobacterial cultures were numerically dominated by two genera that were differentially
isolated on ASM-low and ASM-high: *Nocardioides* and *Mycobacterium*. *Nocardioides* spp. (46
cultures) were exclusively isolated on ASM-low medium (Fig 3 and Supplemental Fig. 2). Other
cultures that were isolated on ASM-low included those classified as *Arthrobacter* (3 cultures),
Marmoricola (2 cultures), *Nakamurella* (2 cultures), *Aeromicrobium* (1 culture), *Blastococcus* (1
culture), and *Patulibacter* (1 culture) (Fig 3 and Supplemental Fig 2). While cultures classified as
Mycobacterium sp. were generally isolated on ASM-high (38 cultures), we did isolate seven
mycobacterial cultures on ASM-low medium—five of which form a phylogenetically distinct
cluster from those isolated on ASM-high (Fig. 3 and Supplementary Fig. 2). Other
actinobacterial cultures isolated on ASM-high included *Jatrophihabitans* (4 cultures),
Conexibacter (1 culture), and *Amycolatopsis* (1 culture).
Interestingly, we isolated what are likely the first members of two novel actinobacterial
lineages on ASM-low. The first such culture—Microtrichales sp. str. AZCC_0197—belongs to
the Microtrichales order of the Acidimicrobiia class. The best 16S rRNA gene sequence match
to an existing isolate is 93.4% identity to *Aquihabitans daechunggensis* str. G128, but str.
AZCC_0197 more closely matched numerous 16S rRNA gene sequences from environmental
clones of uncultured Acidimicrobiia. The second lineage—Frankiales sp. strains AZCC_0102
and AZCC_0072—classified as members of the Frankiales order of the Actinobacteria class
with best matches of <97% nucleotide identity to existing Frankiales isolates (29).
Several of the cultures we isolated are representative of abundant members of the
subsurface soil microbial community at Oracle Ridge. We matched the 16S rRNA gene
sequences from our cultures to the operational taxonomic units (OTUs) present in the Oracle
Ridge soil sample collected at 55 cm. The cultures isolated from these 55 cm subsamples are
representative of 13 OTUs (Fig. 4) that comprise a total of 8.8% of the total amplifiable
microbial community in subsamples collected at the same time and depth. For example, our
Bradyrhizobium isolates match a single *Bradyrhizobium* OTU that was the most abundant OTU
at 55 cm (relative abundance of ~5.7%) (Fig. 4). Similarly, we isolated representatives of most
of the abundant Actinobacteria (Fig. 4) including: two *Mycobacterium* OTUs (the 11[th] and 17[th]
most abundant OTUs overall, and the most abundant and sixth most abundant actinobacterial
OTUs, respectively); *Nocardioides* (the 13[th] most abundant OTU overall, and third most
abundant actinobacterial OTU), and two *Arthrobacter* OTUs (16[th] and 1,271[st] most abundant
OTU overall, and the fifth and 143[rd] most abundant actinobacterial OTUs). The other
Actinobacteria cultured in these experiments represent rarer species in bulk soils. The 16S
rRNA gene sequences from several of our pure cultures did not match any of the 16S rRNA
gene amplicon OTUs amplified from these soils, including *Nakamurella* (2 cultures),
Nocardioides (5 cultures), *Mycobacterium* (1 culture), *Jatrophihabitans* (1 culture), *Patulibacter* (1
culture), *Conexibacter* (1 culture), *Rhizobiales* sp. "GAS113" (1 culture), *Reyranella* (1 culture),
and *Microtrichales* sp. str. AZCC_0197.

Cultures isolated on ASM-low exhibited significantly lower mean nucleic acid fluorescence
than those isolated on ASM-high (Fig. 5; Kruskal-Wallis Rank sum P=6.3 × 10^{-7}). The overall
mean fluorescence was not significantly different across phyla (Kruskal-Wallis Rank sum
P=0.647) but was significant across assignable genera (Kruskal-Wallis Rank sum P=3.0x10^{-6}).
Moreover, the mean nucleic acid fluorescence values within a given genera were generally
similar (Supplementary Fig. 3). For example, *Mycobacterium* isolates had relatively high nucleic
acid fluorescence, regardless of which medium they were isolated on (Supplementary Fig. 3).
In contrast, *Nocardioides* (ASM-low) and *Jatrophihabitans* (ASM-high) displayed relatively low
nucleic acid fluorescence. Interestingly, when cultured on ASM-high, we observed a clear
nucleic acid fluorescence dichotomy across the *Bradyrhizobium* isolates (Supplementary Fig. 3).
185
Discussion:
We designed a proof-of-concept workflow to determine the feasibility of high-throughput
dilution-to-extinction cultivation for the isolation of soil microbes. The method was
conceptually based on an improved high-throughput culturing workflow that was designed to
isolate oligotrophic marine microbes in inert plastic microtiter plates using automated flow
cytometry to detect microbial growth rapidly and at relatively low cell densities (21). However,
unlike aquatic samples, microbial cells in soils are heterogeneously dispersed within, or
attached to, a complex matrix comprised of non-cellular organic matter and minerals. The
inherent complexity of the soil matrix complicates accurate enumeration of viable cells—
especially of small oligotrophic microbes—because mineral and organic matter can interfere
with flow cytometry. To circumvent these issues, we separated cells by gently shaking soils in a
cell extraction buffer containing a dispersing agent and a nonionic surfactant. Cells were
separated from this slurry by density gradient centrifugation (Fig. 1). This procedure allows
cells to be 'floated' on top of a solution of Nycodenz while allowing minerals to pellet through
the Nycodenz (30).
We expected 168 pure cultures across all experiments, based on the proportion of
cultivation chambers displaying measurable growth (214 chambers) and the number of
chambers inoculated (576 chambers). The number of pure cultures we obtained (133) was
within 38% of this value. However, this result does not account for the cultures that were
initially scored as positive for growth but did not successfully subculture. Some of the cultures
that failed to propagate from microtiter plates to larger volumes might have been false
positives, where flow cytometer instrument noise or well-to-well carryover was mistaken for a
low-density culture. The mean culturability we observed for a given experiment (1.4%-11%;
Fig. 2) is comparable to dilution-to-extinction cultivation studies of marine microbes, which
report 0.5%-14.3% culturability (27). Similar to previous observations for soil microbes (25), we
observed the culturability increased by lengthening the incubation time (Fig. 2).
The concentration of heterotrophic growth substrates in the isolation medium was a
determinant of the ability to isolate certain microbial lineages (Fig. 3). Carbon type and
availability are crucial for heterotrophic soil microbes since carbon acts as both a source of
electrons for respiration and carbon for biomass. To accommodate this requirement, many
common microbial growth medium formulations for heterotrophic microbes supply diverse
growth substrates (yeast extract or casein digests, for example), usually at concentrations
much higher than are normally available *in situ*. Two key assumptions made with these

common media formulations are that: 1) microbes will use only the relevant constituents and
any remaining compounds will have minor or no effect on microbial growth; and 2) microbes
grow optimally when nutrient availability is much greater than their half saturation constant
(31). While many microbes have the capacity to grow on these commonly used complex, high-
nutrient formulations, environmental nucleic acid data informs us that the vast majority of
Earth's microbes cannot (32).
Numerous studies have demonstrated that dilute growth media is superior to substrate
rich growth media for the isolation of novel soil microbes (25, 26, 33, 34). However, the
physiological explanation as to why low-nutrient media facilitates the growth of diverse
microbes, or high nutrient concentrations inhibit the growth of some taxa, are unclear. One
possible explanation for these concentration-dependent effects may be that growth medium
formulations applied at high concentrations contain high amounts of inhibitory substances—
substances that are reduced to non-inhibitory levels in dilute medium formulations. For
example, a key amino acid transporter in *Chlamydia trachomatis* can be blocked by non-
essential amino acids, preventing the transport of required amino acids, resulting in growth
inhibition (35). A similar phenomenon was demonstrated in the marine oligotroph '*Candidatus*
Pelagibacter ubique', where alanine was conditionally required for cell division but abolished
growth at higher concentrations (36). Furthermore, reactive oxygen species can be produced
during the autoclaving of nutrient-rich medium that either directly inhibit growth or combine
with organics in the medium to form inhibitory compounds (37, 38). Finally, growth inhibition
may be the result of misbalanced regulation of growth or accumulation of nutrient storage
structures (poly-β-hydroxybutyrate, for example) ultimately leading to cell lysis (39). A better
understanding of the mechanism(s) that enable growth on low nutrient medium—or prevent
growth on high-nutrient medium—may help us design better strategies for isolating
uncultivated lineages. Critically, having a collection of cultures that were isolated on medium
with identical constituents applied at different concentrations is a first step towards an
experimental method capable of addressing these questions.
While several of the taxa we isolated were abundant microbial members of the shallow
subsurface microbial community (Fig. 4), other isolates were rare or not identified in the
cultivation-independent soil microbial community at all. Although the cultivation of microbial
phylotypes that are not observed in cultivation-independent analyses has been observed (40),
dilution-to-extinction favors the cultivation of abundant microbes in a given sample (28), such
that the appearance of rarer taxa or taxa that were not observed in the original sample is
unusual. There are several possible explanations for this observation. First, the absence of a
particular taxon in a soil microbiome analysis may be the result of insufficient sequencing
depth (40). Similarly, the 'universal' primers used in the soil microbiome analysis (41) may have
been biased against some of the divergent lineages we obtained in pure culture, resulting in
either underrepresentation of these phylotypes in the original community or no amplification
at all. Finally, the taxa isolated in these experiments may have had similar buoyancies and
migrated to similar locations in the density gradient during cell extraction effectively skewing
the proportions of microbes that were diluted into microtiter plate wells. At this point, we do
not have sufficient evidence pointing toward one of these scenarios over the other.
Additionally, as is true in any microbial cultivation experiment, there are many taxa that we did
not isolate. In particular, these soils contained high relative abundances of Verrucomicrobia

related to 'Ca. Udaeobacter copiosus' (42), and Acidobacteria (Subgroup 6), which belong to
highly sought-after lineages of uncultured microbes (22). The reasons for not culturing these
(and other) lineages are numerous but may be the result of inappropriate medium composition
(43, 44), toxic compounds in the cell separation constituents, long doubling times (> ~6 days),
or dormancy (reviewed in (45)).
Similar to aquatic oligotrophs (15, 46), we provide evidence that soil oligotrophs contain
reduced nucleic acid content relative to other soil microbes that prefer higher nutrient
concentrations (Fig. 5). Depending on the taxa in question and the effective population size,
microbial genome reduction can be driven by genetic drift or streamlining selection (reviewed
in (47)). Genome streamlining is strongly linked with oligotrophy in free-living aquatic microbes
as a mechanism to reduce the overhead cost of replication in periodically nutrient-limited
environments (reviewed in (14)). However, direct evidence for streamlined genomes in
terrestrial oligotrophs has been elusive. Metagenome-assembled genomes suggest that some
lineages of abundant and ubiquitous uncultured Verrucomicrobia may contain reduced
genome sizes (42), and a more recent study showed that fire-affected warm soils selected for
groups of microbes with significantly smaller genomes than cooler soils (48). However, there
are few definitive ways to identify these taxa as oligotrophs short of culturing them. The
appearance of reduced nucleic acid content in cultures isolated on ASM-low is also an
indication that genome reduction may be a successful life strategy for soil oligotrophs. We
speculate that "streamlined" genome sizes in soil oligotrophs may be larger than those
observed in marine habitats, since success in the complex soil matrix necessitates a greater
variety of gene function to succeed (14).
We were surprised to identify taxa with such distinct growth substrate concentration
preferences from highly oligotrophic soils (0.5% organic carbon). For example, while the
environmental distribution of Nocardioides sp.—which were isolated on ASM-low—indicate a
preference for deeper, more oligotrophic soils (Fig. 4), we also isolated numerous
Mycobacterium spp. from the same samples. The majority of these Mycobacterium isolates
were isolated on ASM-high medium and did not show a clear preference for subsurface soils
(Fig. 4). Interestingly, we observed that the Nocardioides sp. appeared sooner (within 4 weeks)
in the dilution-to-extinction screens, and Mycobacterium sp. growth was only apparent
between 4 and 11 weeks of incubation. These observations suggest that the oligotrophs from
these soil samples may grow relatively rapidly compared to the microbes requiring higher
nutrient concentrations. This finding is contrary to the perception that oligotrophic microbes
grow more slowly than those microbes capable of growth on higher nutrient concentrations
(16), suggesting that oligotrophy in shallow subsurface soils may be characterized by traits that
are distinct from those described for aquatic microbes.
The development of cultivation techniques emphasizing the high-throughput and sensitive
detection of microbial growth on low-nutrient medium has revolutionized the field of aquatic
microbial ecology in that these methods enable the cultivation of microbes that were
previously inaccessible using standard techniques (21, 27, 49–51). Here we show that similar
principles facilitate the cultivation of abundant soil oligotrophs. Moreover, we demonstrate
that in addition to considering the nutritional composition of a given growth medium, another
obstacle to culturing some lineages is the concentration of growth substrates in the growth
medium. Although we do not yet understand the mechanism of substrate-induced growth

inhibition, there is evidence that this phenomenon is widespread and results in the inability to
culture some microbial lineages in the laboratory (12). Future mechanistic studies aimed at
deciphering the physiology of substrate-induced growth inhibition will likely lead to new
cultivation approaches that will allow us to isolate abundant free-living oligotrophic microbes.
311
Materials and Methods:
Soil source: Fresh soil samples were collected from a soil pit on August 16, 2017 at the
Oracle Ridge site in the Catalina Jemez Critical Zone Observatory (coordinates: 32.45 N, -
110.74 W, elevation 2,103 m). We collected ~300 g subsamples from 0-5 cm, 10 cm, 20 cm, 30
cm, 40 cm and 55 cm depths. We analyzed the microbial community composition at each
depth (see *'Soil microbial community analysis,'* below) and conducted cultivation experiments
from the 55 cm soil sample. Soils were kept cool with ice packs for <4 h while in transit to the
laboratory. At the laboratory, the soils were sieved to 2 mm and kept at 4°C for 50 days at
which point cells were separated from mineral soil.
Soil microbial community analysis: Replicate 1.0 g subsamples were weighed into MoBio
PowerSoil DNA extraction kits and extracted according to the manufacturer's instructions. We
amplified 16S rRNA gene fragments using 515F-Y (5'-
TATGGTAATTGTGTGYCAGCMGCCGCGGTAA-3') and 926R (5'-
AGTCAGTCAGGGCCGYCAATTCMTTTRAGT-3') (52). PCR products were purified using the
QIAquick PCR purification kit (Qiagen, Germantown, MD) per manufacturer's specification.
Cleaned products were quantified using Tecan fluorometric methods (Tecan Group,
Mannedorf, Switzerland), normalized, and pooled for Illumina MiSeq sequencing using custom
sequencing primers and the MiSeq Reagent v2 500 cycle Kit (Roche, Branford, CT) following
the manufacturer's protocols. Operational taxonomic units (OTUs) were generated *de novo*
from raw Illumina sequence reads using the UPARSE pipeline (53). Paired-end reads were
trimmed of adapter sequences, barcodes and primers prior to assembly. Sequences of low-
quality, and singletons were discarded, and the remaining sequences were subjected to a
dereplication step and abundances were determined. Chimera filtering of the sequences was
completed during clustering while taxonomy was assigned to the OTUs with mothur (54) using
version 123 of the SILVA 16S ribosomal RNA database (55) as the reference. OTUs and
corresponding taxonomy assignment tables were generated and used in subsequent analyses.
338
Cell separation: Cells were separated from sieved soils using density gradient centrifugation
with Nycodenz. Briefly, we added 0.5 g wet soil to 44.8 ml of cell extraction buffer (137.5 mM
NaCl, 26.78 mM tetrasodium pyrophosphate, and 0.27% (v/v) Tween 80). The soil-buffer slurry
was vortexed for 30 seconds and shaken horizontally on a platform shaker for 2 hours at 4°C.
We layered 15 ml aliquots of this soil-buffer slurry over 10.0 ml of 80% (w/v) Nycodenz solution
in 50 mM tetrasodium pyrophosphate. We used 50 ml Nalgene Oak Ridge high-speed
polycarbonate centrifuge tubes for density gradient centrifugations. Tubes containing the soil-
buffer solution with Nycodenz were centrifuged at 17,000 × *g* for 30 minutes at 16°C. We
extracted 3 × 0.5 mL aliquots from the resulting density gradient at a location of ~25 mm above
the bottom of the tube (coincident with the approximate level of the top of the Nycodenz
solution) to sterile microcentrifuge tubes containing 1.0 ml 137.5 mM NaCl. The
microcentrifuge tubes containing Nycodenz/NaCl were vortexed and centrifuged for 20 min at

17,000 × g. The resulting faint cell pellets were resuspended in 137.5 mM NaCl, pooled and
stored at 4°C.
353
Dilution-to-extinction: An aliquot of cells extracted from the Nycodenz gradient were fixed
with 1.75% (final v/v) formaldehyde and stained with SYBR Green I (final stain concentration
was a 1:4,000 dilution of commercial stock) for 3.5 h at room temperature in the dark. Cells
were enumerated using a Millipore Guava flow cytometer, as described elsewhere (36). We
diluted cells into Artificial Subterranean Medium (ASM) -high or ASM-low nutrient medium
(Supplementary Table 1) to a density of 5 cells mL^{-1} and aliquoted 1.0 ml of the dilute cell
suspension into the wells of 2 ml polytetrafluoroethylene 96-well microtiter plates (Cowie
Technology, New Castle, DE) so that on average each well contains 5 cells. Plates were covered
with plastic lids that allow air circulation and incubated at 16ºC in the dark. We screened the
dilution-to-extinction plates for growth by fixing (1.75% formaldehyde) and staining (1:4,000
dilution of commercial SYBR Green I stock) aliquots for 18h in the dark at room temperature
and counting by flow cytometry (EMD-Millipore Guava EasyCyte), as described previously (21).
We screened plates for growth at 4 and 11 weeks after inoculation. Positive cultures were
defined as cultures that exceeded 1.0 × 10^4 cells ml^{-1}.
368
Culturability estimates and estimates of the number of pure cultures: Culturability estimates
were determined by the equation: $V = -\ln(1-p)/X$, where V is the estimated culturability, p is the
proportion of inoculated cultivation chambers that displayed measurable growth (number of
chambers positive for growth ÷ total number of chambers inoculated), and X is the number of
cells added to each cultivation chamber (28). The number of pure cultures (u) was estimated as
follows: $u=-n(1-p)\ln(1-p)$, where n is the number of inoculated growth chambers (28).
375
Culture transfer and storage: We subcultured positive growth chambers into 25 ml of the
respective growth medium (ASM-high or ASM-low) in acid-washed, sterile polycarbonate
flasks and incubated them at 16ºC. At the time of transfer, we assigned cultures a unique
Arizona Culture Collection (AZCC) number. Flasks were monitored for growth every other
week for 2 months. Flasks displaying growth within two months were cryopreserved in 10%
glycerol and stored at -80ºC. If no growth appeared within two months, the cultures were
discarded and the assigned AZCC number was retired.
383
384 *Mean fluorescence calculations:* We calculated the mean fluorescence of each culture from
385 the subcultures grown in 25 ml volumes 12-15 weeks after inoculating. Culture aliquots were
386 fixed and stained for 15-18 h as described above in 'Dilution-to-extinction.' We manually gated
387 histograms of the intensity of SYBR Green I fluorescence (in arbitrary units) and extracted the
388 mean fluorescence of the gated peak for each culture.
389
Culture identification: Cultures were identified by full length 16S rRNA gene sequencing.
Briefly, we filtered 5-10 ml of cell biomass from 25 mL cultures on to 0.2 µm-pore size Supor
filters and extracted DNA using a Qiagen PowerSoil DNA extraction kit following the
manufacturer's instructions. We amplified full-length 16S rRNA genes from the resulting DNA
using the 27F-1492R primer set (27F: 5'-AGAGTTTGATCMTGGCTCAG-3'; 1492R: 5'-

ACCTTGTTACGACTT-3' (56)). The reaction mix consisted of Promega's GoTaq HotStart 2x
PCR master mix with final concentrations of 0.4 μM 27F and 0.4μM 1492R primers, and 1-11.5
μl of template DNA, in a total reaction volume of 25 μL. The thermocycling profile was 1 × 94°C
for 10 min followed by 36 cycles of: 94°C for 45 s, 50°C for 90 s, 72°C for 90 s; and a single 72°C
extension for 10 min. The resulting amplicons were cleaned and Sanger sequenced from either
the 27F or 1492R primer by Eurofins Genomics (Louisville, KY, USA) using their standard
techniques. Sequences were curated using 4Peaks (57) and Geneious Prime v2019.0.1 (58).
Reads were trimmed and assembled using the moderate setting in Geneious. Those reads that
failed to assemble with these metrics were considered "mixed" cultures and not analyzed
further.
405
Culture taxonomy and determination of taxonomic differences across growth medium
formulations: High-quality full length 16S rRNA gene sequences from the cultures were used to
assign taxonomy and reconstruct phylogenetic relationships. We assigned taxonomy to all
assembled 16S rRNA gene sequences using the SILVA database SINA aligner v128 (59). A
Shapiro-Wilk test of normality was conducted in base R (60) on the distribution of taxa
recovered from both media types. After concluding the data were non-parametric, we
performed a Kruskal-Wallis test in R (Taxa ~ Media Type).
413
Taxonomic selection for phylogenetic reconstruction: To reconstruct a phylogeny of full
length 16S rRNA genes, our culture sequences were compared to NCBI's Microbial Genomes
and environmental sequences using web-blast (61). The top five hits for each sequence, from
either NCBI database, were chosen based on the highest percent coverage and lowest e-value
score. *E. coli* K-12 was used as the outgroup of the Alphaproteobacterial phylogeny, and
Bacillus subtilis was used as the outgroup for the Actinobacterial tree. These sequences aligned
with MAFFT (62) with turn checking enabled to conform sequence orientation to the outgroup.
The alignment was then trimmed using TrimAl (63) with the automated1 setting to optimize
sequence trimming for maximum likelihood phylogenetic analyses. We reconstructed
phylogenetic relationships from this trimmed alignment in the CIPRES Gateway (64).
Maximum-Likelihood (ML) trees were constructed using IQ-TREE with 10,000 ultrafast
bootstrap trees and Bayesian Information Criterion to select the best fit nucleic acid
substitution model (65, 66). After an initial round of ML trees, sequence alignments were
heuristically curated with IQ-TREE to eliminate duplicitous sequences. Finalized ML trees were
then imported into the ARB environment (67), where any duplicate sequences from our AZCC
cultures were added to the ML trees through ARB's quick add parsimony function. Final trees
were visualized with FigTree (68).
431
432   *Environmental contextualization of AZCC isolates:* We matched the AZCC isolate full length
433   16S rRNA gene sequences against a database of the clustered OTUs from the shallow soil
434   depth profile samples (see '*soil microbial community analysis,*' above) using the usearch_global
435   command (69) at a stringency of ≥97% identity, in both strand orientations, with maxaccepts=1
436   and maxrejects=0.
437

438   *Data Availability*: Full length Sanger Sequenced 16S PCR product sequences are
439 available on NCBI GenBank, accession numbers: MK875836 - MK875967. Illumina data are
440 available on the NCBI SRA, accession numbers: SRR9172130-SRR9172198.
441

Acknowledgements: The authors would like to thank Nathan Abramson, Jasper Bloodsworth,
Brenna Bourque, Amanda Howe, Bridget Taylor, and H. James Tripp for assisting with sample
collection, culture maintenance, and DNA extractions relevant to this work. Funding from this
work came from startup funds provided to PC from the University of Arizona's Technology and
Research Initiative Fund (the Water, Environmental, and Energy Solutions initiative), and seed
grants from the Center for Environmentally Sustainable Mining and The University of Arizona
College of Agriculture and Life Sciences. Work at JCVI was supported by P01AI118687.

References:
451

1.   Chaparro JM, Sheflin AM, Manter DK, Vivanco JM. 2012. Manipulating the soil
microbiome to increase soil health and plant fertility. Biol Fertil Soils 48:489–499.
2.   Long PE, Williams KH, Hubbard SS, Banfield JF. 2016. Microbial Metagenomics Reveals
Climate-Relevant Subsurface Biogeochemical Processes. Trends Microbiol 24:600–610.
3.   Fernández-Luqueño F, Valenzuela-Encinas C, Marsch R, Martínez-Suárez C, Vázquez-
Núñez E, Dendooven L. 2011. Microbial communities to mitigate contamination of PAHs
in soil-possibilities and challenges: A review. Environ Sci Pollut Res 18:12–30.
4.   Tiedje JM, Asuming-Brempong S, Nüsslein K. 1999. Opening the black box of soil
microbial diversity. Appl Soil Ecol 13:109–122.
5.   Jobbagy EG, Jackson RB. 2000. The Vertical Distribution of Soil Organic Carbon and Its
Relation to Climate and Vegetation. Ecol Appl 10:423–436.
6.   Fierer N, Schimel JP, Holden PA. 2003. Variations in microbial community composition
through two soil depth profiles. Soil Biol Biochem 35:167–176.
7.   Ajwa HA, Rice CW, Sotomayor D. 1998. Carbon and Nitrogen Mineralization in Tallgrass
Prairie and Agricultural Soil Profiles. Soil Sci Soc Am J 62:942–951.
8.   Jobbágy EG, Jackson RB. 2001. The Distribution of Soil Nutrients with Depth: Global
Patterns and the Imprint of Plants. Biogeochemistry 53:51–77.
9.   Hillel D. 2013. Fundamentals of Soil PhysicsFundamentals of Soil Physics.
10.   Brewer TE, Aronson EL, Arogyaswamy K, Billings SA, Botthoff JK, Campbell AN, Dove
NC, Fairbanks D, Gallery RE, Hart SC, Kaye J, King G, Logan G, Lohse KA, Maltz MR,
Mayorga E, O'Neill C, Owens SM, Packman A, Pett-Ridge J, Plante AF, Richter DD, Silver
WL, Yang WH, Fierer N. 2019. Ecological and genomic attributes of novel bacterial taxa
that thrive in subsurface soil horizons. bioRxiv.
11.   Eilers KG, Debenport S, Anderson S, Fierer N. 2012. Digging deeper to find unique
microbial communities: The strong effect of depth on the structure of bacterial and
archaeal communities in soil. Soil Biol Biochem 50:58–65.
12.   Gao Y, Wu M. 2018. Free-living Bacterial Communities Are Mostly Dominated by
Oligotrophs. bioRxiv.
13.   Cho J-C, Giovannoni SJ. 2004. Cultivation and Growth Characteristics of a Diverse Group
of Oligotrophic Marine Gammaproteobacteria. Appl Environ Microbiol 70:432–440.

14.   Giovannoni SJ, Cameron Thrash J, Temperton B. 2014. Implications of streamlining
483   theory for microbial ecology. ISME J 8:1553–1565.
484   15.   Lauro FM, McDougald D, Thomas T, Williams TJ, Egan S, Rice S, DeMaere MZ, Ting L,
485   Ertan H, Johnson J, Ferriera S, Lapidus A, Anderson I, Kyrpides N, Munk AC, Detter C,
486   Han CS, Brown M V., Robb FT, Kjelleberg S, Cavicchioli R. 2009. The genomic basis of
487   trophic strategy in marine bacteria. Proc Natl Acad Sci 37:15527–15533.
488   16.   Poindexter JS. 1981. Chapter 2, Oligotrophy: Fast and Famine Existence, p. 63–90. *In*
489   Advances in Microbial Ecology Volume 5.
490   17.   Eichorst SA, Kuske CR, Schmidt TM. 2011. Influence of Plant Polymers on the
491   Distribution and Cultivation of Bacteria in the Phylum Acidobacteria . Appl Environ
492   Microbiol 77:586–596.
493   18.   Noell SE, Giovannoni SJ. 2019. SAR11 Bacteria Have a High Affinity and Multifunctional
494   Glycine Betaine Transporter. Environ Microbiol 00:00–00.
495   19.   Carini P, Dupont CL, Santoro AE. 2018. Patterns of thaumarchaeal gene expression in
496   culture and diverse marine environments. Environ Microbiol 20:2112–2124.
497   20.   Santoro AE, Dupont CL, Richter RA, Craig MT, Carini P, McIlvin MR, Yang Y, Orsi WD,
498   Moran DM, Saito MA. 2015. Genomic and proteomic characterization of "*Candidatus*
499   Nitrosopelagicus brevis": An ammonia-oxidizing archaeon from the open ocean. Proc
500   Natl Acad Sci 112:1173–1178.
501   21.   Stingl U, Tripp HJ, Giovannoni SJ. 2007. Improvements of high-throughput culturing
502   yielded novel SAR11 strains and other abundant marine bacteria from the Oregon coast
503   and the Bermuda Atlantic Time Series study site. ISME J 1:361–371.
504   22.   Carini P. 2019. A "Cultural" Renaissance: Genomics Breathes New Life into an Old Craft
505   Paul. mSystems 4:e00092-19.
506   23.   Hu SJ, Van Bruggen AHC, Grünwald NJ. 1999. Dynamics of bacterial populations in
507   relation to carbon availability in a residue-amended soil. Appl Soil Ecol 13:21–30.
508   24.   Saito A, Mitsui H, Hattori R, Minamisawa K, Hattori T. 1998. Slow-growing and
509   oligotrophic soil bacteria phylogenetically close to Bradyrhizobium japonicum. FEMS
510   Microbiol Ecol 25:277–286.
511   25.   Davis KER, Joseph SJ, Janssen PH. 2005. Effects of growth medium, inoculum size, and
512   incubation time on culturability and isolation of soil bacteria. Appl Environ Microbiol
513   71:826–834.
514   26.   Joseph SJ, Hugenholtz P, Sangwan P, Osborne CA, Janssen PH. 2003. Laboratory
515   cultivation of widespread and previously uncultured soil bacteria. Appl Environ Microbiol
516   69:7210–5.
517   27.   Connon SA, Givannoni SJ. 2002. High-throughput methods for culturing
518   microorganisms in very-low-nutrient media. Appl Env Microbiol 68:3878–3885.
519   28.   Button DK, Schut F, Quang P, Martin R, Robertson BR. 1993. Viability and isolation of
520   marine bacteria by dilution culture: theory, procedures, and initial results. Appl Environ
521   Microbiol.
522   29.   Tahon G, Willems A. 2017. Isolation and characterization of aerobic anoxygenic
523   phototrophs from exposed soils from the Sør Rondane Mountains, East Antarctica. Syst
524   Appl Microbiol 40:357–369.
30.   Liu J, Li JQ, Feng L, Cao H, Cui Z. 2010. An improved method for extracting bacteria

from soil for high molecular weight DNA recovery and BAC library construction. J
Microbiol 48:728–733.

31.   Monod J. 1949. The Growth of Bacterial Cultures. Annu Rev Microbiol 3:371–394.

32.   Lloyd KG, Steen AD, Ladau J, Yin J, Crosby L. 2018. Phylogenetically Novel Uncultured
Microbial Cells Dominate Earth Microbiomes. mSystems 3:e00055-18.

33.   Sait M, Hugenholtz P, Janssen PH. 2002. Cultivation of globally distributed soil bacteria
from phylogenetic lineages previously only detected in cultivation-independent surveys.
Environ Microbiol 4:654–666.

34.   Janssen PH, Yates PS, Grinton BE, Taylor PM, Sait M. 2002. Improved culturability of soil
bacteria and isolation in pure culture of novel members of the divisions Acidobacteria,
Actinobacteria, Proteobacteria, and Verrucomicrobia. Appl Environ Microbiol 68:2391–
6.

35.   Braun PR, Al-Younes H, Gussmann J, Klein J, Schneider E, Meyer TF. 2008. Competitive
inhibition of amino acid uptake suppresses chlamydial growth: Involvement of the
chlamydial amino acid transporter BrnQ. J Bacteriol 190:1822–1830.

36.   Carini P, Steindler L, Beszteri S, Giovannoni SJ. 2013. Nutrient requirements for growth
of the extreme oligotroph "Candidatus Pelagibacter ubique" HTCC1062 on a defined
medium. ISME J 7:592–602.

37.   Tanaka T, Kawasaki K, Daimon S, Kitagawa W, Yamamoto K, Tamaki H, Tanaka M,
Nakatsu CH, Kamagata Y. 2014. A Hidden Pitfall in the Preparation of Agar Media
Undermines Microorganism Cultivability. Appl Environ Microbiol 80:7659–7666.

38.   Carlsson J, Nyberg G, Wrethén J. 1978. Hydrogen peroxide and superoxide radical
formation in anaerobic broth media exposed to atmospheric oxygen. Appl Environ
Microbiol 36:223–229.

39.   Baxter M, Sieburth JM. 1984. Metabolic and ultrastructural response to glucose of two
eurytrophic bacteria isolated from seawater at different enriching concentrations. Appl
Environ Microbiol 47:31–38.

40.   Lau JT, Whelan FJ, Herath I, Lee CH, Collins SM, Bercik P, Surette MG. 2016. Capturing
the diversity of the human gut microbiota through culture-enriched molecular profiling.
Genome Med 8:1–10.

41.   Walters W, Hyde ER, Berg-lyons D, Ackermann G, Humphrey G, Parada A, Gilbert JA,
Jansson JK, Caporaso JG, Fuhrman JA, Apprill A, Knight R. 2015. Improved Bacterial 16S
rRNA Gene (V4 and V4-5) and Fungal Internal Transcribed Spacer Marker Gene Primers.
mSystems 1:1–10.

42.   Brewer TE, Handley KM, Carini P, Gilbert JA, Fierer N. 2016. Genome reduction in an
abundant and ubiquitous soil bacterium "Candidatus Udaeobacter copiosus." Nat
Microbiol 2.

43.   Olsen RA, Bakken LR. 1987. Viability of soil bacteria: Optimization of plate-counting
technique and comparison between total counts and plate counts within different size
groups. Microb Ecol 13:59–74.

44.   Aagot N, Nybroe O, Nielsen P, Johnsen K. 2001. An Altered Pseudomonas Diversity Is
Recovered from Soil by Using Nutrient-Poor Pseudomonas-Selective Soil Extract Media.
Appl Environ Microbiol 67:5233–5239.

45.   Lennon JT, Jones SE. 2011. Microbial seed banks: The ecological and evolutionary

implications of dormancy. Nat Rev Microbiol 9:119–130.

46. Swan BK, Tupper B, Sczyrba A, Lauro FM, Martinez-Garcia M, Gonzalez JM, Luo H,
Wright JJ, Landry ZC, Hanson NW, Thompson BP, Poulton NJ, Schwientek P, Acinas SG,
Giovannoni SJ, Moran MA, Hallam SJ, Cavicchioli R, Woyke T, Stepanauskas R. 2013.
Prevalent genome streamlining and latitudinal divergence of planktonic bacteria in the
surface ocean. Proc Natl Acad Sci 110:11463–11468.

47. Batut B, Knibbe C, Marais G, Daubin V. 2014. Reductive genome evolution at both ends
of the bacterial population size spectrum. Nat Rev Microbiol 12:841–850.

48. Sorensen JW, Dunivin TK, Tobin TC, Shade A. 2019. Ecological selection for small
microbial genomes along a temperate-to-thermal soil gradient. Nat Microbiol 4:55–61.

49. Rappe MS, Connon SA, Vergin KL, Giovannoni SJ. 2002. Cultivation of the ubiquitous
SAR11 marine bacterioplankton clade. Nature 418:630–633.

50. Kim S, Kang I, Seo JH, Cho JC. 2019. Culturing the ubiquitous freshwater actinobacterial
acI lineage by supplying a biochemical 'helper' catalase. ISME J.

51. Salcher MM, Schaefle D, Kaspar M, Neuenschwander SM, Ghai R. 2019. Evolution in
action: Habitat-transition leads to genome-streamlining in Methylophilaceae. ISME J
651331.

52. Parada AE, Needham DM, Fuhrman JA. 2016. Every base matters: Assessing small
subunit rRNA primers for marine microbiomes with mock communities, time series and
global field samples. Environ Microbiol 18:1403–1414.

53. Edgar RC. 2013. UPARSE: Highly accurate OTU sequences from microbial amplicon
reads. Nat Methods 10:996–998.

54. Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, Lesniewski RA,
Oakley BB, Parks DH, Robinson CJ, Sahl JW, Stres B, Thallinger GG, Horn DJ Van, Weber
CF. 2009. Introducing mothur꞉ Open-Source , Platform-Independent , Community-
Supported Software for Describing and Comparing Microbial Communities ꞉. Appl
Environ Microbiol 75:7537–7541.

55. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glöckner FO.
2013. The SILVA ribosomal RNA gene database project: improved data processing and
web-based tools. Nucleic Acids Res 41:D590-6.

56. Lane DJ. 1991. Nucleic acid techniques in bacterial systematicsDevelopment and
Application of Nucleic Acid Probes.

57. Griekspoor A, Groothuis T. 4Peaks. 1.8. Nucleobytes, Amsterdam, Netherlands.

58. 2019. Geneious Prime. v2019.0.1. Biomatters, Ltd., Auckland, New Zealand.

59. Pruesse E, Peplies J, Glöckner FO. 2012. SINA꞉ Accurate high-throughput multiple
sequence alignment of ribosomal RNA genes. Bioinformatics 28:1823–1829.

60. R Core Team. 2014. R: A Language and Environment for Statistical Computing. 3.1.1. R
Foundation for Statistical Computing, Vienna, Austria.

61. Johnson M, Zaretskaya I, Raytselis Y, Merezhuk Y, McGinnis S, Madden TL. 2008. NCBI
BLAST: a better web interface. Nucleic Acids Res.

62. Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. Mol Biol Evol.

63. Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. 2009. trimAl: A tool for automated
alignment trimming in large-scale phylogenetic analyses. Bioinformatics 25:1972–1973.

64.   Miller MA, Pfeiffer W, Schwartz T. 2010. Creating the CIPRES Science Gateway for
615   inference of large phylogenetic treesGateway Computing Environments Workshop.
616   65.   Minh BQ, Nguyen MAT, Von Haeseler A. 2013. Ultrafast approximation for phylogenetic
617   bootstrap. Mol Biol Evol 30:1188–1195.
618   66.   Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. 2014. IQ-TREE: A Fast and Effective
619   Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. Mol Biol Evol
620   32:268–274.
621   67.   Ludwig W, Strunk O, Westram R, Richter L, Meier H, Yadhukumar, Buchner A, Lai T,
622   Steppi S, Jobb G, Förster W, Brettske I, Gerber S, Ginhart AW, Gross O, Grumann S,
623   Hermann S, Jost R, König A, Liss T, Lüßmann R, May M, Nonhoff B, Reichel B, Strehlow
624   R, Stamatakis A, Stuckmann N, Vilbig A, Lenke M, Ludwig T, Bode A, Schleifer K. 2004.
625   ARB: a software environment for sequence data. Nucleic Acids Res 32:1363–1371.
626   68.   Rambaut A, Drummond A. 2016. FigTree. 1.4.3.
627   69.   Edgar RC. 2010. Search and clustering orders of magnitude faster than BLAST.
628   Bioinformatics 26:2460–2461.

634
635
Figure 1: Dilution-to-extinction workflow. Soils are collected and brought to the lab where
they are homogenized in cell extraction buffer, layered over a Nycodenz solution, and
centrifuged (A). The cell layer is extracted from the Nycodenz solution and counted with flow
cytometry (B). Counted cells are diluted into growth medium in 96-well microtiter plates to an
average density of 5 cells well^{-1} (C). After incubation, the 96-well microtiter plates are screened
for growth with flow cytometry, and wells displaying growth are subcultured into larger
volumes (D). After incubating the subcultures, flasks displaying growth are identified by 16S
rRNA gene sequencing and molecular phylogeny (E). Aliquots of these identified subcultures
are cryopreserved at -80ºC (F).
645

646
647 **Figure 2: Microbial percent culturability (v) is greater on ASM-low than on ASM-high.** Bar
648 heights are the mean culturability ± standard deviation in 96 well microtiter plates (n=3) as
649 calculated from the initial cell inoculum and the proportion of wells positive for growth. Double
650 asterisks indicate Wilcoxon rank sum test P values ≤ 0.05.

651
652 **Figure 3: ASM-low and ASM-high cultured distinct alphaproteobacterial (a) and**
653 **actinobacterial (b) microbial taxa.** Bar heights are the number of cultures obtained for each
654 taxon and are colored by the medium type on which they were isolated.
655

Figure 4: The cultures isolated in this study are representative of several abundant soil lineages that show dynamic depth distributions in Oracle Ridge soils. Points are the mean relative abundances ± standard deviation (n=3) of 16S rRNA gene sequence OTUs that matched cultured isolates at ≥97% identity. Error bars that are not visible are located behind the symbol. *Mycobacterium* and *Arthrobacter* cultures best-matched more than one OTU in the cultivation-independent surveys. The best-matching OTU number is designated in parentheses.

Figure 5: The mean nucleic acid fluorescence of taxa isolated on ASM-low was significantly lower than for those microbes isolated on ASM-high. Points are the mean $\log_{10}$ fluorescence (in arbitrary units [AU]) of fixed and SYBR green I-stained stationary phase cultures. The mean fluorescence value was obtained from manually-gated histogram plots of fluorescence within the Guava EasyCyte software. Only those cultures that were defined as pure cultures are plotted. The horizontal line in each plot is the mean fluorescence value and the box surrounding the mean is a 95% confidence interval. Shading illustrates the relative distribution of fluorescence values within each medium type.