1 Original Article

3 Title: Leaf anatomy is not correlated to CAM function in a C_3+CAM hybrid species, *Yucca*

4 *gloriosa*

5 Authors: Karolina Heyduk[*1], Jeremy N. Ray[2], Jim Leebens-Mack[2]

6 1 - Department of Ecology and Evolutionary Biology, Yale University, New Haven, CT USA

7 2 - Department of Plant Biology, University of Georgia, Athens, GA USA

9 Running title: Intraspecific variation in leaf anatomy and physiology in a C_3+CAM hybrid

Abstract

● *Background and Aims*: CAM photosynthesis is often considered to be a complex trait,
requiring orchestration of leaf anatomy and physiology for optimal performance. But the
observation of trait correlations is based largely on comparisons between C_3 and strong
CAM species, resulting in a lack of understanding as to how such traits evolve and the
level of intraspecific variability for CAM and associated traits.
● *Methods*: To understand intraspecific variation for traits underlying CAM and how these
traits might assemble over evolutionary time, we conducted detailed time course
physiological screens and measured aspects of leaf anatomy in 24 genotypes of a
C_3+CAM hybrid species, *Yucca gloriosa* (Asparagaceae). Comparisons were made to *Y.*
gloriosa's progenitor species, *Y. filamentosa* (C_3) and *Y. aloifolia* (CAM).
● *Key results*: Based on gas exchange and measurement of leaf acids, *Y. gloriosa* appears to
use both C_3 and CAM, and varies across genotypes in the degree to which CAM can be
upregulated under drought stress. While correlations between leaf anatomy and
physiology exist when testing across all three *Yucca* species, such correlations break
down at the species level in *Y. gloriosa.*
● *Conclusions*: The variation in CAM upregulation in *Y. gloriosa* is a result of its relatively
recent hybrid origin. The lack of trait correlations between anatomy and physiology
within *Y. gloriosa* indicate that the evolution of CAM, at least initially, can proceed
through a wide combination of traits, and more favorable combinations are eventually
selected for in strong CAM plants.

Keywords: *Yucca gloriosa* L., *Yucca aloifolia, Yucca filamentosa*, CAM photosynthesis, leaf
anatomy, hybrid, Asparagaceae

Introduction

A fundamental aim of comparative biology is to elucidate how, when, and why traits

evolve, and the biological consequences of trait evolution. Some traits have simple genetic

architecture: changes may be induced by mutations to single genes or regulatory elements, as in

the case of hair color in mice (Hoekstra *et al.* 2006), flower color and pollinator shifts in

Erythranthe guttata (Bradshaw and Schemske 2003; Yuan *et al.* 2013), and herbicide resistance

in barley (Lee *et al.* 2011). Other traits are more complex, in that they are actually a sum of

phenotypic states orchestrated across an organism. For example, the evolution of C_4

photosynthesis requires both an altered biochemical pathways as well as changes to leaf vein

density and anatomy (Hatch 1987; Christin *et al.* 2013; Sage *et al.* 2014), and burrowing

behavior in field mice relies on changes to separate genetic modules (Weber *et al.* 2013).

Complex traits are unlikely to evolve via a single mutation (Lenski *et al.* 2003), and as a result

one might expect various intermediate phenotypes to exist through the evolutionary progression

from ancestral to derived character states. Species exhibiting intermediate phenotypes could be

instrumental to ordering the sequence of events that led to the evolution of a complex trait.

Intermediate phenotypes also lend insight into the genetic landscape of a complex trait: for

example, genetic linkage can restrict which trait combinations are possible and can affect how

quickly natural selection can act upon them.

Crassulacean acid metabolism (CAM) is an example of a complex plant trait - involving

biochemistry, anatomy, and physiology - whose evolutionary trajectory remains obscured,

despite multiple independent origins (Edwards 2019). CAM is a carbon concentrating

mechanism that works tandemly with the C_3 Calvin Benson cycle to increase the water use

efficiency of plants. The C_3 pathway uses Rubisco, an enzyme which has both carboxylating and

oxygenating functions. Under high temperatures or conditions that promote stomatal closure,

rates of Rubisco oxygenation increase and force plants to undergo photorespiration, an

energetically costly process that fixes no net CO_2. CAM concentrates CO_2 in an effort to reduce

oxygenation via Rubisco and photorespiratory stress. CAM species open their stomata at night,

when cooler temperatures and higher relative humidity reduce evapotranspiration. Incoming CO_2

is initially converted to malate and stored in the vacuole. During the day, the stomata largely

remain closed, and the stored CO_2 is decarboxylated from malate, surrounding Rubisco and the

C_3 machinery with elevated levels of CO_2. The carbon concentrating mechanism of CAM

reduces levels of photorespiration while simultaneously increasing overall water use efficiency.

As a result, CAM plants are often found in hot, arid, or seasonally dry habitats - often, but not

always, where water is limiting.

Since all CAM plants retain and use the entire C_3 machinery, many species fix carbon

through a mixture of both pathways (Winter 2019). Strong CAM plants use CAM for the vast

majority of their carbon uptake, while C_3+CAM species use a mix of both pathways to fix CO_2.

CAM cycling plants fix respired CO_2 nocturnally through the CAM pathway but otherwise have

C_3 physiology. Moreover, plants can vary not only in their ability to use CAM, but also the

degree to which CAM can be modulated under abiotic stress. Both strong CAM and C_3+CAM

can alter the relative contribution of CAM to CO_2 fixation as a response to abiotic stressors.

C_3+CAM species can up-regulate the CAM pathway ("facultative CAM") or downregulate the

contribution of the C_3 pathway, whereas strong CAM species may increase the degree of C_3

carbon fixation when exceptionally well-watered (Hartsock and Nobel 1976). It is unclear how

intermediate phenotypes such as C_3+CAM and CAM cycling fit into the evolutionary trajectory

of CAM, but the prevalence of intermediate CAM species (Winter 2019) suggests that such a

dynamic phenotype can be advantageous under certain situations (i.e., seasonal drought) (Winter

et al. 2008; Herrera 2009).

CAM photosynthesis has evolved at least 60 times independently (Edwards and Ogburn

2012), though this is likely an inaccurate count due to the difficulties associated with surveying

intermediate CAM, particularly facultative forms involving induction of CAM only under

specific conditions. Additionally, attempts to delineate when CAM evolved within extant CAM

lineages are made difficult by a lack of phylogenetic resolution, particularly in very diverse

lineages. However, the repeated origin of CAM suggests that the transition to CAM from a C_3

ancestor may be evolutionarily straightforward (Heyduk, Moreno-Villena, *et al.* 2019). Indeed,

the entire CAM biochemical pathway is present in all C_3 plants as a part of the tricarboxylic acid

cycle, and thus all genes required for CAM are already present in all C_3 angiosperms. Most, if

not all, genes involved in coding the CAM biochemical reactions are found in multiple copies in

angiosperm genomes, allowing a single member of a gene family to be recruited for CAM

(Silvera *et al.* 2014; Ming *et al.* 2015; Yang *et al.* 2017). There is mixed evidence that amino

acid substitutions are required for evolutionary shifts to CAM function (Yang *et al.* 2017;

Goolsby *et al.* 2018), and many recent genomic studies have suggested that changes to regulation

of gene transcription and translation are critical for CAM evolution (Heyduk *et al.* 2018; Yin *et*

al. 2018; Wai *et al.* 2019).

In addition to the importance of expressing the right genes at the right time, specific

anatomical traits have long thought to be required for maximum CAM function (Nelson *et al.*

2005; Nelson and Sage 2008). To be able to store large amounts of CO_2 as malate, CAM plants

require larger vacuoles; indeed CAM plants typically have larger mesophyll cells than their C_3

counterparts (Heyduk, Burrell, *et al.* 2016; Males 2018). Intercellular airspace (IAS) is often

reduced in CAM species (Nelson and Sage 2008; Zambrano *et al.* 2014). One theory is that tight

packing of cells reduces the amount of CO_2 leakage that can occur during the day, when malate

is decarboxylated and results in high concentrations of CO_2 in the cells. Alternatively, reduced

IAS may just be a result of larger cells packed into a leaf whose size may be limited by other

factors, including the need to maintain hydraulic connectivity. Finally, CAM plants are often

described as having thick, succulent leaves. The importance and timing of these anatomical

changes remains unclear: in some systems, species that are C_3+CAM look anatomically like their

C_3 relatives (Silvera *et al.* 2005; Males 2018), whereas other lineages evolve succulent leaf

anatomy prior to CAM (Heyduk, McKain, *et al.* 2016) or coincident with the origin of CAM

(Zambrano *et al.* 2014). As a result, our understanding of the importance of leaf anatomy on

CAM function remains unclear.

One of the greatest challenges in understanding the concerted evolution between CAM

biochemistry and anatomy is a lack of systems in which genetic segregation produces variation

within and among these traits. While comparisons between C_3 and strong CAM species have

helped us define a suite of traits that seem to segregate with photosynthetic pathway, these

comparisons conflate trait differences with evolutionary distance and yield little insight into how

suites of CAM traits have been assembled repeatedly in plant evolutionary history. Are traits

assembled sequentially, such that a certain phenotype must arise (e.g., large cells) before a

secondary phenotype can evolve (e.g., accumulation of malate)? Or are there a number of trait

combinations that can arise in any order and span phenotypic space, but selection repeatedly

favors certain combinations to maximize the efficiency of CAM? These questions can be

addressed by studying intraspecific variation of anatomy and photosynthetic physiology in an

intermediate C_3+CAM species.

To understand how anatomy and physiology are correlated - or not - we measured

anatomical and photosynthetic traits in a C_3+CAM species, *Yucca gloriosa*, a naturally occurring

homoploid hybrid species resulting from a wild cross between *Y. aloifolia* (CAM) and *Y.*

filamentosa (C_3) (Rentsch and Leebens-Mack 2012; Heyduk, Burrell, *et al.* 2016). All three

Yucca species overlap in the southeastern United States, with *Y. filamentosa* having the broadest

range that extends into the northeast and midwest, *Y. aloifolia* being more restricted to the

southeast, and *Y. gloriosa* inhabiting the narrowest natural range, occurring only in the coastal

regions of the Atlantic seaboard between Florida and Virginia. Previous work has demonstrated

contrasting photosynthetic pathways in the parental species and intermediate physiology and

anatomy in the hybrid (Heyduk, Burrell, *et al.* 2016; Heyduk, Ray, *et al.* 2019). Genetic screens

based on microsatellite data have suggested that while *Y. gloriosa* still retains a mixture of both

parental genomes, genotypes are not identical and thus not likely to be F1 hybrids (Rentsch and

Leebens-Mack 2012; Heyduk, Burrell, *et al.* 2016). With genetic variation in mind, here we

expand our previous sampling to measure gas exchange, leaf acid accumulation, and anatomical

characteristics in 24 genotypes of *Y. gloriosa*. We tested whether 1) *Y. gloriosa* genotypes vary

in physiological and anatomical traits related to CAM and in their response to drought stress, and

2) if anatomical traits are closely correlated with the photosynthetic phenotype of a genotype

(Fig. 1). We show that genotypes vary in the degree of CAM used, as well as the level of

upregulation of CAM under drought stress; we further find that there is little correlation between

anatomical traits and photosynthetic phenotypes. The lack of trait correlations within a species

suggest the processes that shape trait distributions between species are uncoupled from those

occurring within a species.

Methods

Genotypes of *Yucca gloriosa* were collected from across its range (Virginia to Florida)

(Fig. 2) [**Supplementary Information**] as ramets, then transplanted to the University of

Georgia Department of Plant Biology greenhouses. Plants were grown in 60:40 soil:sand mix

with once-weekly watering and fertilizer as needed, and maintained until proper rooting was

established and significant new growth was noticeable (minimum 6 months). Beginning in spring

2016, genotypes were randomly assigned to growth chamber experimental runs in sets of four

genotypes. For each experimental run, 3-4 clonal replicates for each of the four genotypes were

acclimated in the growth chamber for four days prior to manipulation. Growth chambers had

12hr days (beginning at 0700 h), with 30 °C/17 °C day/night temperatures and a relative

humidity of 30-40%. Light intensity at leaf level was ~400 µmol m^{-2} s^{-2} and plants were kept

well-watered during the acclimation phase. On the first experimental day ("day 1"), gas

exchange measurements were collected every two hours for a 24 hour period, beginning one

hour after lights turned on, using a LI-6400XT (LI-COR environmental). After day 1, plants

were allowed to dry down, soil moisture information was collected on experimental days [

Supplementary Information], and on day 7 plants were re-measured for gas exchange

identically to day 1. At the end of day 7, plants were re-watered, then measured a final time for

gas exchange on day 9. Experimental runs were conducted in April, September, and November

of 2016, and February, March, April, and August of 2017 on a total of 24 genotypes [

Supplementary Information]. To compare net carbon gain between genotypes, the area under

the LI-COR curve per genotype per treatment was calculated using the auc() function in the

DescTools (Signorell *et al.* 2019) package in R.

Leaf samples for titratable acidity measurements were collected on days 1, 7, and 9 two

hours before lights off and two hours before lights turned back on. Leaf punches were taken in

triplicate at both time points from all individual plants, then were immediately flash frozen and

stored at -80 °C. Leaves were quickly weighed once removed from the freezer and placed into 60

mL of 20% EtOH. Samples were boiled until the volume was reduced by half, at which point the

total volume was returned to 60 mL by adding water pH 7.0. Samples were boiled to half volume

once more, refilled to 60 mL with water, then allowed to cool to room temperature. The room

temperature liquid was cleared of leaf debris and titrated with 0.002 M NaOH to a final pH of

7.0. The umol H+ amount was calculated as (mL of 0.002 M NaOH x 2 mM)/grams of frozen

tissue.

Leaf cross sections were collected in April 2018 and April 2019 from clonal replicates of

the same genotypes measured for gas exchange, with 2 samples collected per genotype from

separate biological replicates when available. Leaves were cut, fixed in formalin, embedded in

paraffin, then sectioned at the University of Georgia Veterinary Hospital Histology Lab. Sections

were stained with Toluidine Blue and mounted on glass slides. For each of the separate plants

sectioned per genotype, 2 images were taken on a Zeiss microscope at 5 and 10x magnification,

taking care to avoid imaging edges or damaged sections. Images were analyzed in ImageJ to

collect measurements of cell size and intercellular airspace (IAS), as well as leaf thickness. For

cell size, the area of five adaxial and abaxial mesophyll cells was measured per image. IAS was

measured as a fraction of intercellular air per total cell area and is reported as a percent of

mesophyll. Leaf thickness was measured in triplicate across each image analyzed for cell size

and IAS. Stomatal density was measured by painting both adaxial and abaxial leaf surfaces with

clear nail polish (collected March 2019), then removing with tape and imaging stomatal

impressions with a Zeiss microscope. Stomatal measurements were taken from two biological

replicates per genotype, when available. Previously collected data on adaxial and abaxial cell

sizes from additional genotypes of *Y. gloriosa* was also included (Heyduk, Burrell, *et al.* 2016);

while IAS was measured on this previous dataset, due to image quality, we suspect IAS may

have been overestimated in the data previously published. IAS was therefore re-analyzed for all

data published in 2016. ANOVA or ANCOVAs were performed, as appropriate, to determine the

effect of *Y. gloriosa* genotype on phenotypic traits; in the case of CO_2 uptake and acid

accumulation, treatment (watered and drought) was included as a factor.

To compare the hybrid to the parental species, previously published data on *Yucca*

filamentosa and *Yucca aloifolia* was included as well (Heyduk, Burrell, *et al.* 2016; Heyduk,

Ray, *et al.* 2019). The parental datasets are smaller, in that a total of 5 genotypes (14 individual

plants) and 7 genotypes (16 individual plants) were measured for various traits for *Y. aloifolia*

and *Y. filamentosa,* respectively. All statistical analyses were conducted in R v. 3.5.1, and raw

data is available at www.github.com/kheyduk/Yucca_physiology. We correlated both raw data -

that is, individual plant traits - as well as genotypic means using cor.test() in R and adjusting

resulting p-values for multiple testing with the Benjamini-Hochberg correction. Correlations

were conducted pairwise on all traits, except in cases where one trait was a subset of another (for

example, nocturnal CO_2 total assimilation is a subset of total daily CO_2 assimilation).

Correlations were conducted on individual values and genotypic means of all three species

together, then separately for just *Y. gloriosa*. No correlations were tested within the parental

species, as the data pulled from earlier work did not have enough replication. For a trait

combination to be reported as significant, it had to be significant when correlated both across

individuals and across genotypic means; for significant correlations, only the across individual

statistics are reported.

Results

Gas exchange and titratable acidity - Genotypes of *Yucca gloriosa* varied in their gas exchange

patterns over a diel cycle. The majority of genotypes had some level of C_3 daytime CO_2 fixation

as well as slight nocturnal CO_2 assimilation under well-watered conditions (Fig. 3). Under

drought stress, overall responses varied. Some genotypes had a nearly total shutdown of gas

exchange during the day under drought stress, whereas others maintained non-zero levels.

Because plants dried down at slightly variable rates [**Supplementary Information**], we

examined the effect of genotype and soil moisture on CO_2 uptake: well-watered plants had only a

slightly significant effect of genotype ($F_{19,51}$=2.16, p<0.05), while drought stressed plants had a

significant effect of soil moisture ($F_{1,48}$=15.02, p<0.001). However, nocturnal CO_2 assimilation

(and thus the level of CAM performed) was not significantly related to soil moisture under either

well-watered ($F_{1,51}$=0.03, p=0.87) or drought stress ($F_{1,50}$=0.64, p=0.43). Instead, a clear GxE

signal was observed via an ANOVA (Type III) assessing interaction of genotype and treatment

(well-watered vs. drought) on nocturnal CO_2 uptake (interaction: $F_{21,131}$=2.36, p<0.01, excluding

re-watered measurements) [**Supplementary Information**]. At night, certain genotypes (e.g., 16

and 1AB, Fig. 3) were able to increase CO_2 assimilation under drought stress relative to well-

watered conditions. No hybrid genotype fully replicated the levels of nocturnal CO_2 assimilated

by *Y. aloifolia*, and many had the ability to use CAM even when well-watered, indicating *Y.*

gloriosa is not strictly facultative CAM but rather weak CAM with drought inducibility.

Nocturnal acid accumulation in the hybrid *Y. gloriosa,* like gas exchange, had a significant

interaction effect between genotype and treatment ($F_{23,120}$=3.73, p<0.001, excluding re-watered

measurements) [**Supplementary Information**]. In general, the majority of genotypes had an

increase in leaf acidity over the night period, indicative of CAM; most genotypes displayed some

level of acid accumulation on all days of the experiment, regardless of water status (Fig. 4).

Inter- and intra-specific response to drought - When compared to the parental genotypes for

which gas exchange measurements are available, many *Y. gloriosa* genotypes had some of the

highest net CO_2 assimilation (as calculated by the area under the gas exchange curves) during

both well-watered and drought conditions (Fig. 5A). However, nighttime net CO_2 assimilation

was intermediate in *Y. gloriosa* compared to parental species, and tended toward the C_3 parent *Y.*

filamentosa (Fig. 5B). When drought stressed, *Y. filamentosa* genotypes show a decrease in

overall CO_2 assimilation (Fig. 5B) under drought stress. *Yucca aloifolia* showed on average a

decrease in nighttime CO_2 assimilation under drought stress (Fig. 5AB), concordant with

previous findings (Heyduk, Burrell, *et al.* 2016). *Yucca gloriosa* genotypes varied in their gas

exchange drought response; certain genotypes increased the amount of CO_2 acquired at night,

whereas others, like *Y. aloifolia*, decreased net nighttime CO_2 acquisition.

Drought-induced response in leaf acid accumulation varied across hybrid genotypes as

well and spanned a larger phenotypic space than either parent. While *Y. filamentosa* never

accumulated significantly levels of leaf acid (well-watered: $t_5=1.71$, $p=0.07$; drought stressed:

$t_5=1.40$, $p=0.11$), *Y. aloifolia* had appreciable levels of acid accumulation over the night period

under well-watered conditions and increased the amount of acid accumulated under drought (Fig.

5B). *Yucca gloriosa* genotypes spanned the range from low levels of acid accumulation to CAM-

like levels under well-watered conditions, and genotypes varied in their ability to increase the

amount of acid stored under drought. A few genotypes responded to drought with significant and

positive increases in leaf acidity on day 7 relative to day 1 (e.g., Y13 and YG). Genotype Y18 is

a notable exception in its lack of acid accumulation and lack of response to drought stress, which

corresponds to its negligible rates of CO_2 assimilation during the dark period (Fig. 3). Genotypes

that had high levels of acid accumulation under well-watered conditions tended to decrease acid

under drought, while those that had lower level well-watered acid accumulation tended to

increase the amount of acid stored in leaves under drought stress (Fig. 5B).

Because CAM can be defined by both acid accumulation and nighttime CO_2 assimilation,

comparing the response of genotypes through both phenotypes can indicate the mode of CAM

employed under drought stress. For example, *Y. aloifolia* reduces the amount of CO_2 assimilated

at night, but typically increases leaf acid accumulation (Fig. 5D), indicating more reliance on

recycling CO_2 while drought stressed. Some genotypes of *Y. gloriosa* decreased reliance on

atmospheric CO_2 and increase acid accumulation with drought stress (upper left quadrant, Fig.

5D). Others responded to drought by increasing both nighttime CO_2 assimilation and leaf acid

accumulation (upper right quadrant, Fig. 5D). A few genotypes were negatively impacted by

drought in that they reduced both leaf acid accumulation and nighttime CO_2 uptake, such that

stress appears to have diminished their CAM capacity (lower left quadrant, Fig. 5D). Finally, a

few genotypes appear to increase the amount of nighttime CO_2 assimilation but *decrease* the

level of acid stored in the leaves (lower right quadrant, Fig. 5D); however in many of these latter

cases the error bars overlap zero, and therefore we cannot reject the expectation that nighttime

CO_2 uptake is coupled with acid accumulation in these genotypes. Regardless, the general

diversity of drought responses in the hybrid *Y. gloriosa* is clear.

Leaf anatomy - All anatomical traits were significantly different between species, based

on ANOVA (Benjamini-Hochberg corrected p-values) [**Supplementary Information**], with

the exception of abaxial stomatal density. Within *Y. gloriosa*, the only anatomical traits

significantly different between genotypes were IAS ($F_{25,21}=3.36$, $p<0.001$) and mean stomatal

density (averaged abaxial and adaxial values) ($F_{4,13}=3.73$, $p<0.01$) [**Supplementary**

Information]. As with physiological traits, anatomical differences between *Y. aloifolia* and *Y.*

filamentosa were stark, while the hybrid largely filled the phenotypic space between. Cell sizes

on both adaxial and abaxial sides of the leaf were larger in *Y. aloifolia* than in either of the other

two species (Fig. 5A). Stomatal density, conversely, was lowest on average in *Y. aloifolia* and

greatest in *Y. filamentosa* (Fig. 6B). Cell sizes and stomatal densities on adaxial and abaxial sides

of the leaf were highly positively correlated (Fig. 6AB) across all individuals (cell size:

$t_{67}=25.19$, $R^2= 0.90$, $p<0.001$; stomata: $t_{46}=7.12$, $R^2=0.52$, $p<0.001$).

Few anatomical traits could predict physiological traits across species (Fig. 7) [

Supplementary Information]. Total CO_2 assimilation across the whole day under both water

and drought stress was correlated to IAS, albeit with a relatively low R^2 in both cases (Fig. 7A).

The amount of nocturnal CO_2 assimilated under both watered and drought conditions had

multiple trait correlations to both other physiological traits as well as leaf anatomy. Nocturnal

CO_2 uptake was correlated to IAS, levels of acid accumulation (both under watered and drought

conditions), the maximum amount of acid held within a leaf at any time point, the mean cell size,

and leaf thickness (Fig. 7B-G). The amount of leaf acids accumulated under drought stress was

correlated to total nocturnal CO_2 assimilation under drought stress ($R^2=0.14$, $p<0.01$). Within *Y.*

gloriosa, nearly all the trait correlations were not significant [**Supplementary Information**].

The only significant correlations for traits in *Y. gloriosa* were between mean cell size and leaf

thickness ($R^2=0.57$, $p<0.001$) and between total CO_2 assimilation under water and drought

conditions ($R^2=0.24$, $p<0.001$).

Discussion

Detailed physiological and anatomical measurements in *Y. gloriosa* have revealed

among-genotype variation in CAM phenotypes, and that anatomical and physiological traits

show a lack of correlation within *Y. gloriosa*. Under drought stress, the levels of daytime CO_2

assimilation were largely driven by environment - that is, soil moisture content - whereas

nocturnal CO_2 assimilation rates and acid accumulation were influenced by a combination of

genotype and environmental effects. Our results reveal a continuum of photosynthetic traits

across *Y. gloriosa* genotypes, including variation in drought response. Anatomical measurements

were largely not predictive of physiological traits within *Y. gloriosa*. In contrast, cell size, IAS,

and leaf thickness were predictive of nocturnal CO_2 uptake in cross species comparisons. These

observations suggest that anatomical characteristics can be decoupled from photosynthetic

physiology of CAM within the homoploid hybrid species *Y. gloriosa.*

Intra- and interspecific variation in anatomy and physiology

The few studies that have linked leaf anatomy to CAM photosynthetic capacity have

provided often contrasting results on how important various anatomical traits are for CAM. In a

study that compared phylogenetically unrelated strong CAM and C_3+CAM species, cell size was

found to be unrelated to CAM (Nelson and Sage 2008). In contrast, a study of *Clusia* species that

ranged from C_3 to CAM with intermediates showed palisade mesophyll cell size was

significantly correlated to the proportion of CO_2 uptake at night (Zambrano *et al.* 2014). Across

the three *Yucca* species examined here, cell size (area) was related to nocturnal CO_2 uptake under

both watered and drought conditions, although such a relationship did not exist at the

intraspecific level within *Y. gloriosa*. All studies use cell size as a proxy for vacuolar size, which

in theory would limit the storage capacity of malate. It is possible that vacuolar size is not

linearly related to cell size (though see Chan and Marshall 2014), and that inconsistent results on

the importance of cell size between studies is related to using anatomical proxies for the true trait

of interest. Alternatively, and probably more likely, studies that control for phylogenetic

distance, such as this one and those conducted across *Clusia* species, reduce noise introduced by

sampling across evolutionarily distant lineages and may provide a more accurate assessment of

anatomical importance.

In addition to cell size, intercellular air space (IAS) is often cited as a critical trait for

CAM, though whether it evolves as a byproduct of tight cell packing (Maxwell *et al.* 1997) or as

a way to reduce CO_2 remains unclear (Borland *et al.* 2018). IAS was strongly correlated to

strength of CAM when measured across unrelated CAM and C_3+CAM species (Nelson and Sage

2008), but had little role in determining strength of CAM when assessed within the genus *Clusia*

(Zambrano et al. 2014). IAS was correlated to nocturnal CO_2 assimilation when tested across all

three species of *Yucca*, but was not correlated to leaf acid accumulation, and showed no

relationship to *any* other traits within *Y. gloriosa*. That IAS is not predictive of physiology in *Y.*

gloriosa - neither nocturnal CO_2 uptake or the amount of leaf acids that accumulate - is

surprising, given that contrasts between C_3 to CAM species have repeatedly shown the latter

have significantly reduced IAS (Heyduk, McKain, *et al.* 2016; Heyduk, Burrell, *et al.* 2016;

Males 2018). Together, the IAS trends across and within *Yucca* species shows that while IAS

may be required for constitutive CAM, there exists a large intermediate space where IAS

predicts very little about photosynthetic functionality.

For many anatomical and physiological traits, *Y. gloriosa* has intermediate values

compared to the two parental species and often occupies a much broader range of trait values. It

is possible that our limited sampling of the parental species, drawn from previous work, reduces

our ability to accurately assess trait space in *Y. aloifolia* and *Y. filamentosa*. However, multiple

genotypes were sampled across the ranges of both parental species, and thus the greater variation

found within the hybrid is likely due to genomic mixing, rather than sampling bias. While trait

values in *Y. gloriosa* were typically intermediate, one notable exception was the transgressive

values of total CO_2 assimilation under both watered and drought stressed conditions. Due to *Y.*

gloriosa's nearly C_3-level of daytime CO_2 fixation coupled with the ability to use low level

CAM, total CO_2 uptake rates far exceed that of either parent, at least in certain genotypes. Such a

mixed photosynthetic strategy may be particularly valuable on the coastal dunes that *Y. gloriosa*

is restricted to, as although rainfall in the southeastern U.S. is not particularly limiting, any water

that does fall likely percolates through the sandy substrate quickly.

Despite the potentially novel phenotypes that *Y. gloriosa* exhibits relative to its parental

species, they are unlikely to underlie the speciation of the hybrid from its progenitors. All three

Yucca species are found across the southeastern coast, although only *Y. aloifolia* and *Y. gloriosa*

grow with any frequency on the coastal dunes. *Y. filamentosa* is typically further away from the

ocean in the coastal scrub, though can be found in exposed sand near brackish inlets (K Heyduk,

unpubl. res.). Homoploid hybrid species can be formed and maintained either through

chromosomal structural rearrangements that form a reproductive barrier between the new species

and its progenitors, or via ecological differentiation, whereby the new combination of traits in the

hybrid allows for habitation of a novel niche relative to the parental species (Gross and

Rieseberg 2005). As the habitats of the *Yucca* species studied here largely overlap, particularly *Y.*

gloriosa and *Y. aloifolia*, the latter at first seems unlikely, despite *Y. gloriosa* being clearly

distinct in total CO_2 assimilation rates. However, flowering time of the three species is markedly

different: *Y. filamentosa* typically flowers earliest, in late May and June, followed by *Y. aloifolia*.

Yucca gloriosa has been noted to flower at the end of the summer and into autumn (Trelease

1902); whether the later flowering time was selected for in order to reduce backcrossing, or was

instead a byproduct of the initial hybridization events, remains unknown. Additionally, other

biotic interactions (e.g., below-ground mutualisms or pollinators) or microhabitat variation are

largely untested as potential drivers of *Yucca* speciation (but see Rentsch and Leebens-Mack

2014). Chromosomal structural rearrangements may explain an apparent lack of backcrossed

individuals, but we do not currently have the genomic data to test this hypothesis.

Abiotic stress regulation of CAM

Genotypes of *Y. gloriosa* used variable levels of CAM, and differentially up-regulated

CAM under drought stress. The differential drought response was a result of two separate axes of

the CAM phenotype: both leaf acid accumulation and nocturnal CO_2 uptake varied by

environment, and could do so in a de-coupled manner (Fig. 5). That is, certain genotypes

increased the amount of leaf acids accumulated based not on increasing atmospheric CO_2 uptake

but instead by presumably re-fixing respired CO_2. Such a response indicates that much of the

required enzymes are present and regulated correctly, but that stomatal aperture responded

negatively to drought at night. Reducing net CO_2 uptake but increasing leaf acid accumulation is

the typical response of *Y. aloifolia* to drought stress as well. In general, the response to drought

stress in *Y. gloriosa* was transgressive relative to parental phenotypes, in that genotypes of *Y.*

gloriosa were able to respond to drought stress in ways that neither parent could. For example,

certain genotypes could *increase* both CO_2 uptake and leaf acid accumulation under drought

stress - this response was not seen in any of the parental genotypes measured here. Other

genotypes occupied a part of trait space where nocturnal CO_2 uptake increased under drought

stress, but leaf acids decreased (Fig. 5D). How incoming CO_2 is processed in these genotypes

remains unclear and warrants additional exploration in these genotypes, especially through

metabolomic and genomic analyses to help pinpoint alternative pathways.

The segregation of CAM drought response in *Y. gloriosa* also presents an ideal system

with which to interrogate the molecular components of drought response in facultative CAM

species. CAM has been touted as a potential trait for increasing food and biofuel crop drought

tolerance through bioengineering (Borland *et al.* 2014, 2015), and early efforts to transform C_3

species to CAM were instrumental in generating an abundance of genomics data for CAM

species. Yet fully committing a C_3 plant to CAM may result in costs that outweigh any gains in

drought tolerance; larger leaves and cells will require more energy and time to produce, and

constitutive CAM usage is not ideal when drought may be intermittent. Instead, drought

tolerance engineering efforts should look to facultative CAM or C_3+CAM, as in *Y. gloriosa*,

which outperforms its parental species in terms of total CO_2 uptake and may result in faster

biomass gains, though this remains to be tested. The natural variation for CAM induction and up-

regulation in *Y. gloriosa*, as well as the uncoupling of various CAM traits, including anatomy,

acid accumulation, and CO_2 uptake, make *Y. gloriosa* ideal for investigating the molecular basis

of particular CAM traits and their regulation via abiotic signaling.

Implications for the evolution of CAM

While *Y. gloriosa* is a hybrid and therefore represents a somewhat atypical avenue for

trait evolution, it still allows us a glimpse into how a trait like CAM might be assembled. The

homoploid nature of *Y. gloriosa* means that the genomic content of the two parental species is

not highly divergent, and that the mix of traits found in *Y. gloriosa* genotypes are not a result of a

highly perturbed genome but more like what may be expected of an intraspecific cross between

phenotypically divergent parents. The mixture of traits within *Y. gloriosa* allows us to speculate

on the genomic architecture underlying the CAM phenotype. It seems unlikely that many of the

traits are genetically linked - that is, the few relationships between traits within *Y. gloriosa*

means the underlying genes are dispersed in physical genomic location and that they are not

necessarily expressed in or regulated by similar pathways. For example, there is nothing in the

genome of *Y. gloriosa* that requires large cells to develop low IAS (or vice versa), or that CAM

activity is in any way linked genetically to leaf thickness. The variation in and lack of association

between traits in *Y. gloriosa* also implies, unsurprisingly, that the CAM phenotype is highly

quantitative, and that recombination can break up many of the underlying traits. The genetic

architecture of CAM does not fully explain why such a mix of traits has remained in *Y. gloriosa*.

The frequent dry down of sandy coastal dunes may be promoting the maintenance of

intermediate traits as a way of mediating a highly variable environment. Alternatively, *Y.*

gloriosa is not a particularly rare species in its native habitat, but its populations are small and

relatively isolated. In such small populations, selection has a weaker effect than drift, which can

lead to less advantageous combinations of traits persisting in a species. Additional research using

common gardens could facilitate our understanding of whether intermediate traits like those

found in *Y. gloriosa* can confer a fitness advantage in some circumstances.

The variation and lack of correlation between traits underlying the CAM phenotype in *Y.*

gloriosa also give insight into how CAM is assembled over evolutionary time. While certain

traits appear fixed when we examine strong C_3 and CAM species, intermediate species are

important for understanding the processes that may have led to trait fixation and correlation

across traits. After all, selection acts not on the species level, but on individual, and indeed there

is a broad phenotypic space within *Y. gloriosa* for the traits examined here that selection could

act upon. That selection seems to recurrently end up on a particular anatomical phenotype in

CAM species - i.e., larger cells, thicker leaves - despite no genetic constraint for such a

correlation suggests there is an optimal combination of traits for CAM efficiency. The pattern of

convergent evolution of trait combinations, paired with intermediate species showing highly

variable trait combinations, implies a funnel shape to the evolutionary trajectory of CAM.

Species can use a degree of CAM without committing to any particular leaf anatomy, meaning

that initial transitions to using C_3+CAM following broad and varied routes. This is in contrast to

the evolution of C_4 photosynthesis, which, like CAM, requires specific anatomical

characteristics. In C_4 lineages, anatomical changes occur prior to the evolution of C_4

biochemistry (McKown and Dengler 2007; Lundgren *et al.* 2019); in some cases, like the

PACMAD clade of grasses, these anatomical changes happen early enough in evolutionary time

that they are thought to have facilitated repeated origins of C_4 (Christin *et al.* 2013). In contrast,

"weak" CAM or C_3+CAM has no anatomical constraints in *Yucca*. There is, however, an upper

bound where further investment in CO_2 fixation by the CAM pathway requires dedicated

anatomy, though the exact threshold of that transition point remains unclear. The funnel shape to

the evolution of CAM, whereby no anatomical constraints impact low levels of CAM function,

means that ordering of events on the evolutionary trajectory from C_3 to CAM will be

exceedingly difficult, as lineages can take various routes through the intermediate zone.

While the lack of correlated traits in an intermediate C_3+CAM hybrid species has

implications for broader questions on the evolution of CAM, future work can elaborate upon the

patterns seen here and help assess how generalizable these results are. Sampling of parental

genotypes and overall range was limited, and thus there may exist greater variation among traits

in the parental C_3 and CAM species as well. Indeed, most studies that examine the correlation of

anatomy to photosynthetic physiology do not sample intraspecific variation, and therefore it

remains a largely unexplored area of CAM. Growth conditions used in this study were based on

previous work in the *Yucca* system, but modulating those conditions may reveal deeper levels of

variation across environmental gradients. Finally, the *Yucca* hybrid system is a single example of

intermediacy between C_3 and CAM, and other C_3+CAM species should continue to be examined

via detailed physiology and anatomy to advance fundamental understanding of how CAM

evolves. Investigations within and between species exhibiting a mix of CAM, C_3, and

intermediate species will continue to provide insights into whether the decoupling of CAM traits

we observe in a hybrid species holds more generally.

Conclusions

Comparisons between C_3 and CAM species have suggested suites of traits are correlated

to maximize the efficiency of each photosynthetic pathway: CAM species have large cells for

storing malate, and the cells are often packed together densely in large, thick leaves to minimize

CO_2 leakage back into the atmosphere; C_3 plants have large amounts of airspace between

significantly smaller cells to facilitate the diffusion of CO_2 to the sites of Rubisco carboxylation.

These trends have been seen repeatedly in independent CAM lineages, but few studies have

examined intermediate C_3+CAM plants, and even fewer have assessed intraspecific variation for

trait correlations. The C_3+CAM hybrid *Y. gloriosa* examined here not only has a greater range of

traits than either of its parental species, but it also lacks many of the trait correlations commonly

associated with the ability to use CAM. Indeed, not a single leaf anatomical trait could predict

the amount of CO_2 acquired via CAM in the hybrid species. The lack of variation within the

intermediate *Y. gloriosa* suggests that the evolutionary trajectory to CAM from C_3 passes

through a stage where many combinations of anatomical and photosynthetic physiology traits

can coexist in a single plant. Furthermore, in *Yucca* at least, anatomical and physiological traits

are not genetically linked, and supports existing hypotheses that suites of leaf traits found

repeatedly in CAM species have been selected for in order to maximize photosynthetic

efficiency.

Funding

This work was supported by the Division of Environmental Biology at the National Science

Foundation [#1442199 to J.L-M.] and the Donnelley Postdoctoral Fellowship at Yale University

[to K.H.].

Acknowledgements

The authors would like to thank the state parks of Georgia, Florida, South Carolina, North

Carolina, and Virginia, as well as the National Park Service, for their assistance in collection and

permitting. Special thanks to Nida Moledina, Dan Tepstov, Grace Manning, Rushi Patel, and

Charmi Patel for measuring thousands of titration samples, the greenhouse staff at UGA

(particularly Mike Boyd and Greg Cousins) for granting us growth chamber access, and Rick

Field and Amanda Cummings for assistance in collecting anatomical samples.

1   **Figure 1**

3   **Figure 1** - Anatomical traits typically associated with C_3 and strong CAM plants, showing

4   possible resulting trait associations in a hybrid between a C_3 and a CAM species.

1   **Figure 2**

3   **Figure 2** - Collection origin of samples used in this study for each of the three *Yucca* species,

4   plotted against mean annual temperature from the Worldclim database. Points are jittered so they

5   do not overlap, see [**Supplementary Information**] for full locality information.

Figure 3

Figure 3 - A) Gas exchange of *Y. gloriosa* genotypes measured every two hours over a 24 hour

period beginning at 1 hour after lights on (8 a.m.). White and grey backgrounds specify daytime

and nighttime measurements, respectively. Mean and standard deviation are shown for days 1

(well watered), 7 (drought stress), and 9 (re-watered). Four samples were omitted due to

potentially inaccurate LiCOR measurements (genotypes 51, 55, 61, and 70).

Figure 4

Figure 4 - Levels of leaf titratable acidity (AM H+ equivalents - PM H+ equivalents to pH 7.0)

across well-watered (D1), drought (D7), and re-watered (D9) time points in 24 genotypes of

Yucca gloriosa. If any given value is not significantly different from 0 (no acid accumulation),

N.S. is shown above the box. Significant difference between time points within a genotype are

indicated by brackets above the boxes.

Figure 5

Species: ● *Y. aloifolia* ◐ *Y. gloriosa* ○ *Y. filamentosa*

Figure 5 - Physiological effect of drought stress on genotypic means (and standard deviation)

across all three species. A) The estimated total CO_2 assimilation, based on the area under the

LiCOR curves across the entire diel cycle, for both hybrid and parental genotypes under well-

watered and drought-stressed conditions. B) The estimated total CO_2 assimilation based on the

area under the LiCOR curves at night only, for both hybrid and parental genotypes under well-

watered and drought-stressed conditions. C) Leaf acid accumulation under well-watered conditions (W) versus drought stressed conditions (D), with genotypic mean and one standard deviation. Dashed line indicates equal values under both conditions; genotypes that fall above or below indicate greater or lower amounts of acid, respectively, were accumulated under drought stress than under well-watered. D) Comparison of the change in nighttime CO_2 assimilation induced by drought stress (xaxis) to the change in leaf acid accumulation induced by drought (yaxis). Quadrants are labeled with the phenotype observed.

Figure 6

Species: ● *Y. aloifolia* ◉ *Y. gloriosa* ○ *Y. filamentosa*

Figure 6 - Mean values of cell sizes (A) and stomatal densities (B) on adaxial and abaxial sides of the leaf per individual plant. In both cases, the dashed line is the regression line from the lm() function in R. R^2 and p-value are reported based on correlation tests in R.

Figure 7

Figure 7 - Scatterplots and regression lines with R^2 and p-values for a subset of traits [

Supplementary Information]. Individual data per plant are plotted, and correlations are shown

for individual plant data, rather than genotypic means [**Supplementary Information**]. A) Total

CO_2 assimilation under watered and drought plotted against intercellular airspace (IAS). B-G)

Nocturnal CO_2 assimilation under watered and drought plotted against IAS (B), mean cell area

(C), leaf thickness (D), leaf acid accumulation under watered (E) and drought (F) treatments, and

against the maximum amount of acid present at any time point (G).

References

Borland AM, Hartwell J, Weston DJ, *et al.* **2014**. Engineering crassulacean acid metabolism to improve water-use efficiency. *Trends in plant science* **19**: 327–338.

Borland AM, Leverett A, Hurtado-Castano N, Hu R, Yang X. **2018**. Functional Anatomical Traits of the Photosynthetic Organs of Plants with Crassulacean Acid Metabolism In: Adams WW III, Terashima I, eds. *The Leaf: A Platform for Performing Photosynthesis*. Cham: Springer International Publishing, 281–305.

Borland AM, Wullschleger SD, Weston DJ, *et al.* **2015**. Climate-resilient agroforestry: physiological responses to climate change and engineering of crassulacean acid metabolism (CAM) as a mitigation strategy. *Plant, cell & environment* **38**: 1833–1849.

Bradshaw HD, Schemske DW. **2003**. Allele substitution at a flower colour locus produces a pollinator shift in monkeyflowers. *Nature* **426**: 176–178.

Chan Y-HM, Marshall WF. **2014**. Organelle size scaling of the budding yeast vacuole is tuned by membrane trafficking rates. *Biophysical journal* **106**: 1986–1996.

Christin P-A, Osborne CP, Chatelet DS, *et al.* **2013**. Anatomical enablers and the evolution of C4 photosynthesis in grasses. *Proceedings of the National Academy of Sciences of the United States of America* **110**: 1381–1386.

Edwards EJ. **2019**. Evolutionary trajectories, accessibility, and other metaphors: the case of C4 and CAM photosynthesis. *The New phytologist*.

Edwards EJ, Ogburn RM. **2012**. Angiosperm Responses to a Low-CO 2 World: CAM and C 4 Photosynthesis as Parallel Evolutionary Trajectories. *International journal of plant sciences* **173**: 724–733.

Goolsby EW, Moore AJ, Hancock LP, De Vos JM, Edwards EJ. **2018**. Molecular evolution of key metabolic genes during transitions to C4 and CAM photosynthesis. *American journal of botany* **105**: 602–613.

Gross BL, Rieseberg LH. **2005**. The ecological genetics of homoploid hybrid speciation. *The Journal of heredity* **96**: 241–252.

Hartsock TL, Nobel PS. **1976**. Watering converts a CAM plant to daytime CO2 uptake. *Nature* **262**: 574–576.

Hatch MD. **1987**. C4 photosynthesis: a unique blend of modified biochemistry, anatomy and ultrastructure. *Biochimica et Biophysica Acta (BBA) - Reviews on Bioenergetics* **895**: 81–106.

Herrera A. **2009**. Crassulacean acid metabolism and fitness under water deficit stress: if not for carbon gain, what is facultative CAM good for? *Annals of botany* **103**: 645–653.

Heyduk K, Burrell N, Lalani F, Leebens-Mack J. **2016**. Gas exchange and leaf anatomy of a C3-CAM hybrid, Yucca gloriosa (Asparagaceae). *Journal of experimental botany* **67**: 1369–1379.

Heyduk K, Hwang M, Albert V, *et al.* **2018**. Altered Gene Regulatory Networks Are Associated With the Transition From C3 to Crassulacean Acid Metabolism in Erycina (Oncidiinae: Orchidaceae).

Frontiers in plant science **9**: 2000.

Heyduk K, McKain MR, Lalani F, Leebens-Mack J. **2016**. Evolution of CAM anatomy predates the origins of Crassulacean acid metabolism in the Agavoideae (Asparagaceae). *Molecular phylogenetics and evolution* **105**: 102–113.

Heyduk K, Moreno-Villena JJ, Gilman IS, Christin P-A, Edwards EJ. **2019**. The genetics of convergent evolution: insights from plant photosynthesis. *Nature reviews. Genetics* **2019**.

Heyduk K, Ray JN, Ayyampalayam S, *et al.* **2019**. Shared expression of Crassulacean acid metabolism (CAM) genes predates the origin of CAM in the genus Yucca. *Journal of experimental botany*.

Hoekstra HE, Hirschmann RJ, Bundey RA, Insel PA, Crossland JP. **2006**. A single amino acid mutation contributes to adaptive beach mouse color pattern. *Science* **313**: 101–104.

Lee H, Rustgi S, Kumar N, *et al.* **2011**. Single nucleotide mutation in the barley acetohydroxy acid synthase (AHAS) gene confers resistance to imidazolinone herbicides. *Proceedings of the National Academy of Sciences of the United States of America* **108**: 8909–8913.

Lenski RE, Ofria C, Pennock RT, Adami C. **2003**. The evolutionary origin of complex features. *Nature* **423**: 139–144.

Lundgren MR, Dunning LT, Olofsson JK, *et al.* **2019**. C4 anatomy can evolve via a single developmental change. *Ecology letters* **22**: 302–312.

Males J. **2018**. Concerted anatomical change associated with crassulacean acid metabolism in the Bromeliaceae. *Functional plant biology: FPB* **45**: 681–695.

Maxwell K, Caemmerer S von, Evans JR. **1997**. Is a Low Internal Conductance to CO2 Diffusion a Consequence of Succulence in Plants with Crassulacean Acid Metabolism? *Australian journal of plant physiology* **24**: 777.

McKown AD, Dengler NG. **2007**. Key innovations in the evolution of Kranz anatomy and C4 vein pattern in Flaveria (Asteraceae). *American journal of botany* **94**: 382–399.

Ming R, VanBuren R, Wai CM, *et al.* **2015**. The pineapple genome and the evolution of CAM photosynthesis. *Nature genetics* **47**: 1435–1442.

Nelson EA, Sage RF. **2008**. Functional constraints of CAM leaf anatomy: tight cell packing is associated with increased CAM function across a gradient of CAM expression. *Journal of experimental botany* **59**: 1841–1850.

Nelson EA, Sage TL, Sage RF. **2005**. Functional leaf anatomy of plants with crassulacean acid metabolism. *Functional plant biology: FPB* **32**: 409.

Rentsch JD, Leebens-Mack J. **2012**. Homoploid hybrid origin of Yucca gloriosa: intersectional hybrid speciation in Yucca (Agavoideae, Asparagaceae). *Ecology and evolution* **2**: 2213–2222.

Rentsch JD, Leebens-Mack J. **2014**. Yucca aloifolia (Asparagaceae) opts out of an obligate pollination mutualism. *American journal of botany* **101**: 2062–2067.

Sage RF, Khoshravesh R, Sage TL. **2014**. From proto-Kranz to C4 Kranz: building the bridge to C4 photosynthesis. *Journal of experimental botany* **65**: 3341–3356.

Signorell A. **2019**. DescTools: Tools for Descriptive Statistics.

Silvera K, Santiago LS, Winter K. **2005**. Distribution of crassulacean acid metabolism in orchids of
Panama: evidence of selection for weak and strong modes. *Functional plant biology: FPB* **32**: 397.

Silvera K, Winter K, Rodriguez BL, Albion RL, Cushman JC. **2014**. Multiple isoforms of
phosphoenolpyruvate carboxylase in the Orchidaceae (subtribe Oncidiinae): implications for the evolution
of crassulacean acid metabolism. *Journal of experimental botany* **65**: 3623–3636.

Trelease W. **1902**. The yucceae. *Missouri Botanical Garden Annual Report* **1902**: 27–133.

Wai CM, Weise SE, Ozersky P, Mockler TC, Michael TP, VanBuren R. **2019**. Time of day and
network reprogramming during drought induced CAM photosynthesis in Sedum album. *PLoS genetics*
15: e1008209.

Weber JN, Peterson BK, Hoekstra HE. **2013**. Discrete genetic modules are responsible for complex
burrow evolution in Peromyscus mice. *Nature* **493**: 402–405.

Winter K. **2019**. Ecophysiology of constitutive and facultative CAM photosynthesis. *Journal of*
experimental botany.

Winter K, Garcia M, Holtum JAM. **2008**. On the nature of facultative and constitutive CAM:
environmental and developmental control of CAM expression during early growth of Clusia, Kalanchöe,
and Opuntia. *Journal of experimental botany* **59**: 1829–1840.

Yang X, Hu R, Yin H, *et al.* **2017**. The Kalanchoë genome provides insights into convergent evolution
and building blocks of crassulacean acid metabolism. *Nature communications* **8**: 1899.

Yin H, Guo H-B, Weston DJ, *et al.* **2018**. Diel rewiring and positive selection of ancient plant proteins
enabled evolution of CAM photosynthesis in Agave. *BMC genomics* **19**: 588.

Yuan Y-W, Sagawa JM, Young RC, Christensen BJ, Bradshaw HD Jr. **2013**. Genetic dissection of a
major anthocyanin QTL contributing to pollinator-mediated reproductive isolation between sister species
of Mimulus. *Genetics* **194**: 255–263.

Zambrano VAB, Lawson T, Olmos E, Fernández-García N, Borland AM. **2014**. Leaf anatomical
traits which accommodate the facultative engagement of crassulacean acid metabolism in tropical trees of
the genus Clusia. *Journal of experimental botany* **65**: 3513–3523.

28