Protein yield is tunable by synonymous codon changes of translation initiation sites

Bikash K. Bhandari[1,†], Chun Shen Lim[1,†], Daniela M. Remus[3], Augustine Chen[1], Craig van
Dolleweerd[3], Paul P. Gardner[1,2,*]

[1]Department of Biochemistry, School of Biomedical Sciences, University of Otago, Dunedin,
New Zealand
[2]Biomolecular Interaction Centre, University of Canterbury, Christchurch, New Zealand
[3]Callaghan Innovation Protein Science and Engineering, University of Canterbury, Christchurch,
New Zealand

[†]These authors contributed equally.
*Corresponding author. Email: paul.gardner@otago.ac.nz

Short Title: Tunable recombinant protein expression

ABSTRACT
Recombinant protein production is a widely used technique, yet half of these experiments fail at
the expression phase. Failures are expected for 'difficult-to-express' proteins, but for others,
codon bias and mRNA folding have been proposed explanations for variable protein
abundance. We question how practical these features are for solving protein expression
failures. We discover that the energetics of RNA structure ensembles, which models the
'accessibility' of translation initiation sites, show the strongest correlation with protein
abundance across species. Importantly, accessibility outperforms other features when predicting
the outcomes of 11,430 recombinant protein production experiments in *Escherichia coli*.
Furthermore, we show that protein level is tunable by synonymous codon changes of the first
few codons that alter accessibility.

INTRODUCTION
Recombinant protein expression has numerous applications in biotechnology and biomedical
research. Despite extensive refinements in protocols over the past three decades, half of the
experiments fail in the expression phase (http://targetdb.rcsb.org/metrics/). Notable problems
are the low expression of 'difficult-to-express' proteins such as those found in, or associated
with, membranes, and the poor growth of the expression hosts, which may relate to toxicity of
heterologous proteins (*1*) (see (*2, 3*) for detailed reviews). Despite these issues, mRNA
abundance can only explain up to 40% of the variation in protein abundance, due to the
complexity of translation and turnover of biomolecules (*4–10*). Furthermore, strong promoters
used in expression vectors do not always lead to a desirable level of protein expression
because of leaky expression (*3*).

For *Escherichia coli*, mainstream models that may explain the lower-than-expected correlation
between mRNA and protein levels are codon-usage and mRNA structure. Codon analysis is

based on the frequency of codon usage in highly expressed proteins using codon adaptation
index (CAI) (*11*) or tRNA adaptation index (tAI) (*12*, *13*), whereas mRNA folding analysis
predicts the stability of mRNA secondary structures. Codon usage bias is thought to correlate
with tRNA abundance, translation efficiency and protein production (*11–17*) but its usefulness
has been questioned (*18–21*). More recent studies show stronger support for models based on
mRNA folding, in which the stability of RNA structures around the Shine-Dalgarno sequence
and translation initiation sites inversely correlates with protein expression (*18*, *19*, *21–24*). We
recently proposed a third model in which the avoidance of inappropriate interactions between
mRNAs and non-coding RNAs has a strong effect on protein expression (*25*). The roles of these
models in protein expression is an active area of research.

The algorithms for gene optimisation sample synonymous protein-coding sequences using
'fitness' models based on CAI, tAI, mRNA folding, and/or G+C content (%) (*26–30*). However,
these 'fitness' models are usually based on some of the above findings that rely on either
endogenous proteins, reporter proteins, or a few heterologous proteins with their synonymous
variants. It is unclear whether these features are generalisable to explain the expression of all
heterologous proteins. To address this question, we studied multiple large datasets across
species in order to extract features that allow us to predict the outcomes of 11,430 experiments
of recombinant protein expression in *E. coli*. With this information, we propose how such
features can be exploited to fine-tune protein expression at a low cost.

RESULTS
Accessibility of translation initiation sites strongly correlates with protein abundance
To identify a better energetic model for mRNA structure that explains protein expression, we
examined an *E. coli* expression dataset of green fluorescent protein (GFP) fused in-frame with a
library of 96-nt upstream sequences (N=244,000) (*21*). We removed the redundancy of these
96-nt upstream sequences by clustering on sequence similarity, giving rise to 14,425
representative sequences. We calculated the accessibility (also known as 'opening energy'
based on unpairing probability) for all the corresponding sub-sequences (see Methods). We
examined the correlation between the opening energies and GFP levels. We found that the
opening energies of translation initiation sites, in particular from the nucleotide positions −30 to
18 (−30:18), shows the highest correlation with protein abundance (Fig 1A; Spearman's
correlation, R_s=−0.65, P<2.2×10^{-16}). This is stronger than the highest correlation between the
minimum free energy −30:30 and protein abundance, which was previously reported as the
highest ranked feature (Fig 1A; R_s=0.51, P<2.2×10^{-16}). To account for multiple-testing, the
P-values were adjusted using Bonferroni's correction and reported to machine precision. The
datasets used and results are summarised in Supplementary Table S1.

We repeated the analysis for a dataset of yellow fluorescent protein (YFP) expression in
Saccharomyces cerevisiae (23). This dataset corresponds to a library of 5′UTR variants, in
which the 10-nt sequences preceding the YFP translation initiation site were randomly
substituted (N=2,041). In this case, the opening energy −7:89 showed a stronger correlation

with protein abundance than that of the minimum free energy −15:50 reported previously (Fig
1B; R_s=−0.55 versus 0.46).
87
To examine the usefulness of accessibility in complex eukaryotes, we analysed a dataset of
GFP expression in *Mus musculus (31).* The reporter library was originally designed to measure
the strength of translation initiation sequence context, in which the 6- and 2-nt sequences
upstream and downstream of the GFP translation initiation site were randomly substituted,
respectively (N=65,536). Here the opening energy −8:11 showed a maximum correlation with
expressed proteins, which again, is stronger than that of the minimum free energy −30:30 (Fig
1C; R_s=−0.28 versus 0.12).
95
96 Taken together, our findings suggest that the accessibility of translation initiation sites strongly
97 correlates with protein abundance across species. Interestingly, our findings also suggest that
98 the Shine-Dalgarno sequence (*32*) at −13:−8 should be accessible to recruit ribosomes.
99
Accessibility predicts the outcome of recombinant protein expression
We investigated how accessibility performs in the real world in prediction of recombinant protein
expression. For this purpose, we analysed 11,430 expression experiments in *E. coli* from the
'Protein Structure Initiative:Biology' (PSI:Biology) (*33–35*). These PSI:Biology targets were
expressed using the pET21_NESG expression vector that harbours the *T7lac* inducible
promoter and a C-terminal His tag (*35*).
106
107 We split the experimental results of the PSI:Biology targets into protein expression 'success' and
108 'failure' groups (N=8,780 and 2,650, respectively; see Supplementary Fig S2). These
109 PSI:Biology targets span more than 189 species and the failures are representative of various
110 problems in heterologous protein expression. Only 1.6% of the targets were *E. coli* proteins,
111 which is negligible (N=179; see Supplementary Fig S2).
112
We calculated the opening energies for all possible sub-sequences of the PSI:Biology targets as
above (Fig 2, positions relative to initiation codons). For each sub-sequence region, we used the
opening energies to predict the expression outcomes and computed the prediction accuracy
using the area under the receiver operating characteristic curve (AUC; see Fig 2C). A closer
look into the correlations between opening energies and expression outcomes, and AUC scores
calculated for the sub-sequence regions reveals a strong accessibility signal of translation
initiation sites (Fig 2B&C, Cambray's GFP and PSI:Biology datasets, respectively). We matched
the correlations and AUC scores by sub-sequence regions and confirmed that sub-sequence
regions that have strong correlations are likely to have high AUC scores (Fig 2D). In contrast,
the sub-sequence regions that have zero correlations are not useful for predicting the
expression outcomes (AUC approximately 0.5).
124
We then asked how accessibility manifests in the endogenous mRNAs of *E. coli*, for which we
studied a proteomics dataset of 3,725 proteins available from PaxDb (*36*). As expected, we
observed a similar accessibility signal, with the region −25:16 correlated the most with protein

abundance (Fig 2E). However, the correlation was rather low (R=−0.17, P<2.2×10^{-16}), which
may reflect the limitation of mass spectrometry to detect lower abundances (*37*, *38*).
Furthermore, the endogenous promoters have variable strength, which gives rise to a broad
range of mRNA and protein levels (*39*, *40*). Taken together, our results show that the
accessibility signal of translation initiation sites is very consistent across various datasets
analysed (Supplementary Fig S1 and Fig 2).
134

Accessibility outperforms other features in prediction of recombinant protein expression
To choose an accessibility region for subsequent analyses, we selected the top 200 regions
from the above correlation analysis on Cambray's dataset (Fig 2B) and used random forest to
rank their Gini importance scores in prediction of the outcomes of the PSI:Biology targets. The
region −24:24 was ranked first, which is nearly identical to the region −23:24 with the top AUC
score (Fig 2C, AUC=0.70). We therefore used the opening energy at the region −24:24 in
subsequent analyses.
142

We asked how the other features perform compared to accessibility in prediction of
heterologous protein expression, for which we analysed the same PSI:Biology dataset. We first
calculated the minimum free energy and avoidance at the regions −30:30 and 1:30, respectively.
These are the local features associated with translation initiation rate. We also calculated CAI
(*11*), tAI (*41*), codon context (CC) (*42*), G+C content, and Iχnos scores (*43*). CC is similar to CAI
except it takes codon-pairs into account, whereas the Iχnos scores are translation elongation
rates predicted using a neural network model trained with ribosome profiling data
(Supplementary Fig S3). These are the global features associated with translation elongation
rate. We built a random forest model to rank the Gini importance scores of these local and
global features. The local features ranked higher than the global features (Fig 3A). We then
calculated and compared the prediction accuracy of these features. The AUC scores for the
local features were 0.70, 0.67 and 0.62 for the opening energy, minimum free energy and
avoidance, respectively, whereas the global features were 0.58, 0.57, 0.54, 0.54 and 0.51 for
Iχnos, G+C content, CAI, CC and tAI, respectively (Fig 3B). The local features outperform the
global features, suggesting that effects on translation initiation are a major predictor of the
outcome of heterologous protein expression. We further examined the local G+C contents
corresponding to the local features (Supplementary Fig S4). The G+C contents in the regions
−24:24 and −30:30 weakly correlate with opening energy and minimum free energy,
respectively. The AUC scores for these local G+C contents are also lower than the
corresponding local features, suggesting that these local G+C contents are not good proxies for
the corresponding local features. Overall, our findings support previous reports that the effects
on translation initiation are rate-limiting (*18*, *24*) which, interestingly, correlate with the binary
outcome of recombinant protein expression (Fig 3C). Importantly, accessibility outperformed all
other features.
167

To identify a good opening energy threshold, we calculated positive likelihood ratios for different
opening energy thresholds using the cumulative frequencies of true negative, false negative,
true positive and false positive derived from the above receiver operating characteristic (ROC)

analysis (Supplementary Fig S5, top panel). Meanwhile, we calculated the 95% confidence
intervals of these positive likelihood ratios using 10,000 bootstrap replicates. We reasoned that
there is an upper and lower bound on translation initiation rate, therefore the relationship
between translation initiation rate and accessibility is likely to follow a sigmoidal pattern. We fit
the positive likelihood ratios into a four-parametric logistic regression model (Supplementary Fig
S5). As a result, we are 95% confident that an opening energy of 10 kcal/mol or below at the
region −24:24 is about two times more likely to belong to the sequences which are successfully
expressed than those that failed.
179
Accessibility can be improved using a simulated annealing algorithm
The above results suggest that accessibility can, in part, explain the low expression problem of
heterologous protein expression. Therefore, we sought to exploit this idea for optimising gene
expression. We developed a simulated annealing algorithm to maximise the accessibility at the
region −24:24 using synonymous codon substitution (see Methods). Previous studies have
found that full-length synonymous codon-substituted transgenes may produce unexpected
results, such as a reduction in mRNA abundance, RNA toxicity, and/or protein misfolding (*25,*
43–45). Therefore, we sought to determine the minimum number of codons required for
synonymous substitutions in order to achieve near-optimum accessibility. For this purpose, we
used the PSI:Biology targets that failed to be expressed. We applied our simulated annealing
algorithm such that synonymous substitutions can happen at any codon of the sequences
except the start and stop codons, although the changes may not necessarily happen to all
codons due to the stochastic nature of our optimisation algorithm (see Methods). Next, we
constrained synonymous codon substitution to the first 14 codons and applied the same
procedure (Supplementary Fig S6A). Therefore, the changes may only occur at any or all of the
first 14 codons. We repeated the same procedure for the first nine and also the first four codons.
Thus a total of four series of codon-substituted sequences were generated. We then compared
the distributions of opening energy −24:24 for these series using the Kolmogorov-Smirnov
statistic (D_{KS}; see Supplementary Fig S6B). The distance between the distributions of the nine
and full-length codon-substituted series was significantly different yet sufficiently close
(D_{KS}=0.087, P=3.3×10^{-8}), suggesting that optimisation of the first nine codons is sufficient in
most cases to achieve an optimum accessibility of translation initiation sites. We named our
software Translation Initiation coding region designer (TIsigner), which by default, allows
synonymous substitutions in the first nine codons.
204
We asked to what extent the existing gene optimisation tools modify the accessibility of
translation initiation sites. For this purpose, we first submitted the PSI:Biology targets that failed
to be expressed to the ExpOptimizer web server from NovoPro Bioscience (see Methods). We
also optimised the PSI:Biology targets using the standalone version of Codon Optimisation
OnLine (COOL) (*29*). We found that both tools increase accessibility indirectly even though their
algorithms are not specifically designed to do so. In fact, a purely random synonymous codon
substitution on these PSI:Biology targets using our own script resulted in similar increases in
accessibility (Supplementary Fig S6C). These results may explain some indirect benefits from

the existing gene optimisation tools (i.e. any change from suboptimal is likely to be an improvement, see below).

Low protein yields can be improved by synonymous codon changes in the vicinity of translation initiation sites

To demonstrate that heterologous protein expression is tunable with minimum effort, we designed and tested a series of GFP reporter gene constructs. We tested 29 plasmids harbouring GFP reporter genes with synonymous changes within the first nine codons (opening energies of 5.56-21.68 kcal/mol; Supplementary Table S2 and Supplementary Methods). GFP expression is controlled by an IPTG inducible *T7lac* promoter. In addition, all plasmids harbour a second reporter gene, i.e. mScarlet-I, which is controlled by the constitutive promoter from the *nptII* gene for aminoglycoside-3'-O-phosphotransferase of *E. coli* transposon Tn5 (*46, 47*). mScarlet-I expression was measured to correct for plasmid copy number and as a proxy for bacterial growth (*48*). As expected, the GFP level significantly correlates with accessibility (i.e., anti-correlates with opening energy, R_s=−0.53, P=3.4×10^{-3}; Fig 6A). Curiously, we observed a diminishing return with opening energies lower than that of the wild-type sequence (11.68 kcal/mol). To investigate this, we simulated a protein production experiment by modelling cell growth, transcription, translation, and turnovers (see Methods). We assumed that opening energies of 12 kcal/mol or below is favourable in this model, based on our analysis of 8,780 PSI:Biology 'success' group (Supplementary Fig S6). Interestingly, our *in silico* coarse-grained model shows a similar protein production trend as the actual experiment (Fig 6B).

We then tested this finding using the luciferase reporter of *Renilla reniformis* (RLuc). Similarly, we designed a series of RLuc variants, but with opening energies below that of the wild-type sequence (5.77-10.38 kcal/mol; Fig 6C and Supplementary Table S2). In addition, we tested commercially designed sequences, in which sequence optimisations were performed in full-length rather than the first 9 codons. We observed that TIsigner (9.9 kcal/mol) and commercially optimised luciferase reporter genes produced significantly higher luminescence than the wild-type (Fig 6C), although RLuc is poorly soluble in the *E. coli* host (Supplementary Fig S8). We also found that the levels of wild-type luciferase and many variants with lower opening energies (5-7 kcal/mol) were not significantly different.

As both wild-type GFP and RLuc genes are strongly expressed in *E. coli*, we asked whether poorly expressed proteins can be improved by increasing accessibility of translation initiation sites. We performed densitometric analysis of previously published Western blots, which include the results of a cell-free expression system using constructs harbouring a wild-type antibody fragment or archaebacterial dioxygenase and its synonymous variants (within the first six codons) (*49*). Indeed, variants with opening energies lower than the wild-type sequences were expressed at higher levels (Fig 6D).

DISCUSSION

Our findings show that the accessibility of translation initiation sites is the strongest predictor of heterologous protein expression in *E. coli.* Whereas previous studies have largely used

minimum free energy models to define the accessibility of a region of interest (*27, 49–52*).
However, Terai and Asai (2020) and ourselves have independently discovered that the opening
energy is a better choice for modelling accessibility (*53, 54*) (see Fig 1A for example). Opening
energy is an ensemble average energy that accounts for suboptimal RNA structures that are not
reported by minimum free energy models by default (*55, 56*). Currently, the modelling of
accessibility using opening energy is largely used for the prediction of RNA-RNA intermolecular
interactions, for example, as implemented in RNAup and IntaRNA (*57, 58*). Our study has
shown that this approach can be used to identify the key accessibility regions that are consistent
across multiple large expression datasets. We have implemented our findings in TIsigner web
server, which currently supports recombinant protein expression in *E. coli* and *S. cerevisiae*
(optimisation regions −24:24 and −7:89, respectively; see Fig 1). An independent yet similar
implementation is available in XenoExpressO web server with the purpose of optimising protein
expression for an *E. coli* cell-free system (*59*). The authors showed that an increase in
accessibility of a 30 bp region from the Shine-Dalgarno sequence enhances the expression
level of human voltage dependent anion channel, which further supports our findings.
271
The strengths of our approaches are five-fold. Firstly, the likelihood of success or failure can be
assessed prior to running an experiment. Users can compare the opening energies calculated
for the input and optimised sequences and the distributions of the 'success' and 'failure' of the
PSI:Biology targets. We also introduced a scoring scheme to score the input and optimised
sequences based upon how likely they are to be expressed (Supplementary Fig S5; also see
Methods). Secondly, optimised sequences can have up to the first nine codons substituted (by
default), meaning that gene optimisation using a standard PCR cloning method is feasible. For
cloning, we propose a nested PCR approach, in which the final PCR reaction utilises a forward
primer designed according to the optimised sequence (*60*) (Supplementary Fig S6D). Thirdly,
the cost of gene optimisation can be reduced dramatically as gene synthesis is replaced with
PCR using our approach. This enables high-throughput protein expression screening using the
optimised sequences, generated at a low cost. Fourthly, tunable expression is possible, i.e.
high, intermediate or even low expression 5′ codon sequences can be designed, allowing for
more control over heterologous protein production, as demonstrated by our experiments (Fig 4).
Finally, our fast, lightweight, coarse-grained simulation approach has opened up new avenues
to study several aspects of gene expression, such as transcription, translation, cellular growth,
and turnovers, which give good proxies to how cellular systems behave.
289
MATERIALS AND METHODS
Sequence features analysis
Datasets used in this study are listed in Supplementary Table S1. Representative sequences
were chosen using CD-HIT-EST (*61, 62*). Minimum free energies, opening energies and
avoidance were calculated using RNAfold, RNAplfold and RNAup from ViennaRNA package
(version 2.4.11), respectively (*55–57, 63–66*). RNAfold was run with default parameters. For
RNAplfold, sub-sequences were generated from the input sequences to calculate opening
energies (using the parameters -W 210 -u 210). For RNAup, we examined the stochastic
interactions between the region 1:30 of each mRNA and 54 non-coding RNAs (using the

parameters -b -o). RNAup reports the total interaction between two RNAs as the sum of energy
required to open accessible sites in the interacting molecules ΔG_u and the energy gained by
subsequent hybridisation ΔG_h (*55*). For the interactions between each mRNA and 54
non-coding RNAs, we chose the most stable mRNA:ncRNA pair to report an inappropriate
mRNA:ncRNA interaction, i.e. the pair with the strongest hybridisation energy, $(\Delta G_h)_{min}$.
304
305 CAI, tAI and CC were calculated using the reference weights from Sharp and Li (*11*), Tuller et al.
306 (*41*) and Ang et al. (*42*), respectively. Translation elongation rate was predicted using Iχnos(*43*)
307 trained with ribosome profiling data (SRR7759806 and SRR7759807) (*67*).
308
Coarse-grained simulation
Our experiments showed a diminishing trend on protein production beyond a certain opening
energy (Fig 4). To explain this, we performed a coarse grained simulation using constructs with
increasing opening energy on a simulated cellular system. Despite being less precise than fine
grained methods such as *ab initio* and molecular dynamics, coarse grained simulations often
give similar results, with an added advantage of being scalable to very large systems.
315
To set the simulation, we binned the opening energies between 2 and 32 in intervals of two, with
each bin representing a 'reporter plasmid construct' whose opening energy is the mean of the
bin. For each construct, the 'technical replicates' were generated by allowing slight variations on
the mean opening energy of the bin. This is to model variation between replicates, and the
discrepancies between the estimated and the actual opening energies *in vivo*. For each round of
transcription, mRNA copies were randomly generated from 30 to 60 plasmid DNA copies (*3*, *68*,
69). We chose an optimum opening energy of 12 kcal/mol or less for translation. However, this
is probabilistic which occasionally allowed protein production from higher opening energy
transcripts. We allowed mRNA to decay probabilistically when a mRNA molecule is translated
for more than 10 rounds.
326
327 We also set a threshold of protein tolerance to be 1,000,000 copies where the copy numbers of
328 endogenous proteins are usually less than 10,000 (*10*), beyond which there is a sporadic death
329 of cells. However, in this simulation, the chances of staying viable and reproducing are higher
330 than death, and cells grow steadily. This threshold also simulated random but low cell deaths in
331 the experiment, without setting an extra variable.
332
To limit the computational complexity, our coarse-grained simulations used lower constants and
iterations. Initialising with 100 cells, the algorithm was set to terminate either after 10,000
iterations or when the total number of cells becomes zero. After termination, the total number of
proteins and cells for each construct were taken from the endpoints. To imitate 'biological
replicates', we repeated the above simulation three times with different random numbers, which
provides slightly different initial conditions for each experiment.
339
340 **TIsigner development**

Finding a synonymous sequence with a maximum accessibility is a combinatorial problem that spans a vast search space. For example, for a protein-coding sequence of nine codons, assuming an average of 3 synonymous codons per amino acid, we can expect a total of 19,682 unique synonymous coding sequences. This number increases rapidly with increasing numbers of codons. Heuristic optimisation approaches are preferred in such situations because the search space can be explored more efficiently to obtain nearly optimal solutions.

To optimise the accessibility of a given sequence, TIsigner uses a simulated annealing algorithm (*70–73*), a heuristic optimisation technique based on the thermodynamics of a system settling into a low energy state after cooling. Simulated annealing algorithms have been used to solve many combinatorial optimisation problems in bioinformatics. For example, we previously applied this algorithm to align and predict non-coding RNAs from multiple sequences (*74*). Other studies use this algorithm to find consensus sequences (*72*), optimise ribosome binding sites (*27*) and predict mRNA foldings (*75*) using minimum free energy models.

According to statistical mechanics, the probability p_i of a system occupying energy state E_i, with temperature T, follows a Boltzmann distribution of the form $e^{-E_i/T}$, which gives a set of probability mass functions along every point i in the solution space. Using a Markov chain sampling, these probabilities are sampled such that each point has a lower temperature than the previous one. As the system is cooled from high to low temperatures ($T \rightarrow 0$), the samples converge to a minimum of E, which in many cases will be the global minimum (*72*). A frequently used Markov chain sampling technique is Metropolis-Hastings algorithm in which a 'bad' move E_2 from initial state E_1 such that $E_2 > E_1$, is accepted if $R(0,1) \geq p_2/p_1$, where $R(0,1)$ is a uniformly random number between 0 and 1.

In our implementation, each iteration consists of a move that may involve multiple synonymous codon substitutions. The algorithm begins at a high temperature where the first move is drastic, synonymous substitutions occur in all replaceable codons. At the end of the first iteration, a new sequence is accepted if the opening energy is smaller than that of the input sequence. However, if the opening energy of a new sequence is greater than that of the input sequence, acceptance depends on the Metropolis-Hastings criteria. The accepted sequence is used for the next iteration, which repeats the above process. As the temperature cools, the moves get milder with fewer synonymous codon changes (Supplementary Fig S6A). Simulated annealing stops upon reaching a near-optimum solution.

For the web version of TIsigner, the default number of replaceable codons is restricted to the first nine codons. However, this default setting can be reset to range from the first four to nine codons, or the full length of the coding sequence. Since the accessibility of a fixed region is optimised, this process only takes O(1) time (Supplementary Fig S7). Furthermore, TIsigner runs multiple simulated annealing instances, in parallel, to obtain multiple possible sequence solutions.

When users select *T7lac* promoter as the 5′UTR, they can adjust 'Expression Score', that is calculated based on the PSI:Biology dataset (see below). This allows them to tune the expression level of a target gene. In contrast, when users input a custom 5′UTR sequence, they only have the option to either maximise or minimise expression.

To implement 'Expression Score', the posterior probabilities of success for input and optimised sequences are evaluated using the following equations from Bayesian statistics:

$$positive\ posterior\ odds\ =\ prior\ odds\ \times\ fitted\ positive\ likelihood\ ratio \qquad (1)$$

$$positive\ posterior\ probability\ =\ \frac{positive\ posterior\ odds}{(1+positive\ posterior\ odds)} \qquad (2)$$

The fitted positive likelihood ratios in equation (1) were obtained from the following 4-parametric logistic regression equation:

$$fitted\ positive\ likelihood\ ratio\ =\ d\ +\ \frac{a-d}{1+(\frac{positive\ likelihood\ ratio}{c})^{b}} \qquad (3)$$

with parameters a, b, c, and d. The prior probability was set to 0.49, which is the proportion of 'Expressed' (N=21,046) divided by 'Cloned' (N=42,774) of the PSI:Biology targets reported as of 28 June 2017 (http://targetdb.rcsb.org/metrics/). Posterior probabilities were scaled as percentages to score the input and optimised sequences.

The presence of terminator-like elements (*76*) in the protein-coding region may result in expression of truncated mRNAs due to early transcription termination. Therefore, we implemented an optional check for putative terminators in the input and optimised sequences by cmsearch (INFERNAL version 1.1.2) (*77*) using the covariance models of terminators from RMfam (*78*, *79*). We also allow users to filter the output sequences for the presence of restriction sites. Restriction modification sites (AarI, BsaI, and BsmBI) are avoided by default.

Besides *E. coli*, users can choose *S. cerevisiae*, *M. musculus* or 'Other' as the expression host. The regions for optimising accessibility are −7:89, −8:11 and −24:89 for *S. cerevisiae*, *M. musculus* and 'Other', respectively (Fig 1 and Supplementary Fig S1). When users choose 'Custom' for expression host, the region for optimising accessibility becomes customisable.

Sequence optimisation
We submitted the PSI:Biology targets that failed to be expressed (N=2,650) to the ExpOptimizer web server from NovoPro Bioscience (https://www.novoprolabs.com/tools/codon-optimization). A total of 2,573 sequences were optimised. The target sequences were also optimised using a local version of COOL (*29*) and TIsigner using default settings. We also ran a random synonymous codon substitution as a control for these 2,573 sequences.

GFP assay

Plasmids were constructed using the MIDAS Golden Gate cloning system (Supplementary Methods) (*80*). BL21(DE3)pLysS competent *E. coli* cells (Invitrogen) were transformed with plasmids and grown overnight on Luria-Bertani (LB) agar plates containing spectinomycin (50 µg/ml) and chloramphenicol (25 µg/ml). Single colonies were picked and inoculated into 3 ml LB broth containing the same antibiotics, and cultures were grown for 18 hours at 37°C, 200 rpm. Cultures were diluted with fresh media at 1:20 and grown at 37°C, 200 rpm, until reaching the mid-logarithmic growth phase (optical densities at 600 nm (OD_{600}) of ~0.3). Of each culture, 20 µl was seeded into 96-well plates containing 180 µl LB broth supplemented with antibiotics and isopropyl-β-D thiogalactopyranoside (IPTG) (1 mM final concentration) per well. Fluorescence intensities and ODs were measured in a black, flat, clear bottom 96-well plate with lid (CELLSTAR, Greiner) using a FLUOstar Omega plate reader (BMG Labtech) equipped with an excitation filter (band pass 485-12) and an emission filter (band pass 520) for GFP and excitation filter (band pass 484) and an emission filter (band pass 610-10) for mScarlet-I. The plate was incubated at 37°C with "meander corner well shaking" at 300 rpm for 7 hours measuring fluorescence and ODs every 10 minutes. Fluorescence was measured in a 2 mm circle recording the average of 8 measurements per well. Average values of technical replicates were calculated and normalised to the mScarlet-I second reporter, and then to the normalised value of the GFP variant with the highest opening energy (21.68 kcal/mol). Normalised fluorescence values were obtained from the average values of biological replicates (Supplementary Table S2).

Luciferase assay

BL21Star(DE3) competent cells (Invitrogen) were transformed with plasmids and grown overnight at 37°C on LB agar plates containing 50 µg/ml spectinomycin. Single colonies were picked and inoculated into 5 ml LB broth (50 µg/ml spectinomycin) and grown for 18 hours at 37°C, 200 rpm. Bacterial cultures were diluted with fresh media at 1:20 and grown at 37°C, 200 rpm, up to a mid-logarithmic phase (OD_{600} of ~0.4). The cultures were split and induced with IPTG at a final concentration of 0.25 mM (or uninduced as controls), and seeded into a white, flat, clear bottom 96-well white plate with lid (Costar, Corning), 150 µl per well, in triplicates. Cells were incubated in a FLUOstar Omega Microplate Reader (BMG LABTECH) for 90 minutes at 25°C, 200 rpm, and OD_{600} was measured every 15 minutes (over 7 cycles). Cells were harvested by centrifugation at 3000 ×g, for 10 minutes, at 20°C. Supernatants were removed. As the substrate can penetrate into cells, 50 µl of coelenterazine h (Promega) was added to living cells to minimise sample processing steps and variability (*81*, *82*). Luminescence was measured (λ_{em} = 475 nm) in a Clariostar microplate reader (BMG LABTECH) at 25°C every 2 minutes (over 11 cycles). Average values of technical replicates were calculated and normalised to the wild-type. Normalised luminescence values were obtained from the average values of biological replicates (Supplementary Table S2).

Statistical analysis

AUC and Gini importance scores were calculated using scikit-learn (version 0.20.2) (*83*). The 95% confidence intervals for AUC scores were calculated using DeLong's method (*84*). Spearman's correlation coefficients and Kolmogorov-Smirnov statistics were calculated using

Pandas (version 0.23.4) (*85*) and scipy (version 1.2.1) (*86, 87*), respectively. Positive likelihood ratios with 95% confidence intervals were calculated using the bootLR package (*88, 89*). The P-values of multiple testing were adjusted using Bonferroni's correction and reported to machine precision. Plots were generated using Matplotlib (version 3.0.2) (*90*) and Seaborn (version 0.9.0) (*91*).

Code and data availability
Our code and data can be found in our GitHub repository (https://github.com/Gardner-BinfLab/TIsigner_paper_2019). These include the scripts and Jupyter notebooks to reproduce our results and figures. The source code of TIsigner is available at https://github.com/Gardner-BinfLab/TISIGNER-ReactJS. The public web version of this tool runs at https://tisigner.com/tisigner. The experimental data, analysis and results are available at https://github.com/bkb3/TIsignerExperiment/tree/master/Jupyter and an interactive version of results are available at https://bkb3.github.io/TIsignerExperiment/.

ACKNOWLEDGEMENTS
We thank Professor Ivo Hofacker for fruitful discussions at the Benasque RNA Meeting, and Dr Ronny Lorenz for helpful discussions about RNAplfold. We are grateful to the members of the Biomolecular Interaction Centre at the University of Canterbury for supporting this research. We thank New Zealand eScience Infrastructure for providing high performance computing resources. This work was supported in part by the Ministry of Business, Innovation and Employment [MBIE Smart Idea grant: UOOX1709 and MBIE Data Science Programmes grant: UOAX1932] and the Royal Society of New Zealand Te Apārangi [Marsden grant: 19-UOO-040].

AUTHOR CONTRIBUTIONS
C.S.L. and P.P.G. conceived the work; C.S.L. contributed RNA accessibility analyses; B.K.B. performed the coarse-grained simulation and developed the TIsigner web server; C.D., D.M.R., and A.C. constructed the plasmids, performed the GFP assay, and the luciferase assay, respectively. C.S.L. and B.K.B. analysed the data and drafted the manuscript. All authors reviewed, edited and approved the manuscript.

COMPETING INTERESTS
The authors declare no competing interests.

REFERENCES

1. A. Kimelman, A. Levy, H. Sberro, S. Kidron, A. Leavitt, G. Amitai, D. R. Yoder-Himes, O. Wurtzel, Y. Zhu, E. M. Rubin, R. Sorek, A vast collection of microbial genes that are toxic to bacteria. *Genome Res.* **22**, 802–809 (2012).

2. A. Berlec, B. Strukelj, Current state and recent advances in biopharmaceutical production in Escherichia coli, yeasts and mammalian cells. *J. Ind. Microbiol. Biotechnol.* **40**, 257–274 (2013).

3. G. L. Rosano, E. A. Ceccarelli, Recombinant protein expression in Escherichia coli:
advances and challenges. *Front. Microbiol.* **5**, 172 (2014).

4. R. de S. Abreu, R. de Sousa Abreu, L. O. Penalva, E. M. Marcotte, C. Vogel, Global
signatures of protein and mRNA expression levels. *Molecular BioSystems* (2009), ,
doi:10.1039/b908315d.

5. G. Hanson, J. Coller, Codon optimality, bias and usage in translation and mRNA decay.
Nat. Rev. Mol. Cell Biol. **19**, 20–30 (2018).

6. C. S. Lim, S. J. T. Wardell, T. Kleffmann, C. M. Brown, The exon–intron gene structure
upstream of the initiation codon predicts translation efficiency. *Nucleic Acids Research*. **46**
(2018), pp. 4575–4591.

7. S. G. Stevens, C. M. Brown, In silico estimation of translation efficiency in human cell lines:
potential evidence for widespread translational control. *PLoS One*. **8**, e57625 (2013).

8. B. Schwanhäusser, D. Busse, N. Li, G. Dittmar, J. Schuchhardt, J. Wolf, W. Chen, M.
Selbach, Global quantification of mammalian gene expression control. *Nature*. **473**,
337–342 (2011).

9. J. A. Bernstein, A. B. Khodursky, P.-H. Lin, S. Lin-Chao, S. N. Cohen, Global analysis of
mRNA decay and abundance in Escherichia coli at single-gene resolution using two-color
fluorescent DNA microarrays. *Proc. Natl. Acad. Sci. U. S. A.* **99**, 9697–9702 (2002).

10. Y. Taniguchi, P. J. Choi, G.-W. Li, H. Chen, M. Babu, J. Hearn, A. Emili, X. S. Xie,
Quantifying E. coli proteome and transcriptome with single-molecule sensitivity in single
cells. *Science*. **329**, 533–538 (2010).

11. P. M. Sharp, W. H. Li, The codon Adaptation Index--a measure of directional synonymous
codon usage bias, and its potential applications. *Nucleic Acids Res.* **15**, 1281–1295 (1987).

12. M. d. Reis, M. d. Reis, Solving the riddle of codon usage preferences: a test for
translational selection. *Nucleic Acids Research*. **32** (2004), pp. 5036–5044.

13. R. Sabi, T. Tuller, Modelling the Efficiency of Codon–tRNA Interactions Based on Codon
Usage Bias. *DNA Research*. **21** (2014), pp. 511–526.

14. G. A. Gutman, G. W. Hatfield, Nonrandom utilization of codon pairs in Escherichia coli.
Proc. Natl. Acad. Sci. U. S. A. **86**, 3699–3703 (1989).

15. C. E. Brule, E. J. Grayhack, Synonymous Codons: Choose Wisely for Expression. *Trends
Genet.* **33**, 283–297 (2017).

16. I. A. Osterman, Z. S. Chervontseva, S. A. Evfratov, A. V. Sorokina, V. A. Rodin, M. P.
Rubtsova, E. S. Komarova, T. S. Zatsepin, M. R. Kabilov, A. A. Bogdanov, M. S. Gelfand,
O. A. Dontsova, P. V. Sergiev, Translation at first sight: the influence of leading codons.
Nucleic Acids Res. **48**, 6931–6942 (2020).

17. M. Verma, J. Choi, K. A. Cottrell, Z. Lavagnino, E. N. Thomas, S. Pavlovic-Djuranovic, P.
Szczesny, D. W. Piston, H. S. Zaher, J. D. Puglisi, S. Djuranovic, A short translational ramp

determines the efficiency of protein synthesis. *Nat. Commun.* **10**, 5774 (2019).

18. G. Kudla, A. W. Murray, D. Tollervey, J. B. Plotkin, Coding-sequence determinants of gene
expression in Escherichia coli. *Science*. **324**, 255–258 (2009).

19. J. B. Plotkin, G. Kudla, Synonymous but not the same: the causes and consequences of
codon bias. *Nature Reviews Genetics*. **12** (2011), pp. 32–42.

20. G. Boël, R. Letso, H. Neely, W. Nicholson Price, K.-H. Wong, M. Su, J. D. Luff, M. Valecha,
J. K. Everett, T. B. Acton, R. Xiao, G. T. Montelione, D. P. Aalberts, J. F. Hunt, Codon
influence on protein expression in E. coli correlates with mRNA levels. *Nature*. **529** (2016),
pp. 358–363.

21. G. Cambray, J. C. Guimaraes, A. P. Arkin, Evaluation of 244,000 synthetic sequences
reveals design principles to optimize translation in Escherichia coli. *Nat. Biotechnol.* **36**,
1005–1015 (2018).

22. M. H. de Smit, J. van Duin, Secondary structure of the ribosome binding site determines
translational efficiency: a quantitative analysis. *Proc. Natl. Acad. Sci. U. S. A.* **87**,
7668–7672 (1990).

23. S. Dvir, L. Velten, E. Sharon, D. Zeevi, L. B. Carey, A. Weinberger, E. Segal, Deciphering
the rules by which 5'-UTR sequences affect protein expression in yeast. *Proc. Natl. Acad.*
Sci. U. S. A. **110**, E2792–801 (2013).

24. T. Tuller, H. Zur, Multiple roles of the coding sequence 5′ end in gene expression regulation.
Nucleic Acids Research. **43** (2015), pp. 13–28.

25. S. U. Umu, A. M. Poole, R. C. Dobson, P. P. Gardner, Avoidance of stochastic RNA
interactions can be harnessed to control protein expression levels in bacteria and archaea.
Elife. **5** (2016), doi:10.7554/eLife.13479.

26. A. Villalobos, J. E. Ness, C. Gustafsson, J. Minshull, S. Govindarajan, Gene Designer: a
synthetic biology tool for constructing artificial DNA segments. *BMC Bioinformatics*. **7**, 285
(2006).

27. H. M. Salis, E. A. Mirsky, C. A. Voigt, Automated design of synthetic ribosome binding sites
to control protein expression. *Nature Biotechnology*. **27** (2009), pp. 946–950.

28. D. Raab, M. Graf, F. Notka, T. Schödl, R. Wagner, The GeneOptimizer Algorithm: using a
sliding window approach to cope with the vast sequence space in multiparameter DNA
sequence optimization. *Syst. Synth. Biol.* **4**, 215–225 (2010).

29. B. K.-S. Chung, D.-Y. Lee, Computational codon optimization of synthetic gene for protein
expression. *BMC Syst. Biol.* **6**, 134 (2012).

30. G. Terai, S. Kamegai, K. Asai, CDSfold: an algorithm for designing a protein-coding
sequence with the most stable secondary structure. *Bioinformatics*. **32**, 828–834 (2016).

31. W. L. Noderer, R. J. Flockhart, A. Bhaduri, A. J. Diaz de Arce, J. Zhang, P. A. Khavari, C. L.
Wang, Quantitative analysis of mammalian translation initiation sites by FACS-seq. *Mol.*

Syst. Biol. **10**, 748 (2014).

32. J. Shine, L. Dalgarno, The 3'-terminal sequence of Escherichia coli 16S ribosomal RNA:
complementarity to nonsense triplets and ribosome binding sites. *Proc. Natl. Acad. Sci. U.*
S. A. **71**, 1342–1346 (1974).

33. L. Chen, R. Oughtred, H. M. Berman, J. Westbrook, TargetDB: a target registration
database for structural genomics projects. *Bioinformatics.* **20**, 2860–2862 (2004).

34. C. Y. Seiler, J. G. Park, A. Sharma, P. Hunter, P. Surapaneni, C. Sedillo, J. Field, R. Algar,
A. Price, J. Steel, A. Throop, M. Fiacco, J. LaBaer, DNASU plasmid and
PSI:Biology-Materials repositories: resources to accelerate biological research. *Nucleic*
Acids Res. **42**, D1253–60 (2014).

35. T. B. Acton, K. C. Gunsalus, R. Xiao, L. C. Ma, J. Aramini, M. C. Baran, Y.-W. Chiang, T.
Climent, B. Cooper, N. G. Denissova, S. M. Douglas, J. K. Everett, C. K. Ho, D. Macapagal,
P. K. Rajan, R. Shastry, L.-Y. Shih, G. V. T. Swapna, M. Wilson, M. Wu, M. Gerstein, M.
Inouye, J. F. Hunt, G. T. Montelione, Robotic cloning and Protein Production Platform of the
Northeast Structural Genomics Consortium. *Methods Enzymol.* **394**, 210–243 (2005).

36. M. Wang, C. J. Herrmann, M. Simonovic, D. Szklarczyk, C. von Mering, Version 4.0 of
PaxDb: Protein abundance data, integrated across model organisms, tissues, and
cell-lines. *Proteomics.* **15**, 3163–3168 (2015).

37. D. L. Tabb, L. Vega-Montoto, P. A. Rudnick, A. M. Variyath, A.-J. L. Ham, D. M. Bunk, L. E.
Kilpatrick, D. D. Billheimer, R. K. Blackman, H. L. Cardasis, Others, Repeatability and
reproducibility in proteomic identifications by liquid chromatography- tandem mass
spectrometry. *J. Proteome Res.* **9**, 761–776 (2009).

38. T. Nilsson, M. Mann, R. Aebersold, J. R. Yates 3rd, A. Bairoch, J. J. M. Bergeron, Mass
spectrometry in high-throughput proteomics: ready for the big time. *Nat. Methods.* **7**,
681–685 (2010).

39. U. Deuschle, W. Kammerer, R. Gentz, H. Bujard, Promoters of Escherichia coli: a hierarchy
of in vivo strength indicates alternate structures. *EMBO J.* **5**, 2987–2994 (1986).

40. F. Delvigne, J. Baert, H. Sassi, P. Fickers, A. Grünberger, C. Dusny, Taking control over
microbial populations: Current approaches for exploiting biological noise in bioprocesses.
Biotechnol. J. **12** (2017), doi:10.1002/biot.201600549.

41. T. Tuller, Y. Y. Waldman, M. Kupiec, E. Ruppin, Translation efficiency is determined by both
codon bias and folding energy. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 3645–3650 (2010).

42. K. S. Ang, S. Kyriakopoulos, W. Li, D.-Y. Lee, Multi-omics data driven analysis establishes
reference codon biases for synthetic gene design in microbial and mammalian cells.
Methods. **102** (2016), pp. 26–35.

43. R. Tunney, N. J. McGlincy, M. E. Graham, N. Naddaf, L. Pachter, L. F. Lareau, Accurate
design of translational output by a neural network model of ribosome distribution. *Nat.*

Struct. Mol. Biol. **25**, 577–582 (2018).

44. T. Ben-Yehezkel, S. Atar, H. Zur, A. Diament, E. Goz, T. Marx, R. Cohen, A. Dana, A.
Feldman, E. Shapiro, T. Tuller, Rationally designed, heterologous S. cerevisiae transcripts
expose novel expression determinants. *RNA Biol.* **12**, 972–984 (2015).

45. P. Mittal, J. Brindle, J. Stephen, J. B. Plotkin, G. Kudla, Codon usage influences fitness
through RNA toxicity. *Proc. Natl. Acad. Sci. U. S. A.* **115**, 8639–8644 (2018).

46. D. S. Bindels, L. Haarbosch, L. van Weeren, M. Postma, K. E. Wiese, M. Mastop, S.
Aumonier, G. Gotthard, A. Royant, M. A. Hink, T. W. J. Gadella Jr, mScarlet: a bright
monomeric red fluorescent protein for cellular imaging. *Nat. Methods*. **14**, 53–56 (2017).

47. R. O. Schlechter, H. Jun, M. Bernach, S. Oso, E. Boyd, D. A. Muñoz-Lintz, R. C. J. Dobson,
D. M. Remus, M. N. P. Remus-Emsermann, Chromatic Bacteria - A Broad Host-Range
Plasmid and Chromosomal Insertion Toolbox for Fluorescent Protein Expression in
Bacteria. *Front. Microbiol.* **9**, 3052 (2018).

48. R. O. Schlechter, D. M. Remus, M. N. P. Remus-Emsermann, Constitutively expressed
fluorescent proteins allow to track bacterial growth and to determine relative fitness of
bacteria in mixed cultures. *Cold Spring Harbor Laboratory* (2020), p. 2020.12.01.399113.

49. D. Voges, M. Watzele, C. Nemetz, S. Wizemann, B. Buchberger, Analyzing and enhancing
mRNA translational efficiency in an Escherichia coli in vitro expression system. *Biochem.*
Biophys. Res. Commun. **318**, 601–614 (2004).

50. S. Bhattacharyya, W. M. Jacobs, B. V. Adkar, J. Yan, W. Zhang, E. I. Shakhnovich,
Accessibility of the Shine-Dalgarno Sequence Dictates N-Terminal Codon Bias in E. coli.
Mol. Cell. **70**, 894–905.e5 (2018).

51. T. Nieuwkoop, N. J. Claassens, J. van der Oost, Improved protein production and codon
optimization analyses in Escherichia coli by bicistronic design. *Microb. Biotechnol.* **12**,
173–179 (2019).

52. J. Pelletier, N. Sonenberg, The involvement of mRNA secondary structure in protein
synthesis. *Biochem. Cell Biol.* **65**, 576–581 (1987).

53. B. K. Bhandari, C. S. Lim, P. P. Gardner, Highly accessible translation initiation sites are
predictive of successful heterologous protein expression. *BioRxiv* (2019), ,
doi:10.1101/726752.

54. G. Terai, K. Asai, Improving the prediction accuracy of protein abundance in Escherichia
coli using mRNA accessibility. *Nucleic Acids Res.* **48**, e81–e81 (2020).

55. U. Mückstein, H. Tafer, J. Hackermüller, S. H. Bernhart, P. F. Stadler, I. L. Hofacker,
Thermodynamics of RNA–RNA binding. *Bioinformatics*. **22**, 1177–1182 (2006).

56. S. H. Bernhart, U. Mückstein, I. L. Hofacker, RNA Accessibility in cubic time. *Algorithms*
Mol. Biol. **6**, 3 (2011).

57. R. Lorenz, S. H. Bernhart, C. Höner Zu Siederdissen, H. Tafer, C. Flamm, P. F. Stadler, I. L.

Hofacker, ViennaRNA Package 2.0. *Algorithms Mol. Biol.* **6**, 26 (2011).

58. M. Mann, P. R. Wright, R. Backofen, IntaRNA 2.0: enhanced and customizable prediction of
RNA–RNA interactions. *Nucleic Acids Res.* **45**, W435–W439 (2017).

59. S. Zayni, S. Damiati, S. Moreno-Flores, F. Amman, I. Hofacker, E.-K. Ehmoser, Enhancing
the cell-free expression of native membrane proteins by in-silico optimization of the coding
sequence – an experimental study of the human voltage-dependent anion channel. *BioRxiv*
(2018), , doi:10.1101/411694.

60. J. Sambrook, D. W. Russell, *Molecular cloning: a laboratory manual. Vol. 3* (CSHL Press,
2001).

61. W. Li, A. Godzik, Cd-hit: a fast program for clustering and comparing large sets of protein or
nucleotide sequences. *Bioinformatics*. **22** (2006), pp. 1658–1659.

62. L. Fu, B. Niu, Z. Zhu, S. Wu, W. Li, CD-HIT: accelerated for clustering the next-generation
sequencing data. *Bioinformatics*. **28**, 3150–3152 (2012).

63. I. L. Hofacker, W. Fontana, P. F. Stadler, L. S. Bonhoeffer, M. Tacker, P. Schuster, Fast
folding and comparison of RNA secondary structures. *Monatshefte für Chemie / Chemical*
Monthly. **125**, 167–188 (1994).

64. S. Bernhart, I. L. Hofacker, P. F. Stadler, Local Base Pairing Probabilities in Large RNAs.
Bioinformatics (available at
http://mescalin.tbi.univie.ac.at/papers/Abstracts/05-10-001abs.html).

65. A. F. Bompfünewerer, R. Backofen, S. H. Bernhart, J. Hertel, I. L. Hofacker, P. F. Stadler, S.
Will, Variations on RNA folding and alignment: lessons from Benasque. *J. Math. Biol.* **56**,
129–144 (2008).

66. R. Lorenz, I. L. Hofacker, P. F. Stadler, RNA folding with hard and soft constraints.
Algorithms Mol. Biol. **11**, 8 (2016).

67. F. Mohammad, R. Green, A. R. Buskirk, A systematically-revised ribosome profiling method
for bacteria reveals pauses at single-codon resolution. *Elife*. **8** (2019),
doi:10.7554/eLife.42591.

68. D. Held, K. Yaeger, R. Novy, "New coexpression vectors for expanded compatibilities in E.
coli" (18, Novagen, 2003).

69. L. Gomes, G. Monteiro, F. Mergulhão, The Impact of IPTG Induction on Plasmid Stability
and Heterologous Protein Expression by Biofilms. *Int. J. Mol. Sci.* **21** (2020),
doi:10.3390/ijms21020576.

70. S. Kirkpatrick, C. D. Gelatt, M. P. Vecchi, Optimization by Simulated Annealing. *Science*.
220 (1983), pp. 671–680.

71. L. Ingber, Adaptive simulated annealing (ASA): Lessons learned (2000), (available at
http://arxiv.org/abs/cs/0001018).

72. J. M. Keith, P. Adams, D. Bryant, D. P. Kroese, K. R. Mitchelson, D. A. E. Cochran, G. H. Lala, A simulated annealing algorithm for finding consensus sequences. *Bioinformatics*. **18**, 1494–1499 (2002).

73. J. Brownlee, *Clever Algorithms: Nature-inspired Programming Recipes* (Jason Brownlee, 2011).

74. S. Lindgreen, P. P. Gardner, A. Krogh, MASTR: multiple alignment and structure prediction of non-coding RNAs using simulated annealing. *Bioinformatics*. **23**, 3304–3311 (2007).

75. P. Gaspar, G. Moura, M. A. S. Santos, J. L. Oliveira, mRNA secondary structure optimization using a correlated stem-loop prediction. *Nucleic Acids Res.* **41**, e73 (2013).

76. Y.-J. Chen, P. Liu, A. A. K. Nielsen, J. A. N. Brophy, K. Clancy, T. Peterson, C. A. Voigt, Characterization of 582 natural and synthetic terminators and quantification of their design constraints. *Nat. Methods*. **10**, 659–664 (2013).

77. E. P. Nawrocki, S. R. Eddy, Infernal 1.1: 100-fold faster RNA homology searches. *Bioinformatics*. **29** (2013), pp. 2933–2935.

78. P. P. Gardner, H. Eldai, Annotating RNA motifs in sequences and alignments. *Nucleic Acids Res.* **43**, 691–698 (2015).

79. I. Kalvari, J. Argasinska, N. Quinones-Olvera, E. P. Nawrocki, E. Rivas, S. R. Eddy, A. Bateman, R. D. Finn, A. I. Petrov, Rfam 13.0: shifting to a genome-centric resource for non-coding RNA families. *Nucleic Acids Res.* **46**, D335–D342 (2018).

80. C. J. van Dolleweerd, S. A. Kessans, K. C. Van de Bittner, L. Y. Bustamante, R. Bundela, B. Scott, M. J. Nicholson, E. J. Parker, MIDAS: A Modular DNA Assembly System for Synthetic Biology. *ACS Synth. Biol.* **7**, 1018–1029 (2018).

81. W. W. Lorenz, M. J. Cormier, D. J. O'Kane, D. Hua, A. A. Escher, A. A. Szalay, Expression of the Renilla reniformis luciferase gene in mammalian cells. *J. Biolumin. Chemilumin.* **11**, 31–37 (1996).

82. M. Fuhrmann, A. Hausherr, L. Ferbitz, T. Schödl, M. Heitzer, P. Hegemann, Monitoring dynamic expression of nuclear genes in Chlamydomonas reinhardtii by using a synthetic luciferase reporter gene. *Plant Mol. Biol.* **55**, 869–881 (2004).

83. F. Pedregosa, G. Varoquaux, A. Gramfort, V. Michel, B. Thirion, O. Grisel, M. Blondel, P. Prettenhofer, R. Weiss, V. Dubourg, J. Vanderplas, A. Passos, D. Cournapeau, M. Brucher, M. Perrot, É. Duchesnay, Scikit-learn: Machine Learning in Python. *J. Mach. Learn. Res.* **12**, 2825–2830 (2011).

84. E. R. DeLong, D. M. DeLong, D. L. Clarke-Pearson, Comparing the areas under two or more correlated receiver operating characteristic curves: a nonparametric approach. *Biometrics*. **44**, 837–845 (1988).

85. W. McKinney, in *Proceedings of the 9th Python in Science Conference* (2010), pp. 51–56.

86. T. E. Oliphant, Python for Scientific Computing. *Computing in Science Engineering*. **9**,

10–20 (2007).

87. K. J. Millman, M. Aivazis, Python for Scientists and Engineers. *Computing in Science*
Engineering. **13**, 9–12 (2011).

88. K. A. Marill, Y. Chang, K. F. Wong, A. B. Friedman, Estimating negative likelihood ratio
confidence when test sensitivity is 100%: A bootstrapping approach. *Stat. Methods Med.*
Res. **26**, 1936–1948 (2017).

89. R Core Team, *R: A Language and Environment for Statistical Computing* (R Foundation for
Statistical Computing, Vienna, 2019; https://www.r-project.org/).

90. Matplotlib: A 2D Graphics Environment - IEEE Journals & Magazine, (available at
https://doi.org/10.1109/MCSE.2007.55).

91. M. Waskom, O. Botvinnik, D. O'Kane, P. Hobson, J. Ostblom, S. Lukauskas, D. C.
Gemperline, T. Augspurger, Y. Halchenko, J. B. Cole, J. Warmenhoven, J. de Ruiter, C.
Pye, S. Hoyer, J. Vanderplas, S. Villalba, G. Kunter, E. Quintero, P. Bachant, M. Martin, K.
Meyer, A. Miles, Y. Ram, T. Brunner, T. Yarkoni, M. L. Williams, C. Evans, C. Fitzgerald,
Brian, A. Qalieh, mwaskom/seaborn: v0.9.0 (July 2018) (2018),
doi:10.5281/zenodo.1313201.

769 **Figures**

770 Translation initiation site

771 More accessible ← → Less structural

773 **A** *Escherichia coli*
Cambray *et al.* GFP dataset (n=14,425)

774 ...U...A̲A̲U̲G̲N̲N̲N...N...N...N GFP fusion

$R_s = -0.65$ $R_s = 0.51$

785 **B** *Saccharomyces cerevisiae*
Dvir *et al.* YFP dataset (n=2,041)

786 A...NNNNNNNNNN̲A̲U̲G...A...C YFP

$R_s = -0.55$ $R_s = 0.46$

797 **C** *Mus musculus*
Noderer *et al.* GFP dataset (n=65,536)

799 ...C...GUNNNNNN̲A̲U̲G̲N̲N̲C̲G̲A̲A̲U̲U...G GFP

$R_s = -0.28$ $R_s = 0.12$

Fig 1. Correlations between the opening energies of translation initiation sites and protein abundance are stronger than that of minimum free energy. (A) For *E. coli*, the opening energy at the region −30:18 shows the strongest correlation with protein abundance (also see Fig 2B or Supplementary Fig S1A, sub-sequence l=48 at position i=18). For this analysis, we used a representative GFP expression dataset from Cambray et al. (2018). The reporter library consists of GFP fused in-frame with a library of 96-nt upstream sequences (N=14,425). The minimum free energy −30:30 shown was determined by Cambray et al. (right panel). **(B)** For *S. cerevisiae*, the opening energy −7:89 shows the strongest correlation with protein abundance (also see Supplementary Fig S1B, sub-sequence l=96 at position i= 89). For this analysis, we used the YFP expression dataset from Dvir et al. (2013). The YFP reporter library consists of 2,041 random decameric nucleotides inserted at the upstream of YFP start codon. The minimum free energy −15:50 was previously shown to correlate the best with protein abundance (right panel). **(C)** For *M. musculus*, the opening energy −8:11 shows the strongest correlation with protein abundance (also see Supplementary Fig S1C, sub-sequence l=19 at position i=11). For this analysis, we used the GFP expression dataset from Noderer et al. (2014). The GFP reporter library consists of 65,536 random hexameric and dimeric nucleotides inserted at the upstream and downstream of GFP start codon, respectively. The minimum free energy −30:30 was shown (right panel). See also Supplementary Table S1. R_s, Spearman's rho. Bonferroni adjusted P-values are statistically significant ($<2.2×10^{-16}$) for the correlations between opening energies and protein abundances shown in the left panels.

Fig 2. Opening energies of regions surrounding the Shine-Dalgarno and start codons are predictive of protein expression in *E. coli*. (A) Schematic representation of a transcript sub-sequence l at position i for the calculation of opening energy. For example, sub-sequence l=10 at position i=10 corresponds to the region 1:10. **(B)** Correlation between the opening energies for the sub-sequences of GFP transcripts and protein abundance. The opening energy at the region −30 to 18 nt (sub-sequence l=48 at position i=18, green crosshair) shows the strongest correlation with protein abundance [R_s=−0.65; N=14,425, GFP expression dataset of Cambray et al. (2018)]. For this dataset, the reporter plasmid used is pGC4750, in which the promoter and ribosomal binding site are oFAB1806 inducible promoter and oFAB1173/BCD7, respectively. **(C)** Prediction accuracy of the expression outcomes of the PSI:Biology targets using opening energy (N=11,430). The opening energy at the region −23:24 (sub-sequence l=47 at position i=24, green crosshair) shows the highest prediction accuracy score (AUC=0.70). For this dataset, the expression vector used is pET21_NESG, in which the promoter and fusion tag are T7lac and C-terminal His tag, respectively. **(D)** Comparison between the correlations and AUC scores by sub-sequence region taken from the above analyses. The sub-sequence regions that have strong correlations are likely to have high AUC scores, whereas the sub-sequence regions that have no correlations are likely not useful in prediction of the expression outcomes. **(E)** Correlation between the opening energies for the sub-sequences of *E. coli* transcripts and protein abundance. The transcripts used for this analysis are protein-coding sequences concatenated with 50 and 10 nt located upstream and downstream, respectively. The opening energy at the region −25:16 (sub-sequence l=41 at position i=16, green crosshair) shows the strongest correlation with protein abundance (R_s=−0.17; N=3,725, PaxDb integrated proteomics dataset). See also Supplementary Table S1. R_s, Spearman's rho.

Fig 3. Accessibility of translation initiation sites is the strongest predictor of heterologous protein expression in *E. coli*. (A) mRNA features ranked by Gini importance for random forest classification of the expression outcomes of the PSI:Biology targets (N=8,780 and 2,650, 'success' and 'failure' groups, respectively). The features associated with translation initiation rate (purple; opening energy −24:24, minimum free energy −30:30, and mRNA:ncRNA avoidance 1:30) have higher scores than the feature associated with translation elongation rate [blue; tRNA adaptation index (tAI), codon context (CC), codon adaptation index (CAI), G+C content (%), and IХnos]. The IХnos scores are translation elongation rates predicted using a neural network model trained with ribosome profiling data (Supplementary Fig S3). **(B)** ROC analysis shows that accessibility (opening energy −24:24) has the highest classification accuracy. The AUC scores with 95% confidence intervals are shown. See also Supplementary Table S1. **(C)** Accessibility (opening energy −24:24) is the best feature in explaining the expression outcomes. Relationships between the features and expression outcomes represented as squared Spearman's rho (R_s^2).

Fig 4. The yields of heterologous protein productions are tunable by synonymous codon changes in the first nine codons. (A) GFP level strongly correlates with accessibility, i.e., anti-correlates with opening energy (R_s=−0.53, P=3.4×10^{-3}; N=29). The protein levels of GFP, *Renilla* luciferase (RLuc), an antibody fragment and an archaebacterial dioxygenase were transformed using z-score method. The GFP and RLuc levels were derived from the average values of at least two and three independent biological replicates, respectively. Black outlines denote wild-type sequences. **(B)** Coarse-grained simulation of a protein production experiment by modelling cell growth, transcription, translation, and turnovers, given that translation initiation sites with opening energies less than or equal to 12 kcal/mol is optimum. The *in silico* model shows a similar trend of protein production as the wet-lab experimental results. Unfilled and filled (purple) circles denote the *in silico* replicates and their corresponding average values, respectively (R_s=−0.75, P=2.8×10^{-9}). **(C)** The expression of RLuc can be improved, despite its poor solubility in *E. coli* (Supplementary Fig S8). Opening energies are shown next to labels. The luciferase activities of commercially designed RLuc reporter genes (full-length sequence optimisation) and TIsigner (9.9 kcal/mol) are significantly higher than the wild-type luciferase (Mann-Whitney U tests, P=9.1×10^{-3}). No significant differences were observed between the commercial designs and TIsigner (9.9 kcal/mol). Error bars denote standard deviation of three independent biological replicates. **(D)** Densitometric analysis of previously published Western

892 blots shows that the yields of an antibody fragment and an archaebacterial dioxygenase can be
893 improved by synonymous codon changes within the first six codons (*49*). A RTS *E. coli* cell-free
894 expression system was used. The processed data are available Supplementary Table S2. AU,
895 arbitrary unit; R_s, Spearman's rho; WT, wild-type.