DYNLRB1 is Essential for Dynein Mediated Transport and Neuronal Survival

Marco Terenzio[1,2,*], Agostina Di Pizio[1], Ida Rishal[1], Letizia Marvaldi[1], Pierluigi Di Matteo[1], Riki Kawaguchi[3], Giovanni Coppola[3], Giampietro Schiavo[4,5,6], Elizabeth M.C. Fisher[4], Mike Fainzilber[1]

[1] Department of Biomolecular Sciences, Weizmann Institute of Science, Rehovot 76100, Israel

[2] Molecular Neuroscience Unit, Okinawa Institute of Science and Technology, Kunigami, Okinawa 904-0412, Japan

[3] Departments of Psychiatry and Neurology, Semel Institute for Neuroscience and Human Behavior, University of California Los Angeles, Los Angeles, CA 90095, U.S.A.

[4] Department of Neuromuscular Diseases, UCL Queen Square Institute of Neurology, Queen Square, London, WC1N 3BG, U.K.

[5] UK Dementia Research Institute, University College London, London WC1E 6BT, U.K.

[6] Discoveries Centre for Regenerative and Precision Medicine, University College London Campus, London WC1N 3BG, U.K.

[*]*Corresponding author:* marco.terenzio@weizmann.ac.il
Tel +972 8 934 4474
Fax +972 8 934 4118

SUMMARY

The cytoplasmic dynein motor complex transports essential signals and organelles from the cell periphery to perinuclear region, hence is critical for the survival and function of highly polarized cells such as neurons. Dynein Light Chain Roadblock-Type 1 (DYNLRB1) is thought to be an accessory subunit required for specific cargos, but here we show that it is essential for general dynein-mediated transport and sensory neuron survival. Homozygous *Dynlrb1* null mice are not viable and die during early embryonic development. Furthermore, heterozygous or adult knockdown animals display reduced neuronal growth, and selective depletion of *Dynlrb1* in proprioceptive neurons compromises their survival. Conditional depletion of *Dynlrb1* in sensory neurons causes deficits in several signaling pathways, including β-catenin subcellular localization, and a severe impairment in the axonal transport of both lysosomes and retrograde signaling endosomes. Hence, DYNLRB1 is an essential component of the dynein complex.

INTRODUCTION

Molecular motors perform a critical function in eukaryotic cells, ensuring the anterograde (kinesins) and retrograde (dyneins) transport of cargoes, organelles and cellular components. Dynein motors are divided into two major classes: axonemal dyneins, which are involved in ciliary and flagellar movement, and cytoplasmic dyneins, minus-end microtubule molecular motors involved in vesicular transport, cell migration, cell division and maintenance of Golgi integrity (Hirokawa et al., 2010; Kardon and Vale, 2009; Reck-Peterson et al., 2018). Cytoplasmic dynein is the main retrograde motor in all eukaryotic cells and is particularly important in highly polarized cells such as neurons, where it carries essential signals and organelles from the periphery to the cell body (Terenzio et al., 2017). Cytoplasmic dynein consists of two heavy chains (HC) associated with intermediate chains (IC), which provide direct binding to dynactin (Karki and Holzbaur, 1995; King et al., 2003; Siglin et al., 2013), four light intermediate chains (LIC) and several light chains (LC) (Reck-Peterson et al., 2018). There are three families of dynein light chains, LC7/road-block, LC8 and Tctx1/rp3, the latter playing a role in the direct binding of cargoes to the dynein motor complex (Olenick and Holzbaur, 2019; Reck-Peterson et al., 2018; Tai et al., 1999).

Dynein Light Chain Roadblock-Type 1 (DYNLRB1), also known as Km23, and DNLC2A (Jiang et al., 2001; Tang, 2002), was first identified in Drosophila during a genetic screen for axonal transport mutants (Bowman et al., 1999). *Roadblock* mutants exhibited defects in intracellular transport, including intra-axonal accumulation of synaptic cargoes, severe axonal degeneration and aberrant chromosome segregation (Bowman et al., 1999). DYNLRB protein sequence is highly conserved among different species; with two isoforms, DYNLRB1 and DYNLRB2, sharing 98% sequence similarity in mammals (Susalka et al., 2002). Mammalian DYNLRB1 has been studied mainly as an adaptor linking specific modules to the dynein complex, including SMAD2 complexes activated by TGFβ receptors (Jin et al., 2005, 2007, 2009, 2012, 2013; Tang, 2002) and Rab6 or N-acetyl-D-glucosamine kinase (NAGK) for interactions with the Golgi compartment (Islam et al., 2015a, 2015b; Wanschers et al., 2008).

In previous work, we identified a role for the dynein complex in regulating axon growth rates (Perry et al., 2016; Rishal et al., 2012). A screen of all dynein accessory subunits now shows that knockdown of *Dynlrb1* reduces neurite outgrowth in cultured sensory neurons. Surprisingly, a complete knockout of *Dynlrb1* was found to be embryonic lethal, suggesting the existence of non-redundant functions between DYNLRB1 and DYNLRB2. Conditional knockout or acute knockdown of *Dynlrb1* compromised survival of proprioceptive neurons, together with changes in subcellular localization of transported cargos, and major impairments in the axonal transport of lysosomes and retrograde signaling endosomes. Thus, DYNLRB1 is an essential component of the dynein complex, and the conditional allele described here enables selective targeting of dynein functions.

RESULTS

Depletion of *Dynlrb1* impairs neurite outgrowth in cultured DRG neurons

We performed an siRNA screen in cultured dorsal root ganglion (DRG) neurons to investigate the effects of downregulating individual dynein subunits on axonal growth. Knockdowns of *Dync1i2*, *Dync1li2*, *Dynlrb1* and *Dctn6* reduced the extent of axon outgrowth in cultured DRG neurons (**Figure 1A**), with *Dynlrb1* knockdown providing the most robust effect (**Figure 1A, B**). We then sought to validate this finding in a knockout mouse model for *Dynlrb1* generated by the European Conditional Mouse Mutagenesis Program (EUCOMM) as part of the International Mouse Knockout Consortium. The gene targeted $Dynlrb1^{tm1a(EUCOMM)Wtsi}$ allele, hereafter referred to as $Dynlrb^{tm1a}$, results in a complete knockout of *Dynlrb1* with concomitant expression of β-galactosidase from the same locus; this allele also allows the subsequent generation of a *floxed* conditional allele ($Dynlrb^{tm1c}$) upon application of Flp recombinase (**Figure S1A**)(Skarnes et al., 2011). Homozygous complete knockout mice ($Dynlrb^{tm1a/tm1a}$) were not viable, and analyses of early pregnancies showed loss of homozygous null embryos before E9.5, while heterozygous animals were viable. The β-galactosidase expression pattern in heterozygous mice showed that DYNLRB1 was highly expressed in DRGs of E12.5 mouse embryos (**Figure S1B**). We then crossed ($Dynlrb^{tm1a/+}$) heterozygous mice with Thy1-yellow fluorescent protein (YFP) mice (Feng et al., 2000) to allow live imaging of

DRG neuron growth in culture. Similarly to *Dynlrb1* knockdown neurons (**Figure 1A, B**), Thy1-YFP- *Dynlrb$^{tm1a/+}$* DRG neurons had significantly reduced axon outgrowth compared to littermate controls (**Figure 1C, D**).

Dynlrb1 deletion affects survival of proprioceptive neurons

We proceeded to cross *Dynlrb$^{tm1a/+}$* heterozygotes with mice expressing the Flp recombinase (Farley et al., 2000) to recombine the FRT sites, thereby removing the LacZ sequence (Sadowski, 1995). This generated a conditional allele, *Dynlrb^{tm1c}* in which *LoxP* sites flank exon 3 of *Dynlrb1* (**Figure S1A**). *Dynlrb^{tm1c}* mice were then crossed with a *RNX3-Cre* driver line (Levanon et al., 2002), to examine the effect of *Dynlrb1* deletion in proprioceptive sensory neurons. In contrast to the full knockout, *RNX3-Dynlrb$^{tm1d/tm1d}$,*, hereafter referred to as *RNX3-Dynlrb1$^{-/-}$*, mice were viable to adulthood. However, these mice displayed abnormal hind limb posture (**Figure 2A**), and an uncoordinated walking pattern with abdomen posture close to the ground (**Movie S1**). A battery of tests of motor activity and proprioception showed that *RNX3-Dynlrb1$^{-/-}$* mice were unable to balance themselves in the rotarod (**Figure 2B**) and exhibited an abnormal walking pattern in the catwalk (**Figures 2C, D**). There was no difference between genotypes in basal motor activity (**Figure S2A**), but *RNX3-Dynlrb1$^{-/-}$* mice displayed lower speed and covered less distance than their wild type counterparts in the open field (**Figure S2B-D**). Overall, these tests reveal a strong impairment of proprioception in adult *RNX3-Dynlrb1$^{-/-}$* mice.

To understand the mechanistic basis of this impairment, we performed histological analyses on *RNX3-Dynlrb1$^{-/-}$* mice. There was no difference between genotypes in neuromuscular junction size or numbers (**Figure S2E, F**), and in total (Tuj1 positive) or nociceptive (CGRP positive) skin innervation (**Figure S2G, H**). However, the number of proprioceptive NFH-positive neurons in DRG ganglia was greatly reduced in *RNX3-Dynlrb1$^{-/-}$* mice (**Figure 2E, F**), while CGRP-positive nociceptive neuron numbers did not differ between genotypes (**Figure 2E, F**). These data reaffirm the specificity of the *RNX3-Cre* driver and indicate that the proprioceptive impairment observed in *RNX3-Dynlrb1$^{-/-}$* mice is due to a loss of this class of sensory neurons.

Acute viral-mediated knockdown of *Dynlrb1* causes motor-balance defects *in vivo*

The *RNX3-Cre* driver is activated early in development. In order to determine whether *Dynlrb1* deletion has similar effects upon acute depletion in adult proprioceptive neurons, wild type mice were injected intrathecally with adeno-associated virus (AAV) harboring an shRNA against *Dynlrb1* or an shControl sequence. We used AAV serotype 9 for neuronal-specific knockdown (Naso et al., 2017; Snyder et al., 2011), and observed transduction rates of 25-35% in DRG ganglia (**Figure S3A, B**). Efficiency of the knockdown was found to be in the range of 40% decrease in *Dynlrb1* mRNA levels in cultured DRG (**Figure S3C**). 32 days after the injection, mice transduced with AAV-shDynlrb1 developed motor problems and abnormal hind limb posture (**Figure S3D**). AAV-shDynlrb1 transduced mice showed significant impairment in rotarod and catwalk analyses when compared with AAV-shControl transduced animals (**Figure S3E-G**). Overall these data show that *Dynlrb1* is critical for adult sensory neurons and its depletion by knockdown recapitulates the phenotype observed in the *RNX3-Dynlrb1^{-/-}* mice.

Gene expression changes in *Dynlrb^{tm1a/+}* DRG neurons

We then sought to obtain insights on the physiological role(s) of DYNLRB1. Since complete depletion of DYNLRB1 is lethal, we focused on *Dynlrb^{tm1a/+}* DRG neurons, which are viable and display reduced neurite outgrowth. RNA-seq of *Dynlrb^{tm1a/+}* versus wild type DRG neurons at 12, 24 and 48 hr in culture showed almost no differences between the two genotypes at 12 and 24 hr, while significant differences were observed at 48 hr (**Figure 3A, B, Table S1**). Bioinformatic analyses of the differentially expressed genes dataset indicated changes in a number of signaling and transcriptional pathways, including the canonical Wnt – β-catenin pathway (**Figure 3B, C, Table S2**). Interestingly, direct interaction between a Dynein intermediate chain and β-catenin has been shown in epithelial cells (Ligon et al., 2001), where the complex tethers microtubules at adherens junctions. Thus, DYNLRB1 might be part of a dynein complex carrying β-catenin from the periphery to the nucleus. Indeed, β-catenin nuclear accumulation at 48 h in culture is reduced in *Dynlrb^{tm1a/+}* DRG neurons compared to their wild type littermates (**Figure 3C, D**).

Depletion of *Dynlrb1* has significant effects on axonal retrograde transport

The broad spectrum of gene expression changes in *Dynlrb*$^{tm1a/+}$ DRG neurons argues for a general role of this subunit in the dynein complex. We therefore investigated the effect of DYNLRB1 depletion on dynein-mediated axonal retrograde transport, using acute depletion of *Dynlrb1* in culture to circumvent the loss of neurons observed in *RNX3-Dynlrb1*$^{-/-}$ mice. To this end, we transduced *Dynlrb*$^{tm1c/tm1c}$ DRG neurons with Adenovirus 5 (Ad5) expressing a Cre-GFP, to induce recombination of the *Dynlrb*^{tm1c} allele to *Dynlrb*^{tm1d} in infected cells. We first optimized the transduction protocol using neurons from the Ai9 mouse (Madisen et al., 2010), which expresses td-Tomato under the control of a LoxP-flanked STOP cassette and showed that 72 hr of incubation with Ad5 Cre-GFP are sufficient to express the Cre and trigger the recombination of LoxP sites in cultured DRG neurons (**Figure S4A, B**). We then tracked axonal retrograde transport of acidic organelles, including lysosomes, late endosomes and autophagosomes in Ad5-Cre: *Dynlrb*$^{tm1d/tm1d}$ DRG neurons using Lysotracker Red. Indeed, Ad5-Cre: *Dynlrb*$^{tm1d/tm1d}$ transduced neurons 96 h in culture showed a significant impairment of axonal retrograde transport of these organelles, both in terms of velocity of individual carriers and the overall fraction of moving carriers (**Figure 4A-C**).

We further examined transport properties using a marker for neurotrophin signaling endosomes, the binding fragment of tetanus neurotoxin (H$_C$T) fragment (Deinhardt et al., 2006; Lalli and Schiavo, 2002). Whilst we could not detect any significant differences in the rates of transport of H$_C$T-positive retrograde axonal carriers in Ad5-Cre: *Dynlrb*$^{tm1d/tm1d}$ neurons (**Figure 4D, E**), the percentage of stationary carriers was significantly increased (**Figure 4F**), as was also found for Lysotracker Red-positive organelles (**Figure 4C**). Thus, depletion of DynlRb1 has significant effects on retrograde transport of a broad spectrum of dynein cargos.

DISCUSSION

The dynein complex has essential roles in all multicellular organisms, most prominently in long distance axonal transport in neurons. Studies on specific roles of dynein carriers have suffered from a lack of appropriate mouse models, since

homozygous deletion or mutation of *Dync1h1* results in early embryonic death (Hafezparast et al., 2003; Harada et al., 1998). We have characterized a new conditional knockout model for the DYNLRB1 subunit of dynein and use this model to show that DYNLRB1 is required for retrograde axonal transport of a broad spectrum of cargos in sensory axons, as well as for neuronal survival. These findings were surprising given the view in the literature that DYNRLB1 should function mainly as a binding adaptor for TGFβ signaling complexes and other specific cargos (Jin et al., 2005, 2007, 2009, 2012, 2013; Tang, 2002), whereas our data suggests that DYNLRB1 is essential for dynein-mediated axonal transport of lysosomes, late endosomes and signaling endosomes.

The hindlimb proprioception phenotype observed in both *RNX3-Dynlrb1$^{-/-}$* mice and *Dynlrb1* viral-mediated knockdown is highly reminiscent of dynein heavy chain mutant models, such as *Dync1h1^{Loa}* and *Dync1h1^{Cra1}* (Dupuis et al., 2009; Hafezparast et al., 2003). Both models show deficits in endosomal axonal retrograde transport (Hafezparast et al., 2003) and proprioception (Dupuis et al., 2009; Ilieva et al., 2008), as we have shown above for knockout or knockdown of *Dynlrb1*. In addition, depletion of *Roadblock* in *Drosophila* was described to cause defects in intra-axonal accumulation of cargoes and severe axonal degeneration (Bowman et al., 1999). The distal enrichment of axonal accumulations observed in *Roadblock* mutants was suggested to arise from axonal transport deficits originating at the synapse (Bowman et al., 1999). Moreover, incorporation of DYNLRB1 and DYNLRB2 into the Dynein-2 complex, which drives retrograde protein trafficking in primary cilia, was recently shown to be crucial for dynein-2 transport function (Tsurumi et al., 2019). While the precise mechanism by which dynein activity is impaired upon loss of DYNLRB1 is yet to be determined, it is interesting to note that this subunit can bind directly to core components of the dynein complex such as the intermediate chains DYNCI1 and DYNCI2 (IC74-1 and IC74-2) (Susalka et al., 2002). The latter directly associate with the dynactin complex, which facilitates interactions of numerous cargos with the main motor complex, and expedites processive transport along microtubules (Karki and Holzbaur, 1995; Reck-Peterson et al., 2018).

Our findings show that *Dynlrb1* is critical not only for embryonic development, but also for maintenance of sensory neurons in adulthood. Deficits in dynein functions have been implicated in the pathogenesis of neurodegenerative diseases such as Amyotrophic Lateral Sclerosis (ALS) (Bilsland et al., 2010; Gershoni-Emek et al., 2016; Hoang et al., 2017; Schiavo et al., 2013; Teuling et al., 2008; Xie et al., 2015), and an increasing number of human neurodegenerative pathologies have been linked to mutations of dynein complex genes (Terenzio et al., 2018). In addition, mutations and/or disruption of the function of the dynactin complex has been shown to be connected to motor neuron degeneration and ALS (Kuźma-Kozakiewicz et al., 2013; LaMonte et al., 2002; Moore et al., 2009; Münch et al., 2004; Puls et al., 2003). The new mouse model described herein will expand the options for studying such roles of dynein, using conditional targeting of Dynlrb1 for both temporal and tissue/cell type specificity in such studies *in vivo*.

Acknowledgments: We thank Nitzan Korem and Nataliya Okladnikov for excellent technical assistance; Vladmir Kiss for professional microscopy support; Dalia Gordon, Indrek Koppel, Nicolas Panayotis, Stefanie Alber and Michael Tsoory for helpful comments and advice, Avraham Yaron for the kind gift of the Aig9 mice and Avri Ben-Ze'ev for the kind gift of a β-catenin antibody. We thank the Medical Research Council Mammalian Genetics Unit, Harwell, UK for providing the *Dynlrb1$^{tm1a(EUCOMM)Wtsi}$* mice.

Funding: This work was supported by funding from the European Research Council (Neurogrowth, to M.F.), the Dr. Miriam and Sheldon G. Adelson Medical Research Foundation (to M.F. and G.C.), a Wellcome Trust Senior Investigator Award (107116/Z/15/Z) (to GS), the European Union's Horizon 2020 Research and Innovation programme under grant agreement 739572 (to GS), a UK Dementia Research Institute Foundation award (to GS), and the Medical Research Council (MRC, to E.M.C.F.). M.T. was supported by a Koshland senior postdoctoral fellowship. M.F. is the incumbent of the Chaya Professorial Chair in Molecular Neuroscience at the Weizmann Institute of Science.

Author contributions: M.F. and M.T. designed the study. M.T., A.D.P., I.R., L.M., P.D.M., and R.K. performed experiments and data analyses. R.K. carried out RNA sequencing. M.F., E.M.C.F., G.C., and G.S. supervised research. M.F. and M.T. wrote the initial manuscript draft. All authors revised the manuscript and approved the final version.

REFERENCES

Ben-Yaakov, K., Dagan, S.Y., Segal-Ruder, Y., Shalem, O., Vuppalanchi, D., Willis, D.E., Yudin, D., Rishal, I., Rother, F., Bader, M., et al. (2012). Axonal transcription factors signal retrogradely in lesioned peripheral nerve. EMBO J. *31*, 1350–1363.

Bilsland, L.G., Sahai, E., Kelly, G., Golding, M., Greensmith, L., and Schiavo, G. (2010). Deficits in axonal transport precede ALS symptoms in vivo. Proc. Natl. Acad. Sci. U.S.A. *107*, 20523–20528.

Bowman, A.B., Patel-King, R.S., Benashski, S.E., McCaffery, J.M., Goldstein, L.S., and King, S.M. (1999). Drosophila roadblock and Chlamydomonas LC7: a conserved family of dynein-associated proteins involved in axonal transport, flagellar motility, and mitosis. J. Cell Biol. *146*, 165–180.

Cooper, A., Grigoryan, G., Guy-David, L., Tsoory, M.M., Chen, A., and Reuveny, E. (2012). Trisomy of the G protein-coupled K+ channel gene, Kcnj6, affects reward mechanisms, cognitive functions, and synaptic plasticity in mice. PNAS *109*, 2642–2647.

Crawley, J.N. (2008). Behavioral Phenotyping Strategies for Mutant Mice. Neuron *57*, 809–818.

Dagan, S.Y., Tsoory, M.M., Fainzilber, M., and Panayotis, N. (2016). COLORcation: A new application to phenotype exploratory behavior models of anxiety in mice. Journal of Neuroscience Methods *270*, 9–16.

Deinhardt, K., Salinas, S., Verastegui, C., Watson, R., Worth, D., Hanrahan, S., Bucci, C., and Schiavo, G. (2006). Rab5 and Rab7 control endocytic sorting along the axonal retrograde transport pathway. Neuron *52*, 293–305.

Dupuis, L., Fergani, A., Braunstein, K.E., Eschbach, J., Holl, N., Rene, F., Gonzalez De Aguilar, J.-L., Zoerner, B., Schwalenstocker, B., Ludolph, A.C., et al. (2009). Mice with a mutation in the dynein heavy chain 1 gene display sensory neuropathy but lack motor neuron disease. Experimental Neurology *215*, 146–152.

Farley, F.W., Soriano, P., Steffen, L.S., and Dymecki, S.M. (2000). Widespread recombinase expression using FLPeR (flipper) mice. Genesis *28*, 106–110.

Feng, G., Mellor, R.H., Bernstein, M., Keller-Peck, C., Nguyen, Q.T., Wallace, M., Nerbonne, J.M., Lichtman, J.W., and Sanes, J.R. (2000). Imaging neuronal subsets in transgenic mice expressing multiple spectral variants of GFP. Neuron *28*, 41–51.

Gershoni-Emek, N., Mazza, A., Chein, M., Gradus-Pery, T., Xiang, X., Li, K.W., Sharan, R., and Perlson, E. (2016). Proteomic Analysis of Dynein-Interacting Proteins in Amyotrophic Lateral Sclerosis Synaptosomes Reveals Alterations in the RNA-Binding Protein Staufen1. Mol. Cell Proteomics *15*, 506–522.

Hafezparast, M., Klocke, R., Ruhrberg, C., Marquardt, A., Ahmad-Annuar, A., Bowen, S., Lalli, G., Witherden, A.S., Hummerich, H., Nicholson, S., et al. (2003). Mutations in dynein link motor neuron degeneration to defects in retrograde transport. Science *300*, 808–812.

Hanz, S., Perlson, E., Willis, D., Zheng, J.-Q., Massarwa, R., Huerta, J.J., Koltzenburg, M., Kohler, M., van-Minnen, J., Twiss, J.L., et al. (2003). Axoplasmic importins enable retrograde injury signaling in lesioned nerve. Neuron *40*, 1095–1104.

Harada, A., Takei, Y., Kanai, Y., Tanaka, Y., Nonaka, S., and Hirokawa, N. (1998). Golgi Vesiculation and Lysosome Dispersion in Cells Lacking Cytoplasmic Dynein. The Journal of Cell Biology *141*, 51–59.

Hirokawa, N., Niwa, S., and Tanaka, Y. (2010). Molecular motors in neurons: transport mechanisms and roles in brain function, development, and disease. Neuron *68*, 610–638.

Hoang, H.T., Schlager, M.A., Carter, A.P., and Bullock, S.L. (2017). DYNC1H1 mutations associated with neurological diseases compromise processivity of dynein-dynactin-cargo adaptor complexes. Proc. Natl. Acad. Sci. U.S.A. *114*, E1597–E1606.

Ilieva, H.S., Yamanaka, K., Malkmus, S., Kakinohana, O., Yaksh, T., Marsala, M., and Cleveland, D.W. (2008). Mutant dynein (Loa) triggers proprioceptive axon loss that extends survival only in the SOD1 ALS model with highest motor neuron death. PNAS *105*, 12599–12604.

Islam, M.A., Sharif, S.R., Lee, H., Seog, D.-H., and Moon, I.S. (2015a). N-acetyl-D-glucosamine kinase interacts with dynein light-chain roadblock type 1 at Golgi outposts in neuronal dendritic branch points. Exp. Mol. Med. *47*, e177.

Islam, M.A., Sharif, S.R., Lee, H., and Moon, I.S. (2015b). N-Acetyl-D-Glucosamine Kinase Promotes the Axonal Growth of Developing Neurons. Mol. Cells *38*, 876–885.

Jiang, J., Yu, L., Huang, X., Chen, X., Li, D., Zhang, Y., Tang, L., and Zhao, S. (2001). Identification of two novel human dynein light chain genes, DNLC2A and DNLC2B, and their expression changes in hepatocellular carcinoma tissues from 68 Chinese patients. Gene *281*, 103–113.

Jin, Q., Ding, W., Staub, C.M., Gao, G., Tang, Q., and Mulder, K.M. (2005). Requirement of km23 for TGFbeta-mediated growth inhibition and induction of fibronectin expression. Cell. Signal. *17*, 1363–1372.

Jin, Q., Ding, W., and Mulder, K.M. (2007). Requirement for the dynein light chain km23-1 in a Smad2-dependent transforming growth factor-?? signaling pathway. Journal of Biological Chemistry *282*, 19122–19132.

Jin, Q., Gao, G., and Mulder, K.M. (2009). Requirement of a dynein light chain in TGFbeta/Smad3 signaling. J. Cell. Physiol. *221*, 707–715.

Jin, Q., Ding, W., and Mulder, K.M. (2012). The TGFβ receptor-interacting protein km23-1/DYNLRB1 plays an adaptor role in TGFβ1 autoinduction via its association with Ras. J. Biol. Chem. *287*, 26453–26463.

Jin, Q., Zhong, Y., and Mulder, K.M. (2013). Requirement for protein kinase A in the phosphorylation of the TGFβ receptor-interacting protein km23-1 as a component of TGFβ downstream effects. Exp. Cell Res. *319*, 897–907.

Johnson, K.R., Lewis, J.E., Li, D., Wahl, J., Soler, A.P., Knudsen, K.A., and Wheelock, M.J. (1993). P- and E-cadherin are in separate complexes in cells expressing both cadherins. Exp. Cell Res. *207*, 252–260.

Kardon, J.R., and Vale, R.D. (2009). Regulators of the cytoplasmic dynein motor. Nat. Rev. Mol. Cell Biol. *10*, 854–865.

Karki, S., and Holzbaur, E.L. (1995). Affinity chromatography demonstrates a direct binding between cytoplasmic dynein and the dynactin complex. J. Biol. Chem. *270*, 28806–28811.

King, S.J., Brown, C.L., Maier, K.C., Quintyne, N.J., and Schroer, T.A. (2003). Analysis of the Dynein-Dynactin Interaction In Vitro and In Vivo. Mol Biol Cell *14*, 5089–5097.

Kuźma-Kozakiewicz, M., Chudy, A., Kaźmierczak, B., Dziewulska, D., Usarek, E., and Barańczyk-Kuźma, A. (2013). Dynactin Deficiency in the CNS of Humans with Sporadic ALS and Mice with Genetically Determined Motor Neuron Degeneration. Neurochem Res *38*, 2463–2473.

Lalli, G., and Schiavo, G. (2002). Analysis of retrograde transport in motor neurons reveals common endocytic carriers for tetanus toxin and neurotrophin receptor p75NTR. J. Cell Biol. *156*, 233–239.

LaMonte, B.H., Wallace, K.E., Holloway, B.A., Shelly, S.S., Ascaño, J., Tokito, M., Winkle, T.V., Howland, D.S., and Holzbaur, E.L.F. (2002). Disruption of Dynein/Dynactin Inhibits Axonal Transport in Motor Neurons Causing Late-Onset Progressive Degeneration. Neuron *34*, 715–727.

Levanon, D., Bettoun, D., Harris-Cerruti, C., Woolf, E., Negreanu, V., Eilam, R., Bernstein, Y., Goldenberg, D., Xiao, C., Fliegauf, M., et al. (2002). The Runx3 transcription factor regulates development and survival of TrkC dorsal root ganglia neurons. EMBO J. *21*, 3454–3463.

Ligon, L.A., Karki, S., Tokito, M., and Holzbaur, E.L. (2001). Dynein binds to beta-catenin and may tether microtubules at adherens junctions. Nat. Cell Biol. *3*, 913–917.

Madisen, L., Zwingman, T.A., Sunkin, S.M., Oh, S.W., Zariwala, H.A., Gu, H., Ng, L.L., Palmiter, R.D., Hawrylycz, M.J., Jones, A.R., et al. (2010). A robust and high-throughput Cre reporting and characterization system for the whole mouse brain. Nat. Neurosci. *13*, 133–140.

Marvaldi, L., Thongrong, S., Kozłowska, A., Irschick, R., Pritz, C.O., Bäumer, B., Ronchi, G., Geuna, S., Hausott, B., and Klimaschewski, L. (2015). Enhanced axon outgrowth and improved long-distance axon regeneration in sprouty2 deficient mice. Dev Neurobiol *75*, 217–231.

Moore, J.K., Sept, D., and Cooper, J.A. (2009). Neurodegeneration mutations in dynactin impair dynein-dependent nuclear migration. PNAS *106*, 5147–5152.

Münch, C., Sedlmeier, R., Meyer, T., Homberg, V., Sperfeld, A.D., Kurt, A., Prudlo, J., Peraus, G., Hanemann, C.O., Stumm, G., et al. (2004). Point mutations of the p150 subunit of dynactin (DCTN1) gene in ALS. Neurology *63*, 724–726.

Naso, M.F., Tomkowicz, B., Perry, W.L., and Strohl, W.R. (2017). Adeno-Associated Virus (AAV) as a Vector for Gene Therapy. BioDrugs *31*, 317–334.

Neufeld-Cohen, A., Tsoory, M.M., Evans, A.K., Getselter, D., Gil, S., Lowry, C.A., Vale, W.W., and Chen, A. (2010). A triple urocortin knockout mouse model reveals an essential role for urocortins in stress recovery. PNAS *107*, 19020–19025.

Olenick, M.A., and Holzbaur, E.L.F. (2019). Dynein activators and adaptors at a glance. J Cell Sci *132*, jcs227132.

Perlson, E., Hanz, S., Ben-Yaakov, K., Segal-Ruder, Y., Seger, R., and Fainzilber, M. (2005). Vimentin-dependent spatial translocation of an activated MAP kinase in injured nerve. Neuron *45*, 715–726.

Perry, R.B.-T., Doron-Mandel, E., Iavnilovitch, E., Rishal, I., Dagan, S.Y., Tsoory, M., Coppola, G., McDonald, M.K., Gomes, C., Geschwind, D.H., et al. (2012). Subcellular knockout of importin β1 perturbs axonal retrograde signaling. Neuron *75*, 294–305.

Perry, R.B.-T., Rishal, I., Doron-Mandel, E., Kalinski, A.L., Medzihradszky, K.F., Terenzio, M., Alber, S., Koley, S., Lin, A., Rozenbaum, M., et al. (2016). Nucleolin-Mediated RNA Localization Regulates Neuron Growth and Cycling Cell Size. Cell Rep *16*, 1664–1676.

Puls, I., Jonnakuty, C., LaMonte, B.H., Holzbaur, E.L.F., Tokito, M., Mann, E., Floeter, M.K., Bidus, K., Drayna, D., Oh, S.J., et al. (2003). Mutant dynactin in motor neuron disease. Nat. Genet. *33*, 455–456.

Reck-Peterson, S.L., Redwine, W.B., Vale, R.D., and Carter, A.P. (2018). The cytoplasmic dynein transport machinery and its many cargoes. Nature Reviews Molecular Cell Biology *19*, 382.

Rishal, I., Kam, N., Perry, R.B.-T., Shinder, V., Fisher, E.M.C., Schiavo, G., and Fainzilber, M. (2012). A motor-driven mechanism for cell-length sensing. Cell Rep *1*, 608–616.

Sadowski, P.D. (1995). The Flp recombinase of the 2-microns plasmid of Saccharomyces cerevisiae. Prog. Nucleic Acid Res. Mol. Biol. *51*, 53–91.

Schiavo, G., Greensmith, L., Hafezparast, M., and Fisher, E.M.C. (2013). Cytoplasmic dynein heavy chain: the servant of many masters. Trends Neurosci. *36*, 641–651.

Siglin, A.E., Sun, S., Moore, J.K., Tan, S., Poenie, M., Lear, J.D., Polenova, T., Cooper, J.A., and Williams, J.C. (2013). Dynein and dynactin leverage their bivalent character to form a high-affinity interaction. PLoS ONE *8*, e59453.

Simcha, I., Shtutman, M., Salomon, D., Zhurinsky, J., Sadot, E., Geiger, B., and Ben-Ze'ev, A. (1998). Differential Nuclear Translocation and Transactivation Potential of β-Catenin and Plakoglobin. J Cell Biol *141*, 1433–1448.

Skarnes, W.C., Rosen, B., West, A.P., Koutsourakis, M., Bushell, W., Iyer, V., Mujica, A.O., Thomas, M., Harrow, J., Cox, T., et al. (2011). A conditional knockout resource for the genome-wide study of mouse gene function. Nature *474*, 337–342.

Snyder, B.R., Gray, S.J., Quach, E.T., Huang, J.W., Leung, C.H., Samulski, R.J., Boulis, N.M., and Federici, T. (2011). Comparison of Adeno-Associated Viral Vector Serotypes for Spinal Cord and Motor Neuron Gene Delivery. Human Gene Therapy *22*, 1129–1135.

Susalka, S.J., Nikulina, K., Salata, M.W., Vaughan, P.S., King, S.M., Vaughan, K.T., and Kevin Pfister, K. (2002). The roadblock light chain binds a novel region of the cytoplasmic dynein intermediate chain. Journal of Biological Chemistry *277*, 32939–32946.

Tai, A.W., Chuang, J.Z., Bode, C., Wolfrum, U., and Sung, C.H. (1999). Rhodopsin's carboxy-terminal cytoplasmic tail acts as a membrane receptor for cytoplasmic dynein by binding to the dynein light chain Tctex-1. Cell *97*, 877–887.

Tang, Q. (2002). A Novel Transforming Growth Factor-beta Receptor-interacting Protein That Is Also a Light Chain of the Motor Protein Dynein. Molecular Biology of the Cell *13*, 4484–4496.

Terenzio, M., Schiavo, G., and Fainzilber, M. (2017). Compartmentalized Signaling in Neurons: From Cell Biology to Neuroscience. Neuron *96*, 667–679.

Terenzio, M., Koley, S., Fisher, E.M.C., and Fainzilber, M. (2018). 11 - The use of mouse models to probe cytoplasmic dynein function. In Dyneins (Second Edition), S.M. King, ed. (Academic Press), pp. 234–261.

Teuling, E., van Dis, V., Wulf, P.S., Haasdijk, E.D., Akhmanova, A., Hoogenraad, C.C., and Jaarsma, D. (2008). A novel mouse model with impaired dynein/dynactin function develops amyotrophic lateral sclerosis (ALS)-like features in motor neurons and improves lifespan in SOD1-ALS mice. Hum. Mol. Genet. *17*, 2849–2862.

Tsurumi, Y., Hamada, Y., Katoh, Y., and Nakayama, K. (2019). Interactions of the dynein-2 intermediate chain WDR34 with the light chains are required for ciliary retrograde protein trafficking. Mol. Biol. Cell *30*, 658–670.

Wanschers, B., van de Vorstenbosch, R., Wijers, M., Wieringa, B., King, S.M., and Fransen, J. (2008). Rab6 family proteins interact with the dynein light chain protein DYNLRB1. Cell Motil. Cytoskeleton *65*, 183–196.

Xie, Y., Zhou, B., Lin, M.-Y., Wang, S., Foust, K.D., and Sheng, Z.-H. (2015). Endolysosomal Deficits Augment Mitochondria Pathology in Spinal Motor Neurons of Asymptomatic fALS Mice. Neuron *87*, 355–370.

Yu, H., Su, Y., Shin, J., Zhong, C., Guo, J.U., Weng, Y.-L., Gao, F., Geschwind, D.H., Coppola, G., Ming, G., et al. (2015). Tet3 regulates synaptic transmission and homeostatic plasticity via DNA oxidation and repair. Nat. Neurosci. *18*, 836–843.

FIGURE LEGENDS

Figure 1. Depletion of DYNLRB1 reduces axon outgrowth in DRG neurons.
(A) siRNA screen for effects of dynein complex subunits on neuronal growth (n > 100 cells for each). Positive hits (*Dync1i2, Dync1li2, Dynlrb1 and Dctn6*) were validated in at least three independent experiments (n > 500 cells each). ** p < 0.01, and *** p < 0.001 (Student's t test and one-way ANOVA). **(B)** Fluorescence images of cultured DRG neurons from adult Thy1-YFP mice transfected with either siControl or siDynlrb1. Neurons were re-plated 24 hr after siRNA transfection, and images were acquired 48h after re-plating. YFP signal in green. Scale bar, 200 μm. **(C)** Fluorescent images of cultured DRG neurons from wild type Thy1-YFP mice and Thy1-YFP- *Dynlrb$^{tm1a/+}$* mice (YFP signal in green). Cells were imaged every hour for a period of 48h. Scale bar, 100 μm. **(D)** Quantification of the time-lapse imaging experiment described in C. Mean ± SEM, *** p < 0.001, n = 3, two-way ANOVA.

Figure 2. Conditional Dynlrb1 depletion affects survival of proprioceptive neurons *in vivo.* **(A)** Abnormal clenched hind-limb phenotype in *RNX-Dynlrb1$^{-/-}$* mice suspended from the tail, compared to the typical limb extension of wild type animals. **(B)** Rotarod test (acceleration set at 20 rpm in 240 sec) showing that *RNX-Dynlrb1$^{-/-}$* mice are unable to balance themselves on the rod. Mean ± SEM, *** p<0.001, n>6 mice per genotype, two-way ANOVA. **(C)** Representative catwalk gait traces for wild type and *RNX-Dynlrb1$^{-/-}$* mice. The position of the paws are shown in different colors as indicated, RF - right front, RH - right hind, LF - left front, LH - left hind. **(D)** Step sequence regularity index and Paw stands from the catwalk gait analyses of C shows impairment in *RNX-Dynlrb1$^{-/-}$* mice compared to wild type. Mean ± SEM, * p<0.05 *** p<0.001, n>6 mice per genotype, one-way ANOVA followed by Tukey's HSD post hoc correction for multiple comparisons. **(E)** Representative DRG ganglia sections from wild type and *RNX-Dynlrb1$^{-/-}$* mice, stained with either NFH (green) for proprioceptive neurons or CGRP (red) for nociceptors. Scale bar, 50 μm. **(F)** Quantification of NFH and CGRP positive neurons from the experiment described in G. Mean ± SEM, * p<0.05, n=3 per genotype, unpaired t-test.

Figure 3. RNA-seq of cultured *Dynlrb*^{tm1a/+} DRG neurons highlights signaling deficits. (A) Bar charts representing the numbers of differentially expressed genes (contrast analysis in EdgeR, FDR<0.1) in wild type and *Dynlrb*$^{tm1a/+}$ DRG neurons at the indicated time points in culture. **(B)** heat map representing the differential expressed genes in wild type and *Dynlrb*$^{tm1a/+}$ mice at 48 hr in culture (Log10(FPKM)). **(C)** Ingenuity pathway analyses on the 48 hr differentially expressed genes dataset, ranked by –log(p values). The Wnt – β–catenin pathway (highlighted in red) was selected for further validation. **(D)** Representative cell bodies of wild type and *Dynlrb*$^{tm1a/+}$ DRG neurons cultured for 48 hr before immunostaining as shown. Scale bar, 20 μm. **(E)** Quantification of nuclear versus cytoplasmic β–catenin in the experiment described in C. Mean ± SEM, * p<0.05 ** p<0.01, n>3 mice per genotype, one-way ANOVA followed by Tukey's HSD post hoc correction for multiple comparisons.

Figure 4. Axonal retrograde transport deficits after *Dynlrb1* depletion. (A) Representative kymographs of Lysotracker Red tracking on wild type and *Dynlrb*$^{tm1c/tm1c}$ DRG neurons transduced for 96h in culture as indicated (Ad5-Cre, Adenovirus5 expressing Cre-GFP) for 96 hr. Diagonal lines indicate retrogradely moving carriers. **(B)** Velocity distributions from the experiment shown in A, Mean ± SEM, n>32 axons per group over 3 independent biological repeats. **(C)** Fraction of moving versus stationary carriers in the experiment described in A. Mean ± SEM, *** p<0.001, n>32 axons per group over 3 independent biological repeats, two-way ANOVA. **(D)** Representative kymographs of TeNT H$_C$T tracking on wild type and *Dynlrb*$^{tm1c/tm1c}$ DRG neurons transduced for 96 hr in culture as indicated. **(E)** Velocity distributions from the experiment shown in D. Mean ± SEM, n>36 axons per group over 3 independent biological repeats. **(F)** Fraction of moving versus stationary carriers in the experiment described in D. Mean ± SEM, ** p<0.01, n>36 axons per group over 3 independent biological repeats, two-way ANOVA.

SUPPLEMENTARY ITEMS AND FIGURE LEGENDS

Figure S1, related to Figure 1. Targeting Vector and Expression Profile of DynIrb1.
(A) Design of the targeting vector of the allele: *DynIrb1$^{tm1a(EUCOMM)Wtsi}$* from the International Knockout Mouse Consortium (IKMC). The unmodified allele creates a knockout due to the insertion of a β-Gal cassette and a stop codon after exon 2. It can be converted to a floxed 'Tm1c' allele by recombining the Frt sites with a *Flp*-driver and to a 'Tm1d' KO allele by recombining the LoxP sites with a *Cre*-driver as shown. **(B)** Lateral and dorsal views of an X-gal-stained E12.5 *DynIrb$^{tm1a/+}$* embryo to profile expression of β-Gal from the DynIrb1 locus. Note the strong expression in the nervous system, including the dorsal root ganglia (DRG, pointed at by an asterisk).

Figure S2, related to Figure 2. Characterization of *RNX3-DynIrb1* Conditional Knockout Mice. (A) Home cage locomotion analyses show no difference between the genotypes, mean ± SEM, n>6 mice per genotype. **(B)** Colorcation heat map representation of wild type, *RNX3-DynIrb1$^{+/-}$* and *RNX3-DynIrb1$^{-/-}$* mice over 10 min in the open field. **(C)** *RNX3-DynIrb1$^{-/-}$* mice spend more time in corners and less time near walls in the open-field compared to the other genotypes. Mean ± SEM, ** $p<0.01$ *** $p<0.001$, n>6 mice per genotype, one-way ANOVA followed by Tukey's HSD post hoc correction for multiple comparisons. **(D)** *RNX3-DynIrb1$^{-/-}$* mice also showed reduced speed and distance travelled in the open-field compared to the other genotypes. Mean ± SEM, *** $p<0.001$, n>6 mice per genotype, two-way ANOVA. **(E)** α-bungarotoxin staining (red) of the neuromuscular junctions (NMJ) in wild type and *RNX3-DynIrb1$^{-/-}$* mice. Scale bar, 50 μm. **(F)** Quantification of NMJ densities in wild type and *RNX3-DynIrb1$^{-/-}$* mice. No significant difference was found between the genotypes, mean ± SEM, n=5 mice per genotype, Unpaired t-test. **(G)** Sensory innervation of the skin in wild type and *RNX3-DynIrb1$^{-/-}$* mice. Tuj1 immunostaining (green) is used to highlight all the nerve endings, while nociceptive neurons are labeled with a CGRP antibody (red). Scale bar, 50 μm . **(H)** Quantification of Tuj1 and CGRP positive innervation of the skin in wild type and *RNX3-DynIrb1$^{-/-}$* mice. No significant difference was found between the genotypes, mean ± SEM, n=4 mice per genotype, unpaired t-test.

Figure S3, related to figure 2. Characterization of *Dynlrb1* knockdown in the DRG *in vivo*. (A) DRG ganglia sections from mice transduced with either AAV9-shControl or AAV9-shDynlrb1. AAV9 infected cells express the Venus reporter (green). Sensory neurons are labeled with β3 tubulin (red). Scale bar, 50 μm. **(B)** Percentage of GFP-positive neurons normalized to β3 tubulin-positive neurons in DRG sections of the experiment described in A. **(C)** Quantitative PCRs on RNA extracted from DRG neuron cultures transduced with AAV9-shControl or AAV9-shDynlrb1 for 7 days. AAV9-shDynlrb1 transduced neurons showed 40% reduction of *Dynlrb1* mRNA compared to shControl. Mean ± SEM, *** $p<0.001$, n=6, unpaired t-test. **(D)** Tail suspension reveals a clenched hind limb phenotype in mice transduced with AAV9-shDynlrb1 compared to AAV9-shControl-treated animals. **(E)** Rotarod tests (acceleration set at 20 rpm in 240 sec) on mice transduced as shown, 32 days after viral injection. AAV9-shDynlrb1-transduced mice show significantly reduced performance in the test. Mean ± SEM, * $p<0.05$, ** $p<0.01$, *** $p<0.001$, n>11 mice per group, two-way ANOVA. **(F)** Representative traces from catwalk gait analyses on mice transduced as described in E. The position of the paws are shown in different colors as indicated, RF – right front, RH - right hind, LF - left front, LH - left hind. **(G)** Catwalk analysis of experiment described in C. The step sequence regularity index as well as the duration of the paw stands are impaired for both front paws in mice transduced with AAV9-shDynlrb1. Mean ± SEM, * $p<0.05$, *** $p<0.001$, n>11 mice per group, unpaired t-test.

Figure S4, related to Figure 4. Transduction Efficiency with Adenovirus-Cre. (A) DRG neurons from Aig9 mice 72 hr after transduction with Cre-GFP Adenovirus. Transduced cells are positive for the antibody α-GFP (green) and td-Tomato (red). Neurons are identified with the cytoskeletal marker NFH (magenta). Scale bar, 100 μm. **(B)** Quantification of relative td-Tomato expression in the experiment described in A. Mean ± SEM, ** $p<0.01$, n = 3, one-way ANOVA followed by Tukey's HSD post hoc correction for multiple comparisons.

Supplementary movie 1, related to Figure 1. Video of a 2 months old *RNX3-Dynlrb1*[-/-] mouse showing the hindlimb clenching as well as the abnormal walk and posture (abdomen close to the ground).

Supplementary Table 1, related to Figure 3. List of differentially regulated genes highlighted by RNA sequencing between wild type and *Dynlrb1*[+/-] mice at 48h. The genes are ranked by FDR. An additional sheet is present, with the list of genes selected by a value of FDR<0.1. Raw and processed data were deposited within the Gene Expression Omnibus (GEO) repository (www.ncbi.nlm.nih.gov/geo), accession number (GSE131455).

Supplementary Table 2, related to Figure 3. List of pathways identified by the Ingenuity Pathway Analysis (QIAGEN Bioinformatics, sheet1). A list of all the genes of the WNT pathway found to be differentially regulated and the value of the change fold is also shown (sheet2).

STAR METHODS

Key Resource Table

Reagent or Resource	Source	Identifier
Antibodies		
NFH (Chicken)	Millipore	Cat Number: AB144P
Tuj1 (Rabbit)	Covance	RRID: AB_2313773
CGRP (Goat)	Abcam	RRID: AB_725807
β-catenin (Mouse)	A. Ben-Ze'ev lab (Johnson et al., 1993; Simcha et al., 1998)	Clone 5H10
GFP (Rabbit)	MBL International	RRID:AB_10597267
Alexa Fluor 488 secondary antibody (Donkey anti-rabbit)	Thermo Fisher Scientific	RRID: AB_2556546
Alexa Fluor 546 secondary antibody (Donkey anti-rabbit)	Thermo Fisher Scientific	RRID: AB_2534016
Alexa Fluor 546 secondary antibody (Donkey anti-goat)	Thermo Fisher Scientific	RRID: AB_2534103
DAPI (4.6-diamino-2-phenolindol dihydrochloride)	Thermo Fisher Scientific	RRID: AB_2629482
LysoTracker™ Red DND-99	Thermo Fisher Scientific	Cat number: L-7528
Alexa555-α-bungarotoxin	Weizmann Institute, Immunohistochemistry facility	N/A
TeNT HC – Alexa555	G. Schiavo Lab (Lalli and Schiavo, 2002)	N/A

Experimental Models: Organism/Strains

Mouse/C57BL/6OlaHsd	Harlan Israel	N/A
Mouse/C57BL/6OlaHsd/*Dynlrb1^{tm1a}*	International Knockout Mouse Consortium	allele: Dynlrb1$^{tm1a(EUCOMM)Wtsi}$
Mouse/FLP1eR - 129 / 129S4/SvJaeSor-Gt(ROSA)26Sor$^{tm1(FLP1)Dym}$/J	(Farley et al., 2000)	N/A
Mouse/Ai9	(Madisen et al., 2010)	N/A
Mouse/R26LacZ/Runx3^{Cre}	(Levanon et al., 2002)	N/A

siRNA screen Dharmacon smartpools

Catalog Number	Gene Symbol	Locus ID	Accession
D-001210-02	Non-targeting siRNA	N/A	N/A
M-043143-01	Dync1i1	13426	NM_010063
M-043171-01	Dync1i2	13427	NM_010064
M-051314-01	Dync1li1	235661	NM_146229
M-061580-01	Dync1li2	234663	NM_001013380
M-062961-01	Dync2li1	213575	NM_172256
M-055779-01	Dynlt1	21648	NM_009342
M-059901-01	Dynlt3	67117	NM_025975
M-044820-01	Dynlrb1	67068	NM_025947
M-050822-01	Dynlrb2	75465	NM_029297
M-063157-01	Dynll1	56455	NM_019682
M-059592-01	Dynll2	68097	NM_026556
M-044821-00	DCTN1	13191	NM_007835
M-056851-00	DCTN2	69654	NM_027151
M-063002-00	DCTN3	53598	NM_016890
M-063074-00	DCTN4	67665	NM_026302
M-063270-00	DCTN5	59288	NM_021608
M-047876-00	DCTN6	22428	NM_011722
M-049777-00	ACTR10	56444	NM_019785

Contact for Reagent and Resources Sharing

Further information and requests for resources and reagents should be directed to Marco Terenzio (marco.terenzio@weizmann.ac.il).

Experimental Model and Subject Details

Animal Subjects

All animal procedures were approved by the IACUC of the Weizmann Institute of Science. The knockout mouse model for *Dynlrb1* was generated by the European Conditional Mouse Mutagenesis Program (EUCOMM) as part of the International

Mouse Knockout Consortium (IKMC, allele: Dynlrb1$^{tm1a(EUCOMM)Wtsi}$) using a KO first targeting vector (MPGS8001_A_B08, IKMC Project 25156) (Skarnes et al., 2011). All mouse strains used were bred and maintained at the Veterinary Resource Department of the Weizmann Institute of Science. For live imaging of DRG outgrowth (**Figure 1A-D**) C57BL/6J YFP16 mouse were used (Feng et al., 2000). All control mice were age-, sex-matched littermates. For testing Adenovirus5 Cre-GFP infectivity Ai9 mice were used (Madisen et al., 2010). Mice utilized for behavioral experiments were kept at 24.0 ± 0.5°C in a humidity-controlled room under a 12-h light–dark cycle with free access to food and water. Experiments were carried out on 2-6 months old male mice, unless specified otherwise.

Generation of the *Dynlrb^{tm1c}* and *RNX3- Dynlrb$^{tm1d/tm1d}$* lines

The *Dynlrb1* mice acquired from the International Mouse Knockout Consortium (IKMC, allele: Dynlrb1$^{tm1a(EUCOMM)Wtsi}$).The gene targeted *Dynlrb1$^{tm1a(EUCOMM)Wtsi}$* allele, hereafter referred to as *Dynlrb^{tm1a}*, results in a complete knockout of *Dynlrb1* with concomitant expression of β-galactosidase from the same locus; this allele also allows the subsequent generation of a *floxed* conditional allele (*Dynlrb^{tm1c}*) upon application of the *Flp* recombinase (Skarnes et al., 2011). *Dynlrb$^{tm1a/+}$* heterozygous mice were crossed with a mouse harboring the *Flp* recombinase in all tissues (FLP1eR - 129/129S4/SvJaeSor-Gt(ROSA)26Sor$^{tm1(FLP1)Dym}$/J) (Farley et al., 2000), which was provided by the veterinary services of the Weizmann Institute of Science to recombine the FRT sites and so remove the LacZ sequence and giving a conditional allele, *Dynlrb^{tm1c}* in which *LoxP* sites flank a likely critical exon, exon 3, of *Dynlrb1* ((Skarnes et al., 2011), Supplementary Figure 1A). We further crossed the *Dynlrb^{tm1c}* mice with the proprioceptive Cre-driver R26LacZ/Runx3^{Cre} (Levanon et al., 2002) to generate the *RNX3- Dynlrb$^{tm1d/tm1d}$* line, which we refer to in the text as *RNX3-Dynlrb1* (Supplementary Figure 1A) (Skarnes et al., 2011), in which *Dynlrb1* is depleted in proprioceptive neurons.

Methods

DRG Culture

Adult mice DRGs were dissociated with 100 U of papain followed by 1 mg/ml collagenase-II and 1.2 mg/ml dispase. The ganglia were then triturated in HBSS, 10 mM Glucose, and 5 mM HEPES (pH 7.35). Neurons were recovered through Percoll, plated on laminin, and grown in F12 medium (ThermoFisher Scientific) as previously described (Ben-Yaakov et al., 2012; Hanz et al., 2003; Perlson et al., 2005).

siRNA Screen

Adult C57BL/6 YFP16 mouse DRG (Feng et al., 2000) were dissociated as previously described. Mouse siGenome smart pools (Dharmacon) were used for transfection of neuronal cultures two hours after plating, using DharmaFect 4 as the transfection reagent. 24 hours after siRNA transfection the neurons were plated in new poly-L-lysine and Laminin treated glass bottom 96 well plates at the same density (80,000-100,000 cells per well) and imaged 72 h after transfection. Images were captured at X10 magnification on ImageXpress Micro (Molecular Devices) and analyzed using the Metamorph software (Molecular Devices). The total outgrowth parameter was considered, defined as the sum of lengths of all processes and branches per cell. Statistical analyses were carried out with Student's t-test and One Way ANOVA using the statistical software GraphPad Prism.

Growth rate analysis

Adult C57BL/6J YFP16 mouse DRG (Feng et al., 2000) were dissociated as previously described and neurons were plated on MatTek glass bottom dishes (MatTek corporation). YFP6 - Wild type and YFP16-*Dynlrb1*$^{+/-}$ neurons were imaged at 1h intervals in a Fluoview (FV10i, Olympus) automated confocal laser-scanning microscope with built-in incubator chamber. Neuronal morphology was subsequently quantified using the Metamorph sofware (Molecular Devices). The total outgrowth parameter was considered, defined as the sum of lengths of all processes and branches per cell. Statistical analyses were carried out with a Two-way ANOVA test using the statistical software GraphPad Prism.

Adenovirus infection of DRG neurons from Ai9 mice

The efficiency transduction of adenovirus 5 Cre-GFP (Ad5 Cre-GFP) and the efficiency of the viral Cre were tested by infecting DRG neurons with Ad5 Cre-GFP at a multiplicity of infection (MOI) of 100. DRG neurons from Ai9 mice (Madisen et al., 2010) were dissected as previously described. The Ai9 Cre reporter mice harbour a LoxP-flanked STOP cassette preventing transcription of a CAG promoter-driven red fluorescent protein variant (td-Tomato). The virus was added directly into the F12 culture medium immediately after plating. Infected cultures were then incubated at 37°C for 48 and 72 h. After each time-point, neurons were washed in PBS, fixed in 4% PFA in PBS for 30 min and immunostained with an α-GFP antibody (1:1000) as previously described. Coverslips were then mounted with Flouromount-G^{TM} and imaged using the ImageXpress (Molecular Devices) automated microscope at 10X magnification or using a confocal laser-scanning microscope (Olympus FV1000, 60x oil-immersion objective Olympus UPLSAPO - NA 1.35). Image quantification was performed using the proprietary software Metamorph (Molecular Devices). Statistical analysis was carried out with a one-way ANOVA followed by Tuckey post-hoc multiple comparison test using the statistical software GraphPad Prism.

Axonal transport experiments

Wild type and *Dynlrb1 Flox$^{-/-}$* DRG ganglia were dissociated as previously described and neurons were plated on MatTek glass bottom dishes (MatTek corporation) in F12 medium (ThermoFisher Scientific) supplemented with 10 μM cytosine β-D-arabinofuranoside hydrochloride (AraC, Sigma-Aldrich, C6645) to inhibit glial proliferation. Adenovirus5 Cre-GFP was added to the neurons 4 h after plating. 96 h after infection with the virus, cells were washed in warm HBSS medium (ThermoFisher Scientific). LysoTracker™ Red DND-99 (ThermoFisher Scientific) (200 nM concentration) or AlexaFluor555-conjugated H$_C$T (40 nM) were added to the cells. Neurons were transferred to a a Fluoview (FV10i, Olympus) automated confocal laser-scanning microscope with built-in incubator chamber and imaged with a 60X water immersion objective. 120 frames at intervals of 5s between each frame were taken for every transport movie. Movies were tracked manually using the Manual Tracking plugin

of the ImageJ software; while the percentage of stationary vs moving carriers was performed on kymographs generated using the Multi Kymograph plugin of ImageJ. Statistical analysis was performed with two-way ANOVA using the statistical software GraphPad Prism.

β-Catenin nuclear accumulation in *Dynlrb1*[+/-] DRG neurons

Wild type and *Dynlrb1*[+/-] DRG ganglia were dissociated as previously described and cells were plated on coverslips in F12 medium (ThermoFisher Scientific). 48 h after plating cells w were washed in PBS, fixed in 4% PFA in PBS for 30 min and immunostained for β−catenin (1:100 dilution), NFH (1:2,000 dilution) and DAPI as described previously. Coverslips were then mounted with Flouromount-G[TM] and imaged using a confocal laser-scanning microscope (Olympus FV1000, 60x oil-immersion objective Olympus UPLSAPO - NA 1.35). Image quantification was performed using ImageJ software. Nuclei were identified using a mask drawn on the DAPI straining and cytoplasm was masked using the NFH staining. Nuclear and cytoplasmic intensity of β−catenin was recorded. Statistical analysis was carried out with a one-way ANOVA followed by Tuckey post-hoc multiple comparison test using the statistical software GraphPad Prism.

Design and production of AAV shDynlrb1

Design of AAV shRNA constructs was based on AAV-shRNA-ctrl (addgene, plasmid #85741, (Yu et al., 2015)). Sh-Ctrl sequence was replaced by a sequence targeting *Dynlrb1* using the BamHI and XbaI restriction sites. The primer ordered for the cloning were:

Forward:

GATCCGATCCAGAATCCAACTGAATAGAAGCTTGTATTCAGTTGGATTCTGGATCTT TTTTT,

Reverse:

CTAGAAAAAAGATCCAGAATCCAACTGAATACAAGCTTCTATTCAGTTGGATTCTG GATCG

Purified adeno associated virus (AAV) was produced in HEK 293t cells (ATCC®), with the AAVpro® Purification Kit (All Serotypes) from TaKaRa (#6666). For each construct 10 plates (15 cm) were transfected with 20 µg of DNA (AAV-plasmid containing our construct of interest + 2 helper plasmids for AAV9) using jetPEI® (Polyplus) in DMEM medium without serum or antibiotics. Medium (DMEM, 20 % FBS, 1 mM sodium pyruvate, penicillin-streptomycin) was added on the following day to a final concentration of 10% FBS and extraction was done 3 days post transfection. Purification was performed according to the manufactures' instructions (TaKaRa, #6666).

Immunohistochemistry of tissue samples

Tissue samples from muscle, skin, DRGs and sciatic nerve were harvested and fixed in 4% PFA in PBS (30 min for skin at RT, overnight at 4°C for muscles, DRG ganglia and sciatic nerve). The tissues were then washed in PBS and equilibrated in 30% w/v sucrose in PBS, embedded in Optimal Cutting Temperature Compound (O.C.T., Tissue-Tek®) and finally cryo-sectioned at a thickness ranging from 10-50 µm. Sections were then re-hydrated in PBS, blocked and permeabilized with 5% goat serum, 1% BSA, 0.1% Triton X-100 in PBS, or donkey serum, 0.1% Triton X-100 in PBS for 1 h and incubated with primary antibody overnight at 4°C. Sections were then washed 3 times in PBS and incubated for 2 h with secondary antibodies from Jackson Immunoresearch (Alexa Fluor 647 or Alexa Fluor 488 conjugated, 1:500 dilution in PBS). Slides were then washed in PBS, mounted with Flouromount-G^{TM}, and imaged using a confocal laser-scanning microscope (Olympus FV1000, 60x oil-immersion objective Olympus UPLSAPO - NA 1.35).

NMJ plate counting

Gastrocnemius muscles from wild type and *RNX3-Dynlrb1$^{-/-}$* mice were sectioned at a thickness of 50 µm. Staining of the neuromuscular junction was achieved as previously described by incubation with Alexa 555-α-bungarotoxin (1:1000 dilution). Sections were then imaged using a confocal laser-scanning microscope (Olympus FV1000, 60x oil-immersion objective Olympus UPLSAPO - NA 1.35) and the number of

positive NMJ-plate per area were manually counted and normalized by the area. Statistical analysis was performed with t-test using the statistical software GraphPad Prism.

Sensory skin innervation

Hind paw skin of wild type and *RNX3-DynIrb1$^{-/-}$* mice was sectioned at a thickness of 20 μm and immunostained as previously described with an α-CGRP and α-TUJ1 antibodies. The nuclear counterstaining DAPI was added to highlight all cells in the tissue. Sections were imaged using a confocal laser-scanning microscope (Olympus FV1000, 60x oil-immersion objective Olympus UPLSAPO - NA 1.35). The number of nerve endings was manually counted and normalized by the area of the skin. Statistical analysis was performed with t-test using the statistical software GraphPad Prism.

DRG neuron count

L4 DRGs from wild type and *RNX3-DynIrb1$^{-/-}$* mice were serially sectioned at a thickness of 20 μm in set of 3. One set for each DRG was then processed for immunostaining as previously described for α-NFH, α-CGRP and α-TUJ1 antibody. NFH and CGRP positive neurons were manually counted in blind. Statistical analysis was performed with t-test using the statistical software GraphPad Prism.

Gene expression analysis by Q-PCR

Wild type DRG ganglia were dissociated as previously described and neurons were plated in F12 medium (ThermoFisher Scientific) supplemented with 10 μM cytosine β-D-arabinofuranoside hydrochloride (AraC, Sigma-Aldrich, C6645) to inhibit glial proliferation. AAV9-shControl and AAV9-shDynIrb1 viruses were added to the neurons 4 h after plating. 7 days after infection, cells were washed in PBS and total RNA was extracted using the Oligotex mRNA Mini Kit (Qiagen). RNA purity and concentration was determined, and 300 ng of total RNA was then used to synthesize cDNA using SuperScript III (Invitrogen). Q-PCR was performed on a ViiA7 System (Applied Biosystems) using PerfeCTa SYBR Green (Quanta Biosciences, Gaithersburg,

USA). Forward/Reverse primers were designed to span exon-exon junction, and the RNA was treated with DNase H to avoid false-positives. Amplicon specificity was verified by melting curve analysis. All Q-PCR reactions were conducted in technical triplicates and the results were averaged for each sample, normalized to 18S levels and analyzed using the comparative ΔΔCt method (Livak and Schmittgen, 2001). The primers (*Mus musculus*) that were used are the following:

18S forward: AAACGGCTACCACATCCAAG

18S reverse: CCTCCAATGGATCCTCGTTA

Dynlrb1 forward: CAACCTCATGCACAACTTCATC (exon 3)

Dynlrb1 reverse: TCTGGATCACAATCAGGAAATAGTC (exon 4)

RNA sequencing

Wild type and *Dynlrb1*$^{+/-}$ DRG ganglia were dissociated as previously described and cells were plated on coverslips in F12 medium (ThermoFisher Scientific). 12, 24 and 48 h after plating cells were washed in PBS and total RNA was extracted using the Oligotex mRNA Mini Kit (Qiagen). RNA purity, integrity (RIN > 7) and concentration was determined. 3 biological repeats for each time point for each genotype was send to the laboratory of Prof. Giovanni Coppola at UCLA (USA).

RNA-sequencing libraries were prepared using the TrueSeq Stranded RNA kit (100ng). Libraries were indexed and sequenced by HiSeq4000 with 75 bp paired-end reads and at least 50M reads were obtained for each sample. Quality control (QC) was performed on base qualities and nucleotide composition of sequences, mismatch rate, mapping rate to the whole genome, repeats, chromosomes, key transcriptomic regions (exons, introns, UTRs, genes), insert sizes, AT/GC dropout, transcript coverage and GC bias to identify problems in library preparation or sequencing. Reads were aligned to the latest mouse mm10 reference genome (GRCm38.75) using the STAR spliced read aligner (ver 2.4.0). Average input read counts were 66.6M and average percentage of uniquely aligned reads were 86.3%. Total counts of read-fragments aligned to known gene regions within the mouse (mm10) ensembl (GRCm38.80) transcript reference

annotation are used as the basis for quantification of gene expression. Fragment counts were derived using HTSeq program (ver 0.6.0). Genes with minimum of 5 counts for at least half of samples were selected. Filtered count data were normalized and examined for sequencing bias by PCA, and PC1 and PC2 of sequencing QC (described above) were regressed out. Differentially expressed transcripts were determined by Bioconductor package EdgeR (ver 3.14.0) at FDR <0.1. Scripts used in the RNA sequencing analyses are available at https://github.com/icnn/RNAseq-PIPELINE.git. Raw and processed data were deposited within the Gene Expression Omnibus (GEO) repository (www.ncbi.nlm.nih.gov/geo), accession number (GSE131455).

The gene list from the 48h time point was filtered according to the FDR (FDR<0.1, **Table S1**) and uploaded to the Ingenuity Pathway Analysis software suite (QIAGEN Bioinformatics). The 10 best high ranking pathways according to their score (-log(p-values)) were further evaluated and the WNT – β-catenin pathway selected for validation (**Table S2**).

Intrathecal injection of AAV9 in mice

Intrathecal (IT) injections of adeno associated virus serotype 9 (AAV9) of 5 µl total volume were performed at the lumbar segment L4 using a sterile 10 µl Hamilton microsyringe fitted with a 30 gauge needle. For all constructs, we obtained titers in the range of 10^12-10^13 viral genomes/ml, which were used undiluted for intrathecal injections.

Behavioral profiling

All assays were performed during the "dark" active phase of the diurnal cycle under dim illumination (~10 lx); the ventilation system in the test rooms provided a ~65 dB white noise background. Every daily session of testing started with a 2 h habituation period to the test rooms. A recovery period of at least 1 day was provided between the different behavioral assays.

Home-cage locomotion test

Wild type and *RNX3-Dynlrb1^{-/-}* mice were monitored over 72 h using the InfraMot system (TSE Systems, Germany) in order to investigate possible alterations of basal motor activity and/or circadian rhythms. The spatial displacement of body-heat images (infra-red radiation) over time was tracked (Cooper et al., 2012; Neufeld-Cohen et al., 2010). The first 24 h were considered as a habituation phase, while the mouse activity of following 48 h (2 consecutive dark/light cycles) was calculated. Statistical analysis was carried out with a one-way ANOVA followed by Tuckey post-hoc multiple comparison test using the statistical software GraphPad Prism.

Open field

The motility of wild type and *RNX3-Dynlrb1^{-/-}* mice was assessed in the open-field arena under 120lx (Cooper et al., 2012; Neufeld-Cohen et al., 2010). The total distance moved (cm), the time spent (wall, corner; s) and mean speeds (cm/s) were recorded using VideoMot2 (TSE System, Germany) or Ethovision XT11 (Noldus Information Technology, the Netherlands). The data were analyzed with the COLORcation software (Dagan et al., 2016). Statistical analysis was carried out with a one-way ANOVA followed by Tuckey post-hoc multiple comparison test or two-way ANOVA (for speed and distance) using the statistical software GraphPad Prism.

Accelerated rotarod

Wild type and *RNX3-Dynlrb1^{-/-}* mice and wild type mice intrathecally injected with AAV9-shControl or AAV9-shDynlrb1 were tested with the ROTOR-ROD system (83x91x61 - SD. Instruments, San Diego) for their balance/coordination ability (Crawley, 2008; Marvaldi et al., 2015). Mice were subjected to 3 trials, with 5 min inter-trial intervals. Rotarod acceleration was set at 20 rpm in 240 s. Latency to fall (s) was recorded and the average of the 3 trials was considered. Statistical analysis was carried out with two-way ANOVA using the statistical software GraphPad Prism.

Catwalk Gait Analysis

Wild type and *RNX3-Dynlrb1^{-/-}* mice and wild type mice intrathecally injected with AAV9-shControl or AAV9-shDynlrb1 were assessed for their motor/balance coordination using the CatWalk as previously described (Perry et al., 2012). Motivation was achieved by a placing the mice home cage at the end of the runaway. The test was repeated 3-5 times for each mouse. Data were collected and analyzed using the Catwalk Ethovision

XT11 software (Noldus Information Technology, The Netherlands). The parameters reported for each animal are the paw stance, defined as the duration in sec of the contact of the paw with the glass surface, and the step sequence regularity index, defined as the number of normal step sequence patterns relative to the total number of paw placements. Statistical analysis was carried out with a one-way ANOVA followed by Tuckey post-hoc multiple comparison test (Wild type and *RNX3-Dynlrb1$^{-/-}$* mice) or t-test (AAV injected mice) using the statistical software GraphPad Prism.

Statistical analysis

Analysis of multiple groups was made using the ANOVA method. The choice between one- or two-way ANOVA was based on the requirements for identification of specific factors' contribution to statistical differences between groups and were followed by the Tukey and the Sidak post hoc analysis tests respectively. For 2-groups analyses, unpaired Student's t test was used. All analyses were performed using GraphPad Prism version 7.00 for MacOS (GraphPad Software, La Jolla, California, USA, https://www.graphpad.com/). The results are expressed as the mean ± standard error of the mean (SEM). All statistical parameters for specific analyses are reported in the figure legends of the paper. Statistically significant P-values are shown as ∗p < 0.05, ∗∗p < 0.01, ∗∗∗p < 0.001 and ∗∗∗∗p < 0.0001.

Figure 1

Figure 2

Figure 3

A
- ▨ downregulated genes
- ☐ upregulated genes

time (hr)		
48	737	944
24	1	1
12	2	3

differentially expressed genes

B

wild type $Dynlrb1^{tm1a/+}$ Log10(FPKM)

D

Neuroprotective Role of THOP1 in Alzheimer's Disease
eNOS Signaling
Wnt/β-catenin Signaling
Role of Macrophages, Fibroblasts and Endothelial Cells in Rheumatoid Arthritis
cAMP-mediated signaling
Role of Osteoblasts, Osteoclasts and Chondrocytes in Rheumatoid Arthritis
Glioblastoma Multiforme Signaling
Complement System
Proline Biosynthesis II (from Arginine)
Inhibition of Matrix Metalloproteases

IPA categories

-log(p-values)

C

DAPI   NFH   β-catenin   DIC

wild type

$Dynlrb1^{tm1a/+}$

D

- ● wild type
- ● $Dynlrb1^{tm1a/+}$

Nuclear/Cyto β-Catenin

*

Figure 4

Supplementary Figure 1 (refers to Figure 1)

A

Dynlrb1^{tm1a} allele

FLPe mediated cassette deletion

Dynlrb1^{tm1c} allele
(transcript expressed as WT)

Cre mediated
cassette deletion

Dynlrb1^{tm1d} KO allele
(frameshift transcript expressed)

B

Supplementary Figure 2 (refers to Figure 2)

Supplementary Figure 3 (refers to Figure 2)

Supplementary Figure 4 (refers to Figure 4)