Light suppresses the acoustically prompted predator-avoidance behaviour in moths

Theresa Hügel[1]*, Holger R. Goerlitz[1]*

[1] Acoustic and Functional Ecology Group, Max Planck Institute for Ornithology, Seewiesen,

Germany

*corresponding authors:

Acoustic and Functional Ecology Group

Max Planck Institute for Ornithology

Eberhard-Gwinner-Str. 11

82319 Seewiesen, Germany

Email: thuegel@orn.mpg.de, hgoerlitz@orn.mpg.de

ORCID:

Theresa Hügel: 0000-0003-0500-1988

Holger Goerlitz: 0000-0002-9677-8073

ABSTRACT

Urbanization leads to increased light exposure of wild animals. Especially nocturnal animals

are affected by artificial light, which shifts balances of animals' interactions. Predator-prey

interactions, for example, can be shifted either in favour of the prey or the predator under

the influence of light. In the interaction between echolocating bats and eared moths, bats

often benefit from moths eliciting less last-ditch behaviour in response to close-by attacking

bats when exposed to light. In contrast, it is unknown to which extent light also affects

moths' negative phonotoaxis, which moths elicit in response to distant bats. To quantify the

overall effect of light on both last-ditch behaviour and negative phonotaxis, we counted over

several nights the moths caught by two light traps, which were alternatingly equipping with

speakers broadcasting ultrasonic simulations of hunting bats. Playbacks did not affect moth

counts, indicating that light not only reduces the last-ditch behaviour of moths, but also their

negative phonotaxis. The increasing light pollution therefore interferes with different

aspects of anti-predator behaviour in moths, and possibly other eared insects. Our study

emphasizes the negative effect of lights on insects and the need to further study the

underlying mechanism.

KEY WORDS

Artificial light at night (ALAN), predator-prey, bats, Noctuidae, light traps, playback

experiment

BACKGROUND

One-fifth of the world's human population cannot see the Milky Way anymore (1), caused by

the substantial increase of artificial light at night (ALAN) over the last decades, causing

increasing light pollution (2,3). Likewise, also plants and animals are affected by increasing

light pollution (4–6). Light pollution can impact a single species' orientation, reproduction or

communication (4), but also affects whole communities, for example by shifting the balance

of trophic interactions such as predator-prey interactions (7–12).

Echolocating bats preying upon eared moths are a global predator-prey system with

important ecological impacts (13–16). Bats and moths are primarily nocturnal and strongly

rely on their sense of hearing to communicate, orientate, hunt or defend themselves. When

bats search for their food, they emit echolocation calls to probe the environment (17,18).

Eared moths are able to hear those calls and to respond with a two-staged anti-predator

flight behaviour (19,20). During stage one, moths show negative phonotaxis, i.e. they fly

away from the calling bat. Negative phonotaxis is elicited when the bat is still far away and

serves the purpose of avoiding detection by the bat. During stage two, moths show last-ditch

behaviour such as erratic manoeuvres or diving to the ground. Last-ditch behaviour occurs

when the bat is nearby and serves the purpose of avoiding capture by the bat. Whether

negative phonotaxis or last-ditch behaviour is elicited depends on the acoustic properties of

the bat's echolocation call received at the moth's ear, most importantly received sound

pressure level, but also call length and call repetition rate, which provide information about

the distance and behaviour of the bat (17). The received sound pressure level of distant bats

is fainter than of nearby bats, due to spherical spreading and atmospheric attenuation (21).

Despite this predominantly acoustic interaction, both bats and moths are strongly affected

by artificially introduced light sources. While some bats may profit from artificial light, for

example by exploiting resulting accumulations of prey animals (22,23), other species are also

negatively affected, for example when commuting or distribution to new habitats is

interrupted or hindered (24–26). Moths are mainly negatively affected by light. Light

strongly attracts moths, which interrupts and reduces their feeding (27,28) and commuting

behaviour (29), resulting in ecological consequences such as reduced pollination (30) or

population decline (27,31). Light also increases moth's predation risk in two ways. First,

moth accumulations around lights attract bats, thus increasing the moth's predation

pressure (22,23). Second, light impedes the moths' anti-predator flight behaviour. Several

studies that compared the sound-induced anti-predator flight of moths between lit and unlit

conditions reported a reduction of anti-predator flight in light. Wakefield et al. (32)

specifically studied last-ditch behaviour, showing that only 24% of moths performed power-

dives under LED illumination compared to 60% in darkness, i.e. LED-illumination inhibited

the last-ditch behaviour in 60% of the moths that reacted in darkness. Similarly, Svensson &

Rydell (33) reported a reduction of ~60% within a radius of 1 meter around the light source

compared to darkness (where all moths reacted), also mostly noticing last-ditch evasive

behaviour. Minnaar et al (34) reported the most extreme effect, showing that the model

without light-induced reduction in anti-predator flight behaviour explained best the

differences in bat diets between lit and unlit conditions. Another set of studies kept the light

exposure constant while manipulating the sound received by the moth, showing that moths

still exhibit some degree of anti-predator flight under illumination. Acharya & Fenton (35)

compared last-ditch behaviour of eared and deafened moths under illumination, showing

that 48% of eared moths exhibited last-ditch behaviour when attacked by bats, whereas

deafened moths did not exhibit any erratic flight. Agee & Webb (36) and Treat (37)

compared the number of caught moths at light traps with and without ultrasonic stimuli. In

both studies, ultrasound reduced the capture rates at the light trap with ultrasound playback

compared to the silent light trap to various degrees depending on stimulus design and moth

family. Across different stimulus types, capture rates at the sounding light trap were

reduced by 51-86 percentage points compared to the capture rate at the silent trap in a

Noctuid species (*Heliothis zea*, (36)), and by 8-49 percentage points for different eared moth

families (with a minimum sample sizes of 37 individuals in total, (37)). In summary, while

some studies confirm that light has a suppressive effect on the anti-predator flight of moths,

ranging from 60-100% suppression when comparing acoustically elicited anti-predator

behaviour in dark and lit conditions, other studies confirm that even under light conditions

there is a repelling effect of the playbacks, ranging from 8-86 percentage points when

comparing attraction to light under ultrasonic and silent conditions. Therefore, the extent of

light suppressing anti-predator flight in moths is unclear. Importantly, however, the majority

of those studies either focused on (32) or naturally observed (33) only last-ditch behaviour.

Even in the studies of Treat (37) and Agee & Webb (36), where the stimuli potentially elicited

both types of evasive flight behaviour, last-ditch behaviour was likely dominating the

outcome due to the study design. Both studies used directional light- and sound sources

which made it likely that moths attracted to the light were always exposed to sufficiently

high sound levels that could elicit last-ditch behaviour. Only Minnaar et al. (34) (indirectly)

investigated both stages of evasive flight by observing the diet of naturally foraging bats

under lit and dark conditions, suggesting that light under these more naturalistic conditions

may completely suppress the antipredator flight throughout the natural moth population.

Yet, a direct quantification of the overall effect of light on both stages of the anti-predator

flight in moths is missing. Here, we designed an experiment to investigate the effects of light

on both last-ditch behaviour and negative phonotaxis. Like Agee & Webb and Treat (36,37),

we compared moth captures at two light traps, while alternatingly simulating bat predation

pressure via ultrasonic playbacks at one light trap. We combined an omnidirectional light

field (attracting moths equally) with a directional sound field. Therefore, sound intensities

around the light trap varied and therefore triggered either last-ditch behaviour (on-axis) or

negative phonotaxis (off-axis) in proximity to the trap.

 We predicted three different outcomes, depending on the effect of light on the two stages

of moths' antipredator flight: (1) No moth captures at the ultrasonic light trap, indicating

that both stages of sound-elicited antipredator flight is functional and not suppressed by

light. (2) Reduced moth capture rates at the ultrasonic light trap, indicating that light partly

suppresses sound-elicited anti-predator flight. (3) Equal captures at both light traps,

indicating that light completely suppresses both stages of sound-elicited antipredator flight.

METHODS

Setup, study site, moth capture and measurement

We compared the number of caught moths between two light traps, one additionally

equipped with a loudspeaker, to examine the effect of light on the evasive flight behaviour

of eared moths. We set up two equal light traps (Sylvania, blacklight, F15W/350BL-T8, USA,

Fig 1A) at 30 m distance to each other, one at a path in a forest (trap A) and one in the forest

itself (trap B), close to the Max Planck Institute for Ornithology, Seewiesen, Germany. Both

traps hung freely at ~1.7 m above ground, radiating light at 360° in the horizontal plane (Fig

1A). We collected data over 15 rainless nights between the 19[th] of July and the 16[th] of

August 2018. Each test night, we alternatingly equipped one of the two traps with two

loudspeakers (Ultrasonic Dynamic Speaker ScanSpeak, Avisoft Bioacoustics, Germany)

broadcasting an ultrasonic stimulus (see below) to simulate echolocating bats. We fixed both

speakers back to back facing into opposite directions and mounted them above one of the

light traps at ~2 m above ground, associating each trap for 7-8 nights with the playback. We

switched the light of the traps and the acoustic playback on in the evening (between 20:30h

and 23:40h) and off the next morning (between 7:00h and 9:40h). After switching off, we

emptied the traps each morning and counted all moths of the ear-possessing families

Noctuidae, Geometridae, and Sphigindae. We measured the individuals' body length along

the middorsal line (from the head to the end of the abdomen), to correct for the fact that

larger moths are more sensitive (38,39). For those individuals whose body length we could

not measure (e.g. due to missing abdomen, N = 137, 15.5%) we either used the mean value

of the same species or, if this was not possible (N = 1), the mean value of the family.

Ultrasonic playback stimulus design and evaluation

We simulated predation pressure by echolocating bats by playing an ultrasonic stimulus

consisting of repeated pure tones. The stimulus was designed based on the acoustic

properties of bat echolocation calls (call frequency, duration, and interval) and the hearing

system of moths. Pulse frequency was 35 kHz, matching most moths' best hearing threshold

between 20-50 kHz (Noctuidae: (39,40); Erebidae: (41); Pyralidae: (42); Geometridae:

(43,44); Sphingidae/Drepanidae: (40)). Pulse duration was 10 ms including 2 ms linear rise

and fall time, i.e., corresponding to the longer calls of European open space bats (45,46), to

optimize detection by moths and information transfer about bat distance (47). The

interpulse interval was 100 ms, matching the call interval of bats' echolocation calls in their

search phase (60 - 200 ms, (45,48)). On-axis sound pressure level was 98 dB SPL re. 20 µPa

RMS at 1 m distance (see below for a detailed description of the sound field). This stimulus

was presented continuously in a loop throughout the night via the loudspeakers using

Avisoft-RECORDER software (Avisoft Bioacoustics, Germany), a sound card (Avisoft

UltraSoundGate 116, Avisoft Bioacoustics) and a laptop (DELL Latitude E6540, Dell

Technologies Inc., USA).

To validate the effect of the stimulus in darkness, without the potential suppressing

influence of light, we exposed free-flying moths in a dark flight room (5.3 m x 3.5 x 3 m^3) to

the stimulus. We caught moths at trap A over the course of four nights and tested them

within 30 hours after capture. We placed moths on the ground of the flight room and

recorded the flight paths of upwards flying moths with an IR-sensitive camera (Sony HDR-

CX560, Sony) under IR illumination (850 nm, Mini IR Illuminator TV6700, ABUS Security-

Center). Using the same audio equipment as described above, we presented the stimulus by

manually triggering when a moth flew in front of the speaker. The moth's distance to the

speaker at stimulus onset was about 1-2 m. We categorized the recorded flight paths as

"reaction", when the flight direction, level of erraticness, or both changed with stimulus

onset (for examples see Supplement Information); as "no reaction", when we did not

observe those changes; or as "ambiguous", if we could not assign the flight path to one of

the other categories. In total, we evaluated 32 flight paths with ten moths (31.2 %) showing

a reaction, twelve moths (37.5%) not showing a reaction, and eleven moths (34.3%) were

categorized as "ambiguous".

Overlap of sound and light field

The range and geometry of the presented sound and light fields differed. While the light was

emitted omnidirectional in the horizontal plane, the sound field was directional (Fig. 1). We

estimated the biologically effective range for the attraction by light and the biologically

relevant sound fields for triggering evasive flight behaviour based on literature values and

own measurements.

Light traps can attract released moths from up to 80 m distance, yet recapture rates

dramatically decrease beyond 15 m, and additionally depend on family (49,50). Family-

specific modelled 5%-recapture rates are 10 ± 6 m (mean ± SEM) for Noctuidae and 23 ±

12 m for Geometridae (49). Note, however, that these data were obtained with a different

light source (6W actinic vs. 15W blacklight in our case), and that the radius of attraction to

light is not necessarily equivalent to the radius over which light interferes with anti-predator

behaviour. Yet, we used these family-specific values as first estimates of the biological

relevant distances for the attraction of moths to our light trap (Fig. 1B, dashed lines).

To estimate the effect of the playback, we measured the playback level of the loudspeaker

from 0° on-axis to 90° off-axis in steps of 5° (for details about calculations see Supplement

Information). On-axis source level was 97 dB SPL re. 20 µPa RMS @ 1 m distance, i.e. within

the lower range of natural bat call levels in the field (100-120 dB peSPL @ 1 m (48), with

corresponding RMS-SPL values being ~3-7 dB lower than the peSPL values (51,52)). Playback

source level was reduced with increasing off-axis angle by up to ~30 dB at 45°, resulting in a

minimum playback level of 70 dB SPL RMS @ 1m. We then calculated the angle-dependent

distances to the loudspeaker where the playback would reach biologically relevant levels of

60 and 80 dB SPL RMS. We chose 60 and 80 dB SPL as approximate thresholds for eliciting

negative phonotaxis and last-ditch behaviour, respectively, based on several lines of

evidence. Triggering of negative phonotaxis and last-ditch behaviour in moths is supposed to

be correlated with the activity of moths' auditory receptor neurons A1 and A2, respectively

(47,53,54). The neuronal thresholds of these neurons vary with frequency and species

(39,47,53,55–57) with A1 being generally about 20 dB more sensitive than A2 (39,53,58). A1

best thresholds range from ~35-55 dB SPL at the position of the moth, and A2 best

thresholds range from ~52-72 dB SPL (20,39,47,53,55,56). Where known, behavioural

thresholds in moths are often ~10 dB higher than neuronal ones (reviewed in (52)). Indeed,

our own data indicate that sound pulses of ~80 dB SPL elicit last-ditch behaviour (59), which

is 8-28 dB above known A2 thresholds (20,39,47,53,55,56). As no behavioural data exists for

the threshold of negative phonotaxis, we chose 60 dB SPL, i.e., 20 dB below the threshold for

last-ditch behaviour, matching the difference between A1 and A2 thresholds. We calculated

the distance-isolines for 60 and 80 dB SPL, which varied with the angle around the

loudspeaker, ranging from 3.6-14.6 m and 0-5.6 m, respectively (Fig. 1B).

Figure 1: (A) Experimental setup. One of the light trap with two loudspeakers attached
above the trap, which broadcast the acoustic stimulus in opposite directions. **(B) Side view**
of the biological effective sound field around the light trap with attached loudspeakers.
Coloured areas indicate areas with minimum sound levels of 60 and 80 dB SPL re. 20 μPa,
which are biological relevant acoustic thresholds for eliciting negative phonotaxis and last-
ditch behaviour, respectively, in eared moths. Schematic indicates position and dimenstion
of the light trap. **(C) Top view of the biological effective sound field (coloured areas) and**
light field (dashed lines). Dashed lines indicate the distance of 5% recapture rate for noctuid
(10 m) and geometrid (23 m) moths, respectively (49), as an indicator of the maximum range
of light attraction.

Statistical analysis

To test for an effect of light on the moths' negative phonotaxis, we fitted linear mixed-effects models to the moth count data as a function of the fixed effects *playback*, *trap*, *moth family*, and *moth body length*, the random factor *date* (to account for repeated measures)*, and the interactions of *playback* and *moth family*, and *playback* and *moth body length*. Count data were log-transformed to approximate a normal distribution. Even though normal distribution was not achieved, we selected the log-transformed model over the alternative Poisson-model, as this showed some signs of overdispersion (dispersion factor = 1.4) making it less reliable than the log-transformed Gaussian-model with an incomplete normal distribution (60). We defined the moth family Noctuidae as intercept, as it had the largest sample size and thus was the most reliable reference. To test which factors significantly contributed to the model fit, we conducted backward model reduction. We compared models with likelihood ratio tests using a chi-squared statistic to find the minimal adequate model. All statistical analyses were conducted in R version 3.3.2 (R Foundation for Statistical Computing, Vienna, Austria) using the packages "lme4"(61), "lmerTest"(62), and "blmeco" (63).

RESULTS

We caught a total of 871 moths over 15 experimental nights (range: 7-115 moths per trap and night, Fig. 2). The minimal adequate model included *playback*, *family* and *size* as fixed and *date* as random factors. Capture success varied between moth families (likelihood ratio test for factor *family*, $\chi^2 = 62.814$, d.f. = 3, $p < 0.001$). The vast majority of 81.2% of the moths belonged to the family Noctuidae, the remaining 18.8 % were composed of

Geometridae (11.4%), Erebidae (6.9%), and Sphingidae (0.5%). Moths of these four families

possess ears, except for Sphingidae where ears might be a derived structure as they are only

found in two subtribes (64–66). We caught more small than large moths (likelihood ratio test

for factor *size*, χ^2 = 89.856, d.f. = 5, p < 0.001). Although capture rates seemed to differ

between traps (total capture of 506 (trap A) vs. 365 (trap B), Fig 2A), *trap* was not a

significant factor of the minimal adequate model (likelihood ratio test, χ^2 = 0.488, d.f. = 1, p =

0.484).

Our playback treatment had no effect on the number of caught moths, i.e. the number of

caught moths did not differ between nights with and without playback (Fig. 2A, likelihood

ratio test of factor *playback*, χ^2 = 1.084, d.f. = 1, p = 0.297). There was also no significant

effect of the interactions between *moth size* and *playback* and of *moth family* and *playback*

on the model fit (likelihood ration tests, χ^2 = 3.159, d.f. = 3, p = 0.367 and χ^2 = 1.328, d.f. = 3,

p = 0.722, respectively).

Figure 2: Number of caught moths per night and playback treatment. **(A) Number of caught**
moths at both light traps without and with playback. Boxplots show median, quartiles and
whiskers (up to 1.5 x inter-quartile range) of the daily counts (dots). **(B) Daily counts of**
caught moths per night and trap for all 15 nights. White letters at the base of the bars
indicate the trap.
273

DISCUSSION

Echolocating bats and eared insects are a textbook example of predator-prey interactions,

which is based on acoustic information (20). Increasing artificial light at night (ALAN) (2,3),

however, severely impacts both bats and moths (e.g. (23,24,28)), with potential cascading

effects on their predator-prey-interactions, population dynamics and ecosystems (11,34).

Light reduces the acoustically triggered last-ditch behaviour of eared moths, making them

more vulnerable to close-by attacking bats. In contrast, it is less clear whether light also

reduces negative phonotaxis in eared moths, which serves as their first line of defence

against distant echolocating bats. To better understand how artificial light at night affects

the overall capability of predator avoidance in eared moths, we compared moth captures at

two light traps. One trap was "silent", while the other trap broadcast bat-like ultrasonic

stimuli that trigger last-ditch behaviour as well as negative phonotaxis under dark

conditions. We caught the same number of moths at both traps. Thus, the presented

playback did not repel the moths, suggesting that the light suppressed both anti-predator

behaviours, negative phonotaxis and last-ditch behaviour.

In contrast to our results, Treat (37) and Agee & Webb (36) both showed a repelling effect of

their playbacks on moth capture rates at the ultrasonic trap. We deem differences in

stimulus design and habituation effects of the moths unlikely to explain these differences.

Treat (37) and Agee & Webb (36) broadcast a variety of different stimuli varying in pulse

rate, frequency, pulse duration, and SPLs (ranges: pulse rate:0.7-155 pulses/s; frequency:

12.5-200 kHz; pulse duration: 2-10ms; SPL: ~60-100 dB SPL @ 1 m distance), which all

elicited evasive flight to various degrees in eared moths. As our stimulus had acoustic

properties within this range and elicited evasive flight in unlit conditions, we doubt that

differences in the stimuli can explain the difference in capture rates. Habituation to the

ultrasound could have affected individuals that flew close to the light traps for extended

time periods. It is more likely, however, that most caught individuals randomly encountered

and approached the traps over larger distances and thus did not perceive our presented

stimulus continuously. Likewise, Agee & Webb (36)also presented their ultrasonic stimuli

over the course of a whole night and were able to detect a repelling effect of their

playbacks.

We suggest that major differences in the geometry and overlap of light- and sound-fields

could explain these contrasting results. In the setups of Treat (37) and Agee & Webb (36),

sound and light fields were almost overlapping. Before entering the light traps in these

setups, approaching moths had to pass high sound pressure levels, which were sufficiently

high to potentially elicit last-ditch behaviour. Both studies indeed caught fewer moths in the

ultrasonic trap compared to the silent trap, suggesting that the playback elicited still some

anti-predator flight (likely last-ditch behaviour) under these light and sound conditions.

Specifically, for stimuli similar to ours, the capture rate for noctuid moths was 35% for a

37.5 kHz pulsed tone stimulus vs. 65% at the silent trap (37), while Agee & Webb (36)

reported ~15% capture rate for 30 kHz pulsed tones vs. 85% at the silent trap for two species

of the families Noctuidae and Pyralidae. In contrast, our setup combined omnidirectional

light fields with directional sound fields, which exposed the moths to different direction-

dependent sound pressure levels (SPLs) while approaching the light. Moths approaching the

trap on-axis were exposed to first low and then increasingly higher SPLs that should first

elicit negative phonotaxis and later last-ditch behaviour. In contrast, moths approaching the

trap off-axis were exposed only to lower SPLs that can only elicit negative phonotaxis. As our

capture rates did not differ between the light traps, i.e., indicating no repelling effect of the

playback on moth flight, we suggest that the light strongly reduced the negative phonotaxis

of eared moths. Even if the moths still exhibited some degree of last-ditch behaviour close to

the trap (as shown by Treat and Agee & Webb, (36,37), these last-ditch manoeuvres might

bring the moth into a position off-axis to the loudspeaker, either to the sides or below the

loudspeaker, where the SPL is lower. Due to the lower SPL, no further last-ditch evasive

flight is elicited, while the light keeps attracting the moth into the trap. Our results are in line

with Minnaar et al. (34), who found that a model assuming 0% efficiency of escape

behaviour in moths best explained bats' diet under lit conditions. We therefore suggest that

light not only suppresses last-ditch behaviour, as previously shown but also negative

phonotaxis.

In summary, our results underline the strong effect of (artificial) light on eared moths,

providing an additional explanation for shifts in predator-prey interaction between bats and

moths (11,12,34). With negative phonotaxis also being suppressed by light, additionally to

last-ditch behaviour, moths are not just unable to escape predators, but also to avoid them.

Similarly, fish are attracted to lit areas, where they are "trapped" and predated upon by

seals (9) and other fish (67). Our results emphasize the need for understanding the

mechanisms underlying animals' attraction to light and other effects of light on animals'

behaviour, such as predator avoidance. This is particularly important given the increasing

light pollution and the general decline of insect populations. Only by understanding

underlying mechanisms, we can apply efficient conservation measures.

Acknowledgements

We thank the IMPRS for support and especially the participants of the course for "Scientific Writing" 2019 for comments on previous versions of the manuscript. We further thank the Max Planck Institute for Ornithology Seewiesen for providing excellent infrastructure and support.

Competing interests

We have no competing interests.

Funding

This study was supported by funds of the Deutsche Forschungsgemeinschaft (Emmy Noether research grants GO2029/2-1 and GO2029/2-2 to H.R.G.) and the International Max Planck Research School for Organismal Biology (student grant to T.H.)

Author contributions

TH and HRG conceived of and designed the study.

TH collected and analysed the data.

TH and HRG wrote the manuscript.

All authors gave final approval for publication and agree to be held accountable for the work performed therein.

REFERENCES

1. Falchi F, Cinzano P, Duriscoe D, Kyba CCM, Elvidge CD, Baugh K, et al. The new world atlas of
artificial night sky brightness. Sci Adv. 2016;2(6):e1600377.

2. Hölker F, Moss T, Griefahn B, Kloas W, Voigt C, Henckel D, et al. The dark side of light: a
transdisciplinary research agenda for light pollution policy. Ecol Soc. 2010;15(4).

3. Fouquet R. Seven centuries of energy services: the price and use of light in the United Kingdom
(1300-2000). Energy Journal. 2006;27:139–77.

4. Longcore T, Rich C. Ecological light pollution. Front Ecol Environ. 2004;2(4):191–8.

5. Davies TW, Smyth T. Why artificial light at night should be a focus for global change research in
the 21st century. Glob Chang Biol. 2018;24(3):872–82.

6. Knop E, Zoller L, Ryser R, Gerpe C, Hörler M, Fontaine C. Artificial light at night as a new threat
to pollination. Nature. 2017;548(7666):206–9.

7. Davies TW, Bennie J, Inger R, Ibarra NH, Gaston KJ. Artificial light pollution: are shifting spectral
signatures changing the balance of species interactions? Glob Chang Biol. 2013
May;19(5):1417–23.

8. Davies TW, Duffy JP, Bennie J, Gaston KJ. The nature, extent, and ecological implications of
marine light pollution. Front Ecol Environ. 2014;12(6):347–55.

9. Yurk H, Trites AW. Experimental attempts to reduce predation by harbor seals on out-migrating
juvenile salmonids. Trans Am Fish Soc. 2000;129(6):1360–6.

10. Miller Colleen R., Barton Brandon T., Zhu Likai, Radeloff Volker C., Oliver Kerry M., Harmon
Jason P., et al. Combined effects of night warming and light pollution on predator–prey
interactions. P Roy Soc P B Biol Sci. 2017;284(1864):20171195.

11. Russo D, Cosentino F, Festa F, De Benedetta F, Pejic B, Cerretti P, et al. Artificial illumination
near rivers may alter bat-insect trophic interactions. Environ Pollut. 2019;252:1671–7.

12. Bailey LA, Brigham RM, Bohn SJ, Boyles JG, Smit B. An experimental test of the allotonic
frequency hypothesis to isolate the effects of light pollution on bat prey selection. Oecologia.
2019;190(2):367–74.

13. Van Toor ML, O'Mara MT, Abedi-Lartey M, Wikelski M, Fahr J, Dechmann DKN. Linking colony
size with quantitative estimates of ecosystem services of African fruit bats. Curr Biol.
2019;29(7):R237–8.

14. Kunz TH, Braun de Torrez E, Bauer D, Lobova T, Flemming TH. Ecosystem services provided by
bats. Ann NY Acad Sci. 2011;1223(1):1–38.

15. Kasso M, Balakrishnan M. Ecological and economic importance of bats (order Chiroptera). ISRN
Ecol. 2013;Article ID 187415:9 pages.

16. Boyles JG, Cryan PM, McCracken GF, Kunz TH. Economic importance of bats in agriculture.
Science. 2011;332(6025):41–2.

17. Schnitzler H-U, Moss CF, Denzinger A. From spatial orientation to food acquisition in
echolocating bats. Trends Ecol Evol. 2003;18:386–94.

18. Fenton MB. Eavesdropping on the echolocation and social calls of bats. Mammal Rev.
2003;33:193–204.

19. Roeder KD. The behaviour of free flying moths in the presence of artificial ultrasonic pulses.
Anim Behav. 1962;10(3–4):300–4.

20. Ter Hofstede HM, Ratcliffe JM. Evolutionary escalation: the bat-moth arms race. J Exp Biol.
2016;219(Pt 11):1589–602.

21. Goerlitz HR. Weather conditions determine attenuation and speed of sound: Environmental
limitations for monitoring and analyzing bat echolocation. Ecol Evol. 2018;8(10):5090–100.

22. Rydell J. Exploitation of insects around streetlamps by bats in Sweden. Funct Ecol.
1992;6(6):744–50.

23. Cravens ZM, Brown VA, Divoll TJ, Boyles JG. Illuminating prey selection in an insectivorous bat
community exposed to artificial light at night. J Appl Ecol. 2018;55(2):705–13.

24. Stone EL, Jones G, Harris S. Street lighting disturbs commuting bats. Curr Biol.
2009;19(13):1123–7.

25. Mathews F, Roche N, Aughney T, Jones N, Day J, Baker J, et al. Barriers and benefits:
implications of artificial night-lighting for the distribution of common bats in Britain and Ireland.
Philos Trans R Soc Lond B Biol Sci. 2015;370(1667).

26. Straka TM, Lentini PE, Lumsden LF, Wintle BA, Ree R van der. Urban bat communities are
affected by wetland size, quality, and pollution levels. Ecol Evol. 2016;6(14):4761–74.

27. van Langevelde Frank, van Grunsven Roy H. A., Veenendaal Elmar M., Fijen Thijs P. M. Artificial
night lighting inhibits feeding in moths. Biol Lett. 2017;13(3):20160874.

28. Macgregor CJ, Pocock MJO, Fox R, Evans DM. Effects of street lighting technologies on the
success and quality of pollination in a nocturnally pollinated plant. Ecosphere.
2019;10(1):e02550.

29. Owens ACS, Lewis SM. The impact of artificial light at night on nocturnal insects: A review and
synthesis. Ecol Evol. 2018;1–22.

30. Macgregor CJ, Evans DM, Fox R, Pocock MJO. The dark side of street lighting: impacts on moths
and evidence for the disruption of nocturnal pollen transport. Glob Chang Biol. 2017;23(2):697–
707.

31. Wilson JF, Baker D, Cheney J, Cook M, Ellis M, Freestone R, et al. A role for artificial night-time
lighting in long-term changes in populations of 100 widespread macro-moths in UK and Ireland:
a citizen-science study. J Insect Conserv. 2018;22(2):189–96.

32. Wakefield A, Jones G, Stone EL, Harris S, Jones G, Harris S. Light-emitting diode street lights
reduce last-ditch evasive manoeuvres by moths to bat echolocation calls. R Soc open sci.
2015;2(8):150291.

33.   Svensson AMA, Rydell J. Mercury vapour lamps interfere with the bat defence of tympanate
435   moths (Operophtera spp.; Geometridae). Animal Behav. 1998;55(1):223–6.

436   34.   Minnaar C, Boyles JG, Minnaar IA, Sole CL, McKechnie AE. Stacking the odds: light pollution may
437   shift the balance in an ancient predator-prey arms race. J Appl Ecol. 2015;52(2):522–31.

438   35.   Acharya L, Fenton MB. Bat attacks and moth defensive behaviour around street lights. Can J
439   Zool. 1999;77(1):27–33.

440   36.   Agee HR, Webb JC. Effects of ultrasound on capture of *Heliothis zea* and *Ostrinia nubilalis*
441   moths in traps equipped with ultraviolet lamps. Ann Entomol Soc Am. 1969;62(6):1248–52.

442   37.   Treat AE. Comparative moth catches by an ultrasonic and a silent light trap. Ann Entomol Soc
443   Am. 1962;55(6):716–20.

444   38.   Surlykke A, Filskov M, Fullard JH, Forrest E. Auditory relationships to size in noctuid moths:
445   bigger ist better. Naturwissenschaften. 1999;86(5):238–41.

446   39.   Ter Hofstede HM, Goerlitz HR, Ratcliffe JM, Holderied MW, Surlykke A. The simple ears of
447   noctuoid moths are tuned to the calls of their sympatric bat community. J Exp Biol.
448   2013;216(21):3954–62.

449   40.   Nakano R, Takanashi T, Surlykke A. Moth hearing and sound communication. J Comp Physiol A.
450   2015;201(1):111–21.

451   41.   ter Hofstede HM, Ratcliffe JM, Fullard JH. Nocturnal activity positively correlated with auditory
452   sensitivity in noctuoid moths. Biol Lett. 2008;4(3):262–5.

453   42.   Skals N, Surlykke A. Hearing and evasive behaviour in the greater wax moth, *Galleria mellonella*
454   (Pyralidae). Physiol Entomol. 2000;25(4):354–62.

455   43.   Rydell J, Skals N, Surlykke A, Svensson M. Hearing and bat defence in geometrid winter moths.
456   Proc Biol Sci. 1997 Jan 22;264(1378):83–8.

457   44.   Surlykke A, Filskov M, Sanderson CG, Lal D, Peters B, Handb, et al. Hearing in geometrid moths.
458   Naturwissenschaften. 1997;84:356–9.

459   45.   Skiba R. Europäische Fledermäuse. 2. Edition. Magdeburg: VerlagsKG Wolf; 2014.

460   46.   Obrist MK, Boesch R, Fluckiger PF. Variability in echolocation call design of 26 Swiss bat species:
461   consequences, limits and options for automated field identification with a synergetic pattern
462   recognition approach. Mammalia. 2004;68:307–22.

463   47.   Gordon, Ter Hofstede HM. The influence of bat echolocation call duration and timing on
464   auditory encoding of predator distance in noctuoid moths. J Exp Biol. 2018;221(6):jeb171561.

465   48.   Holderied MW, von Helversen O. Echolocation range and wingbeat period match in aerial-
466   hawking bats. Proc Royal Soc B. 2003;270:2293–9.

467   49.   Merckx T, Slade EM. Macro-moth families differ in their attraction to light: implications for
468   light-trap monitoring programmes. Insect Conserv Divers. 2014;7(5):453–61.

469   50.   Truxa C, Fiedler K. Attraction to light - from how far do moths (Lepidoptera) return to weak
470   artificial sources of light? Eur J Entomol. 2012;109(1):77–84.

51. Seibert A-M, Koblitz JC, Denzinger A, Schnitzler H-U. Bidirectional echolocation in the bat
Barbastella barbastellus: different signals of low source level are emitted upward through the
nose and downward through the mouth. PLoS One. 2015;10(9):e0135590.

52. Lewanzik D, Goerlitz HR. Continued source level reduction during attack in the low-amplitude
bat *Barbastella barbastellus* prevents moth evasive flight. Funct Ecol. 2017;32(5):1251–61.

53. Madsen BM, Miller LA. Auditory input to motor neurons of the dorsal longitudinal flight
muscles in a noctuid moth (*Barathra brassicae L.*). J Comp Physiol. 1987 Jan 1;160(1):23–31.

54. Roeder KD. Responses of the less sensitive acoustic sense cells in the tympanic organs of some
noctuid and geometrid moths. J Insect Physiol. 1974;20(1):55–66.

55. Surlykke A. Hearing in hooktip moths (Drepanidae: Lepidoptera). J Exp Biol. 2003;206(15):2653–
63.

56. Waters DA, Jones G. The peripheral auditory characteristics of noctuid moths: responses to the
search-phase echolocation calls of bats. J Exp Biol. 1996;199(4):847–56.

57. Zha Y, Chen Q, Lei C. Ultrasonic hearing in moths. Ann Soc Entomol Fr. 2009;45(2):145–56.

58. Goerlitz HR, ter Hofstede HM, Zeale MRK, Gareth J, Holderied MW. An aerial-hawking bat uses
stealth echologcation to counter moth hearing. Curr Biol. 2010;20:1568–72.

59. Hügel T, Goerlitz HR. Species-specific strategies increase unpredictability of escape flight in
eared moths. Funct Ecol. 2019;0(0).

60. Knief U, Forstmeier W. Violating the normality assumption may be the lesser of two evils |
bioRxiv [Internet]. bioRxiv; 2019. Available from: https://doi.org/10.1101/498931

61. Bates D, Maechler M, Bolker B, Walker S. Fitting linear mixed-effects models using lme4. J Stat
Softw. 2015;67(1):1–48.

62. Kuznetsova A, Brockhoff PB, Christensen RHB. lmerTest package: tests in linear mixed effects
models. J Stat Softw. 2017;82(1):1–26.

63. Korner-Nievergelt F, Rpth T, von Felten S, Guelat J, Almasi B, Korner-Nievergelt P. blmeco
[Internet]. [cited 2019 Mar 21]. (R package). Available from: https://rdrr.io/cran/blmeco/

64. Roeder KD, Treat AE, Vandeberg JS. Auditory sense in certain sphingid moths. Science.
1968;159(3812):331–3.

65. Göpfert MC, Wasserthal LT. Auditory sensory cells in hawkmoths: identification, physiology and
structure. J Exp Biol. 1999;202:1579–87.

66. Göpfert MC, Wasserthal LT. Hearing with the mouthparts: behavioural responses and the
structural basis of ultrasound perception in acherontiine hawkmoths. J Exp Biol.
1999;202(8):909–18.

67. Becker A, Whitfield AK, Cowley PD, Järnegren J, Næsje TF. Potential effects of artificial light
associated with anthropogenic infrastructure on the abundance and foraging behaviour of
estuary-associated fishes. J Appl Ecol. 2013;50(1):43–50.