Light suppresses acoustically triggered predator-avoidance behaviours in moths

Theresa Hügel[1]*, Holger R. Goerlitz[1]*

[1] Acoustic and Functional Ecology Group, Max Planck Institute for Ornithology, Seewiesen,

Germany

*corresponding authors:

Acoustic and Functional Ecology Group

Max Planck Institute for Ornithology

Eberhard-Gwinner-Str. 11

82319 Seewiesen, Germany

Email: thuegel@orn.mpg.de, hgoerlitz@orn.mpg.de

ORCID:

Theresa Hügel: 0000-0003-0500-1988

Holger Goerlitz: 0000-0002-9677-8073

ABSTRACT

Urbanization exposes wild animals to increased light. Nocturnal animals in particular are

affected by artificial light. Under its influence, the balance of predator-prey interactions may

shift. In the interaction between echolocating bats and eared moths, bats often benefit from

the effects of light because moths exposed to light resort less often to last-ditch manoeuvres

in response to attacking nearby bats. To what extent increased light affects moths' ability to

engage in negative phonotaxis, a response to distant bats, is unknown. To quantify the

overall effect of light on both last-ditch manoeuvres and negative phonotaxis, we counted

the moths caught by two light traps over several nights. Traps were alternately equipped

with speakers playing back ultrasonic simulations of hunting bats. The playback did not

affect moth counts, indicating that light not only reduces the last-ditch manoeuvres of

moths but also influences whether moths rely on negative phonotaxis. Increasing light

pollution, therefore, interferes with different aspects of anti-predator behaviour in moths

and, possibly, other eared insects. Our study demonstrates the negative effect of light on

insects, clarifies the behavioural changes that result from light exposure and highlights the

need for future research to uncover the mechanism(s) underlying these behavioural

changes.

KEY WORDS

Artificial light at night, predator-prey, bats, Noctuidae, light traps, playback experiment

BACKGROUND

One-fifth of the world's human population can no longer see the Milky Way (1), because of the substantial increase of artificial light at night over the last decades (2,3). Plants and animals are also known to be adversely affected by light pollution (4–6). Light pollution not only impacts a single species' orientation, reproduction or communication (4), it also affects whole communities, for example, by shifting the balance of trophic interactions such as predator-prey interactions (7–12).

Echolocating bats preying upon eared moths represent a global predator-prey system with important ecological impacts (13–16). Bats and moths are primarily nocturnal and rely strongly on their sense of hearing to communicate, orientate, hunt or defend themselves. Bats echolocate to search for food (17,18). Eared moths are able to hear those echolocation calls and to respond with a two-staged anti-predator flight behaviour (19,20). During stage one, moths respond with negative phonotaxis, i.e. they fly away from the calling bat. Negative phonotaxis occurs when the bat is still far away and helps the moth avoid detection. During stage two, moths engage in last-ditch evasive behaviours such as attempting erratic manoeuvres or diving toward the ground. Last-ditch manoeuvres, which occur when the bat is nearby, enable the moth to avoid being captured by the bat. Whether negative phonotaxis or last-ditch manoeuvres are triggered depends on the acoustic properties of the bat's echolocation call as received at the moth's ear (19,21). The probably most important of these properties is received sound pressure level (SPL) (21–23), as received SPL of distant bats is fainter than that of nearby bats due to spherical spreading and atmospheric attenuation (24). In addition, call length and call repetition rate are also important (22,25). All of these properties provide information about the bat's distance and behaviour (17).

Despite their predominantly acoustic interaction, both bats and moths are also

strongly affected by artificially introduced light sources. While some bats may profit from

artificial light because they are able to exploit the resulting accumulation of prey animals

(26,27), other species are negatively affected because they are unable to commute without

interruption to new habitats (28–30). Generally all moths are negatively affected by light.

Light strongly attracts moths, reducing the time they spend feeding (31,32) and interrupting

their commuting behavior (33). Among the ecological consequences is reduced pollination

(34) and population decline (31,35). Light also increases moths' predation risk in two ways.

First, because the accumulation of moths around lights attracts bats, the predation pressure

on moths increases (26,27). Second, light impedes moths' anti-predator flight behaviour.

Several studies compared the sound-triggered anti-predator flights of moths under lit and

unlit conditions, and reported that light reduces anti-predator flight behaviours. Wakefield

et al. (36), who studied last-ditch manoeuvres, showed that only 24% of moths performed

power-dives under LED illumination compared to 60% of moths in the dark; that is, LED-

illumination inhibited last-ditch manoeuvres in 60% of the moths that reacted in darkness.

Similarly, Svensson & Rydell (37) reported a reduction of ~60% in mostly last-ditch evasive

behaviours in moths attempting to escape from bats within a radius of 1 meter around the

light source compared to in moths in the dark (where all moths reacted). Minnaar et al. (38)

reported the most extreme effect: while bat diet (as proxy for moth capture rate) in unlit

conditions was best explained by a model that included anti-predator behaviour, bat diet in

lit conditions was best explained by a model that included 100% reduction in anti-predator

flight behaviour.

In another set of studies (39–41), light exposure was kept constant while the sound

received by the moth was manipulated. Those results showed that moths still exhibited

some degree of anti-predator flight under illumination. Acharya & Fenton (39) compared

last-ditch manoeuvres in eared and deafened moths under illumination, showing that 48% of

eared moths exhibited last-ditch manoeuvres when preyed on by bats, whereas deafened

moths did not. Agee & Webb (40) and Treat (41) compared the number of caught moths at

light traps with and without ultrasonic stimuli. In both studies, capture rates at light traps

with ultrasound playback were lower than capture rates at silent light traps; differences

depended on stimulus design and moth family. Across different stimulus types, capture rates

at light traps with ultrasound were reduced by 51-86 percentage points compared to

capture rates at silent traps in a Noctuid species (*Heliothis zea*, (40)), and by 8-49 percentage

points for different eared moth families (with minimum sample sizes of 37 individuals in

total (41)).

In summary, while the first set of studies shows that light suppresses the sound-

triggered anti-predator flight of moths (ranging from 60-100 percentage points suppression),

the second set of studies shows that sound can still trigger anti-predator flight even in the

light (ranging from 8-86 percentage points). Therefore, how and to what extent light

suppresses anti-predator flight in moths is unclear. Importantly, however, in most of the

studies, only last-ditch manoeuvres are reported (36,37). Even in the studies of Treat (41)

and Agee & Webb (40), where the stimuli may have elicited both stages of evasive flight,

last-ditch manoeuvres likely dominated the outcome due to the study. As both studies used

directional light- and sound sources, it is likely that the moths that were attracted to the

light were always exposed to such high sound pressure levels that last-ditch manoeuvres

were elicited. Only Minnaar et al. (38) investigated both stages of evasive flight, albeit

indirectly. They observed the diet of naturally foraging bats under lit and dark conditions,

suggesting that light under more naturalistic conditions may completely suppress anti-

predator flight throughout all moth populations. Yet, no direct quantification of the overall

effect of light on both stages of the anti-predator flight in moths has been made. Here, we

designed an experiment to investigate the effects of light on both stages of evasive flight,

negative phonotaxis and last-ditch manoeuvres. Like Agee & Webb and Treat (40,41), we

compared moth captures at two light traps, and then simulated bat predation via alternating

ultrasonic playback at one of the traps. We combined an omnidirectional light field that

would attract moths equally with a directional sound field. Therefore, sound intensities

around the light trap varied and triggered either last-ditch manoeuvres (high on-axis SPL) or

negative phonotaxis (low off-axis SPL) in proximity to the trap.

We predicted three different outcomes, depending on the effect of light on the two

stages of moths' anti-predator flight: (1) No moth captures at the ultrasonic light trap,

indicating that both stages of sound-elicited anti-predator flight behaviour are functional

and neither stage is suppressed by light. (2) Reduced moth capture rates at the ultrasonic

light trap, indicating that light partly suppresses sound-elicited anti-predator flight

behaviour. (3) Equal captures at both light traps, indicating that light completely suppresses

both stages of sound-elicited antipredator flight.

METHODS

Setup, study site, moth capture and measurement

We compared the number of caught moths between two light traps, one of which had been

additionally equipped with a loudspeaker, to examine the effect of light on the evasive flight

behaviours of eared moths. We set up two equal-intensity light traps (Sylvania, blacklight,

F15W/350BL-T8, USA, Fig. 1A) at 30 m distance to each other; one was situated at a path in a

forest (trap A) and one was in the forest itself (trap B), close to the Max Planck Institute for

Ornithology, Seewiesen, Germany. Both traps hung freely at ~1.7 m above ground, radiating

light at 360° in the horizontal plane (Fig. 1A). In 2018, we collected data over 15 rainless

nights between July 19 and August 16. The number of nights for data collection was based

on the number of nights sampled in Agee & Webb and Treat (6 nights and 2-12 nights,

respectively (40,41)). Each test night, we equipped one or the other of the two traps with

two loudspeakers (Ultrasonic Dynamic Speaker ScanSpeak, Avisoft Bioacoustics, Glienicke,

Germany); both broadcasting an ultrasonic stimulus (see below) to simulate echolocating

bats. Both speakers were fixed back to back facing in opposite directions and were mounted

above one of the light traps at ~2 m above ground. Thus, each trap was associated for 7-8

nights with the sound of predating bats. In the evening (between 20:30h and 23:40h), we

turned on the light of the traps and the acoustic playback, and the next morning (between

7:00h and 9:40h), we turned off the light and the playback. Every morning we emptied the

traps and counted all moths from the three ear-possessing families Noctuidae, Geometridae

and Erebidae. We measured individuals' body lengths along the middorsal line (from the

head to the end of the abdomen), to correct for the fact that larger moths tend to be more

sensitive to sound than smaller moths (42,43). For statistical analysis, we assigned

individuals to one of six categories, covering sizes from 1 to 3.9 cm in bins of 0.5 cm (for

details, see R-script in Supplemental Information, SI). For those individuals whose body

length we could not measure (e.g., due to a missing abdomen, N = 137, 15.5%), we used

either the mean value of the same species or, if this was not possible (N = 1), the mean value

of the family.

Ultrasonic playback stimulus design and evaluation

We simulated the predation pressure that echolocating bats represent by playing an ultrasonic stimulus consisting of repeated pure tones. The stimulus was designed based on the acoustic properties of bat echolocation calls (call frequency, duration, and interval) and the hearing system of moths. Pulse frequency was 35 kHz to match most moths' best hearing threshold between 20-50 kHz (Noctuidae: (43,44); Erebidae: (45); Pyralidae: (25); Geometridae: (46,47); Sphingidae/Drepanidae: (44)). Pulse duration was 10 ms including 2 ms linear rise and fall time, which corresponds to the longer calls of European open space bats (48,49) and according to which we hoped to optimize detection by moths and information transfer about bat distance (22). The interpulse interval was 100 ms, matching the call interval of bats' echolocation calls in their search phase (60 - 200 ms, (48,50)). On-axis sound pressure level (SPL) was 98 dB SPL re. 20 µPa RMS at 1 m distance (see below for a detailed description of the sound field). This stimulus was presented continuously in a loop throughout the night via the loudspeakers using Avisoft-RECORDER software (Avisoft Bioacoustics), a sound card (Avisoft UltraSoundGate 116, Avisoft Bioacoustics) and a laptop computer.

To validate the effect of the stimulus in the dark, without the potentially suppressing influence of light, we exposed free-flying moths in a dark flight room (5.3 m x 3.5 x 3 m^3) to the stimulus. We caught moths at trap A over the course of four nights and tested them within 30 hours after capture. We placed moths on the ground of the flight room and recorded the flight paths of upward-flying moths with an IR-sensitive camera (Sony HDR-CX560, Sony, Tokio, Japan) under IR illumination (850 nm, Mini IR Illuminator TV6700, ABUS Security-Center, Wetter, Germany). Using the same audio equipment as described above, we presented the stimulus by manually triggering it when a moth flew in front of the speaker.

The moth's distance to the speaker at stimulus onset was about 1-2 m. We categorized the

recorded flight paths as "reaction" when the flight direction, level of erraticness or both

changed with the stimulus onset (for examples, see video in SI); as "no reaction" when we

did not observe those changes; or as "ambiguous" when we could not assign the flight path

to one of the other categories. In total, of the 32 flight paths we evaluated, ten moths (31.2

%) showed a reaction, twelve moths (37.5%) did not show a reaction, and eleven moths

(34.3%) were categorized as "ambiguous".

Overlap of sound and light field

The range and geometry of the presented sound and light fields differed. While the light was

emitted omnidirectionally in the horizontal plane, the sound field was directional (Fig. 1). We

estimated the biologically relevant range for attracting moths by light and the biologically

relevant sound fields for triggering moths' evasive flight based on literature values and our

own measurements.

Light traps can attract released moths from up to 80 m, yet recapture dramatically

decrease beyond 15 m, and additionally depended on family (51,52). Family-specific models

estimated 5%-recapture rate to be 10 ± 6 m (mean ± SEM) for Noctuidae and 23 ± 12 m for

Geometridae (51). Note, however, that these data were obtained with a different light

source than ours (6W actinic vs. 15W blacklight in our case), and that the radius of attraction

to light is not necessarily equivalent to the radius over which light interferes with anti-

predator behaviour. Yet, we used these family-specific values as first estimates of the

biologically relevant distances beyond which moths would not be attracted to our light trap

(Fig. 1B, dashed lines).

To estimate the effect of the playback, we measured the playback level of the

loudspeaker from 0° on-axis to 90° off-axis in steps of 5° (for details about calculations, see

SI). The on-axis source level was 97 dB SPL re. 20 µPa RMS @ 1 m distance, i.e., within the

lower range of natural bat call levels in the field (100-120 dB peSPL @ 1 m (50), with

corresponding RMS-SPL values being ~3-7 dB lower than the peSPL values (53,54)). With

increasing off-axis angle, playback source level was reduced by up to ~30 dB at 45°, resulting

in a minimum playback level of 70 dB SPL RMS @ 1m. We then calculated the distance to the

loudspeaker at which the playback would reach biologically relevant levels of 60 and 80 dB

SPL RMS as a function of angle around the loudspeaker. We chose 60 and 80 dB SPL as

approximate thresholds for eliciting negative phonotaxis and last-ditch manoeuvres,

respectively, based on several lines of evidence. The triggering of negative phonotaxis and

last-ditch manoeuvres in moths is thought to be correlated with the activity of moths'

auditory receptor neurons, A1 and A2 (22,55,56). The neuronal thresholds of these neurons

vary with frequency and species (22,43,55,57–59), with A1 being generally about 20 dB more

sensitive than A2 (43,55,60). The best thresholds of A1 range from ~35-55 dB SPL at the

position of the moth, and the best thresholds of A2 range from ~52-72 dB SPL

(20,22,43,55,57,58). Behavioural thresholds in moths are often unknown, but those that are

known tend to be ~10 dB higher than neuronal thresholds (reviewed in (54)). Indeed, our

own data indicate that sound pulses of ~80 dB SPL elicit last-ditch manoeuvres (61), which is

8-28 dB above known A2 thresholds (20,22,43,55,57,58). We thus defined 80 dB SPL as

threshold which securely elicits last-ditch maneouvres. As no behavioural data exist for the

threshold of negative phonotaxis, we chose 60 dB SPL, which is 20 dB below our threshold

for last-ditch manoeuvres and matches the difference between the A1 and A2 thresholds.

We calculated the distance-isolines for 60 and 80 dB SPLs; these varied with the angle

around the loudspeaker, ranging from 3.6 to14.6 m and 0 to 5.6 m, respectively (Fig. 1B).

Statistical analysis

To test for an effect of light on the moths' negative phonotaxis, we fitted linear mixed-

effects models to the moth count data as a function of the fixed effects *playback*, *trap*, *moth*

family, and *moth body length*, the random factor *date* (to account for repeated measures)*,*

and the interactions of *playback* and *moth family*, and *playback* and *moth body length*. We

modelled error distribution using a negative binomial distribution and tested for

overdispersion using the "DHARMa" package (62). We defined the moth family Noctuidae as

intercept, as it had the largest sample size and thus was the most reliable reference. To test

which factors significantly contributed to the model fit, we conducted backwards model

reduction (63). Hence, the full model was successively reduced, by stepwise removing

covariates, starting wit the covariate with the highest p-value of t-statistics provided by the

model summary. We compared nested models with likelihood ratio tests (type I) using a chi-

squared statistic to find the minimal adequate model. All statistical analyses were conducted

in R version 3.3.2 (R Foundation for Statistical Computing, Vienna, Austria) using the

packages "lme4"(64), "lmerTest"(65), "MASS" (66), and "blmeco" (67). For further details,

see R-script included in SI.

RESULTS

We caught a total of 878 moths over 15 nights (Fig. 2). The minimal adequate model included *playback*, *family* and *size* as fixed factors and *date* as a random factor. Capture success varied among moth families (likelihood ratio test for factor *family*, χ^2 = 59.43, d.f. = 2, p < 0.001). Most of the counted moths (80.6 %) belonged to the family Noctuidae; the remaining 19.4 % was composed of Geometridae (11.3%) and Erebidae (8.1%). We caught more small than large moths (likelihood ratio test for factor *size*, χ^2 = 89.76, d.f. = 5, p < 0.001). Although total capture rates seemed to differ between traps (total capture of 506 (trap A) vs. 365 (trap B) moths), *trap* was not a significant factor of the minimal adequate model (likelihood ratio test, χ^2 = 2.46, d.f. = 1, p = 0.116). That is, the capture rates per night did not significantly differ between the traps (Fig. 2A).

The number of caught moths was similar on nights with and without playback for both traps (Fig. 2A), that is, our playback treatment had no detectable effect on the number of caught moths (likelihood ratio test of factor *playback*, χ^2 = 2.44, d.f. = 1, p = 0.118). The number of caught moths fluctuated strongly between nights (7-116 moths per trap and night), which might be one reason for the low power of our dataset (~38%, see R-script in SI for more information).

We did not detect an interaction between *moth size* and *playback* and between *moth family* and *playback* (likelihood ratio tests, χ^2 = 4.54, d.f. = 3, p = 0.208 and χ^2 = 0.38, d.f. = 2, p = 0.826, respectively), that is, moths of different sizes and families did not react differently to the playback. This missing interaction is likely due to *playback* not being a significant factor of the minimal adequate model.

DISCUSSION

Echolocating bats and eared insects offer a textbook example of predator-prey interactions,

which is based on acoustic information (20). Increasing light pollution (2,3), however,

severely impacts both bats and moths (e.g. (27,28,32), and has potentially cascading effects

on their predator-prey-interactions, population dynamics and ecosystems (11,38). Light

reduces the acoustically triggered last-ditch manoeuvres in eared moths, which in turn

makes them more vulnerable to nearby predating bats. It is less clear, however, whether

light also reduces the ability of eared moths to perform negative phonotaxis, a behaviour

that serves as the moths' first line of defence against distant echolocating bats. To better

understand how artificial light at night affects the eared moths' overall ability to avoid

predators, we compared moth captures at two light traps. One trap was silent, while the

other trap broadcast bat-like ultrasonic stimuli that triggered last-ditch manoeuvres as well

as negative phonotaxis under dark conditions. The number of caught moths did not

significantly differ between traps. Thus, the presented playback did not repel the moths,

suggesting that the light suppressed both anti-predator behaviours, negative phonotaxis and

last-ditch manoeuvres.

In contrast to our results, both Treat (41) and Agee & Webb (40) observed reduced

moth capture rates at the ultrasonic trap, showing that sound repelled eared moths.

Differences in stimulus design and habituation effects of the moths are in our opinion

insufficient to explain the variation in capture rates. Treat (41) and Agee & Webb (40)

broadcast a variety of different stimuli varying in pulse rate, frequency, pulse duration, and

SPLs (ranges: pulse rate: 0.7-155 pulses/s; frequency: 12.5-200 kHz; pulse duration: 2-10 ms;

SPL: ~60-100 dB SPL @ 1 m distance), all of which elicited varying degrees of evasive flight in

eared moths. The acoustic properties of our stimulus was within this range and did elicit

evasive flight under dark control conditions, but did not elicit evasive flight under lit

conditions. Habituation to the ultrasound could have affected individuals that flew close to

the light traps for extended time periods. It is more likely, however, that most caught

individuals randomly encountered and approached the traps over larger distances and thus

did not perceive our presented stimulus continuously. Likewise, Agee & Webb (40) also

presented their ultrasonic stimuli over the course of a whole night and found that ultrasonic

playbacks repelled the moths.

We suggest that major differences in the geometry and overlap of light- and sound-

fields explain these contrasting results. In the setups of Treat (41) and Agee & Webb (40),

sound and light fields almost overlapped. Before entering the light traps in these setups,

approaching moths had to pass through high levels of sound pressure; such levels likely have

been high enough to elicit last-ditch manoeuvres. In both studies, fewer moths were caught

in the ultrasonic traps compared to the silent traps, suggesting that the playback elicited

some anti-predator flight (likely last-ditch manoeuvres) under these light and sound

conditions. Specifically, for stimuli similar to ours, noctuid moths were captured at a rate of

35% for a 37.5 kHz pulsed tone stimulus vs. 65% at the silent trap (41), while in Agee &

Webb (40), the moths were captured at a rate of ~15% for a 30 kHz pulsed tone stimulus vs.

85% at the silent trap for two species of the families Noctuidae and Pyralidae. In contrast,

our setup combined omnidirectional light fields with directional sound fields, a design which

exposed the moths to different direction-dependent sound pressure levels while

approaching the light. Moths approaching the trap on-axis were exposed first to low SPLs

and then to increasingly high SPLs. This spatial pattern of increasing SPL was intended to first

elicit negative phonotaxis and later last-ditch manoeuvress. In contrast, moths approaching

the traps off-axis were exposed to low SPLs that can elicit only negative phonotaxis. As our

capture rates did not differ among the light traps -- indicating that the playback had no

repelling effect on moth flight -- we suggest that the light strongly reduced the negative

phonotaxis of eared moths. Even if the moths still exhibited some degree of last-ditch

manoeuvre close to the trap (as shown by Treat and Agee & Webb, (40,41), these

manoeuvres might have brought the moth into a position off-axis to the loudspeaker, either

to the sides or below, where the SPL is lower. From there, no further last-ditch manoeuvres

was elicted due to the lower SPL, and the light kept attracting the moth into the trap. Our

results are in line with those of Minnaar et al. (38), who found that a model of escape

behaviour in moths assuming 0% efficiency best explained bats' diet under lit conditions. We

therefore suggest that light suppresses not only last-ditch manoeuvres, as previously shown,

but also negative phonotaxis.

In summary, our results underline the strong effect of light on eared moths, providing

an additional explanation for shifts in predator-prey interactions between bats and moths

(11,12,38). Because negative phonotaxis, along with last-ditch manoeuvres, is suppressed by

light, moths are not only unable to escape nearby predators, but they are also unable to

avoid them by flying away. Similarly, fish are attracted to lit areas, where they are "trapped"

and preyed upon by seals (9) and other fish (68). Our results emphasize the need to

understand the mechanism(s) underlying animals' attraction to light and to quantify the

effects of light on animals' behaviour, such as predator avoidance. Light pollution is

increasing and insect populations are declining. Only by understanding the mechanism(s)

underlying these issues can we apply efficient conservation measures.

Ethics

Ethics

The experiments in this study did not require an approval by an ethical committee.

Data accessibility

All data, including R-scripts and supplementary video, is attached via electronic

supplemental material.

Author contributions

TH and HRG conceived of and designed the study. TH collected and analysed the data. TH

and HRG wrote the manuscript. All authors gave final approval for publication and agree to

be held accountable for the work performed therein.

Competing interests

We have no competing interests.

Funding

This study was supported by funds of the Deutsche Forschungsgemeinschaft

(Emmy Noether research grants GO2029/2-1 and GO2029/2-2 to H.R.G.) and the

International Max Planck Research School for Organismal Biology (student grant to T.H.)

Acknowledgements

We thank the IMPRS for support and especially the participants of the course for "Scientific

Writing" 2019 for comments on previous versions of the manuscript. We further thank the

Max Planck Institute for Ornithology Seewiesen for providing excellent infrastructure and

support.

Figure 1: **(A) Experimental setup.** The photo shows one of the light traps with two

loudspeakers attached above the trap, both of which broadcast the acoustic stimulus in

opposite directions. **(B) Side view of the biologically relevant sound field around the light**

trap with attached loudspeakers. Coloured areas indicate areas with minimum sound

pressure levels of 60 and 80 dB SPL re. 20 μPa, which are biologically relevant acoustic

thresholds for eliciting negative phonotaxis and last-ditch manoeuvres, respectively, in eared

moths. Schematic indicates position and dimension of the light trap. **(C) Top view of the**

biologically relevant sound field (coloured areas) and light field (dashed lines). Dashed lines

indicate the distance over which 5% of released noctuid (10 m) and geometrid (23 m) moths

are recaptured at the light trap, respectively (51), as an indicator of the maximum range of

light attraction.

384   **Figure 2: Number of caught moths per night and per playback treatment**. **(A) Number of**

385   **caught moths at both light traps, with and without playback.** Boxplots show median,

386   quartiles and whiskers (up to 1.5 x inter-quartile range) of the daily counts (dots). **(B) Daily**

387   **counts of caught moths per night and per trap for all 15 nights.** White letters at the base of

388   the bars indicate the trap.

REFERENCES

1. Falchi F, Cinzano P, Duriscoe D, Kyba CCM, Elvidge CD, Baugh K, et al. The new world atlas of artificial night sky brightness. Sci Adv. 2016;2(6):e1600377.

2. Hölker F, Moss T, Griefahn B, Kloas W, Voigt C, Henckel D, et al. The dark side of light: a transdisciplinary research agenda for light pollution policy. Ecol Soc. 2010;15(4).

3. Fouquet R. Seven centuries of energy services: the price and use of light in the United Kingdom (1300-2000). Energy Journal. 2006;27:139–77.

4. Longcore T, Rich C. Ecological light pollution. Front Ecol Environ. 2004;2(4):191–8.

5. Davies TW, Smyth T. Why artificial light at night should be a focus for global change research in the 21st century. Glob Chang Biol. 2018;24(3):872–82.

6. Knop E, Zoller L, Ryser R, Gerpe C, Hörler M, Fontaine C. Artificial light at night as a new threat to pollination. Nature. 2017;548(7666):206–9.

7. Davies TW, Bennie J, Inger R, Ibarra NH, Gaston KJ. Artificial light pollution: are shifting spectral signatures changing the balance of species interactions? Glob Chang Biol. 2013;19(5):1417–23.

8. Davies TW, Duffy JP, Bennie J, Gaston KJ. The nature, extent, and ecological implications of marine light pollution. Front Ecol Environ. 2014;12(6):347–55.

9. Yurk H, Trites AW. Experimental attempts to reduce predation by harbor seals on out-migrating juvenile salmonids. Trans Am Fish Soc. 2000;129(6):1360–6.

10. Miller Colleen R., Barton Brandon T., Zhu Likai, Radeloff Volker C., Oliver Kerry M., Harmon Jason P., et al. Combined effects of night warming and light pollution on predator–prey interactions. P Roy Soc P B Biol Sci. 2017;284(1864):20171195.

11. Russo D, Cosentino F, Festa F, De Benedetta F, Pejic B, Cerretti P, et al. Artificial illumination near rivers may alter bat-insect trophic interactions. Environ Pollut. 2019;252:1671–7.

12. Bailey LA, Brigham RM, Bohn SJ, Boyles JG, Smit B. An experimental test of the allotonic frequency hypothesis to isolate the effects of light pollution on bat prey selection. Oecologia. 2019;190(2):367–74.

13. Van Toor ML, O'Mara MT, Abedi-Lartey M, Wikelski M, Fahr J, Dechmann DKN. Linking colony size with quantitative estimates of ecosystem services of African fruit bats. Curr Biol. 2019;29(7):R237–8.

14. Kunz TH, Braun de Torrez E, Bauer D, Lobova T, Flemming TH. Ecosystem services provided by bats. Ann NY Acad Sci. 2011;1223(1):1–38.

15. Kasso M, Balakrishnan M. Ecological and economic importance of bats (order Chiroptera). ISRN Ecol. 2013;Article ID 187415:9 pages.

16. Boyles JG, Cryan PM, McCracken GF, Kunz TH. Economic importance of bats in agriculture. Science. 2011;332(6025):41–2.

17. Schnitzler H-U, Moss CF, Denzinger A. From spatial orientation to food acquisition in echolocating bats. Trends Ecol Evol. 2003;18:386–94.

426   18.   Fenton MB. Eavesdropping on the echolocation and social calls of bats. Mammal Rev.
427   2003;33:193–204.

428   19.   Roeder KD. The behaviour of free flying moths in the presence of artificial ultrasonic pulses.
429   Anim Behav. 1962;10(3–4):300–4.

430   20.   Ter Hofstede HM, Ratcliffe JM. Evolutionary escalation: the bat-moth arms race. J Exp Biol.
431   2016;219(Pt 11):1589–602.

432   21.   Roeder KD. Aspects of the noctuid tympanic nerve response having significance in the
433   avoidance of bats. J Insect Physiol. 1964;10(4):529–46.

434   22.   Gordon, Ter Hofstede HM. The influence of bat echolocation call duration and timing on
435   auditory encoding of predator distance in noctuoid moths. J Exp Biol. 2018;221(6):jeb171561.

436   23.   Svensson M, Rydell J, Brown R. Bat predation and flight timing of winter moths, *Epirrita* and
437   *Operophtera* species (Lepidoptera, Geomentridae). Oikos. 1999;84(2):193–8.

438   24.   Goerlitz HR. Weather conditions determine attenuation and speed of sound: Environmental
439   limitations for monitoring and analyzing bat echolocation. Ecol Evol. 2018;8(10):5090–100.

440   25.   Skals N, Surlykke A. Hearing and evasive behaviour in the greater wax moth, *Galleria mellonella*
441   (Pyralidae). Physiol Entomol. 2000;25(4):354–62.

442   26.   Rydell J. Exploitation of insects around streetlamps by bats in Sweden. Funct Ecol.
443   1992;6(6):744–50.

444   27.   Cravens ZM, Brown VA, Divoll TJ, Boyles JG. Illuminating prey selection in an insectivorous bat
445   community exposed to artificial light at night. J Appl Ecol. 2018;55(2):705–13.

446   28.   Stone EL, Jones G, Harris S. Street lighting disturbs commuting bats. Curr Biol.
447   2009;19(13):1123–7.

448   29.   Mathews F, Roche N, Aughney T, Jones N, Day J, Baker J, et al. Barriers and benefits:
449   implications of artificial night-lighting for the distribution of common bats in Britain and Ireland.
450   Philos Trans R Soc Lond B Biol Sci. 2015;370(1667).

451   30.   Straka TM, Lentini PE, Lumsden LF, Wintle BA, Ree R van der. Urban bat communities are
452   affected by wetland size, quality, and pollution levels. Ecol Evol. 2016;6(14):4761–74.

453   31.   van Langevelde Frank, van Grunsven Roy H. A., Veenendaal Elmar M., Fijen Thijs P. M. Artificial
454   night lighting inhibits feeding in moths. Biol Lett. 2017;13(3):20160874.

455   32.   Macgregor CJ, Pocock MJO, Fox R, Evans DM. Effects of street lighting technologies on the
456   success and quality of pollination in a nocturnally pollinated plant. Ecosphere.
457   2019;10(1):e02550.

458   33.   Owens ACS, Lewis SM. The impact of artificial light at night on nocturnal insects: A review and
459   synthesis. Ecol Evol. 2018;1–22.

460   34.   Macgregor CJ, Evans DM, Fox R, Pocock MJO. The dark side of street lighting: impacts on moths
461   and evidence for the disruption of nocturnal pollen transport. Glob Chang Biol. 2017;23(2):697–
462   707.

35. Wilson JF, Baker D, Cheney J, Cook M, Ellis M, Freestone R, et al. A role for artificial night-time
lighting in long-term changes in populations of 100 widespread macro-moths in UK and Ireland:
a citizen-science study. J Insect Conserv. 2018;22(2):189–96.

36. Wakefield A, Jones G, Stone EL, Harris S, Jones G, Harris S. Light-emitting diode street lights
reduce last-ditch evasive manoeuvres by moths to bat echolocation calls. R Soc open sci.
2015;2(8):150291.

37. Svensson AMA, Rydell J. Mercury vapour lamps interfere with the bat defence of tympanate
moths (Operophtera spp.; Geometridae). Animal Behav. 1998;55(1):223–6.

38. Minnaar C, Boyles JG, Minnaar IA, Sole CL, McKechnie AE. Stacking the odds: light pollution may
shift the balance in an ancient predator-prey arms race. J Appl Ecol. 2015;52(2):522–31.

39. Acharya L, Fenton MB. Bat attacks and moth defensive behaviour around street lights. Can J
Zool. 1999;77(1):27–33.

40. Agee HR, Webb JC. Effects of ultrasound on capture of *Heliothis zea* and *Ostrinia nubilalis*
moths in traps equipped with ultraviolet lamps. Ann Entomol Soc Am. 1969;62(6):1248–52.

41. Treat AE. Comparative moth catches by an ultrasonic and a silent light trap. Ann Entomol Soc
Am. 1962;55(6):716–20.

42. Surlykke A, Filskov M, Fullard JH, Forrest E. Auditory relationships to size in noctuid moths:
bigger ist better. Naturwissenschaften. 1999;86(5):238–41.

43. Ter Hofstede HM, Goerlitz HR, Ratcliffe JM, Holderied MW, Surlykke A. The simple ears of
noctuoid moths are tuned to the calls of their sympatric bat community. J Exp Biol.
2013;216(21):3954–62.

44. Nakano R, Takanashi T, Surlykke A. Moth hearing and sound communication. J Comp Physiol A.
2015;201(1):111–21.

45. ter Hofstede HM, Ratcliffe JM, Fullard JH. Nocturnal activity positively correlated with auditory
sensitivity in noctuoid moths. Biol Lett. 2008;4(3):262–5.

46. Rydell J, Skals N, Surlykke A, Svensson AM. Hearing and bat defence in geometrid winter moths.
Proc Biol Sci. 1997;264(1378):83–8.

47. Surlykke A, Filskov M, Sanderson CG, Lal D, Peters B, Handb, et al. Hearing in geometrid moths.
Naturwissenschaften. 1997;84:356–9.

48. Skiba R. Europäische Fledermäuse. 2. Edition. Magdeburg: VerlagsKG Wolf; 2014.

49. Obrist MK, Boesch R, Fluckiger PF. Variability in echolocation call design of 26 Swiss bat species:
consequences, limits and options for automated field identification with a synergetic pattern
recognition approach. Mammalia. 2004;68:307–22.

50. Holderied MW, von Helversen O. Echolocation range and wingbeat period match in aerial-
hawking bats. Proc Royal Soc B. 2003;270:2293–9.

51. Merckx T, Slade EM. Macro-moth families differ in their attraction to light: implications for
light-trap monitoring programmes. Insect Conserv Divers. 2014;7(5):453–61.

52.   Truxa C, Fiedler K. Attraction to light - from how far do moths (Lepidoptera) return to weak
501   artificial sources of light? Eur J Entomol. 2012;109(1):77–84.

502   53.   Seibert A-M, Koblitz JC, Denzinger A, Schnitzler H-U. Bidirectional echolocation in the bat
503   *Barbastella barbastellus*: different signals of low source level are emitted upward through the
504   nose and downward through the mouth. PLoS One. 2015;10(9):e0135590.

505   54.   Lewanzik D, Goerlitz HR. Continued source level reduction during attack in the low-amplitude
506   bat *Barbastella barbastellus* prevents moth evasive flight. Funct Ecol. 2017;32(5):1251–61.

507   55.   Madsen BM, Miller LA. Auditory input to motor neurons of the dorsal longitudinal flight
508   muscles in a noctuid moth (*Barathra brassicae L.*). J Comp Physiol. 1987 Jan 1;160(1):23–31.

509   56.   Roeder KD. Responses of the less sensitive acoustic sense cells in the tympanic organs of some
510   noctuid and geometrid moths. J Insect Physiol. 1974;20(1):55–66.

511   57.   Surlykke A. Hearing in hooktip moths (Drepanidae: Lepidoptera). J Exp Biol. 2003;206(15):2653–
512   63.

513   58.   Waters DA, Jones G. The peripheral auditory characteristics of noctuid moths: responses to the
514   search-phase echolocation calls of bats. J Exp Biol. 1996;199(4):847–56.

515   59.   Zha Y, Chen Q, Lei C. Ultrasonic hearing in moths. Ann Soc Entomol Fr. 2009;45(2):145–56.

516   60.   Goerlitz HR, ter Hofstede HM, Zeale MRK, Gareth J, Holderied MW. An aerial-hawking bat uses
517   stealth echolocation to counter moth hearing. Curr Biol. 2010;20:1568–72.

518   61.   Hügel T, Goerlitz HR. Species-specific strategies increase unpredictability of escape flight in
519   eared moths. Funct Ecol. 2019;0(0).

520   62.   Hartig F. DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed) Regression
521   Models [Internet]. 2019. Available from: https://CRAN.R-project.org/package=DHARMa

522   63.   Lewis F, Butler A, Gilbert L. A unified approach to model selection using the likelihood ratio test.
523   Methods Ecol Evol. 2011;2(2):155–62.

524   64.   Bates D, Maechler M, Bolker B, Walker S. Fitting linear mixed-effects models using lme4. J Stat
525   Softw. 2015;67(1):1–48.

526   65.   Kuznetsova A, Brockhoff PB, Christensen RHB. lmerTest package: tests in linear mixed effects
527   models. J Stat Softw. 2017;82(1):1–26.

528   66.   Venables WN, Ripley BD. Modern Applied Statistics with S. Fourth Edition. New York: Springer;
529   2002.

530   67.   Korner-Nievergelt F, Rpth T, von Felten S, Guelat J, Almasi B, Korner-Nievergelt P. blmeco
531   [Internet]. [cited 2019 Mar 21]. (R package). Available from: https://rdrr.io/cran/blmeco/

532   68.   Becker A, Whitfield AK, Cowley PD, Järnegren J, Næsje TF. Potential effects of artificial light
533   associated with anthropogenic infrastructure on the abundance and foraging behaviour of
534   estuary-associated fishes. J Appl Ecol. 2013;50(1):43–50.