Cross-continental biogeography of the common bean rhizosphere microbiome reveals a

persistent core membership

Nejc Stopnisek[1,2,3] and Ashley Shade[1,2,3,4,5]

1. Department of Microbiology and Molecular Genetics, Michigan State University, East

Lansing, MI 48840 USA

2. The Plant Resilience Institute, Michigan State University, East Lansing, MI 48840

3. The DOE Great Lakes Bioenergy Research Center, Michigan State University, East

Lansing, MI 48840

4. Program in Ecology, Evolutionary Biology and Behavior, Michigan State University, East

Lansing, MI 48840

5. Department of Plant, Soil and Microbial Sciences, Michigan State University, East Lansing,

MI 48840

Corresponding author: shadeash@msu.edu

Keywords: plant-microbiome, Phaseolus vulgaris, plant health, neutral model, co-occurrence

network, rhizobia, fertilizer, abundance-occupancy, 16S rRNA, ITS, microbial ecology, soil, hub

taxa, connector taxa, plant genotype, sustainable agriculture

Significance statement

Ongoing and magnifying consequences of environmental change present a complex challenge for plants, food security and sustainable agriculture. One mechanism by which plants may promote long-term stress tolerance is via beneficial interactions with the microbiome. However, there are thousands of microbiome members in the rhizosphere, and it is difficult to know which engage with the plant. We applied biogeography and assembly concepts from ecology to discover a core rhizosphere microbiome for the globally important legume, common bean. The core microbiome was not influenced by genotype, local soil properties or agricultural management, and was cosmopolitan among beans grown in North and South America. The insights and core approach provided here advance goals to manage microbiomes to support plant health and resilience.

Abstract

Plants recruit soil microbes that provide nutrients, promote growth and protect against pathogens. However, the full potential of microbial communities for supporting plant health and agriculture is unrealized, in part because rhizosphere members key for plant health are difficult to prioritize. Microbes that ubiquitously associate with a plant species across large spatial scales and varied soil conditions provide a practical starting point for discovering beneficial members. Here, we quantified the structures of bacterial/archaeal and fungal communities in the common bean rhizosphere (*Phaseolus vulgaris*), and assessed its core membership across two evolutionarily-distinct genotypes. Beans were grown in field conditions across five major growing regions in the United States. We discovered a conserved core microbiome of 271 bacterial, archaeal and fungal taxa that were consistently associated with the common bean, despite different soil types, management, climates and genotypes. Neutral models of abundance-occupancy relationships and co-occurrence networks suggest that these core taxa are in intimate relationships with the plant,

rather than important members of the local soil microbiome. We expanded our study to leverage

rhizosphere samples inclusive of eight additional common bean genotypes that were grown in

Colombian soils. Surprisingly, there were 48 persistent bacterial taxa that were detected in all

samples, inclusive of U.S. and Colombian-grown beans. Many of the core taxa were yet-uncultured

and affiliated with Proteobacteria; these taxa are prime targets for functional investigation in

support of sustainable common bean agriculture. More generally, our approach provides insights

into microbial taxa that can be prioritized towards translational studies of plant-microbiome.

management.

Introduction

Current agricultural practices rely on unsustainable levels of agrochemicals that can harm the environment. Agriculture requires more efficient use of available resources, and the naturally-occurring, soil-dwelling microbiota offers potential to contribute to the responsible intensification of agriculture. Selection and breeding of plants for their beneficial associations with microbiota has promise to deliver a new generation of microbe-improved plants (1–7). The ideal outcome of such efforts would achieve a balance of sustainable agriculture with food security. To achieve this, we must understand the relationships between plant genomes and microbiomes. To date, several studies have investigated the influence of plant genotype on the plant microbiome (8–11). Though there is not complete agreement among these studies, they generally suggest that plant genotype has a limited influence on the microbiome relative to the local environmental conditions, especially when grown in field conditions (12–15) (as opposed to greenhouse or controlled environments (9, 10)). There is clearly more work needed to advance goals of microbe-improved plants.

Common bean (*Phaseolus vulgaris* L.) is the most important food legume grown worldwide, and especially for developing economies in South America, Africa and Asia (16). The origin of common bean is nowadays central Mexico, and from there it spread to central and south America around 165,000 years ago (17). This resulted in the development of two major eco-geographically distinct common bean gene pools with partial reproductive isolation (18–20). The Mesoamerican gene pool was distributed from northern Mexico to Colombia and the Andean gene pool ranged from southern Peru to northwestern Argentina. Since 8,000 years ago, each pool was separately and selectively bred, leading to further diversification between them (21–23). Because of pre-existing genetic differences in each gene pool followed by divergent breeding history, common bean presently offers a distinctive model for understanding crop evolution and plant domestication on plant-microbiome interactions.

The objective of this study was to quantify the influences of genotype and cross-continental

biogeography on the common bean root microbiome, inclusive of bacteria, archaea and fungi. We

also aimed to identify core microbial lineages that have cosmopolitan associations with common

bean and therefore potential as targets towards microbe-improved plant breeding. With the

cooperation of U.S. common bean breeding programs, we executed a first-of-its kind study of two

divergent bean genotypes grown under field conditions in five major North American bean

growing regions. We additionally performed a meta-analysis of the bacterial microbiome with bean

genotypes grown in South America (9). Surprisingly and excitingly, we discovered a core bean

rhizosphere microbiome of a handful of members that persistently and consistently associated with

this nutritionally, agronomically, and economically important crop. This core was discovered in

spite of apparent rhizosphere microbiome divergences that were attributable to differences in local

soil conditions and management. We did not detect an influence of plant genotype, suggesting that

this core membership supersedes it.

Results and Discussion

Study design, sequencing summary, and alpha diversity

To assess the influence of plant genotype and environment on the common bean rhizosphere microbiome, we designed a biogeography study of two divergent bean genotypes (from Mesoamerican and Andean gene pools (24, 25)), both grown in the field at five research and extension farms that represent major U.S. bean production regions (**Table 1**) (26). From each growing location, triplicate bean plants grown in three to four plots were harvested at flowering (mid to late July).

We sequenced 39 rhizosphere (attached to the root, see methods) and root-associated soil samples using 16S rRNA gene and ITS1 region, respectively to assess microbiome community structures. We assessed the bacterial and archaeal communities using 16S rRNA gene amplicon sequencing, and the fungal communities using ITS1 amplicon sequencing. After clustering reads to OTUs at 97% similarity threshold and filtering chloroplast and mitochondria, there was 31,255 to 506,166 16S rRNA reads per sample and 22,716 to 252,810 ITS reads. We rarefied samples to 31,255 reads for 16S rRNA gene amplicons and to 22,716 for ITS. With these thresholds, we achieved richness asymptotes for both datasets, suggesting that sequencing efforts were sufficient to capture comparative dynamics and diversity (**SI Appendix, Fig. S1**). The total richness observed at this rarefaction depth was 1,505 fungal and 23,872 bacterial and archaeal OTUs.

As reported in other rhizosphere studies, the total fungal diversity was lower than bacterial /archaeal diversity in the rhizosphere of the common bean (27–29). Richness varied by growing location (ANOVA, F value=12.4, *p*-value<0.0001 and F value=13.1, *p*-value<0.0001 for 16S rRNA and ITS data, respectively, **SI Appendix, Fig. S2**).

Beta diversity: the impact of biogeography and plant genotype

There were differences across growing locations in edaphic soil properties (e.g. pH,

nitrogen species, organic matter) as well as in management practices (e.g. fertilization, irrigation,

crop rotation) and climate (**Table 1**). Growing location, pH and fertilization explained differences

in microbial community structure for bacteria/archaea and fungi, but plant genotype did not (**Fig.**

1AB, **SI Appendix, Dataset S1**). Fungal and bacterial/archaeal communities had synchronous

biogeographic patterns (Procrustes m12 squared: 0.2137, R^2=0.8867 and p-value=0.001),

supporting that both communities are shaped by local edaphic factors. An effect of growing

location on beta-diversity has been previously reported for plant- and soil- associated microbiomes

(11, 13, 14, 30).

We selected two distinct bean genotypes with divergent evolutionary and breeding histories

to assess the impact of plant genotype on the root microbiota, expecting that any signal would be

maximized between these lineages. Briefly, these two genotypes belong to the two major lineages

of common bean that diverged and bottlenecked ∼165,000 years ago, and represent the

Mesoamerican (Eclipse) and Andean (CELRK) gene pools (17). There was a weak interaction

between plant genotype and growing location in explaining fungal community structure

(PERMANOVA, R^2=0.0377, p-value=0.001) (**SI Appendix, Dataset S1**), but no effect of plant

genotype alone on the bacterial and archaeal communities, even when controlling for growing

location. Similarly, plant genotype had a weak but statistically detectable effect on bacterial and

archaeal richness (p-value=0.019) but no measurable effect on fungal richness (p-value=0.95). (**SI**

Appendix, Fig. S2CD). An analysis of differential taxon abundances across genotypes (DESeq2

(31)) detected 20 taxa out of 21,881 total that were distinguishing between CELRK and Eclipse,

but their adjusted p-value were marginally significant (> 0.02; **SI Appendix, Dataset S2**). The

absence of a robust genotype effect on the microbiome is in contrast to a previous study on

common bean (9). However, several studies have reported a relatively weak to no influence of

genotype more generally on the rhizosphere communities of other plants grown in the field (14, 30,

32). The study that observed a genotype effect on the bean root microbiota included wild,

domesticated and landrace common beans of Mesoamerican origin that were grown in a

greenhouse experiment using Colombian field soil (9). The authors attributed those bean genotype

differences to differences in root architecture, which often has a different phenotype in greenhouse

pots than in the field (33). We summarize that, in this study and in agreement with other field

studies, plant genotype has a minor to no measurable impact on rhizosphere microbiome.

A core rhizosphere microbiome across U.S. bean growing regions

The common bean rhizosphere microbiome included the major expected lineages for both

bacteria and fungi (**SI Appendix, Fig. S3AB**), in agreement with other plant rhizosphere studies

(10, 15, 34–37). Together with previous studies, these data provide more evidence that root-

associated microbial taxa are phylogenetically and functionally conserved (38). Proteobacteria,

Acidobacteria, Bacteroidetes, and Actinobacteria collectively comprised on average 73.5% of the

bacteria/archaeal community, Ascomycota dominated the fungal community with a mean total

relative abundance of 53% with notable sample-to-sample variance (range from 16.5% to 84.5%).

Despite differences in the rhizosphere microbiomes across growing locations, we noticed a

large number of OTUs that were shared for the bacterial/archaeal communities (2,173 taxa, mean

31.5%, range 29.5% to 34.7%). There was a smaller but notable overlap for the fungal communities

(70 taxa or mean 4.5%, range from 0.9% to 17.9%; **SI Appendix, Fig. S3CD**). These data

suggested that, despite measured edaphic differences across growing locations and strong

biogeographic signal, the common bean rhizosphere recruited many similar taxa that could be

functionally important for the bean. Therefore, we explored occupancy-abundance distributions of

taxa ((39, 40) and references therein) to infer the core bean microbiome of taxa with an occupancy

of 1 (i.e., found in all samples and across all growing locations; **Fig. 2AB**). Among bacteria and

archaea, 258 phylogenetically diverse taxa were cosmopolitan in the dataset (**Fig. 2C**), including

numerous and abundant Proteobacteria (117 OTUs) with a dominant taxon classified as

Arthrobacter sp. (FM209319.1.1474, mean relative abundance of 1.43%). The bacterial/archaeal

core also contained taxa of interest for potential plant benefits (e.g. *Sphingomonas*, *Rhizobium*,

Bacillus, Streptomyces), as well as some genera that can be associated with disease (e.g. *Ralstonia*).

There were 13 taxa in the fungal core (**Fig. 2D**), and these were largely composed of Ascomycota

(10 OTUs), with dominating taxon OTU823 from the *Phaeosphaeriaceae* family (mean relative

abundance 10.1%). Notably, taxa that were specific to either bean genotype were relatively rare and

inconsistently detected (**Fig. 2**, orange and black points). Together, these results suggest that

common bean, regardless of the genotype, consistently recruits particular taxa.

The neutral expectation of abundance-occupancy distributions is that very abundant taxa

will have high occupancy, while rare taxa will have low (40–43). Taxa that deviate from the neutral

expectation are more strongly influenced by deterministic factors, like environmental conditions,

than by stochastic factors, like drift and dispersal. The neutral model fit (41, 42) of the abundance-

occupancy distribution (solid line, **Fig. 2AB**) identified several taxa that had frequencies either

above or below the 95% confidence intervals of the model (dashed lines). Specifically, 13.7% of

the bacterial/archaeal and 30.4% of fungal taxa, deviated from the neutral expectation (**SI**

Appendix, Dataset S3). One hundred and seventy-one core taxa were predicted above the neutral

model partition; these deterministically-selected taxa are prime candidates for follow-up studies of

their interactions with the host plant. Overall, the bacteria/archaea community had better fit to the

neutral expectation than fungal (R^2 of 0.74 and 0.34, and migration rates (m) of 0.301 and 0.003,

respectively), suggesting that dispersal was relatively less limiting for the bacteria/archaea than for

the fungi. This finding agrees with other work suggesting that fungi are more sensitive to local

climate or more dispersal limited than bacteria (44–47).

A cross-continental core rhizosphere microbiome for common bean

We wanted to better understand if these U.S. core taxa were associated with the bean

rhizosphere across a larger geographical scale, which would suggest the potential for selective plant

recruitment and cosmopolitan distribution of core taxa. Therefore, we compared our U.S. data to a

recently published study of rhizosphere bacteria and archaea from common beans grown in

Colombian soil (9). The Colombian study offered a key contrast because it included eight divergent

bean lineages, including wild (n=2), landrace (n=1), and cultivated genotypes (n=5), grown in soil

from a different continent that has starkly different climate and management from the U.S. growing

regions. To enable direct comparison, we re-analyzed raw reads and compared the datasets by

matching to either the same taxon identifiers when clustered to SILVA database, or 100% identity

by BLAST to *de novo* clustered reads (see Materials and Methods). Surprisingly, 39.6% (3,359

OTUs) of rhizosphere taxa from the Colombian-grown beans were also shared with the U.S. dataset

(**Fig. 3AB**). Both datasets included taxa that were highly represented in the other: 62% of U.S. core

(159 out of 258) were found also in Colombia, and 51% of Colombian core (433 out of 848) were

shared with the U.S. (**Fig. 3B**). Core taxa were again defined stringently with an occupancy of 1,

and 48 taxa were found across all samples, inclusive of both datasets. These core taxa were

composed of many Proteobacteria, with *Rhizobiales* showing the most consistent relative

abundance between the studies (**Fig. 3C,** e.g. 0.187% and 0.138% in Colombia and U.S. dataset,

respectively). None of the cross-continental core taxa were universally detected in very high

abundance, and all but two OTUs (a U.S. *Arthrobacter* sp. and a Colombian *Austrofundulus*

limnaeus with mean relative abundances of 1.43% and 1.01%, respectively) would be classified as

rare by a typical mean relative abundance threshold below 1%, hinting to a potential role of rare

taxa in providing key functions. Notably, only 48% of these cross-continental shared core taxa have

genus classification, suggesting that most of them are under-described in their functional potential

and interactions with plants.

We used a 97% sequence identity to conservatively define the core membership, but we

wanted to explore whether there was more resolved underlying taxonomic structure in the core

microbiome members. Therefore, we also analyzed reads associated with the global core taxa with

the UNOISE3 pipeline (48) to generate predicted biological sequences (zero-radius OTUs –

ZOTUs with100% sequence identity) and provide the maximal possible biological resolution. We

found that the global core taxa consisted of 422 ZOTUs, and that there was a range of 2 to 35

ZOTUs identified within each OTU (**SI Appendix, Fig. S4**). With one exception

(HQ597858.1.1508), all of core OTUs contained at least one ZOTUs that also had an occupancy of

1. In addition, all of the ZOTUs with an occupancy of 1 were also the most abundant ZOTU within

each OTU (**SI Appendix, Fig. S4**). This result is important because it suggests that the same

members constitute the core, even with increased taxonomic resolution. In summary, these results

show that common bean can associate with a core set of rhizosphere microbiome members at taxon

and ecotype levels, across diverse bean genotypes and across continents.

Inter-kingdom network analysis of the bean rhizosphere microbiome

Returning to the U.S. dataset that included both bacterial and fungal data, we wanted to

better understand the potential interactions of core taxa. We were particularly interested in fungal-

bacterial co-occurrences because of reports of their potential benefits for the plant (49–51). We

applied a co-occurrence network analysis to explore these patterns (52). We merged rarified 16S

and ITS datasets, filtered the datasets to include taxa with occupancy greater than or equal to 50%,

and considered only interactions significant at p-value<0.05 and LA score greater or equal to 0.88.

The resulting network included 572 taxa (nodes) and 1,857 statistically significant interactions

(edges) structured among 52 modules. Most of the modules were relatively small, with only six

including more than 10 nodes (**Fig. 4**). The network was scale-free and had small-world

characteristics as indicated by the node degree distribution fitting to the power law model ($R^2=$

0.993), and also had significant deviation of the modularity, length and clustering coefficients from

those calculated from random network (same number of nodes and edges), respectively (**SI**

Appendix, Dataset S4).

The topological role of each taxon within the network was determined by the relationship

between their within-module (Z_i) and among-module (P_i) connectivity scores (**Fig. 4A**) (53). Based

on this, the majority of taxa were peripheral (potentially, specialists; 563 nodes). There were also 3

connectors and 6 module hubs, but no network hubs. Indeed, supporting the conclusions of the

beta-diversity analysis (**Fig. 1**), there was a strong geographic signal in the largest four modules,

and these were comprised mostly of bacterial-bacterial (rather than bacterial-fungal or fungal-

fungal) hypothesized interactions (**Fig. 4B**). Interestingly, cross-kingdom interactions constituted

only a small fraction of all co-occurrences (n=168; bacteria-fungi=156, archaea-bacteria=6 and

archaea-fungi=6).

The analysis identified 26 co-occurrences between core taxa. However, we were surprised

to find that only two bacterial core and no fungal core taxa were also classified as network hubs

(taxa that connect to many other taxa within a modules) or connectors (taxa that connect across

modules). As exceptions, core *Chitinophagaceae* taxon FR749720.1.1492 was a module hub node

and a *Nitrobacter* sp. GDHX01215817.4.1477 was a connector. However, the lack of overlap

between core and hub/connector taxa is important because hubs and connectors are often

hypothesized to play important roles for the plant (50, 54, 55). Our results, inclusive of a dataset of

divergent plant genotypes and broad biogeography, suggest that while hub and connector taxa may

be important for the maintenance of the root microbiome, these taxa are not consistently detected in

the common bean rhizosphere and, by deduction, could not be of universal importance for the host

plant. Our study cannot speak to the potential for functional redundancy among hub or connector

taxa, which could ultimately suggest a functional core among phylogenetically diverse taxa (56).

However, it is expected that more resolved taxonomy also will reveal functional redundancy at the

sub-OTU level among core members. Taken together, these results suggest that core taxa likely are

important for the plant, while hub and connector taxa are important for the integrity of the soil

microbial community and its responses to the local environment.

Signatures of fertilizer on the rhizosphere microbiome

As a legume, common bean can form symbiotic relationships with nitrogen-fixing bacteria (mainly,

Rhizobium sp.). However, as compared to other legumes of agricultural importance (e.g., soy,

chickpeas), common bean is inefficient in that a relatively lower percentage of its total nitrogen can

be attributed to that fixed by symbionts (57). As a consequence, many U.S. growers provide

nitrogen fertilizer to common bean crops. Not only is nitrogen fertilizer an expense to the grower,

but its use also contributes to global emissions of nitrous oxide, a powerful greenhouse gas (58). It

is of interest to decrease use of nitrogen fertilizer in agriculture, but especially in legumes that are

capable of acquiring nitrogen through their symbiotic relationships with bacteria.

Our study included research farms that used different fertilizer types, including

conventional synthetic fertilizer, organic application, and no added fertilizer. Fertilizer type

explained 35% variance of the differences in the microbiome communities across growing

locations (**SI Appendix, Dataset S1**). Additionally, changes in bacterial, archaeal and fungal

communities were more strongly correlated with nitrate concentrations than with total nitrogen or

ammonium concentrations (**SI Appendix, Dataset S1**). Therefore, we wanted to explore the U.S.

dataset for rhizosphere taxa that were indicative of fertilizer type, which would be suggestive of

conditional plant preference or competitive advantage and could ultimately provide hints as to

which taxa are most sensitive to changes in nitrogen management. We assessed differentially

abundant taxa between each fertilizer pair (p-value < 0.001; **SI Appendix, Figure S5**), and also

applied indicator species analysis to identify taxa that were characteristic of particular fertilizer

types (**Fig. 5**) (59). In summary, synthetic fertilizer enriched for 198 taxa, organic for 95 taxa and

no fertilization for 66 taxa (**Fig. 5**). Organic fertilization proportionally enriched for taxa belonging

to the Gammaproteobacteria relative to other treatments, whereas synthetic fertilization was

associated with their depletion, in agreement with previous studies (60, 61). In contrast,

Alphaproteobacteria were enriched in soils without fertilization, but were detected only in

relatively low abundance (**Fig. 5**). Interestingly, some taxa with potential to fix atmospheric

nitrogen (i.e. *Rhizobium*, *Bradyrhizobium*, *Burkholderia-Paraburkolderia*) were more abundant in

the fertilized sites (**Fig. 5**). This is surprising considering the importance of these lineages for

plants in unfertilized soils. *Rhizobium* sp., which have specific symbiotic interactions with common

bean (62, 63), were well represented in all growing locations and not statistically affected by

fertilization type or plant genotype, but instead by soil conditions and management (ANOVA, p-

value=0.019) (**Fig. 5**). Anecdotally, we observed substantial nodulation only on plant root sampled

at Saginaw Valley, MI, the only location where irrigation was not available (**Table 1**).

A caveat to these analyses is that there was not cross-farm replication of each fertilizer type,

and so these results should be considered a first observation towards understanding the selective

consequences of fertilizer type on the bean rhizosphere microbiome. Broadly, these results provide

first insights as to taxa that could be targeted towards microbiome management to decrease reliance

on nitrogen fertilizer.

Prospectus

Legumes like common bean that form symbiotic relationships with nitrogen fixing bacteria are a

crucial component of sustainable agriculture. Growing common bean can both improve soil fertility

(62, 64) and contribute to food security, and its production has reached more than 31 million metric

tons worldwide (http://www.fao.org/faostat/, accessing on July 12, 2019)). However, common bean

yield estimates, production areas, and even nutritional value, are predicted to decrease in this

century due to the impacts of global change (65–67). Thus, breeding or other improvements to

common bean is needed to make this important crop resistant to stressors associated with climate

change while increasing yields to meet higher demands. To date, only traditional plant breeding

approaches have been used for the common bean, exploiting the genetic diversity of the plant, but

the latest knowledge of the plant microbiome in supporting host well-being has opened new

avenues for breeding the next generation of crops with plant microbiome manipulation (68).

Here, we show that common bean genotype has marginal effect on the rhizosphere

microbiome grown in field conditions. We describe for the first time the common bean core

microbiome at continental and cross-continental scales. A majority of the 48 bacterial taxa

cosmopolitan in the common bean rhizosphere are under-described or largely unknown, as only 23

out of 48 have genus classification. But, due to their ubiquitous association with the bean, we

hypothesize that these taxa provide functions that are crucial for common bean health. A next step

is to understand functions associated with these taxa and to determine if and how they contribute to

plant health and productivity under different growth conditions, such as drought or with particular

management strategies, as well as over plant development (32, 69). Additionally, targeting

persistent microbiota that are part of a native assemblage for management and manipulation can be

advantageous, as these are expected to be both host and environment adapted and thus may be

more persistent *in situ* than external or non-native biocontrols (70). In conclusion, this work

provides insights and approaches for prioritizing plant-microbiome members that will advance

goals in microbiome management and microbe-improved crops.

Material and Methods

Sampling and soil physicochemical analysis

We selected two common bean cultivars with very distinct genotypes: Eclipse (24) with

Mesoamerican origin and California Early Light Red Kidney (CELRK) and old kidney bean

landrace with Andean origin (25). The plants were grown at 5 different locations on research

extension farms: Saginaw Valley, Montcalm county (both in Michigan), Scott Bluff county

(Nebraska), Fort Collins (Colorado) and Othello (Washington), each representing an important

U.S. common bean growing region (26). Both bean genotypes were grown in the same locations

except in Michigan, where CELRK was grown at Saginaw Valley and Eclipse at Montcalm county.

Plants (n≥3) were harvested at flowering from replicated plots (n≥3) at each location. The

flowering time point was selected because we wanted to analyze mature microbial communities as

influenced by the plant and not by plant development. Plant roots with attached soil were packed

into bags, stored on ice and shipped immediately to Michigan State University for processing.

From each sample, root attached soil was collected and used as rhizosphere soil and

remaining soil was collected and used as bulk soil. Both soils were sieved through 4 mm sieves to

remove any organic, plant debris and larger soil minerals before soil analysis and DNA isolation.

Soil analysis was done at the Michigan State Soil and Plant Nutrient Laboratory and

included following parameters: pH, phosphorous (P ppm), potassium (K ppm), calcium (Ca ppm),

magnesium (Mg ppm), organic matter content (OM%), nitrate (NO_3^- ppm), ammonium (NH_4^+ ppm)

and total nitrogen (%). Soil analysis was done on bulk soil samples pooled by plant genotype and

plot.

DNA isolation and sequencing

DNA was isolated using DNeasy PowerSoil kit (QIAGEN, US) by following the

manufacturer's recommendations. Quality and quantity of isolated DNA was checked with Qubit

2.0 fluorometer using HS dsDNA Assay kit (ThermoFisher, US). Presence of 16S rRNA gene was

confirmed by PCR using the V4 16S rRNA specific sequencing primer set 515f and 806r (71) and

then visualizing the PCR products by gel electrophoresis. Because samples from Nebraska were

sent in bags by plot and genotype, we pooled all other samples by plot and did not sequence

separately by plant. All 16S rRNA samples were prepared by the Michigan State Genomics Core

Research Support Facility. Their standard protocol included 16S rRNA gene PCR amplification

and library preparation. The ITS samples were amplified in our lab using the primer pair ITS1f and

ITS2 (72) with index adapters as instructed by the Genomics Core

(https://rtsf.natsci.msu.edu/genomics/sample-requirements/illumina-sequencing-sample-

requirements/, June 2019), The Genomics Core used the Illumina TruSeq Nano DNA library

preparation kit for both 16S and ITS libraries. Paired-end, 250-bp reads were generated on an

Illumina MiSeq platform using v2 Standard 500 cycle kit, and the Genomics Core provided

standard Illumina quality control, adaptor, barcode trimming, and sample demultiplexing.

16S rRNA and ITS amplicon processing

For both datasets, raw reads were merged, quality filtered, dereplicated, and clustered into

97% identity operational taxonomic units (OTUs) using the UPARSE pipeline version 11 (73).

Reads not matching to the SILVA database were used for de-novo clustering at 97% identity.

Reference picked and de-novo reads were combined before taxonomy was assigned. Taxonomic

annotations for 16S rRNA OTU representative sequences were assigned in the QIIME 1.19

environment (74) using SILVA rRNA database release v128 (75). ITS OTU representative

sequences were taxonomically annotated using the CONSTAX tool (76) with the UNITE database

(version 7.2) (77). OTUs with unassigned taxonomy at the domain level and OTUs annotated as

mitochondria or chloroplasts were removed.

Statistical analysis

Alpha and beta diversity analyses were performed to datasets subsampled to the minimum

observed quality filtered reads per sample (31,255 for 16S rRNA and 22,716 for ITS). Due to low

read counts one sample (SVERC1) was removed from the 16S rRNA dataset. We report richness as

total number of OTUs clustered at 97% sequence identity. Differences among groups (plant

genotype, growing location, fertilization, pH) were assessed using analysis of variance (ANOVA),

with a Tukey *post hoc* test for multiple comparisons. We used the *protest* function in the vegan

package in R (78) to test for synchrony between bacterial/archaeal and fungal communities.

To quantify the variation in community composition, we calculated pairwise Bray-Curtis

distances. Permutational multivariate analysis (PERMANOVA) using 1,000 random permutations

was used to test hypothesis of beta diversity using *adonis* function in the vegan package in R (78).

Differential abundance and indicator species analysis

We used DESeq2 (31) to determine bacterial and archaeal taxa with differential abundance

across plant genotypes and fertilization types. Thresholds for calling taxa as differentially abundant

between selected groups was set at p-value of 0.05 and 0.001 or lower for plant genotype and

fertilization types, respectively. Taxa that were differentially abundant between fertilization types

were then used for indicator species analysis using *labdsv* package in R (79).

Defining continental and cross-continental core rhizosphere microbiota of common bean

For cross-continental core, we first downloaded and processed raw reads from a study that

investigated the influence of common bean genotype on the rhizosphere microbiome composition

and included many Mesoamerican genotypes but used only one soil type, greenhouse conditions for

growth of the plants, and focused only on the bacterial and archaeal communities (9) (NCBI

BioProject ID PRJEB19467). We processed the raw sequences as described above, and despite

differences in our pipeline and the originally published pipeline, we generated very similar amount

of OTUs (12,209 instead of 12,293 (9)). Because the studies used different sequencing primers, we

identified the identical OTUs by matching the IDs as given by the SILVA database v128 (75) for

all reference clustered OTUs. For the *de-novo* clustered OTUs, we used BLAST (80) to identify

100% matches between the two datasets, across the overlapping V4 region of reads from both

studies. As before, we calculated occupancy of every taxon in the rhizosphere datasets and

identified those with occupancy of 1. Taxa with the same OTU IDs or 100% BLAST match were

designated as the cross-continental core rhizosphere microbiota of the common bean.

Neutral model and distance-decay analysis

To assess the importance of neutral process in the assembly of common bean rhizosphere

communities we applied the Sloan neutral model (81). This model predicts the relationship between

the frequency with which taxa occur in a set of local communities (occupancy) and their mean

abundance across a broader metacommunity. The prediction is that rare taxa will be lost from

individual hosts due to ecological drift, but abundant taxa will be more widespread in a

metacommunity due to higher chance of dispersal and thus be randomly sampled by individual

hosts.

The occupancy of OTUs and their mean relative abundances across the metacommunity

were fitted to the model, using the R code described by Burns et al. (42). We used the following

parameters of neutral model: 95% confidence intervals, the goodness of fit of the neutral model

(R^2), and the estimated migration rate (m).

Co-occurrence network analysis

Global network properties were calculated using the Molecular Ecological Network

Analysis Pipeline (MENAP) (52). The majority was set to 0.5, missing data were kept blank, read

counts were converted by the logarithm, and the Pearson correlation coefficient was used as the

similarity measure. The results were then filtered by keeping pairwise correlations with absolute

LA values greater or equal to 0.88. To determine the modularity of the network we created 100

random networks within MENAP using the same number of nodes and edges as the complete

potentially active network. The general properties of the inferred networks were analyzed using

NetworkAnalyzer in Cytoscape (82, 83).

The connectivity of each node in the network was calculated using within-module

connectivity (Zi) and among-module connectivity (Pi) scores which defined the topological role of

each node (taxon) (52, 53). We classified our nodes as per the four categories describing node

topology: *network hubs* (highly connected nodes within the entire network, Zi > 2.5 and Pi > 0.62),

module hubs (highly connected nodes within modules, Zi > 2.5), *connectors* (nodes that connect

modules, Pi > 0.62), and *peripheral* nodes (nodes connected in modules with few outside

connections, Zi < 2.5 and Pi < 0.62) (53).

Cytoscape v.3.5.1 (82) was used for visualization of significant co-occurrences and editing

the appearance of nodes size, shape and color based on the number of connections (degrees),

taxonomic affiliation and module, respectively.

Acknowledgments

This work was supported by the Plant Resilience Institute at Michigan State University. We

are grateful to Dr. James Kelly, Dr. Phillip Miklas, Dr. James J. Heitholt, Dr. Carlos Urrea Florez,

Dr. Mark A. Brick, Dr. Juan M. Osorno, Dr. Thomas H Smith and other supporting staff at the

research extension farms for their partnership in this study. We would also like to thank the

Michigan State Genomics Core Research and Technology Support Facility and the Institute for

Cyber-Enabled Research High Performance Computing Center for excellent support and service.

Footnotes

Author contributions: N.S. and A.S. designed research, analyzed data and wrote the paper; and N.S.

performed research.

The authors declare no conflict of interest.

Data deposition: The raw sequence data have been deposited in the NCBI Sequence Read Archive

(BioProject accession no. PRJNA524532). All read processing steps, bioinformatic workflows,

and custom scripts are available on GitHub

(https://github.com/ShadeLab/PAPER_Stopnisek_2019_BeanBiogeography).

References

1.   Dessaux Y, Grandclément C, Faure D (2016) Engineering the Rhizosphere. *Trends Plant Sci*

21(3):266–278.

2.   Gopal M, Gupta A (2016) Microbiome Selection Could Spur Next-Generation Plant

Breeding Strategies. *Front Microbiol* 7:1971.

3.   Compant S, Samad A, Faist H, Sessitsch A (2019) A review on the plant microbiome:

Ecology, functions, and emerging trends in microbial application. *J Adv Res*.

doi:10.1016/J.JARE.2019.03.004.

4.   Berg G, Grube M, Schloter M, Smalla K (2014) The plant microbiome and its importance

for plant and human health. *Front Microbiol* 5:1.

5.   Wille L, Messmer MM, Studer B, Hohmann P (2019) Insights to plant-microbe interactions

provide opportunities to improve resistance breeding against root diseases in grain legumes.

Plant Cell Environ 42(1):20–40.

6.   Busby PE, et al. (2017) Research priorities for harnessing plant microbiomes in sustainable

agriculture. *PLOS Biol* 15(3):e2001793.

7.   Kroll S, Agler MT, Kemen E (2017) Genomic dissection of host–microbe and microbe–

microbe interactions for advanced plant breeding. *Curr Opin Plant Biol* 36:71–78.

8.   Peiffer JA, et al. (2013) Diversity and heritability of the maize rhizosphere microbiome

under field conditions. *Proc Natl Acad Sci U S A* 110(16):6548–6553.

9.   Perez-Jaramillo JE, et al. (2017) Linking rhizosphere microbiome composition of wild and

domesticated Phaseolus vulgaris to genotypic and root phenotypic traits. *ISME J*.

doi:10.1038/ismej.2017.85.

10.   Lundberg DS, et al. (2012) Defining the core Arabidopsis thaliana root microbiome. *Nature*

488. doi:10.1038/nature11237.

11.   Xu J, et al. (2018) The structure and function of the global citrus rhizosphere microbiome.

493   *Nat Commun* 9(1):4894.

494   12.   Walters WA, et al. (2018) Large-scale replicated field study of maize rhizosphere identifies

495   heritable microbes. *Proc Natl Acad Sci U S A* 115(28):7368–7373.

496   13.   Edwards J, et al. (2015) Structure, variation, and assembly of the root-associated

497   microbiomes of rice. *Proc Natl Acad Sci U S A* 112(8):E911-20.

498   14.   Wagner MR, et al. (2016) Host genotype and age shape the leaf and root microbiomes of a

499   wild perennial plant. *Nat Commun* 7. doi:10.1038/ncomms12151.

500   15.   Gomes EA, De Sousa SM, Oliveira CA (2018) Root-Associated Microbiome of Maize

501   Genotypes with Contrasting Phosphorus Use Efficiency. 2(3):129.

502   16.   Broughton WJ, et al. (2003) Beans (Phaseolus spp.) – model food legumes. *Plant Soil*

503   252(1):55–128.

504   17.   Schmutz J, et al. (2014) A reference genome for common bean and genome-wide analysis of

505   dual domestications. *Nat Genet* 46(7):707–713.

506   18.   Rendón-Anaya M, et al. (2017) Genomic history of the origin and domestication of common

507   bean unveils its closest sister species. *Genome Biol* 18(1):60.

508   19.   Bitocchi E, et al. (2017) Beans (Phaseolus ssp.) as a Model for Understanding Crop

509   Evolution. *Front Plant Sci* 8:722.

510   20.   Bitocchi E, et al. (2012) Mesoamerican origin of the common bean (Phaseolus vulgaris L.) is

511   revealed by sequence data. *Proc Natl Acad Sci* 109(14):788–796.

512   21.   Kwak M, Gepts P (2009) Structure of genetic diversity in the two major gene pools of

513   common bean (Phaseolus vulgaris L., Fabaceae). *Theor Appl Genet* 118(5):979–992.

514   22.   Bitocchi E, et al. (2013) Molecular analysis of the parallel domestication of the common

515   bean (Phaseolus vulgaris) in Mesoamerica and the Andes. *New Phytol* 197(1):300–313.

23.   Schmutz J, et al. (2014) A reference genome for common bean and genome-wide analysis of

dual domestications. *Nat Publ Gr* 46(7). doi:10.1038/ng.3008.

24.   NDSU Agriculture (2004) *Eclipse Black Bean* (Fargo) Available at:

http://ndsuresearch.eclipticcms.com/files/pdf/Ag Brochures/Eclipse_Black_brochure.pdf

[Accessed June 3, 2019].

25.   McClean P, Myers J (1990) Pedigrees of dry bean cultivars, lines and PIs. *Annu Rep Bean*

Improv Coop 33:23–25.

26.   USDA (2017) *United States Department of Agriculture National Agricultural Statistics*

Service Crop Production 2016 Summary Available at:

https://www.agmrc.org/media/cms/CropProdSu01122017_C27B81FB8F64D.pdf [Accessed

May 10, 2019].

27.   Singer E, Bonnette J, Kenaley SC, Woyke T, Juenger TE (2019) Plant compartment and

genetic variation drive microbiome composition in switchgrass roots. *Environ Microbiol Rep*

11(2):185–195.

28.   Coleman-Derr D, et al. (2016) Plant compartment and biogeography affect microbiome

composition in cultivated and native *Agave* species. *New Phytol* 209(2):798–811.

29.   De Souza RSC, et al. (2016) Unlocking the bacterial and fungal communities assemblages of

sugarcane microbiome. *Sci Rep* 6(June):1–15.

30.   Walters WA, et al. (2018) Large-scale replicated field study of maize rhizosphere identifies

heritable microbes. *Proc Natl Acad Sci* 115(28):7368–7373.

31.   Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and dispersion

for RNA-seq data with DESeq2. *Genome Biol* 15(12):550.

32.   Edwards JA, et al. (2018) Compositional shifts in root-associated bacterial and archaeal

microbiota track the plant life cycle in field-grown rice. *PLOS Biol* 16(2):e2003862.

33.   Rich SM, Watt M (2013) Soil conditions and cereal root system architecture: Review and

considerations for linking Darwin and Weaver. *J Exp Bot*. doi:10.1093/jxb/ert043.

34.   Fitzpatrick CR, et al. (2018) Assembly and ecological function of the root microbiome

across angiosperm plant species. *Proc Natl Acad Sci U S A* 115(6):E1157–E1165.

35.   Hamonts K, et al. (2018) Field study reveals core plant microbiota and relative importance

of their drivers. *Environ Microbiol* 20(1):124–140.

36.   Estendorfer J, et al. (2017) The Influence of Land Use Intensity on the Plant-Associated

Microbiome of Dactylis glomerata L. *Front Plant Sci* 8:930.

37.   Leff JW, Lynch RC, Kane NC, Fierer N (2017) Plant domestication and the assembly of

bacterial and fungal communities associated with strains of the common sunflower,

Helianthus annuus. *New Phytol* 214(1):412–423.

38.   Levy A, et al. (2018) Genomic features of bacterial adaptation to plants. *Nat Genet*

50(1):138–150.

39.   Grady KL, Sorensen JW, Stopnisek N, Guittar J, Shade A (2019) Assembly and seasonality

of core phyllosphere microbiota on perennial biofuel crops. *bioRxiv*:446369.

40.   Shade A, et al. (2018) Macroecology to Unite All Life, Large and Small. *Trends Ecol Evol*

33(10):731–744.

41.   Woodcock S, et al. (2007) Neutral assembly of bacterial communities. *FEMS Microbiol*

Ecol 62(2):171–180.

42.   Burns AR, et al. (2016) Contribution of neutral processes to the assembly of gut microbial

communities in the zebrafish over host development. *ISME J* 10(3):655–664.

43.   Gaston KJ, et al. (2000) Abundance-occupancy relationships. *J Appl Ecol* 37(s1):39–59.

44.   Martiny JBH, Eisen JA, Penn K, Allison SD, Horner-Devine MC (2011) Drivers of bacterial

beta-diversity depend on spatial scale. *Proc Natl Acad Sci U S A* 108(19):7850–4.

45.   Gumiere T, Durrer A, Bohannan BJM, Andreote FD (2016) Biogeographical patterns in

565   fungal communities from soils cultivated with sugarcane. *J Biogeogr* 43(10):2016–2026.

566   46.   Talbot JM, et al. (2014) Endemism and functional convergence across the North American

567   soil mycobiome. *Proc Natl Acad Sci* 111(17):6341–6346.

568   47.   Tedersoo L, et al. (2014) Global diversity and geography of soil fungi. *Science (80-)*

569   346(6213). doi:10.1126/science.1256688.

570   48.   Edgar RC (2016) UNOISE2: improved error-correction for Illumina 16S and ITS amplicon

571   sequencing. *bioRxiv*:081257.

572   49.   Jambon I, Thijs S, Weyens N, Vangronsveld J (2018) Harnessing plant-bacteria-fungi

573   interactions to improve plant growth and degradation of organic pollutants. *J Plant Interact*

574   13(1):119–130.

575   50.   Agler MT, et al. (2016) Microbial Hub Taxa Link Host and Abiotic Factors to Plant

576   Microbiome Variation. *PLOS Biol* 14(1):e1002352.

577   51.   Durá P, et al. (2018) Microbial Interkingdom Interactions in Roots Promote Arabidopsis

578   Survival. *Cell* 175:973–983.

579   52.   Deng Y, et al. (2012) Molecular ecological network analyses. *BMC Bioinformatics*

580   13(1):113.

581   53.   Guimerà R, Nunes Amaral LA (2005) Functional cartography of complex metabolic

582   networks. *Nature* 433(7028):895–900.

583   54.   Trivedi P, et al. (2017) Keystone microbial taxa regulate the invasion of a fungal pathogen in

584   agro-ecosystems. *Soil Biol Biochem* 111:10–14.

585   55.   Shi S, et al. (2016) The interconnected rhizosphere: High network complexity dominates

586   rhizosphere assemblages. *Ecol Lett* 19(8). doi:10.1111/ele.12630.

56.   Lemanceau P, Blouin M, Muller D, Moënne-Loccoz Y (2017) Let the Core Microbiota Be

Functional. *Trends Plant Sci*. doi:10.1016/j.tplants.2017.04.008.

57.   Hardarson G, Atkins C (2003) Optimising biological N 2 fixation by legumes in farming

systems. *Plant Soil* 252:41–54.

58.   Shcherbak I, Millar N, Robertson GP (2014) Global metaanalysis of the nonlinear response

of soil nitrous oxide (N2O) emissions to fertilizer nitrogen. *Proc Natl Acad Sci U S A*

111(25):9199–204.

59.   Dufrêne M, Legendre P (1997) Species assemblages and indicator species: the need for a

flexible asymmetrical approach. *Ecol Monogr* 67(3):345–366.

60.   Ma M, et al. (2018) Effect of long-term fertilization strategies on bacterial community

composition in a 35-year field experiment of Chinese Mollisols. *AMB Express* 8(1):20.

61.   Herzog S, Wemheuer F, Wemheuer B, Daniel R (2015) Effects of Fertilization and

Sampling Time on Composition and Diversity of Entire and Active Bacterial Communities

in German Grassland Soils. *PLoS One* 10(12):e0145575.

62.   Aguilar OM, Riva O, Peltzer E (2004) Analysis of Rhizobium etli and of its symbiosis with

wild Phaseolus vulgaris supports coevolution in centers of host diversification. *PNAS*

101(37):13548–13553.

63.   Tong W, et al. (2018) Genomic insight into the taxonomy of Rhizobium genospecies that

nodulate Phaseolus vulgaris. *Syst Appl Microbiol* 41(4):300–310.

64.   Martínez-Romero E (2003) Diversity of Rhizobium-Phaseolus vulgaris symbiosis: overview

and perspectives. *Plant Soil* 252(1):11–23.

65.   Jaggard KW, Qi A, Ober ES (2010) Possible changes to arable crop yields by 2050. *Philos*

Trans R Soc Lond B Biol Sci 365(1554):2835–51.

66.   Beebe S, Rao I, Mukankusi C, Buruchara R (2012) Improving Resource Use Efficiency and

Reducing Risk of Common Bean Production in Africa, Latin America, and the Caribbean.

CIAT Book Chapter, ed Hershey CH (Centro Internacional de Agricultura Tropical (CIAT),

Cali), pp 1–18.

67.   Hummel M, et al. (2018) Reduction in nutritional quality and growing area suitability of

common bean under climate change induced drought stress in Africa. *Sci Rep* 8(1):16187.

68.   Wallenstein MD (2017) Managing and manipulating the rhizosphere microbiome for plant

health: A systems approach. *Rhizosphere*. doi:10.1016/j.rhisph.2017.04.004.

69.   Xu L, et al. (2018) Drought delays development of the sorghum root microbiome and

enriches for monoderm bacteria. *Proc Natl Acad Sci* 115(18):E4284–E4293.

70.   Bell TH, et al. (2019) Manipulating Wild and Tamed Phytobiomes: Challenges and

Opportunities. *Phytobiomes J*:PBIOMES-01-19-0006-W.

71.   Caporaso JG, et al. (2011) Global patterns of 16S rRNA diversity at a depth of millions of

sequences per sample. *Proc Natl Acad Sci U S A* 108 Suppl 1(Supplement 1):4516–22.

72.   White T, et al. (1990) Amplification and direct sequencing of fungal ribosomal RNA genes

for phylogenetics. *PCR - Protocols and Applications - A Laboratory Manual*, eds Innis M,

Gelfand D, Sninsky J, White T (Academic Press), pp 315–322. 1st Ed.

73.   Edgar RC (2013) UPARSE: highly accurate OTU sequences from microbial amplicon reads.

Nat Methods 10(10):996–998.

74.   Caporaso JG, et al. (2010) QIIME allows analysis of high-throughput community

sequencing data. *Nat Methods* 7(5):335–336.

75.   Quast C, et al. (2013) The SILVA ribosomal RNA gene database project: improved data

processing and web-based tools. *Nucleic Acids Res* 41(Database issue):D590-6.

76.   Gdanetz K, Benucci GMN, Vande Pol N, Bonito G (2017) CONSTAX: a tool for improved

taxonomic resolution of environmental fungal ITS sequences. *BMC Bioinformatics*

18(1):538.

77.   UNITE Community (2017) UNITE USEARCH/UTAX release. Available at:

https://doi.org/10.15156/BIO/587476.

78.   Oksanen J, et al. (2019) vegan: Community Ecology Package. Available at: https://cran.r-

project.org/package=vegan.

79.   Roberts DW (2016) labdsv: Ordination and Multivariate Analysis for Ecology.

80.   Madden T (2013) The BLAST Sequence Analysis Tool. Available at: https://www-ncbi-

nlm-nih-gov.proxy1.cl.msu.edu/books/NBK153387/?report=reader [Accessed June 3, 2019].

81.   Sloan WT, et al. (2006) Quantifying the roles of immigration and chance in shaping

prokaryote community structure. *Environ Microbiol* 8(4):732–740.

82.   Shannon P, et al. (2003) Cytoscape: A Software Environment for Integrated Models of

Biomolecular Interaction Networks. *Genome Res* 13(11):2498–2504.

83.   Doncheva NT, Assenov Y, Domingues FS, Albrecht M (2012) Topological analysis and

interactive visualization of biological networks and protein structures. *Nat Protoc* 7(4):670–

685.

84.   De Cáceres M, Legendre P (2009) Associations between species and groups of sites: Indices

and statistical inference. *Ecology* 90(12):3566–3574.

Figure Legends

Fig. 1: Growing location drives bacterial/archaeal (A) and fungal (B) microbiome structure of the

common bean rhizosphere. The principal coordinate analysis (PCoA) is based on Bray-Curtis

distances. Growing location is indicated by color and plant genotype is indicated by shape shapes

(diamond=CELRK, circle= Eclipse, square=bulk soil). The strength of statistically significant (p-

$value$ < 0.01) explanatory variables are shown as the length of fitted vectors.

Fig. 2: Abundance-occupancy distributions were used to identify core members of the rhizosphere

microbiome for bacteria/archaea (A) and fungi (B). Taxa exclusive to a genotype are indicated in

orange (CELRK) or black (Eclipse), and taxa shared across both genotypes are white. The solid

line represents the fit of the neutral model, and the dashed line is 95% confidence around the model

prediction. Taxa with an occupancy of 1 (i.e., detected in all samples) were considered members of

the core. Relative abundance of these taxa is represented as boxplots, grouped by order and number

of taxa therein (C, D). Panels C and D are color-coded by phyla.

Fig. 3: A cross-continental core rhizosphere microbiome. There were 3361 shared

bacterial/archaeal taxa across U.S. and Colombia rhizosphere samples, suggesting highly similar

recruitment across continental scales (A). Forty-eight taxa were detected in all samples of both

datasets, as depicted by the Venn diagram (B), and included many Proteobacteria (C). Relative

abundance of the 48 U.S.-Colombia core taxa is represented as boxplots (left panel), grouped by

order and dataset (Colombia/U.S.). Number of taxa per order is represented as bars (right panel).

Boxplots and bars are color-coded by phylum.

Fig. 4: Network co-occurrence analysis shows that core rhizosphere microbiome members are

predominantly classified as peripheral taxa that are weakly connected, and generally clustering by

the growing location. The network depicts only clusters of modules that were connected by more

than 6 nodes (A). Nodes shape is representing domain associations (archaeal, bacterial or fungal)

and node size is proportional to its total number of connecting edges (A). The four largest modules,

are generally reflective of community biogeography and distinguished by color (A). The within-

(Zi) and among- (Pi) module connectivity plot was used to identify module (Pi<0.62, Zi>0.2.5) or

network hub taxa (Pi>0.62, Zi>2.5), as well as connector (Pi>0.62, Zi<2.5) and peripheral taxa (B).

The density plots surrounding the Zi-Pi plot represent core (green) and non-core taxa (black) (B).

The relative abundances of taxa within each module is represented in box plots (C).

Fig. 5: Effects of fertilization on the rhizosphere microbiome of the common bean. Proteobacterial

taxa with significant deviation in relative abundance between fertilization treatments were

determined by the indicator species analysis (*indval* (84)). Ninety-five taxa were indicators of

organic fertilizer (manure, urea), 66 were indicators of no fertilizer, and 198 of synthetic fertilizer

(A). Presence-absence of indicator taxa is represented in the panel B, with symbol size indicating

their numbers if present (B) and their relative abundance in panel C. Number of indicator taxa per

taxonomic affiliation is reported in the brackets (C).

Table 1: Description of geographical, managemental and soil properties of the common bean growing locations included in the study.

Sample same	Growing location	Elevation	Bean genotype	Weather (T, precipitation)*	Rotation history	Fertilization (per acre)	Irrigation	pH	Nitrogen (%)	Organic matter (%)
SVREC	Saginaw Valley Research and Extension Center, MI, US	190 m	CELRK	24.5ºC, 5.7 mm	Common bean, wheat, maize,	Synthetic (400lbs of 15N-5 P-13K)	None	7.7 (± 0.1)	0.13 (± 0.00)	2.33 (± 0.06)
MRC	Montcalm Research Center, MI, US	280 m	Eclipse	24ºC, 7.4 mm	Common bean, maize, potato	Synthetic (200lbs of 19N-0P-19K)	Yes	5.9 (± 0.4)	0.10 (± 0.01)	2.13 (± 0.06)
NE	Scottsbluff, NE, US	1200 m	CELRK and Eclipse	27ºC, 2.18 mm	Common bean, maize, common bean	None	Yes	7.9 (± 0.1)	0.07 (± 0.01)	1.39 (± 0.18)
CO	ARDEC, Fort Collins, CO, US	1536 m	CELRK and Eclipse	31ºC, 1.4 mm **	Common bean, maize, barley	Organic (220lbs urea (2016), 3 tons manure (2015))	Yes	8.1 (± 0.2)	0.11 (± 0.00)	1.7 (± 0.15)
WA	WSU, Othello, WA, US	320 m	CELRK and Eclipse	26.3ºC, 0.2 mm	Common bean, maize, wheat	Synthetic (40lbs N/20lbs P/10lbs ZN2O5)	Yes	6.1 (± 0.4)	0.08 (± 0.01)	1.78 (± 0.10)

* Temperature (as max air temperature) and precipitation are averaged for the period between May 1st and July 31st 2017.

** Data available only for the period between Jun 28th and Jul 31st 2017.

Fig.1

Fig. 2

Fig. 3

Fig. 4

Fig. 5