Discovery of a spatially and temporally persistent core microbiome of the common bean

rhizosphere

Nejc Stopnisek[1,2,3] and Ashley Shade[1,2,3,4,5]

1. Department of Microbiology and Molecular Genetics, Michigan State University, East

Lansing, MI 48840 USA

2. The Plant Resilience Institute, Michigan State University, East Lansing, MI 48840

3. The DOE Great Lakes Bioenergy Research Center, Michigan State University, East

Lansing, MI 48840

4. Program in Ecology, Evolutionary Biology and Behavior, Michigan State University, East

Lansing, MI 48840

5. Department of Plant, Soil and Microbial Sciences, Michigan State University, East Lansing,

MI 48840

Corresponding author: shadeash@msu.edu

Keywords: plant-microbiome, Phaseolus vulgaris, plant health, neutral model, co-occurrence

network, rhizobia, fertilizer, abundance-occupancy, 16S rRNA, ITS, microbial ecology, soil, hub

taxa, connector taxa, plant genotype, sustainable agriculture

Abstract

Plants recruit soil microbes that provide nutrients, promote growth and protect against

pathogens[1-3]. However, the full potential of microbial communities for supporting plant health and

agriculture is unrealized [4-6], in part because rhizosphere members key for plant health are difficult

to prioritize[7]. Microbes that ubiquitously associate with a plant species across large spatial scales

and varied soil conditions provide a practical starting point for discovering beneficial members[7].

Here, we quantified the structures of bacterial/archaeal and fungal communities in the common

bean rhizosphere (*Phaseolus vulgaris*), and assessed its core membership across space and time. To

assess a spatial core, two divergent bean genotypes were grown in field conditions across five

major growing regions in the United States, and then also compared to eight genotypes grown in

Colombian soil. To assess a temporal core, we conducted a time course of rhizosphere and

rhizoplane microbiome members over bean development in the field. Surprisingly, there were 48

persistent bacterial taxa that were detected in all samples, inclusive of U.S. and Colombian-grown

beans and over plant development, suggesting cosmopolitan enrichment and time-independence.

Neutral models of abundance-occupancy relationships and co-occurrence networks show that many

of these core taxa are deterministically selected and likely in intimate relationships with the plant.

Many of the core taxa were yet-uncultured and affiliated with Proteobacteria; these taxa are prime

targets in support of translational plant-microbiome management. More generally, this work

reveals that core members of the plant microbiome can have both broad ranges and temporal

persistence with their host, suggesting intimate, albeit possibly opportunistic, interactions.

Main

Common bean (*Phaseolus vulgaris* L.) is the most important food legume grown

worldwide, and especially for developing economies in South America, Africa and Asia[8]. The

origin of common bean is now central Mexico, and from there it spread to central and south

America around 165,000 years ago[9]. This resulted in the development of two major and eco-

geographically distinct common bean gene pools with partial reproductive isolation[10–12]. The

Mesoamerican gene pool was distributed from northern Mexico to Colombia and the Andean gene

pool ranged from southern Peru to northwestern Argentina. Since 8,000 years ago, each pool was

separately and selectively bred, leading to further diversification between them[13–15]. Because of pre-

existing genetic differences in each gene pool followed by divergent breeding history, common

bean presently offers a distinctive opportunity for understanding how the host and the environment

contribute to rhizosphere microbiome assembly.

The objective of this study was to quantify the influences of space, time and genotype on

detecting a core common bean root microbiome in the field, inclusive of bacteria, archaea and

fungi. We aimed to identify core microbial lineages that have cosmopolitan associations with

common bean and therefore potential as targets towards microbe-improved plant breeding. With

the cooperation of U.S. common bean breeding programs, we executed a first-of-its kind study of

two divergent bean genotypes grown under field conditions in five major North American bean

growing regions, including Michigan, Nebraska, Colorado and Washington. These two genotypes

belong to the Mesoamerican (Eclipse genotype) and Andean (California Early Light Red Kidney,

CELRK, genotype) gene pools that represent the major divergences from the wild bean ancestor[9].

For these two divergent genotypes, we also assessed how core members of the root microbiome

changed over plant development. Finally, we performed a comparative analysis of the bacterial

microbiome found in these study and with bean genotypes grown in South America[16,17]. From our

effort that was inclusive of both broad biogeography and plant development, we discovered a core

bean rhizosphere microbiome of 48 members that persistently associated with this nutritionally,

agronomically, and economically important crop. This core was discovered in spite of apparent

rhizosphere microbiome differences that were attributable to local soil conditions and management.

We did not detect an influence of plant genotype, suggesting that this core membership supersedes

it.

Results and Discussion

Study design, sequencing summary, and diversity

To assess the influence of environment and plant genotype on the common bean

rhizosphere microbiome, we designed a biogeography study of two divergent bean genotypes (from

Mesoamerican and Andean gene pools[18,19]), both grown in the field at five research and extension

farms that represent major U.S. bean production regions (**Table 1**)[20]. From each growing location,

triplicate bean plants grown in three to four plots were harvested at flowering (mid to late July).

We sequenced 39 rhizosphere (attached to the root, see methods) and root-associated soil

samples using 16S rRNA gene and ITS1 region, respectively to assess microbiome community

structures. We assessed the bacterial and archaeal communities using 16S rRNA gene amplicon

sequencing, and the fungal communities using ITS1 amplicon sequencing. After clustering reads to

OTUs at 97% similarity threshold and filtering chloroplast and mitochondria, there was 31,255 to

506,166 16S rRNA reads per sample and 22,716 to 252,810 ITS reads. We rarefied samples to

31,255 reads for 16S rRNA gene amplicons and to 22,716 for ITS. With these thresholds, we

achieved richness asymptotes for both datasets, suggesting that sequencing efforts were sufficient

to capture comparative dynamics and diversity (**Supplemental Fig. S1**). The total richness

observed at this rarefaction depth was 1,505 fungal and 23,872 bacterial and archaeal OTUs.

As reported in other rhizosphere studies, the total fungal diversity was lower than bacterial

/archaeal diversity in the rhizosphere of the common bean[21–23]. Richness varied by growing location

(ANOVA, F value=12.4, *p*-value<0.0001 and F value=13.1, *p*-value<0.0001 for 16S rRNA and ITS

data, respectively, **Supplemental Fig. S2**) but was highest at the Montcalm Research Facility

(Michigan, US) for both, bacteria/archaea and fungi. An analysis of community beta diversity

revealed strong biogeographic patterns in community structure explained by location, soil pH and

fertilization, in agreement with other literature[24–27] (**Supplemental Fig. S3**, see **Supplemental**

Text). However, an influence of plant genotype was either week or not detected, which we

partially attribute to plant growth in uncontrolled field conditions, which, in other plant species,

also show low influence of genotype on rhizosphere structure[24,25,28] (**Supplemental Fig. S2BD**, see

Supplemental Text).

A core rhizosphere microbiome is detected across U.S. bean growing regions

The common bean rhizosphere microbiome included the major expected lineages for both

bacteria and fungi (**Supplemental Fig. S4AB**), in agreement with other plant rhizosphere studies[29–]

[34]. Together with previous studies, these data provide more evidence that root-associated microbial

taxa are phylogenetically and functionally conserved[35]. Proteobacteria, Acidobacteria,

Bacteroidetes, and Actinobacteria collectively comprised on average 73.5% of the bacteria/archaeal

community, Ascomycota dominated the fungal community with a mean total relative abundance of

53% with notable sample-to-sample variance (range from 16.5% to 84.5%).

Despite differences in the rhizosphere microbiomes across growing locations, we noticed a

large number of OTUs that were shared among them for the bacterial/archaeal communities (2,173

taxa, mean 31.5%, range 29.5% to 34.7%). There was a smaller but notable overlap for the fungal

communities (70 taxa or mean 4.5%, range from 0.9% to 17.9%; **Supplemental Fig. S4CD**). These

data suggested that, despite measured edaphic differences across growing locations and strong

biogeographic signal, the common bean rhizosphere recruited many similar taxa that could be

functionally important for the bean. Therefore, we explored occupancy-abundance distributions of

taxa ([36,37] and references therein) to infer the core bean microbiome of taxa with an occupancy of 1

(i.e., found in all samples and across all growing locations; **Fig. 1AB**). Among bacteria and

archaea, 258 phylogenetically diverse taxa were cosmopolitan in the dataset (**Fig. 1C**), including

numerous and abundant Proteobacteria (117 OTUs) with a dominant taxon classified as

Arthrobacter sp. (FM209319.1.1474, mean relative abundance of 1.43%). The bacterial/archaeal

core also contained taxa of interest for potential plant benefits (e.g. *Sphingomonas*, *Rhizobium*,

Bacillus, Streptomyces), as well as some genera that can be associated with disease (e.g. *Ralstonia*).

There were 13 taxa in the fungal core (**Fig. 1D**), and these were largely composed of Ascomycota

(10 OTUs), with dominating taxon OTU823 from the *Phaeosphaeriaceae* family (mean relative

abundance 10.1%). Notably, taxa that were unique to either bean genotype were relatively rare and

inconsistently detected (**Fig. 1**, orange and black points). Together, these results suggest that

common bean, regardless of the genotype, consistently recruits particular microbiome taxa.

The neutral expectation of abundance-occupancy distributions is that very abundant taxa

will have high occupancy, while rare taxa will have low[37–40]. Taxa that deviate from the neutral

expectation are more strongly influenced by deterministic factors, like environmental conditions,

than by stochastic factors, like drift and dispersal. The neutral model fit[38,39] of the abundance-

occupancy distribution (solid line, **Fig. 1AB**) identified several taxa that had frequencies either

above or below the 95% confidence intervals of the model (dashed lines). Specifically, 13.7% of

the bacterial/archaeal and 30.4% of fungal taxa, deviated from the neutral expectation

(**Supplemental Dataset S3**). One hundred and seventy-one core taxa were predicted above the

neutral model partition; these deterministically-selected taxa are prime candidates for follow-up

studies of interactions with the host plant. Overall, the bacteria/archaea community had better fit to

the neutral expectation than fungal (R^2 of 0.74 and 0.34, and migration rates (m) of 0.301 and

0.003, respectively), suggesting that dispersal was relatively less limiting for the bacteria/archaea

than for the fungi. This finding agrees with other work suggesting that fungi are more sensitive to

local climate or more dispersal limited than bacteria[41–44].

A cross-continental core rhizosphere microbiome is detected for common bean

We wanted to better understand if these U.S. core taxa were associated with the bean

rhizosphere across a larger geographical scale, which would suggest the potential for selective plant

recruitment and cosmopolitan distribution of core taxa. Therefore, we compared our U.S. data to a

recently published study of rhizosphere bacteria and archaea from common beans grown in

Colombian agricultural soil[16]. The Colombian study offered a key contrast because it included eight

divergent bean lineages, including wild (n=2), landrace (n=1), and cultivated genotypes (n=5),

grown in soil from a different continent that has starkly different climate and management from the

U.S. growing regions. To enable direct comparison, we re-analyzed raw reads and compared the

datasets by matching to either the same taxon identifiers when clustered to SILVA database, or

100% identity by BLAST to *de novo* clustered reads (see Materials and Methods). Surprisingly,

39.6% (3,359 OTUs) of rhizosphere taxa from the Colombian-grown beans were also shared with

the U.S. dataset (**Fig. 2**). Both datasets included taxa that were highly represented in the other: 62%

of U.S. core (159 out of 258) were found also in Colombia, and 51% of Colombian core (433 out of

848) were shared with the U.S. (**Fig. 2A**). Core taxa were again defined stringently with an

occupancy of 1, and 48 taxa were found across all samples, inclusive of both datasets. These core

taxa were composed of many Proteobacteria, with *Rhizobiales* showing the most consistent relative

abundance between the studies (**Fig. 2B,** e.g. 0.187% and 0.138% in Colombia and U.S. dataset,

respectively). Notably, none of this cross-continental core taxa were universally detected in very

high abundance, and all but two OTUs (a U.S. *Arthrobacter* sp. and a Colombian *Austrofundulus*

limnaeus with mean relative abundances of 1.43% and 1.01%, respectively) would be classified as

rare by a typical mean relative abundance threshold below 1%, hinting to a potential role of rare

taxa in providing key functions. Notably, only 48% of these cross-continental shared core taxa have

genus classification, suggesting that most of them are under-described in their functional potential

and interactions with plants.

We also analyzed reads associated with the global core taxa with the UNOISE3 pipeline[45] to

generate predicted biological sequences (zero-radius OTUs – ZOTUs with100% sequence identity)

and provide the maximal possible biological resolution. We found that the global core taxa

consisted of 422 ZOTUs, and that there was a range of 2 to 35 ZOTUs identified within each OTU

(**Supplemental Fig. S5**). With one exception (HQ597858.1.1508), all of core OTUs (clustered at

97% sequence identity) contained at least one ZOTUs that also had an occupancy of 1. In addition,

all of the ZOTUs with an occupancy of 1 were also the most abundant ZOTU within each OTU

(**Supplemental Fig. S5**). This result suggests that the same members constitute the core even with

increased taxonomic resolution.

A recent study considered the effect of the common bean domestication history on the root

microbiome and identified a core set of microbial taxa that are consistently present with these

diverse bean genotypes, including bacterial taxa recruited from agricultural and natural soil from

Colombia[17]. This core also had high representation of *Proteobacteria*, *Acidobacteria*,

Actinobacteria and *Verrucomicrobia*, similar to those observed in the present study. We also re-

analyzed these data and show that 46 (out of 48) of the cross-continental core taxa identified here

have occupancy>0.9 across all three included studies (the present study, Pérez-Jaramillo et al.[16] and

Pérez-Jaramillo et al.[17]). Forty-two of these taxa had the highest possible occupancy of 1, but only

within rhizosphere samples from agricultural soils (**Supplemental Dataset S4**). When

incorporating data from beans grown in forest soils, the average occupancy decreases to 0.77

(median 0.75, min = 0.53). However, 14 taxa still had an occupancy of 1 when including beans

grown in forest soils (**Supplemental Dataset S4**).

In summary, these results show that common bean can associate with a core set of

rhizosphere microbiome members at taxon and ecotype levels, across diverse bean genotypes and

across continents. Additionally, these core taxa are likely enriched by the host in field agriculture,

as suggested by their higher occupancy in agroecosystems.

Core taxa are enriched in the rhizoplane and are consistently detected across bean development

To identify the common bean core taxa over space (a biogeographic core) we sampled

plants across growing locations at the same growth stage (flowering) and focused on the

rhizosphere compartment. However, the question remained whether these core taxa are detected

beyond that particular plant development stage. To answer this question, we conducted a field

experiment to assess the core taxa over time in plant development stage. In the next growing season

(2018), we used the same divergent bean genotypes grown at both Michigan, U.S, locations

(Montcalm and Saginaw Valley; see Material and Methods). We harvested root systems at 5

different plant development stages including flowering stage. We investigated the relative

abundance of the cross-continental core taxa and the U.S.-specific core taxa in both rhizosphere

(soil that could be removed from the root after shaking) and rhizoplane (soil adhered to the root

tissue and removed via vortex in buffer) compartments to determine their ability to closely

associate with the plant tissue. All 48 cross-continental core taxa were detected a year later in these

field, and their relative abundances were significantly higher in the rhizoplane as compared to the

rhizosphere irrespective of the plant development stage, bean genotype and growing location (**Fig.**

3A). Interestingly, the remaining core taxa that were found exclusively in the U.S. dataset at an

occupancy of 1 were equally abundant in the rhizosphere and rhizoplane at both Michigan growing

locations (**Fig. 3A**).

We next asked whether there were enrichments of particular core taxa by plant development

stage, root compartment, or growing location (**Fig. 3B**). On balance, almost all core taxa showed

some growth stage preference, but these trends were specific to each growing location and root

compartment. Despite these nuances, all core taxa were consistently found with high occupancy

inclusive of the plant development series (**Supplemental Fig. S6**).

Together these results suggest that these core taxa are selected by the plant early in the

development stage and maintained. Enrichment of the core taxa in the rhizoplane further supports

the hypothesis that these core taxa engage closely with the host plant.

Core taxa are not hub or connector taxa in an inter-kingdom microbiome network

Network analysis has been proposed to be a useful method to identify important members

of the plant microbiome with beneficial traits[46–48]. Hub taxa, identified by their high connectivity

with many members of the community, are regarded as the most important part of the community

and influence network structure and community stability[5,49]. We applied this method to ask if any of

the core taxa that we identified using abundance-occupancy were also key for co-occurrence

network structure. Additionally, we were interested in identifying fungal-bacterial co-occurrences

because of reports of their potential benefits for the plant[50,51]. To explore these patterns we applied

the molecular ecology network analysis pipeline[52] using samples collected across the U.S. where

we have both 16S rRNA and ITS amplicon data. We merged rarified 16S and ITS datasets, filtered

the datasets to include taxa with occupancy greater than or equal to 50%, and considered only

interactions significant at *p*-value<0.05 and LA score greater or equal to 0.88. The resulting

network included 572 taxa (nodes) and 1,857 statistically significant interactions (edges) structured

among 52 modules. Most of the modules were relatively small, with only six including more than

10 nodes (**Fig. 4**). The network was scale-free and had small-world characteristics as indicated by

the node degree distribution fitting to the power law model (R^2= 0.993), and also had significant

deviation of the modularity, length and clustering coefficients from those calculated from random

network (same number of nodes and edges), respectively (**Supplemental Dataset S5**).

The topological role of each taxon within the network was determined by the relationship

between their within-module (Z_i) and among-module (P_i) connectivity scores (**Fig. 4A**)[53]. Based on

this, the majority of taxa were peripheral (potentially, specialists; 563 nodes). There were also 3

connectors and 6 module hubs, but no network hubs. Indeed, in agreement with the beta-diversity

analysis (**Fig. S3**, Supporting Materials), there was a strong geographic signal in the largest four

modules, and these were comprised mostly of bacterial-bacterial (rather than bacterial-fungal or

fungal-fungal) hypothesized interactions (**Fig. 4B**). Interestingly, cross-kingdom interactions

constituted only a small fraction of all co-occurrences (n=168; bacteria-fungi=156, archaea-

bacteria=6 and archaea-fungi=6).

The analysis identified 26 co-occurrences between core taxa. However, we were surprised

to find that only two bacterial core and no fungal core taxa were also classified as network hubs

(taxa that connect to many other taxa within a modules) or connectors (taxa that connect across

modules). As exceptions, core *Chitinophagaceae* taxon FR749720.1.1492 was a module hub node

and a *Nitrobacter* sp. GDHX01215817.4.1477 was a connector. Our results, inclusive of a dataset

of divergent plant genotypes and broad biogeography, suggest that while hub and connector taxa

may be important for the maintenance of the root microbiome, these taxa are not consistently

detected in the common bean rhizosphere and, by deduction, could not be of universal importance

for the host plant. Our study cannot speak to the potential for functional redundancy among hub or

connector taxa, which could ultimately suggest a functional core among phylogenetically diverse

taxa[54]. Taken together, these results suggest that core taxa likely are important for the plant, while

hub and connector taxa are important for the integrity of the soil microbial community and its

responses to the local environment.

Prospectus

Legumes like the common bean that form symbiotic relationships with nitrogen fixing bacteria are

a crucial component of sustainable agriculture. Growing common bean can both improve soil

fertility[55,56] and contribute to food security, as bean production is >31 million metric tons

worldwide (http://www.fao.org/faostat/, accessing on July 12, 2019)). However, common bean

yield estimates, production areas, and even nutritional value, are predicted to decrease in this

century due to the impacts of global change[57–59]. Thus, breeding or other improvements to common

bean is needed to make this important crop resistant to stressors associated with climate change

while increasing yields to meet higher demands. To date, only traditional plant breeding

approaches have been used for the common bean, exploiting the genetic diversity of the plant, but

the latest knowledge of the plant microbiome in supporting host well-being has opened new

avenues for breeding the next generation of crops with plant microbiome manipulation[4].

Here, we show that common bean genotype has marginal effect on the rhizosphere

microbiome grown in field conditions. We describe for the first time the common bean core

microbiome at multiple spatial and temporal scales. A majority of the 48 core bacterial taxa

cosmopolitan among these common bean rhizospheres are under-described or largely unknown, as

only 23 out of 48 have genus classification. But, due to their ubiquitous association with the bean

over space and time, we hypothesize that these taxa provide functions that are crucial for common

bean health. A next step is to understand functions associated with these taxa and to determine if

and how they contribute to plant health and productivity under different growth conditions, such as

drought or with particular management strategies[28,60]. Additionally, targeting persistent microbiota

that are part of a native assemblage for management and manipulation can be advantageous, as

these are expected to be both host and environment adapted and thus may be more persistent *in situ*

than external or non-native biocontrols[3]. In conclusion, this work provides insights for prioritizing

plant-microbiome members that will advance goals in microbiome management and microbe-

improved crops.

Material and Methods

An expanded description of materials and methods is included in the Supporting Text.

Sampling and soil physicochemical analysis

We selected two common bean cultivars with very distinct genotypes: Eclipse[18] with

Mesoamerican origin and California Early Light Red Kidney (CELRK) and old kidney bean

landrace with Andean origin[19]. The plants were grown at 5 different locations on research extension

farms: Saginaw Valley (SVERC), Montcalm county (MRF) (both in Michigan), Scott Bluff county

(Nebraska) (NE), Fort Collins (Colorado) (CO) and Othello (Washington) (WA), each representing

an important U.S. common bean growing region[20]. Both bean genotypes were grown in the same

locations except in Michigan, where CELRK was grown at Saginaw Valley and Eclipse at

Montcalm county. Plants (n≥3) were harvested at flowering from replicated plots (n≥3) at each

location in summer 2017. The flowering time point was selected because we wanted to analyze

mature microbial communities as influenced by the plant and not by plant development. Plant roots

with attached soil were packed into bags, stored on ice and shipped immediately to Michigan State

University for processing.

From each sample, root attached soil was collected and used as rhizosphere soil and

remaining soil was collected and used as bulk soil. Both soils were sieved through 4 mm sieves to

remove any organic, plant debris and larger soil minerals before soil analysis and DNA isolation

and soil physical-chemical analysis (see Supporting Text). Soil analysis was done on bulk soil

samples pooled by plant genotype and plot.

308   We conducted a second study to assess the dynamics of the core taxa over plant

309 development. The bean genotypes, CELRK and Eclipse were grown at the same sites in Michigan,

310 U.S. as previously (Montcalm county (MRF) and Saginaw Valley (SVERC)) in the summer of

311 2018. Plants root systems were harvested at 5 growth stages, including stage 1: V2 (appearance of

312 second trifoliate), stage 2: V5 (appearance of fifth trifoliate), stage 3: flowering, stage 4: pod

313 filling, and stage 5: senescence/drying). At each sampling time, roots were collected and

314 transported on ice to the laboratory for immediate processing. Rhizosphere soil was collected by

315 gently shaking roots. Remaining attached soil (rhizoplane soil) was collected by first vortexing

316 roots in 1x phosphate buffer solution (PBS) for 4 min and finally removing the supernatant after

317 centrifugation for 10 min at 8000 g and 4°C. Collected soil was immediately frozen in liquid

318 nitrogen and stored at -80°C.

319   16S rRNA gene amplicon and ITS amplicon sequencing was performed at the Michigan

320 State Genomics Core Research Support Facility and analyzed using UPARSE (version 11)[61]. We

321 processed the 16S rRNA gene amplicons using an open-reference clustering first against the

322 SILVA v128 database[62, 63]. ITS OTU representative sequences were taxonomically annotated

323 using the CONSTAX tool[64] with the UNITE database version 7.2[65]. OTUs with unassigned

324 taxonomy at the domain level and OTUs annotated as mitochondria or chloroplasts were removed.

325 Statistical analysis and visualization were performed in R (see Supporting Text). The co-

326 occurrence network and global network properties were calculated using the Molecular Ecological

327 Network Analysis Pipeline (MENAP) [52] and visualized with Cytoscape v.3.5.1 [66]. Comparative

328 analyses with published datasets was performed by downloading and processing study raw reads

329 from NCBI to analyze them identically. Despite differences in our processing pipeline and the

330 dataset's originally published pipeline, we generated very similar numbers of OTUs (12,209

331 compared to 12,293[16]).

Acknowledgments

This work was supported by the Plant Resilience Institute at Michigan State University. We

are grateful to Dr. James Kelly, Dr. Phillip Miklas, Dr. James J. Heitholt, Dr. Carlos Urrea Florez,

Dr. Mark A. Brick, Dr. Juan M. Osorno, Dr. Thomas H Smith and other supporting staff at the

research extension farms for their partnership in this study. We would also like to thank the

Michigan State Genomics Core Research and Technology Support Facility and the Institute for

Cyber-Enabled Research High Performance Computing Center for excellent support and service.

Footnotes

Author contributions: N.S. and A.S. designed research, analyzed data and wrote the paper; and N.S.

performed research.

The authors declare no conflict of interest.

Data deposition: The raw sequence data have been deposited in the NCBI Sequence Read Archive

(BioProject accession no. PRJNA524532). All read processing steps, bioinformatic workflows,

and custom scripts are available on GitHub

(https://github.com/ShadeLab/PAPER_Stopnisek_2019_BeanBiogeography).

References

1. Berendsen, R. L. *et al*. Disease-induced assemblage of a plant-beneficial bacterial

consortium. *ISME J*. 1 (2018). doi:10.1038/s41396-018-0093-1

2. Castrillo, G. *et al*. Root microbiota drive direct integration of phosphate stress and

immunity. *Nature* **543**, 513–518 (2017).

3. Bell, T. H. *et al*. Manipulating Wild and Tamed Phytobiomes: Challenges and Opportunities.

Phytobiomes J. **3**, 3–21 (2019).

4. Wallenstein, M. D. Managing and manipulating the rhizosphere microbiome for plant health:

A systems approach. *Rhizosphere* (2017). doi:10.1016/j.rhisph.2017.04.004

5. Toju, H. *et al*. Core microbiomes for sustainable agroecosystems. *Nat*. *Plants* **4**, 247–257

(2018).

6. Hawkes, C. V. & Connor, E. W. Translating Phytobiomes from Theory to Practice:

Ecological and Evolutionary Considerations. *Phytobiomes* **1**, 57–69 (2017).

7. Shade, A. & Stopnisek, N. Abundance-occupancy distributions to prioritize plant core

microbiome membership. *Current Opinion in Microbiology* (2019).

doi:10.1016/j.mib.2019.09.008

8. Broughton, W. J. *et al*. Beans (Phaseolus spp.) – model food legumes. *Plant Soil* **252**, 55–

128 (2003).

9. Schmutz, J. *et al*. A reference genome for common bean and genome-wide analysis of dual

domestications. *Nat*. *Genet*. (2014). doi:10.1038/ng.3008

10. Rendón-Anaya, M. *et al*. Genomic history of the origin and domestication of common bean

unveils its closest sister species. *Genome Biol*. **18**, 60 (2017).

11. Bitocchi, E. *et al*. Beans (Phaseolus ssp.) as a Model for Understanding Crop Evolution.

Front. *Plant Sci*. **8**, 722 (2017).

12. Bitocchi, E. *et al.* Mesoamerican origin of the common bean (Phaseolus vulgaris L.) is

revealed by sequence data. *Proc. Natl. Acad. Sci.* **109**, 788–796 (2012).

13. Kwak, M. & Gepts, P. Structure of genetic diversity in the two major gene pools of common

bean (Phaseolus vulgaris L., Fabaceae). *Theor. Appl. Genet.* **118**, 979–992 (2009).

14. Bitocchi, E. *et al.* Molecular analysis of the parallel domestication of the common bean

(Phaseolus vulgaris) in Mesoamerica and the Andes. *New Phytol.* **197**, 300–313 (2013).

15. Schmutz, J. *et al.* A reference genome for common bean and genome-wide analysis of dual

domestications. *Nat. Publ. Gr.* **46**, (2014).

16. Perez-Jaramillo, J. E. *et al.* Linking rhizosphere microbiome composition of wild and

domesticated Phaseolus vulgaris to genotypic and root phenotypic traits. *ISME J.* 1–14

(2017). doi:10.1038/ismej.2017.85

17. Pérez-Jaramillo, J. E. *et al.* Deciphering rhizosphere microbiome assembly of wild and

modern common bean (Phaseolus vulgaris) in native and agricultural soils from Colombia.

Microbiome **7**, 114 (2019).

18. NDSU Agriculture. *Eclipse Black Bean*. (2004).

19. McClean, P. & Myers, J. Pedigrees of dry bean cultivars, lines and PIs. *Annu. Rep. Bean*

Improv. Coop **33**, 23–25 (1990).

20. USDA. *United States Department of Agriculture National Agricultural Statistics Service*

Crop Production 2016 Summary. (2017).

21. Singer, E., Bonnette, J., Kenaley, S. C., Woyke, T. & Juenger, T. E. Plant compartment and

genetic variation drive microbiome composition in switchgrass roots. *Environ. Microbiol.*

Rep. **11**, 185–195 (2019).

22. Coleman-Derr, D. *et al.* Plant compartment and biogeography affect microbiome

composition in cultivated and native *Agave* species. *New Phytol.* **209**, 798–811 (2016).

23.   De Souza, R. S. C. *et al*. Unlocking the bacterial and fungal communities assemblages of

sugarcane microbiome. *Sci. Rep.* **6**, 1–15 (2016).

24.   Walters, W. A. *et al*. Large-scale replicated field study of maize rhizosphere identifies

heritable microbes. *Proc. Natl. Acad. Sci.* **115**, 7368–7373 (2018).

25.   Wagner, M. R. *et al*. Host genotype and age shape the leaf and root microbiomes of a wild

perennial plant. *Nat. Commun.* **7**, 12151 (2016).

26.   Xu, J. *et al*. The structure and function of the global citrus rhizosphere microbiome. *Nat.*

Commun. **9**, 4894 (2018).

27.   Edwards, J. *et al*. Structure, variation, and assembly of the root-associated microbiomes of

rice. *Proc. Natl. Acad. Sci.* (2015). doi:10.1073/pnas.1414592112

28.   Edwards, J. A. *et al*. Compositional shifts in root-associated bacterial and archaeal

microbiota track the plant life cycle in field-grown rice. *PLOS Biol.* **16**, e2003862 (2018).

29.   Guttman, D. S. *et al*. Assembly and ecological function of the root microbiome across

angiosperm plant species. *Proc. Natl. Acad. Sci.* (2018). doi:10.1073/pnas.1717617115

30.   Hamonts, K. *et al*. Field study reveals core plant microbiota and relative importance of their

drivers. *Environ. Microbiol.* **20**, 124–140 (2018).

31.   Gomes, E. A., De Sousa, S. M. & Oliveira, C. A. Root-Associated Microbiome of Maize

Genotypes with Contrasting Phosphorus Use Efficiency. **2**, 129 (2018).

32.   Estendorfer, J. *et al*. The Influence of Land Use Intensity on the Plant-Associated

Microbiome of Dactylis glomerata L. *Front. Plant Sci.* **8**, 930 (2017).

33.   Leff, J. W., Lynch, R. C., Kane, N. C. & Fierer, N. Plant domestication and the assembly of

bacterial and fungal communities associated with strains of the common sunflower,

Helianthus annuus. *New Phytol.* **214**, 412–423 (2017).

34.   Lundberg, D. S. *et al*. Defining the core Arabidopsis thaliana root microbiome. *Nature* **488**,

(2012).

35. Levy, A. *et al.* Genomic features of bacterial adaptation to plants. *Nat. Genet.* (2018).

doi:10.1038/s41588-017-0012-9

36. Grady, K. L., Sorensen, J. W., Stopnisek, N., Guittar, J. & Shade, A. Assembly and

seasonality of core phyllosphere microbiota on perennial biofuel crops. *Nat. Commun.* **19**,

(2019).

37. Shade, A. *et al.* Macroecology to Unite All Life, Large and Small. *Trends Ecol. Evol.* **33**,

731–744 (2018).

38. Woodcock, S. *et al.* Neutral assembly of bacterial communities. *FEMS Microbiol. Ecol.* **62**,

171–180 (2007).

39. Burns, A. R. *et al.* Contribution of neutral processes to the assembly of gut microbial

communities in the zebrafish over host development. *ISME J.* **10**, 655–664 (2016).

40. Gaston, K. J. *et al.* Abundance-occupancy relationships. *J. Appl. Ecol.* **37**, 39–59 (2000).

41. Martiny, J. B. H., Eisen, J. A., Penn, K., Allison, S. D. & Horner-Devine, M. C. Drivers of

bacterial beta-diversity depend on spatial scale. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 7850–4

(2011).

42. Gumiere, T., Durrer, A., Bohannan, B. J. M. & Andreote, F. D. Biogeographical patterns in

fungal communities from soils cultivated with sugarcane. *J. Biogeogr.* **43**, 2016–2026

(2016).

43. Talbot, J. M. *et al.* Endemism and functional convergence across the North American soil

mycobiome. *Proc. Natl. Acad. Sci.* **111**, 6341–6346 (2014).

44. Tedersoo, L. *et al.* Global diversity and geography of soil fungi. *Science (80-.).* **346**, (2014).

45. Edgar, R. C. UNOISE2: improved error-correction for Illumina 16S and ITS amplicon

sequencing. *bioRxiv* 81257 (2016). doi:10.1101/081257

46. Agler, M. T. *et al*. Microbial Hub Taxa Link Host and Abiotic Factors to Plant Microbiome

Variation. *PLOS Biol*. **14**, e1002352 (2016).

47. Trivedi, P. *et al*. Keystone microbial taxa regulate the invasion of a fungal pathogen in agro-

ecosystems. *Soil Biol*. *Biochem*. **111**, 10–14 (2017).

48. Shi, S. *et al*. The interconnected rhizosphere: High network complexity dominates

rhizosphere assemblages. *Ecology Letters* **19**, (2016).

49. Carlström, C. I. *et al*. Synthetic microbiota reveal priority effects and keystone strains in the

Arabidopsis phyllosphere. *Nat*. *Ecol*. *Evol*. **3**, 1445–1454 (2019).

50. Durá, P. *et al*. Microbial Interkingdom Interactions in Roots Promote Arabidopsis Survival.

Cell **175**, 973–983 (2018).

51. Wagg, C., Schlaeppi, K., Banerjee, S., Kuramae, E. E. & van der Heijden, M. G. A. Fungal-

bacterial diversity and microbiome complexity predict ecosystem functioning. *Nat*.

Commun. **10**, 4841 (2019).

52. Deng, Y. *et al*. Molecular ecological network analyses. *BMC Bioinformatics* **13**, (2012).

53. Guimerà, R. & Nunes Amaral, L. A. Functional cartography of complex metabolic networks.

Nature **433**, 895–900 (2005).

54. Lemanceau, P., Blouin, M., Muller, D. & Moënne-Loccoz, Y. Let the Core Microbiota Be

Functional. *Trends in Plant Science* (2017). doi:10.1016/j.tplants.2017.04.008

55. Aguilar, O. M., Riva, O. & Peltzer, E. Analysis of Rhizobium etli and of its symbiosis with

wild Phaseolus vulgaris supports coevolution in centers of host diversification. *PNAS* **101**,

13548–13553 (2004).

56. Martínez-Romero, E. Diversity of Rhizobium-Phaseolus vulgaris symbiosis: overview and

perspectives. *Plant Soil* **252**, 11–23 (2003).

57. Jaggard, K. W., Qi, A. & Ober, E. S. Possible changes to arable crop yields by 2050. *Philos*.

Trans. R. Soc. Lond. B. Biol. Sci. **365**, 2835–51 (2010).

58.   Beebe, S., Rao, I., Mukankusi, C. & Buruchara, R. Improving Resource Use Efficiency and

Reducing Risk of Common Bean Production in Africa, Latin America, and the Caribbean. in

CIAT Book Chapter (ed. Hershey, C. H.) 1–18 (Centro Internacional de Agricultura Tropical

(CIAT), 2012).

59.   Hummel, M. *et al.* Reduction in nutritional quality and growing area suitability of common

bean under climate change induced drought stress in Africa. *Sci. Rep.* **8**, 16187 (2018).

60.   Xu, L. *et al.* Drought delays development of the sorghum root microbiome and enriches for

monoderm bacteria. *Proc. Natl. Acad. Sci.* **115**, E4284–E4293 (2018).

61.   Edgar, R. C. UPARSE: highly accurate OTU sequences from microbial amplicon reads. *Nat.*

Methods **10**, 996–998 (2013).

62.   Rideout, J. R. *et al.* Subsampled open-reference clustering creates consistent, comprehensive

OTU definitions and scales to billions of sequences. *PeerJ* **2**, e545 (2014).

63.   Quast, C. *et al.* The SILVA ribosomal RNA gene database project: Improved data

processing and web-based tools. *Nucleic Acids Res.* **41**, (2013).

64.   Gdanetz, K., Benucci, G. M. N., Vande Pol, N. & Bonito, G. CONSTAX: A tool for

improved taxonomic resolution of environmental fungal ITS sequences. *BMC*

Bioinformatics **18**, (2017).

65.   UNITE Community. UNITE USEARCH/UTAX release. (2017).

66.   Shannon, P. *et al.* Cytoscape: a software environment for integrated models of biomolecular

interaction networks. *Genome Res.* **13**, 2498–2504 (2003).

Table and Figure Legends

Table 1: Description of geographical, managemental and soil properties of the common bean

growing locations included in the study.

Sample same	Growing location	Elevation	Bean genotype	Weather (T, precipitation)*	Rotation history	Fertilization (per acre)	Irrigation	pH	Nitrogen (%)	Organic matter (%)
SVREC	Saginaw Valley Research and Extension Center, MI, US	190 m	CELRK	24.5ºC, 5.7 mm	Common bean, wheat, maize	Synthetic (400lbs of 15N-5 P-13K)	None	7.7 (± 0.1)	0.13 (± 0.00)	2.33 (± 0.06)
MRC	Montcalm Research Center, MI, US	280 m	Eclipse	24ºC, 7.4 mm	Common bean, maize, potato	Synthetic (200lbs of 19N-0P-19K)	Yes	5.9 (± 0.4)	0.10 (± 0.01)	2.13 (± 0.06)
NE	Scottsbluff, NE, US	1200 m	CELRK and Eclipse	27ºC, 2.18 mm	Common bean, maize, common bean	None	Yes	7.9 (± 0.1)	0.07 (± 0.01)	1.39 (± 0.18)
CO	ARDEC, Fort Collins, CO, US	1536 m	CELRK and Eclipse	31ºC, 1.4 mm **	Common bean, maize, barley	Organic (220lbs urea (2016), 3 tons manure (2015))	Yes	8.1 (± 0.2)	0.11 (± 0.00)	1.7 (± 0.15)
WA	WSU, Othello, WA, US	320 m	CELRK and Eclipse	26.3ºC, 0.2 mm	Common bean, maize, wheat	Synthetic (40lbs N/20lbs P/10lbs ZN2O5)	Yes	6.1 (± 0.4)	0.08 (± 0.01)	1.78 (± 0.10)

495   * Temperature (as max air temperature) and precipitation are averaged for the period between May

496   1st and July 31st 2017.

497   ** Data available only for the period between Jun 28th and Jul 31st 2017.

499
500

Fig. 1: Abundance-occupancy distributions were used to identify core members of the rhizosphere

microbiome for bacteria/archaea (A) and fungi (B). Taxa exclusive to a genotype are indicated in

orange (CELRK) or black (Eclipse), and taxa shared across both genotypes are white. The solid

line represents the fit of the neutral model, and the dashed line is 95% confidence around the model

prediction. Taxa with an occupancy of 1 (i.e., detected in all samples) were considered members of

the core. Relative abundance of these taxa is represented as boxplots, grouped by order and number

of taxa therein (C, D). Panels C and D are color-coded by phyla.

Fig. 2: A cross-continental core rhizosphere microbiome. There were 3361 shared

bacterial/archaeal taxa across U.S. and Colombia rhizosphere samples, suggesting highly similar

recruitment across continental scales. Forty-eight taxa were detected in all samples of both datasets,

as depicted by the Venn diagram (A), and included many Proteobacteria (B). Relative abundance of

the 48 U.S.-Colombia core taxa is represented as boxplots (left panel), grouped by order and

dataset (Colombia/U.S.). Number of taxa per order is represented as bars (right panel). Boxplots

and bars are color-coded by phylum.

Fig. 3: Relative abundance of core taxa in the root system of the common bean during plant

development by root compartment and growing location. The combined relative abundance of 48

core taxa is significantly higher in the rhizoplane (green) compared to the rhizosphere (orange) and

does not show plant development dependence (A – upper panel). The combined relative abundance

of remaining core taxa specific to the U.S. (n=210), tend to be high throughout the plant

development but are equally abundant in both compartments (A - lower panel). Stars above box

plots represent statistical significances as determined by Wilcoxon test (**** ≤ 0.0001, *** ≤

0.001, ** ≤ 0.01, * ≤ 0.05). Z-score normalized relative abundance of the 48 core taxa across plant

development, compartments and growing location (B). Taxa on the y-axis are arranged by their

classification at the phylum level.

Fig. 4: Network co-occurrence analysis shows that core rhizosphere microbiome members are

predominantly classified as peripheral taxa that are weakly connected, and generally clustering by

the growing location. The network depicts only clusters of modules that were connected by more

than 6 nodes (A). Nodes shape is representing domain associations (archaeal, bacterial or fungal)

and node size is proportional to its total number of connecting edges (A). The four largest modules

are generally reflective of community biogeography and distinguished by color (A). The within-

(Zi) and among- (Pi) module connectivity plot was used to identify module (Pi<0.62, Zi>0.2.5) or

network hub taxa (Pi>0.62, Zi>2.5), as well as connector (Pi>0.62, Zi<2.5) and peripheral taxa (B).

The density plots surrounding the Zi-Pi plot represent core (green) and non-core taxa (black) (B).

The relative abundances of taxa within each module is represented in box plots (C).