Prioritizing persistent microbiome members in the common bean rhizosphere: an integrated

analysis of space, time, and plant genotype

Nejc Stopnisek[1,2,3] and Ashley Shade[1,2,3,4,5]

1. Department of Microbiology and Molecular Genetics, Michigan State University, East

Lansing, MI 48840 USA

2. The Plant Resilience Institute, Michigan State University, East Lansing, MI 48840

3. The DOE Great Lakes Bioenergy Research Center, Michigan State University, East

Lansing, MI 48840

4. Program in Ecology, Evolutionary Biology and Behavior, Michigan State University, East

Lansing, MI 48840

5. Department of Plant, Soil and Microbial Sciences, Michigan State University, East Lansing,

MI 48840

Corresponding author: shadeash@msu.edu

Keywords: plant-microbiome, Phaseolus vulgaris, plant health, neutral model, co-occurrence

network, abundance-occupancy, 16S rRNA, ITS, hub taxa, sustainable agriculture, core

microbiome

Abstract

The full potential of managing microbial communities to support plant health is yet-

unrealized, in part because it remains difficult to ascertain which members are most important for

the plant. However, microbes that consistently associate with a plant species across varied field

conditions and over plant development likely engage with the host or host environment. Here, we

applied abundance-occupancy concepts from macroecology to quantify the core membership of

bacterial/archaeal and fungal communities in the rhizosphere of the common bean (*Phaseolus*

vulgaris). Our study investigated the microbiome membership that persisted over multiple

dimensions important for plant agriculture, including major growing regions, plant development,

annual plantings, and divergent genotypes, and also included re-analysis of public data. We found

48 core bacterial taxa that were consistently detected in all samples, inclusive of all datasets and

dimensions. This suggests reliable enrichment of these taxa to the plant environment and time-

independence of their association with the plant. More generally, this work provides a robust

approach for systematically prioritizing core microbiome memberships in any host or system.

Introduction

Agriculture requires more efficient use of available resources, and the naturally occurring, soil-dwelling microbiota offers potential to contribute to the responsible intensification of agriculture. Selection and breeding of plants for their beneficial associations with microbiota has promise to deliver a new generation of microbe-improved plants [1–6]. The ideal outcome of such efforts would achieve a balance of sustainable agriculture with food security. To achieve this, we must understand the relationships between plants and their associated microbiomes, including the differentiation of key or "core" members that engage with the plant directly from transient or opportunistic members that do not.

Common bean (*Phaseolus vulgaris* L.) is the most important food legume grown worldwide, and especially for developing economies in South America, Africa and Asia [7]. The origin of common bean is central Mexico, and from there it spread to central and to south America around 165,000 years ago [8]. This resulted in the development of two major and eco-geographically distinct common bean gene pools with partial reproductive isolation [9–11]. The Mesoamerican gene pool was distributed from northern Mexico to Colombia and the Andean gene pool ranged from southern Peru to northwestern Argentina. Since 8,000 years ago, each pool was separately and selectively bred, leading to further diversification between them [8, 12, 13]. Because of pre-existing genetic differences in each gene pool followed by divergent breeding history, common bean presently offers a distinctive opportunity for understanding how the host and the environment contribute to rhizosphere microbiome assembly.

The objective of this study was to apply approaches from macroecology to prioritize the persistent members of a core bacterial, archaeal, and fungal root microbiome inclusive of multiple gradients and categories of drivers expected to be important for plant agriculture. With the cooperation of the U.S. Bean coordinated agricultural project (Bean CAP), we executed a first-of-

its kind study of two divergent bean genotypes grown under field conditions in five major North

American bean growing regions, including Michigan, Nebraska, Colorado and Washington. These

two genotypes belong to the Mesoamerican (Eclipse genotype) and Andean (California Early Light

Red Kidney, CELRK genotype) gene pools that represent the major divergences from the wild

bean ancestor [8]. For these two divergent genotypes, we also assessed how core members of the

root microbiome changed over plant development and root compartment. We also used public data

to perform a comparative analysis of the bacterial microbiome found in our study with microbiome

members detected in other bean genotypes grown in South America [14, 15]. From our effort that

was inclusive of both broad biogeography (including the U.S. and Colombia), plant development,

and inter-annual plantings, we discovered a core bean rhizosphere microbiome of 48 members that

persistently associated with this nutritionally, agronomically, and economically important crop.

This core was discovered in spite of apparent microbiome differences that were attributable to local

soil conditions and management. However, we did not detect an influence of plant genotype,

suggesting that this core membership supersedes it.

Material and Methods

Study design, sampling and soil physicochemical analysis

We designed a biogeography study of two divergent bean genotypes (from Mesoamerican

and Andean gene pools [16, 17]), both grown in the field in the summer of 2017 at five research

and extension farms that represent major U.S. bean production regions (**Table 1**) [18]. The research

and extension farms were: Saginaw Valley (SVERC), Michigan (MI), Montcalm county (MRF),

MI, Scott Bluff county, Nebraska (NE), Fort Collins, Colorado (CO), and Othello, Washington

(WA).

Triplicate bean plants were grown in each of three (MI, WA, CO), or four (NE) plots,

totaling 9 or 12 plants per growing location. Plants were harvested at flowering stage (mid to late

July) because we wanted to analyze mature microbial communities and control for potential

differences in the microbiome over early plant development.

We selected two common bean cultivars with very distinct genotypes: Eclipse [16] of

Mesoamerican origin and California Early Light Red Kidney (CELRK), an old kidney bean

landrace of Andean origin [17]. Even though we included only two bean genotypes, we selected

highly divergent genotypes from far ends of the spectrum of the common bean genetic diversity;

these lineages diversified after a biogeographic bottleneck that separated them between Central and

South America [19]. We hypothesized that if there was an effect of bean genotype on the root

microbiome, it should be measurable when comparing these genotypes from divergent lineages.

Both bean genotypes were grown in each growing location, except in Michigan, where CELRK

was grown at Saginaw Valley and Eclipse at Montcalm county.

Plant roots with attached soil were packed into bags, stored on ice and shipped immediately

to Michigan State University for processing. From each sample, root-attached soil was collected by

gently shaking the roots and designated as rhizosphere soil; unattached root-zone soil was also

collected and used as bulk soil. It was expected that the rhizosphere soil contains a subset of

microbiome diversity that is recruited from the bulk soil [20] but we note that these soils were not

"bulk" in the traditional definition, in that they were collected from the agricultural fields and

expected to be proximal to and influenced by the roots of previous crops. Soils were sieved through

4 mm sieves to remove any plant debris and larger soil minerals before soil physical-chemical

analysis. Soil analysis was done at the Michigan State Soil Plant and Nutrient Laboratory on root-

zone soil samples pooled by plant genotype and plot (see Supporting Text for details).

We designed a second temporal study to assess the dynamics of the core taxa over plant

development. The two bean genotypes, CELRK and Eclipse, were grown at the same two sites in

Michigan, U.S., Montcalm county and Saginaw Valley, in the summer of 2018. Plants root systems

were harvested at 5 growth stages, including stage 1: V2 (appearance of second trifoliate), stage 2:

V5 (appearance of fifth trifoliate), stage 3: flowering, stage 4: pod filling, and stage 5:

senescence/drying). At each sampling time, roots were collected and transported on ice to the

laboratory for immediate processing. Rhizosphere soil was collected by gently shaking roots, as

described above. Any remaining, tightly-attached soil was designated as rhizoplane soil, and was

collected by first vortexing roots in 1x phosphate buffer solution (PBS) for 4 min and finally

removing the supernatant after centrifugation for 10 min at 8000 g and 4°C. Collected soil was

immediately frozen in liquid nitrogen and stored at -80°C.

Microbiome sequencing and analysis

There were 31 rhizosphere (attached to the root and pooled by plot) and 8 root-zone/bulk

(one pooled sample per growing location and plant genotype) soil samples sequenced for

microbiome analysis from the biogeography study, and 125 rhizosphere and 127 rhizoplane soil

(individual plants) samples sequenced from the plant development study. DNA extractions,

including negative controls, were performed as per standard soil protocols (see Supporting Text for

details). 16S rRNA gene amplicon and ITS amplicon sequencing was performed at the Michigan

State Genomics Core Research Support Facility. We processed the 16S rRNA gene amplicons

using an open-reference clustering first against the SILVA v128 database [21, 22]. First, raw reads

were merged, quality filtered, dereplicated, and clustered into 97% identity operational taxonomic

units (OTUs) using the UPARSE pipeline (version 11, [23]). Reads not matching to the SILVA

database were used for de-novo clustering at 97% sequence identity. Reference picked and de-novo

reads were combined before taxonomy was assigned [21]. Taxonomic annotations for 16S rRNA

gene OTU representative sequences were assigned in the QIIME 1.19 environment [24] using

SILVA database [22]. ITS OTU representative sequences were taxonomically annotated using the

CONSTAX tool [25] with the UNITE database version 7.2 [26]. OTUs with unassigned taxonomy

at the domain level and OTUs annotated as mitochondria or chloroplasts were removed.

Contaminant OTUs were removed using decontam package in R [27]. Additionally, we performed

zero-radios OTU analysis (aka ZOTUs that are clustered at 100% sequence identity) for reads

detected within the core OTUs. In short, first we subset all reads that were included within the core

OTUs and processed them through the UNOISE3 pipeline [28] to create ZOTUs. Then we

determined each ZOTU's affiliation to their originating core OTU by matching the read identities

between ZOTU and OTU clusters using a customized code (see GitHub repository).

Statistical analysis, including abundance-occupancy analysis to detect a core microbiome

[29], and data visualization were performed in R (see Supporting Text). The co-occurrence network

and global network properties were calculated using the Molecular Ecological Network Analysis

Pipeline (MENAP) [30] and visualized with Cytoscape v.3.5.1 [31]. Comparative analyses with

published datasets from Pérez-Jaramillo et al. [15] (BioProject ID PRJEB26084). was performed

by downloading study raw reads from NCBI and processing as per our data pipeline to analyze

both sets identically. An expanded description of all ecological statistics, MENAP parameters, and

our combined core microbiome analysis with public data are provided in Supporting Text.

Data and code availability

The raw sequence data are deposited in the NCBI Sequence Read Archive (BioProject ID

PRJNA524532). All read processing steps, bioinformatic workflows, R code, and custom scripts

are available on GitHub

(https://github.com/ShadeLab/PAPER_Stopnisek_2019_BeanBiogeography).

Results

Sequencing summary and microbial diversity across growing regions

There were 31,255 to 506,166 and 22,716 to 252,810 reads per sample for 16S rRNA and

ITS biogeography datasets, respectively. We rarefied samples to 31,255 reads for 16S rRNA gene

amplicons and to 22,716 for ITS. With these thresholds, we achieved richness asymptotes for both

datasets, suggesting that sequencing efforts were sufficient to capture comparative dynamics and

diversity (**Fig. S1**). The total richness observed at this rarefaction depth was 1,505 fungal and

23,872 bacterial and archaeal OTUs.

As reported in other rhizosphere studies, the total fungal diversity was lower than bacterial

/archaeal diversity in the rhizosphere of the common bean [32–34]. Richness varied by growing

location (ANOVA, F value=12.4, *p*-value<0.0001 and F value=13.1, *p*-value<0.0001 for 16S

rRNA and ITS data, respectively, **Fig. S2**) but was highest at the Montcalm Research Farm

(Michigan, US) for both, bacteria/archaea and fungi. An analysis of community beta diversity

revealed strong biogeographic patterns in community structure explained by location, soil pH and

fertilization, in agreement with other literature [35–38] (**Fig. S3**, see **Supplemental Text**).

However, an influence of plant genotype was either weak or not detected, which we partially

attribute to plant growth in field conditions, following observations made in other plant species

grown under field conditions [35, 36, 39] (**Fig. S2BD**, see **Supplemental Text).**

The common bean rhizosphere microbiome included the major expected lineages for both

bacteria and fungi (**Fig. S4AB**), in agreement with other plant rhizosphere studies [40–45].

Together with previous studies, these data provide more evidence that root-associated microbial

taxa are phylogenetically and potentially functionally conserved [46]. Proteobacteria,

Acidobacteria, Bacteroidetes, and Actinobacteria collectively comprised on average 73.5% of the

bacteria/archaeal community, Ascomycota dominated the fungal community with a mean total

relative abundance of 53% with notable sample-to-sample variance (range from 16.5% to 84.5%).

A core rhizosphere microbiome is detected across U.S. bean growing regions

We noticed a large number of OTUs that were shared among all growing locations them for

the bacterial/archaeal communities (2,173 taxa, mean 31.5%, range 29.5% to 34.7%). There was a

smaller but notable overlap for the fungal communities (70 taxa or mean 4.5%, range from 0.9% to

17.9%; **Fig. S4CD**). These data suggested that, despite measured edaphic differences across

growing locations and strong biogeographic signal, the common bean rhizosphere recruited many

similar taxa that could be functionally important for the bean. Therefore, we explored abundance-

occupancy distributions of taxa ([47, 48] and references therein) to infer the core bean microbiome

of taxa with an occupancy of 1 (i.e., found in all soil samples, all plots and across all growing

locations; **Fig. 1AB**). Among bacteria and archaea, 258 phylogenetically diverse taxa were

cosmopolitan in the dataset (**Fig. 1C**), including numerous and abundant Proteobacteria (117

OTUs) with a dominant taxon classified as *Arthrobacter* sp. (FM209319.1.1474, mean relative

abundance of 1.43%). The bacterial/archaeal core also contained taxa of interest for potential plant

benefits (e.g. *Sphingomonas*, *Rhizobium*, *Bacillus, Streptomyces*), as well as some genera that can

be associated with disease (e.g. *Ralstonia*). There were 13 taxa in the fungal core (**Fig. 1D**), and

these were largely composed of Ascomycota (10 OTUs), with dominating taxon OTU823 from the

Phaeosphaeriaceae family (mean relative abundance 10.1%). Notably, taxa that were unique to

either bean genotype were relatively rare and inconsistently detected (**Fig. 1**, orange and black

points). Together, these results suggest that common bean consistently recruits particular

microbiome taxa.

Next we wanted to investigate if the prioritized core taxa are indeed selected by the plant

environment or assembled through neutral processes by applying the Sloan neutral model [49, 50].

The neutral expectation of abundance-occupancy distributions is that very abundant taxa will have

high occupancy, while rare taxa will have low [48–51]. Taxa that deviate from the neutral

expectation are more strongly influenced by deterministic factors, like environmental conditions,

than by stochastic factors, like drift and dispersal. The neutral model fit of the abundance-

occupancy distribution (solid line, **Fig. 1AB**) identified several taxa that had frequencies either

above or below the 95% confidence intervals of the model (dashed lines). Specifically, 13.7% of

the bacterial/archaeal and 30.4% of fungal taxa, deviated from the neutral expectation (**Table S3**).

One hundred and seventy-one core taxa were predicted above the neutral model partition; these

deterministically-selected taxa are prime candidates for follow-up studies of interactions with the

host plant. Overall, the bacteria/archaea community had better fit to the neutral expectation than

fungal (R^2 of 0.74 and 0.34, and migration rates (m) of 0.301 and 0.003, respectively), suggesting

that dispersal was relatively less limiting for the bacteria/archaea than for the fungi. This finding

agrees with other work suggesting that fungi are more sensitive to local climate or more dispersal

limited than bacteria [52–55].

A core rhizosphere microbiome is detected for common bean grown on different continents

We wanted to better understand if these U.S. core taxa were associated with the bean

rhizosphere across a larger geographical scale, which would suggest the potential for selective plant

recruitment and cosmopolitan distribution of core taxa. Therefore, we compared our U.S. data to a

recently published study of rhizosphere bacteria and archaea from common beans grown in

Colombian agricultural soil [14]. The Colombian study offered a key contrast because it included

eight divergent bean lineages, including wild (n=2), landrace (n=1), and cultivated genotypes

(n=5), grown in soil from a different continent that has starkly different climate and management

from the U.S. growing regions. To enable direct comparison, we re-analyzed raw reads and

compared the datasets by matching to either the same taxon identifiers when clustered to SILVA

database, or 100% identity by BLAST to *de novo* clustered reads (see Materials and Methods).

Surprisingly, 39.6% (3,359 OTUs) of rhizosphere taxa from the Colombian-grown beans were also

shared with the U.S. dataset (**Fig. 2**). Both datasets included taxa that were highly represented in

the other: 62% of U.S. core (159 out of 258) were found also in Colombia, and 51% of Colombian

core (433 out of 848) were shared with the U.S. (**Fig. 2A**). Core taxa were again defined stringently

with an occupancy of 1, and 48 taxa were found across all samples, inclusive of both datasets. We

refer to this as the "global" core to distinguish the subset from the larger group of core taxa

inclusive to the US only (though note that this descriptor is for simplicity and that this does not

include global representation of bean root samples on Earth). These global core taxa were

composed of many Proteobacteria, with *Rhizobiales* showing the most consistent relative

abundance between the studies (**Fig. 2B,** e.g. 0.187% and 0.138% in Colombia and U.S. dataset,

respectively). Notably, none of this global core taxa were universally detected in very high

abundance, and all but two OTUs (a U.S. *Arthrobacter* sp. and a Colombian *Austrofundulus*

limnaeus with mean relative abundances of 1.43% and 1.01%, respectively) would be classified as

rare by a typical mean relative abundance threshold below 1%, hinting to a potential role of rare

taxa in providing key functions. A similar observation was made with rhizosphere microbiota of 19

herbaceous plant species, in which taxa of low abundance were among those significantly enriched

in the rhizosphere as compared to the bulk soil [56]. Notably, only 48% of these global core taxa

have genus classification, suggesting that most of them are under-described in their functional

potential and interactions with plants (**Table S4**).

We also analyzed reads associated with the global core taxa with the UNOISE3 pipeline

[28] to generate predicted biological sequences (zero-radius OTUs – ZOTUs with100% sequence

identity) and provide the maximal possible biological resolution. We found that the global core

taxa consisted of 422 ZOTUs, and that there was a range of 2 to 35 ZOTUs identified within each

OTU (**Fig. S5**). With one exception (HQ597858.1.1508), all of core OTUs (clustered at 97%

sequence identity) contained at least one ZOTU that also had an occupancy of 1. In addition, all of

the ZOTUs with an occupancy of 1 were also the most abundant ZOTUs within each OTU (**Fig.

S5**). This result suggests that the same members constitute the core even with increased taxonomic

resolution.

A recent study considered the effect of the common bean domestication history on the root

microbiome and identified a core set of microbial taxa that are consistently present with these

diverse bean genotypes, including bacterial taxa recruited from agricultural and natural soil from

Colombia [15]. This core also had high representation of *Proteobacteria*, *Acidobacteria*,

Actinobacteria and *Verrucomicrobia*, similar to those observed in the present study. We also re-

analyzed these data and show that 46 (out of 48) of the global core taxa identified here have

occupancy>0.9 across all three included studies (the present study, Pérez-Jaramillo et al. [14] and

Pérez-Jaramillo et al. [15]). Forty-two of these taxa had the highest possible occupancy of 1, but

only within rhizosphere samples from agricultural soils (**Table S4**). When incorporating data from

beans grown in forest soils, the average occupancy decreases to 0.77 (median 0.75, min = 0.53).

However, 14 taxa still had an occupancy of 1 when including beans grown in forest soils (**Table**

S4).

In summary, these results show that common bean can associate with a core set of

rhizosphere microbiome members at taxon and ecotype levels, across diverse bean genotypes and

across continents. Additionally, these core taxa are likely enriched by the host in managed soils, as

suggested by their higher occupancy in agroecosystems.

Core taxa are enriched in the rhizoplane and are consistently detected across bean development

To identify the common bean core taxa over space (a biogeographic core) we sampled

plants across growing locations at the same growth stage (flowering) and focused on the

rhizosphere compartment. However, the question remained whether these core taxa are detected

beyond that particular plant development stage. To answer this question, we conducted a field

experiment to assess the core taxa over time in plant development stage. In the next growing season

(2018), we used the same divergent bean genotypes grown at both Michigan, U.S, locations

(Montcalm and Saginaw Valley; see Material and Methods). We harvested root systems at 5

different plant development stages including flowering stage. We investigated the relative

abundance of the global core taxa and the U.S.-specific core taxa in both rhizosphere (soil that

could be removed from the root after shaking, n=125 samples) and rhizoplane (soil adhered to the

root tissue and removed via vortex in buffer, n=127 samples) compartments to determine their

ability to closely associate with the plant tissue. The range of the rhizosphere sequencing depth was

7,905-78,436 reads per sample, and for the rhizoplane it was 32-189,433 reads per sample. We

rarefied to 15,000 reads per samples (i.e. samples reaching richness asymptote), resulting in loss of

3 rhizosphere and 7 rhizoplane samples for a final dataset of 122 rhizosphere and 120 rhizoplane

samples. The total richness observed was 36,022 bacterial and archaeal OTUs.

From this development time series, we found that all 48 global core taxa were detected a

year later on these two Michigan farms. The collective relative abundances of the global core taxa

were significantly higher in the rhizoplane as compared to the rhizosphere irrespective of the plant

development stage, bean genotype and growing location (**Fig. 3A**). Interestingly, the remaining US

core taxa that were found exclusively in the U.S. dataset at an occupancy of 1 were equally

abundant in the rhizosphere and rhizoplane at both Michigan growing locations (**Fig. 3A**).

We next asked whether there were enrichments of particular core taxa by plant development

stage, root compartment, or growing location (**Fig. 3B**). On balance, almost all core taxa showed

some growth stage preference, but these trends were specific to each growing location and root

compartment. Despite these nuances, all core taxa were consistently found with high occupancy

inclusive of the plant development series (**Fig. S6**).

Together these results suggest that these core taxa are selected by the plant early in the

development stage and maintained. Enrichment of the core taxa in the rhizoplane further supports

the hypothesis that these core taxa engage closely with the host plant.

Core taxa are not hub or connector taxa in an inter-domain microbiome network

Network analysis has been proposed to be a useful method to identify important members

of the plant microbiome with beneficial traits [57–59]. Hub taxa, identified by their high

connectivity with many members of the community, are regarded as the most important part of the

community and influence network structure and community stability [60, 61]. We applied this

method to ask if any of the core taxa that we identified using abundance-occupancy were also key

for co-occurrence network structure. Additionally, we were interested in identifying fungal-

bacterial co-occurrences because of reports of their potential benefits for the plant [62, 63]. To

explore these patterns we applied the molecular ecology network analysis pipeline (MENAP)

which constructs ecological association network through random matrix theory (RMT) [30]. We

merged rarified 16S and ITS rhizosphere datasets, filtered the datasets to include taxa with

occupancy greater than or equal to 50%, and considered only interactions significant at p-

value<0.05 and RMT threshold of 0.88. The resulting network included 572 taxa (nodes) and 1,857

statistically significant co-occurrences (edges) structured among 52 modules. Most of the modules

were relatively small, with only six including more than 10 nodes (**Fig. 4**). The network was scale-

free (i.e. characteristics of the network are independent of the size of the network) and had small-

world characteristics (i.e. highly clustered) as indicated by the node degree distribution fitting to

the power law model (R^2= 0.993), and also had significant deviation of the modularity, length and

clustering coefficients from those calculated from random network (i.e. same number of nodes and

edges), respectively (**Table S5**).

The topological role of each taxon within the network was determined by the relationship

between their within-module (Z_i) and among-module (P_i) connectivity scores as described in [64].

Based on this, the majority of taxa were peripheral (potentially, specialists; 563 nodes) (**Fig. 4A**).

There were also 3 connectors and 6 module hubs, but no network hubs. Indeed, in agreement with

the beta-diversity analysis (**Fig. S3**, Supporting Materials), there was a strong geographic signal in

the largest four modules, and these were comprised mostly of bacterial-bacterial (rather than

bacterial-fungal or fungal-fungal) hypothesized interactions (**Fig. 4B**). We note that cross-domain

edges constituted only a small fraction of all co-occurrences (n=168; bacteria-fungi=156, archaea-

bacteria=6 and archaea-fungi=6).

The analysis identified 26 co-occurrences between core taxa. However, we were surprised

to find that only two bacterial core and no fungal core taxa were also classified as network hubs

(taxa that connect to many other taxa within a modules) or connectors (taxa that connect across

modules). As exceptions, core *Chitinophagaceae* taxon FR749720.1.1492 was a module hub node

and a *Nitrobacter* sp. GDHX01215817.4.1477 was a connector. Our results, inclusive of a dataset

of divergent plant genotypes and broad biogeography, suggest that while hub and connector taxa

may be important for the maintenance of the root microbiome, these taxa are not consistently

detected in the common bean rhizosphere and, by deduction, could not be of universal importance

for the host plant. Our study cannot speak to the potential for functional redundancy among hub or

connector taxa, which could ultimately suggest a functional core among phylogenetically diverse

taxa [65]. Taken together, these results suggest that core taxa likely are important for the plant,

while hub and connector taxa are important for the integrity of the soil microbial community and its

responses to the local environment.

Discussion

Numerous studies have aimed to detect core microbiome members for various plants and

animals (see references within [29, 66]). Because methods are inconsistent across studies and

because the parameters used to define a core are often arbitrary, there is some question as to how

robust or useful such studies may be. For these reasons, it has become easy to overlook new studies

that claim to have discovered a core microbiome for their system. If every study or design results in

a different core microbiome, how can the research build to move forward?

Here, we applied concepts from macroecology to prioritize, for the first time, a core plant

microbiome that persists at across growing locations on two continents, over plant development,

over two years of annual plantings at two different farms, across highly divergent plant genotypes,

and also across datasets collected by different research groups. Furthermore, core members were

enriched in the rhizoplane, and not just the rhizosphere. These multiple and consistent lines of

evidence provide strong support that these taxa likely engage with the plant and also demonstrate

the robustness of the abundance-occupancy method to discover core members.

A majority of the 48 core bacterial taxa among these common bean rhizospheres are under-

described or largely unknown, as only 23 out of 48 have a genus-level classification. But, due to

their ubiquitous association with the bean over space and time, we hypothesize that these taxa

provide functions that are crucial for common bean health and should be targeted for microbiome

management of in support of common bean productivity and wellness. To test this hypothesis

further research is needed, but plant-beneficial traits have been previously reported for members of

the genera that we identified as part of the bean core microbiome. For example, members

belonging to the genera *Mesorhizobium* and *Rhizobium* are known symbiotic nitrogen fixers of

legumes [67, 68], *Ramlibacter* sp. have the ability to promote P mobilization [69], and *Variibacter*,

Novosphingobium and *Sphingomonas* sp. harbor many specialized genes indicating their

relationship with plant hosts [70–72].

Discovering these core taxa is a first step in a rich line of inquiry to understand host

engagement with them. The next steps are to understand functions associated with these taxa and to

determine how they contribute to plant health and productivity under different growth conditions,

such as drought or with particular management strategies [39, 73]. These steps will include

cultivation dependent and independent approaches aimed to enrich and isolate core members,

assemble or bin genomes from isolates and metagenomes, annotate functional genes on both

chromosomes and plasmids, link functions and activities through transcript or metabolome

analyses, and perform experiments with constructed communities of core members to test

hypotheses about microbiome engagement with and benefits to the plant.

383   To conclude, this work provides robust approaches and general insights for prioritizing core

384 microbiome members, and that also advance goals in plant-microbiome management and microbe-

385 improved crops by providing insights into core member identities and dynamics.

388 **Acknowledgments**

389   This work was supported by the Plant Resilience Institute at Michigan State University. We

390 are grateful to Dr. James Kelly, Dr. Phillip Miklas, Dr. James J. Heitholt, Dr. Carlos Urrea Florez,

391 Dr. Mark A. Brick, Dr. Juan M. Osorno, Dr. Thomas H Smith and other supporting staff at the

392 research extension farms for their partnership in this study. We would also like to thank the

393 Michigan State Genomics Core Research and Technology Support Facility and the Institute for

394 Cyber-Enabled Research High Performance Computing Center for excellent support and service.

395 AS acknowledge support from Michigan State AgBioResearch (Hatch).

397 **Competing Interests**

398 The authors declare no conflict of interest.

400 **Author contributions**

401 N.S. and A.S. designed research, analyzed data and wrote the paper; and N.S. performed research.

403 *Supporting Text, Figures, and Tables accompany this manuscript.*

References

1. Dessaux Y, Grandclément C, Faure D. Engineering the Rhizosphere. *Trends Plant Sci* 2016; **21**: 266–278.

2. Gopal M, Gupta A. Microbiome Selection Could Spur Next-Generation Plant Breeding Strategies. *Front Microbiol* 2016; **7**: 1971.

3. Compant S, Samad A, Faist H, Sessitsch A. A review on the plant microbiome: Ecology, functions, and emerging trends in microbial application. *J Adv Res* 2019; **19**: 29–37.

4. Wille L, Messmer MM, Studer B, Hohmann P. Insights to plant-microbe interactions provide opportunities to improve resistance breeding against root diseases in grain legumes. *Plant Cell Environ* 2019; **42**: 20–40.

5. Busby PE, Soman C, Wagner MR, Friesen ML, Kremer J, Bennett A, et al. Research priorities for harnessing plant microbiomes in sustainable agriculture. *PLOS Biol* 2017; **15**: e2001793.

6. Kroll S, Agler MT, Kemen E. Genomic dissection of host–microbe and microbe–microbe interactions for advanced plant breeding. *Curr Opin Plant Biol* 2017; **36**: 71–78.

7. Broughton WJ, Hernández G, Blair M, Beebe S, Gepts P, Vanderleyden J. Beans (Phaseolus spp.) – model food legumes. *Plant Soil* 2003; **252**: 55–128.

8. Schmutz J, McClean PE, Mamidi S, Wu GA, Cannon SB, Grimwood J, et al. A reference genome for common bean and genome-wide analysis of dual domestications. *Nat Genet* 2014; **46**: 707–713.

9. Rendón-Anaya M, Montero-Vargas JM, Saburido-Álvarez S, Vlasova A, Capella-Gutierrez S, Ordaz-Ortiz JJ, et al. Genomic history of the origin and domestication of common bean unveils its closest sister species. *Genome Biol* 2017; **18**: 60.

10. Bitocchi E, Rau D, Bellucci E, Rodriguez M, Murgia ML, Gioia T, et al. Beans (Phaseolus

ssp.) as a Model for Understanding Crop Evolution. *Front Plant Sci* 2017; **8**: 722.

11.   Bitocchi E, Nanni L, Bellucci E, Rossi M, Giardini A, Zeuli PS, et al. Mesoamerican origin

of the common bean (Phaseolus vulgaris L.) is revealed by sequence data. *Proc Natl Acad*

Sci 2012; **109**: 788–796.

12.   Kwak M, Gepts P. Structure of genetic diversity in the two major gene pools of common

bean (Phaseolus vulgaris L., Fabaceae). *Theor Appl Genet* 2009; **118**: 979–992.

13.   Bitocchi E, Bellucci E, Giardini A, Rau D, Rodriguez M, Biagetti E, et al. Molecular

analysis of the parallel domestication of the common bean (Phaseolus vulgaris) in

Mesoamerica and the Andes. *New Phytol* 2013; **197**: 300–313.

14.   Perez-Jaramillo JE, Carrion VJ, Bosse M, Ferrao LF V, de Hollander M, Garcia AAF, et al.

Linking rhizosphere microbiome composition of wild and domesticated Phaseolus vulgaris

to genotypic and root phenotypic traits. *ISME J* 2017.

15.   Pérez-Jaramillo JE, de Hollander M, Ramírez CA, Mendes R, Raaijmakers JM, Carrión VJ.

Deciphering rhizosphere microbiome assembly of wild and modern common bean

(Phaseolus vulgaris) in native and agricultural soils from Colombia. *Microbiome* 2019; **7**:

114.

16.   NDSU Agriculture. Eclipse Black Bean. 2004. Fargo.

17.   McClean P, Myers J. Pedigrees of dry bean cultivars, lines and PIs. *Annu Rep Bean Improv*

Coop 1990; **33**: 23–25.

18.   USDA. United States Department of Agriculture National Agricultural Statistics Service

Crop Production 2016 Summary. 2017.

19.   Cichy KA, Porch TG, Beaver JS, Cregan P, Fourie D, Glahn RP, et al. A Diversity Panel for

Andean Bean Improvement. *Crop Sci* 2015; **55**: 2149.

20.   Sasse J, Martinoia E, Northen T. Feed Your Friends: Do Plant Exudates Shape the Root

Microbiome? *Trends Plant Sci* 2017.

21.   Rideout JR, He Y, Navas-Molina JA, Walters WA, Ursell LK, Gibbons SM, et al.

Subsampled open-reference clustering creates consistent, comprehensive OTU definitions

and scales to billions of sequences. *PeerJ* 2014; **2**: e545.

22.   Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, et al. The SILVA ribosomal

RNA gene database project: improved data processing and web-based tools. *Nucleic Acids*

Res 2013; **41**: D590-6.

23.   Edgar RC. UPARSE: highly accurate OTU sequences from microbial amplicon reads. *Nat*

Methods 2013; **10**: 996–998.

24.   Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, et al.

QIIME allows analysis of high-throughput community sequencing data. *Nat Methods* 2010;

7: 335–336.

25.   Gdanetz K, Benucci GMN, Vande Pol N, Bonito G. CONSTAX: a tool for improved

taxonomic resolution of environmental fungal ITS sequences. *BMC Bioinformatics* 2017;

18: 538.

26.   UNITE Community. UNITE USEARCH/UTAX release. 2017.

27.   Davis NM, Proctor DiM, Holmes SP, Relman DA, Callahan BJ. Simple statistical

identification and removal of contaminant sequences in marker-gene and metagenomics

data. *Microbiome* 2018; **6**: 226.

28.   Edgar RC. UNOISE2: improved error-correction for Illumina 16S and ITS amplicon

sequencing. *bioRxiv* 2016; 081257.

29.   Shade A, Stopnisek N. Abundance-occupancy distributions to prioritize plant core

microbiome membership. *Curr Opin Microbiol* 2019; **49**: 50–58.

30.   Deng Y, Jiang Y-H, Yang Y, He Z, Luo F, Zhou J. Molecular ecological network analyses.

BMC Bioinformatics 2012; **13**: 113.

31.   Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, et al. Cytoscape: A

Software Environment for Integrated Models of Biomolecular Interaction Networks.

Genome Res 2003; **13**: 2498–2504.

32.   Singer E, Bonnette J, Kenaley SC, Woyke T, Juenger TE. Plant compartment and genetic

variation drive microbiome composition in switchgrass roots. *Environ Microbiol Rep* 2019;

11: 185–195.

33.   Coleman-Derr D, Desgarennes D, Fonseca-Garcia C, Gross S, Clingenpeel S, Woyke T, et

al. Plant compartment and biogeography affect microbiome composition in cultivated and

native *Agave* species. *New Phytol* 2016; **209**: 798–811.

34.   De Souza RSC, Okura VK, Armanhi JSL, Jorrín B, Lozano N, Da Silva MJ, et al. Unlocking

the bacterial and fungal communities assemblages of sugarcane microbiome. *Sci Rep* 2016;

6: 1–15.

35.   Walters WA, Jin Z, Youngblut N, Wallace JG, Sutter J, Zhang W, et al. Large-scale

replicated field study of maize rhizosphere identifies heritable microbes. *Proc Natl Acad Sci*

U S A 2018; **115**: 7368–7373.

36.   Wagner MR, Lundberg DS, del Rio TG, Tringe SG, Dangl JL, Mitchell-Olds T. Host

genotype and age shape the leaf and root microbiomes of a wild perennial plant. *Nat*

Commun 2016; **7**: 12151.

37.   Xu J, Zhang Y, Zhang P, Trivedi P, Riera N, Wang Y, et al. The structure and function of

the global citrus rhizosphere microbiome. *Nat Commun* 2018; **9**: 4894.

38.   Edwards J, Johnson C, Santos-Medellín C, Lurie E, Podishetty NK, Bhatnagar S, et al.

Structure, variation, and assembly of the root-associated microbiomes of rice. *Proc Natl*

Acad Sci U S A 2015; **112**: E911-20.

39.   Edwards JA, Santos-Medellín CM, Liechty ZS, Nguyen B, Lurie E, Eason S, et al.

Compositional shifts in root-associated bacterial and archaeal microbiota track the plant life

cycle in field-grown rice. *PLOS Biol* 2018; **16**: e2003862.

40.   Fitzpatrick CR, Copeland J, Wang PW, Guttman DS, Kotanen PM, Johnson MTJ. Assembly

and ecological function of the root microbiome across angiosperm plant species. *Proc Natl*

Acad Sci U S A 2018; **115**: E1157–E1165.

41.   Hamonts K, Trivedi P, Garg A, Janitz C, Grinyer J, Holford P, et al. Field study reveals core

plant microbiota and relative importance of their drivers. *Environ Microbiol* 2018; **20**: 124–

140.

42.   Gomes EA, De Sousa SM, Oliveira CA. Root-Associated Microbiome of Maize Genotypes

with Contrasting Phosphorus Use Efficiency. 2018; **2**: 129.

43.   Estendorfer J, Stempfhuber B, Haury P, Vestergaard G, Rillig MC, Joshi J, et al. The

Influence of Land Use Intensity on the Plant-Associated Microbiome of Dactylis glomerata

L. *Front Plant Sci* 2017; **8**: 930.

44.   Leff JW, Lynch RC, Kane NC, Fierer N. Plant domestication and the assembly of bacterial

and fungal communities associated with strains of the common sunflower, Helianthus

annuus. *New Phytol* 2017; **214**: 412–423.

45.   Lundberg DS, Lebeis SL, Herrera Paredes S, Yourstone S, Gehring J, Malfatti S, et al.

Defining the core Arabidopsis thaliana root microbiome. *Nature* 2012; **488**.

46.   Levy A, Salas Gonzalez I, Mittelviefhaus M, Clingenpeel S, Herrera Paredes S, Miao J, et al.

Genomic features of bacterial adaptation to plants. *Nat Genet* 2018; **50**: 138–150.

47.   Grady KL, Sorensen JW, Stopnisek N, Guittar J, Shade A. Assembly and seasonality of core

phyllosphere microbiota on perennial biofuel crops. *Nat Commun* 2019; **10**: 4135.

48.   Shade A, Dunn RR, Blowes SA, Keil P, Bohannan BJM, Herrmann M, et al. Macroecology

to Unite All Life, Large and Small. *Trends Ecol Evol* 2018; **33**: 731–744.

49.   Woodcock S, Van Der Gast CJ, Bell T, Lunn M, Curtis TP, Head IM, et al. Neutral

assembly of bacterial communities. *FEMS Microbiol Ecol* 2007; **62**: 171–180.

50.   Burns AR, Stephens WZ, Stagaman K, Wong S, Rawls JF, Guillemin K, et al. Contribution

of neutral processes to the assembly of gut microbial communities in the zebrafish over host

development. *ISME J* 2016; **10**: 655–664.

51.   Gaston KJ, Blackburn TM, Greenwood JJD, Gregory RD, Quinn RM, Lawton JH.

Abundance-occupancy relationships. *J Appl Ecol* 2000; **37**: 39–59.

52.   Martiny JBH, Eisen JA, Penn K, Allison SD, Horner-Devine MC. Drivers of bacterial beta-

diversity depend on spatial scale. *Proc Natl Acad Sci U S A* 2011; **108**: 7850–4.

53.   Gumiere T, Durrer A, Bohannan BJM, Andreote FD. Biogeographical patterns in fungal

communities from soils cultivated with sugarcane. *J Biogeogr* 2016; **43**: 2016–2026.

54.   Talbot JM, Bruns TD, Taylor JW, Smith DP, Branco S, Glassman SI, et al. Endemism and

functional convergence across the North American soil mycobiome. *Proc Natl Acad Sci*

2014; **111**: 6341–6346.

55.   Tedersoo L, Bahram M, Põlme S, Kõljalg U, Yorou NS, Wijesundera R, et al. Global

diversity and geography of soil fungi. *Science (80-)* 2014; **346**.

56.   Dawson W, Hör J, Egert M, van Kleunen M, Pester M. A Small Number of Low-abundance

Bacteria Dominate Plant Species-specific Responses during Rhizosphere Colonization.

Front Microbiol 2017; **8**: 975.

57.   Agler MT, Ruhe J, Kroll S, Morhenn C, Kim S-T, Weigel D, et al. Microbial Hub Taxa Link

Host and Abiotic Factors to Plant Microbiome Variation. *PLOS Biol* 2016; **14**: e1002352.

58.   Trivedi P, Delgado-Baquerizo M, Trivedi C, Hamonts K, Anderson IC, Singh BK. Keystone

microbial taxa regulate the invasion of a fungal pathogen in agro-ecosystems. *Soil Biol*

Biochem 2017; **111**: 10–14.

59.   Shi S, Nuccio EE, Shi ZJ, He Z, Zhou J, Firestone MK. The interconnected rhizosphere:

High network complexity dominates rhizosphere assemblages. *Ecol Lett* 2016; **19**: 926–936.

60.   Carlström CI, Field CM, Bortfeld-Miller M, Müller B, Sunagawa S, Vorholt JA. Synthetic

microbiota reveal priority effects and keystone strains in the Arabidopsis phyllosphere. *Nat*

Ecol Evol 2019; **3**: 1445–1454.

61.   Toju H, Peay KG, Yamamichi M, Narisawa K, Hiruma K, Naito K, et al. Core microbiomes

for sustainable agroecosystems. *Nat Plants* 2018; **4**: 247–257.

62.   Durán P, Thiergart T, Garrido-Oter R, Agler M, Kemen E, Schulze-Lefert P, et al. Microbial

Interkingdom Interactions in Roots Promote Arabidopsis Survival. *Cell* 2018; **175**: 973-

983.e14.

63.   Wagg C, Schlaeppi K, Banerjee S, Kuramae EE, van der Heijden MGA. Fungal-bacterial

diversity and microbiome complexity predict ecosystem functioning. *Nat Commun* 2019; **10**:

1–10.

64.   Guimerà R, Nunes Amaral LA. Functional cartography of complex metabolic networks.

Nature 2005; **433**: 895–900.

65.   Lemanceau P, Blouin M, Muller D, Moënne-Loccoz Y. Let the Core Microbiota Be

Functional. *Trends Plant Sci* . 2017. Elsevier Ltd. , **22**: 583–595

66.   Risely A. Applying the core microbiome to understand host-microbe systems. *J Anim Ecol*

2020.

67.   Andrews M, Andrews ME, Andrews M, Andrews ME. Specificity in Legume-Rhizobia

Symbioses. *Int J Mol Sci* 2017; **18**: 705.

68.   Wang Q, Liu J, Zhu H. Genetic and molecular mechanisms underlying symbiotic specificity

in legume-rhizobium interactions. *Front Plant Sci* . 2018. Frontiers Media S.A. , **9**

69.   Skouri-Panet F, Benzerara K, Cosmidis J, Férard C, Caumes G, De Luca G, et al. In vitro

574   and in silico evidence of phosphatase diversity in the biomineralizing bacterium Ramlibacter

575   tataouinensis. *Front Microbiol* 2018; **8**: 2592.

576   70.   Lee JS, Lee KC, Kim KK, Lee B. Complete genome sequence of the Variibacter

577   gotjawalensis GJW-30T from soil of lava forest, Gotjawal. *J Biotechnol* 2016; **218**: 64–65.

578   71.   Khan AL, Waqas M, Kang SM, Al-Harrasi A, Hussain J, Al-Rawahi A, et al. Bacterial

579   endophyte Sphingomonas sp. LK11 produces gibberellins and IAA and promotes tomato

580   plant growth. *J Microbiol* 2014; **52**: 689–695.

581   72.   Aylward FO, McDonald BR, Adams SM, Valenzuela A, Schmidt RA, Goodwin LA, et al.

582   Comparison of 26 sphingomonad genomes reveals diverse environmental adaptations and

583   biodegradative capabilities. *Appl Environ Microbiol* 2013; **79**: 3724–33.

584   73.   Xu L, Naylor D, Dong Z, Simmons T, Pierroz G, Hixson KK, et al. Drought delays

585   development of the sorghum root microbiome and enriches for monoderm bacteria. *Proc

586   Natl Acad Sci* 2018; **115**: E4284–E4293.

Figure Legends

Fig. 1: Abundance-occupancy distributions were used to identify core members of the rhizosphere

microbiome for bacteria/archaea (A) and fungi (B). Taxa exclusive to a genotype are indicated in

orange (CELRK) or black (Eclipse), and taxa shared across both genotypes are white. The solid

line represents the fit of the neutral model, and the dashed line is 95% confidence around the model

prediction. Taxa with an occupancy of 1 (i.e., detected in all samples) were considered members of

the core. Relative abundance of these taxa is represented as boxplots, grouped by order and number

of taxa therein (C, D). Panels C and D are color-coded by phyla level.

Fig. 2: A global core rhizosphere microbiome. There were 3361 shared bacterial/archaeal taxa

across U.S. and Colombia rhizosphere samples, suggesting highly similar recruitment across

continental scales. Forty-eight taxa were detected in all samples of both datasets, as depicted by the

Venn diagram (A), and included many Proteobacteria (B). Relative abundance of the 48 U.S.-

Colombia core taxa is represented as boxplots (left panel), grouped by order and dataset

(Colombia/U.S.). Number of taxa per order is represented as bars (right panel). Labels on the y-axis

and bars are color-coded by phylum level.

Fig. 3: Relative abundance of core taxa in the root system of the common bean during plant

development by root compartment and growing location. The combined relative abundance of 48

core taxa is significantly higher in the rhizoplane (green) compared to the rhizosphere (orange) and

does not show plant development dependence (A – upper panel). The combined relative abundance

of remaining core taxa specific to the U.S. (n=210), tend to be high throughout the plant

development but are equally abundant in both compartments (A - lower panel). Stars above box

plots represent statistical significances as determined by Wilcoxon test (**** $\leq$ 0.0001, *** $\leq$

0.001, ** ≤ 0.01, * ≤ 0.05). Z-score normalized relative abundance of the 48 core taxa across plant

development, compartments and growing location (B). Taxa on the y-axis are arranged by their

classification at the phylum level.

Fig. 4: Network co-occurrence analysis shows that core rhizosphere microbiome members are

predominantly classified as peripheral taxa that are weakly connected, and generally clustering by

the growing location. The network depicts only clusters of modules that were connected by more

than 6 nodes (A). Nodes shape is representing domain associations (archaeal, bacterial or fungal)

and node size is proportional to its total number of connecting edges (A). The four largest modules

are generally reflective of community biogeography and distinguished by color (A). The within-

(Zi) and among- (Pi) module connectivity plot was used to identify module (Pi<0.62, Zi>0.2.5) or

network hub taxa (Pi>0.62, Zi>2.5), as well as connector (Pi>0.62, Zi<2.5) and peripheral taxa (B).

The density plots surrounding the Zi-Pi plot represent core (green) and non-core taxa (black) (B).

The relative abundances of taxa within each module is represented in box plots (C).

Supporting Figure legends

Fig. S1: Sequencing depth and rarefaction curves for 16S rRNA (A,B) and ITS (C, D) dataset from

samples collected in 2017. The red splitted line represents the rarefaction threshold. Note that we

submitted amplicons for sequencing from the 2017 sampling effort twice which resulted in the

sample to sample variation in read depth (rhizosphere 16S rRNA samples sequenced first followed

by the ITS amplicons with addition of the root-associated samples).

Fig. S2: Alpha diversity indices for the 16S RNA (A, B) and ITS (C, D) datasets. Represented are

rich-ness, Shannon and Pielou indices measured by growing location (A, C) and bean genotype (B,

D). For statistical comparison of the pairs we used ANOVA (A, C) or Wilcoxon test (B, D).

Fig. S3: Growing location drives bacterial/archaeal (A) and fungal (B) microbiome structure of the

common bean rhizosphere. The principal coordinate analysis (PCoA) is based on Bray-Curtis

distances. Growing location is indicated by color and plant genotype is indicated by shape shapes

(diamond=CELRK, circle=Eclipse, square=root zone soil). The strength of statistically significant

(p-value < 0.01) explanatory variables are shown as the length of fitted vectors.

Fig S4: Community composition (A, C) and number of shared taxa between sites represent as Venn

diagrams (B, D). The 16S rRNA dataset is represented in the top panels (A, B) and ITS in the lower

panels (C, D). The bar charts are colored based on the phylum association (phyla represented by

relative abundance < .05 are grouped and labelled as other). For Venn diagrams, samples were

grouped by the growing location and root zone samples were removed.

Fig. S5: Analysis of ZOTUs represented by each identified core OTU. 48 core OTUs were

represented by as few as 4 ZOTUs and by up to 35 ZOTUs. For every OTU we found at least one

ZOTU with occupancy = 1 and all of them, except of 2 ZOTUs, had also the highest relative

abundance among them. Points are color coded by their presence, red representing those with

occupancy < 1 and blue for ZOTUs with occupancy of 1. The OTUs on the x-axis are ordered

alphabetically.

Fig. S6: Occupancy of core OTUs in the development study. Occupancy is represented by color

and size.

Fig. S7: Comparison of relative abundance of the 48 core OTUs between the DNA isolation

methods used in the development study (G=Griffith, P=PowerSoil). Statistical difference,

determined by Wilcoxon test, is represented as star symbol (*<0.05, **<0.01, ***<0.001). The

pints are color coded by sample they derived from.

Fig. S8: The effect of isolation method on alpha and beta diversity. For alpha diversity richness,

Shannon and Pielou indices are presented (A). For the principal coordinates analysis, Bray-Curtis

distance matrix was used. Symbols are colored by samples. Wilcoxon test was used to determine

statistical differences between isolation methods for the alpha diversity metrices (*<0.05).

PERMANOVA was used to determine the effect of isolation method on community structure.

Tables

Table 1: Description of geographical, managemental and soil properties of the common bean

growing locations included in the study.

Sample same	Growing location	Elevation	Bean genotype	Weather (T, precipitation)*	Rotation history	Fertilization (per acre)	Irrigation	pH	Nitrogen (%)	Organic matter (%)
SVREC	Saginaw Valley Research and Extension Center, MI, US	190 m	CELRK	24.5ºC, 5.7 mm	Common bean, wheat, maize	Synthetic (400lbs of 15N-5 P-13K)	None	7.7 (± 0.1)	0.13 (± 0.00)	2.33 (± 0.06)
MRC	Montcalm Research Center, MI, US	280 m	Eclipse	24ºC, 7.4 mm	Common bean, maize, potato	Synthetic (200lbs of 19N-0P-19K)	Yes	5.9 (± 0.4)	0.10 (± 0.01)	2.13 (± 0.06)
NE	Scottsbluff, NE, US	1200 m	CELRK and Eclipse	27ºC, 2.18 mm	Common bean, maize, common bean	None	Yes	7.9 (± 0.1)	0.07 (± 0.01)	1.39 (± 0.18)
CO	ARDEC, Fort Collins, CO, US	1536 m	CELRK and Eclipse	31ºC, 1.4 mm **	Common bean, maize, barley	Organic (220lbs urea (2016), 3 tons manure (2015))	Yes	8.1 (± 0.2)	0.11 (± 0.00)	1.7 (± 0.15)
WA	WSU, Othello, WA, US	320 m	CELRK and Eclipse	26.3ºC, 0.2 mm	Common bean, maize, wheat	Synthetic (40lbs N/20lbs P/10lbs ZN2O5)	Yes	6.1 (± 0.4)	0.08 (± 0.01)	1.78 (± 0.10)

* Temperature (as max air temperature) and precipitation are averaged for the period between May

1st and July 31st 2017.

** Data available only for the period between Jun 28th and Jul 31st 2017.

Supporting Table legends

Table S1: PERMANOVA results for the 16S rRNA and ITS data. Highly correlated or/and

statistically significant values are highlighted in bold.

Table S2: 20 differentially abundant OTUs between the two plant genotypes as identified by using

DESeq2 (Love et al. 2014).

Table S3: Sloan neutral model summary.

Table S4: List of 48 core taxa and their taxonomic classification.

Table S5: Occupancy of core OTUs in agricultural, natural (forest) soils or when combined.

Results are based on the re-analysis of the data from the Pérez-Jaramillo et al. 2019. OTUs with

occupancy of 1 in both soils are highlighted in orange. Additional to the number we used green

gradient shading to represents the occupancy of each OTU.

Table S6: Summary of network properties of actual and random network.

Figures

Fig. 1: Abundance-occupancy distributions were used to identify core members of the rhizosphere

microbiome for bacteria/archaea (A) and fungi (B). Taxa exclusive to a genotype are indicated in

orange (CELRK) or black (Eclipse), and taxa shared across both genotypes are white. The solid

line represents the fit of the neutral model, and the dashed line is 95% confidence around the model

prediction. Taxa with an occupancy of 1 (i.e., detected in all samples) were considered members of

the core. Relative abundance of these taxa is represented as boxplots, grouped by order and number

of taxa therein (C, D). Panels C and D are color-coded by phyla level.

Fig. 2: A global core rhizosphere microbiome. There were 3361 shared bacterial/archaeal taxa

across U.S. and Colombia rhizosphere samples, suggesting highly similar recruitment across

continental scales. Forty-eight taxa were detected in all samples of both datasets, as depicted by the

Venn diagram (A), and included many Proteobacteria (B). Relative abundance of the 48 U.S.-

Colombia core taxa is represented as boxplots (left panel), grouped by order and dataset

(Colombia/U.S.). Number of taxa per order is represented as bars (right panel). Labels on the y-axis

and bars are color-coded by phylum level.

Fig. 3: Relative abundance of core taxa in the root system of the common bean during plant

development by root compartment and growing location. The combined relative abundance of 48

core taxa is significantly higher in the rhizoplane (green) compared to the rhizosphere (orange) and

does not show plant development dependence (A – upper panel). The combined relative abundance

of remaining core taxa specific to the U.S. (n=210), tend to be high throughout the plant

development but are equally abundant in both compartments (A - lower panel). Stars above box

plots represent statistical significances as determined by Wilcoxon test (**** ≤ 0.0001, *** ≤

0.001, ** ≤ 0.01, * ≤ 0.05). Z-score normalized relative abundance of the 48 core taxa across plant

development, compartments and growing location (B). Taxa on the y-axis are arranged by their

classification at the phylum level.

Fig. 4: Network co-occurrence analysis shows that core rhizosphere microbiome members are predominantly classified as peripheral taxa that are weakly connected, and generally clustering by the growing location. The network depicts only clusters of modules that were connected by more than 6 nodes (A). Nodes shape is representing domain associations (archaeal, bacterial or fungal) and node size is proportional to its total number of connecting edges (A). The four largest modules are generally reflective of community biogeography and distinguished by color (A). The within- (Zi) and among- (Pi) module connectivity plot was used to identify module (Pi<0.62, Zi>0.2.5) or network hub taxa (Pi>0.62, Zi>2.5), as well as connector (Pi>0.62, Zi<2.5) and peripheral taxa (B). The density plots surrounding the Zi-Pi plot represent core (green) and non-core taxa (black) (B). The relative abundances of taxa within each module is represented in box plots (C).