Single-particle view of stress-promoters induction dynamics: an interplay between MAPK signaling, chromatin and transcription factors

Victoria Wosika and Serge Pelet[*]

Department of Fundamental Microbiology

University of Lausanne

1015 Lausanne, Switzerland.

*Correspondance : serge.pelet@unil.ch

Abstract

Precise regulation of gene expression in response to environmental changes is crucial for cell survival, adaptation and proliferation. In eukaryotic cells, extracellular signal integration is often carried out by Mitogen-Activated Protein Kinases (MAPK). Despite a robust MAPK signaling activity, downstream gene expression can display a great variability between single cells. Using a live mRNA reporter, we monitored the dynamics of transcription in *Saccharomyces cerevisiae* upon hyper-osmotic shock. The transient activity of the MAPK Hog1 opens a temporal window where stress-response genes can be activated. Here we show that the first minutes of Hog1 activity are essential to control the activation of a promoter. The chromatin repression on a locus slows down this transition and contributes to the variability in gene expression, while binding of transcription factors increases the level of transcription. However, soon after Hog1 activity peaks, negative regulators promote chromatin closure of the locus and transcription progressively stops.

1 Introduction

A crucial function of all cellular life is the ability to sense its surroundings and adapt to
its variations. These changes in the extracellular environment will induce specific
cellular responses, orchestrated by signal transduction cascades which receive cues
from plasma membrane sensors. This information is turned into a biological response
by inducing complex transcriptional programs implicating hundreds of genes[1–3]. Tight
regulation of signaling is thus crucial to ensure the correct temporal modulation of gene
expression, which can otherwise alter the cell physiology[4–6]. Interestingly, single-cell
analyses have revealed that genes regulated by an identical signaling activity can
display a high variability in their transcriptional responses[7–10]. This noise in
transcriptional output questions how signal transduction can faithfully induce different
loci and which molecular mechanisms contribute to the variability in gene expression.

In eukaryotic cells, various environmental stimuli are transduced by the highly
conserved Mitogen-Activated Protein Kinases (MAPK) cascades[11,12]. They control a
wide range of cellular responses from cell proliferation, differentiation or apoptosis. In
Saccharomyces cerevisiae, a sudden increase in the osmolarity of the medium is
sensed by the High Osmolarity Glycerol (HOG) pathway, which leads to the activation
of the MAPK Hog1, a homolog of p38 in mammals[13,14]. Upon hyper-osmotic stress, the
kinase activity of Hog1 promotes the adaptation of the cells to their new environment
by driving an increase in the internal glycerol concentration, thereby allowing to
balance the internal and the external osmotic pressures. In parallel to its cytoplasmic
activity, Hog1 also transiently accumulates into the nucleus to induce the expression
of hundreds of osmostress-responsive genes (Fig. 1a). The MAPK is recruited to
promoter regions by Transcription Factors (TFs) and, in turn, Hog1 recruits chromatin
remodeling complexes, the Pre-Initiation Complex and the RNA Polymerase II (PolII)
to trigger gene expression[15,16]. Once cells have adapted, Hog1 is inactivated and exits
the cell nucleus, transcription stops and chromatin is rapidly reassembled at HOG-
induced gene loci.

Biochemical analyses of this pathway have identified the key players implicated in
gene expression and the central role played by the MAPK in all these steps[15]. In

parallel, single-cell measurements have uncovered the large variability present in their

expression. In particular, translational reporters and RNA-FISH measurements have

identified that slow chromatin remodeling promoted by the MAPK at each individual

locus is generating strong intrinsic noise in the activation of many stress-responsive

genes[9,17].

In order to get deeper insights into the regulation of osmostress-genes expression

kinetics, we aimed at monitoring the dynamics of mRNA production in live single cells.

Phage coat protein-based assays, like the MS2 or PP7 systems, have been used to

visualize mRNA in live single cells[18–20]. These experiments contributed to revealing the

bursty nature of transcription, whereby a set of polymerases simultaneously

transcribing a gene generates a burst in mRNA production, which is followed by a

pause in transcription[21–23].

In this study, we dissect the kinetics of transcription of osmostress-genes. The

production of mRNA is monitored using the PP7 phage coat protein assay. This

reporter allows us to measure with high temporal resolution and in a fully automated

manner, the fluctuations in transcription arising in hundreds of live single cells. This

analysis enables to dissect the contribution of various players to the overall

transcriptional output. We show that the first few minutes of MAPK activity will

determine if a gene is transcribed. We also demonstrate that the chromatin state of a

promoter will control the timing of activation and thus the variability of the transcription,

while the TF binding will influence the level and duration of the mRNA production.

Results

High osmotic pressure is sensed and transduced in the budding yeast *Saccharomyces*

cerevisiae via the HOG signaling cascade, which culminates in the activation of the

MAPK Hog1 (Fig. 1a). Upon activation, this key regulator accumulates in the nucleus

to trigger gene expression in a stress level-dependent manner (Supplementary Figure

1a). The activity of the kinase can be monitored by following its own nuclear

enrichment[24,25]. In parallel to Hog1, the general stress response pathway is induced

by the hyper-osmotic shock and the transcription factor Msn2 also relocate into the

nucleus with dynamics highly similar to the ones observed for Hog1 (Supplementary

Figure 1b and c) [26,27]. Nuclear Hog1 and Msn2 (together with its paralog Msn4) induce

osmostress-genes expression, with approximately 250 genes being up-regulated upon

osmotic shock[1,28,29]. The activity of the pathway is limited to the cellular adaptation

time, which conincide with the nuclear exit of Hog1 and the recovery of the cell size

(Supplementary Figure 1a and d). The fast and transient activity of the osmostress

response as well as the homogenous activation of the MAPK within the population[9,25]

(Supplementary Figure 1e), make this signaling pathway an excellent model to

understand the induction of eukaryotic stress-responsive genes, which are often

accompanied by important chromatin remodeling.

Monitoring the dynamics of osmostress-genes transcription

In order to quantify the dynamics of transcription in live single cells, we use the PP7

system to label the production of messenger RNAs (mRNA)[30]. Briefly, constitutively

expressed and fluorescently labeled PP7 phage coat proteins strongly associate to a

binding partner: an array of twenty-four PP7 mRNA stem-loops (PP7sl). In our settings,

this PP7 reporter construct is placed under the control of a promoter of interest and

integrated in the genome at the *GLT1* locus (Fig. 1b)[30], in a strain bearing a nuclear

tag (Hta2-mCherry) and expressing a fluorescently tagged PP7 protein (PP7ΔFG-

GFPenvy[31,32], abbreviated PP7-GFP, Methods). Upon activation of the promoter, local

accumulation of newly synthesized transcripts at the Transcription Site (TS) leads to

the formation of a bright fluorescent focus due to the enrichment in PP7-GFP

fluorescence above the background signal (Fig. 1c and Supplementary Movie 1). The

fluorescence intensity at the TS is proportional to the number of mRNA being

transcribed and thus to the instantaneous load of RNA polymerases. After termination,

single mRNAs are exported out of the nucleus and their fast diffusion in the cytoplasm

prevents their detection under the selected illumination conditions.

Typically, time-lapses with fifteen-second intervals for twenty-five minutes with six Z-

planes for the PP7-GFP channel on four fields of view were performed. Image

segmentation and quantification were performed automatically, allowing to extract up

to four hundred single-cell traces for each experiment[33]. The mean intensity of the 20

brightest pixels in the nucleus, from which the average cell fluorescence was

subtracted, was used as a measurement of TS intensity and thus as proxy for

transcriptional activity (Fig. 1d, Methods).

Fig. 1d displays the average TS fluorescence from more than 200 cells bearing the

p*STL1*-PP7sl reporter, following the activation of the HOG pathway by various NaCl

concentrations. The HOG-induced *STL1* promoter has been extensively studied at the

population and single cell level[9,17,34,35]. As expected, increasing salt concentrations

lead to a proportionally increasing transcriptional output from the cell population,

whereas no change in TS fluorescence is detected in the control medium.

The hundreds of dynamic measurements acquired with the PP7 reporter form a rich

dataset where multiple features can be extracted from each single-cell trace (Fig. 1e,

Methods). Our automated image segmentation and analysis allow to reliably quantify

the appearance (Start Time) and disappearance (End Time) of the TS (Supplementary

Figure 2 and Method). The maximum intensity of the trace and the integral under the

curve provide estimates of the transcriptional output from each promoter (Fig. 1e). In

addition, transcriptional bursts can be identified by monitoring of strong fluctuations in

the TS intensity.

Validation of the live mRNA reporter assay

The mRNA dynamics measured with the PP7 assay are in close agreement with

previously reported data set[34,36]. Nonetheless, we also verified with a dynamic protein

expression reporter that comparable results can be obtained (Supplementary Figure

3a). The dynamic Protein Synthesis Translocation Reporter (dPSTR) is an assay that

allows the kinetics of gene expression from a promoter of interest. It by-passes the

slow maturation time of fluorescent proteins (FP) by monitoring the relocation of the

fluorescent signal in the nucleus of the cell[37].

For the PP7 assay, as well as the dPSTR and many other expression reporters, an

additional copy of the promoter of interest is inserted in a non-native locus. In order to

address if this modified genomic environment alters the dynamics of gene expression,

we used CRISPR-Cas9 to integrate the PP7sl downstream of the endogenous *STL1*

promoter (Supplementary Figure 4). Interestingly, we observe only minor differences

between the p*STL1* at its endogenous location and at the *GLT1* locus. This observation

strongly suggests that the *STL1* promoter sequence placed at a non-endogenous locus
replicates many of the properties of the endogenous promoter.

Intrinsic noise in osmostress-gene activation
The microscopy images presented in Fig. 1c illustrate the noise that can be observed
in the activation of the p*STL1* promoter upon osmotic stress and which has been
previously reported[9,17]. In order to verify that this noise is not due to a lack of activation
of the MAPK Hog1 in the non-responding cells, we combined the p*STL1*-PP7sl reporter
and the Hog1-mCherry relocation assay in the same strain. As expected, we observe
an absence of correlation between the two measurements (Supplementary Figure 5).
Indeed, cells with similar Hog1 relocation behaviors can display highly variable
transcriptional outputs.

An additional assay to observe this heterogeneity is to monitor the activation of two
STL1 promoters within the same cell. Using a diploid strain where both *GLT1* loci were
modified with either a p*STL1*-24xPP7sl or a p*STL1*-24xMS2sl and expressing PP7-
mCherry and MS2-GFP proteins, we observe an uncorrelated activation of both loci
within each single cell (Supplementary Figure 6 and Supplementary Movie 2). This
observation confirms the high intrinsic noise generated by the *STL1* promoter upon
osmotic stress[9,37]. The highly dynamic measurements provided by the PP7 reporter
allows us to decipher some of the parameters that contribute to this large variability.

High variability in osmostress-genes transcription dynamics
In addition to p*STL1*, five other stress-responsive promoters often used in the literature
to report on the HOG pathway transcriptional activity were selected for this study[34,38].
Each reporter strain differs only by the one thousand base pairs of the promoter
present in front of the PP7sl (800bp for p*STL1*, 660 for p*ALD3*[9,39]); however, each
strain displays a different transcriptional response following a 0.2M NaCl stimulus (Fig.
2a). Because the level of accumulation of the PP7 signal at the transcription site and
the timing of the appearance and disappearance of the TS is different for each tested
promoter, it implies that the promoter sequence dictates multiple properties of the
transcription dynamics. These dynamic measurements are in general agreement with
control experiments performed with the dPSTR assay (Supplementary Figure 3b) and
previously published population-averaged data[34,37]. Importantly, expressing three

times more phage coat proteins did not change the parameters extracted from the PP7

measurements for the two strongest promoters, denoting the absence of titration of

PP7-GFP reporter proteins in our experimental settings (Supplementary Figure 7).

The automated analysis allows to identify the presence or absence of a transcription

site in each single cell, and thus the fraction of cells that induce the promoter of interest

(Fig. 2b). Interestingly, even in absence of stimulus, some promoters display a basal

level of transcription. In the p*GRE2*, p*HSP12* and p*GPD1* reporter strains, an active

transcription site can be detected in 5 to 20% of the cells in the few time points before

the stimulus (Fig. 2c, Supplementary Movie 3). If the period of observation is extended

to a twenty-five-minute time lapse without stimulus, this fraction increases 2 to 3-folds

(Supplementary Figure 8). Upon activation by 0.2M NaCl, the fractions of responding

cells for the three promoters that display basal expression overcomes 85%, while it

remains below 65% for the three promoters without basal induction.

Chromatin state sets the timing of transcription initiation

A key parameter controlled by the promoter sequence is the timing of induction. In Fig.

2d, the time when cells become transcriptionally active (Start Time) is plotted as a

Cumulative Distribution Function (CDF) only for the cells where a TS is detected after

the stimulus, thereby excluding basal expressing cells and non-responding cells.

Treatment with 0.2M NaCl results in a sudden activation of transcription (Fig. 2d). This

contrasts with non-induced samples, where the CDF of the promoters displaying basal

activity rises almost linearly due to stochastic activation during the recording window

(Supplementary Figure 8c).

Upon stress, the promoters displaying basal activity are induced faster than the

promoters that are repressed under log-phase growth, with p*GPD1* being activated the

fastest (~1 min), while p*ALD3* and p*STL1* require more than 4 minutes for activation

(Fig. 2e). However, there is a great variability in transcription initiation between cells of

the same population, since we generally observe 3 to 4 minutes delay between the

10th and 90th percentiles, with the exception of p*GPD1* where the induction is more

uniform and less than 2 min delay is observed (Fig. 2e). Comparison between

individual replicates demonstrates the reliability of our measurement strategy.

Interestingly, we observe a positive correlation between faster transcriptional activation

from p*GPD1*, p*HSP12* and p*GRE2* and the presence of basal expression level. These
promoters also display the highest numbers of responding cells upon a 0.2M NaCl
shock. These results suggest that basal expression is associated with a more
permissive chromatin state, which enables a faster activation and higher probability of
transcription among the cell population.

To test this hypothesis, we disrupted the function of the SAGA chromatin remodeling
complex by deleting *GCN5*[40]. As expected, we observe fewer transcribing cells and a
slower induction of the p*STL1* promoter in this background (Fig. 2f). Less remarkably,
abolishing histone H2AZ variants exchange at +1 and -1 nucleosomes by deleting
HTZ1[41] only results in a reduced percentage of transcribing cells. Conversely,
chromatin state at the *STL1* promoter can be loosened by relieving the glucose
repression using raffinose as a C-source[42]. Interestingly, a fraction of the cells grown
in these conditions displays basal expression from the p*STL1*-PP7 reporter and the
Start Time in raffinose is accelerated by 1 min compared to glucose (Fig. 2g).

The link between the chromatin state under log-phase growth and the ability to induce
stress-responsive genes is cofirmed by these results. A promoter that is tightly
repressed will need more Hog1 activity and thus more time to become transcriptionally
active, therefore displaying a lower fraction of responding cells.

Early Hog1 activity dictates transcriptional competence
The period of Hog1 activity provides a temporal window where transcription can
potentially be initiated. However, the switch to a transcriptionally active state takes
place almost exclusively within the first few minutes after the stimulus. When
comparing the characteristic timing of Hog1 nuclear enrichment to the CDF of Start
Times for cells bearing the p*STL1* reporter (Fig. 3a and Supplementary Figure 9a), we
observe that 90% of the transcribing cells initiate transcription during the first few
minutes of the stress response, while Hog1 nuclear accumulation rises and before it
drops below 80% of its maximum (decay time). A similar behavior is observed for all
the promoters tested, independently of the presence of basal levels (Fig. 3a and b).
For p*ALD3*, which is the slowest promoter tested, 87% of the Start Times are detected
before the decay of Hog1 activity (7 min) while the full adaptation time takes 14 min at
0.2M NaCl.

Interestingly, promoter output also decreases with the time after stimulus. Cells that
start transcribing p*STL1* early display a larger integral over the PP7 signal and a higher
maximum intensity compared to cells that initiate transcription later (Fig. 3c and
Supplementary Figure 9b). A similar behavior is quantified for all tested promoters
(Supplementary Figures 9c and d). These measurements demonstrate that the high
Hog1 activity present in the first minutes of the response is key to determine both the
transcriptional state and overall output of the promoters.

TFs control the dynamics and level of mRNA production
Promoters dictate the timing of transcriptional activation of the ORF and the level at
which the mRNA is produced. To extract the transcriptional level of each promoter, we
use as a proxy the maximum of the PP7 trace of each single cell where a transcription
event could be detected (Fig. 4a). This value represents the maximal loading of
polymerases on the locus during the period of transcription. Similar results are
obtained when comparing the integral below the PP7 trace, which represents the total
transcriptional output from a promoter (Supplementary Figure 10a). As shown in Fig.
4a, each promoter has an intrinsic capability to induce a given level of transcription,
which is independent from the presence of basal transcription or the locus activation
time. Indeed, p*GRE2* displays the lowest level of induction among all tested promoters,
despite the presence of basal transcription and being the second-fastest promoter
activated.

As expected, the recruitment of the RNA polymerases is stimulated by the stress; the
three promoters with basal activities display a higher transcriptional level upon a 0.2M
NaCl stress than in normal growth conditions (Fig. 4b and Supplementary Figure 10b).
Both the general stress transcription factors Msn2 and Msn4 and the TFs activated by
the MAPK Hog1 (Hot1, Sko1, Smp1) contribute to the transcriptional up-
regulation[29,38,43]. Based on studies on synthetic promoters, it has been established that
binding site number and distance from the Transcription Start Site (TSS) influence the
promoter output[44]. Unfortunately, osmostress promoters display a wide diversity in
number and affinity of TF binding sites and no obvious prediction of the transcriptional
activity can be drawn (Supplementary Figure 11). While multiple Msn2/4 binding sites

can be found on the *GPD1* and *STL1* promoter sequences, their activation are only
mildly affected by deletions of these two TFs (Supplementary Figure 12).

Both *GPD1* and *STL1* are primarily Hog1 targets[28,29,38]. However, their requirements
for Hog1 activity is strikingly different. In strains where the MAPK has been anchored
to the plasma membrane to limit its nuclear enrichment[45], p*STL1* induction is virtually
abolished (only 1.5% transcribing cells) while p*GPD1* activity is barely affected
(Supplementary Figure 13). Similarly, deletion of either TF Sko1 or Hot1 profoundly
alter the capacity of p*STL1* to be induced (Fig. 4c -f) while these same mutations have
a weaker effect on the *GPD1* promoter.

Because the induction of the *STL1* promoter requires an efficient chromatin
remodeling, every defect (TF deletion or absence of Hog1 in the nucleus) strongly
alters its capability to induce transcription. In comparison, the p*GPD1* is less perturbed
by these same defects. We postulate that TFs act in a cooperative manner on p*STL1*,
while they act independently of each other on p*GPD1*.

Bursts of PolII transcription in osmostress-gene activation
The PP7 and MS2 systems have allowed to directly visualize transcriptional bursting.
In order to identify bursts arising from osmostress promoters, we sought to detect
strong fluctuations in each single-cell trace. Fluctuations in TS intensities were filtered
to retain only peaks separated by pronounced troughs (Methods). In 20 to 30% of the
traces, two or more peaks are identified (Fig. 5a and b). The total length of the
transcript downstream of the promoter is 8kb (1.5kb for the stem loops + 6.5kb for
GLT1). Based on a transcription speed of 20bp/s[30], the expected lifetime of a transcript
at the TS is 6.6 min. This corresponds well to the mean duration observed for the
p*ALD3*, p*CTT1*, p*STL1* and p*GRE2* reporters (Fig. 5c). However, it is unlikely that the
strong TS intensities recorded are generated by a single transcript, but rather by a
group of RNA PolII that simultaneously transcribe the locus, probably forming convoys
of polymerases[46]. Indeed, single mRNA FISH experiments have shown that following
a 0.2M NaCl stress, the endogenous *STL1* locus produces on average 20 mRNAs per
cell, with some cells producing up to 100[36].

For p*HSP12* and p*GPD1*, the average peak duration is longer than 11 min (Fig. 5c),
suggesting that multiple convoys of polymerases are traveling consecutively through

the ORF. Unfortunately, the long half-lives of the transcripts on the locus prevent a

separation of individual groups of polymerases. However, when we achieve to isolate

individual peaks in the single cell traces, their duration becomes closer to the expected

value of 6.6 min (Fig. 5d). In addition, the output of the transcription estimated by the

maximum intensity of the trace or the integral under the whole curve is equal or lower

for traces with multiple pulses compared to traces where only a single peak is present

(Fig. 5e and Supplementary Figure 14). Together these data strengthen the notion that

these stress-responsive promoters are highly processive, displaying an elevated rate

of transcription once activated. Only brief pauses in the transcription can be observed

in a small fraction of the responding cells.

MAPK activity opens an opportunity window for transcription

We have shown that transcription initiation is dictated by early Hog1 activity. Next, we

want to assess what the determinants of transcription shutoff are and by extension,

the duration of transcriptional activity. In the HOG pathway, the duration of transcription

has been reported to be limited by the cellular adaptation time[34,38]. Therefore, the

duration of transcription is shorter after a 0.1M NaCl stress and longer after a 0.3M

stress, compared to a 0.2M stress (Fig. 6a). For the p*STL1* promoter, the last time

point where a PP7 signal is detected at the TS matches the timing of nuclear exit of

the MAPK at all concentrations tested (Fig. 6b).

In order to challenge this link between Hog1 activity and transcriptional arrest, we

sought to modulate the MAPK activity pattern by controlling the cellular environment in

a dynamic manner. Using a flow channel set-up, we generated a step, a pulse, or a

ramp in NaCl concentrations (Fig. 6c, Methods). These experiments were performed

in a strain carrying the p*STL1*-PP7sl reporter in conjunction with Hog1-mCherry,

allowing to monitor kinase activity and the downstream transcriptional response in the

same cell.

The step stimulus at 0.2M NaCl mimics the experiments performed in wells, where the

concentration of the osmolyte is suddenly increased at time zero and remains constant

throughout the experiments (Supplementary Movie 4). The mean responses at the

population level (Fig. 6c) confirm this relationship between Hog1 adaptation time and

transcription shutoff time. However, at the single cell level, no direct correlation is

observed between these two measurements due to important single cell variability (Fig.
6f).

In the pulse assay, 7 min after the initial 0.2M step, the NaCl concentration is set back
to 0M (Supplementary Movie 5). This shortens the MAPK activity period, as Hog1
leaves the nucleus immediately when cells are brought back in the normal growth
medium. Removing the kinase from the nucleus has a direct impact on the
transcriptional process. First, fewer cells become transcriptionally active. Second, the
active TS sites disappear within a few minutes after the end of the pulse (Fig. 6d and
e). Therefore in this context, we observe a direct correlation between Hog1 activity and
transcription, which is in line with the known role played by MAPKs, and Hog1 in
particular, on multiple steps of the transcriptional process[47,48].

The ramp experiment starts with a pulse at 0.2M NaCl followed by a slow increase of
the NaCl concentration up to 0.6M over the next 20 min (Supplementary Movie 6). This
constant rise in external osmolarity extends the Hog1 activity window by preventing
the adaptation of the cells. More cells can become transcriptionally active and the
transcription shut off is delayed (Fig. 6d and e). However, in these conditions, there is
a clear lack of correlation between Hog1 activity, which is sustained in many cells over
the 30min of the time-lapse, and the transcription output of the p*STL1* that stops much
earlier. Taken together, these experiments demonstrate that the MAPK activity is
required but not sufficient to sustain the transcriptional process. In the ramp
experiment, transcription cannot be sustained throughout the whole Hog1 activity
window, demonstrating that other factors contribute to limiting the duration of the
transcription.

Promoter identity influences the transcription shutoff time
In order to test whether the promoter identity plays a role in the process of transcription
shutoff, we quantified the duration of the transcriptional period for the six promoters
and plotted the cumulative distribution of End Times following a 0.2M NaCl stress (Fig.
7a and b). Interestingly, despite similar cell volume adaptation time for all the
experiments, the promoters display substantially different kinetics of inactivation.
Promoters transcribing at a lower level (p*CTT1* and p*GRE2*) terminate transcription
earlier. This shorter transcriptional window may reflect an inferior recruitment of

transcriptional activators to the promoter, enabling an earlier inhibition of transcription
due to chromatin closure. In addition, promoters with basal activity display an extended
period of transcription after adaptation (Fig. 7a and b). For p*GPD1* and p*GRE2*, this
results in a biphasic decay, where the first phase corresponds to the arrest of Hog1-
induced transcription and the second phase can be associated to the basal
transcription arising from these promoters (Fig. 7b). Note that basal transcription may
even be increased due to a higher basal Hog1 signaling activity post high osmolarity
conditions[49].

Remarkably, p*HSP12* transcription persists beyond the adaptation time, with nearly
30% of the cells displaying an active TS at the end of the experiment. This suggests
that basal expression from this promoter is strongly increased post-stimulus. In
contrast to p*GPD1* and p*GRE2*, p*HSP12* possesses numerous Msn2/4 binding sites.
Although the relocation dynamics of Hog1 and Msn2 are highly similar during the
adaptation phase, Msn2 displays some stochastic secondary pulses[27], that are not
correlated to Hog1 relocation events. This could explain the stronger basal expression
arising from this promoter post-adaptation (Supplementary Figure 1e and f).

To summarize, these measurements demonstrate that the pattern of MAPK activity
provides a temporal window where transcription can take place. When the signaling
cascade is shut off, transcription ceases soon afterward. However, the promoter
identity, and probably its propensity to recruit positive activators, will determine for how
long it can sustain an open chromatin environment favorable to transcription before
Hog1 activity decreases due to cellular adaptation.

Discussion

In this study, we have constructed PP7 reporter strains to monitor the transcription
dynamics of osmostress promoters. The second exogenous copy of the promoter is
integrated at the *GLT1* locus. This strategy provides a similar genomic environment for
all the promoters, in order to compare their specific characteristics. Interestingly, we
saw only minor differences in CDF of Start Times of the p*STL1* when integrated at its
endogenous locus compared to the *GLT1* locus. This observation provides a strong
evidence that TF binding and chromatin state of the duplicated promoter sequences

mimic closely the ones at the native environment of the gene. Note that the signal at the TS is expected to be proportional to the length of the transcribed mRNA. The *GLT1* locus with its 6.5kB length was expected to provide a signal four times stronger than the endogenous *STL1* ORF (1.7kB). The unexpectedly high signal obtained from the PP7 reporter at the endogenous locus may be indicative of global difference in transcription rates between the *GLT1* and *STL1* ORFs alternatively, the smaller *STL1* ORF might enhance transcription efficiency via gene-looping[50,51].

Our data illustrate the complex balance that exists between positive and negative regulators taking place at the stress induced loci. At each locus, positive and negative regulators will control the level and duration of transcription. We have shown that the first few minutes of Hog1 activity are essential to initiate the transcription. Transcription factors, chromatin remodelers such as the SAGA and RSC complexes, together with Hog1 will contribute to open and maintain an accessible chromatin environment at the stress-response loci[35,40]. Once initiated, transcription seems highly processive and only in a small fraction of traces, we are able to detect a pause in transcription. However, it has been shown that PolII recruits additional chromatin remodelers, including the Ino80 complex and Asf1 that will redeposit nucleosomes after acute transcription[52]. These conflicting activities will determine the overall duration of transcription at a locus. Indeed, promoters with lower transcriptional activity, such as p*GRE2* and p*CTT1*, recruit fewer positive activators and will be repressed faster by the negative regulators.

The repression level of a promoter during log-phase growth will determine the speed and the noise of the transcription activation process. Thus, for each promoter, a trade-off has to be found between these two contradictory requirements. For instance, *GPD1*, which is essential for survival to osmotic stress, has an important basal expression level and can thus be rapidly induced upon stress. Interestingly, the chromatin state, encoded in part by the promoter sequence, can be tuned by external growth conditions. Thus, the noise generated by a promoter is not rigidly set by its DNA sequence but fluctuate based on the environment.

In higher eukaryotes, the stress response MAPKs p38 and JNK relocate to the nucleus upon activation[53,54]. Early genes such as c-Fos or c-Jun, are induced within minutes

after activation of signaling[3,55]. Interestingly, these loci display basal expression and

require minimal chromatin modification for their induction[56,57]. Conversely, delayed

primary response and secondary response genes require more chromatin remodeling

to induce their activation[55,58]. These similarities with Hog1-gene transcriptional

regulation suggest a high conservation in the mechanisms used by MAPK in

eukaryotes to regulate the dynamics of gene expression.

Methods

Plasmids and yeast strains

Plasmids and yeast strains used in this study are listed in Supplementary Tables 1 and

2. All strains were constructed in the W303 background. Transformations were

performed with standard lithium-acetate protocols. Gene deletions and gene tagging

were performed with either pFA6a cassettes[59,60] or pGT cassettes[61]. Transformants

were selected with auxotrophy markers (Uracil, Histidine, Leucine, Tryptophan,

Adenine) and gene deletions were performed with antibiotic resistance to

Nourseothricin (NAT) or Kanamycin (KAN). In order to generate the membrane

anchored Hog1, the pGTT-mCherry vector was modified by inserting annealed oligos

encoding the last 30bp of the Ras2 sequence to obtain the pGTT-mCherry-CaaX

plasmid. A strain possessing the Hog1-mCherry:*LEU2* modification was transformed

with the pGTT-mCherry-CaaX plasmid linearized with *Xba*I and *Sac*I to induce a

marker switch and introduce the membrane anchoring motif.

PP7 and MS2 strains construction

The PP7-GFP plasmids are based on the bright and photostable GFPenvy fluorescent

protein[32]. The PP7 protein was derived from Larson et al.[30] (Addgene# 35194) with a

truncation in the capsid assembly domain (PP7ΔFG residues 67–75: CSTSVCGE[31]).

The expression of the PP7 construct is controlled by an *ADH1* promoter and a *CYC1*

terminator. The final construct pVW284 is cloned in a Single Integration Vector URA3

(pSIVu[61]). The PP7-mCherry was cloned by replacing the GFP by the mCherry

sequence. The MS2-GFP was generated by using the original MS2 sequence from

Betrand et al.[18], which also lacks the capsid assembly domain (Addgene# 27117),

inserted into the pVW284. The PP7 stem-loops plasmids are based on the previously

published p*POL1* 24x*PP7sl* integrative plasmid[30] (Addgene #35196). The stress

responsive promoters replace the *POL1* promoter in the original construct using 1kb

(0.8kb for p*STL1*, 0.66kb for p*ALD3*) upstream of the start codon. The p*STL1*-

24xMS2sl was generated by replacing the PP7sl with the MS2 stem loops obtained

from Betrand et al.[18] (Addgene# 31865).

A strain bearing the Hta2-mCherry nuclear marker and expressing the PP7-GFP was

transformed with plasmids containing the different osmostress promoters driving the

PP7sl expression, linearized with a *Not*I digestion and integrated upstream of the *GLT1*

ORF, as previously published[30]. Correct integration into the *GLT1* locus was screened

by colony PCR with primers in the *GLT1* ORF (+600 bp) and in the *TEF* terminator of

the selection marker on genomic DNA extractions. The integrity of the PP7 stem-loops

array was assessed with primers within the *TEF* terminator and in *GLT1* ORF (+250

bp) for all the promoters and deletions, after each transformation performed. For all the

strains used in the study, at least two clones with correct genotypes were isolated and

tested during a salt challenge time-lapse experiment. From the data analysis, the most

frequent phenotype was isolated and the strain selected.

To tag the endogenous locus of *STL1* with the 24x*PP7sl*, the plasmid pSP264 with the

STL1 promoter was modified by replacing the *GLT1* ORF sequence by a 500bp

sequence starting 100 bp after the start codon of *STL1*. The plasmid was digested

*Sac*I-*Not*I and purified over a gel to isolate a fragment that contains the p*STL1*-

24x*PP7sl*-*STL1*$_{100-600}$. A double-strand break was generated in the *STL1* ORF using

Cas9 and a sgRNA targeting the PAM motif GGG 62 bp upstream of the start codon.

The Cas9 and sgRNA are expressed from a 2*µ* plasmid (Addgene # 35464[62]) slightly

modified from the work from Laughery et al. (Addgene# 67639[63]). The purified DNA

fragment containing the stem-loops was used as repair DNA to promote homologous

recombination at the *STL1* locus (Supplementary Figure 4a). The correct size of the

inserted fragment was verified by colony PCR around the PP7sl insert. Multiple positive

clones were screened by microscopy. The results from two transformants are

presented in this work to ensure that potentially undesired DNA alterations by Cas9 do

not affect the response in the HOG pathway.

In order to generate the diploid reporter strain, a MAT**a** strain containing the PP7-
mCherry::*URA3*, p*STL1* 24xPP7sl:*GLT1* and Hta2-tdiRFP:*TRP1* was crossed to a
MATα strain bearing the MS2-GFP::*URA3*, p*STL1* 24xMS2sl:*GLT1* and Hta2-
tdiRFP:*NAT*. Haploid cells were mixed on a YPD plate for a few hours before cells
were resuspended in water and spread with beads on a selection plate (SD-TRP
+NAT).

The plasmids generated for this study are available on Addgene.

Yeast culture

Yeast cells were grown in YPD medium (YEP Broth: CCM0405, ForMedium) for
transformation or in Synthetic Defined (SD) medium (YNB:CYN3801/CSM: DCS0521,
ForMedium) for microscopy experiments. Before time-lapse experiments, cells were
grown at least 24 hours in log-phase. A saturated overnight culture in SD medium was
diluted into fresh SD-full medium to OD_{600} 0.025 in the morning and grown for roughly
8 hours to reach OD_{600} 0.3-0.5. In the evening, cultures were diluted by adding
$(0.5/OD_{600})\mu$l of cultures in 5ml SD-full for an overnight growth that kept cells in log-
phase conditions. Cultures reached an OD_{600} of 0.1-0.3 in the morning of the second
day and were further diluted when necessary to remain below an OD_{600} of 0.4 during
the day. To prepare the samples, these log-phase cultures were further diluted to an
OD_{600} 0.05 and sonicated twice 1min (diploids were not sonicated) before placing 200
μl of culture into the well of a 96-well glass bottom plate (MGB096-1-2LG, Matrical
Bioscience) previously coated with a filtered solution of Concanavalin A diluted to
0.5mg/ml in water (C2010, Sigma-Aldrich)[64]. Cells were let to settle for 30–45 minutes
before imaging. Osmotic shock was performed under the microscope, by adding 100
μl of a three times concentrated SD-full+NaCl stock solutions to the 200 μl of medium
already in the well, to reach the final desired salt concentration.

Microscopy

Images were acquired on a fully automated inverted epi-fluorescence microscope (Ti2-
Eclipse, Nikon) placed in an incubation chamber set at 30°C. Excitation was provided
by a solid-state light source (SpectraX, Lumencor) and dedicated filter sets were used
to excite and detect the proper fluorescence wavelengths with a sCMOS camera
(Flash 4.0, Hamamatsu). A motorized XY-stage was used to acquire multiple fields of

views in parallel and a piezo Z-stage (Nano-Z200, Mad City Labs) allowed fast Z-dimension scanning. Micro-manager was used to control the multidimensional acquisitions[65].

Experiments with PP7sl were acquired with a 60X oil objective. For strains with PP7-GFP and Hta2-mCherry, GFP (40ms, 3% LED power) and RFP (20ms), along with two bright field images were recorded every 15 seconds for the GFP and every minute for the other channels, for a total duration of 25 minutes. Six z-stacks were performed on the GFP channels covering ±1.2 μm from the central plane with 0.4 μm steps. An average bleaching of 32% for the GFP and 26% for the RFP for the whole time-lapse was quantified in a strain without the PP7 stem-loops, to avoid artifacts from the appearance of bright fluorescent foci. For all time-lapse experiments, media addition was performed before time point 4, defined as time zero. All microscopy experiments were performed in duplicate for non-induced control experiment and at least triplicate for the NaCl induced experiments.

Flow chamber experiments

The flow experiments were performed in Ibidi chambers (μ-Slide VI 0.4, Ibidi). Two 50ml Falcon tube reservoirs containing SD-full + 0.5 μg/ml fluorescein-dextran (D3305, ThermoFischer) and SD-full + 0.6 M NaCl were put under a pressure of 30mbar (FlowEZ, Fluigent). The media coming from each reservoir were connected using FEP tubing (1/16" OD x 0.020" ID, Fluigent) to a 3-way valve (2-switch, Fluigent). The concentration of NaCl in the medium was controlled using a Pulse-Width Modulation strategy[66,67]. Periods of 4 seconds were used and within this time, the valve controlled the fraction of time when SD-full versus SD-full + NaCl was flowing. TTL signals generated by an Arduino Uno board and dedicated scripts were used to control precisely the switching of the valve. The fluorescein present in the SD-full medium quantified outside the Cell object provided an estimate of the NaCl concentration in the medium. Some strong fluctuations in this signal were probably generated by dust particles in the imaging oil or FLSN-dextran aggregates in the flow chamber. Following 24hrs log-phase growth, cells bearing the p*STL1*-PP7sl reporter, Hog1-mCherry and Hta2-tdiRFP tags were diluted to OD 0.2, briefly sonicated and loaded in the ibidi

channel previously coated by Concanavalin A. Cells were left to settle in the channel

for 10 minutes before SD-full flow was started.

Raffinose experiment

For the experiments comparing p*STL1*-PP7sl induction in glucose versus raffinose,

cells were grown overnight to saturation in SD-full medium. The cultures were diluted

to OD 0.025 (Glucose) or 0.05 (Raffinose) and grown at 30°C for at least four hours.

In the raffinose medium, the expression level of the PP7-GFP was 2-fold lower than in

glucose. Because of this low fluorescence intensity, cells were imaged with a 40X

objective, and a single Z-plane was acquired. Manual curation of the images was

performed to define the Start Time in more than 250 cells. This experiment was

performed in duplicate.

Data analysis

Time-lapse movies were analyzed in an automated way: cell segmentation, tracking

and feature measurements were performed by the YeastQuant platform[33]. Summary

of the dataset, strains and cell numbers are provided in Supplementary Table 3. All

PP7 experiments were realized in at least two or three fully independent replicate

experiments. A representative experiment was selected for each strain and inducing

conditions, based on cell size and cell adaptation dynamics. The replicates which did

not pass one of these controls were discarded from the replicate analyses. Individual

single-cell traces were filtered based on cell shape and GFP intensity to remove

segmentation errors or other experimental artifacts. In addition, cells in mitosis were

removed from the analysis with a 0.95 filter on the nuclei eccentricity, to remove

artifacts from locus and PP7 signal duplication.

The Hta2 signal combined with the two bright field images allowed to define the

nucleus and cell borders. The GFP z-stacks were converted by a maximum intensity

projection in a single image that was used for quantification. In order to avoid improper

quantification of transcription sites at the nuclear periphery, the Nucleus object defined

by the histone fluorescence was expanded by 5 pixels within the Cell object to define

the ExpNucl object. The transcription site intensity was quantified by the difference

between the mean intensity of the 20 brightest pixels in the ExpNucl (HiPix) and the

median intensity from the whole cell. This provides a continuous trace which is close

to zero in absence of TS and increases by up to few hundred counts when a TS is
present. To identify the presence of a transcription site, a second feature named
ConnectedHiPix was used (Supplementary Figure 2a). Starting from the 20 HiPix, a
morphological opening of the image was performed to remove isolated pixels and
retaining only the ones that clustered together which correspond to the transcription
site. The ConnectedHiPix value was set to the mean intensity of the pixel present in
the largest object remaining after the morphological operation. If no pixel remained
after the morphological operation, the ConnectedHiPix was set to NaN. In each single-
cell trace, ConnectedHiPix values only detected for a single time point were removed.
After this filtering, the first and last time points where a ConnectedHiPix was measured
were defined as transcription initiation (Start Time) and shutoff (End Time),
respectively. Manual curation of Start and End Times from raw microscopy images
was performed to validate this transcription site detection strategy (Supplementary
Figure 2b and c). In order to detect individual transcriptional bursts in the HiPix traces,
the *findpeak* algorithm was used to identify in the trace all the peaks larger than a
threshold of 7 counts within the Start and End Times. Following this first process, a set
of conditions were defined to retain only the more reliable fluctuations: the drop
following the peak has to be larger the fourth of the peak intensity; the intensity of the
following peak has to rise by more than a third of the value at the trough. In addition,
the value of the peak has to be at least one fifth of the maximum intensity of the trace
in order to remove small intensity fluctuations being considered as peaks.

Data Availability
Source data for Figures 1d, 2a, 2f,2g, 3a, 4b, 4c, and 6c and Supplementary Figures
1, 4, 5, 6, 8, 12, 13 are provided with the paper. The raw images and additional features
measurements that support the findings of this study are available from the
corresponding author upon reasonable request.

Code Availability
The image analysis platform has been published previously[33]. A more recent version
of the code can be obtained from the corresponding author. A script to extract
measured parameters from the data is provided as a supplementary file.

References

1. Gasch, A. P. *et al.* Genomic expression programs in the response of yeast cells to environmental changes. *Mol Biol Cell* **11**, 4241–4257 (2000).
2. Roberts, C. J. *et al.* Signaling and circuitry of multiple MAPK pathways revealed by a matrix of global gene expression profiles. *Science* **287**, 873–880 (2000).
3. Ferreiro, I. *et al.* Whole genome analysis of p38 SAPK-mediated gene expression upon stress. *BMC Genomics* **11**, 144 (2010).
4. Berry, D. B. & Gasch, A. P. Stress-activated genomic expression changes serve a preparative role for impending stress in yeast. *Mol Biol Cell* **19**, 4580–4587 (2008).
5. Chen, R. E., Patterson, J. C., Goupil, L. S. & Thorner, J. Dynamic localization of Fus3 mitogen-activated protein kinase is necessary to evoke appropriate responses and avoid cytotoxic effects. *Mol Cell Biol* **30**, 4293–4307 (2010).
6. Formstecher, E. *et al.* PEA-15 Mediates Cytoplasmic Sequestration of ERK MAP Kinase. *Developmental Cell* **1**, 239–250 (2001).
7. Raser, J. M. & O'Shea, Erin K. Control of stochasticity in eukaryotic gene expression. *Science* **304**, 1811–1814 (2004).
8. Colman-Lerner, A. *et al.* Regulated cell-to-cell variation in a cell-fate decision system. *Nature* **437**, 699–706 (2005).
9. Pelet, S. *et al.* Transient activation of the HOG MAPK pathway regulates bimodal gene expression. *Science* **332**, 732–735 (2011).
10. Corrigan, A. M. & Chubb, J. R. Regulation of Transcriptional Bursting by a Naturally Oscillating Signal. *Current Biology* **24**, 205–211 (2014).
11. Roux, P. P. & Blenis, J. ERK and p38 MAPK-Activated Protein Kinases: a Family of Protein Kinases with Diverse Biological Functions. *Microbiology and Molecular Biology Reviews* **68**, 320–344 (2004).
12. Chen, R. E. & Thorner, J. Function and regulation in MAPK signaling pathways: lessons learned from the yeast Saccharomyces cerevisiae. *Biochim Biophys Acta* **1773**, 1311–1340 (2007).
13. Saito, H. & Posas, F. Response to hyperosmotic stress. *Genetics* **192**, 289–318 (2012).
14. Hohmann, S., Krantz, M. & Nordlander, B. Yeast osmoregulation. *Meth Enzymol* **428**, 29–45 (2007).
15. de Nadal, E. & Posas, F. Multilayered control of gene expression by stress-activated protein kinases. *EMBO J* **29**, 4–13 (2010).
16. de Nadal, E., Ammerer, G. & Posas, F. Controlling gene expression in response to stress. *Nat Rev Genet* **12**, 833–845 (2011).
17. Neuert, G. *et al.* Systematic Identification of Signal-Activated Stochastic Gene Regulation. *Science* **339**, 584–587 (2013).
18. Bertrand, E. *et al.* Localization of ASH1 mRNA particles in living yeast. *Mol Cell* **2**, 437–445 (1998).
19. Buxbaum, A. R., Haimovich, G. & Singer, R. H. In the right place at the right time: visualizing and understanding mRNA localization. *Nat Rev Mol Cell Biol* **16**, 95–109 (2015).
20. Urbanek, M. O., Galka-Marciniak, P., Olejniczak, M. & Krzyzosiak, W. J. RNA imaging in living cells – methods and applications. *RNA Biology* **11**, 1083–1095 (2014).
21. Lionnet, T. & Singer, R. H. Transcription goes digital. *EMBO Rep* **13**, 313–321 (2012).
22. Fritzsch, C. *et al.* Estrogen-dependent control and cell-to-cell variability of transcriptional bursting. *Molecular Systems Biology* **14**, e7678 (2018).

23. Zoller, B., Little, S. C. & Gregor, T. Diverse Spatial Expression Patterns Emerge from
Unified Kinetics of Transcriptional Bursting. *Cell* **175**, 835-847.e25 (2018).
24. Reiser, V., Ruis, H. & Ammerer, G. Kinase activity-dependent nuclear export opposes
stress-induced nuclear accumulation and retention of Hog1 mitogen-activated protein
kinase in the budding yeast Saccharomyces cerevisiae. *Mol Biol Cell* **10**, 1147–1161
(1999).
25. Muzzey, D., Gómez-Uribe, C. A., Mettetal, J. T. & van Oudenaarden, A. A systems-level
analysis of perfect adaptation in yeast osmoregulation. *Cell* **138**, 160–171 (2009).
26. Görner, W. *et al.* Nuclear localization of the C2H2 zinc finger protein Msn2p is regulated
by stress and protein kinase A activity. *Genes Dev.* **12**, 586–597 (1998).
27. Hao, N. & O'Shea, E. K. Signal-dependent dynamics of transcription factor translocation
controls gene expression. *Nature Structural & Molecular Biology* **19**, 31–39 (2012).
28. O'Rourke, S. M. & Herskowitz, I. Unique and Redundant Roles for HOG MAPK Pathway
Components as Revealed by Whole-Genome Expression Analysis. *Mol Biol Cell* **15**, 532–
542 (2003).
29. Capaldi, A. P. *et al.* Structure and function of a transcriptional network activated by the
MAPK Hog1. *Nat Genet* **40**, 1300–1306 (2008).
30. Larson, D. R., Zenklusen, D., Wu, B., Chao, J. A. & Singer, R. H. Real-Time Observation
of Transcription Initiation and Elongation on an Endogenous Yeast Gene. *Science* **332**,
475–478 (2011).
31. Chao, J. A., Patskovsky, Y., Almo, S. C. & Singer, R. H. Structural basis for the coevolution
of a viral RNA–protein complex. *Nat. Struct. Mol. Biol.* **15**, 103–105 (2007).
32. Slubowski, C. J., Funk, A. D., Roesner, J. M., Paulissen, S. M. & Huang, L. S. Plasmids
for C-terminal tagging in Saccharomyces cerevisiae that contain improved GFP proteins,
Envy and Ivy. *Yeast* **32**, 379–387 (2015).
33. Pelet, S., Dechant, R., Lee, S. S., van Drogen, F. & Peter, M. An integrated image analysis
platform to quantify signal transduction in single cells. *Integrative biology: quantitative
biosciences from nano to macro* **4**, 1274–1282 (2012).
34. de Nadal, E. *et al.* The MAPK Hog1 recruits Rpd3 histone deacetylase to activate
osmoresponsive genes. *Nature* **427**, 370–374 (2004).
35. Mas, G. *et al.* Recruitment of a chromatin remodelling complex by the Hog1 MAP kinase
to stress genes. *EMBO J* **28**, 326–336 (2009).
36. Li, G. & Neuert, G. Multiplex RNA single molecule FISH of inducible mRNAs in single yeast
cells. *Sci Data* **6**, 94 (2019).
37. Aymoz, D., Wosika, V., Durandau, E. & Pelet, S. Real-time quantification of protein
expression at the single-cell level via dynamic protein synthesis translocation reporters.
Nature Communications **7**, 11304 (2016).
38. Rep, M. *et al.* Osmotic stress-induced gene expression in Saccharomyces cerevisiae
requires Msn1p and the novel nuclear factor Hot1p. *Mol Cell Biol* **19**, 5474–5485 (1999).
39. Alepuz, P. M., de Nadal, E., Zapater, M., Ammerer, G. & Posas, F. Osmostress-induced
transcription by Hot1 depends on a Hog1-mediated recruitment of the RNA Pol II. *EMBO
J* **22**, 2433–2442 (2003).
40. Zapater, M., Sohrmann, M., Peter, M., Posas, F. & de Nadal, E. Selective requirement for
SAGA in Hog1-mediated gene expression depending on the severity of the external
osmostress conditions. *Mol Cell Biol* **27**, 3900–3910 (2007).
41. Wan, Y. *et al.* Role of the histone variant H2A.Z/Htz1p in TBP recruitment, chromatin
dynamics, and regulated expression of oleate-responsive genes. *Mol Cell Biol* **29**, 2346–
2358 (2009).
42. Ferreira, C. & Lucas, C. Glucose repression over Saccharomyces cerevisiae glycerol/H+
symporter gene STL1 is overcome by high temperature. *FEBS Lett* **581**, 1923–1927
(2007).
43. de Nadal, E. & Posas, F. Regulation of gene expression in response to osmostress by the
yeast stress-activated protein kinase Hog1. *Topics in Current Genetics* **20**, 81 (2008).

44. Sharon, E. *et al.* Inferring gene regulatory logic from high-throughput measurements of
thousands of systematically designed promoters. *Nat Biotechnol* **30**, 521–530 (2012).
45. Westfall, P. J., Patterson, J. C., Chen, R. E. & Thorner, J. Stress resistance and signal
fidelity independent of nuclear MAPK function. *Proc Natl Acad Sci USA* **105**, 12212–12217
(2008).
46. Tantale, K. *et al.* A single-molecule view of transcription reveals convoys of RNA
polymerases and multi-scale bursting. *Nat Commun* **7**, 12248 (2016).
47. Pokholok, D. K., Zeitlinger, J., Hannett, N. M., Reynolds, D. B. & Young, R. A. Activated
signal transduction kinases frequently occupy target genes. *Science* **313**, 533–536 (2006).
48. Proft, M. *et al.* The stress-activated Hog1 kinase is a selective transcriptional elongation
factor for genes responding to osmotic stress. *Mol Cell* **23**, 241–250 (2006).
49. Macia, J. *et al.* Dynamic signaling in the Hog1 MAPK pathway relies on high basal signal
transduction. *Science Signaling* **2**, ra13 (2009).
50. O'Sullivan, J. M. *et al.* Gene loops juxtapose promoters and terminators in yeast. *Nat
Genet* **36**, 1014–1018 (2004).
51. Ansari, A. A role for the CPF 3'-end processing machinery in RNAP II-dependent gene
looping. *Genes & Development* **19**, 2969–2978 (2005).
52. Klopf, E. *et al.* Cooperation between the INO80 complex and histone chaperones
determines adaptation of stress gene transcription in the yeast Saccharomyces cerevisiae.
Mol Cell Biol **29**, 4994–5007 (2009).
53. Cavigelli, M., Dolfi, F., Claret, F. X. & Karin, M. Induction of c-fos expression through JNK-
mediated TCF/Elk-1 phosphorylation. *The EMBO Journal* **14**, 5957–5964 (1995).
54. Wood, C. D., Thornton, T. M., Sabio, G., Davis, R. A. & Rincon, M. Nuclear Localization of
p38 MAPK in Response to DNA Damage. *Int. J. Biol. Sci.* 428–437 (2009)
doi:10.7150/ijbs.5.428.
55. Tullai, J. W. *et al.* Immediate-Early and Delayed Primary Response Genes Are Distinct in
Function and Genomic Architecture. *J. Biol. Chem.* **282**, 23981–23995 (2007).
56. Fowler, T., Sen, R. & Roy, A. L. Regulation of Primary Response Genes. *Molecular Cell*
44, 348–360 (2011).
57. O'Donnell, A., Odrowaz, Z. A. & Sharrocks, A. D. Immediate-early gene activation by the
MAPK pathways: what do and don't we know? *Biochemical Society transactions* **40**, 58–
66 (2012).
58. Ramirez-Carrozzi, V. R. *et al.* Selective and antagonistic functions of SWI/SNF and Mi-2β
nucleosome remodeling complexes during an inflammatory response. *Genes Dev.* **20**,
282–296 (2006).
59. Goldstein, A. L. & McCusker, J. H. Three new dominant drug resistance cassettes for gene
disruption in Saccharomyces cerevisiae. *Yeast* **15**, 1541–1553 (1999).
60. Sheff, M. A. & Thorn, K. S. Optimized cassettes for fluorescent protein tagging in
Saccharomyces cerevisiae. *Yeast* **21**, 661–670 (2004).
61. Wosika, V. *et al.* New families of single integration vectors and gene tagging plasmids for
genetic manipulations in budding yeast. *Molecular Genetics and Genomics* **291**, 2231–
2240 (2016).
62. Chee, M. K. & Haase, S. B. New and Redesigned pRS Plasmid Shuttle Vectors for Genetic
Manipulation of Saccharomyces cerevisiae. *G3 (Bethesda)* **2**, 515–526 (2012).
63. Laughery, M. F. *et al.* New vectors for simple and streamlined CRISPR-Cas9 genome
editing in Saccharomyces cerevisiae. *Yeast* **32**, 711–720 (2015).
64. Pelet, S., Aymoz, D. & Durandau, E. Temporal quantification of MAPK induced expression
in single yeast cells. *J Vis Exp* (2013) doi:10.3791/50637.
65. Edelstein, A., Amodaj, N., Hoover, K., Vale, R. & Stuurman, N. Computer control of
microscopes using µManager. *Curr Protoc Mol Biol* **Chapter 14**, Unit14.20 (2010).
66. Unger, M., Lee, S.-S., Peter, M. & Koeppl, H. Pulse Width Modulation of Liquid Flows:
Towards Dynamic Control of Cell Microenvironments. in *15th International Conference on*

miniaturized systems for chemistry and life sciences : Microtas 2011 1567–1569 (Chemical
and Biological Microsystems Society, 2011).
67. Unger, M. P. Interrogating the single cell: computational and experimental methods for
optimal live cell experiments. (ETH Zurich, 2014). doi:10.3929/ethz-a-010350761.
738

End Notes

Acknowledgment

We thank members the Pelet lab and Martin lab and for helpful discussions. Marta

Schmitt, Yves Dusserre, Gaëlle Spack, Joan Jordan and Clémence Varidel for

technical assistance. David Shore and his lab for helpful discussions and reagents.

Marie-Pierre Peli-Gulli and Claudio de Virgilio for plasmids, Tineke Lenstra for

suggesting the PP7ΔFG allele. Agathe Pelet for manual curation of microscopy

images. Eulalia de Nadal, Mariona Nadal-Ribelles and Veneta Gerganova for critically

reading the manuscript. Work in the Pelet lab is supported by SystemsX.ch (IPhD

51PHP0_157354), the Swiss National Science Foundation (SNSF, PP00P3_172900

and 31003A_182431) and the University of Lausanne.

Author Contributions

VW and SP designed the experiments, analyzed the data and wrote the manuscript.

VW established the condition for the PP7 imaging. VW and SP performed the

experiments.

Declaration of Interests

The authors declare no competing interests.

Figure Legends

Fig. 1. Monitoring the dynamics of osmostress-genes transcription.

a. Schematics of the transcriptional response induced by the MAPK Hog1 upon

osmotic stress. Under normal growth conditions, the genomic locus is repressed by

histones set in place by the Ino80 complex and Asf1/Rtt109. In addition, H3K4

methylated histones mediated by Set1 contribute to the further repression of the locus

(upper panel). When Hog1 is active (lower panel), it accumulates in the nucleus with

the transcription factors Msn2/4. Hog1 binds to the transcription factors Hot1 and Sko1,

allowing the remodeling of the chromatin by Rpd3 and the SAGA complex. The

polymerases can be recruited to the locus and the RSC and SWR complexes evict

nucleosomes on the ORF. **b.** Construction of the transcriptional reporter. The promoter

of interest (p*PROM*) is cloned in front of 24 stem-loops (*24xPP7sl*). This construct is

transformed in yeast and integrated in the *GLT1* locus 5'UTR, replacing the

endogenous promoter. Upon induction of the promoter, the mRNA stem-loops are

transcribed and recognized by the fluorescently-tagged PP7 phage coat protein. **c.**

Maximum intensity projections of Z-stacks of microscopy images from the p*STL1*-

PP7sl reporter system in a 0.2M NaCl osmotic stress time-lapse experiment. The

appearance of bright foci (arrow heads) in the nucleus of the cells denotes the active

transcription arising from the promoter. Scale bar represents 5 μm. **d.** Dynamics of the

p*STL1*-PP7sl transcription site intensity (20 brightest pixels in the nucleus minus the

median fluorescence of the cell) following hyperosmotic stress. The mean from 200 to

400 cells is represented by the solid line. The shaded areas represent the s.e.m. **e.**

Analysis of one representative single-cell trace. The raw trace (gray) is smoothed with

a moving average (dark blue) and normalized by subtracting the intensity of the first

time point after the stimulus. Multiple quantitative values can be extracted from this

trace (see Methods). Source data are provided for d.

Fig. 2. Chromatin state dictates the transcription initiation of stress-induced

promoters

a. Dynamics of the transcription site intensity from six different promoters following a

0.2M NaCl stress. The mean of at least 140 cells is represented by the solid line. The

shaded areas represent the s.e.m. **b.** Percentage of cells where a PP7 TS site was

detected. The light shaded area represents the percentage of PP7 positive cells before

the stimulus was added (basal transcription). **c.** The microscopy thumbnails display

cells bearing the p*GPD1*-PP7sl reporter system, where transcription sites (arrow

heads) can be detected before and after the stress of 0.2M NaCl. Scale bar represents

5 μm. **d.** Cumulative distribution function (CDF) of the Start Time for each promoter

considering only the cells that induce transcription after time zero. **e.** 10th, 50th and 90th

percentiles of the Start Times shown for the two to three replicates measured for each

promoter. The number of stars next to each measurement corresponds to the number

of promoters without basal level that are significantly different from the promoter with

basal level (two-sample *t*-test, p<0.05). **f.** Cumulative distribution functions of Start

Times for the p*STL1*-PP7sl strain in wild type, *htz1Δ* or *gcn5Δ* backgrounds. The inset

shows the percentage of PP7 positive cells in each background. **g.** Cumulative

distribution functions of Start Times for the p*STL1*-PP7sl strain grown in glucose or

raffinose. The inset shows the percentage of PP7 positive cells, the light blue bar the

basal positive PP7 cells. Source data are provided for a, f, g.

Fig. 3. Early Hog1 activity dictates promoter activation and output

a. In Hog1 nuclear relocation traces obtained from single cells, the timing of Hog1

nuclear entry (■), maximum enrichment (●), start of the decay in nuclear enrichment

(◆) and Hog1 adaptation (▲) can be identified (upper panel). The median (marker)

and 25th to 75th percentiles (lines) for these measurements are plotted for three

different osmotic stresses (central panel). The cumulative distribution functions of Start

Times for the p*STL1*-PP7sl reporter for these same three concentrations are plotted

(lower panel). **b.** Histograms of Start Times following a 0.2M stress for the five other

promoters tested. The vertical dashed line represents the median decay time of Hog1

measured at 0.2M. The number in the legend indicates the percentage of cells which

have initiated transcription before the median Hog1 decay time. **c.** The population of

p*STL1*-PP7sl positive cells is split in four quartiles based on their Start Time. The

median (●) and 25th to 75th percentiles (line) of the integral of the PP7 trace is plotted

for each quartile. Source data are provided for a.

Fig. 4. Transcription factors control the dynamics and level of mRNA production

a. Violin plots of the trace intensity (maximum of the TS during the transcription period)

for the six promoters after stimulation by 0.2M NaCl. Each dot represents the value

calculated from a single cell. The solid line is the median and the dashed line the mean

of the population. **b.** Comparison between the trace intensity in stimulated (0.2M NaCl)
and unstimulated conditions (0.0M) for the three promoters displaying basal
expression. **c. - f.** Effect of the deletions of the *HOT1* and *SKO1* transcription factor
genes on p*STL1* and p*GPD1* dynamics of transcription (c), cumulative distribution
functions of Start Times (d), the trace intensity (e) and the percentage of responding
cells (f) for the p*STL1*-PP7sl and p*GPD1*-PP7sl reporter strains following a 0.2M NaCl
stress for at least 200 cells. For the p*STL1*-PP7sl *hot1Δ* sample, 349 cells were
analyzed with only 9 displaying a PP7 positive signal. This low number does not allow
to draw a meaningful CDF curve in panel d. Source data are provided for b and c.

Fig. 5. Identification of transcription bursts in stress-induced transcription.

a. Percentage of cells where 1, 2 and 3 or more peaks are identified among the
population of responding cells for the different promoters following a 0.2M NaCl stress.
b. Examples of single-cell traces displaying 1 or 2 peaks for the p*STL1*-PP7sl and the
p*GPD1*-PP7sl reporter strains. **c.** Violin plots representing the Peak Duration. Each dot
represents the value calculated for a single peak. The solid line is the median and the
dashed line the mean of all the peaks measured. **d. - e.** The population of cells was
split between cells displaying one peak and two or more peaks. The Peak Duration (d)
and Trace Intensity (e) are plotted for the p*HSP12*-PP7sl and p*GPD1*-PP7sl strains.
Each dot represents the value calculated for a single peak (d) or a single cell (e). The
solid line is the median and the dashed line the mean of the population.

Fig. 6. Hog1 activity and promoter identity control the shutoff of transcription

a. Violin plots representing the Transcription Period (time difference between End Time
and Start Time) measured for the p*STL1*-PP7sl reporter following 0.1, 0.2 and 0.3M
NaCl stresses. Each dot represents the value calculated from a single cell. The solid
line is the median and the dashed line the mean of the population. **b.** One minus the
cumulative distribution function of End Times for the p*STL1*-PP7sl reporter. The
vertical dashed lines represent the median adaptation time of Hog1 for the three
different stress levels. **c.** Dynamics of the estimated NaCl concentrations in the
medium for the pulse, step and ramp experiment protocols (upper panel, Methods).

Corresponding Hog1 relocation dynamics (middle panel) and p*STL1*-PP7sl
transcription site intensity (lower panel). The mean of at least 180 cells is represented
by the solid line. The shaded areas represent the s.e.m. **d.** Cumulative distribution
function (CDF) of the Start Time for all cells in the pulse, step and ramp experiments.
The CDF at 15 min represents the fraction of responding cells for each condition. **e.**
One minus the cumulative distribution function of End Times only for the responding
cells in the pulse, step and ramp experiments. **f.** Correlation between the Hog1
adaptation time and the PP7 End Time measured in the same cells in the pulse, step
and ramp experiments. The open markers indicate cells where Hog1 has not adapted
at the end of the time lapse. Adaptation time is arbitrarily set to 35 min for this sub-
population. Source data are provided for c.

Fig. 7. Transcription shutoff from different promoters

a. Violin plots representing the Transcription Period measured for the six different
promoters following a 0.2 NaCl stress. Each dot represents the value calculated from
a single cell. The solid line is the median and the dashed line the mean of the
population. **b.** One minus the cumulative distribution function of End Times for the
different promoters. The vertical dashed line represents the median adaptation time of
Hog1 at 0.2M NaCl.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7