Intrinsic neuronal excitability and spontaneous activity underlie cortical abnormalities upon Nr2f1/COUP-TFI deficiency

Isabel del Pino[1°], Chiara Tocco[2*], Elia Magrinelli[2*#], Andrea Marcantoni[3*], Celeste Ferraguto[2], Giulia Tomagra[3], Christian Alfano[2+], Xavier Leinekugel[1&], Andreas Frick[1&] and Michèle Studer[2&@]*

[1]Université de Bordeaux, Inserm, Neurocentre Magendie, Bordeaux, France.

[2]Université Côte d'Azur, CNRS, Inserm, iBV, France

[3]Università di Torino, Dipartimento di Scienza e Tecnologia del Farmaco, Torino, Italy

*co-first authors

[&]co-last senior authors

[°]Present address: Centro de Investigación Príncipe Felipe, Valencia, Spain

[#]Present address: Département des Neurosciences Fondamentales, Université de Lausanne, Switzerland

[+]Present address: GIGA Stem Cells, University of Liège, Liège, Belgium

[@]Corresponding author:

Michèle Studer

iBV - Institut de Biologie Valrose

Univ. Nice Sophia Antipolis

Centre de Biochimie ; UFR Sciences

Parc Valrose, 28 avenue Valrose

06108 Nice Cedex 2

France

Tel.: +33 489150720

e-mail: Michele.studer@unice.fr

ABSTRACT

The formation of functionally different areas in the cerebral cortex results from a
timely regulated interaction between intrinsic genetic mechanisms and electrical
activity. However, whether and how transcriptional regulation influences
spontaneous activity intrinsic to the neocortex during critical periods of
corticogenesis is poorly elucidated. Here, we used mice deficient for cortical *Nr2f1*
(also known as *COUP-TFI*) to investigate how a genetic-determinant of
somatosensory area-identity influences immature neural excitability and network
activity generated within the neocortex. We found that loss of *Nr2f1* alters intrinsic
excitability and morphological features of pyramidal neurons during early postnatal
stages. We also identified voltage-gated ion channels regulated by *Nr2f1* during a
critical time window of somatosensory area specification. Accordingly, *Nr2f1* loss
impacted on spontaneous network activity and synchronization at perinatal stages.
Together, our data suggest that genetic specification of cortical area identity relies
on intrinsic bioelectrical properties regulating patterns of spontaneous activity
intrinsic to the neocortex.

INTRODUCTION

The area- and cell-type specific organization of the mammalian neocortex is a process involving coordinated interactions between cell intrinsic genetic programs and neural activity, an essential feature in specifying the composition and organization of neural circuits during all stages of development (Jabaudon, 2017, Simi and Studer, 2018). Before experience shapes neuronal circuits via sensory-driven activity, the developing cortex is genetically primed to establish patterns of spontaneous activity, which autonomously synchronize large groups of neurons independently of sensory stimulation (Andreae and Burrone, 2018, Kirischuk et al., 2017, Anton-Bolanos et al., 2019). These early patterns of spontaneous activity are thought to set the initial blueprint of cortical network organization. Hence, altering spontaneous activity during development can disrupt formation of functional neural circuits (Kirkby et al., 2013, Li et al., 2013).

In the neocortex, patterns of local spontaneous activity are already present during prenatal and early postnatal stages and contribute to the development of neuronal circuits (Luhmann and Khazipov, 2018). Although cortical areas comprise a variety of circuits, there are nevertheless striking similarities in some aspects of the mechanisms used to generate spontaneous activity. Calcium (Ca^{2+}) transients have been reported already at embryonic day E16, most probably generated in proliferating radial glia cells, as spatially restricted and slowly propagating calcium waves (Kirischuk et al., 2017). Immature neurons express voltage-dependent ion channels, responsible for a highly synchronized spontaneous activity within small neuronal networks (Luhmann et al., 2016). Certain features of immature cells, including high membrane resistance and prominent low-threshold calcium currents, seem to enhance excitability in neurons receiving little afferent sensory input (Moody and Bosma, 2005).

Besides spontaneous activity, the neocortical primordium is also under the control of distinct morphogens that establish proper coordinates along the developing anteroposterior and dorsoventral cortical axes (Greig et al., 2013, O'Leary and Sahara, 2008, Alfano and Studer, 2012). Functioning in a dose-, context- and time-dependent manner, these signaling pathways modulate gradient expression of area patterning genes, such as *Pax6*, *Sp8*, *Emx2* and *Nr2f1* (also known as COUP-TFI), among others. These transcription factors will cooperatively regulate

cell identity specification in cortical progenitors and early postmitotic neurons
building a rough primordial areal map, known as protomap, which will be refined by
sensory-evoked activity during postnatal stages (Simi and Studer, 2018). Alterations
in genetic dosage of these factors alter the size and position of primordial area maps
with severe consequences in cell-type specification and circuit formation (Greig et
al., 2013, Jabaudon, 2017). Although the interaction between patterning genes and
onset of spontaneous activity during primary area specification has been
hypothesized, there is still no clear evidence on how this interaction is perpetrated.

To tackle this question, we used mice deficient for the nuclear receptor *Nr2f1*,
a key transcriptional regulator during cortical development (Armentano et al.,
2007), as a genetic model of area and laminar misspecification. As a member of the
superfamily of steroid hormone receptors, *Nr2f1* represents a well-studied area
patterning gene, which impinges sensory identity predominantly in early
postmitotic cells during neocortical area mapping, and controls neuronal migration
and differentiation (Alfano et al., 2014, Tomassy et al., 2010, Alfano et al., 2011,
Faedo et al., 2008). Haploinsufficiency of the human *NR2F1* gene leads to an
emerging genetic neurodevelopmental disorder called Bosch-Boonstra-Schaaf Optic
Atrophy Syndrome (BBSOAS) (Bertacchi et al., 2019). Patients affected by this
autosomal-dominant disorder exhibit several clinical features, such as intellectual
disability, epileptic seizures, autistic-like traits, affected fine motor coordination and
optic atrophy, among others (Bosch et al., 2014, Chen et al., 2016). In mice, *Nr2f1*
deficiency in the cortex results in impaired primary somatosensory (S1)
specification, delayed radial migration and abnormal laminar specification,
particularly layer V pyramidal neurons, leading to distinct motor and cognitive
impairments (Alfano et al., 2014, Armentano et al., 2007, Tomassy et al., 2010,
Contesse et al., 2019). We employed this mouse model presenting severe area- and
cell-type specific defects within the S1 to investigate whether and how spontaneous
activity intrinsic to the cortex is altered during early stages of neocortical area and
laminar specification. Our data show that glutamatergic pyramidal neurons, and
particularly layer V neurons, undergo changes of intrinsic membrane excitability
along with neuroanatomical and structural alterations within the immature cortex.
Furthermore, we disclosed that altered levels of voltage-gated ion channels, and
particularly HCN1, transcriptionally controlled by Nr2f1, underlie an aberrant state

of neural excitability in the early postnatal neocortex. Finally, we found that alterations observed at the electrophysiological, molecular and morphological levels correlate with altered Ca^{2+} bursts, network firing activity and synchronization events of *Nr2f1* mutant cortices at perinatal stages, before sensory evoked activity has taken place. Together, our data suggest that Nr2f1 controls cortical area patterning through the modulation of spontaneous activity intrinsic to the neocortex. This newly described function is accomplished by the transcriptional modulation of a plethora of channels involved in modulating intrinsic excitability as well as morphological and structural features of layer V pyramidal neurons.

RESULTS

Abnormal intrinsic electrophysiological properties in early postnatal *Nr2f1*-deficient cortical layer V pyramidal neurons

To address whether neural activity intrinsically generated within the neocortex could be regulated by area patterning genes such as *Nr2f1*, we employed a mutant mouse in which *Nr2f1* is exclusively deleted from the cortex without affecting thalamic differentiation (named thereof *Nr2f1cKO*) (Armentano et al., 2007). As intrinsic electrophysiological properties are major determinants of spontaneous neural activity, we first examined the state of intrinsic excitability of layer V glutamatergic pyramidal neurons (LVPNs), described to be abnormally specified in *Nr2f1cKO* mouse somatosensory cortices (Tomassy et al., 2010, Harb et al., 2016), through whole-cell recordings. We employed acute brain slices from *Ctrl* and *Nr2f1cKO* mouse pups during postnatal day (P)5–P8, as critical time points of early cortical circuit establishment. To assess for changes in intrinsic electrophysiological properties in a cell type-specific manner, we selected those neurons that were identified as LVPNs through *post-hoc* biocytin-filling, immunohistochemistry and morphological reconstruction. From 57 neurons recorded, we could identify 19 *Ctrl* and 28 *Nr2f1cKO* as neurons allocated in the layer V (**Figure 1**). All LVPNs from *Ctrl* and mutant brains displayed a slow-adapting regular spiking firing pattern (**Figure 1A**). Multivariate statistical analysis of intrinsic electrophysiological parameters revealed that *Nr2f1cKO* LVPNs clearly segregate from the *Ctrl* population (**Figure 1B)**. Notably, specific changes in passive and active electrophysiological properties

were observed in *Nr2f1cKO* neurons. Mutant LVPNs displayed a significantly more depolarized resting membrane potential, a decreased rheobase and diminished action potential amplitude when compared to *Ctrl* LVPNs. Interestingly, *Nr2f1cKO* LVPNs also exhibited a significantly reduced voltage sag in response to hyperpolarizing current steps, suggesting reduced I_h currents compared to *Ctrl* LVPNs (**Figure 1C**). No significant changes in input resistance, action potential threshold, action potential half-width, inter-spike interval, membrane time constant or input/output function were observed in mutant LVPNs (**Figure 1 – figure supplement 1, table supplement 1**). Taken together, this first functional characterization suggests that Nr2f1 deficiency specific to the developing cortex leads to alterations in intrinsic excitability of immature LVPNs in the prospective somatosensory region.

Reduced dendritic complexity of LVPNs in *Nr2f1* mutant cortices

To assess whether abnormal intrinsic excitability of mutant LVPNs was accompanied by changes in somatodendritic morphology, we reconstructed and analyzed the morphology of biocytin-labelled *Ctrl* and *Nr2f1cKO* P7 LVPNs (**Figure 2A,B**). We first evaluated dendritic arbor geometry and complexity by performing Sholl analysis (**Figure 2C – figure supplement 1A**). Comparison of Sholl profiles showed that *Nr2f1cKO* LVPNs have a significantly reduced proximal dendritic branching (20-100⬚m from the soma) when compared to *Ctrl* LVPNs, whereas minor or no differences were observed in distal regions (**Figure 2 – figure supplement 1A**). Next, to evaluate more specific features of dendrite geometry, we assessed the basal dendrite complexity index (DCI), arbor length, number of primary dendrites, branching points and tips, and oblique formations of basal and apical dendrites (**Figure 2D-H; Figure 2 -figure supplement 1B-D**). Although no significant changes were observed in apical and lateral dendrite geometry between *Ctrls* and *Nr2f1cKOs* (**Figure 2 – figure supplement 1B–D**)*,* our morphological data point to a less elaborated basal dendrite complexity in mutant LVPNs. In particular, the basal DCI was significantly reduced in *Nr2f1cKO* compared to *Ctrl* LVPNs (**Figure 2D–E)**, as a result of a reduced number of primary dendrites, branching points and terminal tips (**Figure 2F–H**). Overall, our data indicate that loss of Nr2f1 function affects the complexity of basal dendrite geometry in LVPNs at early postnatal stages.

Reduced conformation and plasticity properties of the axon initial segment in mutant *Nr2f1* LVPNs

Intrinsic excitability and dendritic changes are often accompanied by variations in structural properties of the axon initial segment (AIS), a specialized structure playing a pivotal role in integration of synaptic inputs and action potential initiation (Yamada and Kuba, 2016, Kole and Stuart, 2012). It has been reported that distinct AIS features, such as length and/or location relative to the soma, are crucial for the homeostatic control of neuronal excitability in layer V pyramidal neurons (Hamada et al., 2016). The observed reduction of basal dendrite complexity, sag response and the presence of more depolarized *Vrest* together with the lack of effect on the input-output relationship (**Figure 1**; **Figure 1 - figure supplement 1)**, led us to hypothesize that structural homeostatic scaling of the AIS could occur in *Nr2f1cKO* LVPNs. This could act as a potential mechanism stabilizing action potential generation, as observed in previous studies (Hamada et al., 2016, Gulledge and Bravo, 2016). To test this hypothesis, we analyzed AIS structure of P7 LVPNs *Ctrl* and *Nr2f1cKO* somatosensory cortices employing the cytoskeletal scaffolding protein Ankyrin G – a widely used marker of the AIS – (Kordeli et al., 1995, Zhou et al., 1998), together with CTIP2, a *bona fide* layer V marker (Arlotta et al., 2005), and NeuN, a neuronal marker labelling the soma (**Figure 3**). Notably, we found a significantly reduced length and diameter of the Ankyrin-positive AIS in *Nr2f1cKO* compared to *Ctrl* LVPNs (**Figure 3A–C**), indicating that altered intrinsic excitability and structural dendritic changes of mutant LVPNs are accompanied with structural changes in AIS conformation (Grubb and Burrone, 2010, Wefelmeyer et al., 2016, Kuba et al., 2010, Kuba, 2010). Decreased length and diameter were maintained in P21 LVPNs (**Figure 3D-F**), suggesting that altered AIS structure observed at P7 is not the result of a delayed neuronal maturation.

Finally, to assess whether *Nr2f1*-deficient LVPNs would still respond to depolarizing input despite their reduced AIS length and diameter, we performed chronic depolarization treatment (+7mM of KCl for 6h) in acute brain slices of *Ctrl* and *Nr2f1cKO* P7 pups (Dumitrescu et al., 2016). While the depolarization treatment reduced, as expected, the relative AIS length of *Ctrl* LVPNs, no significant changes were observed in AIS size between depolarization and control treatment of *Nr2f1*-

215 deficient LVPNs (**Figure 3G, H**). These data unveil that mutant LVPNs are not able to
216 undergo AIS plasticity upon increase in network activity.
217
Altered expression of ion channels and glutamate receptors in postnatal
cortices of *Nr2f1cKO* mutants.
Neuronal intrinsic excitability and AIS functionality of immature neurons are
primarily determined by the expression, subcellular distribution and biophysical
properties of K^+, Na^+ and Ca^{2+} channels. We next conceived that alterations in
intrinsic excitability of *Nr2f1*-deficient LVPNs might correlate with abnormal
expression of voltage gated ion channels. To test this hypothesis, we performed RT-
qPCR for an extensive list of K^+, Na^+ and Ca^{2+} channels specifically expressed in deep
layers, from P7 *Ctrl* and *Nr2f1cKO* whole cortical lysates (**Figure 4; Figure 4 - figure**
supplement 1). Out of 33 ion channels that we could successfully test, we found a
significant reduction in the *mRNA* levels of only *Hcn1*, *Kcnab1*, *Kcnk2*, *Cacna2d1*,
Gria3 and *Grin1* in *Nr2f1cKO* when compared to *Ctrl* brains (**Figure 4A ; Figure 4 -**
figure supplement 1). Changes in expression levels of some relevant voltage-gated
ion channels were subsequently confirmed through western blot analysis in *Ctrl* and
Nr2f1cKO brain lysates. Notably, we found significantly decreased protein levels of
the hyperpolarization-activated cation channel HCN1 (Shah, 2014) in *Nr2f1cKO*
when compared to *Ctrl* cortices (**Figure 4B**). Moreover, double staining of HCN1 and
CTIP2 on P7 cortical sections (**Figure 4C**) confirmed decreased HCN1 protein signal
intensity in CTIP2-expressing LVPNs (**Figure 4D**), supporting reduction of HCN1
protein in layer V somata of *Nr2f1cKO* mutant brains.
Finally, we investigated whether voltage gated ion channels were directly
regulated by Nr2f1. To address this, we searched for evolutionary conserved binding
sites for Nr2f1 on the group of differentially-expressed genes using the ECR browser
tool (Loots and Ovcharenko, 2007, Ovcharenko et al., 2004). We identified conserved
binding sites for Nr2f1 at 25kb of the 5'UTR of *Hcn1*, 3kb of the 5'UTR of *Kcnab1* and
proximal to the 5'UTR of *Kcnd2* loci (**Figure 4E**). To experimentally test direct
regulation of the expression of these ion channels by Nr2f1, we performed
chromatin immunoprecipitation (ChIP) assay on whole newborn *Ctrl* cortices.
Among all the consensus sites identified, binding of Nr2f1 by ChIP unveiled
reproducible binding and significant enrichment only on the *Hcn1* genomic sequence

(**Figure 4F**).

These findings on the molecular landscape of the somatosensory cortex
indicate that while expression levels of several ion channels might be modulated by
Nr2f1 during cortical development, only few ion channels seem to be directly
controlled by Nr2f1. We identified *Hcn1* gene as a direct target, whose protein
product is known to modulate voltage sag and intrinsic excitability (Brager et al.,
2012, Fan et al., 2016), in agreement with our electrophysiological data (**Figure 1**).

Alterations in Ca2$^+$ burst and synchronous activity in early postnatal *Nr2f1*-
deficient cortices.

We next hypothesized that electrophysiological, morphological and structural
alterations together with decreased HCN1 expression levels could translate into
altered patterns of spontaneous cortical network activity (Klueva et al., 2012). To
test this premise, we set up time lapse Ca^{2+} imaging in acute horizontal cortical slices
of *Ctrl* and *Nr2f1cKO* P1 and P4 mouse pups (**Figure 5**). These two stages represent
critical timepoints of early cortical circuit establishment that precede sensory arrival
from thalamic inputs (Corlew et al., 2004). The time course of Ca^{2+} activity from
individual cells was quantified through moving average and further filtered for
bursting cells, thus excluding inactive neurons (**Figure 5A**). When we analyzed the
Ca^{2+} burst frequency (**Figure 5B)**, we observed an overall increase from P1 to P4, in
accordance with previous studies (Corlew et al., 2004). To assess the level of cortical
network maturation, we measured the correlation index of Ca^{2+} bursts for each
experiment, as a way to assess synchronization, by quantifying the number of bursts
across cells in a 3s sliding window (Kirkby et al., 2013). As expected, we observed
increased correlation with network maturation progression (**Figure 5C**).

Next, we determined whether loss of Nr2f1 function would affect Ca^{2+} burst
frequency and synchronization in the developing P1 and P4 cortices. Interestingly,
we observed that Ca^{2+} burst frequency was higher in *Nr2f1cKO* brain slices than in
the corresponding *Ctrl* cohorts at P1, but not at P4 (**Figure 5B**). Moreover, by
comparing the Ca^{2+} burst synchronization events between *Ctrl* and *Nr2f1cKO* brains,
we observed no differences at P1, but instead a decreased synchronization rate in P4
mutant brains (**Figure 5C**). Since synchronization increases with time, these results
suggest an impaired maturation of network complexity in mutant brains. Next, we

pooled experiments by age and genotype and performed a hierarchical clustering using burst frequency and synchronization (**Figure 5D**). The findings indicate that network maturation might be the main factor contributing to differences in *Ctrl* and *Nr2f1cKO* brains since hierarchical clustering provided stronger evidence of data clustering at P1 than at P4.

Our previous data on ion channel transcript differences between the two genotypes support alterations in the endogenous activity of developing sensory areas, most probably affecting the overall frequency at early stages and leading to an asynchronized network at later stages. Taken together, our data indicate that spontaneous network activity and synchronization patterns are altered upon loss of Nr2f1 early after birth, thus deviating from a normal developmental trajectory during neocortical development.

Impaired intrinsic network and bursting activity in cultured *Nr2f1*-deficient cortical neurons.

A recent study shows that subcortical spontaneous activity in the embryonic and postnatal brain contributes to the formation of the S1 cortical map (Anton-Bolanos et al., 2019). We thus aimed to explore whether altered neuronal Ca^{2+} frequency and synchronization events would result exclusively from changes in spontaneous cortical activity intrinsic to the S1 cortex, as opposed to changes in inputs external to S1. To this purpose, we isolated somatosensory cortices from E18.5 *Ctrl* and *Nr2f1cKO* brains, respectively, and cultured the dissociated neurons onto coated multielectrode arrays (MEA). This allowed us to assess network activity during cortical network assembly in an *in vitro Ctrl* and *Nr2f1cKO* embryo-derived model in the absence of extracortical inputs. By recording the firing frequency, number of bursts and their duration at different time points (days in vitro, DIV), we monitored the state of excitability of the neural network (**Figure 6; Figure 6 – figure supplement 1**). Since network maturation correlates with the expression of synchronized events, cross correlation probability between the firing at individual recording sites was used as a quantification parameter of network development *in vitro* (Gavello et al., 2012, Gavello et al., 2018).

Our recordings on primary cultured neuronal networks failed to display any spontaneous electrical activity before 6DIV (*data not shown*). This prompted us to

compare the average firing frequency of the earliest events taking place at 9DIV (**Figure 6A**). We first observed that while *Ctrl* cortical network fired on average at 0.51±0.04 Hz, *Nr2f1cKO* activity was significantly lower (0.30±0.04 Hz) (**Figure 6A, B**). However, despite observing a comparable average number of bursts in *Ctrl* and *Nr2f1cKO* brains (**Figure 6C**), these bursting events were significantly longer in mutant than *Ctrl* neurons (**Figure 6D**), while firing activity was more clustered into bursts in mutant *neurons* (**Figure 6E**). Notably, the firing synchronization index resulted higher in *Nr2f1cKO* than in *Ctrl* neuronal networks (**Figure 6F**). Hence, loss of *Nr2f1* leads to an overall reduced firing frequency, but also to an increased tendency of firing as bursts rather than isolated spikes in 9DIV-derived cortical neurons. Together with the presence of longer burst duration and increased network synchronization, these data point to an involvement of *Nr2f1* in the control of processes that govern neuronal network excitability.

Next, we assessed the rate of firing activity, number of bursts and burst duration in *Ctrl* and *Nr2f1cKO* 15DIV neuronal cultures. At this stage, we observed that the firing frequency of *Ctrl* cortical networks reached a maximum value, which then decreased at 20DIV (*data not shown*). We therefore compared firing parameters measured at 15DIV with those measured at 9DIV to assess whether network maturation was affected *in vitro*. Interestingly, we observed that from 9DIV to 15DIV, the developmental increase in firing activity and in the number of bursts of somatosensory neurons was reduced in *Nr2f1cKO* neuronal networks (**Figure 6 – figure supplement 1A-C**), together with the burst duration, which was significantly decreased in mutant neurons at 15DIV compared to 9DIV (**Figure 6 – figure supplement 1D**). Finally, the percentage of spikes within bursts was also diminished in *Nr2f1cKO* (**Figure 6 – figure supplement 1E**), in accordance with impaired network synchronization, which resulted lower in *Nr2f1cKO* than in *Ctrl* neurons (**Figure 6 – figure supplement 1F**). Taken together, these results indicate that spontaneous network activity generated within the neocortex in the absence of extracortical inputs is affected upon loss of cortical *Nr2f1*. This suggests that *Nr2f1* might be directly or indirectly involved in modulating intrinsic spontaneous network activity in the developing somatosensory cortex.

DISCUSSION

Our work reveals that the area patterning gene *Nr2f1* can influence the emergence of spontaneous activity and intrinsic excitability within the immature somatosensory cortex during early postnatal development. These findings reinforce the role for *Nr2f1* in somatosensory area development, as previously reported (Alfano et al., 2014, Armentano et al., 2007). More importantly, they provide, to our knowledge, first experimental proof for a patterning gene involved in cortical area and cell-type specification that directly determines early patterns of spontaneous network activity intrinsic to the immature cortex. Developmental spontaneous activity has already been described to ultimately influence the formation of long range circuits (Tritsch et al., 2007, Blankenship et al., 2011, Yamamoto and Lopez-Bendito, 2012, Schneggenburger and Rosenmund, 2015, Anton-Bolanos et al., 2019) as well as showing a regionalized pattern in the cortex (Corlew et al., 2004, Uhlen et al., 2015). However, in this study, genetic manipulation targeting exclusively cortical neurons allows to investigate the specific contribution of patterning genes to intrinsic network activity within the neocortex. Our findings provide a primary link between intrinsic genetic determination and spontaneous cortical activity through the regulation of voltage-gated ion channels in pyramidal glutamatergic neurons. Moreover, we also show that the hyperpolarization-activated cation channel HCN1 – known to contribute to intrinsic neural excitability and spontaneous rhythmic activity in the brain (Shah, 2014)– is a potential direct effector of electrical activity regulated by *Nr2f1* during establishment of neuronal synchronicity in the developing neocortex.

Determinants of spontaneous network activity controlled by patterning genes during cortical area identity specification

Normal developmental trajectories in the cerebral cortex are characterized by a sequential process of spontaneous activity patterns in immature networks. In the mouse neocortex, large plateaus of synchronized network activity at birth are followed by early network oscillations and large depolarization potentials along the first postnatal week (Allene et al., 2008, Allene and Cossart, 2010). Multiple aspects that relate to intrinsically regulated maturation of pyramidal neurons might influence the developmental sequence of cortical network activity. Our data provide

a novel characterization at the electrophysiological and molecular levels of intrinsic

excitability features directly or indirectly regulated by the transcriptional regulator

Nr2f1 during the first postnatal week. Our evaluation of how *Nr2f1* expression

influences the molecular determinants relevant for early neuronal network activity

is corroborated by previous transcriptome analyses, in which *Nr2f1* loss in post-

mitotic cortical cells regulate the expression of several activity related genes, such as

Chrna7, Kcnq3, Kcnlp4, Robo1 and *Sv2b* (Alfano et al., 2014). However, the direct

impact of these candidates on immature cortical neuron activity still remains to be

investigated.

Other recent studies also highlighted the role of cell intrinsic and pre-synaptic

factors in determining cortical network development with potential long-term

effects (Boillot et al., 2016, Murase et al., 2016). In addition, a recent report has

shown that the transcription factor Tbr1, known to regulate cortical layer identity,

alters the intrinsic excitability of neocortical neurons, by regulating HCN1

transcription in neonates and increasing sag response in layer VI (Fazel Darbandi et

al., 2018). As Tbr1 is also a key factor for cortical development, it is presumably

regulating spontaneous network activity intrinsic to the neocortex during early

developmental stages, but this still remains to be addressed. Hence, our work sets

the first evidence that spontaneous network activity intrinsic to the cortex is

regulated during early neocortical development by a key area patterning

transcriptional regulator.

Within immature cortical networks, the earliest patterns of correlated

spontaneous activity – e.g. synchronous plateau assemblies – are thought to arise

from voltage-gated intrinsic conductance and gap junction coupling (Pearson et al.,

2004, Roerig and Feller, 2000, Blankenship et al., 2011, Cho and Choi, 2012). Our

work features intrinsic excitability – more precisely, intrinsic bioelectrical

properties of neural membranes – as an important component of neocortical

development, most likely contributing to the abnormal cortical patterning defects

displayed in *Nr2f1* mutants. In this context, it is important to note that *Nr2f1* loss

also affects the expression of many cell-adhesion related genes *(i. e., Cdh4, Cdh11,*

Cdh12, Fat3, and *Ncam2*) (Alfano et al., 2014), which could in turn affect gap-junction

functionality. Interestingly, intrinsic bioelectrical membrane properties prominently

determine the occurrence of membrane potential spikes before gap-junction

coupling takes place (Allene and Cossart, 2010). Hence, it is tempting to speculate

that even earlier forms of activity within the embryonic brain could be responsible

for the abnormal area specification observed in the absence of *Nr2f1* (Armentano et

al., 2007; Alfano et al., 2014).

Nr2f1-dependent *versus* homeostatic changes in intrinsic neural excitability.

Our functional analysis indicates that an abnormally differentiating somatosensory

area displays increased spontaneous network activity at P1 followed by decreased

synchronization within immature neurons at P4. Principal factors for increased

excitability of neural network might be the more depolarized resting membrane

potential of pyramidal glutamatergic neurons, which might contribute to the higher

frequency of Ca^{2+}-events observed at P1. Alternatively, a decreased expression of

voltage sag (I_h) could lead to opposite changes in membrane excitability, which are

strongly age-dependent (Bender and Baram, 2008, Magee, 1998, Williams and

Stuart, 2000, Williams and Stuart, 2003, Berger et al., 2001, Poolos et al., 2002,

Poolos et al., 2006, Wang et al., 2003, Fan et al., 2005, van Welie et al., 2006, Brager

and Johnston, 2007, Chen et al., 2001). Multiple membrane ion channels, which

expression is altered in pyramidal neurons upon loss of *Nr2f1* function, could

contribute to the increase in resting membrane potential in immature neurons, a

property of pyramidal neurons that matures during the two first weeks after birth.

In our *ex vivo* electrophysiological analysis, some membrane properties of *Nr2f1cKO*

LVPNs, such as *Vrest* and *APamp*, resemble those of less mature neurons. However,

other electrophysiological parameters, e.g. R*in*, do not change and therefore are not

in agreement with a delayed maturation in intrinsic properties. In addition, our

analysis on AIS homeostatic plasticity at P21 does not point to a maturation defect in

individual neurons, but rather to a stabilized state of abnormal intrinsic excitability

leading to decreased dendritic complexity and AIS plasticity, as a possible

homeostatic/compensatory mechanism.

Our work using *ex vivo* electrophysiology together with molecular analysis

strongly suggest that *Nr2f1* directly regulates HCN1 expression. It is complex to

precisely dissect changes in membrane-ion channels, priming the start of abnormal

network excitability, from those that are concomitantly changing by intrinsic or

homeostatic plasticity. Since *Nr2f1* directly regulates HCN1 transcription, we

propose that decreased HCN1 levels could act as initiator of the deregulation in intrinsic as well as network activity observed in mutant cortices. Therefore, it is conceivable that the alterations in expression levels we observed in other voltage-gated ion channels at P5–P8 are the product of additional network-activity-driven compensatory mechanisms occurring after P1 or P4. In other words, we hypothesize that changes in ion-channel expression of pyramidal glutamatergic neurons at P5–P8 are due to homeostatic and/or intrinsic plasticity changes following earlier alterations in overall network activity state. For example, experimental evidence from the developing hippocampus suggests that the induction of increased bursting activity and seizures alters persistently HCN channel expression and voltage-dependence of I_h current (Arnold et al., 2019, Chen et al., 2001). Acting as a direct transcriptional regulator of HCN1, we propose that Nr2f1 could influence spontaneous network activity intrinsic to the somatosensory cortex by contributing to the establishment of the bioelectrical state of glutamatergic neurons at the time point, and probably before, the start of cortical network formation. Moreover, intrinsic pacemaker properties through HCN1 expression has been hypothesized to generate and drive distinct synchronized activity patterns in immature networks (Luhmann et al., 2016).

Importantly, genetic or pharmacological dysfunction in HCN channels in mice results in behavioral phenotypes associated to somatosensory-motor coordination, e.g. reduced forelimb reaching accuracy and atypical movements during a single-pellet skill reaching task (Boychuk et al., 2017). This is very analogous to the behavioral phenotypes displayed by *Nr2f1cKO* mice (Tomassy et al., 2010) and symptoms of NR2F1 haploinsufficient patients (Bosch et al., 2014, Chen et al., 2016), suggesting that Nr2f1 might influence cortical maps involving voluntary movements by allowing coordinated skilled behavior *via* the action of HCN1.

Together with the cellular and molecular role of Nr2f1 during intrinsic area patterning, this study has unraveled a key and novel functional role of this transcriptional regulator in cortical intrinsic excitability and spontaneous activity that might affect axonal connectivity and circuit formation necessary for behavioral abnormalities described in mice (Contesse et al., 2019, Flore et al., 2017, Tomassy et al., 2010) and, possibly, in humans (Chen et al., 2016, Bosch et al., 2016), reviewed in (Bertacchi et al., 2019).

ACKNOWLEDGMENTS:

We thank Dr. Marteen Kole and Dr. Nathalie Dehorter for critical reading of the manuscript and members of the Studer, Leinekugel and Frick laboratories for stimulating discussions. We also thank the iBV PRISM Microscopy facility and the Bordeaux Imaging Center (BIC) for their regular support. This work was supported by an ANR (Agence National de la Recherche) grant # ANR-13-BSV4-0011-01 to M.S. and X.L, by the "Fondation Recherche Médicale; Equipe FRM 2015 grant #DEQ20150331750; by an "Investments for the Future" LabEx SIGNALIFE (grant ANR-11- LABX-0028-01) to M.S. and by the "Initiative d´Excellence (IdEx) de l´Université de Bordeaux" fellowship to I.D.P.

Corresponding author statement:

The corresponding author declares non-financial competing interest on behalf of all authors.

MATERIALS & METHODS

Mice

Nr2f1/COUP-TFI*fl/fl* mice were crossed to *Emx1-Cre* mice to inactivate *COUP-TFI/Nr2f1* exclusively in cortical progenitors and their progeny (Armentano et al., 2007). These mice are denominated *Nr2f1cKO* throughout the text. Mice were genotyped as previously described (Armentano et al., 2007). For P21 studies, Control (*Nr2f1 fl/fl)* and mutant *Nr2f1cKO* mice were crossed to *Thy1-eYFP-H* mice to specifically mark LVPNs, as previously reported (Harb et al., 2016). Midday of the day of the vaginal plug was considered as embryonic day 0.5 (E0.5). Control (*Nr2f1 fl/fl)* and mutant *Nr2f1cKO* mice were predominantly bred in a C57BL6 background. Both male and female embryos and pups were used in this study. All mouse experiments were conducted according to national and international guidelines, with authorization #15349 and 15350 by the French Ministry of Education, Research and Innovation (MESRI) and upon request of the local ethics committee – CIEPAL NCE/2019-548 (Nice) and CEEA50 (Bordeaux), and in accordance with the guidelines established by the Italian National Council on Animal Care and approved by the local Animal Care Committee of University of Turin (Authorization DGSAF 0011710-P-26/07/2017).

Ex vivo electrophysiology and analysis

Mouse pups at postnatal stage (P)5–P8 were decapitated and brains were placed on ice-cold modified artificial cerebrospinal fluid (aCSF) containing (in mM): 87 NaCl, 25 $NaHCO_3$, 5 Glucose, 65 Sucrose, 2.5 KCl, 1.25 NaH_2PO_4, 0.5 $CaCl_2$, 7 $MgCl_2$ saturated with 95% CO_2 and 5% O_2. Coronal slices of 300μm were cut in a vibratome (Vibratome 300 Plus, Sectioning Systems) and gently transferred to room temperature aCSF containing (in mM): 125 NaCl, 25 $NaHCO_3$, 25 Glucose, 2.5 KCl, 1.25 NaH_2PO_4, 2 $CaCl_2$, 1 $MgCl_2$ saturated with 95% CO_2 and 5% O_2 (pH 7.4) .

Current-clamp recordings were performed at 32-34°C using a potassium-gluconate based intracellular solution containing (in mM): 120 K-Gluconate, 20 HEPES, 0.5 EGTA, 15 KCl, 4 MgATP and 0.3 NaGTP (pH 7.4) and osmolarity set to 299 mosmol l^{-1}. Biocytin (1.5 – 2 mg ml^{-1}) was added to the intracellular solution for post-hoc immunohistochemistry. Neurons were visualized with an upright microscope (Zeiss Axio Examiner D1) equipped with an Evolve 512 EMCCD Camera

(Photometrics) using a 63X/1.0nA water-immersion objective (Zeiss) and infrared DodT contrast imaging. Recording pipettes (6-10MΩ) were pulled from borosilicate glass using a Narishige P-10 puller. Data was acquired using a Dagan BVC-700A Amplifier, low-pass filtered at 3kHz and sampled at 20kHz with ITC16 (Instrutech) and AxoGraph X software. Membrane voltage was recorded in the bridge mode. To determine the excitability of pyramidal neurons, we performed 800ms square current steps of 10pA from -100pA to +150pA. The rheobase was defined as the minimum somatic current required to elicit an action potential (AP) from the resting membrane potential of the neuron. Input resistance was calculated from voltage responses to -20pA. Threshold potential was defined as dV/dt = 10mV/ms. Inter-spike interval (ISI) was calculated as the difference between first and second AP at the minimal current to elicit >2AP. The amplitude of action potentials was measured from both restingmembrane potential to peak and also from spike threshold (see Table 1). Action potential width was measured at half of the maximum amplitude (APhalf-width or APh-w). Voltage sag was calculated as the difference between steady-state and minimum voltage transient resulting from hyperpolarizing (-150pA) current injections.

Post-hoc immunofluorescence

Brain slices were fixed overnight at 4°C with 4% paraformaldehyde in 0.1M phosphate buffer (PB) pH7.4. The following day slices were washed 3 x 10 min with 0.1M PB and incubated overnight at 4°C with Alexa-Flour®-conjugated streptavidin diluted 1:500 in 0.1M PB. The following day slices were washed 3 x 10 min with 0.1M PB, stained with DAPI and mounted with Mowiol.

KCl *ex vivo* depolarization assay

Ctrl and *Nr2f1cKO* P7 mice were sacrificed by decapitation and brains dissected into ice-cold aCSF saturated with 95% O_2 and 5% CO_2. 300μm-thick slices were produced by vibratome sectioning and then transferred into homemade floating nests immersed into aCSF, kept at room temperature and constantly saturated with 95% O_2 and 5% CO_2. After 15 minutes of acclimatization, slices from both genotypes were split into two groups: an untreated group (**- KCl**) and an experimental group (**+ KCl**). KCl were added to the aCSF of the experimental group whereas control slices were

kept in normal aCSF. After an incubation time of 6 hours(Dumitrescu et al., 2016),

slices were washed for 15 minutes in aCSF and then processed for

immunofluorescence (IF), as described below.

Histology and immunohistochemistry of mouse perinatal cortices

Perinatal mice (P7-8) were anesthetized by using a mixture of Tiletamine-

Zolazepam-Xylazine-Buprenorphine (TZXB) and intracardially perfused with

Phosphate Buffer Saline (PBS) followed by 4% paraformaldehyde (PFA) in PBS. Then,

brains were dissected and post-fixed for 4h at 4°C in 4% PFA and then vibratome-

sectioned to obtain 100⊠m thick, coronal or sagittal floating sections. Overnight

(ON) incubation at 4°C in a solution of 0.5% Triton X-100, 3% BSA, 10% Goat Serum

in PBS was performed to permeabilize the sections and reduce non-specific binding

of the antibodies. For immunofluorescence (IF), sections were incubated for two ONs

at 4°C with indicated primary antibodies in a solution of 0.5% Triton X-100, 10%

Goat Serum in PBS and then ON at 4°C with relative secondary antibodies (see **Key**

resource table) and HOECHST diluted in PBS. Regarding the HCN1 IF, brains were

briefly post-fixed for 30 minutes (min) at 4°C in 4% PFA and then immediately

sectioned with the vibratome to obtain 50μm thick coronal floating sections.

Permeabilization-blocking step was reduced to 2h, primary antibody incubation to

one ON and secondary antibody and HOECHST incubation to 2h. Sections were then

transferred on superfrost plus slides (ThermoScientific), let dry for 30min to 1h and

finally mounted with Mowiol mounting medium.

Microscopy and image analysis

Imaging was performed using a Zeiss 710 confocal microscope equipped with a

405nm diode, an argon ion, a 561nm DPSS and a 647 HeNe lasers. Z-stacks of fixed

cortical sections were imaged using a LD-LCI Plan-Apo 25x/0.8 NA or a LD Plan-Apo

63x/1.4 NA oil immersion lenses. Images from IF experiments were analyzed by

using Fiji-ImageJ Software (Schindelin et al., 2012): cell counter plug-in was used to

assess the number of cells expressing specific proteins, while measurement tools

were used to determine signal intensity or Axonal Initial Segment (AIS) features

such as length and distance from the soma. For the AIS average diameter

measurement, we took advantage of IMARIS software's tool filament tracer. AISs

were semi-automatically reconstructed and an average diameter automatically
measured for every AIS individually.

Morphological characterization of biocytin filled layer V pyramidal neurons
(LVPNs) was analyzed by using IMARIS software. Cell arborization was
reconstructed in a semi-automatic manner, via the filament tracer module.
Automatic length measuring tool was used to calculate the total dendritic length and
the basal dendritic length. Branching points, branching point orders and tips were
manually quantified for both basal and apical dendrites, as well as primary and
lateral dendrites. The branch tip order is an integer value equivalent to the number
of branching points a dendrite undergoes from its somatic origin to the terminal tip
(see also Figure 2D for a graphic representation). Primary dendrites were identified
as directly originating from the soma, lateral dendrites were categorized as
originating from and perpendicular to the apical dendrite whereas total and basal
branch tips were identified as the terminal ends of primary dendrites including or
excluding the apical dendrite arborization respectively.

The basal Dendritic Complexity Index (bDCI) was calculated as follow$s$:

$$bDCI = \frac{\sum \text{basal branch tip orders} + \text{basal branch tips}}{\text{Primary dendrites}} \text{Basal Arbor Length}$$

An automatic detection of Sholl intersections was used to compute Sholl profiles for
every individual cells. Sholl intersections were identified as the number of dendrite
intersections for concentric spheres of increasing radius and having as a center the
centroid of the cell body. Distance between radii was set at 10μm.

RNA extraction and cDNA synthesis
RNA for reverse transcriptase–PCRs (RT–PCRs) was extracted from P0 and P7
dissected neocortices separated from meninges of *Ctrl* and *Nr2f1cKO* mice (n=3) and
was extracted using the RNeasy Mini kit (Qiagen) following manufacturer's
instructions. We synthetized cDNA from 1μg of total RNA using Superscript III First-
Strand Synthesis System for RT–PCR (Invitrogen) following manufacturer's
instructions.

qPCR RNA/cDNA quantification
qPCRs on cDNA was performed using KAPA Sybr Fast (Kapa Biosystems) according

to manufacturer's instructions. Samples were run on a LigthCycler II 480 (Roche)

and the following primers were used: GAPDH Fw: GTATGACTGCACTCACGGCAAA.

Rev: TTCCCATTCTCGGCCTTG; Cacna1A Fw: GGGCCCAAGAGATGTTCCAG. Rev:

TCCACAGACTGGGAGTTGGG; Cacna1B Fw: CACGCAGAGGACCCACGATG. Rev:

CATCACAGCCAGTGTTCCTG; Cacna1G Fw: TCAGCTGCCTGTCAACTCCC. Rev:

CCCATCACCATCCACACTGG; Cacna1H Fw: ACCTACACAGGCCCGGTCAC. Rev:

ATGGGACCTGGAAGGAGGTG; Cacna2D1 Fw: CAGGGAGGGCACTGATCTTC. Rev:

TTGATAGTGACGGGCGAAGG; Cacna2D3 Fw: TTAGGTGTGGCGCTCTCCAG. Rev:

GCCAAGGACACGTCAGGATG; Hcn1 Fw: TGCCACAGCTTTGATCCAGT. Rev:

GCGCATGTCAGCTGGTAACT; Kcnab1 Fw: TCCCATGGAAGAAATCGTTCG. Rev:

CTTCCATGATCTCCATCGCG; Kcnd2 Fw: GGGAAGCCATAGAGGCAGTG. Rev:

GTGGAGTTCTCTCCACACATCTG; Kcng3 Fw: CAACCGCAGTCTGGATGACC. Rev:

CGGTGAACCACCCTATGCAG; Kcnh1: Fw: TTAGTGCCTTTATGGGTGATCC. Rev:

GATGCCCTGACTCTCTCTCC; Kcnh5: Fw: CTTTCCAGACCCAGGATGCC. Rev:

TTGATTCACTGGAGCGCCTG; Kcnj6 Fw: AACTGACGGAGAGGAATGGG. Rev:

TGGGTTGGGTGAATTCTGGG; Kcnk2 Fw: ACGTGGCAGGTGGATCAGAC. Rev:

AGTAGGCCAGCCCAACGAG; Kcnmb4 Fw: TTGGAAAGATGAGATCGGTTCC. Rev:

AAGCAGTGCAGGAGAGCAATC; Kv2.1 Fw: CGTCATCGCCATCTCTCATG; Rev:

CAGCCCACTCTCTCACTAGCAA; Gria1 Fw: CTGGTGGTGGTGGACTGTG. Rev:

TGTCCATGAAGCCCAGGTTG; Gria2 Fw: ATTTGGAATGGTATGGTTGGAG. Rev:

AGGCTCATGAATGGCTTCGAG; Gria3 Fw: TCATTCTCACGGAGGATTCCC. Rev:

CACAGCAAAGCGGAAAGCAC; Gria4 Fw: CCAGAGCCTCCCTGAAGACC. Rev:

TGCGCGCTCTCCTCTCTTTC; Grik1 Fw: AGTTGGTCGCATGCTCTTCG. Rev:

CGAACCTTGATGCCCAATAC; Grik2 Fw: CTGGTGGAGAGTGCTTTGGG. Rev:

CCACACCCTTGCAACCCTG; Grik3 Fw: TCGCCAGATTCAGCCCTTAC. Rev:

GCCCATTCCAAACCAGAAGC; Grik5 Fw: GCCTGCGGTTGGTAGAGGAC. Rev:

TCTGCCTTCCGGTTGATGAG; Grin1 Fw: TCAGAGCACACTGTGGCTGC. Rev:

ATCGGCCAAAGGGACTGAAG; Grin2B Fw: GGAAGTGGGAGAGGGTGGG. Rev:

CAGGACACATTCGAGGCCAC. For each reaction primers were used at 0.2μM using

10nl of total cDNA. Each sample was corrected to the housekeeping gene GAPDH.

Dissection and western blot of mouse perinatal cortices

660 Postnatal pups (P7-8) were sacrificed by decapitation and fresh tissues were

dissected immediately after. Whole cortex samples were collected and lysed in RIPA buffer (10 mM Tris-Cl (pH 8.0), 1 mM EDTA, 1% Triton X-100, 0.1% sodium Deoxycholate, 0.1% SDS, 140 mM NaCl, 1 mM PMSF) implemented with cOmplete protease inhibitors (Roche). Protein quantity was then measured with the Pierce BCA protein assay KIT (ThermoScientific). Equal amounts of proteins from tissue lysates were resolved by reducing SDS-PAGE, transferred to PolyVinylidene DiFluoride (PVDF) membrane and incubated with the indicated antibodies (see supplementary Table 2). Immunoblots were developed using ECL Prime western blotting detection reagent (GE Healthcare). Signals were normalized on actin signal intensity.

Chromatin Immuno-Precipitation (ChIP)

The ChIP protocol was modified from Kuo and Allis 1999 (Methods), using protein sepharose-A resin (SIGMA), prepared by washing in PBS and equilibration buffer (HEPES pH7.4 20mM, EDTA 1mM, NaCl 150mM, Triton 0.8%, SDS 0.1% PMSF 10mg/ml). For each experiment (n=3), we dissected necortices of 8 mice at P0 in ice cold HBSS and washed in DMEM. Samples were crosslinked in 0.9 formaldehyde for 10' and blocked adding 1.1ml of glycine 1.25M for 5'. We washed samples in wash buffer (HEPES pH 7.4 20 mM, NaCl 150mM, Glycine 0.125M, PMSF 10mg/ml) and isolated nuclei in Lysis buffer (HEPES pH 7.4 20mM, NaCl 150mM, Glycine 0.125M, SDS 1% PMSF 10mg/ml). Nuclei were collected (2200 rcf for 5' spin) and sonicated in 1.2 ml Sonication Buffer (HEPES pH 7.4 20mM, NaCl 150mM, Glycine 0.125M, SDS 0.4% PMSF 10mg/ml) 6 times for 5 seconds at 10µm amplitude. DNA fragments were separated from cellular debris by spinning at 14000rcf for 10' and recuperating the surnatant, fragmentation quality was evaluated by gel electrophoresis post-hoc. DNA fragments where split in 3x300 µl samples and precleared with resin Slurry solution (resin/equilibration buffer, 50/50 volume), adding 50ul of Slurry to each and incubating for 1h at 4°C. Surnatant was kept after spinning at 800 rcf for 2' at +4°C and were incubated overnight at +4°C with 3µg of dedicated antibodies: Nr2f1 antibody (Thermo Fisher PA5-21611), GFP antibody (Abcam ab13970, for aspecific binding), no antibody. A fourth samples called Mock was obtained using a blank volume of Sonication buffer. We added 50µl of Slurry to each sample and incubated 3h at +4°C and washed samples 4 times in ice cold

Washing buffer A (HEPES pH 7.4 20 mM, NaCl 150mM, Triton 100 1%, SDS 0.1%

PMSF 10mg/ml), 4 times in ice cold Washing buffer B (Tris pH 8.0 20mM, EDTA 1

mM, LiCl 250mM, NP-40 0.5%, Na-Deoxycholate 0.5%, PMSF 10mg/ml) and 2 times

with TE buffer (Tris HCl pH 8.0 20mM, EDTA 1mM, PMSF 10mg/ml). We eluted DNA

fragments denaturating antibodies adding 300µl of Elution buffer (NaHCO3 50mM,

1% SDS) and incubating in rotation 1h at RT. An Input samples was created here

mixing 300µl of Elution buffer to 10µl of the supernatant from the first washing of

the no-antibody sample. We separated the fragment from the resin ant antibodies

spinning at 8000 rcf for 2' and saving the supernatant, twice. Next, we reversed

crosslinks adding 12µl of NaCl 5M and incubating overnight at +65°C. RNA was

removed by adding 1µl of RNAse-A 10mg/ml and incubated for 30' at +37°C; the

reaction was blocked adding 6µl of EDTA 0.5M. Remaining proteins were digested

adding 12µl of Tris HCl pH 8.0 and 1µl of proteinase K 10mg/ml for 1-2h at +45°C.

Finally, samples were purified by phenol-chloroform extraction and precipitated

into 20µl of H_2Omq.

qPCR of ChIP samples

qPCRs on ChIP was performed using KAPA Sybr Fast (Kapa Biosystems) according to

manufacturer's instructions. Samples were run on a LigthCycler II 480 (Roche) and

the following primers were used: Nr2f1 binding site at Hcn1 locus Fw:

TTTCTGCTTTCCCGTGAGAGC. Rev: AGCCTAGCCTAGGAAGCCAG; Nr2f1 binding site at

Kcnab1 locus Fw: TGTGGTATCAGAGATGAAAGGC. Rev: AAGGGACCTCCCAGCTCTTC;

Nr2f1 binding site at Kcnk2 locus Fw: CCCAAAGCTGGACTTGTGTAC. Rev:

TTCCCCTGATCAAAATTCCATTC; Negative control (Egr1 binding site at Cacna1g

locus) Fw: AGCACCAGCTCAGATCACTCC. Rev: CAGAAGAAACTGTGTCAAGAGC. For

each reaction primers were used at 0.2µM using 0.5µl each sample. For proper

quantification, a calibration curve was included in the reactions using 6 sequential

1:10 solutions of samples obtained from amplicon extraction of PCR performed on

genomic DNA with each set of primers used. For every ChIP experiment the reaction

was carried out on 3 replicates of specific antibody, aspecific antibody, no antibody,

mock and input samples. The calculated concentration of every samples was

normalized to the input, set to 1 and we then calculated the fold change between the

samples with specific antibody against sample with aspecific antibody, which was

normalized to 1. Finally, the fold change of specific/aspecific antibody samples was

corrected by dividing for the fold change of specific/aspecific antibody samples

obtained from the negative control primers. Regarding the qPCR statistical analysis,

we performed paired t.test analysis correcting the P-value for multiple testing using

the Benjamini–Hochberg procedure, false discovery rate = 0.25. Data analysis scripts

were custom-prepared in R (Computing, 2018) and Matlab.

Ca^{2+} imaging from *ex vivo* brain slices preparation and analysis.

Animals were anesthetized by cold and brains were dissected in ice cold high-Mg

ACSF (125mM NaCl, 25mM NaHCO$_3$, 2.5mM KCl, 7mM MgCl$_2$, 2mM CaCl$_2$, 1.25mM

NaH$_2$PO$_4$, 10mM glucose, pH7.4) after decapitation. Horizontal 300 μm slices were

prepared on a Leica Vibratome 100M in ice cold High Mg ACSF under 95% O$_2$, 5%

CO$_2$ bubbling. Selected slices where incubated 1h at 37°C in ACSF (125mM NaCl,

25mM NaHCO$_3$, 2.5mM KCl, 2mM CaCl$_2$, 1.25mM NaH$_2$PO$_4$, 10mM glucose, pH7.4)

0.63% Oregon Green BAPTA™ (ThermoFisher O6807) in DMSO before proceeding to

image acquisition. Slices were loaded on a 35mm diameter petri dish with 1.3mm

glass bottom and immobilized with a custom nylon net. After 30 minutes of

equilibration in the microscope objective holder time lapse images were acquired

over a Spinning Disk microscope (Olympus / Andor technology) with a 40x air

objective, recording for 15 minutes at 150 ms intervals twice over M and S area for

each sample.

Images were automatically corrected for drifting using built-in Fiji

(Schindelin et al., 2012) function and cells profiles were manually defined as

separate ROI on a custom Matlab script. Single ROI were used to acquire BAPTA

fluorescence intensity as the mean of intensity within each area defined in each ROI

for every frame, thus resulting in single cell time series of Ca^{2+} signals. Each of the

these were individually corrected by moving average (Anderson, 1975) to generate a

constant baseline over the progressive signal loss. Actively bursting cells were

selected and considered for further analysis if in at least 40% of frames of an ROI fell

within the first quarter of maximum intensity of the ROI, thus excluding inactive and

fast spiking interneurons, concentrating the analysis on projection neurons. Within

selected timeseries, Ca bursts were detected filtering BAPTA intensity on a threshold

of 3 standard deviations. The bursting frequency of every selected cell was measured

dividing the number of bursts above threshold by the sum of intervals between each

burst. To evaluate coordination of activity, every selected time series in a recording

was vertically evaluated, counting the number of Ca bursts happening in the same 3s

time window across time series and normalizing by the total number of bursts

within the recording, thus obtaining a coordination index, which was then

normalized across in each image by dividing for the total number of Ca burst

multiplied by the total number selected time series, thus obtaining a single

correlation index per image. All time-lapse analysis was performed using a custom

Matlab script.

MEA recordings on primary cultures and analysis of MEA activity.

Cortical neurons were obtained from E18.5 *Ctrl* and *Nr2f1cKO* mouse embryos.

Cortices were rapidly dissected under sterile conditions, kept in cold HBSS (4ºC)

with high glucose, and then digested with papain (0.5 mg/ml) dissolved in HBSS plus

DNAse (0.1 mg/ml), as previously described (Gavello et al., 2018). Isolated cells

were then plated at the final density of 1200 cells/mm^2 onto the MEA (previously

coated with poly-DL-lysine and laminin). Cells were incubated with 1 %

penicillin/streptomycin, 1% glutamax, 2.5% foetal bovine serum, 2% B-27

supplemented neurobasal medium in a humidified 5 % CO_2 atmosphere at 37ºC.

Each MEA dish was covered with a fluorinated ethylene-propylene membrane (ALA

scientific, Westbury, NY, USA) to reduce medium evaporation and maintain sterility,

thus allowing repeated recordings from the same chip.

Multisite extracellular recordings from 60 electrodes were performed using

the MEA-system, purchased from Multi-Channel Systems (Reutlingen, Germany).

Data acquisition was controlled through MC_Rack software (Multi-Channel Systems

Reutlingen, Germany), setting the threshold for spike detection at -30 µV and

sampling at 10 kHz. Experiments were performed in a non-humidified incubator at

37°C and with 5% CO_2, without replacing the culture medium. Before starting the

experiments, cells were allowed to stabilize in the non-humified incubator for 90

seconds; then the spontaneous activity was recorded for 2 minutes.

Bursts analysis was performed using Neuroexplorer software (Nex

Technologies, Littleton, MA, USA) after spike sorting operations. A burst consists of a

group of spikes with decreasing amplitude, thus we set a threshold of at least 3

spikes and a minimum burst duration of 10ms. We set interval algorithm specifications such as maximum interval to start burst (0.17 sec) and maximum interval to end burst (0.3 sec) (Gavello et al., 2012, Gavello et al., 2018). Burst analysis was performed by monitoring the following parameters: mean frequency, number of bursts and mean burst duration. Cross correlation probability values were obtained by means of Neuroexplorer software using ± 2 s and 5 ms bin size. The number of experiments is referred to the number of MEA each of those consisted of 59 recording electrodes.

Statistical tests.

Principal component analysis (PCA) was performed on R software using Factominer and the following parameters: V$rest$, Rin, Rheobase, V$threshold$, APamplitude, APhalf-width, latency, Sag and ISI. In PCA plots, ellipses represent the 95% confidence and the square the center of mass for each cluster. Hierarchical clustering was performed with Euclidean distance, using both average burst frequency and correlation per condition. All statistical analysis scripts were custom-prepared in R (Computing, 2018). Additional statistical analyses were also performed using Graphpad Prism7 and data are presented as mean ± s.e.m. *n* values represent biological replicates from > 3 different brains and > 3 different litters. No statistical tests were used to predetermine sample size but samples sizes were similar as reported in previous publications (Del Pino et al., 2017, Del Pino et al., 2013) or as generally employed in the field. No randomization was used. Whenever possible both genotypes were processed in parallel. *Ex vivo* electrophysiology experiments and analysis of all data were performed blind to genotype. Differences were considered significant when P < 0.05. Data sets were tested for normality (Kolmogorov-Smirnov) or homoscedasticity (Leven test) before performing parametric (Student t-test, one-way or 2-way ANOVA followed by Bonferroni's post-hoc test) or non-parametric tests (Welch's unequal variances t-test or Mann-Whitney test) used to determine p-values. In Figure 5B, P1 and P4 were analyzed independently, testing *Ctrl* and *Nr2f1cKO* frequency and correlation score by Welch Two-samples test.

FIGURE LEGENDS :

Figure 1. Nr2f1 deletion in the cerebral cortex results in abnormal intrinsic excitability of layer V pyramidal neurons. **A**) Layer V pyramidal neurons (LVPNs) identified through fluorescent labelling for biocytin (green) and DAPI (gray) in *Ctrl* (in black) and *Nr2f1cKO* (in red) somatosensory cortex display slow-adapting regular spiking firing pattern. **B**) Individual factor maps (PCA) performed on pyramidal neurons recorded segregate two clusters as distinct populations in *Ctrl* (black) and *Nr2f1cKO* (red). **C**) Mean V*rest* is significantly increased (** P = 0.0013), while Sag (* P = 0.0103) and the rheobase (*** P = 0.0006) are significantly decreased in *Nr2f1cKO* LVPNs compared to *Ctrl* LVPNs. (N = 19 *Ctrl*, N = 28 *Nr2f1cKO*, two-tailed unpaired T-test or U Mann-Whitney test). V, cortical layer V, V*rest*, resting membrane potential; APamp, action potential amplitude. Scale bar: 100 μm. ***See also Figure supplement 1 and table supplement 1.***

Figure 2. Reduced basal dendritic complexity of layer V pyramidal neurons (LVPNs) in Nr2f1 mutant cortices. *R*epresentative images of *Ctrl* **(A)** and *Nr2f1cKO* **(B)** LVPNs. In grayscale, maximum intensity projection of a multi-stack confocal image; in green, correspondent 3D reconstructions. On the right, magnifications of the basal structures. Scale bar: 50 ▯m. **C)** Sholl analysis points to a reduction of the arborization complexity of *Nr2f1cKO* LVPNs (** P = 0.002; **** P < 0.0001). **D**) Basal Dendrite Complexity Index (DCI) formula and schematic representation of primary dendrites, branching points, terminal tips and tip orders. **E-H**) Average basal DCI (** P = 0.0019), number of primary dendrites (* P = 0.0159)**,** basal dendrites branching point (*** P = 0.0004) and terminal tips (*** P = 0.0009) are statistically reduced in mutant cells compared to controls. (N = 15 *Ctrl*, N = 28 *Nr2f1cKO*). P-values are calculated by 2way ANOVA test **(C)**, Mann-Whitney non parametric test **(E)** or Welch's unequal variances t-test **(F-H).** ***See also Figure supplement 1.***

Figure 3. Abnormal structural features of the axon initial segment (AIS) in mutant Nr2f1 layer V pyramidal neurons. **A**) Left panel, coronal sections of P7 somatosensory *Ctrl* and corresponding *Nr2f1cKO* cortices labelled with CTIP2, for layer V identification, Ankyrin G for the AIS and NeuN as a somatic marker. Scale

bar: 100 ▢m. Right panel confocal images of triple stained *Ctrl* and *Nr2f1cKO* LVPNs.

Scale bar: 10 ▢m. **B-C**) Average AIS length (* P=0.0129) and diameter (** P = 0.0043)

are statistically reduced in *Nr2fcKO* neurons compared to *Ctrl* at P7. N=3. **D**)

Confocal images of *Ctrl* (left) and *Nr2f1cKO* (right) LVPNs from P21 somatosensory

cortices. In green Thy1-eYFP reporter gene, in red Ankyrin G. **E-F**) Average AIS

length (**** P < 0.0001) and diameter (**** P < 0.0001) are statistically reduced in

Nr2f1cKO cells compared to controls at P21. N=3. **G**) Confocal images of control and

Nr2f1cKO LVPNs non-treated (-KCl) and treated (+KCl) with high KCl aCSF. In green

CTIP2, in red Ankyrin G. **H**) Average relative length (average length values were

normalized on control values) are significantly reduced in controls after KCl

treatment (**** P ▢▢0.0001). No differences were observed in *Nr2f1cKO* cells

between non-treated and treated populations. (ns P=0.289). P-values are quantified

by the Welch's unequal variance t-test (**B**), Unpaired t-test (**C-F**) or 2way ANOVA test

(**H**).

Figure 4. Nr2f1 regulates the expression of a subset of ion channels. **A**) Data plot

of qPCR analysis on P7 *Ctrl* and *Nr2f1cKO* P7 cortices. N≥3. The red line represents

the threshold. Significantly downregulated genes are depicted in the upper-left

panel. **B**) Western-blot analysis of HCN1 protein shows a significant reduction in

Nr2f1cKO P7 cortices (*P=0.05). N=4. **C**) CTIP2-HCN1 double IF staining of P7

somatosensory cortices. **D**) Dot plot distribution of HCN1 cell signal intensity points

to a reduction of HCN1 in *Nr2f1cKO* LVPNs (**** P < 0.0001). N=3. **E**) Snapshots from

Evolutionary Conserved Regions (ECR) browser showing Nr2f1 binding location and

sites (in pink); intergenic regions (red); UTR (yellow); exons (blue); introns

(orange). **F**) Graph showing the ratio between unspecific and specific antibody (a.b.)

fold changes (f.c.) on Nr2f1 consensus sites. N=3. P-values are calculated by Welch's

unequal variances t-test (B-E), unpaired (A) or paired (F) t-test followed by

Benjamini–Hochberg multiple test correction, false discovery rate = 0.25. ***See also***

figure supplement 1*.*

Figure 5. Altered Ca²⁺-dependent intrinsic activity upon Nr2f1 loss. **A**) Schematic

and representative images of Ca²⁺ BAPTA fluorescent signals in an acute horizontal

slice preparation from somatosensory cortex. On the right, time series of

intracellular signal in manually defined cells, corrected by moving average (Fi).

Bursting events are defined by distinct thresholds. Below, time series matrices of

above threshold Ca^{2+} signals with interneuron unsupervised hierarchical clustering

(left). Interneuron Pearson correlation matrices (right). P1 and P4 were analyzed

independently, *Ctrl* and *Nr2f1cKO* frequency and correlation score quantified by

Welch Two-samples test. hierarchical clustering performed with Euclidean distance,

using average, burst frequency and correlation per condition. Statistical scripts were

custom-prepared in R (Computing, 2018). **B, C**) Distribution of single cell Ca^{2+} burst

frequency (**B**) and burst intra-neuron correlation (**C**) in P1 and P4 brains. **D**)

Unsupervised hierarchical clustering of Ca^{2+} burst frequency and intra-neuron burst

correlation.

Figure 6. Impaired intrinsic network and bursting activity recorded in DIV 9

cultured Nr2f1-deficient cortical neurons. **A**) Spontaneous firing of *Ctrl* (left) and

Nr2f1cKO (right) cortical network was recorded by three representative MEA

electrodes a 9DIV. In insets, representative bursts are shown at expanded scale. **B-F**)

Firing parameters measured in *Ctrl* and *Nr2f1cKO* cortical networks are compared (*

P<0.05; **P<0.01, unpaired T-test). ***See also figure supplement 1.***

SUPPLEMENTAL FIGURES:

Figure 1 – supplement 1. Specific intrinsic excitability features and input-output

relationship remain unlatered in Nr2f1 deficient layer V pyramidal neurons

(LVPNs). **A**) LVPNs in *Ctrl* (in black) and *Nr2f1cKO* (in red) display normal input

resistance and threshold for action potential and inter-spike interval. We observed a

tendency of *Nr2f1cKO* LVPNs to display a reduced half-width of action potential when

compared to *Ctrl* LVPNs (P = 0.066) and a significant decreased AP*amp* (* *P* =

0.04441) measured from resting membrane protential (N = 19 *Ctrl*, N = 28 *Nr2f1cKO*,

two-tailed unpaired T-test or U Mann-Whitney test). **B**) Input/output function is not

significantly different between *Nr2f1cKO* LVPNs and *Ctrl* LVPNs (N = 19 *Ctrl*, N = 28

Nr2f1cKO, two-way ANOVA). R*in*, input resistance; V*threshold*, threshold for action

potential; AP*h-w*, half-width of action potential; ISI, inter-spike interval.

Figure 1 –table supplement 1. Intrinsic electrophysiological properties of layer V

pyramidal neurons in *ex vivo Ctrl* and *Nr2f1cKO* brain slices.

Figure 2 – figure supplement 1. Apical dendritic complexity of layer V pyramidal

neurons (LVPNs) is not affected in Nr2f1 mutant brains. **A**) Sholl analysis shows

no differences in complexity between *Ctrl* and *Nr2f1cKO* LVPNs, relatively to distal

dendrites. (ns P > 0.5) **B-D**) Number of Apical Branching Points (ns P = 0.57), Apical

Terminal Tips (ns P = 0.70) and Lateral dendrites (ns P =0.20) are unaffected by the

loss of *Nr2f1* (N = 12 *Ctrl*, N = 24 *Nr2f1cKO*). P-values are calculated by 2way ANOVA

test (**A**) or Welch's unequal variances t-test (**B-D**).

Figure 4 - figure supplement 1. qPCR quantification of cortical ion channels in

Nr2f1 mutant cortices. Single qPCR quantification of P7 *Ctrl* vs *Nr2f1cKO*

expression of chosen cortically expressed ion channel subunits and Glutamate

receptors. *Ctrl* mean value is normalized to 1 to calculate fold changes in *Nr2f1cKO*

conditions. N≥3. Statistical significance is tested doing paired t-test with P-value

adjusted according to Benjamini–Hochberg procedure. *- P-Value < 0.05. P –

Postnatal day.

Figure 6- figure supplement 1. Impaired intrinsic network and bursting activity

recorded in DIV15 cultured Nr2f1-deficient cortical neurons. **A**) Spontaneous

firing of *Ctrl* (left) and *Nr2f1cKO* (right) cortical network was recorded by three

representative MEA electrodes a 15DIV. In insets, representative bursts are shown

at expanded scale. **B-F**) Firing parameters measured in *Ctrl* and *Nr2f1cKO* cortical

networks are compared (* P<0.05; ***P<0.001, unpaired T-test)

Key resources table. List of antibodies used in this study.

REFERENCES:

ALFANO, C., MAGRINELLI, E., HARB, K., HEVNER, R. F. & STUDER, M. 2014. Postmitotic
control of sensory area specification during neocortical development. *Nat Commun,*
5**,** 5632.
ALFANO, C. & STUDER, M. 2012. Neocortical arealization: Evolution, mechanisms and open
questions. *Dev Neurobiol.*
ALFANO, C., VIOLA, L., HENG, J. I., PIROZZI, M., CLARKSON, M., FLORE, G., DE MAIO, A.,
SCHEDL, A., GUILLEMOT, F. & STUDER, M. 2011. COUP-TFI promotes radial
migration and proper morphology of callosal projection neurons by repressing Rnd2
expression. *Development,* 138**,** 4685-97.
ALLENE, C., CATTANI, A., ACKMAN, J. B., BONIFAZI, P., ANIKSZTEJN, L., BEN-ARI, Y. &
COSSART, R. 2008. Sequential generation of two distinct synapse-driven network
patterns in developing neocortex. *J Neurosci,* 28**,** 12851-63.
ALLENE, C. & COSSART, R. 2010. Early NMDA receptor-driven waves of activity in the
developing neocortex: physiological or pathological network oscillations? *J Physiol,*
588**,** 83-91.
ANDERSON, O. D. 1975. Moving Average Processes. The Statistician.
ANDREAE, L. C. & BURRONE, J. 2018. The role of spontaneous neurotransmission in synapse
and circuit development. *J Neurosci Res,* 96**,** 354-359.
ANTON-BOLANOS, N., SEMPERE-FERRANDEZ, A., GUILLAMON-VIVANCOS, T., MARTINI, F. J.,
PEREZ-SAIZ, L., GEZELIUS, H., FILIPCHUK, A., VALDEOLMILLOS, M. & LOPEZ-
BENDITO, G. 2019. Prenatal activity from thalamic neurons governs the emergence
of functional cortical maps in mice. *Science,* 364**,** 987-990.
ARLOTTA, P., MOLYNEAUX, B. J., CHEN, J., INOUE, J., KOMINAMI, R. & MACKLIS, J. D. 2005.
Neuronal subtype-specific genes that control corticospinal motor neuron
development in vivo. *Neuron,* 45**,** 207-21.
ARMENTANO, M., CHOU, S. J., TOMASSY, G. S., LEINGARTNER, A., O'LEARY, D. D. & STUDER,
M. 2007. COUP-TFI regulates the balance of cortical patterning between
frontal/motor and sensory areas. *Nat Neurosci,* 10**,** 1277-86.
ARNOLD, E. C., MCMURRAY, C., GRAY, R. & JOHNSTON, D. 2019. Epilepsy-Induced Reduction
in HCN Channel Expression Contributes to an Increased Excitability in Dorsal, But
Not Ventral, Hippocampal CA1 Neurons. *eNeuro,* 6.
BENDER, R. A. & BARAM, T. Z. 2008. Hyperpolarization activated cyclic-nucleotide gated
(HCN) channels in developing neuronal networks. *Prog Neurobiol,* 86**,** 129-40.
BERGER, T., LARKUM, M. E. & LUSCHER, H. R. 2001. High I(h) channel density in the distal
apical dendrite of layer V pyramidal cells increases bidirectional attenuation of
EPSPs. *J Neurophysiol,* 85**,** 855-68.
BERTACCHI, M., PARISOT, J. & STUDER, M. 2019. The pleiotropic transcriptional regulator
COUP-TFI plays multiple roles in neural development and disease. *Brain Res,* 1705**,**
75-94.
BLANKENSHIP, A. G., HAMBY, A. M., FIRL, A., VYAS, S., MAXEINER, S., WILLECKE, K. &
FELLER, M. B. 2011. The role of neuronal connexins 36 and 45 in shaping
spontaneous firing patterns in the developing retina. *J Neurosci,* 31**,** 9998-10008.
BOILLOT, M., LEE, C. Y., ALLENE, C., LEGUERN, E., BAULAC, S. & ROUACH, N. 2016. LGI1 acts
presynaptically to regulate excitatory synaptic transmission during early postnatal
development. *Sci Rep,* 6**,** 21769.
BOSCH, D. G., BOONSTRA, F. N., DE LEEUW, N., PFUNDT, R., NILLESEN, W. M., DE LIGT, J.,
GILISSEN, C., JHANGIANI, S., LUPSKI, J. R., CREMERS, F. P. & DE VRIES, B. B. 2016.
Novel genetic causes for cerebral visual impairment. *Eur J Hum Genet,* 24**,** 660-5.
BOSCH, D. G., BOONSTRA, F. N., GONZAGA-JAUREGUI, C., XU, M., DE LIGT, J., JHANGIANI, S.,
WISZNIEWSKI, W., MUZNY, D. M., YNTEMA, H. G., PFUNDT, R., VISSERS, L. E.,
SPRUIJT, L., BLOKLAND, E. A., CHEN, C. A., BAYLOR-HOPKINS CENTER FOR
MENDELIAN, G., LEWIS, R. A., TSAI, S. Y., GIBBS, R. A., TSAI, M. J., LUPSKI, J. R.,

ZOGHBI, H. Y., CREMERS, F. P., DE VRIES, B. B. & SCHAAF, C. P. 2014. NR2F1
mutations cause optic atrophy with intellectual disability. *Am J Hum Genet,* 94**,** 303-9.
BOYCHUK, J. A., FARRELL, J. S., PALMER, L. A., SINGLETON, A. C., PITTMAN, Q. J. & TESKEY, G.
C. 2017. HCN channels segregate stimulation-evoked movement responses in
neocortex and allow for coordinated forelimb movements in rodents. *J Physiol,* 595**,**
247-263.
BRAGER, D. H., AKHAVAN, A. R. & JOHNSTON, D. 2012. Impaired dendritic expression and
plasticity of h-channels in the fmr1(-/y) mouse model of fragile X syndrome. *Cell Rep,*
1**,** 225-33.
BRAGER, D. H. & JOHNSTON, D. 2007. Plasticity of intrinsic excitability during long-term
depression is mediated through mGluR-dependent changes in I(h) in hippocampal
CA1 pyramidal neurons. *J Neurosci,* 27**,** 13926-37.
CHEN, C. A., BOSCH, D. G., CHO, M. T., ROSENFELD, J. A., SHINAWI, M., LEWIS, R. A., MANN, J.,
JAYAKAR, P., PAYNE, K., WALSH, L., MOSS, T., SCHREIBER, A., SCHOONVELD, C.,
MONAGHAN, K. G., ELMSLIE, F., DOUGLAS, G., BOONSTRA, F. N., MILLAN, F.,
CREMERS, F. P., MCKNIGHT, D., RICHARD, G., JUUSOLA, J., KENDALL, F., RAMSEY, K.,
ANYANE-YEBOA, K., MALKIN, E., CHUNG, W. K., NIYAZOV, D., PASCUAL, J. M.,
WALKIEWICZ, M., VELUCHAMY, V., LI, C., HISAMA, F. M., DE VRIES, B. B. & SCHAAF, C.
2016. The expanding clinical phenotype of Bosch-Boonstra-Schaaf optic atrophy
syndrome: 20 new cases and possible genotype-phenotype correlations. *Genet Med.*
CHEN, K., ARADI, I., THON, N., EGHBAL-AHMADI, M., BARAM, T. Z. & SOLTESZ, I. 2001.
Persistently modified h-channels after complex febrile seizures convert the seizure-
induced enhancement of inhibition to hyperexcitability. *Nat Med,* 7**,** 331-7.
CHO, M. W. & CHOI, M. Y. 2012. Spontaneous organization of the cortical structure through
endogenous neural firing and gap junction transmission. *Neural Netw,* 31**,** 46-52.
COMPUTING, R. F. F. S. 2018. R: A Language and Environment for Statistical Computing. R
Foundation for Statistical Computing.
CONTESSE, T., AYRAULT, M., MANTEGAZZA, M., STUDER, M. & DESCHAUX, O. 2019.
Hyperactive and anxiolytic-like behaviors result from loss of COUP-TFI/Nr2f1 in the
mouse cortex. *Genes Brain Behav***,** e12556.
CORLEW, R., BOSMA, M. M. & MOODY, W. J. 2004. Spontaneous, synchronous electrical
activity in neonatal mouse cortical neurones. *J Physiol,* 560**,** 377-90.
DEL PINO, I., BROTONS-MAS, J. R., MARQUES-SMITH, A., MARIGHETTO, A., FRICK, A., MARIN,
O. & RICO, B. 2017. Abnormal wiring of CCK(+) basket cells disrupts spatial
information coding. *Nat Neurosci,* 20**,** 784-792.
DEL PINO, I., GARCIA-FRIGOLA, C., DEHORTER, N., BROTONS-MAS, J. R., ALVAREZ-SALVADO,
E., MARTINEZ DE LAGRAN, M., CICERI, G., GABALDON, M. V., MORATAL, D.,
DIERSSEN, M., CANALS, S., MARIN, O. & RICO, B. 2013. Erbb4 deletion from fast-
spiking interneurons causes schizophrenia-like phenotypes. *Neuron,* 79**,** 1152-68.
DUMITRESCU, A. S., EVANS, M. D. & GRUBB, M. S. 2016. Evaluating Tools for Live Imaging of
Structural Plasticity at the Axon Initial Segment. *Front Cell Neurosci,* 10**,** 268.
FAEDO, A., TOMASSY, G. S., RUAN, Y., TEICHMANN, H., KRAUSS, S., PLEASURE, S. J., TSAI, S. Y.,
TSAI, M. J., STUDER, M. & RUBENSTEIN, J. L. 2008. COUP-TFI coordinates cortical
patterning, neurogenesis, and laminar fate and modulates MAPK/ERK, AKT, and
beta-catenin signaling. *Cereb Cortex,* 18**,** 2117-31.
FAN, J., STEMKOWSKI, P. L., GANDINI, M. A., BLACK, S. A., ZHANG, Z., SOUZA, I. A., CHEN, L. &
ZAMPONI, G. W. 2016. Reduced Hyperpolarization-Activated Current Contributes to
Enhanced Intrinsic Excitability in Cultured Hippocampal Neurons from PrP(-/-)
Mice. *Front Cell Neurosci,* 10**,** 74.
FAN, Y., FRICKER, D., BRAGER, D. H., CHEN, X., LU, H. C., CHITWOOD, R. A. & JOHNSTON, D.
2005. Activity-dependent decrease of excitability in rat hippocampal neurons
through increases in I(h). *Nat Neurosci,* 8**,** 1542-51.
FAZEL DARBANDI, S., ROBINSON SCHWARTZ, S. E., QI, Q., CATTA-PRETA, R., PAI, E. L.,
MANDELL, J. D., EVERITT, A., RUBIN, A., KRASNOFF, R. A., KATZMAN, S., TASTAD, D.,

NORD, A. S., WILLSEY, A. J., CHEN, B., STATE, M. W., SOHAL, V. S. & RUBENSTEIN, J. L. R. 2018. Neonatal Tbr1 Dosage Controls Cortical Layer 6 Connectivity. *Neuron,* 100**,** 831-845.e7.

FLORE, G., DI RUBERTO, G., PARISOT, J., SANNINO, S., RUSSO, F., ILLINGWORTH, E. A., STUDER, M. & DE LEONIBUS, E. 2017. Gradient COUP-TFI Expression Is Required for Functional Organization of the Hippocampal Septo-Temporal Longitudinal Axis. *Cereb Cortex,* 27**,** 1629-1643.

GAVELLO, D., CALORIO, C., FRANCHINO, C., CESANO, F., CARABELLI, V., CARBONE, E. & MARCANTONI, A. 2018. Early Alterations of Hippocampal Neuronal Firing Induced by Abeta42. *Cereb Cortex,* 28**,** 433-446.

GAVELLO, D., ROJO-RUIZ, J., MARCANTONI, A., FRANCHINO, C., CARBONE, E. & CARABELLI, V. 2012. Leptin counteracts the hypoxia-induced inhibition of spontaneously firing hippocampal neurons: a microelectrode array study. *PLoS One,* 7**,** e41530.

GREIG, L. C., WOODWORTH, M. B., GALAZO, M. J., PADMANABHAN, H. & MACKLIS, J. D. 2013. Molecular logic of neocortical projection neuron specification, development and diversity. *Nat Rev Neurosci,* 14**,** 755-69.

GROC, L., PETANJEK, Z., GUSTAFSSON, B., BEN-ARI, Y., HANSE, E. & KHAZIPOV, R. 2002. In vivo blockade of neural activity alters dendritic development of neonatal CA1 pyramidal cells. *Eur J Neurosci,* 16**,** 1931-8.

GRUBB, M. S. & BURRONE, J. 2010. Activity-dependent relocation of the axon initial segment fine-tunes neuronal excitability. *Nature,* 465**,** 1070-4.

GULLEDGE, A. T. & BRAVO, J. J. 2016. Neuron Morphology Influences Axon Initial Segment Plasticity. *eNeuro,* 3.

HAMADA, M. S., GOETHALS, S., DE VRIES, S. I., BRETTE, R. & KOLE, M. H. 2016. Covariation of axon initial segment location and dendritic tree normalizes the somatic action potential. *Proc Natl Acad Sci U S A,* 113**,** 14841-14846.

HARB, K., MAGRINELLI, E., NICOLAS, C. S., LUKIANETS, N., FRANGEUL, L., PIETRI, M., SUN, T., SANDOZ, G., GRAMMONT, F., JABAUDON, D., STUDER, M. & ALFANO, C. 2016. Area-specific development of distinct projection neuron subclasses is regulated by postnatal epigenetic modifications. *Elife,* 5**,** e09531.

JABAUDON, D. 2017. Fate and freedom in developing neocortical circuits. *Nat Commun,* 8**,** 16042.

KIRISCHUK, S., SINNING, A., BLANQUIE, O., YANG, J. W., LUHMANN, H. J. & KILB, W. 2017. Modulation of Neocortical Development by Early Neuronal Activity: Physiology and Pathophysiology. *Front Cell Neurosci,* 11**,** 379.

KIRKBY, L. A., SACK, G. S., FIRL, A. & FELLER, M. B. 2013. A role for correlated spontaneous activity in the assembly of neural circuits. *Neuron,* 80**,** 1129-44.

KLUEVA, J., LIMA, A. D., MEIS, S., VOIGT, T. & MUNSCH, T. 2012. Hyperpolarization-activated cation current contributes to spontaneous network activity in developing neocortical cultures. *Neurosignals,* 20**,** 35-47.

KOLE, M. H. & STUART, G. J. 2012. Signal processing in the axon initial segment. *Neuron,* 73**,** 235-47.

KORDELI, E., LAMBERT, S. & BENNETT, V. 1995. AnkyrinG. A new ankyrin gene with neural-specific isoforms localized at the axonal initial segment and node of Ranvier. *J Biol Chem,* 270**,** 2352-9.

KUBA, H. 2010. Plasticity at the axon initial segment. *Commun Integr Biol,* 3**,** 597-8.

KUBA, H., OICHI, Y. & OHMORI, H. 2010. Presynaptic activity regulates Na(+) channel distribution at the axon initial segment. *Nature,* 465**,** 1075-8.

LI, H., FERTUZINHOS, S., MOHNS, E., HNASKO, T. S., VERHAGE, M., EDWARDS, R., SESTAN, N. & CRAIR, M. C. 2013. Laminar and columnar development of barrel cortex relies on thalamocortical neurotransmission. *Neuron,* 79**,** 970-86.

LOOTS, G. & OVCHARENKO, I. 2007. ECRbase: database of evolutionary conserved regions, promoters, and transcription factor binding sites in vertebrate genomes. *Bioinformatics,* 23**,** 122-4.

LUHMANN, H. J. & KHAZIPOV, R. 2018. Neuronal activity patterns in the developing barrel cortex. *Neuroscience,* 368**,** 256-267.

LUHMANN, H. J., SINNING, A., YANG, J. W., REYES-PUERTA, V., STUTTGEN, M. C., KIRISCHUK, S. & KILB, W. 2016. Spontaneous Neuronal Activity in Developing Neocortical Networks: From Single Cells to Large-Scale Interactions. *Front Neural Circuits,* 10**,** 40.

MAGEE, J. C. 1998. Dendritic hyperpolarization-activated currents modify the integrative properties of hippocampal CA1 pyramidal neurons. *J Neurosci,* 18**,** 7613-24.

MOODY, W. J. & BOSMA, M. M. 2005. Ion channel development, spontaneous activity, and activity-dependent development in nerve and muscle cells. *Physiol Rev,* 85**,** 883-941.

MURASE, S., LANTZ, C. L., KIM, E., GUPTA, N., HIGGINS, R., STOPFER, M., HOFFMAN, D. A. & QUINLAN, E. M. 2016. Matrix Metalloproteinase-9 Regulates Neuronal Circuit Development and Excitability. *Mol Neurobiol,* 53**,** 3477-3493.

O'LEARY, D. D. & SAHARA, S. 2008. Genetic regulation of arealization of the neocortex. *Curr Opin Neurobiol,* 18**,** 90-100.

OVCHARENKO, I., NOBREGA, M. A., LOOTS, G. G. & STUBBS, L. 2004. ECR Browser: a tool for visualizing and accessing data from comparisons of multiple vertebrate genomes. *Nucleic Acids Res,* 32**,** W280-6.

PEARSON, R. A., CATSICAS, M., BECKER, D. L., BAYLEY, P., LUNEBORG, N. L. & MOBBS, P. 2004. Ca(2+) signalling and gap junction coupling within and between pigment epithelium and neural retina in the developing chick. *Eur J Neurosci,* 19**,** 2435-45.

POOLOS, N. P., BULLIS, J. B. & ROTH, M. K. 2006. Modulation of h-channels in hippocampal pyramidal neurons by p38 mitogen-activated protein kinase. *J Neurosci,* 26**,** 7995-8003.

POOLOS, N. P., MIGLIORE, M. & JOHNSTON, D. 2002. Pharmacological upregulation of h-channels reduces the excitability of pyramidal neuron dendrites. *Nat Neurosci,* 5**,** 767-74.

ROERIG, B. & FELLER, M. B. 2000. Neurotransmitters and gap junctions in developing neural circuits. *Brain Res Brain Res Rev,* 32**,** 86-114.

SCHINDELIN, J., ARGANDA-CARRERAS, I., FRISE, E., KAYNIG, V., LONGAIR, M., PIETZSCH, T., PREIBISCH, S., RUEDEN, C., SAALFELD, S., SCHMID, B., TINEVEZ, J. Y., WHITE, D. J., HARTENSTEIN, V., ELICEIRI, K., TOMANCAK, P. & CARDONA, A. 2012. Fiji: an open-source platform for biological-image analysis. *Nat Methods,* 9**,** 676-82.

SCHNEGGENBURGER, R. & ROSENMUND, C. 2015. Molecular mechanisms governing Ca(2+) regulation of evoked and spontaneous release. *Nat Neurosci,* 18**,** 935-41.

SHAH, M. M. 2014. Cortical HCN channels: function, trafficking and plasticity. *J Physiol,* 592**,** 2711-9.

SIMI, A. & STUDER, M. 2018. Developmental genetic programs and activity-dependent mechanisms instruct neocortical area mapping. *Curr Opin Neurobiol,* 53**,** 96-102.

TOMASSY, G. S., DE LEONIBUS, E., JABAUDON, D., LODATO, S., ALFANO, C., MELE, A., MACKLIS, J. D. & STUDER, M. 2010. Area-specific temporal control of corticospinal motor neuron differentiation by COUP-TFI. *Proc Natl Acad Sci U S A,* 107**,** 3576-81.

TRITSCH, N. X., YI, E., GALE, J. E., GLOWATZKI, E. & BERGLES, D. E. 2007. The origin of spontaneous activity in the developing auditory system. *Nature,* 450**,** 50-5.

UHLEN, P., FRITZ, N., SMEDLER, E., MALMERSJO, S. & KANATANI, S. 2015. Calcium signaling in neocortical development. *Dev Neurobiol,* 75**,** 360-8.

VALIULLINA, F., AKHMETSHINA, D., NASRETDINOV, A., MUKHTAROV, M., VALEEVA, G., KHAZIPOV, R. & ROZOV, A. 2016. Developmental Changes in Electrophysiological Properties and a Transition from Electrical to Chemical Coupling between Excitatory Layer 4 Neurons in the Rat Barrel Cortex. *Front Neural Circuits,* 10**,** 1.

VAN WELIE, I., REMME, M. W., VAN HOOFT, J. A. & WADMAN, W. J. 2006. Different levels of Ih determine distinct temporal integration in bursting and regular-spiking neurons in rat subiculum. *J Physiol,* 576**,** 203-14.

WANG, Z., XU, N. L., WU, C. P., DUAN, S. & POO, M. M. 2003. Bidirectional changes in spatial
dendritic integration accompanying long-term synaptic modifications. *Neuron,* 37**,**
463-72.
WEFELMEYER, W., PUHL, C. J. & BURRONE, J. 2016. Homeostatic Plasticity of Subcellular
Neuronal Structures: From Inputs to Outputs. *Trends Neurosci,* 39**,** 656-667.
WILLIAMS, S. R. & STUART, G. J. 2000. Site independence of EPSP time course is mediated by
dendritic I(h) in neocortical pyramidal neurons. *J Neurophysiol,* 83**,** 3177-82.
WILLIAMS, S. R. & STUART, G. J. 2003. Voltage- and site-dependent control of the somatic
impact of dendritic IPSPs. *J Neurosci,* 23**,** 7358-67.
YAMADA, R. & KUBA, H. 2016. Structural and Functional Plasticity at the Axon Initial
Segment. *Front Cell Neurosci,* 10**,** 250.
YAMAMOTO, N. & LOPEZ-BENDITO, G. 2012. Shaping brain connections through
spontaneous neural activity. *Eur J Neurosci,* 35**,** 1595-604.
ZHOU, D., LAMBERT, S., MALEN, P. L., CARPENTER, S., BOLAND, L. M. & BENNETT, V. 1998.
AnkyrinG is required for clustering of voltage-gated Na channels at axon initial
segments and for normal action potential firing. *J Cell Biol,* 143**,** 1295-304.

1181

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 1 – Supplement 1

Figure 2 - Supplement 1

Figure 4 - Supplement 1

Figure 6 - Supplement 1

	Ctrl	**cKO**	p-value	test
V*rest* (mV)	-60.26 ± 1.18	-54.62 ± 1.13	0.0013 (*)	t-test
R*in* (MΩ)	561.2 ± 36.66	494.80 ± 38.64	0.2410	t-test
Latency (ms)	133.5 ± 15.46	145.9 ± 23.59	0.9023	Mann-Whitney
Rheobase (pA)	61.05 ± 9.08	29.64 ± 3.47	0.001(***)	t-test
V*threshold* (mV)	-30.95 ± 0.48	-32.09 ± 0.60	0.1750	t-test
AP*amp(from* V*threshold)* (mV)	66.65 ± 1.62	69.1 ± 1.11	0.2042	t-test
AP*amp(from* V*rest)* (mV)	96.08 ± 1.611	90.72 ± 1.70	0.0441(*)	Mann-Whitney
AP*half-width* (ms)	0.0041±0.0002	0.0037±0.0001	0.0659	t-test
Inter-spike-interval(ISI) (ms)	172.0 ± 19.92	175.1 ± 16.42	0.9529	Mann-Whitney
Sag (mV)	11.64 ± 1.28	7.2 ± 1.051	0.0103(*)	t-test
τ*m* (ms)	28.42 ± 3.26	31.14 ± 3.70	0.6193	Mann-Whitney