Membrane synthesis in *Escherichia coli* is coordinated with growth via post-translational regulation

Marek J Noga[†], Ferhat Büke, Niels JF van den Broek[‡], Nicole Imholz, Nicole Scherer, Gregory

Bokinsky*

Department of Bionanoscience, Kavli Institute of Nanoscience, Delft University of Technology,

Delft, The Netherlands

Materials and correspondence: *e-mail: g.e.bokinsky@tudelft.nl

[†] Current address: Translational Metabolic Laboratory, Department of Laboratory Medicine,

Radboud UMC, Nijmegen, The Netherlands

[‡] Current address: Vaccine Process & Analytical Development, Janssen Vaccines & Prevention

B.V., Leiden, The Netherlands

Abstract

Every growing cell must produce enough membrane to contain itself. However, the mechanisms by which cells couple the rate of membrane synthesis with the rate of growth remain unresolved. By measuring precursors, intermediates, and enzymes of the fatty acid and phospholipid synthesis pathways of *Escherichia coli* during steady-state growth, we find that while transcriptional regulation maintains membrane synthesis capacity at a stable level, the actual rate of phospholipid synthesis is mediated by allosteric control of a single enzyme (PlsB). PlsB activity also strongly influences flux into the synthesis pathway of lipopolysaccharide, the second-most abundant membrane component in Gram negative bacteria. Furthermore, we find that membrane synthesis rates are highly sensitive to changes in fatty acid precursor concentrations, enabling rapid responses to environmental perturbations. Such responses are modulated by the global regulator ppGpp, which we hypothesize acts as an emergency brake on PlsB activity to prevent imbalances between membrane synthesis and synthesis of lipoproteins that tether the outer membrane to the cell. The combination of dynamic post-translational regulation with maintenance of stable synthesis capacity ensures both robustness and responsiveness of bacterial membrane synthesis during growth.

1 Introduction

The viability of growing cells demands that membrane synthesis be tightly orchestrated

with biomass production. The mechanisms that adjust synthesis rates of membrane components

(e.g. phospholipids and lipopolysaccharides) during steady-state growth or in response to

environmental perturbations remain unknown (1). Recently, advances in analytical methods and

systems-level modelling have delivered new insights into how living cells control enzyme

reaction rates. Reaction rates (metabolic fluxes) may be modulated through either changes in

enzyme concentrations mediated by transcriptional regulation, or by adjusting the activities of

enzymes that are already present by post-translational regulation. Two groups inferred

mechanisms of flux control by quantifying metabolic fluxes and concentrations of metabolites

and enzymes during steady-state growth in various conditions (2, 3). Both groups concluded that

pathway fluxes are more often determined by steady-state concentrations of substrates and

inhibitors than by concentrations of enzymes. In other words, steady-state flux is controlled in

many pathways by post-translational regulation of enzyme activity rather than by transcriptional

regulation of enzyme abundance. On the other hand, the global synthesis rate of proteins – the

most abundant form of biomass – is determined in many fast-growing bacteria by ribosome

abundance, which is transcriptionally regulated to balance amino acid supply with consumption

(4). Whether membrane synthesis flux is regulated either via transcriptional control (similar to

protein synthesis) or by post-translational control remains unknown.

The architecture and composition of the cell envelope in Gram-negative bacteria also

demands tight coordination between multiple biosynthesis pathways. The inner membrane and

the inner leaflet of the outer membrane are composed primarily of phospholipids (PL), while the

outer leaflet of the outer membrane is composed of lipopolysaccharides (LPS) (5, 6). PL and

LPS pathways must also be coordinated with protein synthesis, which supplies lipoproteins (e.g.

Lpp) that tether the outer membrane to peptidoglycan (7). In *Escherichia coli,* a model organism

for bacterial fatty acid and PL synthesis (1), efforts to determine the mechanism of PL synthesis

flux regulation have so far proven unsuccessful. Overexpression of PL synthesis enzymes has

been consistently found to have no effect on either PL abundance or on the ratio of zwitterionic

and anionic PL species (8, 9). Perturbations to membrane composition have been achieved only

when PL synthesis enzymes are depleted far below wild-type concentrations, suggesting that

their steady-state concentrations are maintained above levels needed to sustain growth (10–12).

In order to understand how the rate of PL synthesis is coordinated with the rate of growth

(μ), we quantified enzymes, substrates and intermediates of the fatty acid and PL synthesis

pathways in steady-state and dynamic conditions. We find that allosteric regulation of the first

enzyme in the PL synthesis pathway, PlsB, provides μ-dependent regulation of PL synthesis and

influences flux into the LPS synthesis pathway. We furthermore find that the fatty acid and PL

synthesis pathways are highly sensitive to changes in precursor concentrations caused by

internal and external metabolic shifts. This sensitivity enables a rapid response to environmental

changes that can be further modulated by allosteric inhibitors such as the global regulator

guanosine tetraphosphate (ppGpp).

Results

The PL to biomass ratio varies inversely with μ

Lipid precursors of both LPS and PL are synthesized in the cytosol as fatty acyl thioesters covalently attached to the acyl carrier protein (acyl-ACP). PL are synthesized by membrane-bound enzymes from long-chain acyl-ACP and *sn*-glycerol-3 phosphate (G3P). G3P is produced either from glyceraldehyde-3-phosphate, an intermediate of central carbon metabolism, or via glycerol catabolism **(Figure 1A)**. We grew cultures of *E. coli* NCM3722 in 6 defined media that support a 3-fold range of μ (MOPS medium containing 0.2% acetate, succinate, malate, glycerol, glucose, or glucose with Cas-amino acids). Quantities of phosphatidylglycerol (PG), phosphatidylethanolamine (PE), and cardiolipin (CL) per unit biomass (as determined by culture optical density (OD) (13)) decreased with increasing μ **(Figure 1B)**, consistent with previous observations in *E. coli* and other bacteria (14, 15). The PG/PE ratio remained constant across all μ. We quantified steady-state PL flux by multiplying total PE LC/MS counts by μ, which closely approximates the overall PL synthesis rate as PE turnover is slow relative to synthesis (16). PL flux decreased by 2-fold over 3-fold range of μ **(Figure 1C)**. The higher PL to biomass ratio of slow-growing cells likely reflects their higher surface-to-volume ratio (17).

Allosteric regulation of PlsB activity couples PL flux with μ

We first sought to evaluate whether steady-state substrate concentrations determine PL flux. If the PL and fatty acid synthesis rates are modulated by substrate concentrations during steady-state growth, concentrations of these species would necessarily correlate positively with μ and PL flux. We quantified nucleotides, G3P, acyl-ACP, and PL synthesis intermediates using LC/MS (18). No correlation was observed between concentrations of the fatty acid precursor acetyl-CoA and μ (**Figure 1D**). We also found no significant correlation between PL flux and

long-chain acyl-ACP substrates of PL synthesis **(Figure 1E,** Pearson correlation coefficients $r <$

0.4, $p > 0.4$, **Supplemental Table 1).** Although G3P concentrations correlate strongly with PL

flux ($r > 0.99$), growth in glycerol medium increased G3P by 20-fold without affecting PL content

or flux. These trends imply that steady-state PL flux is not modulated by fatty acid or PL

precursor concentrations.

In contrast to PL synthesis substrates, concentrations of the fatty acid initiation and

elongation substrate malonyl-ACP strongly correlate with PL flux **(Figure 1E,** $r = 0.9$,

Supplemental Table 1), suggesting that the fatty acid initiation and elongation rates are

determined by malonyl-ACP concentrations and thus by the rate of malonyl-ACP synthesis by

the ACC complex and FabD. Concentrations of hydroxyl-acyl-ACP intermediates, including

C14:0-OH-ACP, the lipid precursor of LPS, also correlate strongly with PL flux ($r > 0.99$), implying

a simple flux-sensitive coupling mechanism between the PL and LPS synthesis pathways

(Figure 1E, Supplemental Figure 1). As ACC substrates acetyl-CoA and ATP do not correlate

with μ **(Figure 1D, Supplemental Figure 2)**, and as concentrations of ACC protein subunits and

FabD also do not correlate significantly with PL flux (**Supplemental Figure 3, Supplemental**

Table 1), we infer that ACC activity is primarily modulated during steady-state conditions via

allosteric control. Biochemical experiments have identified several regulators of ACC activity,

including nitrogen-responsive regulator GlnB (19) and long-chain acyl-ACP substrates of PL

synthesis (20).

Unlike long-chain acyl-ACP, the concentrations of all PL synthesis intermediates

downstream of PlsB correlate tightly with PL flux **(Figure 1F,** $r > 0.89$-0.99, **Supplemental Table**

1). The contrast between the trends observed in substrates and products of PlsB suggests that

the rates of the reactions following PlsB are primarily determined by substrate concentrations,

further implying that transcriptional regulation of PlsB concentration or post-translational

regulation of PlsB activity is sufficient to regulate flux into the PL pathway. Transcription of *plsB,*

and thus PlsB abundance, is controlled by the μ-sensitive regulator guanosine tetraphosphate

(ppGpp) (21) and by the membrane stress-activated sigma factor RpoE (22). Concentrations of

ppGpp are inversely correlated with μ **(Figure 1D)**, implying that PL flux may be coupled to μ by

transcriptional control of the *plsB* gene by ppGpp. We measured fatty acid and PL synthesis

pathway enzymes in each culture using LC/MS to determine whether enzyme concentrations are

adjusted to couple PL flux with μ. We found no significant correlation between PL synthesis flux

and PlsB concentration **(Figure 1G, Supplemental Table 1,** *p* >0.1). Instead, PlsB

concentrations per unit biomass (OD) are nearly constant across the conditions studied,

indicating that PlsB can accommodate at least a 2-fold range of PL flux without a corresponding

change in concentration. The concentrations of most fatty acid synthesis pathway enzymes also

do not correlate significantly with PL flux **(Supplemental Table 1, Supplemental Figure 3).** The

lack of correlation between PL flux and concentrations of PlsB or its acyl-ACP substrates implies

that PL flux is regulated via allosteric control of PlsB.

μ can also be modulated independently of culture conditions by artificially varying ppGpp

(Figure 1D). Increasing ppGpp above basal concentrations to sub-starvation levels reduces

steady-state μ by reducing ribosomal RNA synthesis (23). Furthermore, PlsB activity is inhibited

at high ppGpp concentrations encountered during starvation (24). We titrated μ by expressing

the catalytic domain of the ppGpp synthesis enzyme RelA (RelA*) from an inducible promoter

(P_{Tet}) in glucose medium. As previously observed (23), RelA*-titrated cultures exhibit ppGpp

concentrations that are elevated 2-fold above wild-type cultures growing at similar rates **(Figure**

1D). Trends in PL abundance, PL flux, acyl-ACP species, PL intermediates, and PL synthesis

enzyme concentrations in ppGpp-limited cultures are highly similar to trends observed as μ is

varied by carbon source **(Figure 1B-C, 1E-G, Supplemental Figures 1, 3).** Concentrations of

PlsB and fatty acid pathway enzymes do not substantially vary in ppGpp-limited cultures **(Figure**

1G, Supplemental Figure 3). The trends in substrate, enzyme, and intermediate concentrations

observed in ppGpp-limited cultures are consistent with regulation of PL flux occurring via

allosteric control of PlsB. Unexpectedly, concentrations of LpxC, which catalyses the committed

step in the LPS pathway (25), decrease 2-fold as PE flux increases **(Supplemental Figure 3).**

Mathematical modelling supports PlsB control of steady-state PL synthesis

To test whether modulation of PlsB activity alone is sufficient to govern PL synthesis flux,

we constructed a simplified differential equation model of fatty acid, LPS initiation, and PL

biosynthesis **(Figure 2A).** The model includes competitive inhibition of ACC by C16:0-ACP as

the sole regulatory interaction. The model also includes a branch point at C14:0-OH-ACP into

LPS synthesis. This branch point is represented by a single step that combines reactions

catalysed by LpxA and LpxC, as LpxC catalyses the first irreversible reaction in the LPS

pathway (25). Concentrations of G3P and C16:1-ACP are fixed in the model to reflect

experimental observations of enzyme saturation and substrate invariance, respectively (see

Supplementary Methods for model details, parameter sets, and sensitivity analysis).

We used the model to predict which enzymes and substrates exert control over PL and

LPS flux. Increasing ACC and PlsB V_{max} parameters by 4-fold each increased PE and LPS

synthesis fluxes in parallel by 2-fold, while changes in the V_{max} parameters of all other enzymes

in the model had little or no effect on simulated PE flux. Surprisingly, simulated LPS flux is also

very strongly influenced by PlsB activity in the absence of LpxC V_{max} regulation due to tight

coupling between PE flux C14:0-OH-ACP concentrations. Changes in the C14:0-OH-ACP

dehydration reduction rate and LPS initiation rate (catalysed in the model by FabZ and LpxC,

respectively) exerted strong and opposite effects on LPS flux without significantly changing PE

flux. Increasing the V_{max} parameters of PL synthesis enzymes downstream of PlsB also did not

change PE synthesis flux, suggesting that the observed variations in PlsC and CdsA

concentrations do not affect PE flux. Of the two substrates considered (acetyl-CoA and C16:1-

ACP), only variations in acetyl-CoA strongly influences PL and LPS flux **(Figure 2B)**.

Simulated variations in acetyl-CoA concentrations and ACC and PlsB enzyme V_{max}

parameters generate changes in intermediate concentrations that closely reproduce most

experimentally-observed trends, supporting our model. However, our steady-state

measurements strongly contradict simulated C16:0-ACP concentration trends caused by

variations in acetyl-CoA and ACC activity. Simulated variations in PlsB V_{max} more closely

approximate observed C16:0-ACP trends **(Figure 2C).** We therefore conclude that PL synthesis

flux is primarily regulated during steady-state growth by allosteric control of PlsB activity.

However, changes in acetyl-CoA concentration or ACC activity may also influence PL flux during

environmental or metabolic perturbations.

ppGpp attenuates excess lipid synthesis triggered by translation inhibition

We set out to evaluate whether a known allosteric regulator of PlsB, ppGpp, is able to

directly regulate PlsB activity during steady-state growth. High concentrations of ppGpp are

known to inhibit PlsB (24), while low concentrations of ppGpp correlate inversely with μ **(Figure**

1D). The notion that ppGpp concentrations might regulate PL synthesis even at the low basal

concentrations present during steady-state growth has been proposed but never tested (26).

Although steady-state PlsB activity clearly decreased in response to ppGpp titration, the mode of

control by ppGpp (transcriptional regulation of a PlsB-regulating process or protein, or direct

post-translational inhibition) cannot be determined from steady-state data. However, we first

wished to observe the effects of high concentrations of ppGpp on fatty acid and PL synthesis

using our assay. Synthesis of high concentrations of ppGpp by RelA (the stringent response) is

triggered by any stress or starvation conditions that lead to the specific biochemical cue of

uncharged tRNA bound to the ribosome. During the stringent response, ppGpp accumulates by

more than 10-fold over basal concentrations and PL synthesis rates are reduced by

approximately half (24, 26), likely due to PlsB inhibition (27).

1    We triggered ppGpp accumulation by adding the tRNA aminoacylation inhibitor mupirocin

2  to glucose cultures of wild-type and Δ*relA E. coli*. Mupirocin inhibits translation in both strains

3  and causes a 10-fold accumulation of ppGpp in the wild-type strain within 1 minute to over 400

4  pmol/OD, reaching >800 pmol/OD after 3 minutes **(Supplemental Figures 4, 5).** To our

5  surprise, mupirocin treatment also transiently increased malonyl-ACP and hydroxyl-ACP at the

6  expense of holo-ACP in both strains **(Figure 3A, Supplemental Figure 6)**. The increase in acyl-

7  ACP is matched by a corresponding increase in PL synthesis intermediates, suggesting that

8  mupirocin diverts a pulse of carbon into the fatty acid pathway that is efficiently transmitted into

9  PL synthesis **(Figure 3B)**. In the wild-type strain, the pulse of fatty acid synthesis activity is

10  rapidly followed by C16:0- and C18:0-ACP accumulation, consistent with ppGpp inhibition of

11  PlsB **(Figure 3A, Supplemental Figure 6)**. C16:0-ACP accumulation is followed in turn by an

12  increase in holo-ACP and a decrease in malonyl-ACP, indicating ACC inhibition **(Figure 3A)**. PL

13  intermediates PA, PS, and PGP are rapidly depleted in the wild-type strain after briefly

14  increasing **(Figure 3B).** The transient carbon influx shifts acyl chains incorporated into PL

15  towards longer-chain fatty acids, likely due to increased malonyl-ACP concentrations favouring

16  fatty acid elongation at the expense of membrane incorporation by PlsB and PlsC

17  **(Supplemental Figure 7).** While suppression of fatty acid synthesis in the wild-type strain can

18  be attributed to ppGpp, it is unclear what attenuates the carbon influx in the Δ*relA* strain;

19  however eventual C14:0-ACP accumulation likely contributes to ACC inhibition. Inhibition of fatty

20  acid elongation also depletes LPS precursor C14:0-OH-ACP, which is expected to decrease

21  LPS synthesis in parallel with PL synthesis **(Figure 3A)**.

22    As the transient increase in fatty acid synthesis after mupirocin treatment occurs in both

23  wild-type and Δ*relA* strains, we hypothesized that it is caused by translation inhibition. We added

24  the ribosome inhibitor chloramphenicol and the transcription initiation inhibitor rifampicin to

25  glycerol cultures of wild-type *E. coli*. Both compounds inhibit translation via mechanisms that

1   suppress ppGpp synthesis. As with mupirocin, chloramphenicol triggered a rapid decrease in

2   holo-ACP and an increase in long-chain unsaturated acyl-ACP species C16:1-ACP and C18:1-

3   ACP **(Figure 3C).** Both antibiotics triggered an increase in PL synthesis intermediates PA and

4   PS that resembled the increase observed in the Δ*relA* strain after mupirocin treatment **(Figure**

5   **3D).**

6   What might cause the transient increase in fatty acid synthesis observed after translation

inhibition? Interestingly, both rifampicin and chloramphenicol increased acetyl-CoA

concentrations by 3-fold **(Figure 3E)**, suggesting a possible cause. Acetyl-CoA also increased in

both the wild-type and the Δ*relA* strain after mupirocin addition in glucose medium before

decreasing ~30% in the wild-type strain **(Supplemental Figures 4, 5).** While it is unclear why

translation inhibition would increase acetyl-CoA, the effects of acetyl-CoA variation on fatty acid

and PL synthesis suggests that both pathways are highly sensitive to its concentration, a finding

that agrees with the sensitivity predicted by our mathematical model. We confirmed the

sensitivity of the fatty acid and PL synthesis pathways to environmental changes using a fast

nutritional upshift. Addition of glucose and amino acids to a glycerol culture also caused a rapid

accumulation of PA and PS species that resemble the increases observed following translation

inhibition **(Supplemental Figure 8).**

Moderate ppGpp concentrations regulate PlsB via post-translational control

In order to clearly discern the effects of ppGpp on the fatty acid and PL synthesis

pathways without complications introduced by translation inhibition, we monitored fatty acid and

PL synthesis pathways after inducing RelA*. The response in the fatty acid and PL synthesis

pathways is consistent with PlsB inhibition by ppGpp followed by ACC activity suppression by

accumulated long-chain acyl-ACP, depleting malonyl-ACP **(Figure 4A)**. PL intermediates also

respond in a manner consistent with PlsB inhibition: LPA species steadily decrease, followed by

PA, PS, and PG **(Figure 4B)**. Addition of chloramphenicol 10 minutes following RelA* induction

causes an increase in unsaturated long-chain acyl-ACP, though the response of the PL pathway

is attenuated. We note that the PS pool is depleted more after ppGpp induction than are the LPA

or PA pools, suggesting an additional regulatory mechanism downstream of PlsB.

We next sought to evaluate whether basal ppGpp concentrations are also capable of

regulating PlsB via post-translational control. As the effects of post-translational control are more

rapidly apparent than those that follow transcriptional regulation, we followed the timescale of

the response of the PL pathway to ppGpp synthesis. We measured ppGpp and PL synthesis

intermediates during RelA* induction to moderate levels. ppGpp increased within 10 minutes of

RelA* induction and reached its elevated steady-state concentration (~150 pmol/OD) by 20

minutes. Concentrations of PL intermediates PA and PS began to decrease within 10 minutes

(Figure 4C). Interestingly, PS decreases further after ~25 minutes, at which point PA increases

slightly, suggesting that PS may be back-converted to PA by PssA (28) **(Figure 4C).** The

immediate decay of intermediate pools indicates that ppGpp inhibits PlsB via post-translational

regulation at moderate concentrations. To verify that the immediate PA and PS depletion is too

fast to be explained by transcriptional regulation, we compared the PA and PS response kinetics

with the kinetics of a ppGpp-driven response known to be mediated by transcriptional control.

Cyclopropyl PL are produced from unsaturated fatty acids of membrane PL by the enzyme Cfa,

expression of which is induced transcriptionally by ppGpp (29). PL cyclopropylation begins 25

minutes after RelA* induction, much slower than the decrease in PA and PS. We conclude that

the rapid decrease in PA and PS 5-10 minutes after RelA* induction is mediated via post-

translational regulation of PlsB.

We used our model to simulate the observed response of PA and PS concentrations

immediately following mild RelA induction (up to 25 minutes post-induction) in order to estimate

the concentrations at which ppGpp inhibits PlsB. As the mode of inhibition is unknown (i.e.

competitive inhibition, mixed inhibition, or indirect inhibition via a ppGpp-responsive mediator),

we used an empirical allosteric inhibition term to obtain an effective K_I parameter (model details

provided in **Supplemental Methods)**. We estimate K_I to be approximately 300 pmol/OD, well

above the concentrations observed during the slowest steady-state growth condition used here

(60 pmol/OD at μ ~ 0.5 /hr). Direct ppGpp inhibition of PlsB may be relevant to PL flux during

extremely slow growth when ppGpp concentrations are above 150 pmol/OD, which may occur at

μ well below 0.5 hr^{-1}.

Discussion

Our comprehensive characterization and modelling of the fatty acid and PL biosynthesis pathways confirms known aspects of PL synthesis regulation and reveals essential principles critical for understanding how membrane synthesis is coupled with µ. Most importantly, we find that steady-state PL flux is coupled with µ via allosteric control of PlsB. PlsB demand controls flux into the fatty acid pathway through ACC via inhibition by PlsB substrates, consistent with metabolic control theory (30). Membrane synthesis is regulated very differently than protein synthesis, as membrane synthesis capacity (i.e. concentrations of PlsB and fatty acid synthesis pathway enzymes) is maintained at stable levels across a 3-fold range of µ. Counterintuitively, increased transcription of the fatty acid and PL synthesis pathways at higher µ (31) stabilizes enzyme concentrations, likely to keep pace with cell volume expansion. Maintaining a constant capacity for membrane synthesis enables the cell to address any change in membrane synthesis demand at the cost of expressing enzymes that remain inhibited at slow or moderate µ (32).

Both our model and our experimental data indicate that ACC and PlsB activities determine the concentrations of LPS precursor C14:0-OH-ACP, indicating a considerable degree of influence of PL synthesis over LPS synthesis in the absence of independent LPS flux regulation. Interestingly, LpxC concentrations decrease in response to higher µ, possibly due to increased degradation by the FtsH protease (33), although we cannot exclude that LpxC concentrations are also modulated between these conditions by transcriptional regulation. We propose that LpxC concentrations are regulated by proteolysis in response to PlsB activity to prevent toxic over-accumulation of LPS that might occur due to abundant precursor C14:0-OH-ACP. Incorporation of our data into models of PL-LPS pathway coordination could lead to further understanding of membrane biogenesis (34).

Although PlsB determines PL flux during steady-state growth, this control is not absolute,

as the fatty acid and PL pathways (and likely the LPS pathway) are highly sensitive to

fluctuations in acetyl-CoA concentrations and ACC activity. The sensitivity of PL flux to central

metabolism and even protein synthesis arrest renders membrane synthesis extremely

responsive to small environmental changes. While this sensitivity facilitates rapid adaptation of

PL flux to changes in μ, the tight connection between protein synthesis, central carbon

metabolism, and membrane biogenesis demands additional allosteric control (including inhibition

by ppGpp) to re-balance the pathway with μ and prevent PL overflow. If protein synthesis is

inhibited in a manner that does not decrease carbon flow into the fatty acid pathway (e.g. by

sudden nitrogen starvation), we hypothesize that continued PL and LPS synthesis would

outpace synthesis of the lipoproteins that tether the outer membrane to the peptidoglycan layer.

Inhibition of PL and LPS pathways by ppGpp would thus prevent production of excess

membrane, enforcing the coupling of PL and lipoprotein synthesis. Strains lacking the ppGpp

response (relaxed strains) have been observed to generate higher quantities of extracellular PL

and LPS, likely as outer membrane vesicles (35).

We estimate that post-translational control of PlsB by ppGpp, reflected by a relatively

high effective K_I (~300 pmol/OD), indicates that ppGpp regulates PL synthesis as an emergency

brake, and is unlikely to couple PL synthesis to growth in conditions that support $\mu > 0.5\ hr^{-1}$.

Although it is generally believed that ppGpp directly inhibits PlsB, multiple groups have been

unable to observe ppGpp inhibition of PlsB *in vitro* when acyl-ACP are used as substrates (36,

37). ppGpp may therefore control PlsB indirectly by modulating activity of a regulator or a cellular

process that interacts with PlsB.

Several candidates for the allosteric regulator of PlsB activity have been discovered. In

immunoprecipitation experiments, PlsB has been found to interact with proteins including ACP

and PssA, as well as several whose roles are unclear (PlsX and YbgC) (38), suggesting that

PlsB forms part of a PL synthesis complex that may couple PlsB activity with μ. As PL synthesis

flux has been observed to oscillate with the cell division cycle in *E. coli* and other bacteria (39),

PlsB may also be regulated by the divisome, membrane deformation, or by associations with the

Tol-Pal machinery. Degradation of PL by phospholipases may also play an important role in

membrane homeostasis during steady-state growth (40), as is known to occur during growth and

division in eukaryotes (41). In addition to identifying the allosteric regulators of PlsB, studies that

integrate connections between PL catabolism and transport with PL synthesis are needed for a

comprehensive understanding of membrane homeostasis.

Acknowledgements

We thank Michael Cashel for providing *E. coli* strain CF7974. We thank Christophe Danelon, Bertus Beaumont, and Frank Bruggeman for invaluable suggestions for improving the manuscript. This research was funded by a grant to GB from the Netherlands Organisation for Scientific Research (ALW Open 824.15.018) and an award from the Frontiers of Nanoscience program.

Author contributions

MN and GB designed the experiments. MN, FB, NvdB, NS, and GB performed experiments. MN, NvdB, and NI developed analytical methods. MN and GB analysed the data. GB supervised the research, constructed the mathematical model, and wrote the manuscript.

Materials and Methods

Culture conditions

Cultures were grown in 250-1000 mL Erlenmeyer flasks filled up to 10% of nominal volume with MOPS minimal medium (42) with 9.5 mM NH_4Cl or $^{15}NH_4Cl$ and 0.2% (w/v) carbon source (acetate, succinate, malate, glycerol, glucose, U-^{13}C-glucose (Cambridge Isotope Laboratories) or glucose supplemented with 0.1% Cas-amino acids). Culture flasks were placed in a Grant Instruments Sub Aqua Pro dual water bath at 37 °C and agitated by stirring with a 12 mm magnetic stir bar (VWR), coupled to a magnetic stir plate (2mag MIXdrive 1 Eco and MIXcontrol 20) set at 1200 rpm. Growth was monitored by optical density measurement at 600 nm using Ultrospec 10 Cell Density Meter (GE Healthcare). Samples for acyl-ACP, lipid analysis

1 and proteomics were collected using cultures without isotopic labelling. Samples for nucleotide

2 phosphate measurements were collected from U-^{15}N-labeled cultures and samples for G3P

3 measurements were collected from U-^{13}C-labeled cultures.

Strains and plasmid pRelA*

All experiments were performed using *Escherichia coli* K-12 strain NCM3722 (CGSC#

12355) and its derivatives. NCM3722 *relA::kan* was constructed by P1 phage transduction using

strain CF7974 (MG1655 Δ*lac* (*rph*$^+$) *relA255*::*kan)* as a donor. Plasmid pRelA* was created by

cloning DNA encoding residues 1-455 from the *E. coli* RelA protein into BglBrick plasmid

pBbS2k (43) (SC101* origin of replication, P$_{Tet}$ promoter, kanamycin resistance). The fluorescent

protein mVenus was fused by restriction-digestion to the C-terminus of RelA via a glycine-serine

linker.

Metabolite sampling

Samples for acyl-ACP, proteomics and lipid analysis were acquired by fast quenching of

1 mL of culture sample into 250 μL of ice-cold 10% trichloroacetic acid. After 10 min incubation

at 0°C cells were pelleted by centrifugation and stored at -80°C until analysis. For nucleotide

phosphates and polar metabolites analysis, samples were acquired by a modified fast vacuum

filtration method (44). 1 mL of culture was collected by vacuum on a pre-wetted 2.5 cm 0.45 μm

HV Durapore membrane filter. After rapid collection the filter was immediately placed upside-

down in quenching solution. For the measurement of nucleotide phosphates, 1 mL of ice-cold 2

M formic acid with 10 μL internal standards mix was used as a quenching solution, which was

subsequently neutralized by 25 μL of 28% ammonium hydroxide. For G3P measurements, 1 mL

of 50:30:20 (v/v/v) mixture of methanol, acetonitrile and water with 0.1% formic acid with 10 μL

internal standard solution (cooled on dry ice) was used as a quenching solution. After 10 min

incubation cells were washed from the filter, transferred to a tube and stored at -80°C until

analysis.

Preparation of internal standards

Isotopically-labeled internal standards (IS) were used to control for sampling and

measurement variation. For acyl-ACP and proteomics assays U-^{15}N *E. coli* whole cell extracts

were prepared using a MOPS minimal medium culture with $^{15}NH_4Cl$ as the sole nitrogen source.

At OD of ~0.5 10% TCA was added 1:4 to the culture to facilitate quenching of metabolism. After

10 min incubation on ice 10 mL single-use IS aliquots were collected by centrifugation and

stored at -80°C until the sample preparation. For the phospholipid measurement U-^{13}C lipid

extract was prepared using a culture grown in minimal MOPS medium with 0.2% U-^{13}C glucose

as the sole carbon source. At OD of ~0.5 10% TCA was added 1:4 to the culture and insoluble

cell material was collected by centrifugation after 10 min incubation on ice. Pellets were

resuspended in mixture consisting of 75 µL MeOH, 10 µL 15 mM citric acid/ 20 mM dipotassium

phosphate buffer and 250 µL of methyl-t-butyl ether per 1 mL of initial culture volume. After

vortexing and 10 min sonication phase separation was induced by addition of 70 µL/1mL of 15

mM citric acid/ 20 mM dipotassium phosphate buffer. After further vortexing, sonication and 10

minutes of incubation at room temperature phases were separated by 10 min centrifugation at

4000 rpm at room temperature. Upper phase was collected to a glass vial and stored at -20°C

until sample preparation.

Instrumentation

All LC/MS runs were performed using Agilent LCMS consisting of binary pump

(G1312B), autosampler (G7167A), temperature-controlled column compartment (G1316A), and

triple quadrupole (QQQ) mass spectrometer (G6460C) equipped with a standard ESI source, all

operated using MassHunter data acquisition software (version 7.0). Mass spectrometer operated

in dynamic MRM mode using transitions generated *in silico* by a script written in Python using

RDkit library using chemical structures of target compound as input. Transitions for targeted

proteomics assays were developed using Skyline (45) based on protein sequences from the

Uniprot database. Details of transition setting are included in **Supplemental Tables 3-7.**

LC/MS quantification of acyl-ACP intermediates

Acyl-ACP were measured using a published method (18) with minor modifications. Lysis

buffer was prepared by suspending appropriate number of frozen U-^{15}N-labeled *E. coli* pellets in

10 mL of 50 mM potassium phosphate buffer, pH 7.2, 6 M urea, 10 mM N-ethyl-maleimide, 5

mM EDTA and 1 mM ascorbic acid. 1 mL of lysis buffer was added to each of TCA-quenched

and pelleted cells and proteins were isolated by chloroform/methanol precipitation as described

previously. Protein pellets were resuspended in 10 µL of digestion buffer (4% 2-octyl-glucoside

in 25 mM potassium phosphate buffer, pH 7.2) and after adding 10 µL of 0.1 mg/mL GluC

protease (Promega) incubated overnight at 37°C. After quenching by addition of 5 µL MeOH,

samples were centrifuged and 10 µL was injected in LC/MS system. Separation was performed

on 2.1 mm x 50 mm 1.7 µm CSH C-18 column (Waters) held at 80°C using a binary gradient:

15% B, 3 minute ramp to 25%, 9 min increase to 95% and 1 minute hold at 95% B before 3

minute re-equilibration at starting conditions (A: 25 mM formic acid, B: 50 mM formic acid) at a

flow rate of 0.6 mL/min.

LC/MS quantification of phospholipids

Phospholipids sample preparation procedure is a combination of an MTBE extraction

method (46) and an established LC/MS method (47) Pelleted *E. coli* were resuspended in

mixture containing 150 µL of MeOH, 250 µL of U-^{13}C *E. coli* extract prepared as described

above and 250 µL MTBE. After vigorous vortexing and sonication 125 µL of 15 mM citric acid/

20 mM dipotassium phosphate buffer was added to homogenized pellets. Following further

vortexing and 10 min incubation at room temperature, liquid phases were separated by

centrifugation for 10 min at 20000g. 500 µL the of upper phase was moved to a new tube and

dried in a vacuum centrifuge (Labconco). Dried lipid films were resuspended in 10 µL 65:30:5

(v/v/v) isopropanol/acetonitrile/H_2O, supplemented with 10 mM acetylacetone. After

resuspension, 5 µL H_2O was added to reduce the organic content of the buffer and 5 µL of

resulting mixture was injected into the LC/MS system. Separation was performed on 2.1 mm x

50 mm 1.7 µm CSH C-18 column (Waters) at 60°C with a flow rate of 0.6 mL/min using the

following binary gradient: 25% B, ramp to 56%B in 6 min followed by linear increase to 80% B in

6 min, 2 min hold at 100% B and 3 min re-equilibration (A: 0.05% NH4OH in water, B: 0.05%

NH_4OH in 80% isopropanol 20% ACN).

LC/MS quantification of nucleotide phosphates

Frozen cell extracts were defrosted by 1-2 minute incubation in a 37°C water bath and

sonicated for 10 minutes in water ice slurry. After 10 minute centrifugation at 20000g, samples

were loaded on 1 mL/30 mg Oasis Wax cartridge (Waters) preconditioned with 1 mL of MeOH

and 1 mL 50 mM ammonium acetate buffer, pH 4.5. After washing with 1 mL ammonium acetate

buffer, analytes were eluted with 200 µL of 2.8% ammonium hydroxide in MeOH:ACN:H_2O

50:30:20 (v:v:v). After addition of 10 µL of 5% trehalose and brief vortexing samples were dried

down in a vacuum centrifuge (Labconco). Dried trehalose-stabilized extracts were re-dissolved

in 20 µL of MeOH:ACN:H_2O 50:30:20 (v:v:v) and moved to an autosampler vial for analysis.

Separation was performed on 2.1 mm x 100 mm 3.5 µm iHilic column (HILICON) or SeQuant

Zic-cHILIC, 2.1mm x 100 mm, 3µm (Merck) at 0.3 mL/min using the following binary gradient:

100% B, ramp to 85%B in 1.5 min followed by 10 min isocratic hold at 85% B and linear

decrease 30% B in 3 minutes with 2 minute hold at 30% B and 8 minute re-equilibration at initial

conditions. (A: 3.75 mM ammonium acetate, 1.25 mM acetic acid, 2 mM acetylacetone in MQ, B:

11.25 mM ammonium acetate, 3.75 mM acetic acid, 2 mM acetylacetone in 80% ACN). Injection

volume was 2 µL.

LC/MS quantification of G3P

Stored metabolite extracts were dried down in a vacuum centrifuge (Labconco), re-

dissolved in 20 µL of MeOH:ACN:H_2O 50:30:20 (v:v:v) and moved to an autosampler vial for

analysis. Separation was performed on 2.1 mm x 100 mm 3.5 µm iHilic column (HILICON) at 0.3

mL/min using the following binary gradient: 100% B, ramp to 80% B in 10 min followed by linear

decrease to 30% B in 3 min, 2 min hold at 30% B and 8 min re-equilibration. Injection volume

was 2 µL.

LC/MS targeted protein quantification

Relative concentrations of enzymes was measured by targeted proteomics using a

modified version of the acyl-ACP assay. Lysis buffer was prepared by suspending appropriate

number of frozen U-[15]N-labeled *E. coli* pellets in 10 mL of 50 mM potassium phosphate buffer,

pH 7.2 and 6 M urea. 1 mL of lysis buffer was added to each of TCA-quenched and pelleted

cells and proteins were isolated by chloroform/methanol precipitation as described previously.

Protein pellets were resuspended in 10 µL of digestion buffer (4% 2-octyl-glucoside in 25 mM

Tris buffer, pH 8.1 supplemented with 1 mM $CaCl_2$ and 5 mM TCEP). Alkylation of cysteine

residues was performed by adding 3 µL of 50 mM iodoacetamide followed by 15 minutes of

incubation in darkness. Digestion was performed by adding 10 µL of 0.2 mg/mL Trypsin Gold

(Promega) and overnight incubation at 37°C. Samples were centrifuged and 10 µL was injected

in LCMS system. Separation was performed on 2.1 mm x 50 mm 1.7 µm CSH C-18 column

(Waters) held at 40°C using a binary gradient: 2% B, 20 minute ramp to 25% B, 4 min increase

to 40% B, 0.5 ramp to 80% and 1 minute hold at 80% B before 3 minute re-equilibration at

starting conditions (A: 25 mL formic acid, B: 50 mM formic acid) at a flow rate of 0.5 mL/min.

Data Analysis

All LC-MS data files were processed in Skyline versions 4.x using target list based on *in silico* generated transition list. Each target compound had matching isotopically-labeled internal standard (IS). Processed data were exported as target compounds and IS peak areas and processed further using a set of Python scripts. Growth rates were obtained from linear fits to log-transformed growth curves. OD-corrected data were obtained by dividing the signal by OD_{600} value interpolated from growth curve at the time of sampling. PE-corrected results were produced by dividing the signal by sum all of signals for all phosphatidyl-ethanolamine species from the same measurement (in case of phospholipids) or matching sample (in case of other assays). In nucleotide phosphate and G3P assay absolute concentrations were estimated based on amounts of internal standards in IS-spike solution assuming RR = 1 implies equimolar amounts of target compound and IS at the moment of quenching. Concentrations of IS in the spike mix are provided in the supplement.

Mathematical modeling

The computational model was constructed and tested using COPASI version 4.24 (48). Full details of the model are described in the **Supplemental Methods**.

References

1.   Dowhan W (2013) A retrospective: use of Escherichia coli as a vehicle to study phospholipid synthesis and function. *Biochim Biophys Acta* 1831(3):471–494.

2.   Chubukov V, et al. (2013) Transcriptional regulation is insufficient to explain substrate-induced flux changes in Bacillus subtilis. *Mol Syst Biol* 9. doi:Artn 70910.1038/Msb.2013.66.

3.   Hackett SR, et al. (2016) Systems-level analysis of mechanisms regulating yeast metabolic flux. *Science (80-)* 354(6311). doi:10.1126/science.aaf2786.

4. Scott M, Gunderson CW, Mateescu EM, Zhang Z, Hwa T (2010) Interdependence of cell
growth and gene expression: Origins and consequences. *Science (80-)*.
doi:10.1126/science.1192588.

5. Kamio Y, Nikaido H (1976) Outer Membrane of Salmonella Typhimurium: Accessibility of
Phospholipid Head Groups to Phospholipase C and Cyanogen Bromide Activated
Dextran in the External Medium. *Biochemistry* 15(12):2561–2570.

6. Funahara Y, Nikaido H (1980) Asymmetric localization of lipopolysaccharides on the outer
membrane of Salmonella typhimurium. *J Bacteriol* 141(3):1463–1465.

7. Suzuki H, et al. (1978) Murein-lipoprotein of Escherichia coli: A protein involved in the
stabilization of bacterial cell envelope. *MGG Mol Gen Genet* 167(1):1–9.

8. Raetz CR, Larson TJ, Dowhan W (1977) Gene cloning for the isolation of enzymes of
membrane lipid synthesis: phosphatidylserine synthase overproduction in Escherichia
coli. *Proc Natl Acad Sci U S A* 74(4):1412–1416.

9. Ohta A, Waggoner K, Louie K, Dowhan W (1981) Cloning of genes involved in membrane
lipid synthesis. Effects of amplification of phosphatidylserine synthase in Escherichia coli.
J Biol Chem.

10. Ganong BR, Raetz CR (1982) Massive accumulation of phosphatidic acid in conditionally
lethal CDP-diglyceride synthetase mutants and cytidine auxotrophs of Escherichia coli. *J
Biol Chem* 257(1):389–394.

11. Heacock PN, Dowhan W (1987) Construction of a lethal mutation in the synthesis of the
major acidic phospholipids of Escherichia coli. *J Biol Chem* 262(27):13044–13049.

12. Increased synthesis of phosphatidylserine decarboxylase in a strain of Escherichia coli
bearing a hybrid plasmid. Altered association of enzyme with the membrane. (1979) *J Biol
Chem* 254(3):627–633.

13. Dennis PP, Bremer H (2008) Modulation of Chemical Composition and Other Parameters
of the Cell at Different Exponential Growth Rates. *EcoSal Plus* 3(1).
doi:10.1128/ecosal.5.2.3.

14. Sud IJ, Schaechter M (1964) Dependence of the content of cell envelopes on the growth
rate of Bacillus megaterium. *J Bacteriol* 88(6):1612–1617.

15. Ballesta JPG, Schaechter M (1972) Dependence of the Rate of Synthesis of Phosphatidylethanolamine and Phosphatidylglycerol on the Rate of Growth of Escherichia coli. *J Bacteriol* 110(1):452–453.

16. Schulman H, Kennedy EP (1977) Relation of turnover of membrane phospholipids to synthesis of membrane derived oligosaccharides of Escherichia coli. *J Biol Chem* 252(12):4250–4255.

17. Harris LK, Theriot JA (2016) Relative rates of surface and volume synthesis set bacterial cell size. *Cell* 165(6):1479–1492.

18. Noga MJ, et al. (2016) Mass-Spectrometry-Based Quantification of Protein-Bound Fatty Acid Synthesis Intermediates from Escherichia coli. *J Proteome Res* 15(10):3617–3623.

19. Gerhardt ECM, et al. (2015) The Bacterial signal transduction protein GlnB regulates the committed step in fatty acid biosynthesis by acting as a dissociable regulatory subunit of acetyl-CoA carboxylase. *Mol Microbiol* 95(6):1025–1035.

20. Davis MS, Cronan J (2001) Inhibition of Escherichia coli acetyl coenzyme A carboxylase by acyl-acyl carrier protein. *J Bacteriol.* doi:10.1128/JB.183.4.1499-1503.2001.

21. Wahl A, My L, Dumoulin R, Sturgis JN, Bouveret E (2011) Antagonistic regulation of dgkA and plsB genes of phospholipid synthesis by multiple stress responses in Escherichia coli. *Mol Microbiol* 80(5):1260–1275.

22. Rhodius VA, Suh WC, Nonaka G, West J, Gross CA (2006) Conserved and variable functions of the σEstress response in related genomes. *PLoS Biol.* doi:10.1371/journal.pbio.0040002.

23. Schreiber G, et al. (1991) Overexpression of the relA gene in Escherichia coli. *J Biol Chem* 266(6):3760–3767.

24. Heath RJ, Jackowski S, Rock CO (1994) Guanosine Tetraphosphate Inhibition of Fatty-Acid and Phospholipid-Synthesis in Escherichia coli Is Relieved by Overexpression of Glycerol-3-Phosphate Acyltransferase (PlsB). *J Biol Chem* 269(42):26584–26590.

25. Sorensen PG, et al. (1996) Regulation of UDP-3-O-[R-3-hydroxymyristoyl]-N-acetylglucosamine Deacetylase in Escherichia coli: THE SECOND ENZYMATIC STEP OF LIPID A BIOSYNTHESIS . *J Biol Chem* 271(42):25898–25905.

1   26.   Merlie JP, Pizer LI (1973) Regulation of phospholipid synthesis in Escherichia coli by
2   guanosine tetraphosphate. *J Bacteriol* 116(1):355–366.

3   27.   Nunn WD, Cronan Jr. JE (1976) Regulation of membrane phospholipid synthesis by the
4   relA gene: dependence on ppGpp levels. *Biochemistry* 15(12):2546–2550.

5   28.   Larson TJ, Dowhan W (1976) Ribosomal-Associated Phosphatidylserine Synthetase from
6   Escherichia coli: Purification by Substrate-Specific Elution from Phosphocellulose Using
7   Cytidine 5′-Diphospho-1,2-diacyl-sn-glycerol. *Biochemistry* 15(24):5212–5218.

8   29.   Eichel J, Chang YY, Riesenberg D, Cronan JE (1999) Effect of ppGpp on Escherichia coli
9   cyclopropane fatty acid synthesis is mediated through the RpoS sigma factor (σ(S)). *J*
10   *Bacteriol* 181(2):572–576.

30.   Hofmeyr JHS, Cornish-Bowden A (2000) Regulating the cellular economy of supply and
demand. *FEBS Lett.* doi:10.1016/S0014-5793(00)01668-9.

31.   Li SJ, Cronan JE (1993) Growth rate regulation of Escherichia coli acetyl coenzyme A
carboxylase, which catalyzes the first committed step of lipid biosynthesis. *J Bacteriol*
175(2):332–340.

32.   Davidi D, Milo R (2017) Lessons on enzyme kinetics from quantitative proteomics. *Curr*
Opin Biotechnol 46:81–89.

33.   Ogura T, et al. (1999) Balanced biosynthesis of major membrane components through
regulated degradation of the committed enzyme of lipid A biosynthesis by the AAA
protease FtsH (HflB) in Escherichia coli. *Mol Microbiol* 31(3):833–844.

34.   Emiola A, Andrews SS, Heller C, George J (2016) Crosstalk between the
lipopolysaccharide and phospholipid pathways during outer membrane biogenesis in
Escherichia coli. *Proc Natl Acad Sci* 113(11):3108–3113.

35.   Crowfoot PD, Esfahani M, Wakil SJ (1972) Relation Between Protein Synthesis and
Phospholipid Synthesis and Turnover in Escherichia coli. *J Bacteriol* 112(3):1408–1415.

36.   Lueking DR, Goldfine H (1975) The involvement of guanosine 5-diphosphate-3-
diphosphate in the regulation of phospholipid biosynthesis in Escherichia coli. Lack of
ppGpp inhibition of acyltransfer from acyl-ACP to sn-glycerol 3-phosphate. *J Biol Chem*
250(13):4911–4917.

37. Nunn WD, Kelly DL, Stumfall MY (1977) Regulation of fatty acid synthesis during the cessation of phospholipid biosynthesis in Escherichia coli. *J Bacteriol* 132(2):526–531.

38. Gully D, Bouveret E (2006) A protein network for phospholipid synthesis uncovered by a variant of the tandem affinity purification method in Escherichia coli. *Proteomics* 6(1):282–293.

39. Carty CE, Ingram LO (1981) Lipid synthesis during the Escherichia coli cell cycle. *J Bacteriol* 145(1):472–478.

40. May KL, Silhavy TJ (2018) The Escherichia coli phospholipase PldA regulates outer membrane homeostasis via lipid signaling. *MBio* 9(2). doi:10.1128/mBio.00379-18.

41. Hermansson M, Hokynar K, Somerharju P (2011) Mechanisms of glycerophospholipid homeostasis in mammalian cells. *Prog Lipid Res* 50(3):240–257.

42. Neidhardt FC, Bloch PL, Smith DF (1974) Culture Medium for Enterobacteria. *J Bacteriol* 119(3):736–747.

43. Lee T, et al. (2011) BglBrick vectors and datasheets: A synthetic biology platform for gene expression. *J Biol Eng* 5(1):12.

44. Link H, Kochanowski K, Sauer U (2013) Systematic identification of allosteric protein-metabolite interactions that control enzyme activity in vivo. *Nat Biotechnol* 31(4):357–361.

45. MacLean B, et al. (2010) Skyline: An open source document editor for creating and analyzing targeted proteomics experiments. *Bioinformatics* 26(7):966–968.

46. Matyash V, Liebisch G, Kurzchalia T V, Shevchenko A, Schwudke D (2008) Lipid extraction by methyl-tert-butyl ether for high-throughput lipidomics. *J Lipid Res* 49(5):1137–1146.

47. Susanto O, et al. (2017) LPP3 mediates self-generation of chemotactic LPA gradients by melanoma cells. *J Cell Sci* 130(20):3455–3466.

48. Hoops S, et al. (2006) COPASI - A COmplex PAthway SImulator. *Bioinformatics* 22(24):3067–3074.

Figures

Figure 1. Characterization of the *E. coli* fatty acid and PL synthesis pathways during steady-

state growth. **A.** *E. coli* fatty acid and PL synthesis pathways highlighting the metabolites and

enzymes discussed in the main text. Fatty acid synthesis is initiated by carboxylation of acetyl-

CoA by acetyl-CoA carboxylase complex (ACC) to produce malonyl-CoA, from which the

malonyl group is transferred to holo-ACP by FabD. Fatty acids are then elongated by iterated

cycles of condensation of acyl-ACP with malonyl-ACP, followed by reduction and dehydration.

sn-glycerol-3-phosphate (G3P) is acylated by PlsB primarily using long-chain acyl-ACP species

C16:0-ACP or C18:1-ACP to yield lysophosphatidic acid (LPA), which is acylated by PlsC using

C16:1-ACP or C18:1-ACP to yield phosphatidic acid (PA). PA is converted by CdsA to (d)CDP-

diacylglycerol ((d)CDP-DAG) using either CTP or dCTP. At the branch point in PL synthesis,

PgsA combines G3P with (d)CDP-DAG to produce phosphatidylglycerol phosphate (PGP).

Dephosphorylation of PGP by PgpA yields PG, which typically comprises 25% of total PL. Two

PG molecules may be further converted to CL, which accounts for ~5% of total PL. (d)CDP-DAG

is also converted by PssA to phosphatidylserine (PS), which is decarboxylated by Psd to yield

PE (70% of total PL). **B, C.** Growth rate (μ)-dependent abundance of total PG, PE, and CL (**B**)

and μ-dependence of PE flux (**C**), normalized to average values across conditions for each

strain. **D.** Steady-state concentrations of acetyl-CoA and ppGpp. **E, F, G.** Concentrations of G3P

and acyl-ACP species **(E)**, PL intermediates **(F)**, and PL synthesis pathway enzymes **(G)** as

functions of PL flux. Concentrations of soluble substrates and enzymes with exclusively soluble

substrates are calculated using cell volume (proportional to OD). For membrane-bound

phospholipid intermediates and for enzymes that react with membrane-bound substrates (all

enzymes following PlsB in the PL synthesis pathway) we calculate concentration within the

membrane volume, proportional to total PE. S/OD, E/OD, S/PE, and E/PE are defined as the

concentration of a substrate S or enzyme E (amount per OD or per PE) relative to its average

concentration across all conditions measured in that strain (either wild-type (WT) or pRelA).

Values are log(2)-transformed. Error bars represent the standard deviation of measurements

from three samples of one culture.

Figure 2. Simulated steady-state fluxes and metabolite concentrations of fatty acid and PL

pathways identifies PlsB as site of flux control during steady-state growth. **A.** Reactions

simulated in the model. For simplicity, only reactions late in the saturated fatty acid pathway are

included, and branching of the PL pathway into PE and PG is not included. Each reaction is

modelled as an irreversible one- or two-substrate Michaelis-Menten reaction. ACC activity is

competitively inhibited by C16:0-ACP. Reactions catalysed by FabI/FabZ and LpxA/LpxC are

considered in the model as single reactions represented by "FabZ" and "LpxC", respectively. **B.**

Response of PE and LPS fluxes to varying V_{max} parameters of pathway enzymes and acetyl-

CoA and C16:1-ACP concentrations. 4-fold variations in C14:0-β-keto-ACP synthesis, FabG,

CdsA, PssA, and Psd V_{max} did not change PE or LPS flux and are therefore not shown. **C.**

Simulated changes in metabolite concentrations in response to variations in PlsB and ACC V_{max}

and acetyl-CoA overlaid on experimentally-measured concentrations (data from **Figure 1**). Line

plots are offset to prevent overlap. Differential equations and parameters are provided in the

1 **Supplemental Methods**.

Figure 3. Response of the fatty acid and PL synthesis pathways to translational inhibition. **A, B.**

Responses of acyl-ACP **(A)** and PL intermediate pools **(B)** to mupirocin treatment. Mupirocin

was added at 0 min (indicated by dashed lines) to glucose cultures of *E. coli* wild-type and Δ*relA*.

C, D. Addition of translation inhibitor chloramphenicol or the transcription inhibitor rifampicin to

glycerol cultures of wild-type *E. coli* (indicated by dashed lines at 0 min) leads to an influx of

carbon into the fatty acid pathway as suggested by a decrease in holo-ACP levels and an

increase in unsaturated long-chain acyl-ACP species **(C).** The pulse of carbon continues further

into the PL synthesis pathway, as indicated by transient increases in total PA and PS **(D).**

Trajectories in **A-D** are each obtained from individual cultures. **E.** Chloramphenicol and

rifampicin both trigger a rapid decrease in ppGpp and accumulation of ACC and FabH substrate

acetyl-CoA. Values depicted in **E** are averages of triplicates from single cultures.

1
Figure 4. PlsB activity is suppressed by moderate to high concentrations of ppGpp via post-

translational inhibition. **A, B.** Response of acyl-ACP **(A)** and PL intermediate pools **(B)** to

maximal induction of RelA*. RelA* expression was induced by addition of 40 ng/mL doxycycline

at -10 minutes (dashed blue line), followed 10 minutes later by chloramphenicol addition (dashed

grey line at 0 min). **C.** Response of PL intermediates and cyclopropyl-PE to mild induction of

RelA* expression by addition of 1 ng/mL doxycycline (dashed grey line at 0 min). Superimposed

are simulated kinetic data assuming allosteric inhibition of PlsB by ppGpp with an empirical

allosteric inhibition constant K_I = 300 pmol/OD. Simulation details are included in the

Supplemental Methods.