Post-translational control of PlsB is sufficient to coordinate membrane synthesis with growth in *Escherichia coli*

Running title: Economy of membrane synthesis in E. coli

Marek J Noga[†], Ferhat Büke, Niels JF van den Broek[‡], Nicole Imholz, Nicole Scherer, Flora Yang, Gregory Bokinsky*

Department of Bionanoscience, Kavli Institute of Nanoscience, Delft University of Technology, Delft, The Netherlands

Materials and correspondence: *e-mail: g.e.bokinsky@tudelft.nl

[†] Current address: Translational Metabolic Laboratory, Department of Laboratory Medicine, Radboudumc, Nijmegen, The Netherlands

[‡] Current address: Vaccine Process & Analytical Development, Janssen Vaccines & Prevention B.V., Leiden, The Netherlands

Abstract

Every cell must produce enough membrane to contain itself. However, the mechanisms by which the rate of membrane synthesis is coupled with the rate of cell growth remain unresolved. By comparing substrate and enzyme concentrations of the fatty acid and phospholipid synthesis pathways of *Escherichia coli* across a 3-fold range of carbon-limited growth rates, we show that the rate of membrane phospholipid synthesis during steady-state growth is determined principally through allosteric control of a single enzyme, PlsB. Due to feedback regulation of the fatty acid pathway, PlsB activity also indirectly controls synthesis of lipopolysaccharide, a major component of the outer membrane synthesized from a fatty acid synthesis intermediate. Surprisingly, concentrations of the enzyme that catalyses the committed step of lipopolysaccharide synthesis (LpxC) do not vary across steady-state growth conditions, suggesting that steady-state lipopolysaccharide synthesis is modulated primarily via indirect control by PlsB. In contrast to steady-state regulation, we find that responses to environmental perturbations are triggered directly via changes in acetyl-CoA concentrations, which enables rapid adaptation. Adaptations are further modulated by ppGpp, which regulates PlsB activity during slow growth and growth arrest. The strong reliance of the membrane synthesis pathway upon post-translational regulation ensures both reliability and responsiveness of membrane synthesis.

Significance

How do bacteria cells grow without breaking their membranes? Although the biochemistry of fatty acid and membrane synthesis is well-known, how membrane synthesis is balanced with growth and metabolism has remained unclear. This is partly due to the many control points that have been discovered within the membrane synthesis pathways. By precisely establishing the contributions of individual pathway enzymes, our results simplify the model of membrane biogenesis in the model bacteria species *Escherichia coli*. Specifically, we find that allosteric control of a single enzyme, PlsB, is sufficient to balance growth with membrane synthesis and to ensure that growing *E. coli* produces sufficient membrane. Identifying the signals that activate and deactivate PlsB will answer the question of how membrane synthesis is synchronized with growth.

Introduction

All cells build and expand their membranes at a pace that must be coordinated with growth, as excess or insufficient membrane production can be fatal. In Gram-negative bacteria, construction of the double membrane from fatty acid precursors demands coordination between phospholipid (PL) and lipopolysaccharide (LPS) synthesis pathways (1, 2) as well as with protein synthesis, which supplies the lipoproteins that tether the outer membrane to the peptidoglycan cell wall (3). Many elements of membrane synthesis regulation that act on either transcription or enzymes activities have been proposed or identified. However, how each of these individual elements are used to regulate membrane synthesis during steady-state growth remains unclear (4).

The question of how membrane construction is coordinated with growth can be approached by considering how the metabolic pathways that synthesize membrane building blocks (PL and LPS) are regulated. Biosynthetic fluxes are regulated either by control of enzyme concentrations or by direct control of enzyme activity. Examples of both forms of regulation can be readily found: in *Escherichia coli,* the steady-state protein synthesis rate is controlled by ribosome concentration, which is transcriptionally regulated to balance amino acid supply with translation demand (5). In contrast, fluxes through central carbon metabolism are controlled instead by concentrations of substrates and inhibitors (i.e. post-translational control) rather than by changes in enzyme concentration (6, 7). Post-translational control is also known to contribute to membrane synthesis regulation (8–13), however, how cells use transcriptional and post-translational control to coordinate membrane synthesis with growth has never been well-defined.

In order to understand how the model Gram-negative species *Escherichia coli* coordinates membrane synthesis with growth, we quantified substrates and enzymes of the fatty acid and PL synthesis pathways in both steady-state and dynamic conditions. We find that post-translational control of the first enzyme in the PL synthesis pathway, PlsB, is sufficient to ensure

1   steady-state regulation of PL synthesis. In contrast, transcriptional regulation maintains stable

2   enzyme concentrations regardless of growth rate. Furthermore, due to feedback regulation of

3   the fatty acid pathway, PlsB exerts strong control over LPS synthesis via its effects on

4   concentrations of an LPS fatty acid precursor.

Results

The PL to biomass ratio varies inversely with μ

Lipid precursors of both LPS and PL are synthesized in the cytosol as fatty acyl thioesters covalently attached to acyl carrier protein (acyl-ACP). Fatty acid synthesis is initiated by carboxylation of acetyl-CoA by the acetyl-CoA carboxylase complex (ACC) to produce malonyl-CoA, from which the malonyl group is transferred to holo-ACP by FabD. Malonyl-ACP is used to elongate fatty acids iterated cycles beginning with condensation with acyl-ACP, reduction, dehydration, and reduction. The first step in PL synthesis is catalysed by the membrane-bound enzyme PlsB, which synthesizes lysophosphatidic acid (LPA) from long-chain acyl-ACP and *sn*-glycerol-3 phosphate (G3P). G3P is produced either from glyceraldehyde-3-phosphate, an intermediate of central carbon metabolism, or via glycerol catabolism **(Figure 1A)**. We grew cultures of *E. coli* NCM3722 in 6 defined media that support a 3-fold range of growth rates (μ). Quantities of major PL species phosphatidylglycerol (PG), phosphatidylethanolamine (PE), and cardiolipin (CL) per unit biomass (as determined by culture optical density (OD) (14)) decrease slightly with increasing μ **(Figure 1B)**, consistent with previous observations in *E. coli* and other bacteria (15, 16). The PG/PE ratio remains constant across all μ. We quantified steady-state PL flux by multiplying total PE LC/MS counts by μ, which closely approximates the total PL synthesis rate as PE turnover is slow relative to synthesis (17) and PG/PE ratio is stable. PL flux increases by 2-fold as μ increases by 3-fold **(Figure 1C)**. The higher PL to biomass ratio of slow-growing cells likely reflects their higher surface-to-volume ratio (18).

Growth regulates PL flux via post-translational control of PlsB

We first sought to identify the control points of PL synthesis by determining correlations between PL flux and concentrations of pathway intermediates during steady-state growth. If PL

and fatty acid synthesis rates are controlled by substrate concentrations, a positive correlation

between substrate concentrations and flux should be observed. We quantified nucleotides, G3P,

acyl-ACP, and PL synthesis intermediates using LC/MS (19). No correlation was observed

between concentrations of the fatty acid precursor acetyl-CoA and μ (**Figure 1D**). We also found

no significant positive correlation between PL flux and PlsB substrates C16:0-ACP or C18:1-

ACP (Pearson correlation coefficients r <0.4, two-tailed test of significance p >0.1), while C18:0-

ACP concentrations correlate negatively with PL flux (r = -0.76, p <0.01, **Figure 1E, Table 1).**

Although G3P concentrations correlate with PL flux (r >0.99, p <0.01), growth in glycerol

medium increases G3P by 20-fold without increasing PL abundance or flux. The absence of any

significant positive correlation between ACP substrate concentrations and PL flux indicates that

steady-state PL synthesis is not controlled by substrate concentrations.

In contrast to PL precursors, concentrations of the fatty acid initiation and elongation

substrate malonyl-ACP correlate with PL flux **(Figure 1E,** r = 0.80, p <0.01, **Table 1)**, while the

concentrations of most saturated ACP species remain relatively constant **(Supplemental Figure**

1) suggesting that the fatty acid initiation and elongation reactions are controlled primarily by

malonyl-ACP concentrations and thus by the rate of malonyl-ACP synthesis by ACC and FabD.

Concentrations of C14:0-OH-ACP, the lipid precursor of LPS, also correlate with PL flux (r =

0.94, p <10^{-4}), suggesting that PL synthesis is naturally coupled with LPS synthesis via

concentrations of C14:0-OH-ACP **(Figure 1E)**.

As with substrates, positive correlations between enzyme concentrations and reaction

flux may indicate that a reaction rate is controlled by enzyme concentrations, i.e. via

transcriptional control. We measured concentrations of fatty acid synthesis pathway enzymes

using LC/MS to determine whether fatty acid flux is coupled with μ by transcription-based

regulation. While concentrations of one of the four ACC subunits (AccA) and several of the fatty

acid synthesis pathway enzymes correlate with PL flux **(Table 1, Supplemental Figure 2),** the

increases are small (30%) relative to the PL flux increase (2-fold). We infer that malonyl-ACP

synthesis and thus the rate of fatty acid synthesis are determined primarily by post-translational

control of ACC. The negative correlation between C18:0-ACP and PL flux is consistent with

long-chain acyl-ACP inhibiting malonyl-CoA synthesis by ACC via negative feedback inhibition

(20).

Unlike long-chain acyl-ACP, the concentrations of all PL synthesis intermediates

downstream of PlsB correlate positively with PL flux **(Figure 1F,** r = 0.61-0.99, **Table 1,**

Supplemental Figure 3). The close match between the increase in concentration and PL flux is

consistent with the rates of these reactions being under substrate control. The contrast between

the trends in concentrations of the substrates and the products of PlsB implies that PlsB activity

alone controls PL synthesis. PlsB activity might be regulated either via transcriptional control of

PlsB concentration or via post-translational control of PlsB activity. Transcription of *plsB* is

controlled by the membrane stress-activated sigma factor RpoE (21) and by the µ-sensitive

regulator guanosine tetraphosphate (ppGpp) (22). Concentrations of ppGpp are inversely

correlated with µ **(Figure 1D)**, implying that PL flux may be coupled to µ by transcriptional

control of the *plsB* gene by ppGpp. However, we found no correlation between PL flux and PlsB

concentration (**Table 1,** p >0.1). Instead, PlsB concentrations are nearly constant across all

conditions studied **(Figure 1G)**. The invariance of PlsB concentrations over a 2-fold range of PL

flux indicates that PlsB activity is regulated mainly via post-translational control. Unexpectedly,

concentrations of LpxC, which catalyses the committed step in the LPS pathway (23), also do

not correlate with PL flux (**Table 1,** p >0.1) and also remain stable **(Figure 1G),** indicating that

flux into the LPS pathway is not varied between steady-state growth conditions by changes in

LpxC concentration.

µ can be modulated independently of nutrient conditions by artificially varying ppGpp.

Titrating ppGpp above basal concentrations, but below concentrations occurring during

starvation, reduces steady-state μ by reducing ribosomal RNA synthesis (24). Furthermore, PlsB

activity is inhibited at high ppGpp concentrations observed during amino acid starvation (25). We

titrated μ by expressing the catalytic domain of the ppGpp synthesis enzyme RelA (RelA*) from

an inducible promoter (P_{Tet}) in glucose medium. As previously observed (24), RelA*-titrated

cultures exhibit ppGpp concentrations that are elevated 2-fold above wild-type cultures growing

at similar rates **(Figure 1D).** Trends in PL abundance, PL flux, acyl-ACP species, PL

intermediates, and PL synthesis enzyme concentrations in ppGpp-limited cultures closely follow

the trends observed when μ is varied by carbon source **(Figure 1B-C, 1E-G, Supplemental**

Figures 1, 2, 3). Notably, despite ppGpp regulation of *plsB* transcription, concentrations of PlsB

also do not substantially vary with increasing concentrations of steady-state ppGpp.

Interestingly, LpxC concentrations decrease with increasing PL flux in these conditions **(Figure**

1G). The trends in substrate, enzyme, and intermediate concentrations observed in ppGpp-

limited cultures are consistent with growth regulating PL flux via post-translational control – and

not transcriptional control – of PlsB.

Mathematical modelling supports PlsB control over steady-state PL and LPS synthesis

To test whether regulation of PlsB activity is sufficient to control steady-state PL

synthesis, we constructed a simplified differential equation model that describes fatty acid, LPS

initiation, and PL biosynthesis **(Figure 2A).** The model includes competitive inhibition of ACC by

both C16:0-ACP and C18:0-ACP as the sole regulatory interactions. The model also includes a

branch point at C14:0-OH-ACP into LPS synthesis. LPS initiation is represented in the model by

a single step that combines reactions catalysed by LpxA and LpxC, as LpxC catalyses the first

irreversible reaction in the LPS pathway. Concentrations of G3P and C16:1-ACP are fixed in the

model to reflect experimental observations of PlsB saturation by G3P and invariance of C16:1-

ACP concentrations, respectively (see **Supplementary Methods** for model details, parameter

sets, and sensitivity analysis).

We used the model to identify enzymes and substrates that exert control over PL and

LPS synthesis. Increasing either ACC or PlsB V_{max} by 4-fold increased simulated PL and LPS

fluxes in parallel by 2-fold **(Figure 2B)**, while changing V_{max} of all other enzymes in the simulated

pathway had little or no effect on simulated PL flux. The insensitivity of PL flux to V_{max} of all fatty

acid and PL synthesis enzymes aside from ACC and PlsB suggests that the experimentally-

observed positive correlations between enzyme concentrations and PL flux do not actually

increase PL flux. The unexpectedly strong control of LPS flux by PlsB is due to the natural

coupling between PL flux and concentrations of the LPS synthesis substrate C14:0-OH-ACP.

Changes in the C14:0-OH-ACP dehydration/reduction V_{max} (catalysed in *E. coli* by FabZ and

FabI, respectively) and LPS initiation V_{max} (catalysed by LpxA and LpxC) exert strong and

opposing control over LPS flux. However, LpxC and FabZ do not exert significant control over

PL synthesis, indicating that substantial flux can be diverted into the LPS pathway without

depleting PL synthesis. Of the two substrates considered in the model (acetyl-CoA and C16:1-

ACP), only acetyl-CoA affects PL and LPS flux **(Figure 2B).**

To determine which of the two enzymes with non-negligible PL flux control – ACC or

PlsB – actually controls PL flux during steady-state growth, we compared the predicted trends in

substrate concentrations caused by simulated V_{max} variations against experimentally-observed

trends. The predicted trends driven by ACC and PlsB V_{max} variation closely reproduce most

experimentally-observed trends in both fatty acid and PL intermediate concentrations **(Figure**

2C). However, variations in ACC and PlsB V_{max} cause opposing trends in predicted

concentrations of PlsB substrates C16:0-ACP and C18:0-ACP, allowing ACC- and PlsB-based

regulation to be distinguished. Increasing ACC V_{max} causes C16:0-ACP and C18:0-ACP

concentrations to increase with PL flux, which contradicts our experimental observations.

However, increasing PlsB V_{max} is predicted to decrease C16:0-ACP and C18:0-ACP

concentrations, a trend that better follows the experimentally-observed trends (**Figure 2C).** We

therefore conclude that PL synthesis is primarily regulated by PlsB during steady-state growth.

Our model also suggests that PlsB control over steady-state LPS flux (achieved by its tight

control over C14:0-OH-ACP concentrations) may be sufficient to couple LPS synthesis with

growth as well.

Translation inhibition causes carbon overflow into fatty acid synthesis

We set out to evaluate whether a known regulator of PL synthesis, ppGpp, is able to

directly regulate PlsB activity during steady-state growth. High concentrations of ppGpp lead to

PlsB inhibition *in vivo* (25), although it remains unclear whether this inhibition acts via a ppGpp-

PlsB interaction or via an indirect route. Low concentrations of ppGpp correlate inversely with μ

(Figure 1D). The notion that ppGpp might directly control PL synthesis even at the low basal

concentrations present during steady-state growth has been proposed but never tested (26).

Acyl-ACP substrates of PlsB increase as ppGpp is titrated upwards with RelA* while the product

of PlsB (LPA) decreases, consistent with ppGpp inhibiting flux into the PL pathway (**Figure 1E,**

1F), and closely following simulated variations in PlsB V_{max} **(Figure 2C).** Therefore, changes in μ

driven by sub-stringent concentrations of ppGpp must somehow influence steady-state PlsB

activity. However, the mode of control by ppGpp (transcriptional control of *plsB*, post-

translational inhibition of PlsB, or indirect control via an unknown regulator of PlsB) cannot be

determined from steady-state data alone.

We first wished to observe the effects of high concentrations of ppGpp on the fatty acid

and PL synthesis pathways. Synthesis of high concentrations of ppGpp by RelA (the stringent

response) is triggered by any stress or starvation conditions that lead to the specific biochemical

cue of uncharged tRNA bound to the ribosome. During the stringent response, ppGpp

accumulates by more than 10-fold over basal concentrations and PL synthesis rates are reduced

by approximately half (25, 26), likely due to PlsB inhibition (27). We triggered ppGpp synthesis

by adding the tRNA aminoacylation inhibitor mupirocin to glucose cultures of wild-type *E. coli.* To

distinguish effects of ppGpp from effects of translation inhibition, mupirocin was also added to a

culture of Δ*relA E. coli*. Mupirocin causes a 10-fold accumulation of ppGpp in the wild-type strain

within 1 minute to over 400 pmol/OD, reaching >800 pmol/OD after 3 minutes **(Supplemental**

Figure 4). Unexpectedly, mupirocin treatment also transiently increased malonyl-ACP and all

hydroxyl-ACP species at the expense of holo-ACP in both strains **(Figure 3A, Supplemental**

Figure 5). The increases in malonyl- and hydroxyl-ACP concentrations are matched by a

corresponding increase in PL synthesis intermediates, suggesting that mupirocin diverts a pulse

of carbon into the fatty acid and PL synthesis pathways **(Figure 3B, Supplemental Figure 6)**. In

the wild-type strain, the pulse of fatty acid synthesis activity is rapidly followed by C14:0-, C16:0-,

and C18:0-ACP accumulation, consistent with ppGpp inhibition of PlsB **(Figure 3A,**

Supplemental Figure 5). C16:0-ACP accumulation is followed in turn by an increase in holo-

ACP and a decrease in malonyl-ACP, consistent with ACC inhibition **(Figure 3A)**. PL

intermediates PA, PS, and PGP are rapidly depleted in the wild-type strain after briefly

increasing **(Figure 3B).** The transient carbon influx briefly shifts the population of the acyl chains

incorporated into PL towards longer-chain fatty acids, likely due to increased malonyl-ACP

concentrations favouring fatty acid elongation over membrane incorporation by PlsB and PlsC

(Supplemental Figure 6). While suppression of fatty acid synthesis in the wild-type strain can

be attributed to ppGpp, it is unclear what attenuates the carbon influx in the Δ*relA* strain;

however C14:0-ACP accumulation after 10 minutes likely contributes to ACC inhibition. Inhibition

of fatty acid elongation also depletes the LPS precursor C14:0-OH-ACP, which is expected to

decrease LPS synthesis in parallel with PL synthesis **(Figure 3A)**.

As malonyl-ACP increases transiently after mupirocin in both wild-type and Δ*relA* strains,

we hypothesized that translation inhibition somehow diverts carbon into lipid synthesis. We

added the ribosome inhibitor chloramphenicol and the transcription initiation inhibitor rifampicin

to glycerol cultures of wild-type *E. coli*. Both compounds inhibit translation via mechanisms that

suppress ppGpp synthesis. As with mupirocin treatment of the Δ*relA* strain, chloramphenicol

triggered a rapid decrease in holo-ACP and an increase in long-chain unsaturated acyl-ACP

species C16:1-ACP and C18:1-ACP **(Figure 3C).** Both antibiotics triggered an increase in PL

synthesis intermediates PA and PS that resembled the response of the Δ*relA* strain to mupirocin

(Figure 3D).

What might cause the transient increase in fatty acid synthesis observed after translation

inhibition? Interestingly, both rifampicin and chloramphenicol increased acetyl-CoA

concentrations in glycerol cultures by 3-fold **(Figure 3E)**, suggesting a possible cause. Acetyl-

CoA also increased in both wild-type and Δ*relA* strains after mupirocin treatment in glucose

medium, before decreasing ~30% in the wild-type strain **(Supplemental Figure 4).** While it is

unclear why translation inhibition would increase acetyl-CoA, the observed response of the fatty

acid pathway is consistent with our mathematical model, which predicts fatty acid flux to be

highly sensitive to changes in acetyl-CoA concentrations **(Figure 2B)**. We confirmed the

sensitivity of the fatty acid and PL synthesis pathways to environmental changes using a fast

nutritional upshift. Addition of glucose and amino acids to a glycerol culture also caused rapid

accumulation of PA and PS species that resemble the increases observed following translation

inhibition **(Supplemental Figure 7).**

Moderate to high concentrations of ppGpp regulate PL synthesis via post-translational

control

In order to clearly discern the effects of ppGpp on the fatty acid and PL synthesis

pathways without complications introduced by translation inhibition, we monitored fatty acid and

PL synthesis pathways immediately after inducing RelA*. The responses of the fatty acid and PL

synthesis pathways are again consistent with PlsB inhibition causing long-chain acyl-ACP to

accumulate, which depletes malonyl-ACP by inhibiting ACC **(Figure 4A)**. PL intermediates also

respond in a manner consistent with PlsB inhibition: LPA species steadily decrease, followed by

PA, PS, and PGP **(Figure 4B)**. Addition of chloramphenicol 10 minutes following RelA* induction

causes an increase in unsaturated long-chain acyl-ACP, though ppGpp appears to attenuate the

response of the PL pathway to translation inhibition.

We next sought to evaluate whether ppGpp concentrations far below those occurring

during the stringent response are also capable of regulating PlsB via post-translational control.

As the effects of post-translational control can be distinguished from transcriptional regulation by

response time, we closely followed the dynamics of the PL pathway after mildly inducing ppGpp

synthesis with a low doxycycline concentration. ppGpp increased within 10 minutes of RelA*

induction and reached its elevated steady-state concentration (~150 pmol/OD) by 20 minutes.

Concentrations of PS begin to decrease within 15 minutes **(Figure 4C)**, consistent with post-

translational inhibition of PL synthesis. To verify that the immediate PS depletion is too fast to be

explained by transcriptional regulation, we compared the PS response kinetics with the kinetics

of a ppGpp-driven response known to be mediated by transcriptional control. Cyclopropyl PL are

produced from unsaturated fatty acids of membrane PL by the enzyme Cfa, expression of which

is induced via transcriptional control by ppGpp (28). Cyclopropyl PL begins to accumulate 25

minutes after RelA* induction, well after the PS decrease is established. We conclude that

ppGpp concentrations well below stringent response concentrations inhibit PlsB via a post-

translational mechanism.

Discussion

Over several decades, biochemical and genetic research has extensively characterized the pathways and the enzymes that produce the building blocks of bacterial membranes (4, 29, 30). Many control mechanisms acting on both transcriptional and post-translational levels have been proposed to maintain membrane homeostasis. For instance, each of the enzymes ACC (31), FabH (32), FabZ (33), FabI (34), LpxC (35), and PlsB (36) have been suggested to contribute to membrane synthesis regulation. However, the existence of a control mechanism does not necessarily indicate that it is used to regulate total membrane synthesis flux during steady-state growth. Furthermore, models derived from biochemical and genetic studies can overlook self-regulating mechanisms that occur automatically in a metabolic pathway, such as the coupling of C14:0-OH-ACP concentrations to PL flux. The primary value of our work is that it reveals the regulatory mechanisms that are actually used by growing *E. coli* to regulate membrane biosynthesis. Both our data and our modelling indicate that only two of the many control mechanisms previously identified are sufficient to regulate total PL biosynthesis during steady-state growth. Specifically, we find that cellular demand for membrane synthesis is communicated to PlsB via an undiscovered post-translational mechanism. In turn, PlsB activity regulates fatty acid synthesis by consuming long-chain acyl-ACP species, relieving feedback inhibition of ACC (20) and increasing fatty acid flux. This demand-level flux regulation is consistent with metabolic control theory (37).

Our measurements of enzyme concentrations indicate that PL flux is regulated primarily via post-translational control of enzyme activity, rather than via transcriptional regulation. Concentrations of pathway enzymes with the greatest control over PL flux (ACC and PlsB) are maintained at nearly invariant levels across a 3-fold range of μ. Although it has been long accepted that PlsB activity is adjusted via post-translational control and not transcriptional control (4), the strong flux control of ACC (31) and the growth-regulated transcription of the

accBC, accA, and *accD* genes (38) inspired the suggestion that PL flux might be coupled with μ

via transcriptional regulation. Instead, we find that transcriptional regulation of fatty acid and PL

synthesis genes stabilizes the concentrations of fatty acid and PL synthesis enzymes across μ.

The correlations between concentrations of several enzymes and PL flux that we observe (e.g.

FabF) are unlikely to increase total PL flux, but rather regulate aspects of membrane metabolism

aside from PL flux, such as membrane fluidity. Maintaining a stable membrane synthesis

capacity enables the cell to quickly respond to any change in membrane demand at the cost of

expressing enzymes that remain less active at slow or moderate μ (39).

Our data are also essential for evaluating proposed models of LPS synthesis regulation.

Most importantly, both our model and our experiments indicate that PL flux – and thus, PlsB

activity – controls concentrations of LPS precursor C14:0-OH-ACP. This tightly links fluxes into

the PL and LPS synthesis pathways, as varying PL synthesis by adjusting either PlsB or ACC

activity will naturally vary LPS synthesis in parallel. Surprisingly, we find that flux into the LPS

synthesis pathway is not adjusted by variations in LpxC concentrations, as LpxC concentrations

remain constant despite a 3-fold change in μ. LPS flux may be varied instead by FabZ,

concentrations of which increase by 50% over the 3-fold range of μ sampled **(Supplemental**

Figure 2). Our model predicts this would attenuate the effects of increasing C14:0-OH-ACP

concentrations caused by elevated fatty acid synthesis and divert flux away from LPS synthesis

without affecting PL flux. LpxC concentrations may be stabilized across μ by degradation by the

FtsH protease (40). Degradation of LpxC by FtsH might be used as an emergency brake to

rapidly halt LPS synthesis during growth arrest or growth transitions to prevent accumulation of

toxic LPS intermediates. A comprehensive characterization and modelling (41) of LPS

substrates and enzymes across several steady-state conditions, as we have done for PL

synthesis, will be necessary to determine how LPS synthesis is varied.

Although PlsB determines PL flux during steady-state growth, this control is not absolute,

as fatty acid and PL flux (and likely LPS flux) are also sensitive to acetyl-CoA concentrations and

ACC activity. The sensitivity of membrane synthesis to protein synthesis arrest demonstrates the

responsiveness of membrane synthesis to carbon metabolism. While this sensitivity facilitates

rapid adaptation of PL flux to environmental changes, the tight connection between protein

synthesis, central carbon metabolism, and membrane biogenesis demands additional regulation

(including inhibition by ppGpp) to re-balance the pathway with μ and prevent PL overflow. If

protein synthesis is inhibited in a manner that does not decrease carbon flow into the fatty acid

pathway (e.g. by nitrogen starvation), continued PL and LPS synthesis would outpace synthesis

of the lipoproteins that tether the outer membrane to the peptidoglycan layer. Inhibition of PL and

LPS pathways by ppGpp would thus prevent production of excess membrane, enforcing the

coupling of PL and lipoprotein synthesis. Consistent with this notion, Δ*relA* strains generate

higher quantities of extracellular PL and LPS, likely as outer membrane vesicles (42).

Identifying the specific signal that controls PlsB will reveal how membrane synthesis is

synchronized with growth. But what controls PlsB? Post-translational control of PlsB by

moderately-high concentrations of ppGpp (≥150 pmol/OD) at least partly contributes to steady-

state membrane synthesis regulation, but likely only during nutritional downshifts, stress, or very

slow steady-state growth ($\mu \ll 0.5$ hr^{-1}) when such concentrations are achieved. Whether basal

ppGpp concentrations observed during fast growth ($\mu > 0.5$ hr^{-1}, ppGpp < 60 pmol/OD) also

regulates PL synthesis via PlsB control remains unclear. *In vivo* experiments such as ours

cannot determine the mechanism by which ppGpp inhibits PlsB. Although it has been

demonstrated that high concentrations of ppGpp directly inhibits PlsB (25), multiple groups have

been unable to observe ppGpp inhibition of PlsB *in vitro* when acyl-ACP are used as substrates

(43, 44). ppGpp may therefore inhibit PlsB indirectly via a regulator or a cellular process that

interacts with PlsB. Immunoprecipitation experiments found several proteins that interact with

PlsB including ACP and PssA, as well as several whose roles are unclear (PlsX and YbgC) (45),

suggesting that PlsB forms part of a PL synthesis complex that may couple PlsB activity with μ.

As PL synthesis flux has been observed to oscillate with the cell division cycle in *E. coli* and

other bacteria (46), PlsB may also be regulated by the divisome or by septum formation.

Degradation of PL by phospholipases may also play an important role in membrane

homeostasis during steady-state growth (47), as is known to occur during growth and division in

eukaryotes (48). In addition to identifying the allosteric regulators of PlsB, studies that integrate

connections between PL catabolism and transport with PL synthesis are needed for a

comprehensive understanding of membrane homeostasis.

Acknowledgements

We thank Michael Cashel for providing *E. coli* strain CF7974. We thank Christophe Danelon, Bertus Beaumont, and Frank Bruggeman for invaluable suggestions for improving the manuscript. We thank the National BioResource Project (NIG, Japan) for sharing the LpxC overexpression plasmid. This research was funded by a grant to GB from the Netherlands Organisation for Scientific Research (ALW Open 824.15.018) and an award from the Frontiers of Nanoscience program.

Conflict of Interest

The authors declare that they have no conflict of interest.

Author contributions

MN and GB designed the experiments. MN, FB, NvdB, NS, FY, and GB performed experiments. MN, NvdB, and NI developed analytical methods. MN, FY, and GB analysed the data. GB supervised the research, constructed the mathematical model, and wrote the manuscript.

Materials and Methods

Culture conditions

Cultures were grown in 250-1000 mL Erlenmeyer flasks filled up to 10% of nominal volume with MOPS minimal medium (49) with 9.5 mM NH_4Cl or $^{15}NH_4Cl$ and 0.2% (w/v) carbon source (acetate, succinate, malate, glycerol, glucose, U-^{13}C-glucose (Cambridge Isotope Laboratories) or glucose supplemented with 0.1% Cas-amino acids). Culture flasks were placed in a Grant Instruments Sub Aqua Pro dual water bath at 37 °C and agitated by stirring with a 12

mm magnetic stir bar (VWR), coupled to a magnetic stir plate (2mag MIXdrive 1 Eco and

MIXcontrol 20) set at 1200 rpm. Growth was monitored by optical density measurement at 600

nm using Ultrospec 10 Cell Density Meter (GE Healthcare). Samples for acyl-ACP, lipid analysis

and proteomics were collected using cultures without isotopic labelling. Samples for nucleotide

phosphate measurements were collected from U-^{15}N-labeled cultures and samples for G3P

measurements were collected from U-^{13}C-labeled cultures.

Strains and plasmid pRelA*

All experiments were performed using *Escherichia coli* K-12 strain NCM3722 (CGSC#

12355) and its derivatives. NCM3722 *relA::kan* was constructed by P1 phage transduction using

strain CF7974 (MG1655 Δ*lac* (*rph*+) *relA255*::*kan*) as a donor. Plasmid pRelA* was created by

cloning DNA encoding residues 1-455 from the *E. coli* RelA protein into BglBrick plasmid

pBbS2k (50) (SC101* origin of replication, P$_{Tet}$ promoter, kanamycin resistance). The fluorescent

protein mVenus was fused by restriction-digestion to the C-terminus of RelA via a glycine-serine

linker.

Metabolite sampling

Samples for acyl-ACP, proteomics and lipid analysis were acquired by fast quenching of

1 mL of culture sample into 250 µL of ice-cold 10% trichloroacetic acid. After 10 min incubation

at 0°C cells were pelleted by centrifugation and stored at -80°C until analysis. For nucleotide

phosphates and polar metabolites analysis, samples were acquired by a modified fast vacuum

filtration method (51). 1 mL of culture was collected by vacuum on a pre-wetted 2.5 cm 0.45 µm

HV Durapore membrane filter. After rapid collection the filter was immediately placed upside-

down in quenching solution. For the measurement of nucleotide phosphates, 1 mL of ice-cold 2

M formic acid with 10 µL internal standards mix was used as a quenching solution, which was

subsequently neutralized by 25 µL of 28% ammonium hydroxide. For G3P measurements, 1 mL

of 50:30:20 (v/v/v) mixture of methanol, acetonitrile and water with 0.1% formic acid with 10 µL

internal standard solution (cooled on dry ice) was used as a quenching solution. After 10 min

incubation cells were washed from the filter, transferred to a tube and stored at -80°C until

analysis.

Preparation of internal standards

Isotopically-labeled internal standards (IS) were used to control for sampling and

measurement variation. For acyl-ACP and proteomics assays U-^{15}N *E. coli* whole cell extracts

were prepared using a MOPS minimal medium culture with ^{15}NH$_4$Cl as the sole nitrogen source.

At OD of ~0.5 10% TCA was added 1:4 to the culture to facilitate quenching of metabolism. After

10 min incubation on ice 10 mL single-use IS aliquots were collected by centrifugation and

stored at -80°C until the sample preparation. For the phospholipid measurement U-^{13}C lipid

extract was prepared using a culture grown in minimal MOPS medium with 0.2% U-^{13}C glucose

as the sole carbon source. At OD of ~0.5 10% TCA was added 1:4 to the culture and insoluble

cell material was collected by centrifugation after 10 min incubation on ice. Pellets were

resuspended in mixture consisting of 75 µL MeOH, 10 µL 15 mM citric acid/ 20 mM dipotassium

phosphate buffer and 250 µL of methyl-t-butyl ether per 1 mL of initial culture volume. After

vortexing and 10 min sonication phase separation was induced by addition of 70 µL/1mL of 15

mM citric acid/ 20 mM dipotassium phosphate buffer. After further vortexing, sonication and 10

minutes of incubation at room temperature phases were separated by 10 min centrifugation at

4000 rpm at room temperature. Upper phase was collected to a glass vial and stored at -20°C

until sample preparation.

Instrumentation

All LC/MS runs were performed using Agilent LCMS consisting of binary pump

(G1312B), autosampler (G7167A), temperature-controlled column compartment (G1316A), and

triple quadrupole (QQQ) mass spectrometer (G6460C) equipped with a standard ESI source, all

operated using MassHunter data acquisition software (version 7.0). Mass spectrometer operated

in dynamic MRM mode using transitions generated *in silico* by a script written in Python using

RDkit library using chemical structures of target compound as input. Transitions for targeted

proteomics assays were developed using Skyline (52) based on protein sequences from the

Uniprot database.

LC/MS quantification of acyl-ACP intermediates

Acyl-ACP were measured using a published method (19) with minor modifications. Lysis

buffer was prepared by suspending appropriate number of frozen U-^{15}N-labeled *E. coli* pellets in

10 mL of 50 mM potassium phosphate buffer, pH 7.2, 6 M urea, 10 mM N-ethyl-maleimide, 5

mM EDTA and 1 mM ascorbic acid. 1 mL of lysis buffer was added to each of TCA-quenched

and pelleted cells and proteins were isolated by chloroform/methanol precipitation as described

previously. Protein pellets were resuspended in 10 µL of digestion buffer (4% 2-octyl-glucoside

in 25 mM potassium phosphate buffer, pH 7.2) and after adding 10 µL of 0.1 mg/mL GluC

protease (Promega) incubated overnight at 37°C. After quenching by addition of 5 µL MeOH,

samples were centrifuged and 10 µL was injected in LC/MS system. Separation was performed

on 2.1 mm x 50 mm 1.7 µm CSH C-18 column (Waters) held at 80°C using a binary gradient:

15% B, 3 minute ramp to 25%, 9 min increase to 95% and 1 minute hold at 95% B before 3

minute re-equilibration at starting conditions (A: 25 mM formic acid, B: 50 mM formic acid) at a

flow rate of 0.6 mL/min.

LC/MS quantification of phospholipids

Phospholipids sample preparation procedure is a combination of an MTBE extraction

method (53) and an established LC/MS method (54) Pelleted *E. coli* were resuspended in

mixture containing 150 µL of MeOH, 250 µL of U-^{13}C *E. coli* extract prepared as described

above and 250 µL MTBE. After vigorous vortexing and sonication 125 µL of 15 mM citric acid/

20 mM dipotassium phosphate buffer was added to homogenized pellets. Following further

vortexing and 10 min incubation at room temperature, liquid phases were separated by

centrifugation for 10 min at 20000g. 500 µL the of upper phase was moved to a new tube and

dried in a vacuum centrifuge (Labconco). Dried lipid films were resuspended in 10 µL 65:30:5

(v/v/v) isopropanol/acetonitrile/H_2O, supplemented with 10 mM acetylacetone. After

resuspension, 5 µL H_2O was added to reduce the organic content of the buffer and 5 µL of

resulting mixture was injected into the LC/MS system. Separation was performed on 2.1 mm x

50 mm 1.7 µm CSH C-18 column (Waters) at 60°C with a flow rate of 0.6 mL/min using the

following binary gradient: 25% B, ramp to 56%B in 6 min followed by linear increase to 80% B in

6 min, 2 min hold at 100% B and 3 min re-equilibration (A: 0.05% NH4OH in water, B: 0.05%

NH_4OH in 80% isopropanol 20% ACN).

LC/MS quantification of nucleotides

Frozen cell extracts were defrosted by 1-2 minute incubation in a 37°C water bath and

sonicated for 10 minutes in water ice slurry. After 10 minute centrifugation at 20000g, samples

were loaded on 1 mL/30 mg Oasis Wax cartridge (Waters) preconditioned with 1 mL of MeOH

and 1 mL 50 mM ammonium acetate buffer, pH 4.5. After washing with 1 mL ammonium acetate

buffer, analytes were eluted with 200 µL of 2.8% ammonium hydroxide in MeOH:ACN:H_2O

50:30:20 (v:v:v). After addition of 10 µL of 5% trehalose and brief vortexing samples were dried

down in a vacuum centrifuge (Labconco). Dried trehalose-stabilized extracts were re-dissolved

in 20 µL of MeOH:ACN:H_2O 50:30:20 (v:v:v) and moved to an autosampler vial for analysis.

Separation was performed on 2.1 mm x 100 mm 3.5 µm iHilic column (HILICON) or SeQuant

Zic-cHILIC, 2.1mm x 100 mm, 3µm (Merck) at 0.3 mL/min using the following binary gradient:

100% B, ramp to 85%B in 1.5 min followed by 10 min isocratic hold at 85% B and linear

decrease 30% B in 3 minutes with 2 minute hold at 30% B and 8 minute re-equilibration at initial

conditions. (A: 3.75 mM ammonium acetate, 1.25 mM acetic acid, 2 mM acetylacetone in MQ, B:

11.25 mM ammonium acetate, 3.75 mM acetic acid, 2 mM acetylacetone in 80% ACN). Injection

volume was 2 μL.

LC/MS quantification of G3P

Stored metabolite extracts were dried down in a vacuum centrifuge (Labconco), re-

dissolved in 20 μL of MeOH:ACN:H_2O 50:30:20 (v:v:v) and moved to an autosampler vial for

analysis. Separation was performed on 2.1 mm x 100 mm 3.5 μm iHilic column (HILICON) at 0.3

mL/min using the following binary gradient: 100% B, ramp to 80% B in 10 min followed by linear

decrease to 30% B in 3 min, 2 min hold at 30% B and 8 min re-equilibration. Injection volume

was 2 μL.

LC/MS targeted protein quantification

Relative concentrations of enzymes was measured by targeted proteomics using a

modified version of the acyl-ACP assay. Lysis buffer was prepared by suspending appropriate

number of frozen U-^{15}N-labeled *E. coli* pellets in 10 mL of 50 mM potassium phosphate buffer,

pH 7.2 and 6 M urea. To improve the detection of peptides from PlsB and LpxC, U-^{15}N-labeled

E. coli NCM3722 strains overexpressing PlsB and LpxC were used as internal standards. LpxC

was overexpressed using the corresponding plasmid from the ASKA library (55) (National

BioResource Project (NIG, Japan)). 1 mL of lysis buffer was added to each of TCA-quenched

and pelleted cells and proteins were isolated by chloroform/methanol precipitation as described

previously. Protein pellets were resuspended in 10 μL of digestion buffer (4% 2-octyl-glucoside

in 25 mM Tris buffer, pH 8.1 supplemented with 1 mM $CaCl_2$ and 5 mM TCEP). Alkylation of

cysteine residues was performed by adding 3 μL of 50 mM iodoacetamide followed by 15

minutes of incubation in darkness. Digestion was performed by adding 10 μL of 0.2 mg/mL

Trypsin Gold (Promega) and overnight incubation at 37°C. Samples were centrifuged and 10 μL

was injected in LCMS system. Separation was performed on 2.1 mm x 50 mm 1.7 μm CSH C-18

column (Waters) held at 40°C using a binary gradient: 2% B, 20 minute ramp to 25% B, 4 min

increase to 40% B, 0.5 ramp to 80% and 1 minute hold at 80% B before 3 minute re-equilibration

at starting conditions (A: 25 mL formic acid, B: 50 mM formic acid) at a flow rate of 0.5 mL/min.

Data Analysis

All LC-MS data files were processed in Skyline versions 4.x using target list based on *in*

silico generated transition list. Each target compound had matching isotopically-labeled internal

standard (IS). Processed data were exported as target compounds and IS peak areas and

processed further using a set of Python scripts. Growth rates were obtained from linear fits to

log-transformed growth curves. OD-corrected data were obtained by dividing the signal by OD_{600}

value interpolated from growth curve at the time of sampling. PE-corrected results were

produced by dividing the signal by sum all of signals for all phosphatidyl-ethanolamine species

from the same measurement (in case of phospholipids) or matching sample (in case of other

assays). In nucleotide phosphate and G3P assay absolute concentrations were estimated based

on amounts of internal standards in IS-spike solution assuming RR = 1 implies equimolar

amounts of target compound and IS at the moment of quenching. Correlations in Table 1 were

calculated from log(2) normalized concentrations and PL fluxes using the Descriptive Statistics

function (OriginPro v. 2015).

Mathematical modeling

The computational model was constructed and tested using COPASI version 4.24 (56).

Full details of the model are described in the **Supplemental Methods**.

References

1.   Kamio Y, Nikaido H. 1976. Outer Membrane of Salmonella Typhimurium: Accessibility of

Phospholipid Head Groups to Phospholipase C and Cyanogen Bromide Activated

Dextran in the External Medium. Biochemistry 15:2561–2570.

2. Funahara Y, Nikaido H. 1980. Asymmetric localization of lipopolysaccharides on the outer

membrane of Salmonella typhimurium. J Bacteriol 141:1463–1465.

3. Suzuki H, Nishimura Y, Yasuda S, Nishimura A, Yamada M, Hirota Y. 1978. Murein-

lipoprotein of Escherichia coli: A protein involved in the stabilization of bacterial cell

envelope. MGG Mol Gen Genet 167:1–9.

4. Dowhan W. 2013. A retrospective: use of Escherichia coli as a vehicle to study

phospholipid synthesis and function. Biochim Biophys Acta 1831:471–494.

5. Scott M, Gunderson CW, Mateescu EM, Zhang Z, Hwa T. 2010. Interdependence of cell

growth and gene expression: Origins and consequences. Science (80-).

6. Chubukov V, Uhr M, Le Chat L, Kleijn RJ, Jules M, Link H, Aymerich S, Stelling J, Sauer

U. 2013. Transcriptional regulation is insufficient to explain substrate-induced flux

changes in Bacillus subtilis. Mol Syst Biol 9.

7. Hackett SR, Zanotelli VRT, Xu W, Goya J, Park JO, Perlman DH, Gibney PA, Botstein D,

Storey JD, Rabinowitz JD. 2016. Systems-level analysis of mechanisms regulating yeast

metabolic flux. Science (80-) 354.

8. Raetz CR, Larson TJ, Dowhan W. 1977. Gene cloning for the isolation of enzymes of

membrane lipid synthesis: phosphatidylserine synthase overproduction in Escherichia

coli. Proc Natl Acad Sci U S A.

9. Ohta A, Waggoner K, Louie K, Dowhan W. 1981. Cloning of genes involved in membrane

lipid synthesis. Effects of amplification of phosphatidylserine synthase in Escherichia coli.

J Biol Chem.

10. Ganong BR, Raetz CR. 1982. Massive accumulation of phosphatidic acid in conditionally

lethal CDP-diglyceride synthetase mutants and cytidine auxotrophs of Escherichia coli. J

Biol Chem 257:389–394.

11. Heacock PN, Dowhan W. 1987. Construction of a lethal mutation in the synthesis of the

major acidic phospholipids of Escherichia coli. J Biol Chem 262:13044–13049.

12. 1979. Increased synthesis of phosphatidylserine decarboxylase in a strain of Escherichia
coli bearing a hybrid plasmid. Altered association of enzyme with the membrane. J Biol
Chem 254:627–633.

13. Jackson BJ, Gennity JM, Eugene PK. 1986. Regulation of the Balanced Synthesis of
Membrane Phospholipids. J Biol Chem 261:13464–13468.

14. Dennis PP, Bremer H. 2008. Modulation of Chemical Composition and Other Parameters
of the Cell at Different Exponential Growth Rates. EcoSal Plus 3.

15. Sud IJ, Schaechter M. 1964. Dependence of the content of cell envelopes on the growth
rate of Bacillus megaterium. J Bacteriol 88:1612–1617.

16. Ballesta JPG, Schaechter M. 1972. Dependence of the Rate of Synthesis of
Phosphatidylethanolamine and Phosphatidylglycerol on the Rate of Growth of Escherichia
coli. J Bacteriol 110:452–453.

17. Schulman H, Kennedy EP. 1977. Relation of turnover of membrane phospholipids to
synthesis of membrane derived oligosaccharides of Escherichia coli. J Biol Chem
252:4250–4255.

18. Harris LK, Theriot JA. 2016. Relative rates of surface and volume synthesis set bacterial
cell size. Cell 165:1479–1492.

19. Noga MJ, Cerri M, Imholz N, Tulinski P, Sahin E, Bokinsky G. 2016. Mass-Spectrometry-
Based Quantification of Protein-Bound Fatty Acid Synthesis Intermediates from
Escherichia coli. J Proteome Res 15:3617–3623.

20. Davis MS, Cronan J. 2001. Inhibition of Escherichia coli acetyl coenzyme A carboxylase
by acyl-acyl carrier protein. J Bacteriol.

21. Rhodius VA, Suh WC, Nonaka G, West J, Gross CA. 2006. Conserved and variable
functions of the σEstress response in related genomes. PLoS Biol.

22. Wahl A, My L, Dumoulin R, Sturgis JN, Bouveret E. 2011. Antagonistic regulation of dgkA
and plsB genes of phospholipid synthesis by multiple stress responses in Escherichia coli.
Mol Microbiol 80:1260–1275.

23. Sorensen PG, Lutkenhaus J, Young K, Eveland SS, Anderson MS, Raetz CRH. 1996.

Regulation of UDP-3-O-[R-3-hydroxymyristoyl]-N-acetylglucosamine Deacetylase in Escherichia coli: THE SECOND ENZYMATIC STEP OF LIPID A BIOSYNTHESIS . J Biol Chem 271:25898–25905.

24. Schreiber G, Metzger S, Aizenman E, Roza S, Cashel M, Glaser G. 1991. Overexpression of the relA gene in Escherichia coli. J Biol Chem 266:3760–3767.

25. Heath RJ, Jackowski S, Rock CO. 1994. Guanosine Tetraphosphate Inhibition of Fatty-Acid and Phospholipid-Synthesis in Escherichia coli Is Relieved by Overexpression of Glycerol-3-Phosphate Acyltransferase (PlsB). J Biol Chem 269:26584–26590.

26. Merlie JP, Pizer LI. 1973. Regulation of phospholipid synthesis in Escherichia coli by guanosine tetraphosphate. J Bacteriol 116:355–366.

27. Nunn WD, Cronan Jr. JE. 1976. Regulation of membrane phospholipid synthesis by the relA gene: dependence on ppGpp levels. Biochemistry 15:2546–2550.

28. Eichel J, Chang YY, Riesenberg D, Cronan JE. 1999. Effect of ppGpp on Escherichia coli cyclopropane fatty acid synthesis is mediated through the RpoS sigma factor (σ(S)). J Bacteriol 181:572–576.

29. Raetz CRH. 1978. Enzymology, Genetics, and Regulation of Membrane Phospholipid Synthesis in Escherichia coli. Microbiol Rev 42:614–659.

30. Cronan JE. 2003. Bacterial Membrane Lipids: Where Do We Stand? Annu Rev Microbiol 57:203–224.

31. Davis MS, Solbiati J, Cronan JE. 2000. Overproduction of acetyl-CoA carboxylase activity increases the rate of fatty acid biosynthesis in Escherichia coli. J Biol Chem 275:28593–28598.

32. Heath RJ, Rock CO. 1996. Inhibition of β-ketoacyl-acyl carrier protein synthase III (FabH) by acyl-acyl carrier protein in Escherichia coli. J Biol Chem 271:10996–11000.

33. Zeng D, Zhao J, Chung HS, Guan Z, Raetz CRH, Zhou P. 2013. Mutants resistant to LpxC inhibitors by rebalancing cellular homeostasis. J Biol Chem 288:5475–5486.

34. Heath RJ, Rock CO. 1995. Enoyl-acyl carrier protein reductase (fabI) plays a determinant role in completing cycles of fatty acid elongation in Escherichia coli. J Biol Chem.

35. Schäkermann M, Langklotz S, Narberhaus F. 2013. Ftsh-mediated coordination of lipopolysaccharide biosynthesis in Escherichia coli correlates with the growth rate and the alarmone (p)ppGpp. J Bacteriol 195:1912–1919.

36. Cronan JE. 1978. Molecular Biology of Bacterial Membrane lipids. Annu Rev Biochem 47:163–189.

37. Hofmeyr JHS, Cornish-Bowden A. 2000. Regulating the cellular economy of supply and demand. FEBS Lett.

38. Li SJ, Cronan JE. 1993. Growth rate regulation of Escherichia coli acetyl coenzyme A carboxylase, which catalyzes the first committed step of lipid biosynthesis. J Bacteriol 175:332–340.

39. Davidi D, Milo R. 2017. Lessons on enzyme kinetics from quantitative proteomics. Curr Opin Biotechnol.

40. Ogura T, Inoue K, Tatsuta T, Suzaki T, Karata K, Young K, Su LH, Fierke CA, Jackman JE, Raetz CRH, Coleman J, Tomoyasu T, Matsuzawa H. 1999. Balanced biosynthesis of major membrane components through regulated degradation of the committed enzyme of lipid A biosynthesis by the AAA protease FtsH (HflB) in Escherichia coli. Mol Microbiol 31:833–844.

41. Emiola A, Andrews SS, Heller C, George J. 2016. Crosstalk between the lipopolysaccharide and phospholipid pathways during outer membrane biogenesis in *Escherichia coli*. Proc Natl Acad Sci 113:3108–3113.

42. Crowfoot PD, Esfahani M, Wakil SJ. 1972. Relation Between Protein Synthesis and Phospholipid Synthesis and Turnover in Escherichia coli. J Bacteriol 112:1408–1415.

43. Lueking DR, Goldfine H. 1975. The involvement of guanosine 5-diphosphate-3-diphosphate in the regulation of phospholipid biosynthesis in Escherichia coli. Lack of ppGpp inhibition of acyltransfer from acyl-ACP to sn-glycerol 3-phosphate. J Biol Chem 250:4911–4917.

44. Nunn WD, Kelly DL, Stumfall MY. 1977. Regulation of fatty acid synthesis during the cessation of phospholipid biosynthesis in Escherichia coli. J Bacteriol 132:526–531.

45. Gully D, Bouveret E. 2006. A protein network for phospholipid synthesis uncovered by a

variant of the tandem affinity purification method in Escherichia coli. Proteomics 6:282–
293.

46.   Carty CE, Ingram LO. 1981. Lipid synthesis during the Escherichia coli cell cycle. J
Bacteriol 145:472–478.

47.   May KL, Silhavy TJ. 2018. The Escherichia coli phospholipase PldA regulates outer
membrane homeostasis via lipid signaling. MBio 9.

48.   Hermansson M, Hokynar K, Somerharju P. 2011. Mechanisms of glycerophospholipid
homeostasis in mammalian cells. Prog Lipid Res.

49.   Neidhardt FC, Bloch PL, Smith DF. 1974. Culture Medium for Enterobacteria. J Bacteriol
119:736–747.

50.   Lee T, Krupa RA, Zhang F, Hajimorad M, Holtz WJ, Prasad N, Lee S, Keasling JD,
Khosla C, Keasling J, Lee S, Chou H, Ham T, Lee T, Keasling J, Atsumi S, Liao J, Raab
R, Tyo K, Stephanopoulos G, Pieper D, Reineke W, Martin V, Pitera D, Withers S,
Newman J, Keasling J, Pitera D, Paddon C, Newman J, Keasling J, Alper H, Fischer C,
Nevoigt E, Stephanopoulos G, Ajikumar P, Xiao W, Tyo K, Wang Y, Simeon F, Leonard
E, Mucha O, Phon T, Pfeifer B, Stephanopoulos G, Dejong J, Liu Y, Bollon A, Long R,
Jennewein S, Williams D, Croteau R, Yan Y, Kohli A, Koffas M, Li M, Elledge S, Shao Z,
Zhao H, Zhao H, Gibson D, Young L, Chuang R, Venter J, Hutchison C, Smith H,
Keasling J, Lee S, Kim H, Park J, Park J, Kim T, Martin C, Nielsen D, Solomon K, Prather
K, Canton B, Labno A, Endy D, Knight T, Anderson J, Dueber J, Leguia M, Wu G, Goler
J, Arkin A, Keasling J, Shetty R, Endy D, Knight T, Tao L, Jackson R, Cheng Q, Anthony
J, Anthony L, Nowroozi F, Kwon G, Newman J, Keasling J, Lee S, Newman J, Keasling J,
Lee S, Keasling J, Lee S, Keasling J, Guzman L, Belin D, Carson M, Beckwith J,
Hajimorad M, Gray P, Keasling J, Lee S, Chou H, Pfleger B, Newman J, Yoshikuni Y,
Keasling J, Gorke B, Stulke J, Skulj M, Okrslar V, Jalen S, Jevsevar S, Slanc P, Strukelj
B, Menart V, Carapuca E, Azzoni A, Prazeres D, Monteiro G, Mergulhao F, Lee C, Ow D,
Oh S, Lee C, Kim J, Shin S, Hwang S, Tao L, Jackson R, Rouviere P, Cheng Q, Lee C,
Lee S, Shin S, Hwang S, Friehs K, Antoine R, Locht C, Lutz R, Bujard H, Manen D, Xia G,
Caro L, Grabherr R, Bayer K, Glass R, Kues U, Stahl U, Redding-Johanson A, Batth T,
Chan R, Krupa R, Szmidt H, Adams P, Keasling J, Lee T, Mukhopadhyay A, Petzold C,
Dunlop M, Dossani Z, Szmidt H, Chu H, Lee T, Keasling J, Hadi M, Mukhopadhyay A,

Leguia M, Brophy J, Densmore D, Anderson J, Horton R, Cai Z, Ho S, Pease L,
Pushnova E, Geier M, Zhu Y, Mason G, Provero P, Vaira A, Accotto G. 2011. BglBrick
vectors and datasheets: A synthetic biology platform for gene expression. J Biol Eng 5:12.

51.   Link H, Kochanowski K, Sauer U. 2013. Systematic identification of allosteric protein-
metabolite interactions that control enzyme activity in vivo. Nat Biotechnol 31:357–361.

52.   MacLean B, Tomazela DM, Shulman N, Chambers M, Finney GL, Frewen B, Kern R,
Tabb DL, Liebler DC, MacCoss MJ. 2010. Skyline: An open source document editor for
creating and analyzing targeted proteomics experiments. Bioinformatics 26:966–968.

53.   Matyash V, Liebisch G, Kurzchalia T V, Shevchenko A, Schwudke D. 2008. Lipid
extraction by methyl-tert-butyl ether for high-throughput lipidomics. J Lipid Res 49:1137–
1146.

54. Susanto O, Koh YWH, Morrice N, Tumanov S, Thomason PA, Nielson M, Tweedy L,
Muinonen-Martin AJ, Kamphorst JJ, Mackay GM, Insall RH. 2017. LPP3 mediates self-
generation of chemotactic LPA gradients by melanoma cells. J Cell Sci 130:3455–3466.

55. Kitagawa M, Ara T, Arifuzzaman M, Ioka-Nakamichi T, Inamoto E, Toyonaga H, Mori H.
2005. Complete set of ORF clones of Escherichia coli ASKA library (A complete set of E.
coli K-12 ORF archive): unique resources for biological research. DNA Res 12:291–299.

56. Hoops S, Gauges R, Lee C, Pahle J, Simus N, Singhal M, Xu L, Mendes P, Kummer U.
2006. COPASI - A COmplex PAthway SImulator. Bioinformatics 22:3067–3074.

Figure and Tables

Figure 1. Characterization of the *E. coli* fatty acid and PL synthesis pathways during steady-

state growth. **A.** The fatty acid and PL synthesis pathways. Long-chain fatty acids are

transferred from C16:0-ACP, C18:0-ACP, or C18:1-ACP (highlighted in blue) to *sn*-glycerol-3-

phosphate (G3P) by PlsB to yield lysophosphatidic acid (LPA), which is acylated with C16:1-

ACP or C18:1-ACP to generate phosphatidic acid (PA). PA is converted to (d)CDP-

diacylglycerol ((d)CDP-DAG) with either CTP or dCTP. At the branch point in PL synthesis,

(d)CDP-DAG is converted to phosphatidylglycerol phosphate (PGP), which is dephosphorylated

to yield PG (~25% of total PL). (d)CDP-DAG is also a precursor of phosphatidylserine (PS),

which is decarboxylated to yield PE (~70% of total PL). PG is further converted to CL (~5% of

total PL). C14:0-OH-ACP (highlighted in red) is a precursor for the synthesis of

lipopolysaccharide (LPS), the major component of the outer leaflet of the outer membrane. **B, C.**

Growth rate (μ)-dependent abundance of PG, PE, and CL per OD (**B**) and μ-dependence of PL

flux (**C**), normalized to average values for each strain. **D.** Steady-state concentrations of acetyl-

CoA and ppGpp. **E, F, G.** Correlations of G3P and acyl-ACP concentrations **(E)**, PL intermediate

concentrations **(F)**, and PlsB and LpxC concentrations **(G)** with PL flux. Concentrations of

soluble substrates and enzymes are calculated using cell volume (proportional to OD), while PL

intermediate concentrations are calculated using membrane volume, which is proportional to

total PE. All concentrations and fluxes are normalized and log(2)-transformed for comparison. All

data points and error bars represent the average and standard deviation, respectively, of

measurements of three samples from one culture. For WT PL and PL intermediates, three

independent biological replicates of each condition are depicted; for ACP species and PlsB, two

independent biological replicates; for nucleotides, LpxC, and G3P, data from one experiment per

condition is shown. For pRelA, one measurement per inducer concentration is shown.

Figure 2. Simulated PL synthesis and experimental data identifies PlsB as site of PL synthesis

control during steady-state growth. **A.** Reactions simulated in the model. For simplicity, only

reactions late in the saturated fatty acid pathway are included, and branching of the PL pathway

into PE and PG is not included. Each reaction is modelled as an irreversible one- or two-

substrate Michaelis-Menten reaction. ACC is competitively inhibited by C16:0-ACP and C18:0-

ACP. Reactions catalysed by FabI/FabZ and LpxA/LpxC are considered in the model as single

reactions represented in the diagram by "FabZ" and "LpxC", respectively. **B.** Response of PE

and LPS fluxes to varying V_{max} of pathway enzymes and acetyl-CoA concentrations. 4-fold

variations in V_{max} of all other reactions tested did not change PE or LPS flux (not shown). **C.**

Simulated changes in metabolite concentrations in response to variations in PlsB and ACC V_{max}

and acetyl-CoA (line plots) overlaid on experimentally-measured concentrations (scatter plots,

data from **Figure 1**). Line plots are offset to prevent overlap. Differential equations and

parameters of the simulation are provided in the **Supplemental Methods**.

Figure 3. Responses of the fatty acid and PL synthesis pathways to translation inhibition. **A, B.**
Responses of acyl-ACP **(A)** and PL intermediates **(B)** to mupirocin. Mupirocin was added at 0
min (indicated by dashed lines) to glucose cultures of *E. coli* wild-type and Δ*relA* NCM3722.
Each point represents one measurement from a time series collected from an independent
culture. Two biological replicate series for each strain are depicted. **C, D.** Addition of translation
inhibitor chloramphenicol or the transcription inhibitor rifampicin to glycerol cultures of wild-type
E. coli (indicated by dashed lines at 0 min) causes an influx of carbon into the fatty acid pathway,
as suggested by a decrease in holo-ACP and an increase in unsaturated long-chain acyl-ACP
species **(C).** The pulse of carbon continues into the PL synthesis pathway, as indicated by
transient increases in total PA and PS **(D).** Each trajectory in **C-D** is obtained from a single
culture. **E.** Chloramphenicol and rifampicin both trigger a rapid decrease in ppGpp and
accumulation of ACC and FabH substrate acetyl-CoA. Points and error bars in **E** are averages
and standard deviation of triplicate samples, respectively, from individual experiments.

Figure 4. PlsB activity is suppressed by moderate to high concentrations of ppGpp via post-translational inhibition. **A, B.** Response of acyl-ACP **(A)** and PL intermediate pools **(B)** to maximal overexpression of ppGpp synthesis enzyme RelA*. RelA* expression was induced by addition of 40 ng/mL doxycycline at -10 minutes (dashed blue line), followed 10 minutes later by chloramphenicol addition (dashed grey line at 0 min). **C.** Response of concentrations of PL intermediates and cyclopropyl-PE to mild RelA* overexpression, triggered by addition of 1 ng/mL doxycycline (dashed grey line at 0 min). Each time series represents one independent biological replicate.

Substrate	N	Pearson correlation coefficient (r)	Significance (p, two-tailed)
malonyl-ACP	12	0.80	<0.01
G3P*	5	0.99	<0.01
C14:0-OH-ACP	12	0.94	$<10^{-4}$
C14:0-ACP	12	-0.16	0.6
C16:0-ACP	12	-0.42	0.2
C18:0-ACP	12	-0.76	<0.01
C16:1-ACP	12	0.42	0.2
C18:1-ACP	12	-0.38	0.2
LPA	18	0.68	<0.01
PA	18	0.95	$<10^{-4}$
(d)CDP-DAG	18	0.89	$<10^{-4}$
PS	18	0.91	$<10^{-4}$
PGP**	15	0.61	0.015

1 *G3P: calculated with measurement in glycerol medium excluded.

2 **PGP: calculated with measurements in glucose + cas-amino acids excluded.

Enzyme	N	Pearson correlation coefficient (r)	Significance (p, two-tailed)
LpxC	6	-0.4	0.4
PlsB	12	-0.4	0.2
PlsC	12	0.63	0.03
CdsA	12	0.90	$<10^{-4}$
PssA	12	-0.4	0.2
Psd	12	0.2	0.6
PgsA	12	-0.7	0.01
PgpA	12	0.9	$<10^{-4}$
AccA	12	0.70	0.01
AccB	12	0.38	0.2
AccC	12	0.50	0.1
AccD	12	0.21	0.5
AcpP	12	0.04	0.9
FabA	12	0.29	0.4
FabB	12	0.28	0.4
FabD	12	0.35	0.3
FabF	12	0.76	<0.01
FabG	12	0.68	0.02
FabH	6	0.49	0.3
FabI	12	0.82	<0.001
FabZ	12	0.77	<0.01
GpsA	12	-0.69	0.01

1   **Table 1.** Pearson correlation coefficients and significance determined for PL flux and species

2   abundance during steady-state growth in wild-type *E. coli*. *N* indicates the number of

3   independent measurements used to calculate the correlation coefficients. Shaded cells indicate

4   *p* < 0.05.