The Gene Expression Deconvolution Interactive Tool (GEDIT):

Accurate Cell Type Quantification from Gene Expression Data

Brian B. Nadel[1], David Lopez[1], Dennis J. Montoya[1], Hannah Waddel[3], Misha M. Khan[4], Serghei Mangul[5,6], Matteo Pellegrini[1,2]

[1]Bioinformatics Interdepartmental Degree Program, Molecular Biology Institute, Department of Molecular Cellular and Developmental Biology, and Institute for Genomics and Proteomics, University of California Los Angeles, Los Angeles, CA

[2]Department of Dermatology, David Geffen School of Medicine, University of California Los Angeles, Los Angeles, CA

[3]Department of Mathematics, University of Utah, Salt Lake City, UT

[4]Departments of Biology and Computer Science, Swarthmore College, Swarthmore, PA

[5] Department of Clinical Pharmacy, USC School of Pharmacy

[6] The Quantitative and Computational Biology, USC Dornsife College of Letters, Arts and Sciences, University of Southern California

Abstract

The cell type composition of heterogeneous tissue samples can be a critical variable in both clinical and laboratory settings. However, current experimental methods of cell type quantification (e.g. cell flow cytometry) are costly, time consuming, and can introduce bias. Computational approaches that infer cell type abundance from expression data offer an alternate solution. While these methods have gained popularity, most are limited to predicting hematopoietic cell types and do not produce accurate predictions for stromal cell types. Many of these methods are also limited to particular platforms, whether RNA-seq or specific microarrays. We present the Gene Expression Deconvolution Interactive Tool (GEDIT), a tool that overcomes these limitations, compares favorably with existing

methods, and provides superior versatility. Using both simulated and experimental data, we

extensively evaluate the performance of GEDIT and demonstrate that it returns robust results under a

wide variety of conditions. These conditions include a variety of platforms (microarray and RNA-seq),

tissue types (blood and stromal), and species (human and mouse). Finally, we provide reference data

from eight sources spanning a wide variety of stromal and hematopoietic types in both human and

mouse. This reference database allows the user to obtain estimates for a wide variety of tissue

samples without having to provide their own data. GEDIT also accepts user submitted reference data,

thus allowing the estimation of any cell type or subtype, provided that reference data is available.

Author Summary

The Gene Expression Deconvolution Interactive Tool (GEDIT) is a robust and accurate tool

that uses gene expression data to estimate cell type abundances. Extensive testing on a variety of

tissue types and technological platforms demonstrates that GEDIT provides greater versatility than

other cell type deconvolution tools. GEDIT utilizes reference data describing the expression profile of

purified cell types, and we provide in the software package a library of reference matrices from various

sources. GEDIT is also flexible and allows the user to supply custom reference matrices. A GUI

interface for GEDIT is available at http://webtools.mcdb.ucla.edu/, and source code and reference

matrices are available at https://github.com/purebrawn/GEDIT.

Introduction

Cell type composition is an important variable in biological and medical research. In laboratory

experiments, cell sample heterogeneity can act as a confounding variable. Observed changes in gene

expression may result from changes in the abundance of underlying cell populations, rather than

changes in expression of any particular cell type [1]. In clinical applications, the cell type composition

of tissue biopsies can inform treatment. For example, in cancer, the number and type of infiltrating

immune cells has been shown to correlate highly with prognosis ([2], [3], [4]). Moreover, patients with

a large number of infiltrating T cells are more likely to respond positively to immunotherapy [5].

For many years, cell flow cytometry via FACS sorting has been the standard method of cell

type quantification. More recently, single cell RNA-seq methods such as 10x Chromium, Drop-Seq,

and Seq-Well have become available [6],[7]. However, both approaches suffer from significant

limitations. FACS sorting is cumbersome and expensive, and some sample types require hours of

highly skilled labor to generate data. Similarly, single cell RNA-seq methods remain expensive for

large sample studies. Additionally, cell types such as neurons, myocytes, and adipocytes are difficult

for these technologies to capture due to cell size and morphology.

Both FACS sorting and single cell methods have the potential to introduce bias, as these

technologies require that tissue samples be dissociated into single cell suspensions. Many stromal cell

types are tightly connected to one another in extracellular matrices. The procedures necessary to

create single cell suspensions can damage some cells, while others remain in larger clusters that are

not captured or sequenced. Consequently, subtle differences in sample preparation can produce

dramatically different results [8]. While FACS sorting and single cell methods can produce pure

samples of each cell type, the observed cell counts may not accurately represent the cell type

abundances in the original sample. Tools like Cell Population Mapping and MuSiC utilize single cell

reference data to perform bulk deconvolution, but requires that single cell data be available for all the

cell types of interest, which is not always the case [9,10].

During the past few years, digital means of cell type quantification, often referred to as cell type

deconvolution or decomposition, have become a popular complement to FACS sorting and single cell

approaches. However, these methods produce approximate results that are often limited to use on

particular cell types or platforms. For example, ImmuneQuant can estimate cell type fractions for

immune cells only [11]. xCell can produce estimates for the 64 cell types supported by the tool, but it

does not allow the inclusion of additional cell types or subtypes [12]. CIBERSORT is specifically

designed for data generated from microarrays, and provides reference data only for hematopoietic cell

types [13].

To overcome some of the limitations of existing cell abundance estimation tools, we present

the Gene Expression Deconvolution Interactive Tool (GEDIT). GEDIT utilizes gene expression data to

accurately predict cell type composition of tissue samples. We have assembled a library of reference

data from 11 distinct sources and use these data to generate thousands of synthetic mixtures. In order

to produce optimal results, these synthetic mixtures are used to test and refine the approaches and

parameters used by GEDIT. We compare the performance of GEDIT relative to other tools using three

sets of mixtures containing known cell type proportions: 12 *in vitro* mixtures of immune cells

sequenced on microarrays, six RNA-seq samples collected from ovarian cancer ascites, and eight

RNA-seq samples collected from blood. We also use GEDIT to deconvolute two sets of human tissue

samples: 21 skin samples from patients with skin diseases, and 17,382 samples of varied tissues from

the GTEx database. Lastly, we apply GEDIT to the Mouse Body Atlas, a collection of samples

collected from various mouse tissues and cell types. We find that GEDIT compares favorably to other

cell type deconvolution tools and is effective across a broad range of datasets and conditions.

Results

Reference Data

Reference data profiling the expression of purified cell types is a requirement for reference-

based deconvolution. Methods that do not directly require reference data, such as non-negative matrix

factorization, still require knowledge of expression profiles or marker genes in order to infer the identity

of the predicted components. For this study, we have assembled or downloaded a set of 11 reference

matrices, each containing the expression profiles of eight to 29 cell types (Table 1). These data

sources span multiple platforms, including bulk RNA-seq, microarray, and single-cell RNA-seq.

Complete details on the sources and assembly of these matrices are described in the methods [13–

23].

Matrix	Species	Reference	Platform	# of Cell Types	Cell Types
Human Skin Signatures	Human	(Swindell et al. 2013)	Multi-Microarray	21	Immune
Human Body Atlas	Human	(Su et al. 2004)	Affymetrix U133A/GNF1H	13	Immune
Human Primary Cell Atlas	Human	(Mabbott et al. 2013)	Affymetrix U133 Plus 2.0	26	Immune and Stromal
BLUEPRINT*	Human	(Martens and Stunnenberg 2013)	Bulk RNA-Seq	8	Immune
ENCODE*	Human	(ENCODE Project Consortium 2004)	Bulk RNA-Seq	29	Mostly Stromal
LM22	Human	(Newman et al. 2015)	Affymetrix Microarray	22	Immune
10x Single Cell Dataset*	Human	(Zheng et al. 2017)	Single Cell RNA-Seq	9	Immune
ImmunoStates	Human	(Vallania et. al., 2018)	Multi-Microarray	20	Immune
Tabula Muris	Mouse	(The Tabula Muris Consortium, 2018)	Single Cell RNA-seq	12	Immune and Stromal
Mouse Body Atlas	Mouse	(Lattin et al, 2008)	Affymetrix Mouse Genome 430 2.0 Array	20	Immune and Stromal
ImmGen	Mouse	(Heng et al, 2008)	Affymetrix Gene 1.0 ST	137	Immune with many subtypes

109   Table 1. Library of Reference Data. Asterisk denotes matrices assembled from source data as part of
110   this project. All matrices are compatible with GEDIT and available on the GitHub repository
111   (https://github.com/BNadel/GEDIT).
112

GEDIT Algorithm

GEDIT requires as input two matrices of expression values. The first is expression data is

collected from the mixtures that will be deconvoluted; each column represents one mixture, and each

row corresponds to a gene. The second matrix contains reference data, with each column

representing a purified reference profile and each row corresponding to a gene. In a multi-step

process, GEDIT utilizes the reference profiles to predict the cell type proportions of each submitted

mixture (Figure 1).

Figure 1. The GEDIT pipeline. The input matrices are quantile normalized then reduced to matrices
containing only signature genes. Next, a row-scaling step serves to control for the dominating effect of
highly expressed genes. Lastly, linear regression is performed, and predictions of cell type
abundances are reported to the user.

Input	Description	Allowed Values	Default Value
RefMat	Matrix of purified cell types	N by M matrix; N is number of genes, M is number of cell types	NA
MixMat	Matrix of mixtures to be deconvoluted	N by P matrix; N is number of genes, P is number of mixtures	NA
SigMeth	Method of signature gene selection	Entropy, MeanRat, MeanDiff, ZScore, fsRat, fsDiff	Entropy
NumSigs	Average number of signature genes per cell type	[1, 10,000]	50
MinSigs	Minimum number of signatures per cell type	[1,NumSigs]	=NumSigs
RowScale	Extent of per-row normalization	[0.0,1.0]	0

125
126

Table 2. GEDIT inputs include two matrices and four parameter settings. RefMat is an expression
matrix documenting the expression profiles of each cell type to be estimated. MixMat is an expression
matrix documenting expression values for each sample to be deconvoluted. SigMeth determines the
method by which signature genes are selected. NumSigs determines the total number of signature
genes, whereas MinSigs sets the minimum number of signature genes for each cell type. RowScale
refers to the extent to which expression vectors are transformed to lessen the dominating effect of
highly expressed genes, with a value of 0.0 representing the most extreme transformation. Default
values were determined by evaluating performance on a set of synthetic mixtures (Figure 3).
135

Synthetic Mixture Generation and Parameter Testing

We generated a large number of synthetic mixtures *in silico* to test the efficacy of GEDIT and

to assess how accuracy varies as a function of four parameter choices (SigMeth, NumSigs, MinSigs,

RowScale, described in Table 2). We produced a total of 10,000 simulated mixtures of known

proportions using data from four reference matrices: BLUEPRINT, The Human Primary Cell Atlas, 10x

Single Cell, and Skin Signatures. We then ran GEDIT on these simulated mixtures and evaluated its

performance while varying four parameter settings (Figure 2) and other design choices. Based on

these results, we selected default values for each parameter (SigMeth = Entropy, NumSigs = 50,

MinSigs = 50, RowScale = 0.0). Full details on the generation of these simulations are described in the

supplementary materials.

Figure 2. Effect of GEDIT parameter choices on accuracy of predictions in simulated experiments.
10,000 simulated mixtures were generated, each using one of four reference matrices, with either four,
five, six, or ten cell types being simulated. Deconvolution was performed using a separate expression
matrix than the one used to generate the mixtures. When not otherwise noted, we use the following
parameters: signature selection method = entropy; number of signatures = 50; row scaling = 0.0; and

number of fixed genes = number of signatures.
153

Preprocessing and Quantile Normalization

The first step in the GEDIT pipeline is to render the two matrices comparable. This is done by

first excluding all genes that are not shared between the two matrices. Genes that have no detected

expression in any reference cell type are also excluded, as they contain no useful information for

deconvolution. Both matrices are then quantile normalized, such that each column follows the same

distribution as every other; this target distribution is the starting distribution of the entire reference

matrix.

Signature Gene Selection

GEDIT next identifies signature genes. Gene expression experiments can simultaneously

measure tens of thousands of genes, but many of these genes are uninformative for deconvolution.

Specifically, genes with similar expression levels across all cell types are of little use, as observed

expression values in the mixtures offer no insight into cell frequencies. Genes that are highly

expressed in a subset of cell types are more informative, and we refer to these as signature genes. By

using only signature genes, rather than the entire expression matrix, the problem of deconvolution

becomes more tractable and less computationally intensive. Moreover, identification of signature

genes can be valuable to researchers for other applications (e.g. cell type assignment for scRNA-seq

data).

In order to identify the best signature genes in a given reference matrix, GEDIT calculates a

signature score for each gene. By default, this score is computed using the concept of information

entropy. Information entropy quantifies the amount of information in a probability distribution, with

highly uniform distributions having the highest entropy. The expression vector for each gene (i.e. the

set of expression values across all cell types in the reference) is divided by its sum, such that the

entries can be interpreted as probabilities. Information entropy is then calculated according to its

mathematical definition (see Methods), and genes with the lowest entropy are selected as signature

genes. Entropy is minimized when expression is detected only in a single cell type and maximized

when expression values are equal across all cell types. Thus, by selecting genes with low entropy, we

favor genes that are expressed in a cell type specific manner. By default, 50 signature genes are

selected for each cell type in the reference matrix. We chose 50 signature genes, and entropy as our

scoring method, because it returned optimal results when run on 10,000 synthetic mixtures (see

Figure 2).

We also evaluated the effect of accepting more signature genes for some cell types than

others, depending on how many genes have low entropy. In this scheme, on average 50 signature

genes are used per cell type. However, a fourth parameter is used, which specifies the minimum

number of signature genes per cell type. After these have been selected, remaining signature genes

are added based only on lowest entropy, regardless of cell type of maximal expression. We found that

this parameter had minimal effect on accuracy, when applied to synthetic mixtures (Figure 2c).

Therefore, this option is not used by default, though it can be specified by the user.

Row Scaling

One complication in the application of linear regression to gene expression data is the

drastically different scale at which some genes are expressed. For example, CD14 and THEMIS

(Figure 3) have both been identified as signature genes: CD14 for monocytes and THEMIS for CD4+

T cells. However, CD14 is expressed at much higher levels in most cell types and will have a larger

impact on the estimation of cell type composition, relative to THEMIS. In other words, the possible

penalty resulting from a poor fit of CD14 is much larger than the penalty from a poor fit of THEMIS.

Figure 3. The "row scaling" transformation, as implemented by GEDIT. CD14 and THEMIS are two
examples of signature genes with drastically different magnitudes of expression. CD14 is a signature
gene for monocytes, and THEMIS for CD4+ T cells. The original expression vectors are transformed,
such that all values fall between 0.0 and 1.0, equalizing the effect of genes with varying magnitudes of
expression.
206

In order to equalize the effect of each signature gene on the linear regression, we implement a

transformation that we term row scaling. Specifically, the range of all observed values for a particular

gene (including reference cell types and samples) is adjusted such that the maximum value is 1.0 and

the minimum value is 0.0. As a result, all genes have a comparable influence on the calculation of the

linear regression solution, regardless of overall magnitude of expression. This transformation can be

modulated by adjusting the row scaling parameter. By default, the value of this parameter is 0.0, and

the transformation is applied as described above. Values between 0.0 and 1.0 are also allowed, which

reduces the extent of the transformation (see Methods for details).

Linear Regression:

Non-negative linear regression was performed using the glmnet package in R. The glmnet

function is used with lower.limits=0, alpha=0, lambda=0, intercept=FALSE. These settings perform a

linear regression where all coefficients are non-negative, and with no regularization and no intercept

term.

GEDIT Compares Favorably with Other Tools

In order to assess the performance of GEDIT relative, we perform an experiment comparing

GEDIT to 4 other deconvolution tools (CIBERSORT, DeconRNASeq, dtangle and xCell;

[12,13,24,25]). The authors have tried to carry out the comparison in an unbiased manner, but it must

be noted that this has been carried out in parallel with the development of the tool. Non-deconvolution

tools like MCP-counter, SAVANT, and the DCQ algorithm are excluded from this study because they

do not predict cell type fractions [26–28].Tools that require single cell data, such as MuSiC and CPM,

are also excluded, as this study is limited to tools that operate on bulk expression data [9,10]). See

Table 3 for a summary of current bulk deconvolution methods.

| Tool | Publication | Custom Reference | Approach | Output | Reference Data Provided with Tool | | |
					Number of Datasets	Cell Types	Species
GEDIT	Nadel et. al., 2020	Yes	Deconvolution	Predicted Fractions	11	Immune and Stromal	Human, Mouse
Cibersort	Newman et. al., 2015	Yes, if marker genes specified	Deconvolution	Predicted Fractions	1	Immune	Human
xCell	Aran et. al., 2017	No	Marker Genes	Predicted Fractions	5	Immune and Stromal	Human

dtangle	Hunt et. al., 2018	Yes, if marker genes specified	Deconvolution	Predicted Fractions	0	N/A	N/A
DeconRNASeq	Gong et. al., 2013	Yes	Deconvolution	Predicted Fractions	0	N/A	N/A
DCQ/ImmQuant	Altboum et. al., 2014; Frishberg et. al., 2016	Yes	Deconvolution	Scores	3	Immune	Human, Mouse
Cibersort (absolute mode)	Newman et. al., 2015	Yes, if marker genes specified	Deconvolution	Scores	1	Immune	Human
SaVant	Lopez et. al., 2017	Yes, if marker genes specified	Marker Genes	Scores	12	Immune and Stromal	Human, Mouse
MCP-Counter	Becht et. al., 2016	No	Marker Genes	Scores	N/A	Immune and Stromal	Human

Table 3. High level characteristics of current cell type estimation tools. Some tools accept custom
references, which allows the tool to estimate the abundance of cell types not present in the default
reference. Tools listed here take one of two approaches: they either perform deconvolution (most
commonly regression) or calculate a score based on intensity of marker gene expression. Depending
on the tool, the output can be interpreted as fractions corresponding to the abundance of each cell
type, or as scores for each cell type that cannot necessarily be compared in an inter-cellular manner.
238

To perform this study, we utilize three datasets for which cell type fractions have been

estimated using orthogonal methods. Two of these datasets were used in a recent benchmarking

study [29]. Both are profiled using RNA-seq, and represent samples collected either from human

cancer ascites or human blood [30,31]. In both cases, cell type fractions have been evaluated by

FACS sorting. The final dataset was prepared *in vitro* and consists of six cell types that were

physically mixed together (in known proportions) to prepare 12 mixtures. These mixtures were then

profiled using an Illumina HT12 BeadChip microarray. Adding to the previous benchmarking study, we

also explore the effect of using four separate reference datasets: The Human Primary Cell Atlas,

LM22, ImmunoStates, and a reference constructed from BLUEPRINT data. For each dataset, all tools

(except xCell) were run four times, each time using a different reference matrix.

Compared to the other tools, GEDIT produces the most robust and consistently accurate

results (Figure 4). For many tools, the quality of predictions varies greatly depending on the cell type,

dataset, or choice of reference matrix. When results are averaged across the four possible reference

choices, GEDIT produces the minimum error and maximum correlation for all three datasets. This

result suggests that GEDIT is the best choice when researchers are using novel references matrices

that have not been curated or tested.

Figure 4. Performance of five deconvolution tools when run on a set of 26 samples from three
sources. Error and correlation between actual and predicted cell type fractions; calculated for each of
the five cell types represented, for each of the three data sources analyzed, and for all predictions
regardless of cell type or data source. Underlying cell type fractions are evaluated via FACS (ascites
and blood) or by controlled mixing of purified cell types (CellMix).
262

The optimal choice of reference matrix varies greatly depending on the exact combination of

tool, dataset, and cell type. While using LM22 often produces the most accurate results, there are

many exceptions. For instance, DeconRNASeq and GEDIT produce their best results for the blood

dataset when using the BLUEPRINT reference. For the ascites data, several tools prefer

ImmunoStates as the optimal reference choice. The best choice of reference is highly dependent on

the nature of the input data and on the tool being used. In practice, researchers may wish to perform

deconvolution multiple times--in each case using a separate reference matrix--and compare results for

consistency.

Skin Expression Data

We further validate GEDIT by using it to deconvolute a set of skin biopsies from humans with a

variety of skin diseases [13]. The exact cell type composition of these samples is unknown, but we

have reasonable expectations based on skin and disease biology. For example, macrophages are

known to be abundant in granulomas of leprosy legions, and Steven-Johnson Syndrome produces

blisters that fill with large numbers of monocytes [32,33]. We find that, in all cases, predictions made

by GEDIT conform well with these biological expectations. Keratinocytes are highly predicted in most

cases, as one would expect with skin samples (Figure 5). Deviations from this pattern correspond with

disease biology. Monocytes are highly predicted in Stevens-Johnson syndrome, as are macrophages

in the three leprosy samples, and T cells in the Mycosis Fungoides (T cell lymphoma) sample.

283
284 Figure 5. GEDIT predictions for 21 samples of various skin diseases. GEDIT correctly identifies
285 keratinocytes and subcutaneous adipose as the most common cell. Deviations from this pattern
286 correspond to disease biology. SJS represents blister fluid from Steven Johnson Syndrome, and is

287 predominantly immune cells. LL and RR represent two forms of leprosy, which result in large numbers
288 of macrophages. MF is a T Cell Lymphoma.
289

Application of GEDIT to Mouse Data

Unlike tools specifically designed for human data, GEDIT can be used to decompose data from

any organism for which reference data is available. Here, we demonstrate the efficacy of GEDIT when

applied to the Mouse Body Atlas, a collection of tissue and cell type samples collected from mice [22].

As reference data, we assembled a matrix of 12 cell types using single cell data from the Tabula Muris

[20]. GEDIT correctly infers the identity of purified cell types, including six samples that consist of

either pure NK cells, B cells, T cells, or granulocytes. An entry for macrophages is not available in the

reference used, but most macrophage samples are identified as monocytes, which is the most similar

cell type present in the reference matrix. For more complex tissues, GEDIT predicts cell type fractions

that correspond to the biology of the samples. Hepatocytes are predicted to be highly prevalent in the

liver sample (84%) and are not predicted in any other sample (less than 5% in all cases). Similar

patterns hold for keratinocytes in the epidermis, epithelial cells in two intestinal samples and cardiac

muscle cells in heart and muscle samples.

Figure 6. GEDIT predictions on 30 samples collected from various mouse tissues and cell types

(mouse body atlas [22]). Predictions largely conform with tissue and cell biology.

Deconvolution of GTEx Database

To assess the use of GEDIT across very large datasets, we applied the tool to 17,382 GTEx

RNA-seq samples collected from various tissues. However, no single reference contained all relevant

cell types. For example, none of the available references contain both myocytes and adipocytes

(Supplementary Figure 1). Therefore, we predicted proportions three times using three separate

references (BlueCode, Human Primary Cell Atlas, Skin Signatures). We then combined these outputs

by taking their median value. This allowed us to produce predictions spanning a larger number of cell

types than are present in any one reference matrix (Figure 6).

Figure 7. GEDIT cell type predictions when applied to 17,382 samples from the GTEx database. Here,

predictions have been averaged for each tissue of origin.

These predictions largely conform to biological expectations. For example, immune cells are

predicted to have high abundance in blood and spleen, adipocytes in adipose tissue, Shwann cells in

nerve and heart, and keratinocytes in skin. Each of these patterns matches expectations of which cell

types should be present in these tissues. Neither cardiac myocytes nor smooth muscle are highly

abundant in GTEx muscle samples. This is likely because the GTEx samples are collected from

skeletal muscle, which is known to have an expression profile that is distinct from that of cardiac and

smooth muscle.

GEDIT Availability

GEDIT can be run online at http://webtools.mcdb.ucla.edu/. Source code, associated data, and

relevant files are available on GitHub at https://github.com/BNadel/GEDIT. We provide access to the

tool, a set of varied reference data, and two sample mixture matrices. The website automatically

produces a heatmap of predicted proportions for the user, as well as a .tsv file. The user also has

access to the parameter choices of GEDIT (signature gene selection method, number of signature

genes, row scaling).

Methods

GEDIT Algorithm

Signature Gene Selection

During signature gene selection, we automatically exclude genes with zero detected

expression in half or more of cell types. Further, we treat all remaining expression values of zero as

the lowest observed non-zero value in the matrix. Implementing this change has minimal effect on

most genes but helps to reduce the scores of very lowly expressed genes. Such lowly expressed

genes are highly susceptible to experimental noise and are generally poor signature genes. Moreover,

including zeros can result in mathematical errors (e.g. dividing by zero, taking the log of zero). We

consider this transformation valid, since values of zero generally do not mean zero expression, but

rather an expression level below the detection limit of the technology used.

For any given gene, a scoring method takes as input the vector of the expression values

across all reference cell types, and outputs a score. A gene is considered a potential signature gene in

cell type X if it is expressed more highly in X than any other cell type. For each cell type, we keep only

the N genes with the highest scores, where N is the NumSigs parameter.

Information entropy (H) is calculated using the following formula:

355
$$H = -\sum_{1}^{i}[p_i * \log_2(p_i)] \qquad (1)$$

where p_i is the probability of the i^{th} observation. To apply this to expression values, we convert

the vector of expression values into a vector of probabilities by dividing by its sum. In an equal mixture

of each cell type, the i^{th} probability can be interpreted as the fraction of transcripts originating from the

i^{th} cell type.

Row Scaling

During this step, we apply a transformation on the expression values for each gene. Each gene

has measured expression in N purified cell types and M samples. Each of these values, X_{old}, is

transformed according to the following formula:

$$\square_{\square\square\square} = (\square_{\square\square\square} - \square\square\square)/(\square\square\square - \square\square\square) * \square\square\square^{\square} \quad (2)$$

Where Min is the minimum of all M + N original values, Max is the maximum of those values,

and p is a tunable parameter with natural range p $\in [0.0, 1.0]$. This procedure produces values

between the range of 0.0 and Max^{p}.

375 Reference Data

BLUEPRINT Reference Dataset

35 gene counts files were downloaded from the BLUEPRINT database, all collected from

venous blood [17]. This included entries for CD14-positive, CD16-negative classical monocytes (5

samples), CD38-negative naive B cells (1), CD4-positive, alpha-beta T cell (8), central memory CD4-

positive, alpha-beta T cell (2), cytotoxic CD56-dim natural killer cell (2), macrophage (4), mature

neutrophil (10), and memory B Cell (1). When two or more transcripts appeared for a single gene, the

transcript with the highest average expression was selected and others were excluded. Genes with no

detected expression in any sample were also excluded, and then each sample was quantile

normalized. Samples generally clustered by cell type, but we excluded one CD4-positive alpha-beta T

cell. Replicates for each cell type were then collapsed into a single entry by taking the median value

for each gene.

ENCODE Reference Dataset

106 transcript quantification files were downloaded from the ENCODE database [18]. These

included all RNA-seq experiments collected from adult primary cells, excluding four with warnings.

Warnings indicated that three samples suffered from low replicate concordance and one sample from

low read depth, and these samples were excluded. All samples were processed by the Gingeras Lab

at Cold Spring Harbor and mapped to GRCH38.

The samples were quantile normalized and clustered. In cases where multiple transcripts were

measured for a single gene, the expression of that gene was calculated as the sum of all transcripts.

At this time, 18 additional samples were excluded as they did not cluster with their replicates. Based

on sample descriptions and data clustering, we found that the remaining 88 samples represented 28

unique cell types. We produced an expression profile for each cell type by merging all samples of that

cell type via median average. For example, a cluster of 19 samples were labelled as endothelial cells

(collected from various body locations) and were merged into a single entry termed canonical

endothelial cells. This dataset spans a wide range of stromal cell types (e.g. smooth muscle,

fibroblast, epithelial), but contains only a single entry for blood cells, which are labelled mononuclear

cells.

We also combined the ENCODE and BLUEPRINT reference matrices into a single reference

matrix, which we call BlueCode. We combined, then quantile normalized, the columns of both

matrices. Possible batch effects in this combined matrix have not been fully evaluated.

10x Reference Dataset

We obtained single cell expression data for nine varieties of immune cells from the 10x website

[19]. This included at least 2446 cells for each cell type, and at least 7566 cells for all cells other than

CD14 monocytes. For each cell type, expression values for all cells were mean averaged to form an

expression profile.

Tabula Muris Reference Dataset

We downloaded from the Tabula Muris single cell data for 12 clusters of mouse cell types. For

each cluster, we averaged all cells of that cluster to produce a reference profile for the corresponding

cell type.

Other Reference Datasets

Other datasets used in this project were obtained from their corresponding publications or

GEO repositories. This includes a reference matrix of human skin signatures, the Human Body Atlas,

the Human Primary Cell Atlas, LM22, ImmunoStates, the Mouse Body Atlas, and ImmGen [13–

16,20,22,23].

Skin Diseases Data

We obtained expression data from 21 skin biopsies, collected from human patients with a

variety of skin diseases. These data originally came from a wide range of sources and platforms, and

were compiled into a single dataset by previous work [34]**.**

GTEx Data

GTExX data for 17,382 samples were obtained from the GTExX database

(https://gtexportal.org/). We ran GEDIT on all samples three times, each time using a different

reference matrix (BlueCode, the Human Primary Cell Atlas, and Skin Signatures). For each cell type,

we calculated our initial estimate as the median estimate across the three sets of predictions (or fewer,

if that cell type is missing from one to two of the reference matrices). Lastly, for each sample we

divided the vector of predictions by its sum, such that the final predictions sum to 100%.

Multi-Tool Performance Evaluation

***In Vitro* Immune Cell Mixture**

Combinations of six immune cells (Neutrophils, Monocytes, Natural Killer Cells, B cells, and

CD4 and CD8 T Cells) were mixed together and sequenced using an affymetrix array. Whole blood

from healthy human donors was supplied with informed consent through a sample sharing agreement

with the UCLA/CFAR Virology Core Lab (grant number 5P30 AI028697). CD4+ T cells, CD8+ T cells,

B cells, and NK cells were isolated using Stem Cell Technologies (Vancouver, BC, Canada)

RosetteSep negative selection. Neutrophils were positively selected through the EasySep approach,

according to the manufacturer's specifications. Cells were then counted by hemocytometer and added

at defined percentages to a total cell count of two million cells to create six different mixtures.

Subsequently cells were processed for RNA isolation by AllPrep DNA/RNA. Illumina HT12 BeadChip

microarray was performed by the UCLA Neuroscience Genomics Core. Data was normalized by

quantile normalization through R 'normalize.quantiles' function (R Core Team, 2013).

RNA-seq Mixtures Used for Tool Evaluation

We also obtained two datasets used in a recent benchmarking study [29]. The first dataset is

composed of three RNA-seq samples, each with two technical replicates that represent biopsies of

ovarian cancer ascites [31]. The second dataset is composed of RNA-seq collected from the blood of

healthy individuals, some of whom recently received an influenza vaccine [30]. These data were

downloaded from the GitHub for the benchmarking paper, which also contained FACS estimates for

six cell types for the ascites data (B cells, dendritic cells, NK cells, T cells, macrophages, neutrophils)

and five cell types for the blood data (B cells, dendritic cells, T cells, monocytes, natural killer cells).

However, since dendritic cells were never present at more than 3.5% abundance, we did not evaluate

performance for this cell type.

Tools

We installed and ran GEDIT, CIBERSORT, DeconRNASeq and dtangle on the hoffman2

computational cluster at UCLA. xCell was run using the online interface at https://xcell.ucsf.edu/. The

default choice for genes signatures (xCell =64) was used. The RNA-seq option was selected for the 2

RNA-seq datasets (blood and ascites), but not for the *in vitro* dataset, which was sequenced on

microarray.

xCell produces 67 output scores, seven of which were used in this study. These were the

entries labelled "B-Cells", "Macrophages", "Monocytes", "NK cells", "Neutrophils", "CD4+ T cells" and

"CD8+ T Cells". As suggested by the xCell authors, the outputs for CD4 and CD8 T cell subtypes were

summed to produce a final output for total T cells.

Reference Data

We evaluated the performance of the four reference-based tools (GEDIT, CIBERSORT,

DeconRNASeq and dtangle) using each of four choices of reference matrix (LM22, ImmunoStates,

BLUEPRINT, and the Human Primary Cell Atlas).The BLUEPRINT and Human Primary Cell Atlas

reference matrices differ from ImmunoStates and LM22 in that they contain tens of thousands of

genes, many of which should not be considered signature genes. This contrasts to ImmunoStates and

LM22; each reference matrix contains fewer than 600 genes, which have been specifically identified

as signature genes by previous work [13,20]. We include both forms of reference matrices in order to

evaluate the input requirements of the tools studied.

Depending on the choice of reference matrix, reference-based tools often produce multiple

outputs for some cell types, each representing a cell sub-type. This includes B cells (naïve and

memory), Monocytes (CD14 and CD16), NK cells (resting and active) and T cells (many subtypes

including varieties of CD4 and CD8). In each case, the outputs for each sub-type were summed in

order to produce a total score for each greater cell type.

Conclusion:

GEDIT is an expression-based cell type quantification tool that offers unprecedented flexibility

and accuracy in a wide variety of contexts. Using both simulated and experimental data, we

demonstrate that GEDIT produces high-quality predictions for multiple platforms, species, and a

diverse range of cell types, outperforming other tools in many cases. We include in the software

package a comprehensive library of reference data, which facilitates application of GEDIT to a wide

range of tissue types in both human and mouse. GEDIT can also accept reference data supplied by

the user, which can be derived from bulk RNA-seq, scRNA-seq, or microarray experiments. GEDIT

represents a competitive addition to the suite of existing tissue decomposition tools while maintaining

for users flexibility and performance robustness.

As part of this project, we perform a study in which we compare the performance of several

deconvolution tools using multiple metrics. Unlike previous evaluation studies, we explore the effect of

reference choice by running tools multiple times with reference data from different sources. We find

that, while choice of optimal reference is a complicated problem, the performance of GEDIT is highly

robust to choice of reference, relative to other tools.

When extensively applied to several large public datasets, GEDIT produces predicted cell type

fractions that conform with biological expectations. When used to decompose skin biopsies,

keratinocytes are found to be the most abundant cell type and variations in the abundance of other cell

types conform to expected immune responses across diseases. Similarly, cell type predictions of

GTEx samples are concordant with our expectations of the dominant cell types across tissues.

Schwann cells, keratinocytes, adipose cells, and immune cells are found to be most abundant in

nerve, skin, adipose tissue, and blood, respectively.

Single cell RNA-seq is an emerging approach to study the composition of cell types within a

sample. Due to biases associated with the capture of different cell types, these methods are not

always capable of accurately quantifying cell type populations [8]. However, the pure reference

profiles produced by existing methods can be used by GEDIT to generate accurate estimates of cell

type populations. Thus, GEDIT circumvents some of the biases associated with the preparation of

samples for both scRNA-seq and FACS. GEDIT is freely available, and therefore an extremely

economical option to researchers, particularly those who profile expression data for other purposes.

GEDIT produces accurate results when tested on mixtures of human immune cells. Compared

to other tools, GEDIT produces the lowest error in majority of scenarios in the studied mixtures.

GEDIT provides increased flexibility over previously developed tools, as we provide a set of reference

matrices for varied cell types for both mouse and human datasets.

GEDIT provides unique advantages compared to other tools, especially in terms of cell type,

species and platform flexibility, and constitutes a useful addition to the existing set of tools for tissue

decomposition. Our efficient decomposition methodology has been extensively optimized and we find

that it performs robustly across a broad range of tissues in both mouse and human datasets. Our

future work will extend reference matrices to facilitate application of GEDIT on varied bulk gene

expression datasets.

Availability of Source Code and Requirements

- Project name: GEDIT

- Project Home Page: https://github.com/BNadel/GEDIT

- Programming Languages: Python 2.0, R

- Other requirements: numpy, glmnet

- Operating Systems: Linux

- License: MIT

Availability of Data and Materials

All data used in this paper are freely available on GitHub

(https://github.com/purebrawn/GEDIT), as well as their original sources. Code for DeconRNASeq was

obtained as an R package from the CRAN repository. Code for CIBERSORT was obtained by

requesting it via the web portal (https://cibersort.stanford.edu/download.php), and code for dtangle

from the project's GitHub page (https://github.com/BNadel/GEDIT).

Reference data is also available from their original sources. Most datasets can be found on

project website pages or from public databases. These include BLUEPRINT (http://www.blueprint-

epigenome.eu/), ENCODE (https://www.encodeproject.org), the Human Primary Cell Atlas

(http://biogps.org/dataset/BDS_00013/primary-cell-atlas/), LM22 (http://cibersort.stanford.edu/ or

GEO:GSE65136), 10x Genomics (https://support.10xgenomics.com/single-cell-gene-

expression/datasets), Tabula Muris (https://tabula-muris.ds.czbiohub.org/), the Mouse Body Atlas

(GEO:GSE10246), and ImmGen (http://www.immgen.org/Databrowser19/DatabrowserPage.html).

Some reference matrices were obtained as supplementary files from the publications listed in Table 1.

Expression values for the blood and ascites RNA-seq datasets were obtained from the GitHub

repository https://github.com/grst/immune_deconvolution_benchmark, and are also available at at

https://figshare.com/s/711d3fb2bd3288c8483a and GEO: GSE64655). The *in vitro* mixture of immune

cells was prepared by our lab, and available on our GitHub page.

Acknowledgments

We acknowledge the Biomedical Big Data Grant (5T32LM012424-03) for supporting Brian Nadel during the course of this research. We also acknowledge the Bruins-in-Genomics Summer Undergraduate Research Program for supporting Hannah Waddel and Misha Khan during the summer of 2017, when they contributed to this work. We also thank Lana Martin for her help with editing and proofreading the manuscript.

References

1. Bolen CR, Uduman M, Kleinstein SH. Cell subset prediction for blood genomic studies. BMC Bioinformatics. 2011;12: 258.

2. Gentles AJ, Newman AM, Liu CL, Bratman SV, Feng W, Kim D, et al. The prognostic landscape of genes and infiltrating immune cells across human cancers. Nat Med. 2015;21: 938–945.

3. Li B, Severson E, Pignon J-C, Zhao H, Li T, Novak J, et al. Comprehensive analyses of tumor immunity: implications for cancer immunotherapy. Genome Biol. 2016;17: 174.

4. Fridman WH, Pagès F, Sautès-Fridman C, Galon J. The immune contexture in human tumours: impact on clinical outcome. Nat Rev Cancer. 2012;12: 298–306.

5. Şenbabaoğlu Y, Gejman RS, Winer AG, Liu M, Van Allen EM, de Velasco G, et al. Tumor immune microenvironment characterization in clear cell renal cell carcinoma identifies prognostic and immunotherapeutically relevant messenger RNA signatures. Genome Biol. 2016;17: 231.

6. Gierahn TM, Wadsworth MH, Hughes TK, Bryson BD, Butler A, Satija R, et al. Seq-Well: portable, low-cost RNA sequencing of single cells at high throughput. Nature Methods. 2017. pp. 395–398. doi:10.1038/nmeth.4179

7. Macosko EZ, Basu A, Satija R, Nemesh J, Shekhar K, Goldman M, et al. Highly Parallel Genome-wide Expression Profiling of Individual Cells Using Nanoliter Droplets. Cell. 2015;161: 1202–1214.

8. Hines WC, Su Y, Kuhn I, Polyak K, Bissell MJ. Sorting out the FACS: a devil in the details. Cell Rep. 2014;6: 779–781.

9. Frishberg A, Peshes-Yaloz N, Cohn O, Rosentul D, Steuerman Y, Valadarsky L, et al. Cell composition analysis of bulk genomics using single-cell data. Nat Methods. 2019;16: 327–332.

10. Wang X, Park J, Susztak K, Zhang NR, Li M. Bulk tissue cell type deconvolution with multi-subject single-cell expression reference. Nat Commun. 2019;10: 380.

11. Frishberg A, Brodt A, Steuerman Y, Gat-Viks I. ImmQuant: a user-friendly tool for inferring immune cell-type composition from gene-expression data. Bioinformatics. 2016;32: 3842–3843.

12. Aran D, Hu Z, Butte AJ. xCell: digitally portraying the tissue cellular heterogeneity landscape. Genome Biol. 2017;18: 220.

13. Newman AM, Liu CL, Green MR, Gentles AJ, Feng W, Xu Y, et al. Robust enumeration of cell subsets from tissue expression profiles. Nat Methods. 2015;12: 453–457.

14. Swindell WR, Johnston A, Voorhees JJ, Elder JT, Gudjonsson JE. Dissecting the psoriasis
transcriptome: inflammatory- and cytokine-driven gene expression in lesions from 163 patients.
BMC Genomics. 2013;14: 527.

15. Su AI, Wiltshire T, Batalov S, Lapp H, Ching KA, Block D, et al. A gene atlas of the mouse and
human protein-encoding transcriptomes. Proc Natl Acad Sci U S A. 2004;101: 6062–6067.

16. Mabbott NA, Baillie JK, Brown H, Freeman TC, Hume DA. An expression atlas of human primary
cells: inference of gene function from coexpression networks. BMC Genomics. 2013;14: 632.

17. Martens JHA, Stunnenberg HG. BLUEPRINT: mapping human blood cell epigenomes.
Haematologica. 2013;98: 1487–1489.

18. ENCODE Project Consortium. The ENCODE (ENCyclopedia Of DNA Elements) Project. Science.
2004;306: 636–640.

19. Zheng GXY, Terry JM, Belgrader P, Ryvkin P, Bent ZW, Wilson R, et al. Massively parallel digital
transcriptional profiling of single cells. Nat Commun. 2017;8: 14049.

20. Vallania F, Tam A, Lofgren S, Schaffert S, Azad TD, Bongen E, et al. Leveraging heterogeneity
across multiple datasets increases cell-mixture deconvolution accuracy and reduces biological
and technical biases. Nat Commun. 2018;9: 4735.

21. Consortium TTM, The Tabula Muris Consortium, Coordination O, Coordination L, Organ collection
and processing, Library preparation and sequencing, et al. Single-cell transcriptomics of 20
mouse organs creates a Tabula Muris. Nature. 2018. pp. 367–372. doi:10.1038/s41586-018-
0590-4

22. Lattin JE, Schroder K, Su AI, Walker JR, Zhang J, Wiltshire T, et al. Expression analysis of G
Protein-Coupled Receptors in mouse macrophages. Immunome Res. 2008;4: 5.

23. Heng TSP, Painter MW, Immunological Genome Project Consortium. The Immunological
Genome Project: networks of gene expression in immune cells. Nat Immunol. 2008;9: 1091–
1094.

24. Gong T, Szustakowski JD. DeconRNASeq: a statistical framework for deconvolution of
heterogeneous tissue samples based on mRNA-Seq data. Bioinformatics. 2013;29: 1083–1085.

25. Hunt GJ, Freytag S, Bahlo M, Gagnon-Bartsch JA. dtangle: accurate and robust cell type
deconvolution. Bioinformatics. 2018 [cited 17 Jan 2019]. doi:10.1093/bioinformatics/bty926

26. Becht E, Giraldo NA, Lacroix L, Buttard B, Elarouci N, Petitprez F, et al. Estimating the population
abundance of tissue-infiltrating immune and stromal cell populations using gene expression.
Genome Biol. 2016;17: 218.

27. Lopez D, Montoya D, Ambrose M, Lam L, Briscoe L, Adams C, et al. SaVanT: a web-based tool
for the sample-level visualization of molecular signatures in gene expression profiles. BMC
Genomics. 2017;18: 824.

28. Altboum Z, Steuerman Y, David E, Barnett-Itzhaki Z, Valadarsky L, Keren-Shaul H, et al. Digital
cell quantification identifies global immune cell dynamics during influenza infection. Mol Syst Biol.
2014;10: 720.

29. Sturm G, Finotello F, Petitprez F, Zhang JD, Baumbach J, Fridman WH, et al. Comprehensive
evaluation of transcriptome-based cell-type quantification methods for immuno-oncology.

Bioinformatics. 2019;35: i436–i445.

30. Hoek KL, Samir P, Howard LM, Niu X, Prasad N, Galassie A, et al. A Cell-Based Systems Biology
Assessment of Human Blood to Monitor Immune Responses after Influenza Vaccination. PLOS
ONE. 2015. p. e0118528. doi:10.1371/journal.pone.0118528

31. Schelker M, Feau S, Du J, Ranu N, Klipp E, MacBeath G, et al. Estimation of immune cell content
in tumour tissue using single-cell RNA-seq data. Nat Commun. 2017;8: 2032.

32. de Sousa JR, Lucena Neto FD, Sotto MN, Quaresma JAS. Immunohistochemical characterization
of the M4 macrophage population in leprosy skin lesions. BMC Infect Dis. 2018;18: 576.

33. Lin C-C, Chen C-B, Wang C-W, Hung S-I, Chung W-H. Stevens-Johnson syndrome and toxic
epidermal necrolysis: risk factors, causality assessment and potential prevention strategies.
Expert Rev Clin Immunol. 2020;16: 373–387.

34. Inkeles MS, Scumpia PO, Swindell WR, Lopez D, Teles RMB, Graeber TG, et al. Comparison of
molecular signatures from multiple skin diseases identifies mechanisms of immunopathogenesis.
J Invest Dermatol. 2015;135: 151–159.

Cell Type	Monocytes	Neutrophils	B Cells	NK Cells	CD4+ T cells	Macrophages
CD14	338.4	163.9	18.9	16.9	19.2	105.9
THEMIS	9.7	11.6	8.4	13.2	52.0	8.7

Mixture 1	Mixture 2
22.3	95.0
50.3	20.3

Row Scaling Transformation

Cell Type	Monocytes	Neutrophils	B Cells	NK Cells	CD4+ T cells	Macrophages
CD14	1.0	.46	.01	0.0	.01	.28
THEMIS	.03	.07	0.0	.11	1.0	.01

Mixture 1	Mixture 2
.02	.24
.96	.27

PROSTATE
BLADDER
ESOPHAGUS
THYROID
PANCREAS
STOMACH
KIDNEY
SALIVARY GLAND
VAGINA
MUSCLE
ADRENAL GLAND
LUNG
LIVER
SMALL INTESTINE
PITUITARY
TESTIS
BRAIN
BLOOD VESSEL
UTERUS
OVARY
CERVIX UTERI
FALLOPIAN TUBE
COLON
HEART
NERVE
ADIPOSE TISSUE
BREAST
SKIN
SPLEEN
BLOOD

Endothelial
Smooth Muscle
Other
Fibroblasts
Schwann Cells
Keratinocyte
Epithelial
Adipocyte
Total Immune

Color Key and Histogram

Count
100
0

0.2 0.4 0.6 0.8

Predicted Abundance