Title: Freezing and water availability structure the evolutionary diversity of trees across the Americas

R. A. Segovia[1,2], R. T. Pennington[3,4], T. R. Baker[5], F. Coelho de Souza[5], D. M. Neves[6], C. C. Davis[7], J. J. Armesto[2,8], A. T. Olivera-Filho[6], K. G. Dexter[1,3]

School of GeoSciences, University of Edinburgh, United Kingdom.
Instituto de Ecología y Biodiversidad (www.ieb-chile.cl), Santiago, Chile.
Tropical Diversity Section, Royal Botanic Garden Edinburgh, United Kingdom.
Department of Geography, University of Exeter, United Kingdom.
School of Geography, University of Leeds, Leeds, United Kingdom.
Department of Botany, Federal University of Minas Gerais, Brazil.
Department of Organismic and Evolutionary Biology, Harvard University, Cambridge, United States.
Departamento de Ecología, Universidad Católica de Chile, Santiago and Facultad Ciencias Naturales and Oceanográficas, Universidad de Concepción, Chile.

* ricardo.segovia@ed.ac.uk

Abstract

The historical course of evolutionary diversification shapes the current distribution of biodiversity, but the main forces constraining diversification are unclear. We unveil the evolutionary structure of tree species diversity across the Americas to assess whether an inability to move (dispersal limitation) or to evolve (niche conservatism) is the predominant constraint in plant diversification and biogeography. We find a fundamental divide in tree lineage composition between tropical and extratropical environments, defined by the absence versus presence of freezing temperatures. Within the Neotropics, we uncover a further evolutionary split between moist and dry forests. Our results demonstrate that American tree lineages, though broadly distributed geographically, tend to retain their ancestral environmental relationships and that phylogenetic niche conservatism is the primary force structuring the distribution of tree biodiversity.

Main text

A central challenge in biogeography and macroevolution is to understand the primary forces that drove the diversification of life. Was diversification confined within continents, and characterized by adaptation of lineages to different major environments (i.e., biome switching), or did lineages tend to disperse across great distances, but retain their ancestral environmental niche (i.e., phylogenetic niche conservatism)? Classically, the attempts to define biogeographic regions based on shared plant and animal distributions lend support to the first hypothesis, that large-scale patterns may be explained by regionally confined evolutionary diversification, rather than long-distance dispersal *(1-3)*. However, recent studies of the distribution of plant lineages at global scales have documented high levels of inter-continental dispersal (e.g., *4-8*), and revealed that lineages tend to retain their ancestral biomes when dispersing *(9,10)*. These latter findings suggest that environmental associations of lineages may be the primary force organizing the course of diversification, but we lack studies comparing the degree of evolutionary similarity between species assemblages at broad scales to adequately resolve this debate.

      With high mountain chains running north to south across latitudes and a mosaic of contrasting environments, the Americas represent a natural laboratory to investigate the evolutionary forces behind the modern distribution of biodiversity. Here, we examine the phylogenetic composition of angiosperm tree assemblages across the Americas as a means to determine whether dispersal limitation or phylogenetic niche conservatism had a greater impact on the present-day evolutionary structure of biodiversity. If lineages tend to retain their environmental niche as they diversify across space, we would expect major evolutionary groups to be restricted to specific environmental regimes. This leads to the prediction that lineage composition of assemblages from extratropical regions in both hemispheres should be more similar to each other than to assemblages occurring in intervening tropical regions. In addition, we would predict that assemblages from dry tropical environments should show greater similarity in tree lineage composition to each other than to assemblages from moist environments with which they may be spatially contiguous *(11)*. Alternatively, if diversification is spatially restricted and biome switching is common, the major evolutionary grouping of assemblages should be segregated geographically. Thus, we would predict assemblages from South America (which was physical isolated through the Cenozoic) to constitute one group and assemblages from North and Central America another.

To test the relative importance of phylogenetic niche conservatism versus dispersal limitation, we analyzed data from $\sim 10{,}000$ tree assemblages with a new temporally-calibrated, genus-level phylogeny that includes 1,358 genera ($\sim 90\%$ of tree genera sampled per assemblage). We assessed similarity in lineage composition among assemblages using clustering analyses and ordinations based on shared phylogenetic branch length. Next, we identified the indicator lineages for each major group in the clustering analysis and explored the geographic and environmental correlates of the distribution of the main evolutionary clusters. Finally, we estimated the unique evolutionary diversity (i.e. sum of phylogenetic branches of lineages restricted to individual groups) versus shared evolutionary diversity (i.e. sum of shared phylogenetic branches) across evolutionary groups (for details see SM).

We show that the evolutionary lineage composition of American tree assemblages is structured primarily by phylogenetic niche conservatism. The two principal groups have a tropics-extratropics structure (Fig. 1). The extratropical group is not geographically segregated, but includes temperate tree assemblages from North America and southern South America, connected by a

Figure 1 The geographic, evolutionary and environmental relationships of the two principal evolutionary groups (from $K{=}2$ clustering analysis). A) Geographic distribution of angiosperm tree assemblages and their affiliation with either the tropical (n = 7145) or extratropical (n = 2792) evolutionary group; **B)** Distribution of assemblages over elevation and latitude showing that the extratropical group is largely restricted to high elevations at low latitudes; **C & D)** Distribution of assemblages over the first two axes of an ordination based on evolutionary composition with assemblages in **C** colored according to group affiliation and in **D** as to whether or not they experience freezing temperatures in a regular year (from (*31*)).

high-elevation corridor in low latitudes (Fig. 1 a,b). The tropics-extratropics structure of tree evolutionary diversity shows a strong correspondence (97% match, Fig. S1) with the absence vs. occurrence of freezing temperatures within a typical year (see Fig. 1 c,d). We observe that most evolutionary diversity, measured as summed phylogenetic branch length, occurs within the tropics, but that there is unique evolutionary diversity restricted to the extratropics ($\sim 10\%$ of the total, Fig. 2b, S3a). Ordination and indicator clade analyses revealed that the tropics-extratropics segregation is associated with the distribution of specific clades, such as the Fagales, which includes the oaks (*Quercus*), beeches (*Fagus*), coihues (*Nothofagus*) and their relatives (Fig. 3, Table S1, S2).

Cluster analyses of *K*=3 and *K*=4 groups are also supported as additional informative splits (Fig. S2), and each of their major groups capture substantial unique evolutionary diversity (Fig. 2 b, Fig. S3, Table S2). In *K*=3, the main extratropical cluster grouped assemblages from North America and extreme southern South America, while the remaining assemblages from temperate

Figure 2. The geographic, evolutionary and environmental relationships among four evolutionary groups (from *K*=4 clustering analysis). A) Geographic distribution of angiosperm tree assemblages and their affiliation with one of the four evolutionary groups; **B)** Euler Diagram representing the amount of evolutionary history, quantified as phylogenetic diversity (in millions of years), restricted to each cluster versus that shared between clusters; **C)** Distribution of assemblages over extremes of temperature (minimum temperature of coldest month) and water availability (maximum climatological water deficit, CWD). Lines represent the 95th quantile of the density of points for each group.

southern South America and the Andean tropics grouped with assemblages from the arid or semiarid tropics and subtropics (Fig. S4). The third group was formed by the moist tropics (Fig. S4). For $K=4$, the extratropics were split into a largely temperate North American group and a second group that joins subtropical sites in South and Central America with southern temperate forests and high elevation sites in the Andes (Fig. 2 a). In the tropics there is one group uniting assemblages found in ever-moist and warm conditions, and a second one of assemblages that extend into drier areas (Fig. 2 c), including most tropical dry forest assemblages (Fig. 2 a; Table S3). Hereafter, we refer to the four groups of assemblages in $K=4$ as the Northern Extratropical, Southern Extratropical, Tropical Moist and Tropical Dry groups.

Our results demonstrate that the tropics-extratropics evolutionary structure of tree diversity is principally associated with the environmental threshold of the presence versus absence of freezing temperatures (Fig. 1a,b, S1). This pattern is consistent with evidence documenting that only angiosperm lineages that were able to evolve traits to avoid freezing-induced embolism radiated into high latitudes [12]. In addition, we found that a unique, sizeable portion of the total evolutionary diversity of angiosperm trees is restricted to extratropical environments, as the fossil record corroborates [13,14]. Collectively, this evidence suggests that the phylogenetic conservatism of lineages from the extratropics has a major relevance for the diversification of angiosperm trees in the Americas. Kerkhoff *et al.* [15] estimated that in the extratropical region (defined by them as areas north of 23°N and south of 23°S) angiosperm ancestors produced extratropical descendants at least 90% of the time. Considering that some areas subjected to regular freezing at high elevations in equatorial latitudes may be better classified as part of the extratropics, as demonstrated here by our results, extratropical phylogenetic conservatism could even be greater than found by Kerkhoff *et al.* [15].

We suggest that the extratropical conservatism has a major importance in the biogeography of the Americas. The relatively recent uplift of the Andes would have created novel environments, with regular freezing temperatures, at low latitudes, allowing extratropiocal lineages to move from both north and south to equatorial latitudes [16,17]. Fossil pollen demonstrates the arrival in the northern Andes of tree genera from temperate forests in the northern hemisphere, including *Juglans* (Juglandaceae), *Alnus* (Betulaceae) and *Quercus* (Fagaceae), at about 2.2 Ma, 1.0 Ma and 300 Ka respectively, and the arrival of southern genera, including *Weinmannia* (Cunoniaceae) and *Drymis* (Winteraceae), during the late Pliocene and Pleistocene (1.5–3.2 Ma) [16,18].

Likewise, phylogenetic evidence shows recent diversification in the Andes of lineages that seem to have originated in the extratropics, including *Lupinus* (Fabaceae) *(19)*, Adoxaceae/Valerianceae *(20,21)* and *Gunnera* (Gunneraceae) *(22)*.

Figure 3. Phylogenetic ordination of tree assemblages based on their evolutionary lineage composition. Colors in the main plot represent the groups from $K=4$ clustering analyses and the different symbols represent major vegetation formations. The subset plot shows the clades most strongly associated with the first two axes of the evolutionary ordination.

Our results also point to a moist versus dry evolutionary divide within the Neotropics. The Tropical Moist Group holds the greatest amount of evolutionary diversity, both overall and unique to it, despite occupying the most restricted extent of climatic space of any of the $K=4$ groups (Fig. 2 b,c). The Tropical Dry Group, in contrast, extends across a broader climatic space, but holds less evolutionary diversity (Fig. 2 b,c). This asymmetry in the accumulation of diversity may reflect phylogenetic conservatism for a putatively moist and hot ancestral angiosperm niche *(23)*, or could result from a favorable environment that can be occupied by any angiosperm lineage, even those that also occur in cooler or drier conditions *(24,25)*. Regardless, the similarity in the lineage composition of the extensive but discontinuously distributed tropical dry forests *(11)* indicates their separate evolutionary history. Although tropical dry forest inhabiting taxa have often been described as more dispersal-limited than those from rain forests (e.g., *11*), dispersal over evolutionary time-scales seems to have been sufficient to maintain this floristic cohesion. Such evolutionary isolation of the dry forest flora has previously been suggested by studies in

Fabaceae *(11,26)*, and is shown here to be evident at the evolutionary scale of all angiosperm tree species.

Our results also help to clarify the contentious evolutionary status of savanna and Chaco regions in the Neotropics. We find that the southern savannas (the Cerrado region of Brazil) are more evolutionary related to tropical moist forests than dry forests (Fig. 2 a, Fig. S4), as previously suggested *(26,27)*. However, northern tropical savannas (i.e., Llanos of Venezuela and Colombia and those in Central America) are split in their evolutionary linkages between the Tropical Moist and Tropical Dry groups (Fig. 3, Table S1). This may reflect the distinct ecology of many northern savannas (e.g., the Llanos are hydrological savannas; *28*) and suggest a divergent evolutionary history for northern and southern savannas. Our results may also help to resolve the debates around the evolutionary affinities of the Chaco (e.g., *29,30*), by showing that this geographically defined region houses a mix of extratropical and tropical lineages (Fig. 2). Furthermore, our analyses consistently point to evolutionary links between assemblages in seasonally dry and seasonally cold areas (Fig. 2, S4). For example, when we consider $K=3$ evolutionary groups, a single 'dry and cool' group coalesces, with the other two groups being the tropical moist forest group and a largely northern, extratropical group (Fig. S4).

We show that the evolutionary structure of tree diversity in the Americas is determined primarily by the presence versus absence of freezing temperatures, dividing tropical from extratropical regions. Within the tropics we find further subdivision among lineages experiencing moist versus seasonally-dry conditions. These findings clearly demonstrate that phylogenetic niche conservatism is the primary force organizing the diversification and, therefore, the biogeography of angiosperm trees. Tree species that can inhabit areas experiencing freezing temperatures and/or environments subjected to seasonal water stress belong to a restricted set of phylogenetic lineages, which gives a unique evolutionary identity to extratropical forests and tropical dry forests in the Americas. While our study is restricted to New World trees, we suggest that plant biodiversity globally may be evolutionarily structured following a tropics-extratropics pattern, while diversity within the tropics may be structured primarily around a moist-dry pattern. These findings advocate strongly for integrating the concept of extratropical conservatism and tropical-dry conservatism into our understanding of global macroevolutionary trends and biogeographic patterns.

Materials and Methods

Tree assemblage dataset

Our tree assemblage dataset was derived by combining the NeoTropTree (NTT)
database *(32)* with selected plots from the Forest Inventory and Analysis (FIA)
Program of the U.S. Forest Service *(33)*, accessed on July 17th, 2018 via the
BIEN package *(34)*. We excluded from the latter any sites that had less than
five angiosperm genera. Sites in the NTT database are dened by a single
vegetation type within a circular area of 5-km radius and contains records of
tree and tree-like species, i.e., freestanding plants with stems that can reach over
3m in height (see www.neotroptree.info and *(35)* for details). Each FIA plot
samples trees that are ≥ 12.7 cm diameter at breast height (dbh) in four
subplots (each being 168.3 m2) that are 36.6 m apart. We aggregated plots from
the FIA dataset within 10 km diameter areas, to parallel the spatial structure of
the NTT database. This procedure produced a total dataset of 9,937 tree
assemblages distributed across major environmental and geographic gradients in
the Americas.

Genera phylogenetic tree

We obtained sequences of the *rbc*L and *mat*K plastid gene for 1,358 angiosperm
tree genera, from Genbank (www.ncbi.nlm.nih.gov/genbank/), building on
previous large-scale phylogenetic efforts for angiosperm trees in the Neotropics
(36,37). Sequences were aligned using the MAFFT software *(38)*. 'Ragged ends'
of sequences that were missing data for most genera were manually deleted from
the alignment.

We estimated a maximum likelihood phylogeny for the genera in the
RAxML v8.0.0 software *(39)*, on the CIPRES web server (www.phylo.org). We
constrained the tree to follow the order-level phylogeny in Gastauer *et al.* (2017)
(40), which is based on the topology proposed by the Angiosperm Phylogeny
Group IV. We concatenated the two chloroplast markers following a General
Time Reversible (GTR) + Gamma (G) model of sequence evolution. We
included sequences of *Nymphaea alba* (Nymphaeaceae) as an outgroup.

We temporally calibrated the maximum likelihood phylogeny using the
software treePL *(41)*. We implemented age constraints for 320 internal nodes
(family-level or higher, from *(42)*) and for 123 genera stem nodes (based on ages
from a literature survey, Table S4). The rate smoothing parameter (lambda)
was set to 10 based on a cross-validation procedure. The final dated tree can be

found in Supplementary Information.

Phylogenetic distance analysis and clustering

We used the one complement of the Phylo-Sorensen Index (i.e., $1 -$ Phylo-Sorensen) to build a matrix of phylogenetic dissimilarities between plots based on genera presence-absence data. The Phylo-Sorensen index sums the total branch length of shared clades between sites *(43)* relative to the sum of branch lengths of both sites:

$$Complement\ of\ Phylo - Sorensen\ ij = 1 - BLij/0.5 * (BLi + BLj)$$

where BLij is the sum of branch lengths shared between plots i and j, and BLi and BLj are the sum of branch length of tips within plots i and j, respectively. Thus, if all branches are shared between two plots, the dissimilarity measure takes on a value of 0. If no branches are shared between plots (i.e. the plots comprise two reciprocally monophyletic clades), the dissimilarity measure will take on a value of 1. This metric was estimated using the phylosor.query() function in the PhyloMeasures *(44)* package for R.

We used *K-means* clustering to explore the main groups, in terms of (dis)similarity in the tree assemblage dataset, according to the Phylo-Sorensen dissimilarity measures. The *K-means* clustering algorithm requires the number of clusters (K) to be specified in advance. In order to estimate the best value for K, the optimal number of clusters to parsimoniously explain the variance in the dataset, we used the Elbow Method and an approach based on the average Silhouette width (Fig. S2). Based on these results, we selected K=2 (Fig. 1), K=3 (Fig. S5) and K=4 (Fig. 2) for further analysis and interpretation. No geographic or environmental data were used to inform the clustering analyses. The K-means clustering was carried out with the kmeans() function in base R (R Core Development Team, 2016). We assessed the robustness of the *K-means* clustering results using a silhouette analysis with functions in the "cluster" package *(45)*. In order to assess variation in group fidelity, we classified individual sites as to whether the silhouette widths were larger or smaller than 0.2. In this way, we could detect areas of geographic, environmental and compositional space where clustering results were strongly or weakly supported.

In addition, we performed an evolutionary ordination of tree assemblages based on their phylogenetic lineage composition, following protocols developed by Pavoine (2016) *(46)*. We specifically used an evolutionary PCA, implemented with the evopca() function in the "adiv" package *(47)*, with a Hellinger

transformation of the genus by site matrix, as this is a powerful approach to   254
detect phylogenetic patterns along gradients, while also allowing positioning of   255
sites and clades in an ordination space *(46)*. The first two axes explained 9.6%   256
and 6.7% of the variation in the data, with subsequent axes each explaining   257
<5.5%.   258

Correspondence between clustering results and environmental   259
variables   260

We tested the correlation between our *K=2* clustering result and eight different   261
delimitations of the tropics, as per Feeley and Stroud (2018) *(31)*. These   262
delimitations were: C1) all areas between 23.4°S and 23.4°N; C2) all areas with   263
a net positive energy balance; C3) all areas where mean annual temperature   264
does not co-vary with latitude; C4) all areas where temperatures do not go below   265
freezing in a typical year; C5) all areas where the mean monthly temperature is   266
never less than 18°C; C6) all areas where the mean annual "biotemperature"   267
≥ 24 °C; C7) all areas where the annual range of temperature is less than the   268
average daily temperature range; C8) all areas where precipitation seasonality   269
exceeds temperature seasonality. We calculated the correspondence between our   270
binary clustering (i.e. tropical vs. extratropical) and each of these delimitations   271
as the proportion of sites where the delimitations matched.   272

To assess the environmental space occupied by different groups from our   273
clustering analyses, we obtained estimates of mean annual temperature, mean   274
annual precipitation and minimum temperature of the coldest month from the   275
Worldclim dataset *(48)* and Maximum Climatological Water Deficit (CWD)   276
from Chave *et al.* (2014) *(49)*. We estimated the density of the distribution of   277
sites in the environmental space using ellipses containing 95% of the sites with   278
the kde() function from *"ks"* package *(50)*.   279

Shared versus Unique "Phylogenetic Diversity" (PD)   280

As the Phylo-Sorensen estimation of evolutionary (dis)similarity cannot   281
distinguish variation associated to differences in total phylogenetic diversity   282
(PD), or phylogenetic richness versus variation associated to phylogenetic   283
turnover per se, we measured the shared and unique PD associated with each   284
group for the *K=2*, *K=3* and *K=4* clustering analyses. First, we estimated the   285
association of genera with each group by an indicator species analysis following   286
de Caceres *et al.* (2009) *(51)*. Specifically, we used the multipatt() function in   287
the R Packages indicspecies *(52)* to allow genera to be associated with more   288

than one group (when $K > 2$). The output of the *multipatt ()* function includes the stat index, which is a function of the specificity (the probability that a surveyed site belongs to the target site group given the fact that the genus has been found) and fidelity (the probability of finding the genus in sites belonging to the given site group). We constructed pruned phylogenies including those genera with specificity greater than 0.6 to a group, or combination of groups, to estimate the total PD found in each group or combination of groups. Then, we subtracted these totals from the total for the complete, unpruned phylogeny to determine the amount of phylogenetic diversity restricted to each group, or combination of groups. Finally, we estimated the PD shared across all groups as that which was not restricted to any particular group or any combination of groups. We fit these different PD totals as areas in a Euler diagram with the euler() function in the *"eulerr"* package *(53)* for the K=2 and K=3 clustering, and with the Venn() fuction in the *"venn"* package *(54)* for the K=4 clustering.

Indicator lineages for clusters

In order to further characterise the composition of the evolutionary groups, we conducted an indicator analysis to determine the clades most strongly associated with each group. We created a site x node matrix (see function used in Appendix 1), which consists of a presence/absence matrix for each internal node in the phylogeny and ran an indicator analysis for the nodes. We selected the highest-level, independent (i.e. non-nested) nodes with the highest stat values to present in Tables S1 and S2. The indicator node analysis was carried out with function *multipatt()* in the R Package indicspecies *(52)*.

References

1. Wallace, A.R. (1876). The geographical distribution of animals: With a study of the relations of living and extinct faunas as elucidating the past changes of the earth's surface. Harper; brothers.

2. A. Takhtajan, Floristic regions of the world, vol. 581 (University of California Press, Berkeley, CA, 1986).

3. B. G. Holt, J.-P. Lessard, M. K. Borregaard, S. A. Fritz, M. B. Araujo, D. Dimitrov, P.- H. Fabre, C. H. Graham, G. R. Graves, K. A. Jonsson, D. Nogues-Bravo, Z. Wang, R. J. Whittaker, J. Fjeldsa, C. Rahbek, An Update of Wallace's Zoogeographic Regions of the World. Science 339, 74-78 (2013).

4. R. Morley, Interplate dispersal paths for megathermal angiosperms. Perspect. Plant Ecol 6, 5-20 (2003).

5. R. T. Pennington, C. W. Dick, The role of immigrants in the assembly of the south american rainforest tree flora. Philos. TR Soc. B 359, 1611–1622 (2004).

6. S. Renner, Plant dispersal across the tropical atlantic by wind and sea currents. Int. J. Plant. Sci. 165, S23–S33 (2004).

7. M. Lavin, B. P. Schrire, G. Lewis, R. T. Pennington, A. Delgado-Salinas, M. Thulin, C. E. Hughes, A. B. Matos, M. F. Wojciechowski, Metacommunity process rather than continental tectonic history better explains geographically structured phylogenies in legumes. Philos. TR Soc. B 359, 1509–1522 (2004).

8. J. F. Slik, J. Franklin, V. Arroyo-Rodrıguez, R. Field, S. Aguilar, N. Aguirre, J. Ahumada, S.-I. Aiba, L. F. Alves, K. Anitha, *et al.*, Phylogenetic classification of the worlds tropical forests. PNAS USA 115, 1837–1842 (2018).

9. M. D. Crisp, M. T. K. Arroyo, L. G. Cook, M. A. Gandolfo, G. J. Jordan, M. S. McGlone, P. H. Weston, M. Westoby, P. Wilf, H. P. Linder, Phylogenetic biome conservatism on a global scale. Nature 458, 754-U90 (2009).

10. E. Gagnon, J. J. Ringelberg, A. Bruneau, G. P. Lewis, C. E. Hughes, Global succulent biome phylogenetic conservatism across the pantropical caesalpinia group (leguminosae). New Phytol. 222, 1994-2008 (2018).

11. R. T. Pennington, M. Lavin, A. Oliveira-Filho, Woody plant diversity, evolution, and ecology in the tropics: perspectives from seasonally dry tropical forests. Annu. Rev. Ecol. Evol. Syst. 40, 437–457 (2009).

12. A. E. Zanne, D. C. Tank, W. K. Cornwell, J. M. Eastman, S. A. Smith, R. G. FitzJohn, D. J. McGlinn, B. C. O'Meara, A. T. Moles, P. B. Reich, D. L. Royer, D. E. Soltis, P. F.Stevens, M. Westoby, I. J. Wright, L. Aarssen, R. I. Bertin, A. Calaminus, R. Govaerts, F. Hemmings, M. R. Leishman, J. Oleksyn, P. S. Soltis, N. G. Swenson, L. Warman, J. M. Beaulieu, Three keys to the radiation of angiosperms into freezing environments. Nature 506, 89-92 (2014).

13. C. Jaramillo, A. Cardenas, Global warming and neotropical rainforests: a historical perspective. Annu. Rev. Earth Planet. Sci. 41, 741–766 (2013).

14. P. E. Jardine, G. J. Harrington, J. A. Sessa, J. Daskova, Drivers and constraints on floral latitudinal diversification gradients. J. Biogeogr. 45, 1408-1419 (2018).

15. A. J. Kerkhoff, P. E. Moriarty, M. D. Weiser, The latitudinal species richness gradient in New World woody angiosperms is consistent with the tropical conservatism hypothesis. PNAS USA 111, 8125–8130 (2014).

16. T. Van der Hammen, J. Werner, H. Van Dommelen, Palynological record of the upheaval of the northern andes: a study of the pliocene and lower quaternary of the colombian eastern cordillera and the early evolution of its high-andean biota. Rev. Palaeobot. Palynol. 16, 1–122 (1973).

17. M. J. Donoghue, A phylogenetic perspective on the distribution of plant diversity. PNAS USA 105, 11549-11555 (2008).

18. H. Hooghiemstra, Quaternary and upper-pliocene glaciations and forest development in the tropical andes:

evidence from a long high-resolution pollen record from the sedimentary basin of bogota, colombia. Palaeogeogr. Palaeoclimatol. Palaeoecol. 72, 11–26 (1989).

19. C. Hughes, R. Eastwood, Island radiation on a continental scale: exceptional rates of plant diversification after uplift of the andes. PNAS USA 103, 10334–10339 (2006).

20. B. R. Moore, M. J. Donoghue, Correlates of diversification in the plant clade dipsacales: Geographic movement and evolutionary innovations. Am. Nat. 170, S28-S55 (2007).

21. C. D. Bell, A. Kutschker, M. T. Arroyo, Phylogeny and diversification of valerianaceae (dipsacales) in the southern andes. Mol. Phylogenet. Evol. 63, 724–737 (2012).

22. C. D. Bacon, F. J. Velasquez-Puentes, L. F. Hinojosa, T. Schwartz, B. Oxelman, B. Pfeil, M. T. Arroyo, L. Wanntorp, A. Antonelli, Evolutionary persistence in Gunnera and the contribution of southern plant groups to the tropical andes biodiversity hotspot. PeerJ 6, e4388 (2018).

23. J. Wiens, M. Donoghue, Historical biogeography, ecology and species richness. Trends Ecol. Evol. 19, 639-644 (2004).

24. E. N. Honorio Coronado, K. G. Dexter, R. T. Pennington, J. Chave, S. L. Lewis, M. N. Alexiades, E. Alvarez, A. A. de Oliveira, I. L. Amaral, A. Araujo-Murakami, E. J. M. M. Arets, G. A. Aymard, C. Baraloto, D. Bonal, R. Brienen, C. Ceron, F. Cornejo Valverde, A. Di Fiore, W. Farfan-Rios, T. R. Feldpausch, N. Higuchi, I. Huamantupa-Chuquimaco, S. G. Laurance, W. F. Laurance, G. Lopez-Gonzalez, B. S. Marimon, B. H. Marimon-Junior, A. Monteagudo Mendoza, D. Neill, W. Palacios Cuenca, M. C. Penuela Mora, N. C. A. Pitman, A. Prieto, C. A. Quesada, H. Ramirez Angulo, A. Rudas, A. R. Ruschel, N. Salinas Revilla, R. P. Salomao, A. S. de Andrade, M. R. Silman, W. Spironello, H. ter Steege, J. Terborgh, M. Toledo, L. Valenzuela Gamarra, I. C. G. Vieira, E. Vilanova Torre, V. Vos, O. L. Phillips, Phylogenetic diversity of Amazonian tree communities. Divers. Distrib. 21, 1295-1307 (2015).

25. A. Esquivel-Muelbert, T. R. Baker, K. G. Dexter, S. L. Lewis, H. ter Steege, G. Lopez Gonzalez, A. Monteagudo Mendoza, R. Brienen, T. R. Feldpausch, N. Pitman, *et al.*, Seasonal drought limits tree species across the neotropics. Ecography 40, 618–629 (2017).

26. B. Schrire, M. Lavin, G. Lewis, Plant Diversity and Complexity Patterns: Local, Regional and Global Dimensions, B. H. Friis, I., ed. (Biologiske Skrifter, 2004), vol. 55, pp. 375–422.

27. M. F. Simon, R. Grether, L. P. de Queiroz, C. Skema, R. T. Pennington, C. E. Hughes, Recent assembly of the Cerrado, a neotropical plant diversity hotspot, by in situ evolution of adaptations to fire. PNAS USA 106, 20359-20364 (2009).

28. R. T. Pennington, G. P. Lewis, J. A. Ratter, Neotropical savannas and seasonally dry forests (CRC press, 2006), pp. 17–45.

29. DRYFLOR, Plant diversity patterns in neotropical dry forests and their conservation implications. Science 353, 1383–1387 (2016).

30. T. Kuemmerle, M. Altrichter, G. Baldi, M. Cabido, M. Camino, E. Cuellar, R. L. Cuellar, J. Decarre, S. Dıaz, I. Gasparri, *et al.*, Forest conservation: remember gran chaco. Science 355, 465–465 (2017).

31. K. J. Feeley, J. T. Stroud, Where on earth are the tropics? Front. Biogeogr. 10, 1-7 (2018).

32. A. Oliveira-Filho, NeoTropTree, Flora arborea da Regiao Neotropical: Um banco de dados envolvendo biogeografia, diversidade e conservacao, Belo Horizonte: Universidade Federal de Minas Gerais, http://www.neotroptree.info (2017).

33. E. Burrill, A. Wilson, J. Turner, S. Pugh, J. Menlove, G. Christiansen, B. Conkling, W. David, The Forest Inventory and Analysis Database: database description and user guide version 8.0 for Phase 2. U.S., Department of Agriculture, Forest Service, www.fia.fs.fed.us/library/database-documentation (2018).

34. B. S. Maitner, B. Boyle, N. Casler, R. Condit, J. Donoghue, S. M. Duran, D. Guaderrama, C. E. Hinchliff, P. M. Jørgensen, N. J. Kraft, *et al.*, The BIEN R package: A tool to access the botanical information and ecology network (BIEN) database. Methods Ecol. Evol. 9, 373–379 (2018).

35. P. L. Silva de Miranda, A. T. Oliveira-Filho, R. T. Pennington, D. M. Neves, Baker, T. R., K. G. Dexter, Using tree species inventories to map biomes and assess their climatic overlaps in lowland tropical South America. Global Ecol. Biogeogr. 27, 899-912 (2018).

36. C. Baraloto, O. J. Hardy, C. T. Paine, K. G. Dexter, C. Cruaud, L. T. Dunning, M.-A. Gonzalez, J.-F. Molino, D. Sabatier, V. Savolainen, *et al.*, Using functional traits and phylogenetic trees to examine the assembly of tropical tree communities. J. Ecol. 100, 690–701 (2012).

37. K. Dexter, J. Chave, Evolutionary patterns of range size, abundance and species richness in amazonian angiosperm trees. PeerJ 4, e2402 (2016).

38. K. Katoh, D. M. Standley, Mafft multiple sequence alignment software version 7: improvements in performance and usability. Mol. Biol. Evol. 30, 772–780 (2013).

39. A. Stamatakis, P. Hoover, J. Rougemont, A rapid bootstrap algorithm for the RAxML web servers. Syst. Biol. 57, 758–771 (2008).

40. M. Gastauer, M. Neto, J. A. Alves, Updated angiosperm family tree for analyzing phylogenetic diversity and community structure. Acta Bot. Brasil. 31, 191–198 (2017).

41. S. A. Smith, B. C. OMeara, treePL: divergence time estimation using penalized likelihood for large phylogenies. Bioinformatics 28, 2689–2690 (2012).

42. S. Magallon, S. Gomez-Acevedo, L. L. Sanchez-Reyes, T. Hernandez-Hernandez, A metacalibrated time-tree documents the early rise of flowering plant phylogenetic diversity. New Phytol. 207, 437–453 (2015).

43. J. A. Bryant, C. Lamanna, H. Morlon, A. J. Kerkhoff, B. J. Enquist, J. L. Green, Microbes on mountainsides: contrasting elevational patterns of bacterial and plant diversity. PNAS USA 105, 11505–11511 (2008).

44. C. Tsirogiannis, B. Sandel, Phylomeasures: a package for computing phylogenetic biodiversity measures and their statistical moments. Ecography 39, 709–714 (2016).

45. M. Maechler, P. Rousseeuw, A. Struyf, M. Hubert, K. Hornik, cluster: Cluster Analysis Basics and Extensions (2019). R package version 2.1.0 — For new features, see the 'Changelog' file (in the package source).

46. S. Pavoine, A guide through a family of phylogenetic dissimilarity measures among sites. Oikos 125, 1719–1732 (2016).

47. S. Pavoine, adiv: Analysis of Diversity (2018). R package version 1.2.

48. R. J. Hijmans, S. E. Cameron, J. L. Parra, P. G. Jones, A. Jarvis, Very high resolution interpolated climate surfaces for global land areas. Int. J. Climatol. 25, 1965–1978 (2005).

49. J. Chave, M. Rejou-Mechain, A. Burquez, E. Chidumayo, M. S. Colgan, W. B. Delitti, A. Duque, T. Eid, P. M. Fearnside, R. C. Goodman, *et al.*, Improved allometric models to estimate the aboveground biomass of tropical trees. Global Change Biol. 20, 3177–3190 (2014).

50. T. Duong, ks: Kernel Smoothing (2018). R package version 1.11.3.50.

51. M. D. Caceres, P. Legendre, Associations between species and groups of sites: indices and statistical inference. Ecology 90, 3566–3574 (2009).

52. M. De Caceres, F. Jansen, Indicspecies: relationship between species and groups of sites (2016). R package.

53. J. Larsson, eulerr: area-proportional Euler and Venn diagrams with ellipses (2018). R package version 3.1.

54. A. Dusa, venn: Draw Venn Diagrams (2018). R package version 1.7.

Supplementary Materials References

55. M. L. Bueno, K. G. Dexter, R. T. Pennington, V. Pontara, D. M. Neves, J. A. Ratter, A. T. de Oliveira-Filho, The environmental triangle of the cerrado domain: Ecological factors driving shifts in tree species composition between forests and savannas. J. Ecol. 106, 2109–2120 (2018).

56. K. E. Armstrong, G. N. Stone, J. A. Nicholls, E. Valderrama, A. A. Anderberg, J. Smedmark, L. Gautier, Y. Naciri, R. Milne, J. E. Richardson, Patterns of diversification amongst tropical regions compared: a case study in Sapotaceae. Front. Genet. 5, 362 (2014).

57. F. A. Carvalho, S. S. Renner, Molecular phylogeny, biogeography and an e-monograph of the papaya family (Caricaceae) as an Example of Taxonomy in the Electronic Age (Springer, 2015), pp. 49–81.

58. M. S. Appelhans, P. J. Keßler, E. Smets, S. G. Razafimandimbison, S. B. Janssens, Age and historical biogeography of the pantropically distributed Spathelioideae (Rutaceae, Sapindales). J. Biogeogr. 39, 1235–1250 (2012).

59. T. R. Baker, R. T. Pennington, S. Magallon, E. Gloor, W. F. Laurance, M. Alexiades, E. Alvarez, A. Araujo, E. J. Arets, G. Aymard, *et al.*, Fast demographic traits promote high diversification rates of Amazonian trees. Ecol. Lett. 17, 527–536 (2014).

60. C. D. Bell, M. J. Donoghue, Dating the Dipsacales: comparing models, genes, and evolutionary implications. Am. J. Bot. 92, 284–296 (2005).

61. L. Bardon, C. Sothers, G. T. Prance, P.-J. G. Male, Z. Xi, C. C. Davis, J. Murienne, R. Garcıa-Villacorta, E. Coissac, S. Lavergne, *et al.*, Unraveling the biogeographical history of Chrysobalanaceae from plastid genomes. Am. J. Bot. 103, 1089–1102 (2016).

62. A. Bruneau, M. Mercure, G. P. Lewis, P. S. Herendeen, Phylogenetic patterns and diversification in the Caesalpinioid legumes. Botany 86, 697–718 (2008).

63. L. Cai, Z. Xi, K. Peterson, C. Rushworth, J. Beaulieu, C. C. Davis, Phylogeny of Elatinaceae and the tropical Gondwanan origin of the Centroplacaceae (Malpighiaceae, Elatinaceae) clade. PLoS One 11, e0161881 (2016).

64. S. A. Catalano, J. C. Vilardi, D. Tosto, B. O. Saidman, Molecular phylogeny and diversification history of Prosopis (Fabaceae: Mimosoideae). Biol. J. Linn. Soc. 93, 621–640 (2008).

65. S.-W. Chin, J. Shaw, R. Haberle, J. Wen, D. Potter, Diversification of almonds, peaches, plums and cherries–molecular systematics and biogeographic history of Prunus (Rosaceae). Mol. Phylogenet. Evol. 76, 34–48 (2014).

66. D. M. Crayn, M. Rossetto, D. J. Maynard, Molecular phylogeny and dating reveals an Oligo-Miocene radiation of dry-adapted shrubs (former Tremandraceae) from rainforest tree progenitors (Elaeocarpaceae) in Australia. Am. J. Bot. 93, 1328–1342 (2006).

67. L. A. Eserman, G. P. Tiley, R. L. Jarret, J. H. Leebens-Mack, R. E. Miller, Phylogenetics and diversification of morning glories (tribe Ipomoeeae, Convolvulaceae) based on whole plastome sequences. Am. J. Bot. 101, 92–103 (2014).

68. E. M. Gardner, P. Sarraf, E. W. Williams, N. J. Zerega, Phylogeny and biogeography of Maclura (Moraceae) and the origin of an anachronistic fruit. Mol. Phylogenet. Evol. 117, 49–59 (2017).

69. A. Harris, Q.-Y. Xiang, D. T. Thomas, Phylogeny, origin, and biogeographic history of Aesculus L.(Sapindales)–an update from combined analysis of DNA sequences, morphology, and fossils. Taxon 58,

108–126 (2009).

70. J. Heckenhauer, R. Samuel, P. S. Ashton, B. Turner, M. H. Barfuss, T.-S. Jang, E. M. Temsch, J. Mccann, K. A. Salim, A. Attanayake, *et al.*, Phylogenetic analyses of plastid DNA suggest a different interpretation of morphological evolution than those used as the basis for previous classifications of Dipterocarpaceae (Malvales). Bot. J. Linn. Soc. 185, 1–26 (2017).

71. L. Li, J. Li, J. G. Rohwer, H. van der Werff, Z.-H. Wang, H.-W. Li, Molecular phylogenetic analysis of the Persea group (Lauraceae) and its biogeographic implications on the evolution of tropical and subtropical Amphi-Pacific disjunctions. Am. J. Bot. 98, 1520–1536 (2011).

72. A. R. Mast, E. F. Milton, E. H. Jones, R. M. Barker, W. R. Barker, P. H. Weston, Time calibrated phylogeny of the woody Australian genus Hakea (Proteaceae) supports multiple origins of insect-pollination among bird-pollinated ancestors. Am. J. Bot. 99, 472–487 (2012).

73. I. Michalak, L.-B. Zhang, S. S. Renner, Trans-Atlantic, trans-Pacific and trans-Indian Ocean dispersal in the small Gondwanan Laurales family Hernandiaceae. J. Biogeogr. 37, 1214-1226 (2010).

74. M. L. Milner, P. H. Weston, M. Rossetto, M. D. Crisp, Biogeography of the Gondwanan genus Lomatia (Proteaceae): vicariance at continental and intercontinental scales. J. Biogeogr. 42, 2440–2451 (2015).

75. L. C. Moyle, M. S. Olson, P. Tiffin, Patterns of reproductive isolation in three Angiosperm genera. Evolution 58, 1195–1208 (2004).

76. A. N. Muellner, T. D. Pennington, A. V. Koecke, S. S. Renner, Biogeography of Cedrela (Meliaceae, Sapindales) in central and South America. Am. J. Bot. 97, 511–518 (2010).

77. S. S. Renner, L.-B. Zhang, J. Murata, A chloroplast phylogeny of Arisaema (Araceae) illustrates Tertiary floristic links between Asia, North America, and East Africa. Am. J. Bot. 91, 881–888 (2004).

78. S. S. Renner, G. W. Grimm, G. M. Schneeweiss, T. F. Stuessy, R. E. Ricklefs, Rooting and dating maples (Acer) with an uncorrelated-rates molecular clock: implications for North American/Asian disjunctions. Syst. Biol. 57, 795–808 (2008).

79. S. S. Renner, J. S. Strijk, D. Strasberg, C. Thebaud, Biogeography of the Monimiaceae (Laurales): a role for East Gondwana and long-distance dispersal, but not West Gondwana. J. Biogeogr. 37, 1227–1238 (2010).

80. A. Rockinger, A. S. Flores, S. S. Renner, Clock-dated phylogeny for 48% of the 700 species of Crotalaria (Fabaceae–Papilionoideae) resolves sections worldwide and implies conserved flower and leaf traits throughout its pantropical range. BMC Evol. Biol. 17, 61 (2017).

81. J. G. Rodrigues, J. A. Lombardi, M. B. Lovato, Phylogeny of Cissus (Vitaceae) focusing on South American species. Taxon 63, 287–298 (2014).

82. H. Sauquet, P. H. Weston, C. L. Anderson, N. P. Barker, D. J. Cantrill, A. R. Mast, V. Savolainen, Contrasted patterns of hyperdiversification in Mediterranean hotspots. PNAS USA 106, 221–225 (2009).

83. J. V. Schneider, G. Zizka, Phylogeny, taxonomy and biogeography of Neotropical Quiinoideae (Ochnaceae sl). Taxon 66, 855–867 (2017).

84. O. Schwery, R. E. Onstein, Y. Bouchenak-Khelladi, Y. Xing, R. J. Carter, H. P. Linder, As old as the mountains: the radiations of the Ericaceae. New Phytol. 207, 355–367 (2015).

85. J. E. Smedmark, A. A. Anderberg, Boreotropical migration explains hybridization between geographically distant lineages in the pantropical clade Sideroxyleae (Sapotaceae). Am. J. Bot. 94, 1491–1505 (2007).

86. J. E. Smedmark, T. Eriksson, B. Bremer, Divergence time uncertainty and historical bio geography reconstruction–an example from Urophylleae (Rubiaceae). J. Biogeogr. 37, 2260–2274 (2010).

87. M. H. Terra-Araujo, A. D. de Faria, A. Vicentini, S. Nylinder, U. Swenson, Species tree phylogeny and

biogeography of the Neotropical genus Pradosia (Sapotaceae, Chrysophylloideae). Mol. Phylogenet. Evol. 87, 1–13 (2015).

88. A. H. Thornhill, S. Y. Ho, C. Kulheim, M. D. Crisp, Interpreting the modern distribution of Myrtaceae using a dated molecular phylogeny. Mol. Phylogenet. Evol. 93, 29–43 (2015).

89. N. Thomas, J. J. Bruhl, A. Ford, P. H. Weston, Molecular dating of Winteraceae reveals a complex biogeographical history involving both ancient Gondwanan vicariance and long distance dispersal. J. Biogeogr. 41, 894–904 (2014).

90. E. A. Tripp, L. A. McDade, A rich fossil record yields calibrated phylogeny for Acanthaceae (Lamiales) and evidence for marked biases in timing and directionality of intercontinental disjunctions. Syst. Biol. 63, 660–684 (2014).

91. P. C. van Welzen, K. Pruesapan, I. R. Telford, J. J. Bruhl, Historical biogeography of Breynia (Phyllanthaceae): what caused speciation? J. Biogeogr. 42, 1493–1502 (2015).

92. T. N. Vasconcelos, C. E. Proenca, B. Ahmad, D. S. Aguilar, R. Aguilar, B. S. Amorim, K. Campbell, I. R. Costa, P. S. De-Carvalho, J. E. Faria, *et al.*, Myrteae phylogeny, calibration, biogeography and diversification patterns: increased understanding in the most species rich tribe of Myrtaceae. Mol. Phylogenet. Evol. 109, 113–137 (2017).

93. Q.-Y. J. Xiang, D. T. Thomas, Q. P. Xiang, Resolving and dating the phylogeny of Cornales–effects of taxon sampling, data partitions, and fossil calibrations. Mol. Phylogenet. Evol. 59, 123–138 (2011).

94. M.-Q. Yang, D.-Z. Li, J. Wen, T.-S. Yi, Phylogeny and biogeography of the amphi-pacific genus Aphananthe. PLoS One 12, e0171405 (2017).

95. T. Yang, L.-M. Lu, W. Wang, J.-H. Li, S. R. Manchester, J. Wen, Z.-D. Chen, Boreotropical range expansion and long-distance dispersal explain two amphi-pacific tropical disjunctions in Sabiaceae. Mol. Phylogenet. Evol. 124, 181–191 (2018).

Supplementary Materials

Freezing and water availability structure the evolutionary diversity of trees across the Americas

R. A. Segovia (ricardo.segovia@)ed.ac.uk), R. T. Pennington, T. R. Baker, F. Coelho de Souza, D. M. Neves, C. C. Davis, J. J. Armesto, A. T. Olivera-Filho & K. G. Dexter

Table S1 Indicator clades for *K*=2 groups. Specificity, fidelity and indicator statistic (stat) of internal nodes associated for the top nodes with the highest indicator statistic. Clades names are based on their taxonomic composition.

Cluster	Clades	Specificity	Fidelity	stat
Tropical	Gentianales	0.9419787	0.9885234	0.965
	Fabaceae, tribe Mimoseae	0.9339992	0.9680896	0.951
	Fabaceae, subfam. Caesalpinioideae	0.9366716	0.9588523	0.948
	Euphorbiaceae and Peraceae	0.9639994	0.9303009	0.947
	Putranjivaceae, Phyllanthaceae, Picrodendraceae and Ochnaceae	0.9792854	0.9143457	0.946
Extratropical	Fagales	0.886934	0.85745	0.872
	Fagales *plus* Cucurbitales	0.875902	0.85745	0.867
	Ulmaceae	0.987098	0.62106	0.783
	Sapindaceae, subfam. Hippocastanoideae	0.943887	0.612106	0.76

Table S2 Indicator clades for *K*=4 groups. Specificity, fidelity and indicator statistic (stat) of internal nodes associated for the top nodes with the highest indicator statistic. Clade names are based on their taxonomic composition.

Cluster	Clades	Specificity	Fidelity	stat
Tropical Moist	*Xylopia*, *Fusaea* and *Duguetia* (Annonaceae)	0.849969	0.700198	0.771
	Couepia and *Hirtella* (Chrysobalanaceae)	0.757572	0.711399	0.734
	Burseraceae, tribes Protieae and Canarieae	0.794147	0.678014	0.734
	Ochnaceae	0.746723	0.706128	0.726
	Myristicaceae	0.851393	0.591917	0.71
	Calophyllaceae	0.7516	0.668351	0.709
Tropical Dry	Bignoniaceae, tribes Bignonieae and Tecomeae	0.778906	0.117489	0.303
	Lasiocarpus and *Ptilochaeta* (Malpighiaceae)	0.853331	0.072646	0.249
	Cactaceae, tribe Trichocereeae	0.864368	0.047982	0.204
Southern Extratropical	*Prosopis*, *Piptadeniopsis* (Fabaceae, tribe Mimoseae)	0.733744	0.402597	0.544
	Fabaceae, tribe Caesalpinieae	0.670481	0.301587	0.45
	Vallea and *Aristotelia* (Elaeocarpaceae)	0.854485	0.223665	0.437
	Cactaceae, tribes Pachycereeae and Notocactea	0.732889	0.248196	0.426
	Scrophulariaceae	0.597146	0.30303	0.425
Northern Extratropical	Sapindaceae, subfam. Hippocastanoideae	0.83986	0.69118	0.762
	Ulmaceae	0.75668	0.70459	0.73
	Oleaceae, tribe Oleeae	0.80497	0.62332	0.708
	Juglandaceae	0.76212	0.53677	0.64

Table S3. Affiliation of principal vegetation formations in the tropics with the two main tropical groups from the K=4 clustering analysis. Vegetation formations were taken from the NeoTropTree dataset, which categorises formations first based on physiognomy (savanna vs. forest) and then segregates the forests based on phenology. Following (*35*) and (*54*), we consider deciduous tropical forests to represent the tropical dry forest biome, while semideciduous forests are more related floristically to the tropical moist forest biome. Semideciduous forests share many tree species with evergreen forests and relatively few with more fully deciduous forests (*35,54*). We further divided the savannas based on geography, as our analyses showed evident differences in group affiliation between savannas in the Cerrado Domain of Brazil versus those further north (i.e. Llanos of Venezuela and Colombia and those in Central America).

	Tropical dry	Tropical moist
Evergreen Forests	15% (501)	85% (2948)
Semideciduous Forests	10% (167)	90% (1530)
Deciduous Forests	75% (868)	25% (285)
Southern Savannas (Cerrado)	8% (56)	92% (657)
Northern Savannas	54% (65)	46% (56)

Table S4. Stem ages for genera nodes used to callibrate the phylogenetic tree, and the reference of their source. References (*55-94*).

	Genus	stem age (Myr)	Reference
1	*Acer*	60	Renner et al 2008, Systematic Biology
2	*Acioa*	19.1	Bardon et al 2016, American Journal of Botany
3	*Aesculus*	65	Harris et al 2009, Taxon
4	*Anaxagorea*	90.44	Baker et al. 2014, Ecology Letters (and references herein)
5	*Andira*	17.51	Baker et al. 2014, Ecology Letters (and references herein)
6	*Antiaris*	34	Gardner et al 2017, Molecular Phylogenetics and Evolution
7	*Aphananthe*	71.5	Yang et al 2007, PLOS ONE
8	*Aphanocalyx*	46	Bruneau et al 2008, Botany
9	*Artocarpus*	51	Rockinger et al 2017, BMC Evolutionary Biology
10	*Atuna*	20.5	Bardon et al 2016, American Journal of Botany
11	*Avicennia*	70.09	Tripp & McDade 2014, Systematic Biology
12	*Bagassa*	67	Gardner et al 2017, Molecular Phylogenetics and Evolution
13	*Bocageopsis*	5.98	Baker et al. 2014, Ecology Letters (and references herein)
14	*Brosimum*	48	Baker et al. 2014, Ecology Letters (and references herein)
15	*Caesalpinia*	48.3	Bruneau et al 2008, Botany
16	*Carapa*	29.5	Baker et al. 2014, Ecology Letters (and references herein)
17	*Cassia*	45	Bruneau et al 2008, Botany
18	*Castilla*	22	Baker et al. 2014, Ecology Letters (and references herein)
19	*Cecropia*	44	Baker et al. 2014, Ecology Letters (and references herein)
20	*Cedrela ///toona*	48.4	Muellner et al 2010, American Journal of Botany
21	*Cedrelopsis*	18.94	Appelhans et al 2012, Journal of Biogeography
22	*Centroplacus*	69	Cai et al 2016, PlosONE
23	*Cercis*	47.3	Bruneau et al 2008, Botany
24	*Chrysobalanus*	24.2	Bardon et al 2016, American Journal of Botany
25	*Cissus*	67.99	Rodrigues et al 2014, Taxon
26	*Clarisia*	70	Gardner et al 2017, Molecular Phylogenetics and Evolution
27	*Coceveiba*	72	Baker et al. 2014, Ecology Letters (and references herein)
28	*Cornus*	74.03	Xiang et al 2011, Molecular Phylogenetics and Evolution
29	*Couepia*	21.6	Bardon et al 2016, American Journal of Botany
30	*Crudia*	45	Bruneau et al 2008, Botany
31	*Cylicomorpha*	35.5	Antunes & Renner 2012, Molecular Phylogenetics and Evolution
32	*Cynometra*	12.93	Baker et al. 2014, Ecology Letters (and references herein)
33	*Dacryodes*	38	Baker et al. 2014, Ecology Letters (and references herein)
34	*Dactyladenia*	15.9	Bardon et al 2016, American Journal of Botany
35	*Dactylocladus*	39	Moyle 2004, Evolution
36	*Dialium*	10.9	Baker et al. 2014, Ecology Letters (and references herein)
37	*Dicymbe*	12	Baker et al. 2014, Ecology Letters (and references herein)
38	*Diplotropis*	20.27	Baker et al. 2014, Ecology Letters (and references herein)
39	*Dipterix*	26.44	Baker et al. 2014, Ecology Letters (and references herein)
40	*Dipterocarpus*	47.7	Heckenhauer et al 2017, Botanical Journal of the Linnean Society
41	*Drimys*	56.76	Thomas et al 2014, Journal of Biogeography
42	*Dryobalanops*	43.3	Heckenhauer et al 2017, Botanical Journal of the Linnean Society
43	*Duguetia*	30.64	Baker et al. 2014, Ecology Letters (and references herein)
44	*Dycorynia*	10.9	Baker et al. 2014, Ecology Letters (and references herein)
45	*Embothrium*	39.3	Sauquet et al 2009, PNAS
46	*Eperua*	12.32	Baker et al. 2014, Ecology Letters (and references herein)
47	*Ficus*	58	Gardner et al 2017, Molecular Phylogenetics and Evolution
48	*Froesia*	39.4	Schneider & Zizka 2017, Taxon
49	*Fusaea*	30.64	Baker et al. 2014, Ecology Letters (and references herein)
50	*Glochidion*	31.51	van Welzen et al 2015, Journal of Biogeography
51	*Glycosmis*	32.54	Appelhans et al 2012, Journal of Biogeography
52	*Guarea*	14.8	Baker et al. 2014, Ecology Letters (and references herein)
53	*Guatteria*	55.83	Baker et al. 2014, Ecology Letters (and references herein)
54	*Gyrocarpus*	72	Michalak et al 2010. Journal of Biogeography
55	*Hakea*	12.8	Mast et al 2012, American Journal of Botany

Genus	stem age (Myr)	Reference
56 *Harrisoinia*	57.99	Appelhans et al 2012, Journal of Biogeography
57 *Helicostylis*	28	Baker et al. 2014, Ecology Letters (and references herein)
58 *Hennecartia*	15.58	Renner et al 2010, Journal of Biogeography
59 *Hernandia*	76	Michalak et al 2010. Journal of Biogeography
60 *Hevea*	85	Baker et al. 2014, Ecology Letters (and references herein)
61 *Hopea*	21.6	Heckenhauer et al 2017, Botanical Journal of the Linnean Society
62 *Hymenaea*	24	Bruneau et al 2008, Botany
63 *Inga*	10	Baker et al. 2014, Ecology Letters (and references herein)
64 *Ipomoea*	34.97	Eserman et al 2014, American Journal of Botany
65 *Iryanthera*	19	Baker et al. 2014, Ecology Letters (and references herein)
66 *Jacaratia*	27.5	Antunes & Renner 2012, Molecular Phylogenetics and Evolution
67 *Lacunaria*	20.3	Schneider & Zizka 2017, Taxon
68 *Lomatia*	70.8	Milner et al 2015, Journal of Biogeography
69 *Lonchocarpus*	15.07	Baker et al. 2014, Ecology Letters (and references herein)
70 *Lonicera*	43.37	Bell & Donoghue 2005, American Journal of Botany
71 *Maclura*	85	Gardner et al 2017, Molecular Phylogenetics and Evolution
72 *Macrolobium*	32	Baker et al. 2014, Ecology Letters (and references herein)
73 *Magnistipula*	19	Bardon et al 2016, American Journal of Botany
74 *Malmea*	19.99	Baker et al. 2014, Ecology Letters (and references herein)
75 *Manilkara*	32	Armstrong et al 2014, Frontiers in Genetics
76 *Maranthes*	20.5	Bardon et al 2016, American Journal of Botany
77 *Meliosma*	67.34	Yang et al 2018, Molecular Phylogenetics and Evolution
78 *Mimusops*	35	Armstrong et al 2014, Frontiers in Genetics
79 *Mouriri*	90	Renner 2004, American Journal of Botany
80 *Myrtae tribe*	58.96	Vasconcelos et al 2017, Molecular Phylogentics and Evolution
81 *Neocarya*	25.6	Bardon et al 2016, American Journal of Botany
82 *Ormosia*	40.62	Baker et al. 2014, Ecology Letters (and references herein)
83 *Otoba*	17	Baker et al. 2014, Ecology Letters (and references herein)
84 *Parashora*	22.9	Heckenhauer et al 2017, Botanical Journal of the Linnean Society
85 *Parkia*	45.5	Baker et al. 2014, Ecology Letters (and references herein)
86 *Peltogyne*	28.8	Baker et al. 2014, Ecology Letters (and references herein)
87 *Persea*	55.3	Li et al 2018, American Journal of Botany
88 *Peumus*	55.66	Renner et al 2010, Journal of Biogeography
89 *Poecilanthe*	40.99	Baker et al. 2014, Ecology Letters (and references herein)
90 *Poulsenia*	22	Baker et al. 2014, Ecology Letters (and references herein)
91 *Pourouma*	44	Baker et al. 2014, Ecology Letters (and references herein)
92 *Pradosia*	47.5	Terra-Araujo et al 2015, Molecular Phylogenetics and Evolution
93 *Prosopis SA*	28.96	Catalano et al 2008, Botanical Journal of the Linnean Society
94 *Protium*	52.5	Baker et al. 2014, Ecology Letters (and references herein)
95 *Prunus*	60.7	Chin et al 2014, Molecular Phylogentics and Evolution
96 *Pseudolmedia*	36	Baker et al. 2014, Ecology Letters (and references herein)
97 *Pseudowintera*	45.18	Thomas et al 2014, Journal of Biogeography
98 *Pseudoxandra*	15.09	Baker et al. 2014, Ecology Letters (and references herein)
99 *Pterocarpus*	16.66	Baker et al. 2014, Ecology Letters (and references herein)
100 *Quiina*	29	Schneider & Zizka 2017, Taxon
101 *Rhododendron*	58	Schwery et al 2015, New Phytologist
102 *Richea*	22.31	Schwery et al 2015, New Phytologist
103 *Sambucus*	45.49	Bell & Donoghue 2005, American Journal of Botany
104 *Sideroxylon*	74	Smedmark & Anderberg 2007, American Journal of Botany
105 *Slonaea*	79	Crayn et al 2006. American Journal of Botany
106 *Sorocea*	59	Baker et al. 2014, Ecology Letters (and references herein)
107 *Spathelia*	19.21	Appelhans et al 2012, Journal of Biogeography
108 *Swartzia*	45.96	Baker et al. 2014, Ecology Letters (and references herein)
109 *Tachigali*	4.65	Baker et al. 2014, Ecology Letters (and references herein)
110 *Tasmania*	69.98	Thomas et al 2014, Journal of Biogeography
111 *Tepualia*	24.9	Thornhill et al 2015, Molecular Phylogenetics and Evolution
112 *Theobroma*	11.6	Richardson et al 2015, Front. Ecol. Evol.
113 *Unonopsis*	7.94	Baker et al. 2014, Ecology Letters (and references herein)
114 *Urophyllum*	27.1	Smedmark et al 2010, Journal of Biogeography
115 *Vallea*	48	Crayn et al 2006. American Journal of Botany
116 *Vateria*	15.4	Heckenhauer et al 2017, Botanical Journal of the Linnean Society
117 *Vatica*	18.3	Heckenhauer et al 2017, Botanical Journal of the Linnean Society
118 *Viburum*	71.18	Bell & Donoghue 2005, American Journal of Botany
119 *Virola*	17	Baker et al. 2014, Ecology Letters (and references herein)
120 *Vitellariopsis*	26	Armstrong et al 2014, Frontiers in Genetics
121 *Vouacapoua*	48.69	Baker et al. 2014, Ecology Letters (and references herein)
122 *Xylopia*	49.98	Baker et al. 2014, Ecology Letters (and references herein)
123 *Zygia*	17.82	Baker et al. 2014, Ecology Letters (and references herein)

Fig. S1. Match between tropics vs. extratropics groups from K=2 clustering and eight delimitations of the tropics following Feeley & Stroud [2018] *(31):* C1) all areas that occur between 23.4°S and 23.4°N; C2) all areas with a net positive energy balance; C3) all areas where mean annual temperature does not vary with latitude; C4) all areas where temperatures do not go below freezing in a typical year; C5) all areas where the mean monthly temperature is never less than 18°C; C6) all areas where the mean annual "biotemperature" ≥ 24°C; C7) all areas where the annual range of temperature is less than the average daily temperature range; C8) all areas where precipitation seasonality exceeds temperature seasonality.

Fig. S2. Selection of number of clusters. A) Elbow criterion, explained variance from clustering as a function of number of groups; B) Silhouette criterion, average silhouette width for each site as a function of number of groups.

65   **Fig. S3 Shared versus unique Phylogenetic Diversity for *K*=2 and *K*=3 clustering analyses.**
Euler Diagrams showing the amount of unique phylogenetic diversity in each cluster and the
phylogenetic diversity shared between clusters (in millions of years). A) *K*=2 clustering and B)
K=3 clustering.

70   **Fig. S4. Clustering *K* = 3.** A) Location of the 9937 angiosperm tree assemblages in three
     evolutionary groups; B) Ordination of tree assemblages based on evolutionary lineage
     composition; C) Maximum Climatological Water Deficit (CWD) versus minimum temperature
     of the coldest month. Lines represent the 95th quantile of the density of points for each group. In
     each panel, symbol type indicates major vegetation type.

Fig. S5 Clustering validation *K*= 2. A) Silhouette width for each assemblage in our dataset. The colors represent the clusters and the red dashed line the average silhouette width. B) shows the distribution of sites with silhouette > 0.2, which indicates that they are strongly associated with their given cluster. C) shows the geographic distribution of sites with silhouette width ≤ 0.2. D) shows sites with silhouette width > 0.2 in a latitude-elevation plot and E) shows sites with silhouette width ≤ 0.2 in a latitude-elevation plot.

85 **Fig. S6. Clustering validation *K*= 3.** A) Silhouette width for each assemblage in our dataset. The colors represent the clusters and the red dashed line the average silhouette width. B) shows the distribution of sites with silhouette width bigger than 0.2, which indicates that they are strongly associated with their given cluster. C) shows the geographic distribution of sites with silhouette width < 0.2. D) shows sites with silhouette width > 0.2 in a latitude-elevation plot and
90 E) shows sites with silhouette width < 0.2 in a latitude-elevation plot.

Fig. S7. Clustering validation *K*= 4. A) Silhouette width for each assemblage in our dataset. The colors represent the clusters and the red dashed line the average silhouette width. B) shows the distribution of sites with silhouette width bigger than 0.2, which indicates that they are strongly associated with their givencluster. C) shows the geographic distribution of sites with silhouette width < 0.2. D) shows sites with silhouette width > 0.2 in a latitude-elevation plot and E) shows sites with silhouette width < 0.2 in a latitude-elevation plot.