Commensal bacteria differentially shape the nutritional requirements of

Drosophila during juvenile growth

Jessika Consuegra[1*], Théodore Grenier[1*], Patrice Baa-Puyoulet[2], Isabelle Rahioui[2],

Houssam Akherraz[1], Hugo Gervais[1], Nicolas Parisot[2], Pedro da Silva[2], Hubert

Charles[2], Federica Calevro[2], François Leulier[1§].

1. Institut de Génomique Fonctionnelle de Lyon, Univ Lyon, CNRS UMR5242, Ecole

Normale Supérieure de Lyon, Université Claude Bernard Lyon 1. 46, allée d'Italie F-

69364 Lyon, France.

2. Univ Lyon, INSA-Lyon, INRA, BF2I, UMR0203, F-69621, Villeurbanne, France

* These authors contribute equally to this work

§ Corresponding author. François Leulier, francois.leulier@ens-lyon.fr

Abstract

The interplay between nutritional and microbial environments is one of the decisive environmental inputs that determine juvenile growth trajectory. Nutritional deficiencies contribute to developmental delays, and an immature gut microbiota is a hallmark of pathologies related to childhood undernutrition. However, how commensal bacteria modulate the impact of nutrition on juvenile growth remains elusive. Here, using gnotobiotic *Drosophila melanogaster* larvae independently associated with two major commensal strains, *Acetobacter pomorum*^{WJL} (Ap^{WJL}) and *Lactobacillus plantarum*^{NC8} (Lp^{NC8}), we performed a large-scale, systematic nutritional screen based on larval growth in 40 different and precisely controlled nutritional environments. We combined these results with *in silico* metabolic network reconstruction to define the biosynthetic capacities of Drosophila germ-free larvae and the two commensals. We first establish that Ap^{WJL} or Lp^{NC8} differentially fulfills the nutritional requirements of the ex-GF larva and parsed such difference down to individual amino acids, vitamins, other micronutrients and trace metals. We found that Drosophila commensal bacteria not only fortify the host's diet with essential nutrients but in specific instances, they functionally compensate for host auxotrophies even if the bacteria fail to synthetize the missing nutrient. Our systematic work reveals that beyond the molecular dialogue engaged between the host and its commensal partners, Drosophila engage into an integrated nutritional network with its facultative commensals centered around the sparing and utilization of nutrients. We thus uncover a novel facet of the facultative nutritional mutualism engaged between Drosophila and its commensal bacteria, which allows the juvenile host to better cope with changes in nutrients availability during the critical phase of post-natal growth; hence ensuring optimal host fitness.

Introduction

Nutrition is arguably the major environmental factor that determines to what extent an organism can realize its genetically encoded growth potential [1]. The attributes of nutrition are defined by the quantity [2], quality [3] and bioavailability [4] of different nutrients in the diet. Nutrients are classified as non-essential or essential [3] based on the organism's biosynthetic capacities. Diets deficient in essential nutrients cause important growth and maturation delays or even growth arrest, or "stunting" characterized by low height-for-age score [5]. In addition, some nutrients are conditionally-essential (or semi-essential). These nutrients can be synthesized by the organism, but insufficiently so under certain metabolically demanding conditions such as juvenile growth. Therefore, these semi-essential nutrients must also be retrieved from diets. Deficient consumption of conditionally-essential nutrients can also be detrimental for growth [3].

The intricate relationship between nutrition and growth is transformed by the microbial environment. In a classic human twin study, Smith *et al* unequivocally demonstrated that the gut microbiota composition of the juvenile subject suffering from stunting is significantly different from that of the healthy twin. When the faecal microbiota from the discordant twins were transplanted into genetically identical germ-free (GF) mice, the recipients of the microbiota from the stunted human twin performed poorly in terms of growth gain and weight recovery compared to the recipients of the microbiota of the healthy twin [6]. Furthermore, functional analyses of gut microbiota from children experiencing strong acute malnutrition showed significant under-representation in pathways of amino acid biosynthesis and uptake; carbohydrate utilization and B-vitamin metabolism [7]. Diets supplemented with nutrients favoring the growth of bacteria enriched in these underrepresented pathways increase plasma

biomarkers and levels of mediators of growth, bone formation, neurodevelopment, and

immune function in children with moderate acute malnutrition [7]. These studies clearly

show that the microbial environment strongly impacts how the organism responds to

changes in its nutritional environment, and vice versa.

Diverse animal models are employed to decipher the physiological, ecological,

genetic and molecular mechanisms underpinning host/microbiota/diet interactions.

Among them, *Drosophila melanogaster* (Dm) is frequently chosen to study the impact

of the nutritional environment on growth and development thanks to its short growth

period as well as easy and cost-effective rearing conditions. During the juvenile phase

of the Drosophila life cycle, the larva feeds constantly and increase its body mass ~200

times until entry into metamorphosis [8]. However, the pace and duration of larval

growth can be altered by the nutritional context and the microbial environment [9–11].

Like other animals, Drosophila live in constant association with commensal microbes,

including bacteria and yeast [12]. The impact of the microbial environment can be

systematically assessed by generating gnotobiotic flies associated with a defined set

of commensals [13–15]. Lab-reared flies typically carry bacterial strains from only four

to eight species. The microbiota from wild flies are more complex. Nevertheless, they

are usually dominated by members from *Acetobacteraceae* and *Lactobacilli* [16–22].

Most bacterial strains from these dominant species are easy to culture in the lab, and

some have even been genetically engineered for functional studies of host-microbes

interactions [23–25]. These fly commensals are facultative symbionts that are

constantly horizontally acquired [26–28]. Even though recent experimental evidence

show that wild bacterial isolates can persistently colonize the adult crop [22,29], the

larval gut commensals are in fact transient; they constantly shuttle between the larval

gut and the food substrate to establish a mutualistic cycle with the host [30,31]. Thus,

the Drosophila larval growth is an attractive model to study the strong interplay

between the Drosophila commensals and the nutritional environment.

We and others have previously shown that GF larvae raised in nutritional

challenges show important developmental delays, and a single gut commensal strain

can accelerate Drosophila development during nutritional challenge [20,25].

Specifically, *Acetobacter pomorum*[WJL] (Ap[WJL]) modulates developmental rate and final

body size through the insulin/insulin-like growth factor signaling (IIS) pathway, and the

intact acetate production machinery is critical [25]. *Lactobacillus plantarum*[WJL] or *L.*

plantarum[NC8] (Lp[NC8]) promote host juvenile growth and maturation partly through

enhanced expression of intestinal peptidases upon sensing of bacterial cell wall

components by *Drosophila* enterocytes [20,23,32]. Interestingly, the growth-promoting

effect of these bacteria was only observed under nutrient scarcity, suggesting that

besides the molecular dialogue engaged between the commensal bacteria and their

host, bacteria-mediated growth promotion on low diets also includes some nutritional

compensation, as already reported for bacteria-derived thiamine provision to the host

[33]. However, how the presence of such bacteria specifically alters the host's

nutritional environment and satisfies the host's nutritional requirements remains

unexplored. To do so, we systematically assessed the microbial contribution to larval

growth in 40 different and strictly controlled nutritional contexts, and we conducted this

large-scale screen based on chemically defined holidic diets (HDs).

Holidic diets comprise a mixture of pure chemical ingredients that satisfy the

different physiological requirements of the Drosophila host [34,35]. By altering the

concentration of each or a combination of ingredient, one can exactly tailor the

experiments by generating specific nutrient deficiencies or excess [36]. The first

development of HD supporting the growth of Drosophila can be traced back to the 40s

[37], and have been used to assess the direct impact of the nutritional environment on

axenic larvae in the 1950s [38,39]. In this study, we adopted a recently developed

FLYAA HD where the amino acids concentrations are calculated so that they match

the amino acid ratios found in the translated exome of the fly [40]. The FLYAA HD is

optimal for both fecundity and lifespan of adults, and was recently used to investigate

the links between nutrition and lifespan [40–43], fecundity [40–42,44], food choice

behavior [45], nutrient sensing [46], growth and maturation [33,40–42,47–49]. Exome-

based HD can efficiently support larval growth albeit not to the optimal growth and

maturation rate obtained with classic oligidic laboratory fly food [34]. Using the

chemically defined HD, we aimed to deconstruct in a systematic manner the microbial

contribution to the host's nutritional requirements down to individual nutrients.

To do so, we first needed to establish the biosynthetic capacities of GF larvae

and two major commensals of Drosophila: Ap^{WJL} and Lp^{NC8} on HD. We further

complemented the *in vivo* study with *in silico* metabolic network reconstruction based

on the genome sequences of *D. melanogaster*, Ap^{WJL} and Lp^{NC8}. In recent years,

metabolic approaches based on genome-driven network reconstructions have been

applied to predict the potential metabolic dependencies and metabolic exchanges

between hosts and associated microbes [50–55]. The mutualistic association between

the pea aphid *Acyrthosiphon pisum* and its obligate intracellular symbiont *Buchnera*

aphidicola was the first symbiotic association for which genomic information were

available on both partners, and is a case study for a comprehensive survey of

integrated host-symbiont metabolic interactions. In this model, decades of nutritional

experiments using HDs and aposymbiotic aphids were reinterpreted in the light of

newly available genomic data, thus changing the traditional paradigm that proposed a

clear separation between the pathways of the host and its symbionts and revealing a

particularly integrated metabolic network between the insect and its obligate

endosymbionts [56,57]. Indeed, the metabolic coevolution between the pea aphid and

Buchnera aphidicola led to the emergence of shared metabolic pathways, such as that

for the biosynthesis of branched-chain [58] and aromatic amino acids [59,60], in which

the enzymes are encoded by both the host and the bacterial genomes. Such highly

integrated metabolic pathways have been subsequently discovered in other obligate

mutualistic associations and can involve an eukaryotic host and all its bacterial

symbionts in the case of multi-partner symbioses [61]. The availability of the genomes

of *D. melanogaster* and of its two major commensal bacterial species provides the

unique opportunity to extend these studies from obligate mutualistic bacterial

endosymbioses to facultative host-commensal bacteria mutualisms. Here we report

that association of GF larvae with Ap^{WJL} or Lp^{NC8} modifies the nutritional requirements

of ex-GF larvae in a specific manner each. We show that Ap^{WJL} and Lp^{NC8} not only

modify the nutritional environment of their host by fortifying diets with essential

nutrients but also functionally compensate for host auxotrophies despite not

synthetizing the missing nutrient.

Results and Discussion

Metabolic network reconstruction of the host (*D. melanogaster*) and its two commensals, Ap^{WJL} and Lp^{NC8}, was automatically generated using the CycADS pipeline [62]. The resulting BioCyc metabolism databases are available at http://artsymbiocyc.cycadsys.org for visualization and analysis purposes. We generated the enriched functional annotations of all the predicted proteins from the complete genomes of *D. melanogaster* (Dm, RefSeq GCF_000001215.4 release 6), *A. pomorum* strain DM001 (Ap^{WJL}, accession NCBI-Taxonomy 945681 version 1.0.1) and *L. plantarum* subsp. *plantarum* NC8 (Lp^{NC8}, NCBI Bioproject PRJNA67175). From the genomic analyses, we inferred all pathways allowing production of the organic compounds that are present in the exome-based FLYAA holidic diet (HD) developed by Piper et al [40]: fly's essential and non-essential amino acids (EAA^{Fly} (n=10) and NEAA^{Fly} (n=10)), B-vitamins (n=7), cholesterol (n=1), and Nucleic Acids and Lipid precursors (NAL, n=4).

***Drosophila melanogaster* biosynthetic capabilities inferred from *in silico* metabolic network reconstruction**

Although a BioCyc metabolic reconstruction of *D. melanogaster* is already publicly available (https://biocyc.org/FLY), we have reconstructed an improved and more comprehensive BioCyc database using CycADS [62]. This metabolic reconstruction comprises 22,189 protein-encoding genes including 5,061 enzymes and 156 transporters associated with 1,610 compounds assembled in a network of 331 pathways (*versus* the 227 pathways found in BioCyc). Like other metazoans, Dm possesses the gene repertoire to produce all the NEAA^{Fly}, but is unable to produce the

EAA^{Fly} (Fig 1A and Table S1). Dm can also produce myo-inositol, inosine and uridine,

but is unable to synthesize vitamins from simple precursors (Fig 1B and Table S2).

Ap^{WJL} biosynthetic capabilities inferred from *in silico* metabolic network

reconstruction

According to our metabolic reconstruction, Ap^{WJL} database comprises 4,268

protein-encoding genes including 1,326 enzymes and 46 transporters associated with

1,306 compounds assembled in a network of 313 pathways. The reconstructed

metabolic network reveals that Ap^{WJL} is a complete autotroph for all amino acids, (Fig

1A and Table S1). Ap^{WJL} also possesses the genetic potential to produce the DNA

bases inosine and uridine and five of the seven vitamins present in the HD: biotin,

folate, pantothenate, riboflavin and thiamine. The biosynthesis of nicotinate from Asp

seems impossible since L-aspartate oxidase (1.4.3.16), the enzyme necessary to

catalyze the first step of the pathway is absent (Fig 1B and Table S2), and further

TBLASTN searches failed to retrieve any 1.4.3.16 activity based on Ap^{WJL} genome. In

the case of pyridoxine, Ap^{WJL} cannot catalyze the first two reactions of its biosynthetic

pathway (Table S2). A direct phosphorylation of Thr would also be needed to complete

the pathway (using the 3.1.3.3 activity, Table S2). However, no evidence for such an

activity in *Acetobacter sp.* was found in genomic data or in the literature. Pyridoxine is

reported as non-essential for acetic acid bacteria [63]. Finally, Ap^{WJL} cannot produce

choline and myo-inositol. In summary, Ap^{WJL} genome analysis predicts that it is able

to synthetize all amino acids and five B-vitamins (biotin, folate, pantothenate, riboflavin

and thiamine). Ap^{WJL} cannot synthesize choline, myo-inositol and nicotinate.

Lp^{NC8} biosynthetic capabilities inferred from *in silico* metabolic network reconstruction

Metabolic reconstruction from Lp^{NC8} genome generated a database that includes 2,868 protein-encoding genes, consisting of 973 enzymes and 74 transporters associated with 1,154 compounds, all assembled in a network of 246 pathways. Interestingly, although the genome of Lp^{NC8} is about 30 % smaller than Ap^{WJL}'s genome, it encodes ~40% of supplementary transporters. This reflects the importance of metabolic exchanges for survival and adaptation of *L. plantarum* strains to a broad spectrum of environments [64].

From a genomic perspective (Fig 1A and Table S1), Lp^{NC8} is able to produce most amino acids from glucose or inner precursors. Exceptions are Phe, sulfur-containing (Cys, Met) and branched-chain amino acids (BCAA, Ile, Leu, Val). Indeed, synthesis of Phe seems to be precluded in Lp^{NC8} since the prephenate dehydratase activity (4.2.1.51) is not encoded by its genome (Table S1). Moreover, Lp^{NC8} has lost the capability to reduce sulfate in sulfite, a necessary step for the integration of sulfur in Cys biosynthesis [65]. Biosynthesis of Met seems to be feasible in Lp^{NC8}, but is probably limited by sulfite production rate and consequently by Cys availability. As previously reported, Lp^{NC8} is unable to produce the three BCAA, as four enzymatic activities of this pathway are lacking (1.1.1.86, 4.2.1.9, 2.3.3.13 and 4.2.1.33) [66]. However, the last transamination step common to each specific BCAA pathway is conserved, suggesting rather than importing the amino acids themselves, Lp^{NC8} may import 2-oxo-carboxylic acids, the direct precursors of Ile, Leu and Val to fulfill the BCAA requirements. Furthermore, Lp^{NC8} genome encodes a complete Arg biosynthesis pathway, yet Arg is known to be limiting [67] or essential to certain *L. plantarum* strains [66,68]. Arg auxotrophies are currently little understood; they are

usually attributed to regulatory effects or to the presence of certain deleterious

mutations in the biosynthetic pathway [68]. We further estimated the Arg biosynthetic

capability of Lp^{NC8} by analyzing the sequence of functional gene cluster *argCJBDF* and

searched for possible mutations using an autotrophic strain of *L. plantarum* (*L.*

plantarum CCM 1904, [69]) as a reference. No stop codon, frame-shifts nor deletions

were identified in the genes of the operon. This suggests that Lp^{NC8} can be an Arg

autotroph, but does not exclude the possibility that Arg may become limiting in specific

conditions. Lp^{NC8} has the minimal gene requirements to produce Ala and Asp (Table

S1), but the biosynthesis of Ala and Asp invokes secondary metabolic routes.

Therefore, when possible, uptake of Ala and Asp from the culture medium should be

favored instead of *de-novo* biosynthesis. Similarly, biosynthesis of Thr is directly linked

to Asp and Cys and is probably very limited in Lp^{NC8}. Accordingly, all the Thr

interconnecting pathways are lost, thus limiting its production or uptake from the diet

for protein synthesis.

Regarding vitamin biosynthesis, Lp^{NC8} is able to produce folate, riboflavin and

thiamine, as well as all DNA bases including uridine and inosine (Fig 1B and Table

S2). Lp^{NC8} cannot produce thiamine through the canonical pyrimidine pathway

because it lacks the 4.1.99.17 enzyme, but it seems possible through the pyrimidine

salvage pathway (2.1.7.49). Unlike Ap^{WJL}, pyridoxine biosynthesis seems feasible in

Lp^{NC8} using an alternative route involving Thr phosphorylation. Conversely,

biosynthesis of nicotinate seems to be impossible. Indeed, the two first steps of its

biosynthetic pathway (1.4.3.16 and 2.5.1.72) are not encoded by Lp^{NC8} genome.

Similarly, Lp^{NC8} has lost the capability to produce pantothenate as two enzymatic

activities of the pathway (2.1.2.11 and 6.3.2.1) are not present in the genome. Finally,

like Ap^{WJL}, Lp^{NC8} likely does not produce choline and myo-inositol.

In summary, our metabolic reconstruction suggests that Lp^{NC8} can synthesize

all amino acids except for Phe, Cys, Ile, Leu and Val. Synthesis of Arg, Thr, Ala and

Asp can easily become limiting in the contexts of high nutritional demand. Thus, Lp^{NC8}

is expected to extract these amino acids from the diet, or its growth would be altered

when they are absent from the diet. Met synthesis seems to be possible, but is

dependent on Cys abundance. Finally, Lp^{NC8} can produce DNA bases, folate,

riboflavine, thiamine and perhaps pyridoxine, but not biotin, nicotinate, pantothenate,

choline and myo-inositol.

Collectively, our *in silico* metabolic network reconstruction shows that

D. melanogaster and its commensals have differential biosynthetic capacities. Indeed,

some of the complete biosynthetic pathways are only present in one organism while

others are present in two or all the three partners. However, for incomplete biosynthetic

pathways we did not detect potentially shared pathways between a commensal and its

host as observed between obligate mutualistic partners (Fig 1-2).

Experimental validation of commensal bacteria auxotrophies using holidic diets

In order to test experimentally the metabolic potential of Dm and its

commensals, predicted by our in silico analysis (see above), we adopted the exome

based FLYAA holidic diet (HD) [40]. We systematically removed a single component

at a time to generate 39 different fly nutritional substrates (henceforth named HDΔX,

X being the nutrient omitted), plus one complete HD medium. This medium can also

be prepared in a liquid version by omitting agar and cholesterol from the recipe. Liquid

HD can then be used to assess bacterial growth in 96-well plates, increasing the

experimental throughput.

We first assessed Ap^{WJL} and Lp^{NC8} growth in each of the 40 different liquid HD

for 72 hours, using maximal optical densities (OD_{Max}) as a readout (Fig 3A and Table

S3). In the complete HD, both Ap^{WJL} and Lp^{NC8} grow well (Fig 3A, first line). On the

deficient media, Ap^{WJL} can grow in HDΔSucrose, presumably using acetate from the

acetate buffer as a carbon source. Also, its growth is not altered in the absence of any

of EAA^{Fly}, Vitamins or NAL. However, while Ap^{WJL} growth is not impacted by the lack

of most $NEAA^{Fly}$, it grows poorly in HDΔAla, HDΔCys and HDΔGlu. In addition, Ap^{WJL}

fails to grow in HDΔCu, HDΔFe and HDΔMg (Fig 3A, first column and Table S3). The

broad growth capacity of Ap^{WJL} in HDs correlates well with the wide range of

environmental niches the genus *Acetobacter* can colonize. Acetobacter species are

found in sugar-rich niches such as flowers and fruits, but also in poorer niches such as

soil and water, where they need to synthesize all nutrients required for their own growth

[70]. These findings are corroborated by our genome-based *in silico* predictions (Fig

1). Furthermore, the *in silico* reconstruction predicted that Ap^{WJL} would not be able to

synthesize choline, myo-inositol and pyridoxine, but we observed that Ap^{WJL} grows in

their absence. Choline is an important precursor of phosphatidylcholine (PC), which is

a major component of Acetobacter membranes and plays an important role to confer

acetic acid tolerance. Despite its importance, PC is not essential for Acetobacter

growth. Indeed, mutants precluding PC synthesis show a shift towards increased

membrane content of phosphatidylethanolamine (PE) and phosphatidylglycerol (PG)

and do not show any growth defect [71]. Similarly, Ap^{WJL} likely does not need myo-

inositol for its growth since inositol compounds are absent from the membrane of most

bacteria [72]. Regarding nicotinate and pyridoxine, the biosynthesis pathways of these

two nutrients are only partially understood and the presence of alternative pathways

and enzymatic activities may be possible in Ap^{WJL} (see above). It is thus possible that

Ap[WJL] is capable of synthesizing nicotinate and pyridoxine through elusive biochemical

pathways permitting growth in their absence. Further investigation will be required to

identify such biochemical pathway(s). Interestingly, Ap[WJL] growth was only precluded

in the absence of some metal ions: Cu, Fe, and Mg. Metal ions are important co-factors

required for enzymatic activities [73]. Specifically, in acetic acid bacteria, Cu is an

important co-factor of the energy-producing cytochromes of the respiratory chain [74]

making it essential for Ap[WJL] growth.

We detected far more nutritional auxotrophies for Lp[NC8] on HDs (Fig 3A, second

column and Table S3). Lp[NC8] fails to grow in HDΔSucrose, since sucrose is the only

suitable carbon source for Lactobacilli in the liquid HD. Also, Lp[NC8] growth is precluded

in the absence of 9 amino acids, including 6 EAA[Fly] (Arg, Ile, Leu, Phe, Thr, Val) and

3 NEAA[Fly] (Ala, Asp, Cys). It also grows poorly in media lacking the EAA[Fly] Lys, Met

and Trp and the NEAA[Fly] Asn and Glu. Moreover, Lp[NC8] does not grow in HDΔBiotin

and HDΔPantothenate. However, it grows in absence of nicotinate, which is predicted

to be essential according to our *in silico* analysis (see above). Finally, Lp[NC8] growth is

not affected by the lack of any NAL. In the absence of certain metal ions such as Ca,

Cu, Mg, Zn , OD_{Max} is increased. On the contrary, OD_{Max} is significantly reduced in

HDΔMn. These relatively elevated nutritional requirements of Lp[NC8] are expected as *L.*

plantarum is a species adapted to nutrient-rich environments [75]. Hence, many *L.*

plantarum strains have lost the capacity to synthetize various nutrients that can easily

be found in their natural habitats [64,75]. The inability of *L. plantarum* to synthetize

important nutrients such as BCAA (Ile, Leu and Val) or the B-vitamin pantothenate was

previously identified by both genome analyses [66] and growth studies in chemically

defined minimal media [67,76,77]. Moreover, it is known that *L. plantarum* needs Mn

to resist to oxidative stress [78], which explains its poor growth in HDΔMn. Our

experimental data only partially correlate with the results of our genomic analysis. *In*

silico predicted auxotrophies to Ile, Leu, Val, Phe, Cys, pantothenate and biotin were

confirmed *in vivo*. The identified Arg auxotrophy was not surprising because, as

mentioned above, Arg is often described as essential to *L. plantarum* in high metabolic

demanding conditions even though all genes necessary for Arg biosynthesis are

present. However, auxotrophies of Lp^{NC8} to Thr, Ala, Asp, were not expected (Fig 3A,

denoted by "*"), even though these amino acids were predicted to be limiting (see

above). As mentioned previously, bacterial growth in liquid HD was assessed in 96-

well plates using a microplate reader (see methods). Every cycle includes an agitation

step to homogenize the solution to improve OD reading accuracy. This agitation step

may oxygenate the media and thus negatively affects Lp^{NC8} growth in scarcity

conditions since *L. plantarum* strains are aerotolerant but optimal growth is achieved

under microaerophilic or anaerobic conditions [79]. To retest these unexpected

auxotrophies, we assessed Lp^{NC8} growth in liquid HDΔThr, HDΔAla, HDΔAsp in 15-mL

closed falcon tubes without aeration. After 72 hours of incubation we determined CFU

counts in each media (Fig 3B). As predicted by our genomic analyses, Lp^{NC8} was able

to grow in each of the three deficient media in static conditions to the same extent as

in the complete HD (Fig 3B). Therefore, Lp^{NC8} auxotrophies observed for Thr, Ala and

Asp in 96-well plates are likely an artifact due to excessive oxygenation. This could

also explain the poor growth of Lp^{NC8} in the absence of the EAA^{Fly} Lys, Met and Trp

and the NEAA^{Fly} Asn and Glu. The ability of Lp^{NC8} to grow in HDΔCholine, HDΔMyo-

inositol, HDΔNicotinate and HDΔPyridoxine do not correlate with *in silico* data. As for

Ap^{WJL} (see above), Lp^{NC8} growth probably does not require choline or myo-inositol. A

previous study quantified choline and inositol compounds in *L. plantarum* cell extracts

and found them to be extremely low and therefore most likely due to contaminations

from the medium rather than components of *L. plantarum* biomass [80]. Moreover, pyridoxine is a precursor of pyridoxal-5-phosphate, a cofactor necessary for amino acid converting reactions. Teusink et al. [66] showed that *L. plantarum*[WCSF1] requires exogenous sources of pyridoxine only in a minimal medium lacking amino acids. Since the HDΔPyridoxine contains all amino acids, it is likely that pyridoxine is not essential for Lp[NC8] growth in these conditions. Finally, the capacity of Lp[NC8] to grow in HDΔNicotinate could be related with the presence of an alternative pathway of nicotinate biosynthesis. Indeed, this possibility has been previously reported in the genera *Lactobacillus* [64] which would explain the capacity to grow in absence of exogenous nicotinate.

Altogether, the complete HD is a suitable nutritional environment that allows the two prevalent Drosophila commensals, Ap[WJL] and Lp[NC8] to grow. Growth capacities in deficient media vary from one bacterium to another and are dictated by their individual genetic repertoires.

Germ free larvae exhibit 22 auxotrophies while developing on holidic diets

We next sought to establish the nutritional requirements of GF larvae by assessing larval developmental timing (DT) in complete HD and in each of the 39 deficient HD (see methods). DT is expressed as D_{50}, which represents the day when 50% of the larvae population has entered metamorphosis in a specific nutritional condition. In agreement with previous studies [38,39], GF larvae fail to develop in all HDΔEAA[Fly], all HDΔVitamins, HDΔCholine HDΔCholesterol, HDΔZn and HDΔMg (Fig 4A, first column). Over 60 years ago, Sang et al reported that Zn was dispensable for GF larval development in a classic study [38]. We suspect that the casein in the medium used in Sang et al inadvertently provided trace amount of Zn, which could

account for the discrepancy between our observation and that of Sang et al. Also in

accordance with previous studies [38,39,49], GF larvae were able to reach pupariation

in HDΔNEAA^{Fly} (ΔAla, ΔCys, ΔGln, ΔGlu, ΔGly, ΔPro), HDΔUridine, HDΔMyo-inositol,

and HDΔMn at the same rate as on a complete HD (Fig 4A). The absence of sucrose,

Tyr, inosine, Ca, Cu and Fe did not prevent pupae emergence, but increased their

developmental timing significantly (Fig 4A first column, Table S4). Surprisingly, GF

larvae were able to reach pupariation, albeit late, in HDΔSucrose. Indeed, all the HDs

developed to date include carbohydrates (either sucrose or fructose) as a carbon

source [34]. Larval development in the absence of carbohydrates suggests that GF

larvae may use other components of the HD as carbon source. In summary, GF *yw*

larvae show 22 auxotrophies while developing on sterile holidic diets.

Our observations correlate well with our *in silico* predictions (Fig 1) with one

exception: GF larvae did not reach pupariation in HDΔAsn. This result was surprising

because Asn is described as a NEAA in Drosophila and other animals [81]. To test

whether Asn auxotrophy was specific to the *yw* fly line used in our assays, we assessed

larval DT in two other Dm reference lines, DGRP-*RAL-25210* [82] and *w^{1118}*. Unlike

yw, both *w^{1118}* and *RAL-25210* larvae were able to develop in GF condition in HDΔAsn

albeit with a severe developmental delay (Fig 4B). Therefore, a complete Asn

auxotrophy is specific to our *yw* strain, which explains why we did not detect this

auxotrophy in our *in silico* analyses that were based on the genome sequence of the

Drosophila Reference Genome strain (Bloomington stock #2057). We therefore

sequenced the coding region of the enzyme AsnS that converts Asp to Asn in *yw* flies,

and did not detect any non-synonymous mutation (Fig S1). Further studies may thus

be required to determine the origin of the Asn auxotrophy in our *yw* line on HD.

However, these results indicate that Asn is not an EAA per se but remains a limiting

NEAA, an observation that also applies to Tyr.

The molecular dialogue engaged between commensals and their host is at play

on complete holidic diet

We then investigated if and how the association with commensals affects the

nutritional requirements of GF larvae during juvenile growth and maturation. To this

end, we mono-associated GF embryos with Ap^{WJL} or Lp^{NC8} and measured D_{50} and egg-

to-pupa survival in complete and deficient HDs (Fig 4A second and third columns,

respectively and Table S4, S5). On a complete HD mono-association with either Ap^{WJL}

or Lp^{NC8} accelerated larval DT with a mean D_{50} of 8.4 and 7.7 days, respectively,

whereas GF mean D_{50} 10.1 days (Fig 3A, first line).

These growth-promoting effects upon mono-association with either Ap^{WJL} or

Lp^{NC8} have been previously reported on complex diets and insights on the underlying

molecular mechanisms were provided [20,25]. Shin *et al.* showed that when the

associated larvae grow on a low casamino-acid semi-oligidic diet, the pyrroloquinoline

quinone–dependent alcohol dehydrogenase (PQQ-ADH) activity of Ap^{WJL} modulates

the developmental rate and body size through insulin/insulin-like growth factor

signaling (IIS). PQQ-ADH disruption in the Ap^{WJL}::Tn*pqq* mutant severely reduces

acetic acid production, which has been proposed to alter the regulation of

developmental and metabolic homeostasis upon mono-association [25]. Lp^{NC8}

promotes host juvenile growth and maturation on a low yeast-based oligidic diet, partly

through enhanced expression of intestinal peptidases upon sensing of bacterial cell

walls components by Drosophila enterocytes [20,23]. Deletion of the *dlt* operon, which

encodes the molecular machinery involved in the D-alanylation of teichoic acids, leads

to bacterial cell wall alteration with a complete loss of D-alanylation of teichoic acids

and consequently cell walls purified from the Lp$^{NC8}\Delta dlt_{op}$ mutant trigger a reduced

expression of peptidases in enterocytes [23]. Therefore, we first probed the importance

of these molecular mechanisms on commensals mediated larval growth promotion on

complete HD. To this end, we tested in our HD setting the loss of function mutants

Ap^{WJL}::Tnpqq [25] and Lp$^{NC8}\Delta dlt_{op}$ [23]. In complete HD, only the Lp$^{NC8}\Delta dlt_{op}$ mutant

failed to support larval growth, reminiscent of the previous observation on the low-

yeast oligidic diets (Fig S2). Surprisingly in complete HD, the Ap^{WJL}::Tnpqq mutant

actually triggered an enhanced growth promotion as compared to its WT reference

strain. Shin *et al.* reported that Ap^{WJL}::Tnpqq associated larvae experienced growth

delay, which can be rescued by acetic acid provision [25]. Therefore, the acetic acid-

based buffer in HD may explain why Ap^{WJL}::Tnpqq no longer behaves as a loss of

function mutant in this setting; however, how it actually surpasses the WT strain on HD

remains elusive. Collectively, these results establish that sensing bacterial cell walls

containing D-alanylated teichoic acids is also an important feature of the intrinsic

growth promoting ability of Lp^{NC8} in a complete chemically defined HD. Thus, the

previously reported molecular sensing mechanism that mediates the growth-promoting

effect of Lp^{NC8} during chronic undernutrition is also at play in synthetic diets.

Association with Ap^{WJL} fulfills 19 of the 22 auxotrophies of GF larvae

Association with Ap^{WJL} sustained larval development in the absence of 19 out of 22

GF essential nutrients (Fig 4A second column). Ap^{WJL} associated larvae reached

pupariation in the absence of each EAA^{Fly} (though their development was slower than

on complete HD), Asn, vitamins, choline and Zn. Association with Ap^{WJL} also rescued

the developmental delay observed in GF larvae in HDΔTyr, HDΔinosine, HDΔCu and

HDΔFe. The only nutritional requirements of GF larvae that were not fulfilled by Ap^{WJL}

were cholesterol, pantothenate and Mg.

Association with Lp^{NC8} fulfills 12 of the 22 auxotrophies of GF larvae

Compared to Ap^{WJL}, mono-association with Lp^{NC8} compensated for a reduced

number of the GF larvae nutritional deficiencies (12 out of 22; Fig 4A third column).

Lp^{NC8} associated larvae reached pupariation in the absence of some EAA^{Fly} (HDΔHis,

HDΔLys, HDΔMet, HDΔPhe, HDΔThr), Asn, certain vitamins (HDΔBiotin, HDΔFolate,

HDΔNicotinate, HDΔRiboflavine, HDΔThiamine), and Zn. Moreover, Lp^{NC8} rescued the

developmental delay observed in GF on HDΔTyr, HDΔinosine, HDΔCu and HDΔFe.

Bacteria need to be metabolically active in order to fulfill larval nutritional

requirements

Bacteria were grown in rich medium before association with larvae. Therefore, they

may have accumulated nutrients that can be used by the larvae to fulfill their nutritional

requirements. To test for the nutritional input brought by the initial bacterial inoculum,

we associated GF larvae with 10X heat-killed (HK) bacteria and measured D_{50} in

complete and deficient HDs (Fig 4A fourth and fifth columns). In most cases, the D_{50}

of larvae in HK and GF conditions were similar, and they both show striking delay

compared to the mono-associated larvae. Therefore, bacteria need to be metabolically

active to fulfill the larval nutritional requirements on HDs. However, we found some

exceptions. In HDΔAsn, HDΔBiotin, HDΔFolate, HDΔCu and HDΔFe, addition of HK

bacteria allowed the larvae to develop, though not as fast as in association with living

bacteria. These results suggest that larvae only require a very small amount of these

nutrients, which can be sufficiently derived from the inert bacterial inoculum.

The ability of bacteria to compensate nutritional deficiencies does not always correlate with the ability of bacteria to synthesize the nutrient

Next, based on the *in silico* predictions and the experimentally revealed

auxotrophies, we correlated the ability of each commensal bacteria to synthesize a

nutrient to their ability to fulfill the larval requirements in this nutrient. We identified 4

distinct scenarios:

<u>Case 1: the bacteria synthesize a nutrient and fulfill the related larval auxotrophy</u>

In most of the tested conditions, when the bacteria can synthesize a nutrient, they can

also fulfill the related nutritional requirements of the GF larvae. For Ap^{WJL} this includes

all EAA^{Fly}, Asn and most vitamins (except pantothenate). For Lp^{NC8}, the correlation

between the nutritional complementation of ex-GF larva and the ability of Lp^{NC8} to

synthesize the missing nutrient is more limited and only applies to the requirements of

His, Lys, Met, Thr, Asn and most vitamins (except pyridoxine). Nonetheless, these

results suggest that bacteria can actively supply the nutrients lacking in the HD to the

larvae. This phenomenon is reminiscent of previous observations using conventional

and gnotobiotic hosts, in which microbial sparing of riboflavin or thiamine by differential

commensals have been proposed [33,83]. Exceptions to this case seem to be Ap^{WJL}

on HDΔPantothenate and Lp^{NC8} on HDΔTrp. Specifically, Ap^{WJL} can produce

pantothenate and grows in HDΔPantothenate and similarly Lp^{NC8} can produce Trp and

grows in HDΔTrp. However, neither supported larval development on the respective

depleted HD. It is therefore probable that Ap^{WJL} and Lp^{NC8} produce enough

pantothenate and Trp, respectively, to sustain their own growth in the depleted HD but

not sufficiently or in a manner inaccessible to the larvae and thus fail to fulfill larval

requirements for these nutrients.

Case 2: the bacteria do not synthesize a nutrient and they cannot fulfill larval

requirements

Expectedly, we observed that when bacteria do not synthesize a nutrient, they

do not fulfill ex-GF larvae requirements for this nutrient. For instance, Lp^{NC8} cannot

produce the BCAA, nor grow in their absence, and thus it cannot fulfill larval

requirements for these amino acids. In some depleted diets such as HDΔCholesterol,

HDΔCholine and HDΔPyridoxine, bacteria were able to grow (Fig 3A) even though they

cannot synthetize these nutrients (See above and Table S2), and they failed to fulfill

the larvae requirements of these specific nutrients. This is observed for Ap^{WJL} and

Lp^{NC8} on HDΔCholesterol. The likely explanation is that cholesterol is an animal sterol

but is dispensable for bacterial growth [72,84].   Similarly, on HDΔCholine and

HDΔPyridoxine, Lp^{NC8} is also unable to fulfill larval requirements (Fig 3A) as it cannot

synthetize these compounds.

Case 3: the bacteria do not synthesize a nutrient but they can fulfill larval requirements

In most cases, we observe growth rescue by bacteria provision of the missing

nutrients, but there are interesting exceptions. Ap^{WJL} is unable to synthesize *de-novo*

choline and pyridoxine (See above and Table S2). Surprisingly, it fulfills larval

auxotrophies on HDΔCholine and HDΔPyridoxine. This could be achieved either by a

functional compensation mechanism or via alternative biosynthetic pathways of these

components that are undetected by our metabolic reconstruction. In the case of

choline, the former seems to be the most plausible explanation, as Ap^{WJL} may

synthesize other compounds that Drosophila can use to replace choline. As stated

before, Acetobacter mutants precluding phosphatidylcholine (PC) synthesis shift their

membrane composition towards increased content of phosphatidylethanolamine (PE)

and phosphatidylglycerol (PG) [71]. PE and PG have been reported to be part of the

phospholipidic repertoire of Drosophila membranes [85], in which PE represents

approximately 50% of their lipid composition [86]. We posit that ex-GF larvae growing

on HDΔCholine may capitalize on ethanolamine or glycerol produced by Ap^{WJL} to

compensate for the lack of choline in their diet. In the case of pyridoxine, Ap^{WJL} may

fulfill larval requirements by either a functional compensation or through pyridoxine

biosynthesis by non-canonical pathways (see above). We reach the same conclusion

regarding nicotinate. Both Ap^{WJL} and Lp^{NC8} grow well on HDΔNicotinate and also can

fulfill the larval requirements in this vitamin even though our metabolic reconstruction

predicts that they cannot synthesize it (see above).

Moreover, Lp^{NC8} cannot grow in the absence of Phe (Fig 2A). The genomic

analyses point to the possible loss of the gene coding for the enzyme prephenate

dehydratase (4.2.1.51), the penultimate step on Phe biosynthesis. Yet Lp^{NC8} can fulfill

larval requirements for Phe (Fig 4A). We wondered if the Phe auxothrophy we

observed in 96-well plates (Fig 3A) was due to the oxygenation generated by the

agitation through OD readings, as for Thr, Ala and Asp (Fig 3B). To test this, we set

cultures of Lp^{NC8} in HDΔPhe in static 15-mL closed falcon tubes and assessed bacterial

growth after 3 days of culture. In contrast to agitation, Lp^{NC8} grows in HDΔPhe ~1×10^2

times in static conditions (Fig 4C), whereas in the complete media (Fig 3B), Lp^{NC8}

grows ~2×10^4 times. Therefore, these results indicate that the rescue of larvae

developmental timing by Lp^{NC8} in HDΔPhe is still mediated by bacterial nutrient supply.

However, the poor growth of Lp^{NC8} in HDΔPhe suggests the existence of an alternative

pathway for Phe biosynthesis in absence of the prephenate dehydratase (Fig 1A). As

suggested by Hadadi et al. [87], Phe might be produced from L-arogenate using a

derivative catalysis through the 2.5.1.47 activity which is encoded in Lp^{NC8} by the *cysD*

gene (nc8_2167).

A second such interesting case is larval developmental rescue by Lp^{NC8} in

HDΔCys. Lp^{NC8} is auxotroph for Cys (Fig 3A), even in static conditions (Fig 4C). Lp^{NC8}-

associated larvae develop faster than GF larvae, though GF larvae are not auxotroph

to Cys (Fig4A). This beneficial effect of Lp^{NC8} on ex-GF larvae development in HDΔCys

is similar to what is observed on a complete HD (Fig 4A, first row) and therefore

probably reflects the basal nutrient-independent growth-promoting effect of Lp^{NC8}, that

we previously reported and which relies on the molecular dialog between Lp^{NC8} and its

host (Fig S2) [23]. Taken together, our results indicate that Lp^{NC8} is able to grow in

HDΔCys only in the presence of Dm larvae. To test this hypothesis, we assessed Lp^{NC8}

growth in solid HDΔCys in the absence (Medium Load) and the presence (Niche Load)

of Dm larvae (Fig 4D). Without larvae (ML), Lp^{NC8} grew one log above the inoculum

level in solid HDΔCys (Fig 4D, Medium load). This minimal growth in solid HDΔCys

could be due to the Cys reserves from Lp^{NC8} growth in rich media prior to inoculation,

or from contaminants in the agar and cholesterol added to prepare the solid HD.

Interestingly, in the presence of larvae in the HDΔCys (NL), Lp^{NC8} CFU counts

dramatically increased over time, reaching ~10^8 CFU/tube at day 6 (net growth: ~3×10^4

times inoculum. Fig 4D). These results suggest that in HDΔCys larvae support Lp^{NC8}

growth, probably by supplying Cys or a precursor/derivative, which in turn promotes

larval development and maturation. This observation extends the recent demonstration

that Dm and *L. plantarum* engage in a mutualistic symbiosis, whereby Dm benefits the

growth of *L. plantarum* in their shared nutritional environment [30]. Here we discover

that Cys is a Dm symbiotic factors also referred to as "bacteria maintenance factors"

[30].

Case 4: mineral and metal traces

We observed that both Ap^{WJL} and Lp^{NC8} could compensate for Cu, Fe and Zn

deficiencies (Fig 4A, second and third column). Requirements in Cu and Fe were also

fulfilled by HK bacteria (Fig 4A, fourth and fifth column) although larvae associated with

HK bacteria in these conditions developed much slower than larvae associated with

living bacteria. This suggests that the inert bacterial inoculum contains traces of Cu

and Fe accumulated during the overnight growth in rich medium prior to the inactivation

and inoculation. These accumulated quantities allowed the larvae to develop when Cu

and Fe were not supplied in the HD. Surprisingly, Zn requirements were fulfilled by

living bacteria only (Fig 4A). We hypothesize that bacteria may concentrate

contaminating traces of these elements in the HD and make them more available to

the larva. Alternatively, this could be an interesting case of functional complementation

that requires further investigation. Indeed, Zn is an important enzymatic cofactor in the

biosynthesis of several metabolites by the larva [88]. In the absence of Zn, these

compounds would not be produced by the GF larva, instead they could be produced

by the bacteria and supplied to the ex-GF larvae similarly to the nutritional

complementation we observed above (examples in Case 3). Interestingly, a link

between Zn response and the microbiota of Drosophila have been described in

previous studies. Expression of the Zn transporter *zip-3* is higher in GF Drosophila

adults midguts than in their conventionally raised (CR) counterparts [27]. Moreover,

the genes encoding Metallothioneins B and C (*MtnB* and *MtnC*) are more expressed

in flies harboring a microbiota than in GF flies [89]. Metallothioneins are intracellular

proteins that can store Zn. Their expression, as well as expression of Zn transporters

such as *zip-3*, is regulated by intracellular levels of Zn [90]. Altogether, these results

suggest that commensal bacteria may play an important role in the uptake of metals

(especially Zn) by Drosophila larvae. This idea is reminiscent of recent reports in

Caenorhabditis elegans, whereby a commensal bacteria promotes worm development

upon Fe scarcity by secreting a scavenging siderophore [91].

Commensal bacteria spare amino acids essential to larval development

Despite the interesting exceptions detailed above, our data establish that in

general, bacterial commensals complement the nutritional requirements of their host

by synthesizing and supplying essential nutrient. Bacteria can actively excrete amino

acids in their environment when they are produced in excess as intracellular by-

products of metabolic reactions [92]. Moreover, the bacterial cell wall is rich in D-amino

acids, and it undergoes an important turnover [93,94]. In certain bacterial species, D-

amino acids accumulate in the supernatant during growth and act as a signal to

undergo stationary phase [95]. Thus, D-amino acids may also contribute to larval

nutrition. Indeed, it has been previously shown that D-amino acids (D-Arg, D-His, D-

Lys, D-Met, D-Phe, and D-Val) can support growth of GF larvae probably through the

action of amino acid racemases [47]. We thus hypothesized that Ap^{WJL} and Lp^{NC8} could

provide amino acids to their host by releasing them in the HD. To directly tested this

hypothesis, we cultured Ap^{WJL} and Lp^{NC8} in liquid HD lacking each EAA^{Fly} and

quantified the production of the corresponding missing EAA^{Fly}. We focused on EAA^{Fly}

whose deficiency could be compensated by the commensals in our DT experiments

(Fig. 4A). In these assays, Ap^{WJL} was cultured under agitation and Lp^{NC8} cultures were

grown in both agitated and static conditions (See Methods). After three days, we

quantified the amino acids concentration from bacterial supernatants using high

pressure liquid chromatography (HPLC). We quantified amino acid production by

Ap^{WJL} under agitation while growing in HDΔArg, HDΔHis, HDΔIle, HDΔLeu, HDΔLys,

HDΔMet, HDΔPhe, HDΔThr, and HDΔVal and observed accumulation all missing AAs

expect for Lys and Met (Fig 5A). For Lp[NC8], we analyzed the supernatants of HDs that

support Lp[NC8] growth under agitation (Fig 3A): HDΔHis, HDΔLys and HDΔMet. We also

analyzed supernatants from static conditions, HDΔHis, HDΔLys, HDΔMet, HDΔPhe

and HDΔThr. Surprisingly, from all tested conditions we only detected His

accumulation in the supernatant of Lp[NC8] grown on HDΔHis under agitation (Fig 5B).

We did not detect Lys and Met in Ap[WJL] culture supernatant or Lp[NC8] culture under

agitation supernatant nor His, Lys, Met, Phe or Thr in Lp[NC8] static culture supernatants.

However, Ap[WJL] or Lp[NC8] can both fulfill larval requirements in a HD lacking these

nutrients (Fig 4A). We only analyzed supernatants after 72h of growth, it is therefore

possible that we missed the peak of accumulation of the targeted AA, which may have

taken place at another time point during the growth phase. Also, Ap[WJL] and Lp[NC8] may

only secrete precursors or catabolites of these AA that we did not target in our analysis.

Such AA derivatives may also be used by the larvae to compensate for the lack of the

cognate amino acids in the diets. However, we detected Arg, His, Ile, Leu, Phe, Thr

and Val production by Ap[WJL] and His by Lp[NC8], a production which correlates with the

respective ability of Ap[WJL] and Lp[NC8] to compensate for the lack of these AAs in the

respective depleted HD. Of note, the concentration of newly synthetized amino acids

accumulating in the supernatant is low compared to their concentration in a complete

HD (20-150 µM in the former vs 1-5 mM in the latter). However, the bacterial supply of

amino acids to the larvae is probably a continuous process, which can be stimulated

upon uptake and transit through the larval intestine. Thus, amino acids are directly

supplied to the larvae and will fulfill its nutritional requirements without the need to

accumulate in the surrounding media. Altogether, our results show that Ap[WJL] and

Lp[NC8] are able to synthetize and excrete some EAA[Fly] in their supernatants. These

results confirm our hypothesis that Dm commensal bacteria Ap^{WJL} and Lp^{NC8} produce

these EAA^{Fly} while growing on HDΔEAA^{Fly}. When associated to Dm larvae, Ap^{WJL} and

Lp^{NC8} will therefore supply these amino acids to the larvae, allowing larval development

on these deficient media as observed upon mono-associations (Fig 4A).

Conclusion

In this study, we have unraveled the interactions between the microbial and the

nutritional environment of Drosophila, as well as the functional importance of these

interactions for Drosophila juvenile growth. We systematically characterized, both *in*

silico and *in vivo*, the biosynthetic capacities of growing GF larvae and two

representative commensal strains of bacteria (Ap^{WJL} and Lp^{NC8}). We show that both

commensals, each in its unique manner, alleviate the nutritional constraints in the

environment to accelerate host growth and maturation in diets depleted of essential

nutrients (Fig 6). The capacity of a bacterium to fulfill the larval requirements in a

specific nutrient correlated with its metabolic activity and in most cases its ability to

produce the missing nutrient. In contrast to obligate symbioses, our results highlight

the clear separation between the metabolic pathways of the host and its commensals

and reveal a particularly integrated nutritional network between the insect and its

facultative commensals around the sparing and utilization of nutrients.

Importantly, we further demonstrate that active nutrient provision by metabolite

biosynthesis not only evoke the canonical pathways, but also non-canonical pathways

such as the ones described here, for example nicotinate by both Ap^{WJL} and Lp^{NC8},

pyridoxine by Ap^{WJL}, and Phe by Lp^{NC8}. Interestingly, we also detected two cases

where nutrient compensation is not explained by a direct nutrient supply: (1) the

compensation of choline deficiency by Ap^{WJL} and (2) the compensation of Zn deficiency

by both Ap^{WJL} and Lp^{NC8}. We propose the existence of functional compensation

mechanisms whereby Ap^{WJL} would complement choline deficiency by synthesizing and

providing precursors and/or functional analogues of choline, such as ethanolamine or

glycerol and both commensals can compensate Zn deficiency by uptaking,

concentrating and sparing contaminant traces of Zn to the host.

Previous works have shown that the molecular dialogue between the host and its commensal partners invokes the sensing of commensal bacteria cell wall and the induction of transcriptomic rewiring of the digestive potential in the Drosophila enterocytes. Here we show that in addition, the metabolic activities of live Drosophila's commensal bacteria correct host auxotrophies. These results reveal a novel facet of the facultative nutritional mutualism engaged between Drosophila and its commensal bacteria, which supports the host's nutritional versatility and allows the juvenile host to better cope with changes in nutrients availability during the critical phase of post-natal growth; hence ensuring optimal host fitness. Our work lays the foundational blueprint for further mechanistic studies to investigate if and how commensals regulate the synthesis and release of essential nutrients for the host and if the host influences this regulatory process. Dissecting how commensals functionally compensate for nutrients that they cannot produce also constitute an attractive perspective for future investigations.

In some cases, the *in silico* prediction of bacterial biosynthetic capabilities were incongruent with our *in vivo* assessment of bacterial auxotrophies (Table S6). Such seeming discrepancy served as an entry point for us to discover novel phenomena and interactions that would have been missed had we only adopted a single approach. One such interesting example is Asn auxotrophy unique to the Dm *yw* line in GF conditions. Another one is the larval provision of Cys (or its derivatives) to Lp^{NC8} to maintain a mutualistic nutritional exchange between host and commensal. Previously, combination of *in silico* and *in vivo* approaches has been successfully used for bacteria [66] but not applied to complex symbiotic systems such as host-commensals nutritional interactions. Indeed, reports characterize these interactions *in silico* [96] but they were not confirmed *in vivo*. Our work closes this gap and emphasizes the importance of

703　using parallel systematic *in silico* and *in vivo* approaches for the understanding of the

704　intricate relationships between the microbial and the nutritional environments and their

705　impact on animal juvenile growth.

Materials and Methods

In silico reconstruction of the biosynthetic potential of *D. melanogaster* and its commensals

We used the Cyc Annotation Database System (CycADS) [62], an automated annotation management system, to integrate protein annotations from the complete genomes of *Drosophila melanogaster* (RefSeq GCF_000001215.4 release 6), *A. pomorum* strain DM001 (accession NCBI-Taxonomy 945681 version 1.0.1) and *L. plantarum* subsp. *plantarum* NC8 (NCBI Bioproject PRJNA67175). CycADS collects protein annotation results from different annotation methods including KAAS [97], PRIAM [98], Blast2GO [99,100], and InterProScan [101] in order to obtain EC numbers and GO annotations. All annotation information was then processed in the CycADS SQL database and automatically extracted to generate appropriate input files to build the three BioCyc databases using the Pathway Tools software v22.5 [102]. The BioCyc databases and their associated metabolic networks are available in the EcoCyc database [103]. From the genomic analyses, we inferred the biosynthetic capabilities of the three organisms and manually inspected all pathways allowing production of the organic compounds that are present in the exome-based FLYAA holidic diet (HD) developed by Piper et al [40]: fly's essential and non-essential amino acids (EAA[Fly] (n=10) and NEAA[Fly] (n=10)), B-vitamins (n=7), cholesterol (n=1), and Nucleic Acids and Lipid precursors (NAL, n=4). For each gap found in biosynthetic pathways or non-conventional enzymatic catalysis, TBLASTN [104] searches were performed in the three genomes to look for unpredicted protein activities. Alternative pathways were searched in the literature or using the BioCyc "Metabolic Route Search" tool [105].

Drosophila diets, stocks and breeding

Drosophila stocks were reared as described previously [32]. Briefly, flies were

kept at 25°C with 12/12-hour dark/light cycles on a yeast/cornmeal medium containing

50 g/L of inactivated yeast, 80 g/L of cornmeal, 7.4 g/L of agar, 4 mL/L of propionic

acid and 5.2 g/L of nipagin. Germ-free stocks were established as described previously

[89] and maintained in yeast/cornmeal medium supplemented with an antibiotic

cocktail composed of kanamycin (50 µg/mL), ampicillin (50 µg/mL), tetracycline (10

µg/mL) and erythromycin (5 µg/mL). Axenicity was tested by plating fly media on

nutrient agar plates. Drosophila *yw* flies were used as the reference strain in this work.

Other Drosophila lines used include DGRP-*RAL*-25210 [82] and *w*^{1118} [106].

Experiments were performed on Holidic Diet (HD) without preservatives.

Complete HD, with a total of 8 g/L of amino acids, was prepared as described by Piper

at al. using the fly's exome matched amino acid ratios (FLYAA) [40]. Briefly, sucrose,

agar, amino acids with low solubility (Ile, Leu and Tyr) as well as stock solutions of

metal ions and cholesterol were combined in an autoclavable bottle with milli-Q water

up to the desired volume, minus the volume of solutions to be added after autoclaving.

After autoclaving at 120°C for 15 min, the solution was allowed to cool down at room

temperature to ~60 °C. Acetic acid buffer and stock solutions for the essential and non-

essential amino acids, vitamins, nucleic acids and lipids were added. Single nutrient

deficient HD was prepared following the same recipe excluding the nutrient of interest

(named HD∆X, X being the nutrient omitted). Tubes used to pour the HD were

sterilized under UV for 20 min. HD was stored at 4°C until use, for no longer than one

week.

Bacterial strains and growth conditions

A. pomorum[WJL] (Ap[WJL]) [25], *L. plantarum*[NC8] (Lp[NC8]) [107], Ap[WJL]::Tn*pqq* [25]

and Lp[NC8]Δ*dlt*op [23] were used in this study. *A. pomorum* strains were cultured in 10

mL of Mannitol Broth (Bacto peptone 3 g/L, yeast extract 5 g/L, D-mannitol 25 g/L) in

50 mL flask at 30°C under 180 rpm during 24h. *L. plantarum* strains were cultured in

10 mL of MRS broth (Carl Roth, Germany) in 15 mL culture tubes at 37°C, without

agitation, overnight. Liquid or solid cultures of Ap[WJL]::Tn*pqq* were supplemented with

kanamycin (Sigma-Aldrich, Germany) at a final concentration of 50 µg/mL. CFU counts

were performed for all strains on MRS agar (Carl Roth, Germany) plated using the

Easyspiral automatic plater (Intersciences, Saint Nom, France). The MRS agar plates

were then incubated for 24-48h at 30°C for Ap[WJL] or 37°C for Lp[NC8]. CFU counts were

done using the automatic colony counter Scan1200 (Intersciences, Saint Nom,

France) and its counting software.

Bacterial growth in liquid HD

To assess bacterial growth in the fly nutritional environment we developed a

liquid HD comprising all HD components except agar and cholesterol. Liquid HD was

prepared as described for HD. Single nutrient deficient liquid HD was prepared

following the same recipe excluding the nutrient of interest. After growth in rich media,

PBS washed Ap[WJL] or Lp[NC8] were inoculated at a final concentration of ~ 10^6 CFU/mL

in 200 µL of either complete liquid HD or nutrient deficient liquid HD. Cultures were

incubated in 96-well microtiter plates (Nunc™ Edge 2.0. Thermo Fisher Scientific) at

30°C for 72h. Growth was monitored using an SPECTROstar[Nano] (BMG Labtech

GmbH, Ortenberg, Germany) by measuring the optical density at 600 nm (OD_{600}) every

30 minutes. Heatmap in Fig 3A represents the maximal OD detected during the 72h of

growth (average of three replicates). The whole experiment was repeated at least

twice. Fig 3A was created using the imagesc function on MATLAB (version 2016b.

MathWorks, Natick, Massachusetts). Lp^{NC8} growth in static conditions was performed

in 10 mL of liquid HD in 15-mL falcon tubes inoculated at a final concentration of ~10^4

CFU/mL. Cultures were incubated at 30°C for 72h. After incubation, cultures were

diluted in PBS and plated on MRS agar as described above. Growth is expressed as

$$\frac{\text{CFU}(t = 72h) - \text{CFU}\,(t = 0)}{\text{CFU}(t = 0)}.$$

Bacterial growth in solid HD

Bacterial CFUs in HDΔCys were assessed in presence (Niche Load: NL) or

absence (Medium Load: ML) of Dm larvae. Microtubes containing 400 μL of HD and

0.75–1 mm glass microbeads were inoculated with ~10^4 CFUs of Lp^{NC8}. For NL, 5 first-

instar larvae, collected from eggs laid on HDΔCys, were added. The tubes were

incubated at 30°C for 0, 3 or 6 days. After incubation, 600μL of PBS were added

directly into the microtubes. Samples were homogenized with the Precellys 24 tissue

homogenizer (Bertin Technologies, Montigny-le-Bretonneux, France). Lysates

dilutions (in PBS) were plated on MRS and CFU counts were assessed as described

above.   Growth   is   expressed   as

$$\frac{\text{CFU}(t = 72h) - \text{CFU}\,(t = 0)}{\text{CFU}(t = 0)}.$$

Developmental timing determination

Axenic adults were placed in sterile breeding cages overnight to lay eggs on

sterile HD. The HD used to collect embryos always matched the experimental

condition. Fresh axenic embryos were collected the next morning and seeded by pools

of 40 in tubes containing the HD to test. For the mono-associated conditions a total of

~10^7 CFUs of Ap^{WJL} or ~10^8 CFUs of Lp^{NC8}, washed on PBS, were inoculated on the

substrate and the eggs. Inoculation of Ap^{WJL} was limited to ~10^7 CFUs because higher

inoculums decreased egg-to-pupa survival. For heat killed (HK) conditions, washed

cells of Ap^{WJL} or Lp^{NC8} were incubated 3h at 65°C. Once at room temperature, embryos

were inoculated with ~10^8 CFUs and ~10^9 CFUs, respectively. In the germ-free

conditions, bacterial suspensions were replaced with sterile PBS. Tubes were

incubated at 25°C with 12/12-hour dark/light cycles. The emergence of pupae was

scored every day until all pupae had emerged. The experiment was stopped when no

pupae emerged after 30 days. Each gnotobiotic or nutritional condition was inoculated

in five replicates. Means, standard error of the mean and statistical tests (Dunn's test

of multiple comparisons) are detailed in Table S4. Since larvae are cannibalistic and

can find missing nutrients by eating their siblings [108,109] therefore we excluded

replicates with low egg-to-pupa survival (<25%,i.e n<10). Moreover, we considered

that larvae failed to develop in one condition if the mean egg-to-pupa survival of the

five replicates was inferior to 25% (for details in egg-to-pupae survival, see Table S5).

D_{50} was determined using D50App (http://paulinejoncour.shinyapps.io/D50App) as

described previously [23]. The whole experiment was repeated at least twice. D_{50}

heatmap represent the average of the five replicates of each gnotobiotic and nutritional

condition. Fig 4A was done using the imagesc function on MATLAB (version 2016b.

MathWorks, Natick, Massachusetts).

DNA extraction and AsnS locus analyses

Genomic DNA from 2 adults *yw* flies was extracted as previously described [110].

Briefly, flies were grinded in microtubes containing 0.75–1 mm glass microbeads and

500 µL of lysis buffer (Tris-HCl 10 mM, EDTA 1 mM, NaCl 1 mM at pH 8.2) using the

Precellys 24 tissue homogenizer (Bertin Technologies, Montigny-le-Bretonneux,

France). Then, we added Proteinase K (PureLink Genomic DNA extraction kit,

Invitrogen) at a final concentration of 200 µg/mL and incubated the samples at 56°C

under 700 rpm agitation for 1h. The samples were centrifuged at 10000 g for 2 min

and we collected the supernatant. *AsnS* coding sequence was amplified by PCR (Q5

Pol High Fidelity M0491S, New England BioLabs) using the primers AsnS_F

(CGGGCCGCTTCGTTAAAAA) and AsnS_R (TGGAATTCCTCAGACTTGCCA) with

a Veriti Thermal Cycler (Applied BioSystems, USA). PCR products were purified using

the NucleoSpin Gel and PCR Cleanup kit (Macherey-Nagel, Germany) following

manufacturer's instructions. Sequencing was done by Sanger sequencing (Genewiz,

Germany) using the following primers: AsnS_F, AsnS_R, AsnS1

(AGGATTATGGAAAGGATCTTCTGCA), AsnS2 (CTCCGGTCGGATTTGCATCA),

AsnS3 (TAATGCCAAAGGGGTCTCGG) and AsnS4

(GTGCGCCAGCTGCATTTATC). The whole coding sequence was then assembled

and analyzed using Geneious (version 10.1.3. Biomatters Ltd. New Zealand) by

mapping on the reference *D. melanogaster* genome (RefSeq GCF_000001215.4

release 6).

Amino acid quantification by HPLC

After growth in rich media, PBS washed Ap^{WJL} or Lp^{NC8} were inoculated in

triplicates at a final concentration of ~10^6 CFU/mL into 10 mL of each liquid HDΔEAA^{Fly}

shown to support their growth (Fig 1A, B) and in which they fulfill larval requirements.

For Ap^{WJL} this includes liquid HDΔArg, HDΔHis, HDΔIle, HDΔLeu, HDΔLys, HDΔMet,

HDΔPhe, HDΔThr, HDΔVal in agitated conditions. For Lp^{NC8} this includes liquid

HDΔHis, HDΔLys and HDΔMet in agitated conditions and liquid HDΔHis, HDΔLys,

HDΔMet, HDΔPhe and HDΔThr in static conditions. For agitated conditions cultures

were done in 50 mL flasks and incubated at 30°C under 180 rpm. Static conditions

were performed in 15-mL falcon tubes at 30°C. Samples were taken at time 0h and

72h. Samples were centrifuged (5000 rpm, 5 min). Supernatants were collected and

stored at -20°C until use.

Amino acid quantification was performed by HPLC from the supernatants

obtained at 0h and 72h. Samples were crushed in 320 µl of ultra-pure water with a

known quantity of norvaline used as the internal standard. Each sample was submitted

to a classical protein hydrolysis in sealed glass tubes with Teflon-lined screw caps (6N

HCl, 115°C, during 22h). After air vacuum removal, tubes were purged with nitrogen.

All samples were stored at -20°C, and then mixed with 50 µL of ultra-pure water for

amino acids analyses. Amino acid analysis was performed by HPLC (Agilent 1100;

Agilent Technologies, Massy, France) with a guard cartridge and a reverse phase C18

column (Zorbax Eclipse-AAA 3.5 µm, 150 × 4.6 mm, Agilent Technologies). Prior to

injection, the sample was buffered with borate at pH 10.2, and primary or secondary

amino acids were derivatized with ortho-phthalaldehyde (OPA) or 9-fluorenylmethyl

chloroformate (FMOC), respectively. The derivatization process, at room temperature,

was automated using the Agilent 1313A autosampler. Separation was carried out at

40°C, with a flow rate of 2 mL/min, using 40 mM NaH_2PO_4 (eluent A, pH 7.8, adjusted

with NaOH) as the polar phase and an acetonitrile/methanol/water mixture (45/45/10,

v/v/v) as the non-polar phase (eluent B). A gradient was applied during

chromatography, starting with 20% of B and increasing to 80% at the end. Detection

was performed by a fluorescence detector set at 340 and 450 nm of excitation and

emission wavelengths, respectively (266/305 nm for proline). These conditions do not

allow for the detection and quantification of cysteine and tryptophan, so only 18 amino

acids were quantified. For this quantification, norvaline was used as the internal

standard and the response factor of each amino acid was determined using a 250

pmol/µl standard mix of amino acids. The software used was the ChemStation for LC

3D Systems (Agilent Technologies).

Acknowledgements

We would like to thank Dali Ma for critical reading and editing of the manuscript

and valuable suggestions, members of the Ribeiro and Piper labs for advice on how

implementing effectively holidic diets preparations, the ArthroTools platform of the SFR

Biosciences (UMS3444/US8) for fly equipment and facility and the Bloomington Stock

Centre for fly lines. Research in F.L's lab is supported by the ENS de Lyon, CNRS and

the FINOVI foundation. Research in P.d.S and F.C labs are supported by INRA and

INSA Lyon. J.C is funded by a postdoctoral fellowship from the "Fondation pour la

Recherche Médicale" (FRM, SPF20170938612). T.G is funded by a Ph.D fellowship

from ENS de Lyon.

Figure legends

Fig 1. *In silico* **metabolic network reconstruction of Dm, Ap^{WJL} and Lp^{NC8}**

(A) Amino acids biosynthetic pathways. (B) Vitamins and cofactors biosynthetic

pathways. Left panels, *Drosophila melanogaster*. Central panels, *Acetobacter*

pomorum^{WJL}. Right panels, *Lactobacillus plantarum^{NC8}*. Color code: blue,

biosynthesized amino acid or vitamin; brown, limited amino acid or vitamin

biosynthesis (biosynthesis of the metabolite may be possible, but it is limited and/or

requires secondary metabolic pathways); black, non-biosynthesized amino acid or

vitamin; grey, pathway intermediary metabolite. Red cross: non-functional pathway

(lack of key enzyme(s)). Orange nods, major metabolic pathways. Abbreviations: α-

cglu: α-keto-glutarate. AceCoA: Acetyl-CoA. Ant: Antranilate. Aro: Arogenate. Cho:

chorismate. Cit: Citrate. Cysta: Cystathionine. Dihyn-P3: 7,8-Dihydroneopterin-3'-P3.

Dm-ribi: 6,7-Dimethyl-8-ribityllumazine. Ery-4P: Erythrose-4P. Fum: Fumarate. Glc:

Glucose. Gly-3P: Glycerate-3P. Homocys: Homocysteine. Homoser: Homoserine. Ind:

Indole. Orn: Ornithine. Oxa: Oxaloacetate. P-ra-imi: 1-(5'-Phospho-ribosyl)-5-

aminoimidazole. Phoser: Phosphoserine. Pre: Prephenate. Pyn-P: Pyridoxine

phosphate. Pyr: Pyruvate. Rib-5P: Ribose-5P.

Fig 2. Dm, Ap^{WJL} and Lp^{NC8} have differential biosynthetic capacities of nutrients

contained in the holidic diet

Venn diagram represents the number of HD nutrients that can be synthesized by each

organism. The list of corresponding metabolites is provided. Green dotted circles:

biosynthesis of this metabolite by Lp^{NC8} may be possible, but it is limited and/or

requires secondary metabolic pathways.

Fig 3. Ap^{WJL} and Lp^{NC8} auxotrophies detected in liquid fly Holidic Diet (HD)

(A) Heat map representing the mean maximal optical density (OD$_{Max}$) reached by

Ap^{WJL} or Lp^{NC8} after 72h of culture. Each line shows growth in a different version of

liquid HD: complete HD (first line) or HD lacking nutrient X (ΔX, lines below). EAA^{Fly}:

Fly's Essential Amino Acids, NEAA^{Fly}: Fly's Non-Essential Amino Acids, NAL: Nucleic

Acids and Lipids. Cultures were made in 96-wells plate under agitation. Asterisks (*)

pinpoint contradictions with *in silico* analysis that are explained in panel B. (B) Net

growth of Lp^{NC8} in four versions of liquid HD: complete HD, HDΔThr, HDΔAla and

HDΔAsp, in static conditions. Plot shows means with standard error based on three

replicates by assay. Each dot represents an independent replicate.

Fig 4. Ap^WJL and Lp^NC8 can differentially fulfill their host's nutritional

requirements

(A) Heat map representing the mean D_{50} (*i.e.* day when 50% of the larvae population

has entered metamorphosis) of GF larvae (first column) and larvae associated with

Ap^WJL, Lp^NC8, Ap^WJL_HK and Lp^NC8_HK (columns 2, 3, 4, 5 respectively). Each line shows

D_{50} in a different version of HD: complete HD (first line) or HD lacking nutrient X (ΔX,

lines below). EAA^Fly: Fly's Essential Amino Acids, NEAA^Fly: Fly's Non-Essential Amino

Acids, NAL: Nucleic Acids and Lipids. White means larvae did not reach pupariation in

these conditions. Means, standard errors of the mean and statistical tests (Dunn's test

of multiple comparisons) are detailed in Table S4. (B) D_{50} of *yw*, DGRP-*RAL*-25210

and *w*^1118 larvae on HDΔAsn. Boxplots show minimum, maximum and median. Each

dot shows an independent replicate. GF *yw* larvae did not reach pupariation. For the

other two lines, we performed Kruskal-Wallis test followed by post-hoc Dunn's tests to

compare each gnotobiotic condition to GF. ns: non-significant, **: p-value<0,005, ***:

p-value<0,0005, ****: p-value<0,0001. (C) Growth of Lp^NC8 in liquid HDΔPhe and liquid

HDΔCys, in static conditions, 3 days after inoculation. Plot shows mean with standard

error. Each dot shows an independent replicate. (D) Growth of Lp^NC8 in solid HDΔCys,

in absence and in presence of larvae (Medium Load and Niche Load, respectively), 3

days and 6 days after inoculation. Plot shows mean with standard error based on five

replicates by assay. Each dot represents an independent replicate. We performed two-

ways ANOVA followed by post-hoc Sidak's test. ns: non-significant, **: p-value<0,005.

Fig 5. Ap^WJL and Lp^NC8 can accumulate EAA^Fly during growth

(A) HPLC measured concentration of Arg, His, Ile, Leu, Phe, Thr and Val in the supernatant of an Ap^WJL culture in HDΔArg, HDΔHis, HDΔIle, HDΔLeu, HDΔPhe, HDΔThr, HDΔVal, respectively, 72h after inoculation. Plot shows mean with standard error. Each dot shows an independent replicate. (B) HPLC measured concentration of His in the supernatant of a Lp^NC8 culture in HDΔHis, 72h after inoculation. Plot shows mean with standard error. Each dot shows an independent replicate.

Fig 6. Commensal bacterial differentially shape the nutritional requirements of their juvenile host

For each gnotobiotic condition, essential nutrients are represented in black and non-essential nutrients in color. Color code: blue, this nutrient can be synthesized by the bacteria; red: this nutrient cannot be synthesized by the bacteria, suggesting a mechanism of functional compensation. In purple: this nutrient may be synthesized by the bacteria through still unknown metabolic pathways.

Supporting Information

Table S1: Inference from genomic analysis of the biosynthetic capabilities for amino acids production in *D. melanogaster, A. pomorum*^WJL and *L. plantarum*^NC8. f / i: targeted amino acid biosynthesis is feasible / impossible in a depleted medium.

Table S2: Inference from genomic analysis of the biosynthetic capabilities for vitamins production in *D. melanogaster, A. pomorum*^WJL and *L. plantarum*^NC8. f / i: targeted vitamin biosynthesis is feasible / impossible in a depleted medium.

Table S3: OD_{Max} of Ap^{WJL} and Lp^{NC8} grown in 39 HDs. Mean and Standard Error to the Mean (SEM) of OD_{Max} reached by Ap^{WJL} or Lp^{NC8} grown in complete liquid HD (first line) or liquid HD lacking nutrient X (ΔX, lines below) during 72h of growth.

Table S4: D_{50} of larvae in 40 HDs and 5 gnotobiotic conditions. Mean and Standard Error of the Mean (SEM) of D_{50} (*i.e.* day when 50% of the larvae population has entered metamorphosis) of GF larvae or larvae associated with Ap^{WJL}, Lp^{NC8}, Ap^{WJL}_{HK} and Lp^{NC8}_{HK}. n: number of independent replicates for each condition. For each gnotobiotic condition, we performed Kruskal-Wallis test followed by post-hoc Dunn's test to compare each nutritional environment to complete HD.

Table S5: Egg-to-pupa survival in 40 HDs and 5 gnotobiotic conditions. Mean and Standard Error of Mean (SEM) of egg-to-pupa survival of GF larvae or larvae associated with Ap^{WJL}, Lp^{NC8}, Ap^{WJL}_{HK} and Lp^{NC8}_{HK}. n: number of independent replicates for each condition. For each gnotobiotic condition, we performed Kruskal-Wallis test followed by post-hoc Dunn's test to compare each nutritional environment to complete HD.

Table S6: Comparison of *in silico* predictions with *in vivo* auxotrophies and bacterial complementation of larval nutritional deficiencies. Can partner A synthesize nutrient X?: prediction from *in silico* metabolic reconstruction (from Fig 1, Table S1 and Table S2). Can partner A grow in the absence of nutrient X?: auxotrophy observed *in vivo* (from Fig 3A-B). Can bacterial partner A promote larval growth on HD ΔX?: *in vivo* complementation of ex-GF larvae requirements (from Fig 4A), y: yes (green), n: no

1004 (red). Hypothesis to explain contradiction: why the different approaches do not always

1005 lead to the same conclusion. NA: Non-applicable.

1006

1007 **References**

1008

1. Lifshitz F. Nutrition and Growth. J Clin Res Pediatr Endocrinol. 2009;1: 157–163.
doi:10.4274/jcrpe.v1i4.39

2. Shanley DP, Kirkwood TBL. Calorie restriction and aging: a life-history analysis.
Evolution. 2000;54: 740–750. doi:10.1111/j.0014-3820.2000.tb00076.x

3. Chipponi JX, Bleier JC, Santi MT, Rudman D. Deficiencies of essential and
conditionally essential nutrients. Am J Clin Nutr. 1982;35: 1112–1116.
doi:10.1093/ajcn/35.5.1112

4. Fairweather-Tait SJ, Southon S. Bioavailability of nutrients. In: Caballero B, editor.
Encyclopedia of Food Sciences and Nutrition (Second Edition). Oxford: Academic
Press; 2003. pp. 478–484. doi:10.1016/B0-12-227055-X/00096-1

5. Stewart CP, Iannotti L, Dewey KG, Michaelsen KF, Onyango AW. Contextualising
complementary feeding in a broader framework for stunting prevention. Matern
Child Nutr. 2013;9: 27–45. doi:10.1111/mcn.12088

6. Smith MI, Yatsunenko T, Manary MJ, Trehan I, Mkakosya R, Cheng J, et al. Gut
microbiomes of Malawian twin pairs discordant for kwashiorkor. Science.
2013;339: 548–554. doi:10.1126/science.1229000

7. Gehrig JL, Venkatesh S, Chang H-W, Hibberd MC, Kung VL, Cheng J, et al.
Effects of microbiota-directed foods in gnotobiotic animals and undernourished
children. Science. 2019;365. doi:10.1126/science.aau4732

8. Tennessen JM, Thummel CS. Coordinating growth and review maturation —
insights from *Drosophila*. Curr Biol CB. 2011;21: R750–R757.
doi:10.1016/j.cub.2011.06.033

9. Douglas AE. The *Drosophila* model for microbiome research. Lab Anim. 2018;47:
157–164. doi:10.1038/s41684-018-0065-0

10. Erkosar B, Storelli G, Defaye A, Leulier F. Host-intestinal microbiota
mutualism: "learning on the fly." Cell Host Microbe. 2013;13: 8–14.

doi:10.1016/j.chom.2012.12.004

11.   Lee K-A, Lee W-J. *Drosophila* as a model for intestinal dysbiosis and chronic

1037   inflammatory diseases. Dev Comp Immunol. 2014;42: 102–110.

1038   doi:10.1016/j.dci.2013.05.005

1039   12.   Broderick NA, Lemaitre B. Gut-associated microbes of *Drosophila*

1040   *melanogaster*. Gut Microbes. 2012;3: 307–321. doi:10.4161/gmic.19896

1041   13.   Bakula M. The persistence of a microbial flora during postembryogenesis of

1042   *Drosophila melanogaster*. J Invertebr Pathol. 1969;14: 365–374.

1043   14.   Koyle ML, Veloz M, Judd AM, Wong AC-N, Newell PD, Douglas AE, et al.

1044   Rearing the fruit fly *Drosophila melanogaster* under axenic and gnotobiotic

1045   conditions. J Vis Exp JoVE. 2016; doi:10.3791/54219

1046   15.   Ma D, Storelli G, Mitchell M, Leulier F. Studying host-microbiota mutualism in

1047   *Drosophila*: harnessing the power of gnotobiotic flies. Biomed J. 2015;38: 285–

1048   293. doi:10.4103/2319-4170.158620

1049   16.   Brummel T, Ching A, Seroude L, Simon AF, Benzer S. *Drosophila* lifespan

1050   enhancement by exogenous bacteria. Proc Natl Acad Sci. 2004;101: 12974–

1051   12979. doi:10.1073/pnas.0405207101

1052   17.   Cox CR, Gilmore MS. Native microbial colonization of *Drosophila*

1053   *melanogaster* and its use as a model of *Enterococcus faecalis* pathogenesis.

1054   Infect Immun. 2007;75: 1565–1576. doi:10.1128/IAI.01496-06

1055   18.   Ren C, Webster P, Finkel SE, Tower J. Increased internal and external

1056   bacterial load during *Drosophila* aging without life-span trade-off. Cell Metab.

1057   2007;6: 144–152. doi:10.1016/j.cmet.2007.06.006

1058   19.   Ryu J-H, Kim S-H, Lee H-Y, Bai JY, Nam Y-D, Bae J-W, et al. Innate immune

1059   homeostasis by the homeobox gene *Caudal* and commensal-gut mutualism in

1060   *Drosophila*. Science. 2008;319: 777–782. doi:10.1126/science.1149357

1061   20.   Storelli G, Defaye A, Erkosar B, Hols P, Royet J, Leulier F. *Lactobacillus*

1062   *plantarum* promotes *Drosophila* systemic growth by modulating hormonal signals

1063   through TOR-dependent nutrient sensing. Cell Metab. 2011;14: 403–414.

1064   doi:10.1016/j.cmet.2011.07.012

1065   21.   Adair KL, Wilson M, Bost A, Douglas AE. Microbial community assembly in

1066   wild populations of the fruit fly *Drosophila melanogaster*. ISME J. 2018;12: 959.

1067   doi:10.1038/s41396-017-0020-x

1068   22.   Pais IS, Valente RS, Sporniak M, Teixeira L. *Drosophila melanogaster*

1069 establishes a species-specific mutualistic interaction with stable gut-colonizing
1070 bacteria. PLoS Biol. 2018;16. doi:10.1371/journal.pbio.2005710

1071 23. Matos RC, Schwarzer M, Gervais H, Courtin P, Joncour P, Gillet B, et al. D-
1072 Alanylation of teichoic acids contributes to *Lactobacillus plantarum*-mediated
1073 *Drosophila* growth during chronic undernutrition. Nat Microbiol. 2017;2: 1635–
1074 1647. doi:10.1038/s41564-017-0038-x

1075 24. Schretter CE, Vielmetter J, Bartos I, Marka Z, Marka S, Argade S, et al. A gut
1076 microbial factor modulates locomotor behaviour in *Drosophila*. Nature. 2018;563:
1077 402. doi:10.1038/s41586-018-0634-9

1078 25. Shin SC, Kim S-H, You H, Kim B, Kim AC, Lee K-A, et al. *Drosophila*
1079 microbiome modulates host developmental and metabolic homeostasis via insulin
1080 signaling. Science. 2011;334: 670–674. doi:10.1126/science.1212782

1081 26. Blum JE, Fischer CN, Miles J, Handelsman J. Frequent replenishment
1082 sustains the beneficial microbiome of *Drosophila melanogaster*. mBio. 2013;4:
1083 e00860-00813. doi:10.1128/mBio.00860-13

1084 27. Broderick NA, Buchon N, Lemaitre B. Microbiota-induced changes in
1085 *Drosophila melanogaster* host gene expression and gut morphology. mBio.
1086 2014;5. doi:10.1128/mBio.01117-14

1087 28. Winans NJ, Walter A, Chouaia B, Chaston JM, Douglas AE, Newell PD. A
1088 genomic investigation of ecological differentiation between free-living and
1089 *Drosophila*-associated bacteria. Mol Ecol. 2017;26: 4536–4550.
1090 doi:10.1111/mec.14232

1091 29. Obadia B, Güvener ZT, Zhang V, Ceja-Navarro JA, Brodie EL, Ja WW, et al.
1092 Probabilistic invasion underlies natural gut microbiome stability. Curr Biol CB.
1093 2017;27: 1999-2006.e8. doi:10.1016/j.cub.2017.05.034

1094 30. Storelli G, Strigini M, Grenier T, Bozonnet L, Schwarzer M, Daniel C, et al.
1095 *Drosophila* perpetuates nutritional mutualism by promoting the fitness of its
1096 intestinal symbiont *Lactobacillus plantarum*. Cell Metab. 2018;27: 362-377.e8.
1097 doi:10.1016/j.cmet.2017.11.011

1098 31. Ma D, Leulier F. The importance of being persistent: The first true resident gut
1099 symbiont in Drosophila. PLoS Biol. 2018;16: e2006945.
1100 doi:10.1371/journal.pbio.2006945

1101 32. Erkosar B, Storelli G, Mitchell M, Bozonnet L, Bozonnet N, Leulier F. Pathogen
1102 virulence impedes mutualist-mediated enhancement of host juvenile growth via

inhibition of protein digestion. Cell Host Microbe. 2015;18: 445–455.

doi:10.1016/j.chom.2015.09.001

33.   Sannino DR, Dobson AJ, Edwards K, Angert ER, Buchon N. The *Drosophila*

melanogaster gut microbiota provisions thiamine to its host. mBio. 2018;9.

doi:10.1128/mBio.00155-18

34.   Piper MD. Using artificial diets to understand the nutritional physiology of

Drosophila melanogaster. Curr Opin Insect Sci. 2017;23: 104–111.

doi:10.1016/j.cois.2017.07.014

35.   Staats S, Lüersen K, Wagner AE, Rimbach G. *Drosophila melanogaster* as a

versatile model organism in food and nutrition research. J Agric Food Chem.

2018;66: 3737–3753. doi:10.1021/acs.jafc.7b05900

36.   Lüersen K, Röder T, Rimbach G. *Drosophila melanogaster* in nutrition

research—the importance of standardizing experimental diets. Genes Nutr.

2019;14. doi:10.1186/s12263-019-0627-9

37.   Schultz J, St Lawrence P, Newmeyer D. A chemically defined medium for the

growth of *Drosophila melanogaster*. Anat Rec. 1946;96: 540.

38.   Sang JH. The quantitative nutritional requirements of *Drosophila*

melanogaster. J Exp Biol. 1956;33: 45–72.

39.   Hinton T, Noyes DT, Ellis J. Amino acids and growth factors in a chemically

defined medium for *Drosophila*. Physiol Zool. 1951;24: 335–353.

40.   Piper MDW, Soultoukis GA, Blanc E, Mesaros A, Herbert SL, Juricic P, et al.

Matching dietary amino acid balance to the *in silico*-translated exome optimizes

growth and reproduction without cost to lifespan. Cell Metab. 2017;25: 610–621.

doi:10.1016/j.cmet.2017.02.005

41.   Jang T, Lee KP. Comparing the impacts of macronutrients on life-history traits

in larval and adult *Drosophila melanogaster*: the use of nutritional geometry and

chemically defined diets. J Exp Biol. 2018; jeb.181115. doi:10.1242/jeb.181115

42.   Piper MDW, Blanc E, Leitão-Gonçalves R, Yang M, He X, Linford NJ, et al. A

holidic medium for *Drosophila melanogaster*. Nat Methods. 2014;11: 100–105.

doi:10.1038/nmeth.2731

43.   Troen AM, French EE, Roberts JF, Selhub J, Ordovas JM, Parnell LD, et al.

Lifespan modification by glucose and methionine in *Drosophila melanogaster* fed a

chemically defined diet. Age Dordr Neth. 2007;29: 29–39. doi:10.1007/s11357-

006-9018-4

44.   Sang JH, King RC. Nutritional requirements of axenically cultured *Drosophila*

melanogaster adults. J Exp Biol. 1961;38: 793–809.

45.   Leitão-Gonçalves R, Carvalho-Santos Z, Francisco AP, Fioreze GT, Anjos M,

Baltazar C, et al. Commensal bacteria and essential amino acids control food

choice behavior and reproduction. PLOS Biol. 2017;15: e2000862.

doi:10.1371/journal.pbio.2000862

46.   Mishra D, Thorne N, Miyamoto C, Jagge C, Amrein H. The taste of

ribonucleosides: Novel macronutrients essential for larval growth are sensed by

Drosophila gustatory receptor proteins. PLOS Biol. 2018;16: e2005570.

doi:10.1371/journal.pbio.2005570

47.   Geer BW. Utilization of D-amino acids for growth by *Drosophila melanogaster*

larvae. J Nutr. 1966;90: 31–39. doi:10.1093/jn/90.1.31

48.   Lee W-C, Micchelli CA. Development and characterization of a chemically

defined food for *Drosophila*. PLOS ONE. 2013;8: e67308.

doi:10.1371/journal.pone.0067308

49.   Rapport EW, Stanley-Samuelson D, Dadd RH. Ten generations of *Drosophila*

melanogaster reared axenically on a fatty acid-free holidic diet. Arch Insect

Biochem Physiol. 1983;1: 243–250. doi:10.1002/arch.940010307

50.   i5K Consortium. The i5K Initiative: advancing arthropod genomics for

knowledge, human health, agriculture, and the environment. J Hered. 2013;104:

595–600. doi:10.1093/jhered/est050

51.   Baa-Puyoulet P, Parisot N, Febvay G, Huerta-Cepas J, Vellozo AF, Gabaldón

T, et al. ArthropodaCyc: a CycADS powered collection of BioCyc databases to

analyse and compare metabolism of arthropods. Database J Biol Databases

Curation. 2016;2016. doi:10.1093/database/baw081

52.   Cottret L, Milreu PV, Acuña V, Marchetti-Spaccamela A, Stougie L, Charles H,

et al. Graph-based analysis of the metabolic exchanges between two co-resident

intracellular symbionts, *Baumannia cicadellinicola* and *Sulcia muelleri*, with their

insect host, *Homalodisca coagulata*. PLoS Comput Biol. 2010;6.

doi:10.1371/journal.pcbi.1000904

53.   Magnúsdóttir S, Heinken A, Kutt L, Ravcheev DA, Bauer E, Noronha A, et al.

Generation of genome-scale metabolic reconstructions for 773 members of the

human gut microbiota. Nat Biotechnol. 2017;35: 81–89. doi:10.1038/nbt.3703

54.   Ankrah NYD, Chouaia B, Douglas AE. The cost of metabolic interactions in

symbioses between Insects and Bacteria with reduced genomes. mBio. 2018;9. doi:10.1128/mBio.01433-18

55.   Opatovsky I, Santos-Garcia D, Ruan Z, Lahav T, Ofaim S, Mouton L, et al. Modeling trophic dependencies and exchanges among insects' bacterial symbionts in a host-simulated environment. BMC Genomics. 2018;19: 402. doi:10.1186/s12864-018-4786-7

56.   Wilson ACC, Ashton PD, Calevro F, Charles H, Colella S, Febvay G, et al. Genomic insight into the amino acid relations of the pea aphid, *Acyrthosiphon pisum*, with its symbiotic bacterium *Buchnera aphidicola*. Insect Mol Biol. 2010;19 Suppl 2: 249–258. doi:10.1111/j.1365-2583.2009.00942.x

57.   Russell CW, Bouvaine S, Newell PD, Douglas AE. Shared metabolic pathways in a coevolved insect-bacterial symbiosis. Appl Environ Microbiol. 2013;79: 6117–6123. doi:10.1128/AEM.01543-13

58.   Viñuelas J, Febvay G, Duport G, Colella S, Fayard J-M, Charles H, et al. Multimodal dynamic response of the *Buchnera aphidicola* pLeu plasmid to variations in leucine demand of its host, the pea aphid *Acyrthosiphon pisum*. Mol Microbiol. 2011;81: 1271–1285. doi:10.1111/j.1365-2958.2011.07760.x

59.   Rabatel A, Febvay G, Gaget K, Duport G, Baa-Puyoulet P, Sapountzis P, et al. Tyrosine pathway regulation is host-mediated in the pea aphid symbiosis during late embryonic and early larval development. BMC Genomics. 2013;14: 235. doi:10.1186/1471-2164-14-235

60.   Simonet P, Gaget K, Parisot N, Duport G, Rey M, Febvay G, et al. Disruption of phenylalanine hydroxylase reduces adult lifespan and fecundity, and impairs embryonic development in parthenogenetic pea aphids. Sci Rep. 2016;6: 34321. doi:10.1038/srep34321

61.   Hansen AK, Moran NA. The impact of microbial symbionts on host plant utilization by herbivorous insects. Mol Ecol. 2014;23: 1473–1496. doi:10.1111/mec.12421

62.   Vellozo AF, Véron AS, Baa-Puyoulet P, Huerta-Cepas J, Cottret L, Febvay G, et al. CycADS: an annotation database system to ease the development and update of BioCyc databases. Database J Biol Databases Curation. 2011;2011: bar008. doi:10.1093/database/bar008

63.   Rao MRR, Stokes JL. Nutrition of the acetic acid bacteria. J Bacteriol. 1953;65: 405–412.

64. Martino ME, Bayjanov JR, Caffrey BE, Wels M, Joncour P, Hughes S, et al.
Nomadic lifestyle of *Lactobacillus plantarum* revealed by comparative genomics of
54 strains isolated from different habitats. Environ Microbiol. 2016;18: 4974–4989.
doi:10.1111/1462-2920.13455

65. Berndt C, Lillig CH, Wollenberg M, Bill E, Mansilla MC, de Mendoza D, et al.
Characterization and reconstitution of a 4Fe-4S adenylyl sulfate/phosphoadenylyl
sulfate reductase from *Bacillus subtilis*. J Biol Chem. 2004;279: 7850–7855.
doi:10.1074/jbc.M309332200

66. Teusink B, van Enckevort FHJ, Francke C, Wiersma A, Wegkamp A, Smid EJ,
et al. *In silico* reconstruction of the metabolic pathways of *Lactobacillus plantarum*:
comparing predictions of nutrient requirements with those from growth
experiments. Appl Environ Microbiol. 2005;71: 7253–7262.
doi:10.1128/AEM.71.11.7253-7262.2005

67. Saguir FM, de Nadra MCM. Improvement of a chemically defined medium for
the sustained growth of *Lactobacillus plantarum*: nutritional requirements. Curr
Microbiol. 2007;54: 414–418. doi:10.1007/s00284-006-0456-0

68. Bringel F, Hubert J-C. Extent of genetic lesions of the arginine and pyrimidine
biosynthetic pathways in *Lactobacillus plantarum*, *L. paraplantarum*, *L. pentosus*,
and *L. casei*: prevalence of CO(2)-dependent auxotrophs and characterization of
deficient arg genes in *L. plantarum*. Appl Environ Microbiol. 2003;69: 2674–2683.
doi:10.1128/aem.69.5.2674-2683.2003

69. Bringel F, Frey L, Boivin S, Hubert JC. Arginine biosynthesis and regulation in
Lactobacillus plantarum: the *carA* gene and the *argCJBDF* cluster are divergently
transcribed. J Bacteriol. 1997;179: 2697–2706.

70. Matsushita K, Toyama H, Tonouchi N, Okamoto-Kainuma A, editors. Acetic
acid bacteria: ecology and physiology [Internet]. Springer Japan; 2016. Available:
https://www.springer.com/gp/book/9784431559313

71. Hanada T, Kashima Y, Kosugi A, Koizumi Y, Yanagida F, Udaka S. A gene
encoding phosphatidylethanolamine N-methyltransferase from *Acetobacter aceti*
and some properties of its disruptant. Biosci Biotechnol Biochem. 2001;65: 2741–
2748. doi:10.1271/bbb.65.2741

72. Sohlenkamp C, Geiger O. Bacterial membrane lipids: diversity in structures
and pathways. FEMS Microbiol Rev. 2016;40: 133–159.
doi:10.1093/femsre/fuv008

73.   Chandrangsu P, Rensing C, Helmann JD. Metal homeostasis and resistance

in bacteria. Nat Rev Microbiol. 2017;15: 338–350. doi:10.1038/nrmicro.2017.15

74.   Rosenberg E, DeLong EF, Lory S, Stackebrandt E, Thompson F, editors. The

Prokaryotes: *Alphaproteobacteria* and *Betaproteobacteria* [Internet]. 4th ed. Berlin

Heidelberg: Springer-Verlag; 2014. Available:

https://www.springer.com/gp/book/9783642301964

75.   Siezen RJ, Tzeneva VA, Castioni A, Wels M, Phan HTK, Rademaker JLW, et

al. Phenotypic and genomic diversity of *Lactobacillus plantarum* strains isolated

from various environmental niches. Environ Microbiol. 2010;12: 758–773.

doi:10.1111/j.1462-2920.2009.02119.x

76.   Teusink B, Wiersma A, Molenaar D, Francke C, Vos WM de, Siezen RJ, et al.

Analysis of growth of *Lactobacillus plantarum* WCFS1 on a complex medium using

a genome-scale metabolic model. J Biol Chem. 2006;281: 40041–40048.

doi:10.1074/jbc.M606263200

77.   Wegkamp A, Teusink B, De Vos W m., Smid E j. Development of a minimal

growth medium for *Lactobacillus plantarum*. Lett Appl Microbiol. 2010;50: 57–64.

doi:10.1111/j.1472-765X.2009.02752.x

78.   Groot MNN, Klaassens E, de Vos WM, Delcour J, Hols P, Kleerebezem M.

Genome-based in silico detection of putative manganese transport systems in

Lactobacillus plantarum and their genetic analysis. Microbiology. 2005;151: 1229–

1238. doi:10.1099/mic.0.27375-0

79.   Vos P, Garrity G, Jones D, Krieg NR, Ludwig W, Rainey FA, et al., editors.

Bergey's Manual of Systematic Bacteriology: Volume 3: The Firmicutes [Internet].

2nd ed. New York: Springer-Verlag; 2009. Available:

https://www.springer.com/gp/book/9780387950419

80.   Ikawa M. Nature of the lipids of some lactic acid bacteria. J Bacteriol. 1963;85:

772–781.

81.   Patterson MK, Orr GR. Asparagine biosynthesis by the Novikoff Hepatoma

isolation, purification, property, and mechanism studies of the enzyme system. J

Biol Chem. 1968;243: 376–380.

82.   Mackay TFC, Richards S, Stone EA, Barbadilla A, Ayroles JF, Zhu D, et al.

The *Drosophila melanogaster* Genetic Reference Panel. Nature. 2012;482: 173–

178. doi:10.1038/nature10811

83.   Wong AC-N, Dobson AJ, Douglas AE. Gut microbiota dictates the metabolic

response of *Drosophila* to diet. J Exp Biol. 2014;217: 1894–1901.

doi:10.1242/jeb.101725

84.   Huang Z, London E. Cholesterol lipids and cholesterol-containing lipid rafts in

bacteria. Chem Phys Lipids. 2016;199: 11–16.

doi:10.1016/j.chemphyslip.2016.03.002

85.   Hammad LA, Cooper BS, Fisher NP, Montooth KL, Karty JA. Profiling and

quantification of *Drosophila melanogaster* lipids using liquid chromatography/mass

spectrometry: Lipid composition of *Drosophila melanogaster*. Rapid Commun

Mass Spectrom. 2011;25: 2959–2968. doi:10.1002/rcm.5187

86.   Jones HE, Harwood JL, Bowen ID, Griffiths G. Lipid composition of subcellular

membranes from larvae and prepupae of *Drosophila melanogaster*. Lipids.

1992;27: 984–987.

87.   Hadadi N, MohammadiPeyhani H, Miskovic L, Seijo M, Hatzimanikatis V.

Enzyme annotation for orphan and novel reactions using knowledge of substrate

reactive sites. Proc Natl Acad Sci U S A. 2019;116: 7298–7307.

doi:10.1073/pnas.1818877116

88.   Navarro JA, Schneuwly S. Copper and zinc homeostasis: lessons from

Drosophila melanogaster. Front Genet. 2017;8. doi:10.3389/fgene.2017.00223

89.   Combe BE, Defaye A, Bozonnet N, Puthier D, Royet J, Leulier F. *Drosophila*

microbiota modulates host metabolic gene expression via IMD/NF-κB signaling.

PLoS ONE. 2014;9. doi:10.1371/journal.pone.0094729

90.   Hennigar SR, Kelley AM, McClung JP. Metallothionein and zinc transporter

expression in circulating human blood cells as biomarkers of zinc status: a

systematic review. Adv Nutr. 2016;7: 735–746. doi:10.3945/an.116.012518

91.   Qi B, Han M. Microbial siderophore enterobactin promotes mitochondrial iron

uptake and development of the host via interaction with ATP synthase. Cell.

2018;175: 571-582.e11. doi:10.1016/j.cell.2018.07.032

92.   Goffin P, Bunt B van de, Giovane M, Leveau JHJ, Höppener-Ogawa S,

Teusink B, et al. Understanding the physiology of *Lactobacillus plantarum* at zero

growth. Mol Syst Biol. 2010;6: 413. doi:10.1038/msb.2010.67

93.   Typas A, Banzhaf M, Gross CA, Vollmer W. From the regulation of

peptidoglycan synthesis to bacterial growth and morphology. Nat Rev Microbiol.

2011;10: 123–136. doi:10.1038/nrmicro2677

94.   Boothby D, Daneo-Moore L, Higgins ML, Coyette J, Shockman GD. Turnover

of bacterial cell wall peptidoglycans. J Biol Chem. 1973;248: 2161–2169.

95.   Lam H, Oh D-C, Cava F, Takacs CN, Clardy J, de Pedro MA, et al. D-amino

acids govern stationary phase cell wall re-modeling in Bacteria. Science.

2009;325: 1552–1555. doi:10.1126/science.1178123

96.   Newell PD, Chaston JM, Wang Y, Winans NJ, Sannino DR, Wong ACN, et al.

In vivo function and comparative genomic analyses of the *Drosophila* gut

microbiota identify candidate symbiosis factors. Front Microbiol. 2014;5.

doi:10.3389/fmicb.2014.00576

97.   Moriya Y, Itoh M, Okuda S, Yoshizawa AC, Kanehisa M. KAAS: an automatic

genome annotation and pathway reconstruction server. Nucleic Acids Res.

2007;35: W182-185. doi:10.1093/nar/gkm321

98.   Claudel-Renard C, Chevalet C, Faraut T, Kahn D. Enzyme-specific profiles for

genome annotation: PRIAM. Nucleic Acids Res. 2003;31: 6633–6639.

doi:10.1093/nar/gkg847

99.   Conesa A, Götz S, García-Gómez JM, Terol J, Talón M, Robles M. Blast2GO:

a universal tool for annotation, visualization and analysis in functional genomics

research. Bioinforma Oxf Engl. 2005;21: 3674–3676.

doi:10.1093/bioinformatics/bti610

100.   Conesa A, Götz S. Blast2GO: A comprehensive suite for functional analysis in

plant genomics. Int J Plant Genomics. 2008;2008: 619832.

doi:10.1155/2008/619832

101.   Jones P, Binns D, Chang H-Y, Fraser M, Li W, McAnulla C, et al. InterProScan

5: genome-scale protein function classification. Bioinforma Oxf Engl. 2014;30:

1236–1240. doi:10.1093/bioinformatics/btu031

102.   Karp PD, Latendresse M, Paley SM, Krummenacker M, Ong QD, Billington R,

et al. Pathway Tools version 19.0 update: software for pathway/genome

informatics and systems biology. Brief Bioinform. 2016;17: 877–890.

doi:10.1093/bib/bbv079

103.   Karp PD, Weaver D, Paley S, Fulcher C, Kubo A, Kothari A, et al. The EcoCyc

Database. EcoSal Plus. 2014;6. doi:10.1128/ecosalplus.ESP-0009-2013

104.   Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment

search tool. J Mol Biol. 1990;215: 403–410. doi:10.1016/S0022-2836(05)80360-2

105.   Latendresse M, Krummenacker M, Karp PD. Optimal metabolic route search

based on atom mappings. Bioinforma Oxf Engl. 2014;30: 2043–2050.

doi:10.1093/bioinformatics/btu150

106.   Teixeira L, Ferreira Á, Ashburner M. The Bacterial Symbiont *Wolbachia*

Induces Resistance to RNA Viral Infections in *Drosophila melanogaster*. PLoS

Biol. 2008;6. doi:10.1371/journal.pbio.1000002

107.   Axelsson L, Rud I, Naterstad K, Blom H, Renckens B, Boekhorst J, et al.

Genome sequence of the naturally plasmid-free *Lactobacillus plantarum* strain

NC8 (CCUG 61730). J Bacteriol. 2012;194: 2391–2392. doi:10.1128/JB.00141-12

108.   Ahmad M, Chaudhary SU, Afzal AJ, Tariq M. Starvation-induced dietary

behaviour in *Drosophila melanogaster* larvae and adults. Sci Rep. 2015;5.

doi:10.1038/srep14285

109.   Vijendravarma RK, Narasimha S, Kawecki TJ. Predatory cannibalism in

Drosophila melanogaster larvae. Nat Commun. 2013;4: 1789.

doi:10.1038/ncomms2744

110.   Green MR, Sambrook J. Molecular Cloning: A Laboratory Manual, Fourth

Edition. 4th edition. Cold Spring Harbor, N.Y: Cold Spring Harbor Laboratory

Press,U.S.; 2012.

1357

Figure 1

Figure 2

Figure 3

Figure 4

A Microbial environment

B Asn essentiality is genotype-dependent

C Lp^NC8 growth in static conditions

D Lp^NC8 growth in solid HDΔCys

Figure 5

Figure 6