Mycobacterium riyadhense clinical isolates provide insights into ancestry

and adaptive evolution in tuberculosis

Qingtian Guan[1], Musa Garbati[2,a], Sara Mfarrej[1], Talal AlMutairi[2], Alicia Smyth[3&], Albel

Singh[4&], Shamsudeen Fagbo[5&], John A. Browne[3], Muhammad Amin urRahman[2], Alya

Alruwaili[2], Anwar Hoosen[2,b], Chie Nakajima[6,7], Yasuhiko Suzuki[6,7], Apoorva Bhatt[4],

Stephen V. Gordon[3,6], Faisal AlAsmari[2], A. Pain[1,6*]

[1]Pathogen Genomics Laboratory, BESE Division, King Abdullah University of Science

and Technology (KAUST), Thuwal-Jeddah, Kingdom of Saudi Arabia; [2]King Fahad

Medical City (KFMC), Riyadh, Saudi Arabia; [3]UCD School of Veterinary Medicine,

University College Dublin, Dublin, D04 W6F6, Ireland; [4]Institute of Microbiology and

Infection, School of Biosciences, University of Birmingham, Edgbaston, Birmingham,

UK;[5]One Health Unit, Executive Directorate for Surveillance and Response, National

Center for Disease Prevention and Control, Riyadh, Saudi Arabia; [6]Global Institution for

Collaborative Research and Education, Hokkaido University, Kita 20 Nishi 10, Kita-ku,

Sapporo, Japan;[7]Research Center for Zoonosis Control, Hokkaido University, Kita 20

Nishi 10, Kita-ku, Sapporo, Japan.

[a]Current address: Infectious Diseases Unit, Department of Medicine, University of

Maiduguri, Nigeria

[&]These authors contributed equally to the work

[b]Current address: Pathcare/Vermaak & Partners Pathologists, Pretoria, South Africa.

*For correspondence
Arnab Pain (arnab.pain@kaust.edu.sa)

Abstract

Mycobacterium riyadhense is a novel non-tuberculous mycobacterium (NTM) that is increasingly isolated from clinical cases with tuberculosis (TB)-like symptoms in humans in various parts of the world. To elucidate the genomic hallmarks that define the relationship between *M. riyadhense* and other mycobacterial species, including members of the *Mycobacterium tuberculosis* complex (MTBC), eight clinical isolates of *M. riyadhense* were analysed. *M. riyadhense* shares a large number of conserved orthologues with the MTBC and contains linear and circular plasmids carrying type IV and type VII secretion systems. We provide evidence that *M. riyadhense* is likely the 'missing link' in the evolution of *M. tuberculosis*, sharing a common ancestor with members of the MTBC that has evolved through a process of genome reduction, expansion of toxin/antitoxin systems, and ultimately host adaptation. We conclude that *M. riyadhense* is an emerging mycobacterial pathogen that can serve as an experimental model to study the evolution and pathogenesis of tubercle bacilli.

Introduction

Non-tuberculous mycobacteria (NTMs), including *Mycobacterium riyadhense* (MR), are ubiquitous, naturally occurring environmental bacteria commonly found in water and soil (Falkinham, 2013)(J. Van Ingen et al., 2009). A wide range of animal and environmental

sources (aquaria, swimming pools) act as reservoirs for NTMs, and several human

disease outbreaks caused by exposure to environmental NTMs have been described

(3)(4)(5)(6). With the ability to cause infections in both immunocompromised (7) and

immunocompetent   (8) individuals, *M. riyadhense* has positioned itself as a clinically

important pulmonary pathogen since its discovery in 2009 (9) as a cause of pulmonary

infection. The clinical and radiologic characteristics of pulmonary infection caused by *M.*

riyadhense are indistinguishable from those caused by *Mycobacterium tuberculosis*, the

most important human pathogen of the *M. tuberculosis* complex (MTBC), which cause

TB in humans and animals (8)(9).

Similar to *M. tuberculosis*, *M. riyadhense* grows at 37°C and requires 2~3 weeks (9) to

form visible colonies on agar media. However, unlike *M. tuberculosis*, which is a

common worldwide pathogen transmitted directly from human to human with no known

environmental reservoirs (10), *M. riyadhense* infections are rare and are transmitted to

patients via contact with contaminated water (10) and soil (11), with no evidence of

human-to-human transmission yet reported. Infections with *M. riyadhense* have been

reported in Asia and Europe including, Bahrain, South Korea, France, Italy and Germany

(12)(13), (13), (8), although most of the recent cases originated in patients from Saudi

Arabia (14). Indeed, the very first case of acute *M. riyadhense* infection was initially

misdiagnosed as a case of *M. tuberculosis* infection in a Saudi hospital using

commercially available diagnostic tests (15).

It is postulated that *M. tuberculosis* evolved from a free-living environmental ancestor

into an obligate pathogen (16)(17). Our current knowledge indicates that *Mycobacterium*

canettii is the most closely related obligate pathogenic species to the MTBC (18)(19).

Infections with *M. canettii* are rare and found solely in people from the Horn of Africa,

with no environmental reservoir described to date (20).

Previous phylogenetic studies have suggested that *Mycobacterium kansasii* (21),

Mycobacterium marinum (22), *Mycobacterium lacus* (23), *Mycobacterium decipiens* (24),

Mycobacterium shinjukuense (25) and *M. riyadhense* (23)(26) are closely related to the

free-living ancestor of the MTBC based on a single marker gene (e.g., *hsp65*, 16S). In

this evolutionary context, a comprehensive understanding of the genomic hallmarks that

drive the close phylogenomic relationship of *M. riyadhense* with the MTBC and NTMs is

of particular scientific interest.

In this study, we report the characterization of eight clinical isolates of *M. riyadhense*,

compared their genomes to members of the MTBC, and provided comprehensive insight

into the adaptive changes associated with the evolution of the MTBC from environmental

bacteria. We examined the lipid profiles of both rough and smooth variants of *M.*

riyadhense, compared them to those of other related mycobacteria, and analysed the

transcriptional response of immunity-related host genes in a murine macrophage infection

model with *M. riyadhense*, *M. kansasii* and *M. bovis* BCG.

Using genome information, we developed a simple PCR-based diagnostic test for the

rapid and accurate identification of *M. riyadhense* to minimize the risk of misdiagnosis in

a clinical setting.

Our analyses provide a comprehensive description of the hallmarks of *M. riyadhense* that

make it one of the closest known environmental relatives of the MTBC and that can serve

to illuminate studies into the evolution and pathogenesis in the MTBC.

Methods and Materials

Clinical reports and bacterial strains

Eight *M. riyadhense* strains were collected in Riyadh, Saudi Arabia, between June 2011

and March 2016 (Table 1) from patients with a positive culture for *M. riyadhense* isolated

from the microbiology laboratory at the King Fahad Medical City (KFMC) in Riyadh,

Saudi Arabia. The patient and sample data collected included demographic and clinical

characteristics, age, sex, clinical features at presentation, presence of comorbidities,

including HIV co-infection, antimicrobial susceptibility, initial and modified therapy,

where applicable; and treatment outcome. Once an isolate was suspected to be an NTM,

the samples were sent to the reference laboratory for full identification and antimicrobial

susceptibility testing. Radiographic and pathologic data were also collected.

Culturing, DNA isolation and sequencing of bacteria

The *M. riyadhense* strains were grown on Lowenstein Jensen (LJ) slants at 37 °C for two

weeks, DNA was extracted using a phenol-chloroform protocol (27), and the quality was

measured by Qubit. Twenty micrograms of high-molecular-weight (HMW) DNA from

the eight *M. riyadhense* strains was sequenced using a PacBio RSII sequencer (Pacific

Biosciences, Menlo Park, USA) with a 10 kb library. An NEBNext Ultra II DNA library

preparation kit (New England BioLabs, Massachusetts USA) was used to prepare

libraries according to the manufacturer's instructions, and sequences from each library

were generated for all *M. riyadhense* strains using the Illumina HiSeq 4000 platform

(Illumina, San Diego, United States).

Genome assembly and annotation

The Illumina short reads were trimmed, and low-quality reads were removed by

Trimmomatic (28). Eight consensus genomes based on each strain were assembled into

contigs with the PacBio long reads using the Canu assembler (29). After assembly, the

draft genome was then corrected with short Illumina reads using Pilon (30) software. The

circularity of assemblies was checked by Gepard (31), and assemblies were annotated by

Prokka (32). A circular map of the chromosome was compared with that of *M.*

tuberculosis H37Rv and visualized with BRIG (33). The genome of the *M. riyadhense*

MR226 strain was used as a high-quality reference in this study.

Comparison of chromosomal and plasmid gene contents of *M. riyadhense* to those of

various *Mycobacterium* species

DNA sequences of 152 mycobacterial species and 77 mycobacterial plasmids were

obtained from the NCBI genome database and independently annotated by Prokka (32).

The predicted protein sequences from the chromosome and each of the two plasmids

(pMRLP01 and pMR01) of the *M. riyadhense* MR226 strain were then compared with

the annotated genes from the rest of the mycobacterial species using orthoMCL (34) with

a 50% identity cutoff and an inflation parameter of 1.5. The obtained orthologues were

visualized with the heatmap package in R (35). A focused OrthoMCL comparison was

performed between (1) *M. riyadhense*, *M. marinum*, *M. kansasii*, *M. szulgai* and

M. tuberculosis and (2) *M. riyadhense* and five species from the MTBC, namely,

M. tuberculosis, *M. bovis*, *M. canettii*, *M. mungi*, and *M. africanum*.

SNP calling and phylogeny based on SNPs

The corrected Illumina reads were mapped using BWA (36) on to the *M. riyadhense*

MR226 genome assembly. Picard tool (37) was used to clean SAM files, fix mate-pair

information and mark duplicates. SNPs were called for two iterations and filtered

(QD<2.0, FS>60.0, SOQ>4.0, ReadPosRankSum<-8.0) with Genome Analysis Toolkit

(GATK) (38). An alignment file was generated by SVAMP (39), and phylogeny was

generated by RaxML (40) with the TVM model.

Phylogeny of *M. riyadhense*

The AMPHORA2 (41) pipeline was used to identify protein sequences from 31

conserved genes in the pangenome datasets and the *Mycobacterium* genomes available at

NCBI or JGI as at 1st January 2018. A total of 148 species were selected after applying a

filter of at least 10 marker genes to be detected in each of the assemblies. *Nocardia*

abscessus was used as the out-group. 12 genes, which have only one copy in each of the

species (i.e., genes *rplA*, *rplB*, *rplF*, *rplM*, *rplN*, *rplP*, *rplS*, *rplT*, *rpmA*, *rpsJ*, *rpsM*, and

rpsS), were concatenated, aligned and trimmed. The phylogenetic tree was then

constructed using RaxML with the Dayhoff model, which is calculated with a script

provided in AMPHORA2 (41).

A whole-genome phylogenetic tree was constructed with the MTBC species (*M.*

tuberculosis, *M. bovis*, *M. canettii*, *M. mungi*, *Mycobacterium orygis*, *M. africanum*) and

with *M. kansasii*, *M. marinum*, *M. shinjukuense*, *Mycobacterium leprae*,

Mycobacterium smegmatis, *Mycobacterium parmense*, *Mycobacterium avium* and

Mycobacterium abscessus. The one-to-one orthologues of each species were obtained

using OrthoMCL and concatenated, then aligned with Muscle (42) and trimmed with

TrimAL (43). The concatenated sequences were composed of 906 genes encoding

296,124 amino acids and were used to build a phylogenetic tree with the LG+G+F model,

which was selected by ModelGenerator. The phylogenetic tree was generated by RaxML

(40).

Toxin/antitoxin systems, *mce/mce*-associated genes and ESX systems in *M.*

***riyadhense* and other mycobacteria**

One hundred fifty-eight toxin/antitoxin (T/A) proteins belonging to the VapBC, RelEF,

HigBA, MazEF, ParDE and UCAT families were downloaded from the NCBI protein

database. We identified the *M. tuberculosis* T/A orthologues from all of the 148 species

by OrthoMCL (34), and the orthologue groups were also examined by Blast+ 2.4.0. The

same pipeline was also applied for the MCE family, PhoPR, PE/PPE, PE-PGRS, and

ESX1-5.

Infection of the RAW 246.7 cell line with *M. riyadhense*, *M. kansasii* and *M. bovis*

BCG Denmark

The murine macrophage RAW264.7 cell line obtained from the American Type Culture

Collection (ATCC, Manassas, USA) was cultured in Dulbecco's modified Eagle's

medium (DMEM) (ThermoFisher Scientific, Waltham, USA) supplemented with 10%

FCS, streptomycin, and penicillin. *M. riyadhense*, *M. kansasii* (subtype I), and *M. bovis*

BCG Denmark strains were grown in Middlebrook 7H9 liquid medium after

single-colony isolation from LJ slants or 7H10 agar. 7H9 was supplemented with 10%

albumin, dextrose and catalase (ADC), while 7H10 was supplemented with oleic acid,

albumin, dextrose and catalase (OADC) in addition to 0.2% glycerol. Ready to use LJ

slants were provided by Saudi Prepared Media Laboratory (SPML, Riyadh, Saudi Arabia).

Before the infection, all bacterial cultures were centrifuged at 1,000xg for 10 minutes. The

supernatant was discarded, and 10-15 3 mm glass beads were added to the pellet and then

vortexed for 1 minute to break up clumps. Six ml of DMEM was added to the pellet and left

to rest for 5 minutes. The upper 5 ml of the suspension was then removed to a fresh 15 ml

falcon tube, which was then centrifuged for an additional 3 minutes at 200xg to remove

remaining bacterial clumps. The supernatant was then taken and passaged using a 26G

hypodermal syringe approximately 15 times to further break up any clumps of bacteria.

The optical density of the culture was measured again before the infection experiment.

RAW264.7 cells were seeded at 2 x 10^5 cells per well in 24-well flat-bottom tissue

culture plates 24 hours prior, reaching 5 x 10^5 cells per well. The DMEM over the cells

was removed, and the cells were washed once with phosphate-buffered saline (PBS). The

cells were infected by applying 1 ml of DMEM containing the prepared mycobacteria

from the former steps at an appropriate concentration to reach an MOI of 5:1, with

DMEM alone used for the control wells. The culture plates were then returned to the

incubator at 37°C with 5% CO_2 for 3 hours to allow for bacterial uptake by the

RAW264.7 cells. The supernatant was removed after 3 hours, and the infected cells were

washed with PBS to remove extracellular bacteria. Subsequently, the cells were

incubated in fresh DMEM with 10% FCS for 24 hours and 48 hours. For harvesting, 400

μl of TRIzol (ThermoFisher Scientific, Waltham, USA) was added to the wells at each

time point, and the adherent cells were scraped out and stored at -80°C for RNA

extraction. Each bacterial infection was performed in triplicate, in addition to the

non-infected controls. RNA was isolated from the samples using the Direct-zol[TM] RNA

Miniprep kit (Zymo Research, Irvine, USA) according to the manufacturer's instructions.

An Agilent RNA 6000 Nano kit was used to check the quality and quantity of the total

RNA. The NanoString murine nCounter Myeloid Innate Immunity Gene Expression

Panel (NanoString Technologies, USA) was used to assess transcript abundance across

infections and timepoints using the nCounter MAX Analysis System (NanoString

Technologies, Seattle, USA). The counts obtained were normalised using the nSolver[TM]

Advanced Analysis plugin (NanoString Technologies, Seattle, USA) using the geNorm

algorithm, and differential gene (DE) expression was analysed using multivariate linear

regression in nSolver^{TM} software, with 0.05 as the p-value cutoff.

Thin layer chromatography (TLC) analysis of lipooligosaccharides (LOSs) in *M.*

riyadhense*, *M. kansasii* and *M. marinum

For TLC analysis, mycobacterial strains were grown at 30°C (*M. marinum*) or 37°C

(*M. smegmatis*, *M. kansasii*, *M. riyadhense*) on LJ slants, and after sufficient incubation,

grown cells were collected and washed once with PBS. Apolar and polar lipids were

extracted from the cell pellets using methods described by Dobson et al. (G. Dobson et al.,

1985). Polar lipids were analysed by 2D-TLC using solvent system E, which is designed

to separate phospholipids and LOSs (G. Dobson et al., 1985). Glycolipids were visualized

by charring following staining with either molybdophophoric acid (MPA) or

alpha-napthol (for glycolipids).

Diagnostic PCR markers for *M. riyadhense*

To develop diagnostic markers for *M. riyadhense* for potential use in clinical diagnosis as

well as environmental and animal studies, unique regions within the *M. riyadhense*

reference genome compared to that of 152 other mycobacterial species were detected

using Shustring (45). These regions were also examined by OrthoMCL (34) and Blastn

(46). *mr_00036*, *mr_00263*, *mr_00606*, and *mr_01005* were selected as the amplification

targets. Two primers for each gene were designed in this study:

MRDP-MR_00036-F (5'-TTCGTTGTCGGTTTCGTCGC-3') and MRDP-MR_00036-R

(5'-GCGTCAGCTCCACCGAAAAC-3');

MRDP-MR_00263-F (5'-CCACCGCTGTTGGCGA-3') and MRDP-MR_00263-R

(5'-TTCGTCCCGTTGATCCCGTT-3');

MRDP-MR_00606-F   (5'-AACCTGCCCGATACGCACTT-3')   and

MRDP-MR_00606-R (5'-ACTGTTCCTCCGTGGGGTTG-3');

MRDP-MR_01005-F   (5'-GACTGTGGGGTAACGGTGGA-3')   and

MRDP-MR_01005-R (5'-CCGGTGATGTCGCCTACTCC-3').

PCR was performed in a 25 µl reaction volume with 12.5 µl of GoTaq ® Green Master

Mix (Promega, USA), 1 µl of 100 ng/µl gDNA, 1 µl with 10 nmol of forward and reverse

primers, 3 µl of dimethyl sulfoxide (DMSO) and 19 µl of nuclease-free water. The PCR

mixture was denatured for 5 minutes at 94°C; followed by 35 cycles of amplification

involving a denaturation step at 94°C for 30 seconds, a primer annealing step at 59°C for

45 seconds, and a primer extension step at 72°C for 45 seconds; and a final extension step

at 72°C for 7 minutes. The ITS-F/mycom-2 primer set (47), which is a *Mycobacterium*

genus-specific primer set, was used as a control, with amplification conditions as

described previously (47). The products were electrophoresed in a 2% agarose gel for 60

minutes and visualized.

Results

Clinical manifestation and culture characteristics of *M. riyadhense* isolates

Between April 2011 and March 2017, eight clinical cases of infection with *M. riyadhense* were recorded in male patients aged from 8-82 years (Table 1). In addition to HIV/AIDS, most of the patients had multiple comorbidities, such as pulmonary and/or systemic hypertension, malignancies, and diabetes mellitus (DM), with the lung being the major site of disease (Table 1).

Bronchoalveolar lavage, endotracheal, sputum and lymph node biopsy specimens grew *M. riyadhense* colonies when cultured in solid and liquid media using LJ agar and *Mycobacteria* growth indicator tube (MGIT) broth, respectively, albeit with varying times to positivity. Although no consistent findings existed for chest imaging, three out of the eight patients presented with upper lobe consolidation, cavitation, ground-glass opacities, 'tree-in-bud' appearance, hilar lymphadenopathy and pleural effusion (Figure supplementary 1).

The isolates were subjected to susceptibility testing for antibiotics commonly used to treat both typical and atypical infections. Isolates showed 100% susceptibility to rifampin

(RIF), rifabutin (RFB), ethambutol (EMB), clarithromycin (CLR), linezolid (LZD),

amikacin (AMK), moxifloxacin (MOXI), and trimethoprim-sulfamethoxazole

(TMP-SMX). Three out of the eight isolates were resistant to ciprofloxacin (CIP).

Assembly and annotation of the *M. riyadhense* MR226 genome

The comparison of different assemblies of all sequenced *M. riyadhense* strains is listed in

Supplementary Table 1. We obtained chromosomes of all eight isolates in single

contiguous sequences for genome comparison at a high resolution. The *M. riyadhense*

MR226 strain contains a 6,243,587 bp chromosome, a linear plasmid (pMRLP01) of

550,247 bp and a circular plasmid (pMR01) of 94,344 bp. The circular nature of the

chromosome and the pMR01 plasmid were demonstrated through Gepard (31).

As expected of a free-living opportunistic pathogen, the chromosome of *M. riyadhense* is

significantly larger than the chromosomes of the MTBC (Table 2). The number of genes

unique in a species showed that the members of the MTBC have a considerably lower

percentage of unique genes than *M. riyadhense* and other closely related NTMs (Table 2).

The comparison of the annotated protein-coding genes from the *M. riyadhense* MR226

strain to the genome assemblies of 152 *Mycobacterium* species (including 77 known

mycobacterial plasmids) shows that *M. riyadhense* forms a tight cluster with the MTBC

and some NTMs (Figure supplementary 2A, highlighted in box a). A total of 335 genes

were identified as unique from the orthologue group comparison, being present in only all

sequenced *M. riyadhense* isolates, including MR226, and the vast majority of these genes

belong to the PE/PPE family, which are thought to be involved in antigen variation and

are widely spread across the slow-growing species within the *Mycobacterium* genus.

Linear plasmids were first described in 1989 in maize (which has a linear

mitochondrion)(48) and have also been found in *Actinomycetales*, including *Streptomyces*

(49), *Rhodococcus* (50) and *Mycobacterium* species, such as *Mycobacterium xenopi*,

Mycobacterium branderi and *Mycobacterium celatum*. They are often accompanied by a

circular plasmid in the same host (51). The linear plasmid pMRLP01 in *M. riyadhense*

contains a pair of partitioning genes (*parA*/*parB*) that are involved in active segregation

and thus stabilize the inheritance of the plasmid (52). The latter are known to contribute

to genome evolution by active DNA transfer and exchange (53). As is often the case for

both circular and linear large plasmids, a relatively higher proportion of pMRLP01 genes

(51%) have no known function compared to those of the main chromosome (26%). This

reinforces the notion that plasmids are an important route by which new genes are

introduced into the genome in *Mycobacteria*. Of the 443 predicted protein-coding genes

of pMRLP01, 118 have at least one orthologue on the main chromosome. Furthermore,

we observed several protein-coding genes in pMRLP01 that have orthologues in the

genomes of *Mycobacterium tusciae* JS617, *Mycobacterium aromaticivorans* JCM 16368,

Mycobacterium llatzerense, *Mycobacterium obuense*, *Mycobacterium novocastrense* and

Mycobacterium holsaticum (Figure supplementary 2B). This finding indicates that

pMRLP01 can be readily exchanged with other environmental mycobacterial species.

It is well known that plasmids are important "vehicles" for the exchange of genetic

material between bacteria or between chromosomes and extra-chromosomal plasmids. In

this study, we further identified a circular plasmid termed pMR01 in *M. riyadhense*

(Figure supplementary 2C, Figure supplementary 3). When compared with the circular

plasmids of other species, such as pRAW in *M. marinum* (54), pMAH135 (55) and

pMA100 (56) of *M. avium*, pMyong1 from *Mycobacterium yongonense* (57), pMK12478

(58) from *M. kansasii* and several plasmids from *Mycobacterium chimaera* (59), a high

similarity was observed. These plasmids all harbour both a TS4 and TS7, which are

necessary for conjugation (54)(60), and facilitate the exchange of genetic material

between different species of slow-growing mycobacteria (54). We therefore speculate

that pMR01 is a novel conjugative plasmid.

In mycobacteria, five type VII secretion systems have been described,　named ESX-1 to

ESX-5 (61),(62). An ESX-P5 locus on pMR01, which shows high similarity to the ESX-5

loci on pMK12478, pRAW and pMAH135, is markedly different from the ESX-5 system

found on the main *M .riyadhense* chromosome. ESX-5 is involved in the secretion of PE

and PPE proteins in *M. tuberculosis* and is involved in modulating the host immune

responses to maintain a persistent infection (63). The potential transmissibility of pMR01

and other pMR01-like plasmids may therefore play a role in the evolution of the ESX

systems in mycobacteria.

The progressive alignments (Figure supplementary 4A) of the assembled chromosomes

and plasmids (Figure supplementary 4B) of each *M. riyadhense* strain show that the

chromosomes are relatively conserved; however, the linear plasmids present in all eight

sequenced isolates are diverse from both structural and similarity perspectives, while the

pRAW-like plasmids are present in only the MR226, MR193 and MR222 strains.

The SNP-based phylogeny of the sequenced *M. riyadhense* isolates based on 43,136

polymorphic sites (Supplementary Table 2, Figure supplementary 5) indicates the

presence of two different clades of *M. riyadhense* among the clinical isolates sequenced

in this study. The nucleotide diversity between the MR222 clade is greater than the

diversity between *M. tuberculosis* strains (64), while it is smaller than that seen between

M. canettii strains (18), and the variation between the MR226 clade is comparable to the

SNP variation in *M. tuberculosis* strains.

Regions of difference (RDs) in *M. riyadhense*

The RDs were originally described as genomic regions present in virulent *M. bovis* and *M.*

tuberculosis but absent from the *M. bovis* BCG genome (65). RD loci were subsequently

described across the MTBC (66) and contain functions believed to contribute to

pathogenicity (67)(68)(69) and evolution of MTBC species (70). *M. riyadhense* was

found to harbour most of the RD loci (RD1, RD3-R11, R13-RD16) that are also intact in

M. tuberculosis, while 2 of the RDs show unique deletions, RD2^{riyadh} (Figure

supplementary 6A) and RD12^{riyadh} (Figure supplementary 6B).

RD2 was originally described as deleted in BCG vaccine strains.   Subsequently, it was

shown that disruption of RD2 in *M. tuberculosis* leads to decreased proliferation *in vivo*

and impaired modulation of the innate immune response (68). The RD2 locus also has a

deletion from the *M. riyadhense* genome. It is a larger deletion than the originally

described RD2^{BCG} as RD2^{riyadh} contains 29 genes (*rv1971~rv2000*, location

2,216,498~2,246,766); eight genes within this locus (*rv1978*, *rv1979c*, *rv1980c*, *rv1981c*,

rv1983, *rv1984*, *rv1987*, *rv1988*) have orthologues elsewhere in the *M. riyadhense*

genome (*mr_05764*, *mr_05852*, *mr_02310*, *mr_02993*, *mr_00486*, *mr_02995*, *mr_02325*,

mr_02349, *mr_01747*), suggesting possible functional redundancy.

The RD12 locus shows deletions across MTBC members, including *M. bovis*,

Mycobacterium caprae and *M. orygis* (71), but it is present in other MTBC members. *M.*

canettii isolates (except group B (72)) also show an independent deletion at the RD12

locus named RD12^{can} (3,479,430~3,491,866, *rv3111~rv3126*), which is distinct from

RD12^{bovis} (3,484,740~3,487,515, *rv3117~rv3121*). We identified another unique deletion

at the RD12 locus in *M. riyadhense*, designated RD12^{riyadh}, which encompasses a larger

region than RD^{can} and RD12^{bovis}, encompassing *rv3108-rv3127* (3,477,171~3,492,150)

(Figure supplementary 6B). It is intriguing that multiple mycobacteria show independent

deletion events at the RD2 and RD12 loci, suggesting selective forces play a role in this

variation.

Comparative phylogeny of *M. riyadhense* with other Mycobacteria

The phylogenetic tree shows that the slow-growers and rapid-growers are separated into

two different clades and that fast-growers evolved from slow-growers (Figure 1, the

detailed figure is shown in Figure supplementary 7). The overall topology of our tree is

remarkably similar to that of previously published phylogenetic trees (23). In the tree,

M. riyadhense is located within the same clade as the obligate and opportunistic causal

organisms of mycobacterial diseases in humans that include the MTBC, *M. marinum*, *M.*

kansasii, *M. leprae* and related host-restricted mycobacteria with reduced genomes and

decreased survivability in the environment. Notably, *M. shinjukuense*, *M. lacus* and *M.*

riyadhense form a sub-clade that is phylogenetically close to the MTBC.

The *PE/PPE* family *mce* and *mce*-associated genes are known to be important for host

adaptation (73) and pathogenicity (74). The *PE/PPE* family genes are enriched in the

MTBC members but also in *M. riyadhense* MR226 (278) and other pathogenic species,

such as *M. kansasii* (228) and *Mycobacterium ulcerans* (200). The number of *mce-* or

mce-associated genes has not significantly changed across mycobacterial genomes,

indicating that this group of genes is under an evolutionary constraint and plays

functional roles bridging both environmental and obligate pathogen lifestyles. Our results

agree with previous finding that during their evolution, the ESX systems were derived

from the ancestor ESX-4, as shown in Figure supplementary 7 at the root node, and then

ESX-3, ESX-1, ESX-5 and ESX-2 evolved by horizontal transfer (75).

A comparative phylogenetic map based on 1,301 conserved proteins (Figure

supplementary 8) reveals this downsizing of the genome and the dynamic changes in

genome components. Certain functional categories of genes are relatively enriched during

the evolution of MTBC, including protein metabolism, regulation and cell signalling, cell

division and cell cycle.

M. riyadhense shares a larger number of orthologues (3,122) with *M. tuberculosis* than

with *M. kansasii* (2,978), *M. marinum* (2,962) and *M. szulgai* (2,724) among the

environmental mycobacteria that are closely related to the MTBC (Figure 2A). A total of

134 orthologues are exclusively shared between *M. riyadhense* and *M. tuberculosis*,

while the number of orthologues exclusively shared between *M. tuberculosis* and *M.*

kansasii (30), *M. marinum* (48) and *M. szulgai* (18) is less (Figure 2A). It is notable that

genes from the phage-derived regions of RD3 and RD11 are shared exclusively between

M. riyadhense and *M. tuberculosis*.

The comparative analysis of the orthologue groups of *M. riyadhense* and the MTBC is

revealing. Firstly, 385 orthologue groups present across all MTBC species are absent

from *M. riyadhense*. Secondly, 221 orthologues uniquely present in *M. riyadhense* are not

found amongst the MTBC (Figure 2B). This latter group of *M. riyadhense* unique

orthologues likely depict the constraints imposed by the free-living biology of *M.*

riyadhense which necessitates maintaining a broad functional repertoire to secure

environmental survival. In contrast, the obligate MTBC species have lost genes involved

in environmental survival but gained a large number of genes required for survival in the

in vivo environment.

A hallmark of *M. tuberculosis* infection is the ability to survive long-term in host

granulomas and develop a latent stage of infection. The molecular mechanisms and

cellular components that are involved in the persistence of *M. tuberculosis* are still poorly

understood, but several T/A systems have been implicated in the pathogenicity of *M.*

tuberculosis (76). T/A systems were first found on plasmids or plasmid-derived

chromosomal loci where they promoted plasmid maintenance in bacterial populations

(77), but when compared to other mycobacteria, the MTBC are remarkable for the

extensive expansion of T/A systems. We thus compared the 79 pairs of T/A systems

(belonging to the HigAB, MazEF, ParDE, RelEF, VapBC and UCAT families) in *M.*

tuberculosis with the T/A pairs found in other members of the MTBC and NTMs. Based

on the presence of 49 out of the 79 T/A orthologue pairs (Figure 3), *M. riyadhense*

appears more closely related to the MTBC   than to other mycobacteria, including *M.*

lacus, M. shinjukuense and *M. decipiens*. It can be hypothesized that the shared

component of T/A pairs play a functional role in the pathogenicity or persistence of *M.*

riyadhense infection in a way similar to that described for the importance of T/A pairs in

M. tuberculosis in vivo biology.

M. riyadhense strains produce a distinct pattern of LOSs

The LOSs are an important class of glycolipids that have previously been linked to

diverse mycobacterial phenotypes, such as colony morphology, secretion of PE/PPE

family proteins, and the pathogenicity of *M. marinum* (78). Notably, the presence or

absence of LOS from *M. canettii* causes a transition from smooth to rough colony

variants, respectively, with rough variants showing increased virulence; MTBC strains

lack LOS, and it has been suggested that the removal of LOS was a key event in the

evolution of the MTBC species towards their current obligate pathogen status (79). It is

noteworthy that we observed both smooth (MR210, MR22, MR226, MR244, MR246,

and MR1023) (Figure 4A) and rough (MR193 and MR206) (Figure 4B) morphologies in

M. riyadhense strains. We therefore sought to first examine whether the genetic

machinery for the production of LOS is present in the *M. riyadhense* genome and then to

follow up on the genome-level predictions with lipid analyses of rough and smooth

variants using TLC.

We observed that the *wecE* and *galE6* LOS genes are absent from the *M. riyadhense*

genome (Table 3, Supplementary Table 3). These genes are linked with the removal of

LOS II* and the production of LOS IV (78). Thus, their absence is likely to cause an

accumulation of LOS II* and the lack of fully formed LOS IV, which have previously

been shown to increase the pathogenicity of *M. marinum* (78). Furthermore, both the *pks5*

and *pap* genes in the LOS locus are intact in *M. riyadhense*, as is the case in *M. canettii*,

but not in *M. tuberculosis,* where the former is truncated and the latter deleted (79). This

finding indicates that *pks5* recombination and *pap* deletion occurred in a common

ancestor of the MTBC after its differentiation from both *M. riyadhense* and *M. canettii.*

Remarkably, the *M. riyadhense* LOS gene locus layout is dissimilar to that in *M. canettii*,

M. tuberculosis, *M. kansasii* and *M. marinum*: indeed, exclusive   rearrangements of this

locus in *M. riyadhense* were observed (Figure 4C).

To correlate rough vs. smooth colony morphology with LOS production, we extracted

polar lipids from the strains and analysed them by 2D-TLC using solvent system E (G.

Dobson et al., 1985), which is designed to separate phospholipids and LOSs. Charring of

the TLC plates with alpha-napthol revealed glycolipids, including the accumulation of a

species that migrated at a position similar to that of LOS III. This lipid was seen in only

smooth strains; species with migration patterns similar to LOS I and LOS II were

observed, while LOS IV was absent. This result was not unexpected because all *M.*

riyadhense strains lack a functional *wecA* gene, which is required for the extension of

LOS II to LOS IV (Figure 4). Additionally, the relative levels of the predominant LOS

species in *M. riyadhense* seem to be quite high when compared to those seen in other

LOS-producing mycobacteria (Figure 4 (D)(E)). Conversely, the rough strains did not

produce any glycolipids that migrated in the positions corresponding to LOSs (Figure 4

(F)(H)). Overall, *M. riyadhense* exhibits a LOS production phenotype distinct from that

of other LOS-producing mycobacteria.

PE-PGRS33* locus and type VII secretion system of *M. riyadhense

The *pe-pgrs33* (*rv1818c*) locus encodes the exported protein PE_PGRS33 and plays an

important role in the pathogenesis of *M. tuberculosis* (80). A previous study (81) showed

that *pe-pgrs33* is present in all MTBC members but not in *M. canettii*, which implies a

specific *pe_pgrs33* insertion event in the ancestor of MTBC strains. Genome comparison

of *M. riyadhense* with *M. tuberculosis*, *M. kansasii*, *M. marinum* and *M. canettii* provides

additional evidence that *M. riyadhense* is the missing link of the *pe-pgrs33*

deletion/insertion event. Our phylogeny strongly suggests that the deletion of *pe-pgrs33*

from *M. kansasii* and *M. marinum* occurred before the divergence of environmental

mycobacteria and the smooth tubercle bacilli (STB)/MTBC clade (Figure supplementary

9).

All 5 ESX systems (ESX1-ESX5) were found in the *M. riyadhense* genome (Figure

supplementary 10) but with minor modifications. Hence, the *eccA* and *eccB* genes are

absent from the ESX-2 system, while the *espACD* operon, which is essential for secretion

of virulence factors via ESX1, is also missing in *M. riyadhense*. The overall gene

arrangement of the ESX1-ESX5 loci is very similar in both *M. riyadhense* and

M. tuberculosis (61). This conserved synteny reinforces the previous results that

phylogenetically *M. riyadhense* may represent an ancestral state to MTBC. As noted

before, the pMR01 plasmid also contains an extra ESX-P5 locus, which could indicate a

role for this plasmid in mediating pathogenicity (Figure supplementary 4).

Transcriptional response of murine macrophage cells upon *M. riyadhense* infection

Our genomic analysis of *M. riyadhense* revealed a range of genes and potential systems

that could play a role in host-pathogen interactions. We therefore sought to assess the

initial interaction of *M. riyadhense* with macrophages, using the RAW264.7 cell line of

murine origin as our experimental model. As comparator strains in our analysis, we

performed parallel infections with *M. kansasii*, an opportunistic pathogen that also

contains an orthologous ESX-1 system (Figure 1, Figure supplementary 10), and *M. bovis*

BCG, the live TB vaccine that is attenuated through deletion of the ESX-1 system. These

comparisons allowed us to explore whether *M. riyadhense* triggered innate immune

responses were more similar to those triggered by *M. kansasii* or BCG or were

intermediate between the two.

To analyse the innate immune responses of the macrophages in the intracellular presence

or absence of these mycobacterial isolates, the transcriptional response was analysed

using a 754 probe NanoString Murine Myeloid Innate Immunity panel V2 (82)   at 3, 24

and 48 hours post infection. These analyses revealed an expected commonality in the

responses to infection with all mycobacteria, such as induction of proinflammatory genes

through TLR signalling (e.g., upregulation of *IL1B*, *TNFA*, *CCL4*, *PTGS2* and *CXCL2*,

albeit to different absolute levels, Figure 5, Figure supplementary 11A and 11B). Distinct

responses triggered by BCG infection compared to *M. riyadhense* and *M. kansasii*

included, for example, upregulation of *MARCO* by BCG infection (Figure supplementary

11B); *MARCO* is involved in pathogen uptake via trehalose dimycolate, a lipid that is

known to show variation in structure between the MTBC and *M. kansasii* (83)(84).

CCL24 and *CXCL14*, which are involved in the attraction of immune cells to the site of

infection (85), were upregulated to higher levels with *M. riyadhense* and *M. kansasii* than

with BCG at 24 and 48 hours.

Overall, our analysis of the macrophage transcriptional profiles showed that the response

to infection with *M. riyadhense* and *M. kansasii* triggered more similar macrophage

transcriptional responses in comparison to the response produced by BCG infection

(Figure supplementary 11).

Developing a rapid PCR-based diagnostic marker for *M. riyadhense*

Due to the issues previously encountered in diagnosing *M. riyadhense* infections (8)(9),

correct and prompt identification of cases upon presentation at healthcare units is of

paramount importance. We therefore sought to translate our knowledge on the genome

sequences into a PCR diagnostic test that could be used in a clinical microbiology setting

to distinguish *M. riyadhense* from other mycobacteria, including the members of the

MTBC.

By identifying unique K-mers ranging in size from 11 bp to 4,209 bp (Figure

supplementary 12) in the assembled genome compared to the genomes of 152 other

mycobacterial species, a primer set was developed targeting the *mr_00036*, *mr_00263*,

mr_00606 and *mr_01005* genes. The MRDP primer pair MRDP-F/MRDP-R amplified a

single product of 372 bp from each of the eight isolates of *M. riyadhense* (Figure 6) but

not from other mycobacterial species, including *M. tuberculosis*, *M. bovis*, *M. kansasii*, *M.*

marinum, *M. szulgai*, *M. avium* and *Mycobacterium angelicum*. This result shows that the

MRDP-F/MRDP-R primers are highly specific to *M. riyadhense* and form the basis for a

simple diagnostic PCR that can inform appropriate treatment protocols.

Discussion

M. riyadhense has become a clinically relevant NTM species globally (8)(86). Contrary

to prior publications on *M. riyadhense* have been based primarily on clinical case reports,

here we present the largest and most comprehensive genomic study undertaken to date on

clinical *M. riyadhense* isolates. Our comparative analysis of *M. riyadhense* genomes with

the MTBC and a large collection of NTMs provide unequivocal evidence that *M.*

riyadhense is one of the closest known environmental mycobacteria species to the MTBC

and forms a phyloclade with *M. lacus*, *M. decipiens* and *M. shinjukuense*.

Our analyses of *M. riyadhense* reveal that expansion of T/A pairs, modification of

secretion systems, alterations in cell wall lipids, and plasmid-mediated horizontal

acquisition of new functionality all played key roles in the evolution of the MTBC. Our

work adds new insight into the evolution of the MTBC from free-living environmental

bacteria to their obligate pathogen biology.

The eight new *M. riyadhense* strains sequenced in this study originated from pulmonary

infections (with some having additional extra-pulmonary involvement), which fulfilled

the American Thoracic Society/ Infectious Diseases Society of America (ATS/IDSA)

criteria for NTM infection (87).

Our study shows that *M. riyadhense* shares a larger number of orthologues with *M.*

tuberculosis than *M. kansasii* and *M. marinum*, notably in the T/A gene family (Figure

supplementary 6). Forty-nine pairs of T/A gene orthologues were found in *M. riyadhense*,

far greater than the number of orthologues observed in any of the other NTMs. The

expansion of T/A genes among the MTBC offers additional proof   that gene acquisition

through lateral gene transfer played a key role in the development of pathogenicity; this

agrees with previous findings (22)(88).

M. riyadhense strains appeared as both smooth and rough colony forms when grown on

solid LJ media. We observed the presence of LOS I and LOS II and the absence of LOS

IV. These biochemical observations agree with the genomic prediction that *M.*

riyadhense strains lack a functional *wecA* gene, which is required for the extension of

LOS II to LOS IV. *In silico* RD analysis *M. riyadhense* revealed independent RD2^{riyadh}

and RD12^{riyadh} deletions when compared to *M. canettii* and other members of the MTBC.

Frequent and independent deletion events in these specific genomic regions across these

groups of closely related mycobacteria suggest a role for natural selection that might be

important for pathogenicity.

All the 5 ESX type VII secretion systems present in *M. tuberculosis*, and known to be

involved in virulence and pathogenicity, were found in *M. riyadhense* with very similar

gene arrangement. An additional ESX-P5 system was also found on the circular plasmid

pMR01. pMR01-like plasmids have been shown in many different NTMs, including *M.*

kansasii, *M. marinum*, *M. chimaera* and *M. avium*, indicating that these groups of

plasmids containing both TS4 and TS7 are conjugative plasmids. The extensive presence

of these plasmids may also explain the origin of the type VII secretion system (54).

ESAT-6 and CFP-10, which are secreted through ESX-1, are critical to phagosome

perturbation and manipulation of host macrophages induced by *M. tuberculosis* (89).

Linked to this, our transcriptome analysis revealed that *M. riyadhense* and *M. kansasii*

trigger similar overall macrophage responses after infection when compared to *M. bovis*

BCG (as a representative of the MTBC). These data reinforce the phylogenetic

relationship of *M. riyadhense* with the MTBC species and its transitional status between

an opportunistic and an obligate pathogen.

The clinical presentation of our cases was mainly indistinguishable from disease caused

by *M. tuberculosis*, as reported earlier (7)(8), but with a negative *M. tuberculosis* PCR.

Due to the relatively recent emergence of *M. riyadhense* as an important clinical

pathogen, and coupled with its misdiagnosis as *M. tuberculosis* by commercially

available kits, we developed an accurate set of diagnostic markers based on the genomic

datasets generated in this study. The primer sets accurately detect *M. riyadhense* in a

mixed cocktail of closely related mycobacteria: it   can serve as part of an accurate and

fast diagnostic protocol in clinical settings thus reducing the need for strict isolation,

laborious contact tracing as well as the inappropriate use of TB antimicrobials. . These

primers can be used in a global epidemiological survey of cases of *M. riyadhense*

infections in humans, animals and the environment. This may thus provide a more

complete picture of the epidemiology of *M. riyadhense* following a broad One Health

approach. It would, for example, be of clinical interest to see if *M. riyadhense* infections

occur in Africa and South America, for which no reports are available, or if *M.*

riyadhense is uniquely endemic in the Arabian Peninsula. Indeed, a limitation of this and

previous similar studies is that all isolates available for genomic studies are clinical. *M.*

riyadhense is an environmental pathogen that is most likely be found in animals sharing

the same niche with humans that were infected from the environment. As the natural

environmental reservoir of *M. riyadhense* remains unknown, we believe that systematic

screening of relevant environmental samples with the MRDP established in this study

may help to establish the natural habitat of this bacterium.

In conclusion, our study provides unprecedented insights into the ancestry and adaptive

evolution of the MTBC relative to extant NTM species and places *M. riyadhense* as one

of the closest environmental relatives to the MTBC. Our work provides compelling data

supporting the use of *M. riyadhense* as a novel mycobacterium model for the study of

virulence evolution and pathogenesis in MTBC.

References

1. Falkinham JO. 2013. Ecology of nontuberculous mycobacteria-where do human
infections come from? Semin Respir Crit Care Med 34:95–102.
2. Van Ingen J, Bendien SA, De Lange WCM, Hoefsloot W, Dekhuijzen PNR,
Boeree MJ, Van Soolingen D. 2009. Clinical relevance of non-tuberculous
mycobacteria isolated in the Nijmegen-Arnhem region, The Netherlands. Thorax
64:502–506.
3. Ding L-W, Lai C-C, Lee L-N, Huang L-M, Hsueh P-R. 2005. Lymphadenitis
caused by non-tuberculous mycobacteria in a university hospital in   Taiwan:
predominance of rapidly growing mycobacteria and high recurrence rate. J Formos
Med Assoc 104:897–904.
4. Panwalker AP, Fuhse E. 1986. Nosocomial Mycobacterium gordonae
pseudoinfection from contaminated ice machines. Infect Control 7:67–70.
5. Carbonne A, Brossier F, Arnaud I, Bougmiza I, Caumes E, Meningaud JP, Dubrou
S, Jarlier V, Cambau E, Astagneau P. 2009. Outbreak of nontuberculous

mycobacterial subcutaneous infections related to multiple mesotherapy injections.

J Clin Microbiol 47:1961–1964.

6.   Singh J, O'Donnell K, Ashouri N, Adler-Shohet FC, Nieves D, Tran MT, Arrieta

A, Tran L, Cheung M, Zahn M. 2018.　926. Outbreak of Invasive Nontuberculous

Mycobacterium (NTM) Infections Associated With a Pediatric Dental Practice .

Open Forum Infect Dis 5:S29–S29.

7.   Garbati, Hakawi AM. 2014. Mycobacterium Riyadhense Lung Infection in a

Patient with HIV/AIDSSub-Saharan African Journal of Medicine.

8.   Choi JI, Lim JH, Kim SR, Lee SH, Park JS, Seo KW, Jeon JB, Jeong J. 2012.

Lung infection caused by Mycobacterium riyadhense confused with

Mycobacterium tuberculosis: The first case in Korea. Ann Lab Med 32:298–303.

9.   van Ingen J, Al-Haijoj SAM, Boeree M, Al-Rabiah F, Enaimi M, de Zwaan R,

Tortoli E, Dekhuijzen R, van Soolingen D. 2009. Mycobacterium riyadhense sp.

nov., a non-tuberculous species identified as Mycobacterium tuberculosis complex

by a commercial line-probe assay. Int J Syst Evol Microbiol 59:1049–1053.

10.   King HC, Khera-Butler T, James P, Oakley BB, Erenso G, Aseffa A, Knight R,

Wellington EM, Courtenay O. 2017. Environmental reservoirs of pathogenic

mycobacteria across the Ethiopian biogeographical landscape. PLoS One 12.

11.   Narendrula-Kotha R, Nkongolo KK. 2017. Microbial response to soil liming of

damaged ecosystems revealed by pyrosequencingand phospholipid fatty acid

analyses. PLoS One 12.

12.   Godreuil S, Marchandin H, Michon AL, Ponsada M, Chyderiotis G, Brisou P, Bhat

A, Panteix G. 2012. Mycobacterium riyadhense pulmonary infection, France and

Bahrain. Emerg Infect Dis.

13.   Van der Werf MJ, K??dm??n C, Katalini??-Jankovi?? V, Kummik T, Soini H,

Richter E, Papaventsis D, Tortoli E, Perrin M, van Soolingen D, ??olnir-Dov??

M, ??stergaard Thomsen V. 2014. Inventory study of non-tuberculous

mycobacteria in the European Union. BMC Infect Dis 14.

14.   Althawadi S, Johani S, Fernandez GM, Al-ghafli H, Enani RMA, Abdul K,

Medical A. 2017. Mycobacterium riyadhense 23:2015–2017.

15.   Tortoli E, Pecorari M, Fabio G, Messinò M, Fabio A. 2010. Commercial DNA

709   probes for mycobacteria incorrectly identify a number of less frequently

710   encountered species. J Clin Microbiol.

711   16.   Murty DS PK. 2014. A Brief review on Ecology and Evolution of Mycobacteria.

712   Mycobact Dis 04.

713   17.   Jang J, Becq J, Gicquel B, Deschavanne P, Neyrolles O. 2008. Horizontally

714   acquired genomic islands in the tubercle bacilli. Trends Microbiol 16:303–308.

715   18.   Supply P, Marceau M, Mangenot S, Roche D, Rouanet C, Khanna V, Majlessi L,

716   Criscuolo A, Tap J, Pawlik A, Fiette L, Orgeur M, Fabre M, Parmentier C, Frigui

717   W, Simeone R, Boritsch EC, Debrie AS, Willery E, Walker D, Quail MA, Ma L,

718   Bouchier C, Salvignol G, Sayes F, Cascioferro A, Seemann T, Barbe V, Locht C,

719   Gutierrez MC, Leclerc C, Bentley SD, Stinear TP, Brisse S, Médigue C, Parkhill J,

720   Cruveiller S, Brosch R. 2013. Genomic analysis of smooth tubercle bacilli

721   provides insights into ancestry and pathoadaptation of Mycobacterium tuberculosis.

722   Nat Genet 45:172–179.

723   19.   Van Soolingen D, Hoogenboezem T, De Haas PEW, Hermans PWM, Koedam

724   MA, Teppema KS, Brennan PJ, Besra GS, Portaels F, Top J, Schouls LM, Van

725   Embden JDA. 1997. A Novel Pathogenic Taxon of the Mycobacterium

726   tuberculosis Complex, Canetti: Characterization of an Exceptional Isolate from

727   Africa. Int J Syst Bacteriol 47:1236–1245.

728   20.   Blouin Y, Hauck Y, Soler C, Fabre M, Vong R, Dehan C, Cazajous G, Massoure

729   P-L, Kraemer P, Jenkins A, Garnotel E, Pourcel C, Vergnaud G. 2012.

730   Significance of the Identification in the Horn of Africa of an Exceptionally Deep

731   Branching Mycobacterium tuberculosis Clade. PLoS One 7:e52841.

732   21.   Wang J, McIntosh F, Radomski N, Dewar K, Simeone R, Enninga J, Brosch R,

733   Rocha EP, Veyrier FJ, Behr MA. 2015. Insights on the emergence of

734   Mycobacterium tuberculosis from the analysis of Mycobacterium kansasii.

735   Genome Biol Evol 7:856–870.

736   22.   Stinear TP, Seemann T, Harrison PF, Jenkin G a, Davies JK, Johnson PDR,

737   Abdellah Z, Arrowsmith C, Chillingworth T, Churcher C, Clarke K, Cronin A,

Davis P, Goodhead I, Holroyd N, Jagels K, Lord A, Moule S, Mungall K,

Norbertczak H, Quail M a, Rabbinowitsch E, Walker D, White B, Whitehead S,

Small PLC, Brosch R, Ramakrishnan L, Fischbach M a, Parkhill J, Cole ST. 2008.

Insights from the complete genome sequence of Mycobacterium marinum on the

evolution of Mycobacterium tuberculosis. Genome Res 18:729–41.

23.   Tortoli E, Fedrizzi T, Meehan CJ, Trovato A, Grottola A, Giacobazzi E, Serpini

GF, Tagliazucchi S, Fabio A, Bettua C, Bertorelli R, Frascaro F, De Sanctis V,

Pecorari M, Jousson O, Segata N, Cirillo DM. 2017. The new phylogeny of the

genus Mycobacterium: The old and the news. Infect Genet Evol 56:19–25.

24.   Brown-Elliott BA, Simmer PJ, Trovato A, Hyle EP, Droz S, Buckwalter SP,

Borroni E, Branda JA, Iana E, Mariottini A, Nelson J, Matteelli A, Toney NC,

Scarparo C, de Man TJB, Vasireddy R, Gandhi RT, Wengenack NL, Cirillo DM,

Wallace RJ, Tortoli E. 2018. Mycobacterium decipiens sp. nov., a new species

closely related to the Mycobacterium tuberculosis complex. Int J Syst Evol

Microbiol 68:3557–3562.

25.   Saito H, Iwamoto T, Ohkusu K, Otsuka Y, Akiyama Y, Sato S, Taguchi O,

Sueyasu Y, Kawabe Y, Fujimoto H, Ezaki T, Butler R. 2011. Mycobacterium

shinjukuense sp. nov., a slowly growing, non-chromogenic species isolated from

human clinical specimens. Int J Syst Evol Microbiol 61:1927–1932.

26.   Fedrizzi T, Meehan CJ, Grottola A, Giacobazzi E, Fregni Serpini G, Tagliazucchi

S, Fabio A, Bettua C, Bertorelli R, De Sanctis V, Rumpianesi F, Pecorari M,

Jousson O, Tortoli E, Segata N. 2017. Genomic characterization of

Nontuberculous Mycobacteria. Sci Rep 7.

27.   Belisle JT, Sonnenberg MG. 1998. Isolation of genomic DNA from mycobacteria.

Methods Mol Biol 101:31–44.

28.   Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: A flexible trimmer for

Illumina sequence data. Bioinformatics 30:2114–2120.

29.   Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM. 2017.

Canu: Scalable and accurate long-read assembly via adaptive κ-mer weighting and

repeat separation. Genome Res 27:722–736.

30.   Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, Cuomo CA,
769   Zeng Q, Wortman J, Young SK, Earl AM. 2014. Pilon: An integrated tool for
770   comprehensive microbial variant detection and genome assembly improvement.
771   PLoS One 9.

772   31.   Krumsiek J, Arnold R, Rattei T. 2007. Gepard: A rapid and sensitive tool for
773   creating dotplots on genome scale. Bioinformatics 23:1026–1028.

774   32.   Seemann T. 2014. Prokka: Rapid prokaryotic genome annotation. Bioinformatics
775   30:2068–2069.

776   33.   Alikhan NF, Petty NK, Ben Zakour NL, Beatson SA. 2011. BLAST Ring Image
777   Generator (BRIG): Simple prokaryote genome comparisons. BMC Genomics 12.

778   34.   Li L, Stoeckert CJ, Roos DS. 2003. OrthoMCL: Identification of ortholog groups
779   for eukaryotic genomes. Genome Res 13:2178–2189.

780   35.   Ihaka R, Gentleman R. 1996. R: A Language for Data Analysis and Graphics. J
781   Comput Graph Stat 5:299–314.

782   36.   Li H, Durbin R. 2009. Fast and accurate short read alignment with
783   Burrows-Wheeler transform. Bioinformatics 25:1754–1760.

784   37.   Broad Institute. 2016. Picard tools. https://broadinstitute.github.io/picard/.

785   38.   Alkan C, Coe BP, Eichler EE. 2011. GATK toolkit. Nat Rev Genet 12:363–76.

786   39.   Naeem R, Hidayah L, Preston MD, Clark TG, Pain A. 2014. SVAMP: Sequence
787   variation analysis, maps and phylogeny. Bioinformatics 30:2227–2229.

788   40.   Stamatakis A, Ludwig T, Meier H. 2005. RAxML-III: A fast program for
789   maximum likelihood-based inference of large phylogenetic trees. Bioinformatics
790   21:456–463.

791   41.   Wu M, Scott AJ. 2012. Phylogenomic analysis of bacterial and archaeal sequences
792   with AMPHORA2. Bioinformatics 28:1033–1034.

793   42.   Edgar RC. 2004. MUSCLE: a multiple sequence alignment method with reduced
794   time and space complexity. BMC Bioinformatics 5:113.

795   43.   Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. 2009. trimAl: A tool for
796   automated alignment trimming in large-scale phylogenetic analyses.
797   Bioinformatics 25:1972–1973.

44. G.Dobson., D.E.Minnikin., S.M.Minnikin., M.Parlett., M.Goodfellow., M.Ridell.,

M.Magnusson. 1985. Systematic analysis of complex mycobacterial lipids., p.

237–265. *In* Chemical Methods in Bacterial Systematics. Academic Press, London,

United Kingdom.

45. Haubold B, Pierstorff N, Möller F, Wiehe T. 2005. Genome comparison without

alignment using shortest unique substrings. BMC Bioinformatics 6.

46. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden

TL. 2009. BLAST+: Architecture and applications. BMC Bioinformatics 10.

47. Park H, Jang H, Kim C, Chung B, Chang CL, Soon Kew Park, Song S. 2000.

Detection and identification of mycobacteria by amplification of the internal

transcribed spacer regions with genus- and species-specific PCR primers. J Clin

Microbiol 38:4080–4085.

48. Leon P, Virginia W, Patricia B. 1989. Molecular analysis of the linear 2.3 kb

plasmid of maize mitochondria: Apparent capture of tRNA genes. Nucleic Acids

Res 17:4089–4099.

49. Kinashi H. 2011. Giant linear plasmids in Streptomyces: A treasure trove of

antibiotic biosynthetic clusters. J Antibiot (Tokyo).

50. Crespi M, Messens E, Caplan AB, van Montagu M, Desomer J. 1992. Fasciation

induction by the phytopathogen Rhodococcus fascians depends upon a linear

plasmid encoding a cytokinin synthase gene. EMBO J 11:795–804.

51. Picardeau M, Vincent V. 1997. Characterization of large linear plasmids in

mycobacteria. J Bacteriol 179:2753–2756.

52. Surtees JA, Funnell BE. 2003. Plasmid and Chromosome Traffic Control: How

ParA and ParB Drive Partition. Curr Top Dev Biol 56:145–180.

53. Zrimec J, Lapanje A. 2018. DNA structure at the plasmid origin-of-Transfer

indicates its potential transfer range. Sci Rep 8.

54. Ummels R, Abdallah AM, Kuiper V, Aâjoud A, Sparrius M, Naeem R, Spaink HP,

Bitter W, Pain A, Bitter W. 2014. Identification of a novel conjugative plasmid in

mycobacteria that requires both type IV and type VII secretion. MBio 5.

55. Uchiya KI, Takahashi H, Nakagawa T, Yagi T, Moriyama M, Inagaki T, Ichikawa
K, Nikai T, Ogawa K. 2015. Characterization of a novel plasmid, pMAH135, from
Mycobacterium avium subsp. hominissuis. PLoS One 10.

56. da Silva Rabello MC, Matsumoto CK, de Almeida LGP, Menendez MC, de
Oliveira RS, Silva RM, Garcia MJ, Leão SC. 2012. First description of natural and
experimental conjugation between mycobacteria mediated by a linear plasmid.
PLoS One 7.

57. Kim B-J, Kim B-R, Lee S-Y, Seok S-H, Kook Y-H, Kim B-J. 2013.
Whole-Genome Sequence of a Novel Species, Mycobacterium yongonense DSM
45126 T. Genome Announc 1:604–13.

58. Veyrier F, Pletzer D, Turenne C, Behr MA. 2009. Phylogenetic detection of
horizontal gene transfer during the step-wise genesis of Mycobacterium
tuberculosis. BMC Evol Biol 9.

59. van Ingen J, Kohl TA, Kranzer K, Hasse B, Keller PM, Katarzyna Szafrańska A,
Hillemann D, Chand M, Schreiber PW, Sommerstein R, Berger C, Genoni M,
Rüegg C, Troillet N, Widmer AF, Becker SL, Herrmann M, Eckmanns T, Haller S,
Höller C, Debast SB, Wolfhagen MJ, Hopman J, Kluytmans J, Langelaar M,
Notermans DW, ten Oever J, van den Barselaar P, Vonk ABA, Vos MC, Ahmed N,
Brown T, Crook D, Lamagni T, Phin N, Smith EG, Zambon M, Serr A, Götting T,
Ebner W, Thürmer A, Utpatel C, Spröer C, Bunk B, Nübel U, Bloemberg G V.,
Böttger EC, Niemann S, Wagner D, Sax H. 2017. Global outbreak of severe
Mycobacterium chimaera disease after cardiac surgery: a molecular
epidemiological study. Lancet Infect Dis 17:1033–1041.

60. Morgado SM, Marín MA, Freitas FS, Fonseca EL, Vicente ACP. 2017. Complete
plasmid sequence carrying type IV-like and type VII secretion systems from an
atypical mycobacteria strain. Mem Inst Oswaldo Cruz 112:514–516.

61. Houben ENG, Korotkov K V., Bitter W. 2014. Take five - Type VII secretion
systems of Mycobacteria. Biochim Biophys Acta - Mol Cell Res 1843:1707–1716.

62. Abdallah A, Pittius G, Nicolaas C, DiGiuseppe Champion P, Cox J, Luirink J,
Vandenbroucke-Grauls C, Appelmelk B, Bitter W. 2007. Type VII secretion -
Mycobacteria show the way. Nat Rev Microbiol 5:883–891.

63. Weerdenburg EM, Abdallah AM, Mitra S, De Punder K, Van der Wel NN, Bird S,
Appelmelk BJ, Bitter W, Van der Sar AM. 2012. ESX-5-deficient Mycobacterium
marinum is hypervirulent in adult zebrafish. Cell Microbiol 14:728–739.

64. Jia X, Yang L, Dong M, Chen S, Lv L, Cao D, Fu J, Yang T, Zhang J, Zhang X,
Shang Y, Wang G, Sheng Y, Huang H, Chen F. 2017. The Bioinformatics
Analysis of Comparative Genomics of Mycobacterium tuberculosis Complex
(MTBC) Provides Insight into Dissimilarities between Intraspecific Groups
Differing in Host Association, Virulence, and Epitope Diversity. Front Cell Infect
Microbiol 7.

65. Mahairas GG, Sabo PJ, Hickey MJ, Singh DC, Stover CK. 1996. Molecular
analysis of genetic differences between Mycobacterium bovis BCG and virulent M.
bovis. J Bacteriol 178:1274–1282.

66. Gordon S V., Brosch R, Billault A, Garnier T, Eiglmeier K, Cole ST. 1999.
Identification of variable regions in the genomes of tubercle bacilli using bacterial
artificial chromosome arrays. Mol Microbiol 32:643–655.

67. Van Ingen J, De Zwaan R, Dekhuijzen R, Boeree M, Van Soolingen D. 2009.
Region of difference 1 in nontuberculous Mycobacterium species adds a
phylogenetic and taxonomical character. J Bacteriol 191:5865–5867.

68. Kozak RA, Alexander DC, Liao R, Sherman DR, Behr MA. 2011. Region of
difference 2 contributes to virulence of mycobacterium tuberculosis. Infect Immun
79:59–66.

69. Ru H, Liu X, Lin C, Yang J, Chen F, Sun R, Zhang L, Liu J. 2017. The Impact of
Genome Region of Difference 4 (RD4) on Mycobacterial Virulence and BCG
Efficacy. Front Cell Infect Microbiol 7.

70. Brosch R, Gordon S V., Marmiesse M, Brodin P, Buchrieser C, Eiglmeier K,
Garnier T, Gutierrez C, Hewinson G, Kremer K, Parsons LM, Pym AS, Samper S,

van Soolingen D, Cole ST. 2002. A new evolutionary scenario for the

Mycobacterium tuberculosis complex. Proc Natl Acad Sci 99:3684–3689.

71. Alexander KA, Laver PN, Michel AL, Williams M, van Helden PD, Warren RM,

van Pittius NCG. 2010. Novel mycobacterium tuberculosis complex pathogen, M.

Mungi. Emerg Infect Dis 16:1296–1299.

72. M Cristina G, Brisse S, Brosch R, Fabre M, Omaïs B, Marmiesse M, Supply P,

Vincent V. 2005. Ancient origin and gene mosaicism of the progenitor of

Mycobacterium tuberculosis. PLoS Pathog 1:0055–0061.

73. Delogu G, Brennan MJ. 2001. Comparative Immune Response to PE and PE _

PGRS Antigens of Mycobacterium tuberculosis Comparative Immune Response to

PE and PE _ PGRS Antigens of Mycobacterium tuberculosis. Infect Immun

69:5606–5611.

74. Isom GL, Davies NJ, Chong ZS, Bryant JA, Jamshad M, Sharif M, Cunningham

AF, Knowles TJ, Chng SS, Cole JA, Henderson IR. 2017. MCE domain proteins:

Conserved inner membrane lipid-binding proteins required for outer membrane

homeostasis. Sci Rep 7.

75. Newton-Foot M, Warren RM, Sampson SL, Van Helden PD, Gey Van Pittius NC.

2016. The plasmid-mediated evolution of the mycobacterial ESX (Type VII)

secretion systems. BMC Evol Biol 16.

76. Slayden RA, Dawson CC, Cummings JE. 2018. Toxin-antitoxin systems and

regulatory mechanisms in Mycobacterium tuberculosis. Pathog Dis.

77. Cooper TF, Heinemann JA. 2000. Postsegregational killing does not increase

plasmid stability but acts to mediate the exclusion of competing plasmids. Proc

Natl Acad Sci 97:12643–12648.

78. Van Der Woude AD, Sarkar D, Bhatt A, Sparrius M, Raadsen SA, Boon L,

Geurtsen J, Van Der Sar AM, Luirink J, Houben ENG, Besra GS, Bitter W. 2012.

Unexpected link between lipooligosaccharide biosynthesis and surface protein

release in Mycobacterium marinum. J Biol Chem 287:20417–20429.

79. Boritsch EC, Frigui W, Cascioferro A, Malaga W, Etienne G, Laval F, Pawlik A,

Le Chevalier F, Orgeur M, Ma L, Bouchier C, Stinear TP, Supply P, Majlessi L,

Daffé M, Guilhot C, Brosch R. 2016. Pks5-recombination-mediated surface

remodelling in Mycobacterium tuberculosis emergence. Nat Microbiol 1.

80.   Cohen I, Parada C, Acosta-Gío E, Espitia C. 2014. The PGRS domain from

PE_PGRS33 of Mycobacterium tuberculosis is target of humoral immune response

in mice and humans. Front Immunol 5:1–9.

81.   Boritsch EC, Supply P, Honoré N, Seeman T, Stinear TP, Brosch R. 2014. A

glimpse into the past and predictions for the future: The molecular evolution of the

tuberculosis agent. Mol Microbiol.

82.   Geiss GK, Bumgarner RE, Birditt B, Dahl T, Dowidar N, Dunaway DL, Fell HP,

Ferree S, George RD, Grogan T, James JJ, Maysuria M, Mitton JD, Oliveri P,

Osborn JL, Peng T, Ratcliffe AL, Webster PJ, Davidson EH, Hood L. 2008. Direct

multiplexed measurement of gene expression with color-coded probe pairs. Nat

Biotechnol 26:317–325.

83.   Bowdish DME, Sakamoto K, Kim MJ, Kroos M, Mukhopadhyay S, Leifer CA,

Tryggvason K, Gordon S, Russell DG. 2009. MARCO, TLR2, and CD14 are

required for macrophage cytokine responses to mycobacterial trehalose dimycolate

and Mycobacterium tuberculosis. PLoS Pathog 5.

84.   Fujita Y, Naka T, McNeil MR, Yano I. 2005. Intact molecular characterization of

cord factor (trehalose 6,6′-dimycolate) from nine species of mycobacteria by

MALDI-TOF mass spectrometry. Microbiology 151:3403–3416.

85.   Stavrum R, Stavrum AK, Valvatne H, Riley LW, Ulvestad E, Jonassen I, Aßmus J,

Doherty TM, Grewal HMS. 2011. Modulation of transcriptional and inflammatory

responses in murine macrophages by the Mycobacterium tuberculosis Mammalian

cell entry (Mce) 1 complex. PLoS One 6.

86.   Katoch VM. 2004. Infections due to non-tuberculous mycobacteria (NTM). Indian

J Med Res.

87.   Griffith DE, Aksamit T, Brown-Elliott B a., Catanzaro A, Daley C, Gordin F,

Holland SM, Horsburgh R, Huitt G, Iademarco MF, Iseman M, Olivier K, Ruoss S,

von Reyn CF, Wallace RJ, Winthrop K. 2007. An Official ATS/IDSA Statement:

Diagnosis, Treatment, and Prevention of Nontuberculous Mycobacterial Diseases.

Am J Respir Crit Care Med 175:367–416.

88. Becq J, Gutierrez MC, Rosas-Magallanes V, Rauzier J, Gicquel B, Neyrolles O,

Deschavanne P. 2007. Contribution of horizontally acquired genomic islands to the

evolution of the tubercle bacilli. Mol Biol Evol 24:1861–1871.

89. Jamwal S V., Mehrotra P, Singh A, Siddiqui Z, Basu A, Rao KVS. 2016.

Mycobacterial escape from macrophage phagosomes to the cytoplasm represents

an alternate adaptation mechanism. Sci Rep 6.

Data Availability

The *M. riyadhense* dataset is available at European Nucleotide Archive (ENA) under the

study accession no. PRJEB32162. The assemblies are available with DOI

10.5281/zenodo.2873972.

Funding

Work in AP's laboratory is supported by the KAUST faculty baseline fund

(BAS/1/1020-01- 01).

Conflicts of Interest

The authors have no conflicts of interest to declare.

Acknowledgement

We would like to thank members of the Biological Core Lab (BCL) of King Abdullah

University of Science and Technology for sequencing the bacteria on the Illumina HiSeq

4000 and PacBio RSII platforms. The research protocol was approved by the Institutional

Review Board of King Fahad Medical City (Riyadh, Saudi Arabia; #16-345) and the

Institutional Biosafety and Bioethics Committee of King Abdullah University of Science

and Technology (Jeddah, Saudi Arabia; #18IBEC23).

Author Contributions Statement

AP conceived the comparative genomics part of the study, obtained the funding and

supervised the work; MG and AH conceived the clinical part of the study; MG and FA

wrote the clinical part of the text; FA, MG, TA, SF, AH, MAR, AR, and TS collected the

microbiological and clinical information; and AP, QG, SG, AB, and FA designed the

experiments. QG performed the data analysis and prepared the initial draft of the

manuscript, followed by edits from AP, SG, MG, SF and AB. QG, SM, ASm and JB

performed the transcriptome experiment. ASi and AB performed the TLC analysis. CN

and YS provided materials for the diagnostic markers and intellectual advice. All authors

have commented on various sections of the manuscript, which were finally curated and

incorporated in the final version by QG and AP.

998   Figure.1

B

1000 Figure.2

Figure.3

Figure.4

1007

1008

1009

1010

1011

1012

1027   Figure.5

1028

1029

1030

1031

1032

1033

1034

1035

1036

1037   Figure.6

Captions for the figures:

Figure.1: A phylogenetic tree of 148 species of mycobacteria showing close

relationship of *M. riyadhense* and MTBC. Twelve marker genes from the

chromosome were used for generating the diagram and were selected by

Amphora2, Dayhoff model selected by Modelgenerator applied to generate

this phylogeny using RaxML. For details please see in the Materials and

Methods. Circles on each branch indicates the bootstrap values (above 80%,

1,000 replicates). The relevant section of the phylogenetic tree has been

blown-up for better illustration. The complete high definition version of the

phylogenetic tree is available as Figure supplementary 7.

1048

1049 Figure.2: Comparison of the shared and unique gene orthologs amongst

1050 selected mycobacteria. Venn diagrams represent the overlap of gene

1051 orthologs between the (A) *M. riyadhense*, *M. tuberculosis* H37Rv, *M.*

1052 *marinum* M, *M. kansasii* 12478 and *M. szulgai* and (B) *M. riyadhense*

1053 MR226 and five species within the MTBC.

1054

Figure.3: A hierarchical clustering of the presence (black) and absence

(white) of *M. tuberculosis* H37Rv toxin/antitoxin orthologs in *M. riyadhense*,

M. marinum, *M. kansasii*, *M. shinjukuense*, *M. lacus*, *M. decipiens* and

MTBC species. The orange colored blocks represent if only one of the toxin

or antitoxin gene ortholog is present in a given pair of the toxin/antitoxin

system. The overall presence and absence toxin/antitoxin gene systems in *M.*

riyadhense are highlighted by the red outline. The name of the

toxin/antitoxin system are shown for each row on the right.

1063

Figure.4: LOS systems in *M. riyadhense* and other related mycobacteria and

2D-TLC analysis of polar lipids extracted from selected *M. riyadhense*

strains. (A) Genetic locus map of the LOS biosynthesis gene cluster from *M.*

riyadhense, *M. canettii* A and *M. tuberculosis* H37Rv (drawn to scale). The

arrows showing the direction of transcription and the genes are colored

according to the orthologous relationships. (B) Rough-dry colony

morphology (MR193) and (C) smooth morphology (MR226) of *M.*

riyadhense. Polar lipids from two known LOS producers, *M. marinum* (D)

and *M. kansasii* (E) are included to show the migration pattern of LOS

species in System E. (F) 2D-TLC analysis of polar lipids extracted from

select *M. riyadhense* rough strain or smooth (G) strain. A separate staining

with alpha-napthol also confirmed that this was a glycolipid species from the

same (H) rough and (I) smooth strain. (D), (E), (F) and (G) were charred

after staining with MPA, while (H) and (I) were charred after staining with

alpha napthol. LOS III from *M. riyadhense* is indicated by a solid arrow.

1079

Figure. 5 A circular heatmap demonstrating the host gene expression values

from RAW264.7 infections with *M. riyadhense* (MR), *M. kansasii* (MK) and

M. bovis BCG Denmark (Denm) over 3, 24 and 48 hours as determined by

the Myeloid Innate Immunity Gene Expression Panel (NanoString

Technologies, USA). Data represent Log2FC expression values compared to

the uninfected control from each time point. The gene sets of each clusters

are: A: Angiogenesis; B: Antigen Presentation; C: Cell Cycle and Apoptosis;

D: Cell Migration and Adhesion; E: Chemokine signaling; F: Complement

Activation; G: Cytokine Signaling; H: Differentiation and Maintenance of

Myeloid Cells; I: ECM remodeling; J: Fc Receptor Signaling; K: Growth

Factor Signaling; L: Interferon Signaling; M: Lymphocyte activation; N:

Metabolism; O: Pathogen Response; P: T-cell Activation and Checkpoint

Signaling; Q: TH1 Activation; R: TH2 Activation; S: TLR signaling. The

circles 1-9 represent RAW264.7 transcriptome response upon infections of 1:

Denm-3h; 2: Denm-24h; 3: Denm-48h; 4: MR-3h; 5: MK-3h; 6: MR-24h; 7:

MK-24h; 8: MR-48h; 9: MK-48h. The cluster dendrogram in the middle

show the hierarchical clustering of the individual gene expression datasets.

1097

Figure.6: Development of a rapid PCR-based diagnostic test for detection of

M. riyadhense. Agarose gel (2%) electrophoresis patterns of the PCR

products are shown as part of the diagnostic test for *M. riyadhense*.

Lane M: DNA Marker, Lane 1-8: Varies *M. riyadhense* strains (From left to

right: MR193, MR206, MR210, MR222, MR226, MR244, MR246,

MR1023), Lane 9-16: Varies *Mycobacterium* species (From left to right: *M.*

tuberculosis, *M. bovis*, *M. africanum*, *M. microti*, *M. oryis*, *M. kansasii*, *M.*

szulgai and *M. angelicum*) template, Template cocktail, mycobacterium

species (*M. tuberculosis*, *M. bovis*, *M. kansasii*, *M. marinum*, *M. szulgai*, *M.*

avium and *M. angelicum*) with (Lane 17, +) and without (Lane 18,-) *M.*

riyadhense MR226 gDNA template. Lane 19: Non-template control (NTC).

Upper Panel: MRDP (*M. riyadhense* diagnostic marker) set. Lower Panel B:

Mycobacterium genus specific primer ITS-T and mycom-R amplified with

mycobacterial gDNA.

1112
1113

1114

1115

1116

1117

1118   Tables:

Table 1 Clinical characteristics of the studied population

Smooth/Rough	Auramine rhodamine stain	Ziehl Neelsen stain	Therapy	
			Initial medication/duration	Continuous medication/duration
R	4+	2+	NA	NA
R	NA	1+	IMP/2M CLR/2M EMB/2M INH/1M PYZ/1M RIF/1M	CLR/1M TMT/4M
S	NA	Negative	EMB/18M CLR/18M	NA
S	NA	Negative	CLR/1M EMB/1M IMP/1M	EMB/2M IMP/2M
S	NA	Negative	NA	NA
S	NA	2+	CLR/1M RIF+INH/1M EMB/1M	MXF/4M
S	NA	Negative	RIF/1M CLR/21M EMB/1M	NA
S	NA	Negative	NA	NA

1119

1120

							Table.2 Comparison of general features of genome in Mycobacteria

Features	M. tuberculosis	M. africanum	M. bovis	M. canettii	M. riyadhense	M. kansasii	M. marinum
Genome Size (M)	4.41	4.39	4.26	4.48	Chromosome: 6.25 Plasmid pMRLP01: 0.55 Plasmid pMR01: 0.09	Chromosome: 6.43 Plasmid pMK12478: 0.14	Chromosome: 6.34 Plasmid pRAW: 0.11
GC content (%)	65.6	65.6	65.6	65.6	Chromosome: 65.35 Plasmid pMRLP01: 67.36 Plasmid pMR01: 65.69	Chromosome: 66.2 Plasmid pMK12478: 65.8	Chromosome: 65.8 Plasmid pRAW: 64.1
CDS	3,906	4,045	4,042	4,064	Chromosome: 5,490 Plasmid pMRLP01: 443 Plasmid pMR01: 88	Chromosome: 5,631 Plasmid pMK12478: 152	Chromosome: 5,149 Plasmid pRAW: 94
Gene Density(bp)	1101	1085	1032	1044	Chromosome: 1126 Plasmid pMRLP01: 1239 Plasmid pMR01: 1072	Chromosome: 1142 Plasmid pMK12478: 954	Chromosome: 1176 Plasmid pRAW: 1166
Conserved protein with assigned function	2855 (73%)	3078 (76%)	3072 (76%)	3125 (77%)	Chromosome: 4079 (74%) Plasmid pMRLP01: 217 (49%) Plasmid pMR01: 22 (25%)	Chromosome: 4468 (79%) Plasmid pMK12478:74 (49%)	Chromosome: 4091 (79%) Plasmid pRAW: 38 (40%)
Unique CDS	62 (1.59%)	25 (0.62%)	40 (0.99%)	83 (2.04%)	Chromosome:1059(19.29%) Plasmid pMRLP01:293 (66.14%) Plasmid pMR01:32 (36.36%)	Chromosome: 1,080 (19.18%) Plasmid pMK12478: 61 (40.13%)	Chromosome: 993 (19.29%) Plasmid pMK12478: 42 (44.68%)
tRNA	45	45	47	45	Chromosome: 51 Plasmid pMRLP01:1	Chromosome: 46 Plasmid: NA	Chromosome: 47 Plasmid: NA
rRNA	3	3	3	3	Chromosome: 3 Plasmid: NA	Chromosome: 3 Plasmid: NA	Chromosome: 3 Plasmid: NA
Transposase /integrase	53	68	56	98	Chromosome:110 Plasmid pMR01:1 Plasmid pMRLP01: 72	Chromosome:79 Plasmid pMK12478:3	Chromosome:41 Plasmid pRAW:4
Reference	1	2	3	4	MR226 This Study	5	6

1121

1122

1123

1124

1125   Supplementary Files:

1137

1138

1139

1140

1141

1142   Figure supplementary 1

1143

1144

1145

1146

1147

1148

1149

1150   Figure supplementary 2A

1151

1152

1153

1154   Figure supplementary 2B

1155

1156   Figure supplementary 2C

1157

1158

1159

1160

1161

1162

1163

1164

1165

1166

1167

1168

1169

1170

1171

1172

1173

1174

1175

1176   Figure supplementary 3

1177

1186       Figure supplementary 4A
1187

Figure supplementary 4B

1202

1203

1204

1205

1206
Figure supplementary 5

1207

1208

1209

1210

1211

1212

1213

1214

1215

1216
Figure supplementary 6A

1217

1218

1219

1220

1221

1222

1223

1224

1225

1226   Figure supplementary 6B

1227

1228

1229

1230

1231

1232

1233

1234

1235

1236

1237 Figure supplementary 7

1247   Figure supplementary 8

1257 Figure supplementary 9

1258
1259
1260

1261

1262

1263

1264

1265

1266

1267   Figure supplementary 10

1268
1269   Figure supplementary 11A

1270

1271   Figure supplementary 11B

1272

1273

1274

1275

1276

1277

1278

1279

1280

1281

1282

1283

1284

1285

1286

1287

1288   Figure supplementary 12

1289 Captions for the figure supplementary:

1290

Figure supplementary 1: Axial enhanced CT scan image of the chest from

the anonymous patient infected with *M. riyadhense* MR193. Multifocal

cavitating consolidation in both lungs predominantly involving the left upper

lobe associated with ill-defined ground-glass centrilobular nodules and

tree-in-bud appearance on both lungs were observed. The findings were

suggestive of tuberculosis.

1297

Figure supplementary 2**:** A comparison heatmap of predicted protein-coding

gene orthologs from *M. riyadhense* MR226 and other mycobacteria. The

orthologs were determined by ortho-MCL. The black and white spaces

denote presence and absences of an ortholog in a given species respectively.

(A) Comparison of chromosome-encoded genes in 152 mycobacterial

species. The box with red outline highlights the closely related species with

M. riyadhense based on ortholog groups clustering. Species from left to right

within box a are: *M. riyadhense* (published assembly GCA_002101845.1),

M. marinum, *M. angelicum*, *M. tuberculosis*, *M. pseudoshottsii*, *M. liflandii*,

M. gastri, *M. kansasii*, *M. ulcerans*, *M. shinjukuense*, *M. lacus*, *M. canettii*,

M. microti, *M. bovis*, *M. orygis*, *M. africanum*, *M. mungi*, *M. bovis BCG*.

The red arrows indicate the genomic regions that are absent in the assembly

GCA_002101845.1 while present in our MR226 assembly. (B) Comparison

of the linear plasmid pMRLP01-encoded genes in 152 mycobacterial species

genome assemblies. The box b with red outline highlights a region which

shared orthologs with pMRLP01 in *Mycobacterium tusciae*, *Mycobacterium*

aromaticivorans, *Mycobacterium llatzerense*, *Mycobacterium obuense*,

Mycobacterium novocastrense and *Mycobacterium holsaticum*; (C)

Comparison of the circular plasmid pMR01-encoded 77 genes in 152

mycobacterial species genome assemblies. The box c with red outline

highlights the cluster of the pRAW-like plasmids.

1319

Figure supplementary 3: Circular map of pMR01. The circles show from

outside to inside (1-12) BlastN results against the pMR01 of various

Mycobacterium plasmids. The corresponding plasmids used are listed on the

right panel. The innermost circle (13) represents pMR01 with the location of

the predicted protein coding genes. Red colour indicates the location of the

members of the *pe*/*ppe* gene family, blue indicates the *esx*-related genes and

all the other predicted genes are colored in grey.

1327

1328 Figure supplementary 4: Multiple alignment of 8 *M. riyadhense* assemblies

1329 using progressive Mauve. (A) The alignment of the 8 assemblies and (B) the

1330   alignment of the linear and circular plasmids from 8 strains. Circular

1331   plasmid are present in 3 strains and highlighted by boxes with black outlines.

1332

1333   Figure supplementary 5: Phylogenetic tree of *M. riyadhense* clinical isolates

1334   used in this study. The *M. riyadhense* phylogenetic tree was constructed with

1335   SNP data from 8 datasets called by GATK pipeline by RaxML with the

1336   TVM model. The circles on each branch indicates the bootstrap values

1337   (above 80%, 1,000 replicates).

1338

Figure supplementary 6: Genome alignments comparing selected Region of

Differences (RDs) in mycobacteria. The Artemis Comparison Tool (ACT)

was used to compare and visualize the annotated genome sequences against

the chosen mycobacteria. (A) The RD2 region of *M. riyadhense* MR226, *M.*

tuberculosis H37Rv and *M. bovis* BCG Pasteur (B) The RD12 region of *M.*

tuberculosis, *M. riyadhense* MR226 and *M. canettii* CIPT 140010059.

1345

Figure supplementary 7: A phylogenetic tree of 148 species of mycobacteria

showing close relationship of *M. riyadhense* and MTBC. Twelve marker

genes from the chromosome were used for generating the diagram and were

selected by Amphora2, Dayhoff model selected by Modelgenerator applied

to generate this phylogeny using RaxML. For details please see in the

Materials and Methods. The slow-growing mycobacteria are colored in light

green while the rapid-growing mycobacteria are colored in light purple.

Circles on each branch indicates the bootstrap values (above 80%, 1,000

replicates).

1355

Figure supplementary 8: The relative distribution of the functions of the

predicted protein-encoding genes were normalized by the genome size of

each genome and compared to the *M. riyadhense* MR226 strain. The

phylogenomic position of *M. riyadhense* within the *Mycobacterium* genus

was determined by concatemer sequences of 906 shared single copy genes

covering 296,124 amino acid positions. *Nocardia abscessus* was used as the

out group. The concatenated sequences were used to build the phylogenetic

tree with the LG+G+F model, which was selected by ModelGenerator. The

phylogenetic tree was generated by RaxML with 1,000 iterations and the

bootstrap values are shown above each branch.

1366

1367   Figure supplementary 9: Genetic locus map of the *pe-pgrs33* gene cluster

1368   from *M. marinum*, *M. kansasii*, *M. riyadhense*, *M. canettii* and *M.*

1369   *tuberculosis* (drawn to scale). The genes are shown with arrows and are

1370   colored according to the orthologs. The deletion event was highlighted with

1371   red arrows while the insertion event in green arrows.

1372

1373   Figure supplementary 10: Comparison of different gene clusters that encode

1374   type VII secretion systems in the *M. riyadhense* MR226 strain. The genes

1375   are shown with arrows and are colored according to the orthologs. The color

1376   codes for the figure are presented in the key. The black arrows indicate

1377   region-specific genes.

1378

Figure supplementary 11: TLR and NFkB pathway responses across the

three infections. Clustering of Nanostring data from infections with *M.*

riyadhense (MR), *M. kansasii* (MK) and *M. bovis* BCG Denmark (Denm)

over 3, 24 and 48 hrs showed commonality and variation in the

transcriptional responses. Panel A and B show selected genes involved in the

TLR (A) and NFkB (B) responses from the overall 754 gene panel used in

this study.

1386

1387   Figure supplementary 12: Basic statistics of Unique K-mers of *M.*

1388   *riyadhense* across 152 mycobacteria genome assemblies.

1389

1390

Table 1. Comparison of *M. riyadhense* strain	MR210	MR222
	6,528,955	6,533,138
	6,034,715	5,960,707
	2	7
	0	0
	P	P
	A	P
	56	42
	5,800	5,852
	5,420	5,353

Supplementary Table 2. The SNPs analysis of different samples using MR226 as the reference

	MR193	MR206	MR210	MR222	MR244	MR246	MR1023
Non-coding SNPS	2,550	160	2,478	2,727	160	192	165
Synonymous SNPs	16,574	1,107	15,757	16,959	1,250	1,138	1,120
Non-synonymous SNPs	3,446	227	3,161	3,287	266	250	232

1391

1392

Supplementary Table 3. *M. marinum* LOS locus genes orthologs shared with *M. riyadhense* strains

	MR1023	MR193	MR206	MR210	MR222	MR226	MR244	MR246
MMAR_1008	MR1023_04688	MR193_04455	MR206_04685	MR210_04486	MR222_04444	MR_04617	MR244_04622	MR246_04625
MMAR_2302	MR1023_03305	MR193_03158	MR206_03308	MR210_03179	MR222_03113	MR_03260	MR244_03260	MR246_03263
MMAR_2303	MR1023_03304	MR193_03157	MR206_03307	MR210_03178	MR222_03112	MR_03259	MR244_03259	MR246_03262
MMAR_2304	MR1023_03303	MR193_03156	MR206_03306	MR210_03177	MR222_03111	MR_03258	MR244_03258	MR246_03261
MMAR_2305	MR1023_03302	MR193_03155	MR206_03305	MR210_03176	MR222_03110	MR_03257	MR244_03257	MR246_03260
MMAR_2306	MR1023_03301	MR193_03154	MR206_03304	MR210_03175	MR222_03109	MR_03256	MR244_03256	MR246_03259
MMAR_2307	MR1023_03295	MR193_03148	MR206_03298	MR210_03167	MR222_03103	MR_03250	MR244_03250	MR246_03253
MMAR_2308								
MMAR_2309	MR1023_03299	MR193_03152	MR206_03302	MR210_03173	MR222_03107	MR_03254	MR244_03254	MR246_03257
MMAR_2310	MR1023_03298	MR193_03151	MR206_03301	MR210_03172	MR222_03106	MR_03253	MR244_03253	MR246_03256
MMAR_2311	MR1023_03297	MR193_03150	MR206_03300	MR210_03171	MR222_03105	MR_03252	MR244_03252	MR246_03255
MMAR_2312				MR210_03166				
MMAR_2313 (*losA*)		MR193_03128						
MMAR_2314	MR1023_00361	MR193_03129	MR206_00366	MR210_00381	MR222_00356	MR_00367	MR244_00362	MR246_00362
MMAR_2315								
MMAR_2316	MR1023_03023	MR193_02856	MR206_03023	MR210_01388,MR210_02888	MR222_02834	MR_02982	MR244_02976	MR246_02979
MMAR_2317	MR1023_05374	MR193_05138	MR206_05362	MR210_05150	MR222_05107	MR_05289	MR244_05292	MR246_05296
MMAR_2318		MR193_03130						
MMAR_2319								
MMAR_2320 *wecE*1								
MMAR_2321		MR193_03145						
MMAR_2322		MR193_03146						
MMAR_2323								
MMAR_2324								
MMAR_2325		MR193_03147						
MMAR_2326								
MMAR_2327	MR1023_03300	MR193_03153	MR206_03303	MR210_03174	MR222_03108	MR_03255	MR244_03255	MR246_03258
MMAR_2328								
MMAR_2329	MR1023_03443	MR193_03296	MR206_03447	MR210_03317	MR222_03253	MR_03400	MR244_03395	MR246_03398
MMAR_2330	MR1023_00875,MR1023_02733	MR193_00637	MR206_00875,MR206_02735	MR210_00864,MR210_02597	MR222_00847	MR_00870,MR_02697	MR244_00866,MR244_0	MR246_00867,MR246_02692
MMAR_2331								
MMAR_2332	MR1023_03970	MR193_03792	MR206_03973	MR210_03161	MR222_03768	MR_03910	MR244_03914	MR246_03918
MMAR_2333	MR1023_03293	MR193_03123	MR206_03296	MR210_03160	MR222_03101	MR_03248	MR244_03248	MR246_03251
MMAR_2334				MR210_03159				
MMAR_2335		MR193_04282		MR210_03158				
MMAR_2336 (*galE6*)								
MMAR_2337	MR1023_03292	MR193_03122	MR206_03295	MR210_03156	MR222_03100	MR_03247	MR244_03247	MR246_03250
MMAR_2338				MR210_03153				
MMAR_2339	MR1023_04010	MR193_03832	MR206_04012	MR210_03873	MR222_03809	MR_03949	MR244_03953	MR246_03957
MMAR_2340 (*pks*1	MR1023_03273	MR193_03102	MR206_03274	MR210_03130	MR222_03081	MR_03228	MR244_03228	MR246_03231
MMAR_2341 (*fadD25*)	MR1023_03277	MR193_03106	MR206_03912	MR210_03134	MR222_03085	MR_03232	MR244_03232	MR246_03235
MMAR_2342	MR1023_03276	MR193_03105	MR206_03277	MR210_03133	MR222_03084	MR_03231	MR244_03231	MR246_03234
MMAR_2343	MR1023_03271	MR193_03100	MR206_03272	MR210_03128	MR222_03079	MR_03226	MR244_03226	MR246_03229
MMAR_2344	MR1023_03270	MR193_03099	MR206_03271	MR210_03127	MR222_03078	MR_03225	MR244_03225	MR246_03228
MMAR_2345	MR1023_03268	MR193_03097	MR206_03269	MR210_03125	MR222_03076	MR_03223	MR244_03223	MR246_03226
MMAR_2346	MR1023_03266	MR193_03095	MR206_03267	MR210_03123	MR222_03073	MR_03221	MR244_03221	MR246_03224
MMAR_2347								
MMAR_2348	MR1023_03264	MR193_03093	MR206_03265	MR210_03121	MR222_03071	MR_03219	MR244_03219	MR246_03222
MMAR_2349	MR1023_03278,MR1023_03285	MR193_03107,MR193_03	MR206_03281,MR206_03288	MR210_03135,MR210_03142	MR222_03086,MR222_0	MR_03233,MR_03240	MR244_03233,MR244_0	MR246_03236,MR246_03243
MMAR_2350	MR1023_02109	MR193_02040	MR206_02111	MR210_02028		MR_02086	MR244_02076	MR246_02077
MMAR_2351								
MMAR_2352	MR1023_01970	MR193_01915	MR206_01970	MR210_01904	MR222_01910	MR_01948	MR244_01936	MR246_01937
MMAR_2353	MR1023_03275,MR1023_03279	MR193_03104,MR193_03	MR206_03276,MR206_03282	MR210_03132,MR210_03136	MR222_03083,MR222_0	MR_03230,MR_03234	MR244_03230,MR244_0	MR246_03233,MR246_03237
MMAR_2354	MR1023_02623	MR193_02508	MR206_02626	MR210_02487	MR222_02487	MR_02589	MR244_02580	MR246_02583
MMAR_2403								
MMAR_2404								
MMAR_2405 (*cpdB*)								
MMAR_2406								
MMAR_2555 (*papA3*)								
MMAR_4419								
MMAR_5170	MR1023_00312	MR193_00322	MR206_00317	MR210_00331	MR222_00307	MR_00318	MR244_00313	MR246_00313