1 Insights into ancestry and adaptive evolution of the *Mycobacterium tuberculosis* complex

2 from analysis of the emerging pathogen *Mycobacterium riyadhense*

Authors

Qingtian Guan[1], Musa Garbati[2,a], Sara Mfarrej[1], Talal AlMutairi[2], Alicia Smyth[3&], Albel

Singh[4&], Shamsudeen Fagbo[5&], John A. Browne[3], Muhammad Amin urRahman[2], Alya

Alruwaili[2], Anwar Hoosen[2,b], Chie Nakajima[6,7], Yasuhiko Suzuki[6,7], Apoorva Bhatt[4],

Stephen V. Gordon[3,6], Faisal AlAsmari[2], A. Pain[1,6*]

[1]*Pathogen Genomics Laboratory, BESE Division, King Abdullah University of Science*

and Technology (KAUST), Thuwal-Jeddah, Kingdom of Saudi Arabia; [2]*King Fahad*

Medical City（KFMC), Riyadh, Saudi Arabia; [3]*UCD School of Veterinary Medicine,*

University College Dublin, Dublin, D04 W6F6, Ireland; [4]*Institute of Microbiology and*

Infection, School of Biosciences, University of Birmingham, Edgbaston, Birmingham,

UK;[5]*One Health Unit, Executive Directorate for Surveillance and Response, Saudi*

Center for Disease Prevention and Control, Riyadh, Saudi Arabia; [6]*Global Institution for*

Collaborative Research and Education, Hokkaido University, Kita 20 Nishi 10, Kita-ku,

Sapporo, Japan;[7]Research Center for Zoonosis Control, Hokkaido University, Kita 20

Nishi 10, Kita-ku, Sapporo, Japan.

Key words: *Mycobacterium riyadhense*; Evolution of *Mycobacterium tuberculosis*

complex; Phylogeny; Virulence factors; Diagnostic marker.

[a]Current address: Infectious Diseases Unit, Department of Medicine, University of

Maiduguri, Nigeria

[&]These authors contributed equally to the work

[b]Current address: Pathcare/Vermaak & Partners Pathologists, Pretoria, South Africa.

*For correspondence
Arnab Pain (arnab.pain@kaust.edu.sa)

Current evolutionary scenarios posit the emergence of *Mycobacterium tuberculosis*,

the deadliest bacterial pathogen for humans globally, from an environmental

saprophyte through a cumulative process of genome adaptation. *Mycobacterium*

***riyadhense* is a novel non-tuberculous mycobacterium (NTM) that is being**

increasingly isolated from human clinical cases with tuberculosis (TB)-like

symptoms in various parts of the world. We provide evidence here that *M.*

***riyadhense* is likely a 'missing link' in the evolution of *Mycobacterium tuberculosis*.**

To elucidate the genomic hallmarks that define the evolutionary relationship

between *M. riyadhense* and other mycobacterial species, including members of the

Mycobacterium tuberculosis* complex (MTBC), eight clinical isolates of *M. riyadhense

were sequenced and analysed. We show, among other features, that *M. riyadhense*

shares a large number of conserved orthologues with the MTBC; contains linear

and circular plasmids carrying type IV and type VII secretion systems; and shows

expansion of toxin/anti-toxin pairs. We conclude that *M. riyadhense* is an emerging

mycobacterial pathogen that shares a common ancestor with members of the MTBC

and that can serve as an experimental model to study the evolution and pathogenesis

of tubercle bacilli.

The *Mycobacterium tuberculosis* complex (MTBC) is a group of genetically related

pathogens that cause tuberculosis (TB) in mammalian species. The hallmark member,

Mycobacterium tuberculosis, is the single most deadly pathogen, causing over 1.6 million

deaths globally in 2017. Current evolutionary scenarios posit the evolution of the MTBC

from an environmental saprophyte through a cumulative process of genome adaptation.

Such scenarios envisage intermediate mycobacterial species with increasing pathogenic

potential for humans, the vestiges of which should be present in extant mycobacterial

species. Comparative genomic analyses between the MTBC members and opportunistic

mycobacterial pathogens may therefore reveal the key evolutionary steps involved in

emergence of the MTBC, as well as illuminating virulence mechanisms across

mycobacterial pathogens as a whole.

Non-tuberculous mycobacteria (NTMs), including *Mycobacterium riyadhense* (MR), are

ubiquitous, naturally occurring environmental bacteria commonly found in water and

soil[1,2]. A wide range of animal and environmental sources (aquaria, swimming pools) act

as reservoirs for NTMs, and several human disease outbreaks caused by exposure to

environmental NTMs have been described[3,4,5,6]. With the ability to cause infections in

both immunocompromised[7] and immunocompetent[8] individuals, *M. riyadhense* has

positioned itself as a clinically important pulmonary pathogen since its discovery in 2009

[2] as a cause of pulmonary infection. The clinical and radiologic characteristics of

pulmonary infection caused by *M. riyadhense* are indistinguishable from those caused by

Mycobacterium tuberculosis, the most important human pathogen of the *M. tuberculosis*

complex (MTBC), which cause TB in humans and animals[2,8].

Similar to *M. tuberculosis*, *M. riyadhense* grows at 37°C and requires 2~3 weeks[2] to form

visible colonies on agar media. However, unlike *M. tuberculosis*, which is a common

worldwide pathogen transmitted directly from human to human with no known

environmental reservoirs[9], *M. riyadhense* infections are rare and are transmitted to

patients via contact with contaminated water[9] and soil[10], with no evidence of

human-to-human transmission yet reported. Infections with *M. riyadhense* have been

reported in Asia and Europe including, Bahrain, South Korea, France, Italy and Germany

[8,11,12], although most of the recent cases originated in patients from Saudi Arabia[13].

Indeed, the very first case of acute *M. riyadhense* infection was initially misdiagnosed as

a case of *M. tuberculosis* infection in a Saudi hospital using commercially available

diagnostic tests[14].

It is postulated that *M. tuberculosis* evolved from a free-living environmental ancestor

into an obligate pathogen[15,16]. Our current knowledge indicates that *Mycobacterium*

canettii is the most closely related obligate pathogenic species to the MTBC[17,18].

Infections with *M. canettii* are rare and found solely in people from the Horn of Africa,

with no environmental reservoir described to date[19].

Previous phylogenetic studies have suggested that *Mycobacterium kansasii*[20],

Mycobacterium marinum[21], *Mycobacterium lacus*[22], *Mycobacterium decipiens*[23],

Mycobacterium shinjukuense[24] and *M. riyadhense*[22,25] are closely related to the

free-living ancestor of the MTBC based on a single marker gene (e.g., *hsp65*, 16S). In

this evolutionary context, a comprehensive understanding of the genomic hallmarks that

drive the close phylogenomic relationship of *M. riyadhense* with the MTBC and NTMs is

of particular scientific interest.

In this study, we report the characterization of eight clinical isolates of *M. riyadhense*,

compared their genomes to members of the MTBC, and provided comprehensive insight

into the adaptive changes associated with the evolution of the MTBC from environmental

bacteria. We examined the lipid profiles of both rough and smooth variants of *M.*

riyadhense, compared them to those of other related mycobacteria, and analysed the

transcriptional response of immunity-related host genes in a murine macrophage infection

model with *M. riyadhense*, *M. kansasii* and *M. bovis* BCG.

Using genome information, we developed a simple PCR-based diagnostic test for the

rapid and accurate identification of *M. riyadhense* to minimize the risk of misdiagnosis in

a clinical setting.

Our analyses provide a comprehensive description of the hallmarks of *M. riyadhense* that

make it one of the closest known environmental relatives of the MTBC and that can serve

to illuminate studies into the evolution and pathogenesis in the MTBC.

RESULTS

Clinical manifestation and culture characteristics of *M. riyadhense* isolates

Between April 2011 and March 2017, eight clinical cases of infection with *M. riyadhense*

were recorded in male patients aged from 8-82 years (**Table 1**). In addition to HIV/AIDS,

most of the patients had multiple comorbidities, such as pulmonary and/or systemic

hypertension, malignancies, and diabetes mellitus (DM), with the lung being the major

site of disease (**Table 1**).

Bronchoalveolar lavage, endotracheal, sputum and lymph node biopsy specimens grew *M.*

riyadhense colonies when cultured in solid and liquid media using LJ agar and

Mycobacteria growth indicator tube (MGIT) broth, respectively, albeit with varying

times to positivity. Although no consistent findings existed for chest imaging, three out of

the eight patients presented with upper lobe consolidation, cavitation, ground-glass

opacities, 'tree-in-bud' appearance, hilar lymphadenopathy and pleural effusion

(**Supplementary Fig. 1**).

The isolates were subjected to susceptibility testing for antibiotics commonly used to

treat both typical and atypical infections. Isolates showed 100% susceptibility to rifampin

(RIF), rifabutin (RFB), ethambutol (EMB), clarithromycin (CLR), linezolid (LZD),

amikacin (AMK), moxifloxacin (MOXI), and trimethoprim-sulfamethoxazole

(TMP-SMX). Three out of the eight isolates were resistant to ciprofloxacin (CIP).

Assembly and annotation of the *M. riyadhense* MR226 genome

The comparison of different assemblies of all sequenced *M. riyadhense* strains is listed in

Supplementary Table 1. We obtained chromosomes of all eight isolates in single

contiguous sequences for genome comparison at a high resolution. The *M. riyadhense*

MR226 strain contains a 6,243,587 bp chromosome, a linear plasmid (pMRLP01) of

550,247 bp and a circular plasmid (pMR01) of 94,344 bp. The circular nature of the

chromosome and the pMR01 plasmid were demonstrated through Gepard[26].

As expected of a free-living opportunistic pathogen, the chromosome of *M. riyadhense* is

significantly larger than the chromosomes of the MTBC (**Table 2**). The number of genes

unique in a species showed that the members of the MTBC have a considerably lower

percentage of unique genes than *M. riyadhense* and other closely related NTMs (**Table**

2).

The comparison of the annotated protein-coding genes from the *M. riyadhense* MR226

strain to the genome assemblies of 152 *Mycobacterium* species (including 77 known

mycobacterial plasmids) shows that *M. riyadhense* forms a tight cluster with the MTBC

and some NTMs (**Supplementary Fig. 2a**, highlighted in box a). A total of 335 genes

were identified as unique from the orthologue group comparison, being present in only all

sequenced *M. riyadhense* isolates, including MR226, and the vast majority of these genes

belong to the PE/PPE family, which are thought to be involved in antigen variation and

are widely spread across the slow-growing species within the *Mycobacterium* genus.

Linear plasmids were first described in 1989 in maize (which has a linear

mitochondrion)[27] and have also been found in *Actinomycetales*, including *Streptomyces* [28],

Rhodococcus[29] and *Mycobacterium* species, such as *Mycobacterium xenopi*,

Mycobacterium branderi and *Mycobacterium celatum*. They are often accompanied by a

circular plasmid in the same host[30]. The linear plasmid pMRLP01 in *M. riyadhense*

contains a pair of partitioning genes (*parA*/*parB*) that are involved in active segregation

and thus stabilize the inheritance of the plasmid[31]. The latter are known to contribute to

genome evolution by active DNA transfer and exchange[32]. As is often the case for both

circular and linear large plasmids, a relatively higher proportion of pMRLP01 genes

(51%) have no known function compared to those of the main chromosome (26%). This

reinforces the notion that plasmids are an important route by which new genes are

introduced into the genome in *Mycobacteria*. Of the 443 predicted protein-coding genes

of pMRLP01, 118 have at least one orthologue on the main chromosome. Furthermore,

we observed several protein-coding genes in pMRLP01 that have orthologues in the

genomes of *Mycobacterium tusciae* JS617, *Mycobacterium aromaticivorans* JCM 16368,

Mycobacterium llatzerense, *Mycobacterium obuense*, *Mycobacterium novocastrense* and

Mycobacterium holsaticum (**Supplementary Fig. 2B**). This finding indicates that

pMRLP01 can be readily exchanged with other environmental mycobacterial species,

with implications for horizontal gene transfer in mycobacteria.

It is well known that plasmids are important "vehicles" for the exchange of genetic

material between bacteria or between chromosomes and extra-chromosomal plasmids. In

this study, we further identified a circular plasmid termed pMR01 in *M. riyadhense*

(**Supplementary Fig. 2C**, **Supplementary Fig. 3**). When compared with the circular

plasmids of other species, such as pRAW in *M. marinum*[33], pMAH135[34] and pMA100[35]

of *M. avium*, pMyong1 from *Mycobacterium yongonense*[36], pMK12478[37] from *M.*

kansasii and several plasmids from *Mycobacterium chimaera*[38], a high similarity was

observed. These plasmids all harbour both a TS4 and TS7, which are necessary for

conjugation[33,39], and facilitate the exchange of genetic material between different species

of slow-growing mycobacteria[33]. We therefore speculate that pMR01 is a novel

conjugative plasmid.

In mycobacteria, five type VII secretion systems have been described, named ESX-1 to

ESX-5[40,41]. An ESX-P5 locus on pMR01, which shows high similarity to the ESX-5 loci

on pMK12478, pRAW and pMAH135, is markedly different from the ESX-5 system

found on the main *M. riyadhense* chromosome. ESX-5 is involved in the secretion of PE

and PPE proteins in *M. tuberculosis* and is involved in modulating the host immune

responses to maintain a persistent infection[42]. The potential transmissibility of pMR01

and other pMR01-like plasmids may therefore play a role in the evolution of the ESX

systems in mycobacteria.

The progressive alignments (**Supplementary Fig. 4A**) of the assembled chromosomes

and plasmids (**Supplementary Fig. 4B**) of each *M. riyadhense* strain show that the

chromosomes are relatively conserved; however, the linear plasmids present in all eight

sequenced isolates are diverse from both structural and similarity perspectives, while the

pRAW-like plasmids are present in only the MR226, MR193 and MR222 strains.

The SNP-based phylogeny of the sequenced *M. riyadhense* isolates based on 43,136

polymorphic sites (**Supplementary Table 2**, **Supplementary Fig. 5**) indicates the

presence of two different clades of *M. riyadhense* among the clinical isolates sequenced

in this study. The nucleotide diversity between the MR222 clade is greater than the

diversity between *M. tuberculosis* strains[43], while it is smaller than that seen between *M.*

canettii strains[17], and the variation between the MR226 clade is comparable to the SNP

variation in *M. tuberculosis* strains.

Regions of difference (RDs) in *M. riyadhense*

The RDs were originally described as genomic regions present in virulent *M. bovis* and *M.*

tuberculosis but absent from the *M. bovis* BCG genome[44]. RD loci were subsequently

described across the MTBC[45] and contain functions believed to contribute to

pathogenicity[46,47,48] and evolution of MTBC species[49]. *M. riyadhense* was found to

harbour most of the RD loci (RD1, RD3-R11, R13-RD16) that are also intact in

M. tuberculosis, while 2 of the RDs show unique deletions, RD2[riyadh] (**Supplementary**

Fig. 6A) and RD12[riyadh] (**Supplementary Fig. 6B**).

RD2 was originally described as deleted in BCG vaccine strains.   Subsequently, it was

shown that disruption of RD2 in *M. tuberculosis* leads to decreased proliferation *in vivo*

and impaired modulation of the innate immune response[47]. The RD2 locus also has a

deletion from the *M. riyadhense* genome. It is a larger deletion than the originally

described RD2[BCG] as RD2[riyadh] contains 29 genes (*rv1971~rv2000*, location

2,216,498~2,246,766); eight genes within this locus (*rv1978*, *rv1979c*, *rv1980c*, *rv1981c*,

rv1983, *rv1984*, *rv1987*, *rv1988*) have orthologues elsewhere in the *M. riyadhense*

genome (*mr_05764*, *mr_05852*, *mr_02310*, *mr_02993*, *mr_00486*, *mr_02995*, *mr_02325*,

mr_02349, *mr_01747*), suggesting possible functional redundancy.

The RD12 locus shows deletions across MTBC members, including *M. bovis*,

Mycobacterium caprae and *M. orygis*[50], but it is present in other MTBC members. *M.*

canettii isolates (except group B[51]) also show an independent deletion at the RD12 locus

named RD12[can] (3,479,430~3,491,866, *rv3111~rv3126*), which is distinct from RD12[bovis]

(3,484,740~3,487,515, *rv3117~rv3121*). We identified another unique deletion at the

RD12 locus in *M. riyadhense*, designated RD12[riyadh], which encompasses a larger region

than RD[can] and RD12[bovis], encompassing *rv3108-rv3127* (3,477,171~3,492,150)

(**Supplementary Fig. 6B**). It is intriguing that multiple mycobacteria show independent

deletion events at the RD2 and RD12 loci, suggesting selective forces play a role in this

variation.

Comparative phylogeny of *M. riyadhense* with other Mycobacteria

The phylogenetic tree shows that the slow-growers and rapid-growers are separated into

two different clades and that fast-growers evolved from slow-growers (**Fig. 1**, the

detailed figure is shown in **Supplementary Fig. 7**). The overall topology of our tree is

remarkably similar to that of previously published phylogenetic trees[22]. In the tree,

M. riyadhense is located within the same clade as the obligate and opportunistic causal

organisms of mycobacterial diseases in humans that include the MTBC, *M. marinum*, *M.*

kansasii, *M. leprae* and related host-restricted mycobacteria with reduced genomes and

decreased survivability in the environment. Notably, *M. shinjukuense*, *M. lacus* and *M.*

riyadhense form a sub-clade that is phylogenetically close to the MTBC.

The *PE*/*PPE* family *mce* and *mce*-associated genes are known to be important for host

adaptation[52] and pathogenicity[53]. The *PE*/*PPE* family genes are enriched in the MTBC

members but also in *M. riyadhense* MR226 (278) and other pathogenic species, such as

M. kansasii (228) and *Mycobacterium ulcerans* (200). The number of *mce-* or

mce-associated genes has not significantly changed across mycobacterial genomes,

indicating that this group of genes is under an evolutionary constraint and plays

functional roles bridging both environmental and obligate pathogen lifestyles. Our results

agree with previous finding that during their evolution, the ESX systems were derived

from the ancestor ESX-4, as shown in **Supplementary Fig. 7** at the root node, and then

ESX-3, ESX-1, ESX-5 and ESX-2 evolved by horizontal transfer[54].

A comparative phylogenetic map based on 1,301 conserved proteins (**Supplementary Fig. 8**) reveals this downsizing of the genome and the dynamic changes in genome components. Certain functional categories of genes are relatively enriched during the evolution of MTBC, including protein metabolism, regulation and cell signalling, cell division and cell cycle.

M. riyadhense shares a larger number of orthologues (3,122) with *M. tuberculosis* than with *M. kansasii* (2,978), *M. marinum* (2,962) and *M. szulgai* (2,724) among the environmental mycobacteria that are closely related to the MTBC (**Fig. 2A**). A total of 134 orthologues are exclusively shared between *M. riyadhense* and *M. tuberculosis*, while the number of orthologues exclusively shared between *M. tuberculosis* and *M. kansasii* (30), *M. marinum* (48) and *M. szulgai* (18) is less (**Fig. 2A**). It is notable that genes from the phage-derived regions of RD3 and RD11 are shared exclusively between *M. riyadhense* and *M. tuberculosis*.

The comparative analysis of the orthologue groups of *M. riyadhense* and the MTBC is revealing. Firstly, 385 orthologue groups present across all MTBC species are absent from *M. riyadhense*. Secondly, 221 orthologues uniquely present in *M. riyadhense* are not found amongst the MTBC (**Fig. 2B**). This latter group of *M. riyadhense* unique

orthologues likely depict the constraints imposed by the free-living biology of *M.*

riyadhense which necessitates maintaining a broad functional repertoire to secure

environmental survival. In contrast, the obligate MTBC species have lost genes involved

in environmental survival but gained a large number of genes required for survival in the

in vivo environment.

A hallmark of *M. tuberculosis* infection is the ability to survive long-term in host

granulomas and develop a latent stage of infection. The molecular mechanisms and

cellular components that are involved in the persistence of *M. tuberculosis* are still poorly

understood, but several T/A systems have been implicated in the pathogenicity of *M.*

tuberculosis[55]. T/A systems were first found on plasmids or plasmid-derived

chromosomal loci where they promoted plasmid maintenance in bacterial populations[56],

but when compared to other mycobacteria, the MTBC are remarkable for the extensive

expansion of T/A systems. We thus compared the 79 pairs of T/A systems (belonging to

the HigAB, MazEF, ParDE, RelEF, VapBC and UCAT families) in *M. tuberculosis* with

the T/A pairs found in other members of the MTBC and NTMs. Based on the presence of

49 out of the 79 T/A orthologue pairs (**Fig. 3**), *M. riyadhense* appears more closely

related to the MTBC than to other mycobacteria, including *M. lacus, M. shinjukuense* and

M. decipiens. It can be hypothesized that the shared component of T/A pairs play a

functional role in the pathogenicity or persistence of *M. riyadhense* infection in a way

similar to that described for the importance of T/A pairs in *M. tuberculosis in vivo*

biology.

M. riyadhense strains produce a distinct pattern of LOSs

The LOSs are an important class of glycolipids that have previously been linked to

diverse mycobacterial phenotypes, such as colony morphology, secretion of PE/PPE

family proteins, and the pathogenicity of *M. marinum*[57]. It is noteworthy that we observed

both smooth (MR210, MR22, MR226, MR244, MR246, and MR1023) (**Fig. 4A**) and

rough (MR193 and MR206) (**Fig. 4B**) morphologies in *M. riyadhense* strains. We

therefore sought to first examine whether the genetic machinery for the production of

LOS is present in the *M. riyadhense* genome and then to follow up on the genome-level

predictions with lipid analyses of rough and smooth variants using TLC.

We observed that the *wecE* and *galE6* LOS genes are absent from the *M. riyadhense*

genome (**Table 3**, **Supplementary Table 3**). These genes are linked with the removal of

LOS II* and the production of LOS IV[57]. Thus, their absence is likely to cause an

accumulation of LOS II* and the lack of fully formed LOS IV, which have previously

been shown to increase the pathogenicity of *M. marinum*[57]. Furthermore, both the *pks5*

and *pap* genes in the LOS locus are intact in *M. riyadhense*, as is the case in *M. canettii*,

but not in *M. tuberculosis,* where the former is truncated and the latter deleted[58]. This

finding indicates that *pks5* recombination and *pap* deletion occurred in a common

ancestor of the MTBC after its differentiation from both *M. riyadhense* and *M. canettii.*

Remarkably, the *M. riyadhense* LOS gene locus layout is dissimilar to that in *M. canettii*,

M. tuberculosis, *M. kansasii* and *M. marinum*: indeed, exclusive rearrangements of this

locus in *M. riyadhense* were observed (**Fig. 4C**).

To correlate rough vs. smooth colony morphology with LOS production, we extracted

polar lipids from the strains and analysed them by 2D-TLC using solvent system E[59],

which is designed to separate phospholipids and LOSs. Charring of the TLC plates with

alpha-napthol revealed glycolipids, including the accumulation of a species that migrated

at a position similar to that of LOS III. This lipid was seen in only smooth strains; species

with migration patterns similar to LOS I and LOS II were observed, while LOS IV was

absent. This result was not unexpected because all *M. riyadhense* strains lack a functional

wecA gene, which is required for the extension of LOS II to LOS IV (**Fig. 4**).

Additionally, the relative levels of the predominant LOS species in *M. riyadhense* seem

to be quite high when compared to those seen in other LOS-producing mycobacteria (**Fig.**

4 (D)(E)). Conversely, the rough strains did not produce any glycolipids that migrated in

the positions corresponding to LOSs (**Fig. 4 (F)(H)**).

PE-PGRS33 locus and type VII secretion system of M. riyadhense

The *pe-pgrs33* (*rv1818c*) locus encodes the exported protein PE_PGRS33 and plays an

important role in the pathogenesis of *M. tuberculosis*[60]. A previous study[61] showed that

pe-pgrs33 is present in all MTBC members but not in *M. canettii*, which implies a

specific *pe_pgrs33* insertion event in the ancestor of MTBC strains. Genome comparison

of *M. riyadhense* with *M. tuberculosis*, *M. kansasii*, *M. marinum* and *M. canettii* provides

additional evidence that *M. riyadhense* is the missing link of the *pe-pgrs33*

deletion/insertion event. Our phylogeny strongly suggests that the deletion of *pe-pgrs33*

from *M. kansasii* and *M. marinum* occurred before the divergence of environmental

mycobacteria and the smooth tubercle bacilli (STB)/MTBC clade (**Supplementary Fig.**

9).

All 5 ESX systems (ESX1-ESX5) were found in the *M. riyadhense* genome

(**Supplementary Fig. 10**) but with minor modifications. Hence, the *eccA* and *eccB* genes

are absent from the ESX-2 system, while the *espACD* operon, which is essential for

secretion of virulence factors via ESX1, is also missing in *M. riyadhense*. The overall

gene arrangement of the ESX1-ESX5 loci is very similar in both *M. riyadhense* and

M. tuberculosis[40]. This conserved synteny reinforces the previous results that

phylogenetically *M. riyadhense* may represent an ancestral state to MTBC. As noted

before, the pMR01 plasmid also contains an extra ESX-P5 locus, which could indicate a

role for this plasmid in mediating pathogenicity (**Supplementary Fig. 4**).

Transcriptional response of murine macrophage cells upon *M. riyadhense* infection

Our genomic analysis of *M. riyadhense* revealed a range of genes and potential systems

that could play a role in host-pathogen interactions. We therefore sought to assess the

initial interaction of *M. riyadhense* with macrophages, using the RAW264.7 cell line of

murine origin as our experimental model. As comparator strains in our analysis, we

performed parallel infections with *M. kansasii*, an opportunistic pathogen that also

contains an orthologous ESX-1 system (**Fig. 1**, **Supplementary Fig. 10**), and *M. bovis*

BCG, the live TB vaccine that is attenuated through deletion of the ESX-1 system. These

comparisons allowed us to explore whether *M. riyadhense* triggered innate immune

responses were more similar to those triggered by *M. kansasii* or BCG or were

intermediate between the two.

To analyse the innate immune responses of the macrophages in the intracellular presence

or absence of these mycobacterial isolates, the transcriptional response was analysed

using a 754 probe NanoString Murine Myeloid Innate Immunity panel V2[62]   at 3, 24 and

48 hours post infection. These analyses revealed an expected commonality in the

responses to infection with all mycobacteria, such as induction of proinflammatory genes

through TLR signalling (e.g., upregulation of *IL1B*, *TNFA*, *CCL4*, *PTGS2* and *CXCL2*,

albeit to different absolute levels, (**Fig. 5**, **Supplementary Fig. 11A and 11B**). Distinct

responses triggered by BCG infection compared to *M. riyadhense* and *M. kansasii*

included, for example, upregulation of *MARCO* by BCG infection (**Supplementary Fig.**

11B); *MARCO* is involved in pathogen uptake via trehalose dimycolate, a lipid that is

known to show variation in structure between the MTBC and *M. kansasii*[63,64]. *CCL24* and

CXCL14, which are involved in the attraction of immune cells to the site of infection[65],

were upregulated to higher levels with *M. riyadhense* and *M. kansasii* than with BCG at

24 and 48 hours.

Overall, our analysis of the macrophage transcriptional profiles showed that the response

to infection with *M. riyadhense* and *M. kansasii* triggered more similar macrophage

transcriptional responses in comparison to the response produced by BCG infection

(**Supplementary Fig. 11**).

Developing a rapid PCR-based diagnostic marker for *M. riyadhense*

Due to the issues previously encountered in diagnosing *M. riyadhense* infections[2,8],

correct and prompt identification of cases upon presentation at healthcare units is of

paramount importance. We therefore sought to translate our knowledge on the genome

sequences into a PCR diagnostic test that could be used in a clinical microbiology setting

to distinguish *M. riyadhense* from other mycobacteria, including the members of the

MTBC.

By identifying unique K-mers ranging in size from 11 bp to 4,209 bp (**Supplementary**

Fig. 12) in the assembled genome compared to the genomes of 152 other mycobacterial

species, a primer set was developed targeting the *mr_00036*, *mr_00263*, *mr_00606* and

mr_01005 genes. The MRDP primer pair MRDP-F/MRDP-R amplified a single product

of 372 bp from each of the eight isolates of *M. riyadhense* (**Fig. 6**) but not from other

mycobacterial species, including *M. tuberculosis*, *M. bovis*, *M. kansasii*, *M. marinum*,

M. szulgai, *M. avium* and *Mycobacterium angelicum*. This result shows that the

MRDP-F/MRDP-R primers are highly specific to *M. riyadhense* and form the basis for a

simple diagnostic PCR that can inform appropriate treatment protocols.

DISCUSSION

M. riyadhense has become a clinically relevant NTM species globally[8,66]. Contrary to

prior publications on *M. riyadhense* that have been based primarily on clinical case

reports, here we present the largest and most comprehensive genomic study undertaken to

date on clinical *M. riyadhense* isolates. The eight new *M. riyadhense* strains sequenced in

this study originated from pulmonary infections with some having additional

extra-pulmonary involvement, which fulfilled the American Thoracic Society/ Infectious

Diseases Society of America (ATS/IDSA) criteria for NTM infection[67].

Our comparative analysis of *M. riyadhense* genomes with the MTBC and a large

collection of NTMs provide unequivocal evidence that *M. riyadhense* is one of the closest

known environmental mycobacteria species to the MTBC and forms a phyloclade with

M. lacus, *M. decipiens* and *M. shinjukuense*. Indeed, while our manuscript was in

preparation, independent work by[68] also showed the close phylogenetic relationship of *M.*

riyadhense to the MTBC, suggesting that it forms part of an MTB-associated phylotype.

Our analyses of multiple *M. riyadhense* isolates complements and extends the findings of

Sapriel and Brosch by revealing that expansion of T/A pairs, modification of secretion

systems, alterations in cell wall lipids, and plasmid-mediated horizontal acquisition of

new functionality all played key roles in the evolution of the MTBC. Our work hence

adds new insight into the evolution of the MTBC from free-living environmental bacteria

to obligate pathogens.

Our study shows that *M. riyadhense* shares a larger number of orthologues with

M. tuberculosis than *M. kansasii* and *M. marinum*, notably in the T/A gene family

(**Supplementary Fig. 6**). Forty-nine pairs of T/A gene orthologues were found in *M.*

riyadhense, far greater than the number of orthologues observed in any of the other

NTMs. The expansion of T/A genes among the MTBC offers additional evidence that

that suggested original acquisition of T/A modules into mycobacteria through lateral gene

transfer played a key role in the development of pathogenicity[21,69].

M. riyadhense strains appeared as both smooth and rough colony forms when grown on

solid LJ media. The observation of smooth and rough colony variants is seen in other

mycobacterial pathogens where it is linked to presence or absence of LOS; for example,

the presence or absence of LOS from *M. canettii* causes a transition from smooth to

rough colony variants, respectively, with rough variants showing increased virulence;

MTBC strains lack LOS, and it has been suggested that the removal of LOS was a key

event in the evolution of the MTBC species towards their current obligate pathogen status

[58]. In *M. riyadhense* smooth colony variants we observed the presence of LOS I and LOS

II but the absence of LOS IV. These biochemical observations agree with the genomic

prediction that *M. riyadhense* strains lack a functional *wecA* gene, which is required for

the extension of LOS II to LOS IV. Overall, our results show that *M. riyadhense* exhibits

a LOS production phenotype distinct from that of other LOS-producing mycobacteria.

The ESX Type VII secretion systems are key elements of mycobacterial virulence. All

the 5 ESX type VII secretion systems present in *M. tuberculosis*, and known to be

involved in virulence and pathogenicity, were found in *M. riyadhense* with very similar

gene arrangement. An additional ESX-P5 system was also found on the circular plasmid

pMR01. pMR01-like plasmids have been shown in many different NTMs, including *M.*

kansasii, *M. marinum*, *M. chimaera* and *M. avium*, indicating that these groups of

plasmids containing both TS4 and TS7 are conjugative plasmids. The extensive presence

of these plasmids may also explain the origin of the type VII secretion system [33]. ESAT-6

and CFP-10, which are secreted through ESX-1, are critical to phagosome perturbation

and manipulation of host macrophages induced by *M. tuberculosis*[70]. Linked to this, our

transcriptome analysis revealed that *M. riyadhense* and *M. kansasii* trigger similar overall

macrophage responses after infection when compared to *M. bovis* BCG (as a

representative of the MTBC). These data reinforce the phylogenetic relationship of

M. riyadhense with the MTBC species and its transitional status between an opportunistic

and an obligate pathogen.

The clinical presentation of our cases was by and large indistinguishable from disease

caused by *M. tuberculosis*, as reported earlier[7,8], but with a negative *M. tuberculosis* PCR.

Due to the relatively recent emergence of *M. riyadhense* as an important clinical

pathogen, and coupled with its misdiagnosis as *M. tuberculosis* by commercially

available kits, we developed an accurate set of diagnostic markers based on the genomic

datasets generated in this study. The primer sets accurately detect *M. riyadhense* in a

mixed cocktail of closely related mycobacteria: it can serve as part of an accurate and fast

diagnostic protocol in clinical settings thus reducing the need for strict isolation,

laborious contact tracing as well as the inappropriate use of TB antimicrobials. These

primers can be used in a global epidemiological survey of cases of *M. riyadhense*

infections in humans, animals and the environment. This may thus provide a more

complete picture of the epidemiology of *M. riyadhense* following a broad One Health

approach. It would, for example, be of clinical interest to see if *M. riyadhense* infections

occur in Africa and South America, for which no reports are available, or if *M.*

riyadhense is uniquely endemic in the Arabian Peninsula. Indeed, a limitation of this and

previous similar studies is that all isolates available for genomic studies are clinical. *M.*

riyadhense is an environmental pathogen that is most likely be found in animals sharing

the same niche with humans that were infected from the environment. As the natural

environmental reservoir of *M. riyadhense* remains unknown, we believe that systematic

screening of relevant environmental samples with the MRDP established in this study

may help to establish the natural habitat of this bacterium.

In conclusion, our study provides unprecedented insights into the ancestry and adaptive

evolution of the MTBC relative to extant NTM species and places *M. riyadhense* as one

of the closest environmental relatives to the MTBC. Our work provides compelling data

supporting the use of *M. riyadhense* as a novel mycobacterium for the study of virulence,

evolution and pathogenesis in the MTBC.

METHODS

Clinical reports and bacterial strains

Eight *M. riyadhense* strains were collected in Riyadh, Saudi Arabia, between June 2011 and March 2016 (**Table 1**) from patients with a positive culture for *M. riyadhense* isolated from the microbiology laboratory at the King Fahad Medical City (KFMC) in Riyadh, Saudi Arabia. The patient and sample data collected included demographic and clinical characteristics, age, sex, clinical features at presentation, presence of comorbidities, including HIV co-infection, antimicrobial susceptibility, initial and modified therapy, where applicable; and treatment outcome. Once an isolate was suspected to be an NTM, the samples were sent to the reference laboratory for full identification and antimicrobial susceptibility testing. Radiographic and pathologic data were also collected.

Culturing, DNA isolation and sequencing of bacteria

The *M. riyadhense* strains were grown on Lowenstein Jensen (LJ) slants at 37 °C for two weeks, DNA was extracted using a phenol-chloroform protocol [71], and the quality was measured by Qubit. Twenty micrograms of high-molecular-weight (HMW) DNA from the eight *M. riyadhense* strains was sequenced using a PacBio RSII sequencer (Pacific Biosciences, Menlo Park, USA) with a 10 kb library. An NEBNext Ultra II DNA library preparation kit (New England BioLabs, Massachusetts USA) was used to prepare libraries according to the manufacturer's instructions, and sequences from each library

were generated for all *M. riyadhense* strains using the Illumina HiSeq 4000 platform

(Illumina, San Diego, United States).

Genome assembly and annotation

The Illumina short reads were trimmed, and low-quality reads were removed by

Trimmomatic [72]. Eight consensus genomes based on each strain were assembled into

contigs with the PacBio long reads using the Canu assembler [73]. After assembly, the draft

genome was then corrected with short Illumina reads using Pilon [74] software. The

circularity of assemblies was checked by Gepard [26], and assemblies were annotated by

Prokka [75]. A circular map of the chromosome was compared with that of *M. tuberculosis*

H37Rv and visualized with BRIG [76]. The genome of the *M. riyadhense* MR226 strain

was used as a high-quality reference in this study.

Comparison of chromosomal and plasmid gene contents of *M. riyadhense* to those of

various *Mycobacterium* species

DNA sequences of 152 mycobacterial species and 77 mycobacterial plasmids were

obtained from the NCBI genome database and independently annotated by Prokka [75]. The

predicted protein sequences from the chromosome and each of the two plasmids

(pMRLP01 and pMR01) of the *M. riyadhense* MR226 strain were then compared with

the annotated genes from the rest of the mycobacterial species using orthoMCL [77] with a

50% identity cutoff and an inflation parameter of 1.5. The obtained orthologues were

visualized with the heatmap package in R [78]. A focused OrthoMCL comparison was

performed between (1) *M. riyadhense*, *M. marinum*, *M. kansasii*, *M. szulgai* and

M. tuberculosis and (2) *M. riyadhense* and five species from the MTBC, namely,

M. tuberculosis, *M. bovis*, *M. canettii*, *M. mungi*, and *M. africanum*.

SNP calling and phylogeny based on SNPs

The corrected Illumina reads were mapped using BWA[79] on to the *M. riyadhense* MR226

genome assembly. Picard tool[80] was used to clean SAM files, fix mate-pair information

and mark duplicates. SNPs were called for two iterations and filtered (QD<2.0, FS>60.0,

SOQ>4.0, ReadPosRankSum<-8.0) with Genome Analysis Toolkit (GATK)[81]. An

alignment file was generated by SVAMP[82], and phylogeny was generated by RaxML[83]

with the TVM model.

Phylogeny of *M. riyadhense*

The AMPHORA2[84] pipeline was used to identify protein sequences from 31 conserved

genes in the pangenome datasets and the *Mycobacterium* genomes available at NCBI or

JGI as at 1st January 2018. A total of 148 species were selected after applying a filter of at

least 10 marker genes to be detected in each of the assemblies. *Nocardia abscessus* was

used as the out-group. 12 genes, which have only one copy in each of the species (i.e.,

genes *rplA*, *rplB*, *rplF*, *rplM*, *rplN*, *rplP*, *rplS*, *rplT*, *rpmA*, *rpsJ*, *rpsM*, and *rpsS*), were

concatenated, aligned and trimmed. The phylogenetic tree was then constructed using

RaxML with the Dayhoff model, which is calculated with a script provided in

AMPHORA2 [84].

A whole-genome phylogenetic tree was constructed with the MTBC species (*M.*

tuberculosis, *M. bovis*, *M. canettii*, *M. mungi*, *Mycobacterium orygis*, *M. africanum*) and

with *M. kansasii*, *M. marinum*, *M. shinjukuense*, *Mycobacterium leprae*,

Mycobacterium smegmatis, *Mycobacterium parmense*, *Mycobacterium avium* and

Mycobacterium abscessus. The one-to-one orthologues of each species were obtained

using OrthoMCL and concatenated, then aligned with Muscle[85] and trimmed with

TrimAL[86]. The concatenated sequences were composed of 906 genes encoding 296,124

amino acids and were used to build a phylogenetic tree with the LG+G+F model, which

was selected by ModelGenerator. The phylogenetic tree was generated by RaxML[83].

Toxin/antitoxin systems, *mce/mce*-associated genes and ESX systems in *M.*

***riyadhense* and other mycobacteria**

One hundred fifty-eight toxin/antitoxin (T/A) proteins belonging to the VapBC, RelEF,

HigBA, MazEF, ParDE and UCAT families were downloaded from the NCBI protein

database. We identified the *M. tuberculosis* T/A orthologues from all of the 148 species

by OrthoMCL [77], and the orthologue groups were also examined by Blast+ 2.4.0. The

same pipeline was also applied for the MCE family, PhoPR, PE/PPE, PE-PGRS, and

ESX1-5.

Infection of the RAW 246.7 cell line with *M. riyadhense*, *M. kansasii* and *M. bovis*

BCG Denmark

The murine macrophage RAW264.7 cell line obtained from the American Type Culture

Collection (ATCC, Manassas, USA) was cultured in Dulbecco's modified Eagle's

medium (DMEM) (ThermoFisher Scientific, Waltham, USA) supplemented with 10%

FCS, streptomycin, and penicillin. *M. riyadhense*, *M. kansasii* (subtype I), and *M. bovis*

BCG Denmark strains were grown in Middlebrook 7H9 liquid medium after

single-colony isolation from LJ slants or 7H10 agar. 7H9 was supplemented with 10%

albumin, dextrose and catalase (ADC), while 7H10 was supplemented with oleic acid,

albumin, dextrose and catalase (OADC) in addition to 0.2% glycerol. Ready to use LJ

slants were provided by Saudi Prepared Media Laboratory (SPML, Riyadh, Saudi Arabia).

Before the infection, all bacterial cultures were centrifuged at 1,000xg for 10 minutes. The

supernatant was discarded, and 10-15 3 mm glass beads were added to the pellet and then

vortexed for 1 minute to break up clumps. Six ml of DMEM was added to the pellet and left

to rest for 5 minutes. The upper 5 ml of the suspension was then removed to a fresh 15 ml

falcon tube, which was then centrifuged for an additional 3 minutes at 200xg to remove

remaining bacterial clumps. The supernatant was then taken and passaged using a 26G

hypodermal syringe approximately 15 times to further break up any clumps of bacteria.

The optical density of the culture was measured again before the infection experiment.

RAW264.7 cells were seeded at 2 x 10^5 cells per well in 24-well flat-bottom tissue

culture plates 24 hours prior, reaching 5 x 10^5 cells per well. The DMEM over the cells

was removed, and the cells were washed once with phosphate-buffered saline (PBS). The

cells were infected by applying 1 ml of DMEM containing the prepared mycobacteria

from the former steps at an appropriate concentration to reach an MOI of 5:1, with

DMEM alone used for the control wells. The culture plates were then returned to the

incubator at 37°C with 5% CO_2 for 3 hours to allow for bacterial uptake by the

RAW264.7 cells. The supernatant was removed after 3 hours, and the infected cells were

washed with PBS to remove extracellular bacteria. Subsequently, the cells were

incubated in fresh DMEM with 10% FCS for 24 hours and 48 hours. For harvesting, 400

μl of TRIzol (ThermoFisher Scientific, Waltham, USA) was added to the wells at each

time point, and the adherent cells were scraped out and stored at -80°C for RNA

extraction. Each bacterial infection was performed in triplicate, in addition to the

non-infected controls. RNA was isolated from the samples using the Direct-zol[TM] RNA

Miniprep kit (Zymo Research, Irvine, USA) according to the manufacturer's instructions.

An Agilent RNA 6000 Nano kit was used to check the quality and quantity of the total

RNA. The NanoString murine nCounter Myeloid Innate Immunity Gene Expression

Panel (NanoString Technologies, USA) was used to assess transcript abundance across

infections and timepoints using the nCounter MAX Analysis System (NanoString

Technologies, Seattle, USA). The counts obtained were normalised using the nSolver[TM]

Advanced Analysis plugin (NanoString Technologies, Seattle, USA) using the geNorm

algorithm, and differential gene (DE) expression was analysed using multivariate linear

regression in nSolver[TM] software, with 0.05 as the p-value cutoff.

Thin layer chromatography (TLC) analysis of lipooligosaccharides (LOSs) in _M._

riyadhense, _M. kansasii_ and _M. marinum_

For TLC analysis, mycobacterial strains were grown at 30°C (_M. marinum_) or 37°C

(_M. smegmatis_, _M. kansasii_, _M. riyadhense_) on LJ slants, and after sufficient incubation,

grown cells were collected and washed once with PBS. Apolar and polar lipids were

extracted from the cell pellets using methods described by Dobson _et al_[59]. Polar lipids

were analysed by 2D-TLC using solvent system E, which is designed to separate

phospholipids and LOSs[59]. Glycolipids were visualized by charring following staining

with either molybdophophoric acid (MPA) or alpha-napthol (for glycolipids).

Diagnostic PCR markers for _M. riyadhense_

To develop diagnostic markers for *M. riyadhense* for potential use in clinical diagnosis as

well as environmental and animal studies, unique regions within the *M. riyadhense*

reference genome compared to that of 152 other mycobacterial species were detected

using Shustring[87]. These regions were also examined by OrthoMCL[77] and Blastn[88].

mr_00036, *mr_00263*, *mr_00606*, and *mr_01005* were selected as the amplification

targets. Two primers for each gene were designed in this study:

MRDP-MR_00036-F (5'-TTCGTTGTCGGTTTCGTCGC-3') and MRDP-MR_00036-R

(5'-GCGTCAGCTCCACCGAAAAC-3');

MRDP-MR_00263-F (5'-CCACCGCTGTTGGCGA-3') and MRDP-MR_00263-R

(5'-TTCGTCCCGTTGATCCCGTT-3');

MRDP-MR_00606-F   (5'-AACCTGCCCGATACGCACTT-3')   and

MRDP-MR_00606-R (5'-ACTGTTCCTCCGTGGGGTTG-3');

MRDP-MR_01005-F   (5'-GACTGTGGGGTAACGGTGGA-3')   and

MRDP-MR_01005-R (5'-CCGGTGATGTCGCCTACTCC-3').

PCR was performed in a 25 µl reaction volume with 12.5 µl of GoTaq ® Green Master

Mix (Promega, USA), 1 µl of 100 ng/µl gDNA, 1 µl with 10 nmol of forward and reverse

primers, 3 µl of dimethyl sulfoxide (DMSO) and 19 µl of nuclease-free water. The PCR

mixture was denatured for 5 minutes at 94°C; followed by 35 cycles of amplification

involving a denaturation step at 94°C for 30 seconds, a primer annealing step at 59°C for

45 seconds, and a primer extension step at 72°C for 45 seconds; and a final extension step

at 72°C for 7 minutes. The ITS-F/mycom-2 primer set[89], which is a *Mycobacterium*

genus-specific primer set, was used as a control, with amplification conditions as

described previously[89]. The products were electrophoresed in a 2% agarose gel for 60

minutes and visualized.

REFERENCES

1.   Falkinham, J. O. Ecology of nontuberculous mycobacteria-where do human
infections come from? *Semin. Respir. Crit. Care Med.* **34**, 95–102 (2013).

2.   van Ingen, J. *et al.* Mycobacterium riyadhense sp. nov., a non-tuberculous species
identified as Mycobacterium tuberculosis complex by a commercial line-probe
assay. *Int. J. Syst. Evol. Microbiol.* **59**, 1049–1053 (2009).

3.   Ding, L.-W., Lai, C.-C., Lee, L.-N., Huang, L.-M. & Hsueh, P.-R. Lymphadenitis
caused by non-tuberculous mycobacteria in a university hospital in   Taiwan:
predominance of rapidly growing mycobacteria and high recurrence rate. *J.*
Formos. Med. Assoc. **104**, 897–904 (2005).

4.   Panwalker, A. P. & Fuhse, E. Nosocomial Mycobacterium gordonae
pseudoinfection from contaminated ice machines. *Infect. Control* **7**, 67–70 (1986).

5.   Carbonne, A. *et al.* Outbreak of nontuberculous mycobacterial subcutaneous
infections related to multiple mesotherapy injections. *J. Clin. Microbiol.* **47**,
1961–1964 (2009).

6.   Singh, J. *et al.*   926. Outbreak of Invasive Nontuberculous Mycobacterium (NTM)
Infections Associated With a Pediatric Dental Practice . *Open Forum Infect. Dis.* **5**,
S29–S29 (2018).

7.   Garbati & Hakawi, A. M. *Mycobacterium Riyadhense Lung Infection in a Patient*

697   *with HIV/AIDS*. *Sub-Saharan African Journal of Medicine* **1**, (2014).

698   8.   Choi, J. I. *et al.* Lung infection caused by Mycobacterium riyadhense confused

699   with Mycobacterium tuberculosis: The first case in Korea. *Ann. Lab. Med.* **32**,

700   298–303 (2012).

701   9.   King, H. C. *et al.* Environmental reservoirs of pathogenic mycobacteria across the

702   Ethiopian biogeographical landscape. *PLoS One* **12**, (2017).

703   10.   Narendrula-Kotha, R. & Nkongolo, K. K. Microbial response to soil liming of

704   damaged ecosystems revealed by pyrosequencingand phospholipid fatty acid

705   analyses. *PLoS One* **12**, (2017).

706   11.   Godreuil, S. *et al.* Mycobacterium riyadhense pulmonary infection, France and

707   Bahrain. *Emerging Infectious Diseases* **18**, 176–178 (2012).

708   12.   Van der Werf, M. J. *et al.* Inventory study of non-tuberculous mycobacteria in the

709   European Union. *BMC Infect. Dis.* **14**, (2014).

710   13.   Althawadi, S. *et al.* Mycobacterium riyadhense. **23**, 2015–2017 (2017).

711   14.   Tortoli, E., Pecorari, M., Fabio, G., Messinò, M. & Fabio, A. Commercial DNA

712   probes for mycobacteria incorrectly identify a number of less frequently

713   encountered species. *Journal of Clinical Microbiology* **48**, 307–310 (2010).

714   15.   Murty DS, P. K. A Brief review on Ecology and Evolution of Mycobacteria.

715   *Mycobact. Dis.* **04**, (2014).

716   16.   Jang, J., Becq, J., Gicquel, B., Deschavanne, P. & Neyrolles, O. Horizontally

717   acquired genomic islands in the tubercle bacilli. *Trends Microbiol.* **16**, 303–308

718   (2008).

719   17.   Supply, P. *et al.* Genomic analysis of smooth tubercle bacilli provides insights into

720   ancestry and pathoadaptation of Mycobacterium tuberculosis. *Nat. Genet.* **45**,

721   172–179 (2013).

722   18.   Van Soolingen, D. *et al.* A Novel Pathogenic Taxon of the Mycobacterium

723   tuberculosis Complex, Canetti: Characterization of an Exceptional Isolate from

724   Africa. *Int. J. Syst. Bacteriol.* **47**, 1236–1245 (1997).

19.   Blouin, Y. *et al.* Significance of the Identification in the Horn of Africa of an
Exceptionally Deep Branching Mycobacterium tuberculosis Clade. *PLoS One* **7**,
e52841 (2012).

20.   Wang, J. *et al.* Insights on the emergence of Mycobacterium tuberculosis from the
analysis of Mycobacterium kansasii. *Genome Biol. Evol.* **7**, 856–870 (2015).

21.   Stinear, T. P. *et al.* Insights from the complete genome sequence of
Mycobacterium marinum on the evolution of Mycobacterium tuberculosis.
Genome Res. **18**, 729–41 (2008).

22.   Tortoli, E. *et al.* The new phylogeny of the genus Mycobacterium: The old and the
news. *Infect. Genet. Evol.* **56**, 19–25 (2017).

23.   Brown-Elliott, B. A. *et al.* Mycobacterium decipiens sp. nov., a new species
closely related to the Mycobacterium tuberculosis complex. *Int. J. Syst. Evol.*
Microbiol. **68**, 3557–3562 (2018).

24.   Saito, H. *et al.* Mycobacterium shinjukuense sp. nov., a slowly growing,
non-chromogenic species isolated from human clinical specimens. *Int. J. Syst. Evol.*
Microbiol. **61**, 1927–1932 (2011).

25.   Fedrizzi, T. *et al.* Genomic characterization of Nontuberculous Mycobacteria. *Sci.*
Rep. **7**, (2017).

26.   Krumsiek, J., Arnold, R. & Rattei, T. Gepard: A rapid and sensitive tool for
creating dotplots on genome scale. *Bioinformatics* **23**, 1026–1028 (2007).

27.   Leon, P., Virginia, W. & Patricia, B. Molecular analysis of the linear 2.3 kb
plasmid of maize mitochondria: Apparent capture of tRNA genes. *Nucleic Acids*
Res. **17**, 4089–4099 (1989).

28.   Kinashi, H. Giant linear plasmids in Streptomyces: A treasure trove of antibiotic
biosynthetic clusters. *Journal of Antibiotics* **64**, 19–25 (2011).

29.   Crespi, M., Messens, E., Caplan, A. B., van Montagu, M. & Desomer, J.
Fasciation induction by the phytopathogen Rhodococcus fascians depends upon a
linear plasmid encoding a cytokinin synthase gene. *EMBO J.* **11**, 795–804 (1992).

30.   Picardeau, M. & Vincent, V. Characterization of large linear plasmids in
mycobacteria. *J. Bacteriol.* **179**, 2753–2756 (1997).

31.   Surtees, J. A. & Funnell, B. E. Plasmid and Chromosome Traffic Control: How
ParA and ParB Drive Partition. *Curr. Top. Dev. Biol.* **56**, 145–180 (2003).

32.   Zrimec, J. & Lapanje, A. DNA structure at the plasmid origin-of-Transfer
indicates its potential transfer range. *Sci. Rep.* **8**, (2018).

33.   Ummels, R. *et al.* Identification of a novel conjugative plasmid in mycobacteria
that requires both type IV and type VII secretion. *MBio* **5**, (2014).

34.   Uchiya, K. I. *et al.* Characterization of a novel plasmid, pMAH135, from
Mycobacterium avium subsp. hominissuis. *PLoS One* **10**, (2015).

35.   da Silva Rabello, M. C. *et al.* First description of natural and experimental
conjugation between mycobacteria mediated by a linear plasmid. *PLoS One* **7**,
(2012).

36.   Kim, B.-J. *et al.* Whole-Genome Sequence of a Novel Species, Mycobacterium
yongonense DSM 45126 T. *Genome Announc* **1**, 604–13 (2013).

37.   Veyrier, F., Pletzer, D., Turenne, C. & Behr, M. A. Phylogenetic detection of
horizontal gene transfer during the step-wise genesis of Mycobacterium
tuberculosis. *BMC Evol. Biol.* **9**, (2009).

38.   van Ingen, J. *et al.* Global outbreak of severe Mycobacterium chimaera disease
after cardiac surgery: a molecular epidemiological study. *Lancet Infect. Dis.* **17**,
1033–1041 (2017).

39.   Morgado, S. M., Marín, M. A., Freitas, F. S., Fonseca, E. L. & Vicente, A. C. P.
Complete plasmid sequence carrying type IV-like and type VII secretion systems
from an atypical mycobacteria strain. *Mem. Inst. Oswaldo Cruz* **112**, 514–516
(2017).

40.   Houben, E. N. G., Korotkov, K. V. & Bitter, W. Take five - Type VII secretion
systems of Mycobacteria. *Biochim. Biophys. Acta - Mol. Cell Res.* **1843**,
1707–1716 (2014).

41.   Abdallah, A. *et al.* Type VII secretion - Mycobacteria show the way. *Nat. Rev.*
Microbiol. **5**, 883–891 (2007).

42.   Weerdenburg, E. M. *et al.* ESX-5-deficient Mycobacterium marinum is
hypervirulent in adult zebrafish. *Cell. Microbiol.* **14**, 728–739 (2012).

43.   Jia, X. *et al.* The Bioinformatics Analysis of Comparative Genomics of

Mycobacterium tuberculosis Complex (MTBC) Provides Insight into

Dissimilarities between Intraspecific Groups Differing in Host Association,

Virulence, and Epitope Diversity. *Front. Cell. Infect. Microbiol.* **7**, (2017).

44.   Mahairas, G. G., Sabo, P. J., Hickey, M. J., Singh, D. C. & Stover, C. K.

Molecular analysis of genetic differences between Mycobacterium bovis BCG and

virulent M. bovis. *J. Bacteriol.* **178**, 1274–1282 (1996).

45.   Gordon, S. V. *et al.* Identification of variable regions in the genomes of tubercle

bacilli using bacterial artificial chromosome arrays. *Mol. Microbiol.* **32**, 643–655

(1999).

46.   Van Ingen, J., De Zwaan, R., Dekhuijzen, R., Boeree, M. & Van Soolingen, D.

Region of difference 1 in nontuberculous Mycobacterium species adds a

phylogenetic and taxonomical character. *J. Bacteriol.* **191**, 5865–5867 (2009).

47.   Kozak, R. A., Alexander, D. C., Liao, R., Sherman, D. R. & Behr, M. A. Region of

difference 2 contributes to virulence of mycobacterium tuberculosis. *Infect. Immun.*

79, 59–66 (2011).

48.   Ru, H. *et al.* The Impact of Genome Region of Difference 4 (RD4) on

Mycobacterial Virulence and BCG Efficacy. *Front. Cell. Infect. Microbiol.* **7**,

(2017).

49.   Brosch, R. *et al.* A new evolutionary scenario for the Mycobacterium tuberculosis

complex. *Proc. Natl. Acad. Sci.* **99**, 3684–3689 (2002).

50.   Alexander, K. A. *et al.* Novel mycobacterium tuberculosis complex pathogen, M.

Mungi. *Emerg. Infect. Dis.* **16**, 1296–1299 (2010).

51.   M Cristina, G. *et al.* Ancient origin and gene mosaicism of the progenitor of

Mycobacterium tuberculosis. *PLoS Pathog.* **1**, 0055–0061 (2005).

52.   Delogu, G. & Brennan, M. J. Comparative Immune Response to PE and PE _

PGRS Antigens of Mycobacterium tuberculosis Comparative Immune Response to

PE and PE _ PGRS Antigens of Mycobacterium tuberculosis. *Infect. Immun.* **69**,

5606–5611 (2001).

53.   Isom, G. L. *et al.* MCE domain proteins: Conserved inner membrane lipid-binding

815   proteins required for outer membrane homeostasis. *Sci. Rep.* **7**, (2017).

816   54.   Newton-Foot, M., Warren, R. M., Sampson, S. L., Van Helden, P. D. & Gey Van

817   Pittius, N. C. The plasmid-mediated evolution of the mycobacterial ESX (Type VII)

818   secretion systems. *BMC Evol. Biol.* **16**, (2016).

819   55.   Slayden, R. A., Dawson, C. C. & Cummings, J. E. Toxin-antitoxin systems and

820   regulatory mechanisms in Mycobacterium tuberculosis. *Pathogens and Disease* **76**,

821   (2018).

822   56.   Cooper, T. F. & Heinemann, J. A. Postsegregational killing does not increase

823   plasmid stability but acts to mediate the exclusion of competing plasmids. *Proc.*

824   *Natl. Acad. Sci.* **97**, 12643–12648 (2000).

825   57.   Van Der Woude, A. D. *et al.* Unexpected link between lipooligosaccharide

826   biosynthesis and surface protein release in Mycobacterium marinum. *J. Biol. Chem.*

827   **287**, 20417–20429 (2012).

828   58.   Boritsch, E. C. *et al.* Pks5-recombination-mediated surface remodelling in

829   Mycobacterium tuberculosis emergence. *Nat. Microbiol.* **1**, (2016).

830   59.   G.Dobson. *et al.* Systematic analysis of complex mycobacterial lipids. in *Chemical*

831   *Methods in Bacterial Systematics* 237–265 (Academic Press, London, United

832   Kingdom., 1985).

833   60.   Cohen, I., Parada, C., Acosta-Gío, E. & Espitia, C. The PGRS domain from

834   PE_PGRS33 of Mycobacterium tuberculosis is target of humoral immune response

835   in mice and humans. *Front. Immunol.* **5**, 1–9 (2014).

836   61.   Boritsch, E. C. *et al.* A glimpse into the past and predictions for the future: The

837   molecular evolution of the tuberculosis agent. *Molecular Microbiology* **93**,

838   835–852 (2014).

839   62.   Geiss, G. K. *et al.* Direct multiplexed measurement of gene expression with

840   color-coded probe pairs. *Nat. Biotechnol.* **26**, 317–325 (2008).

841   63.   Bowdish, D. M. E. *et al.* MARCO, TLR2, and CD14 are required for macrophage

842   cytokine responses to mycobacterial trehalose dimycolate and Mycobacterium

843   tuberculosis. *PLoS Pathog.* **5**, (2009).

64. Fujita, Y., Naka, T., McNeil, M. R. & Yano, I. Intact molecular characterization of
cord factor (trehalose 6,6′-dimycolate) from nine species of mycobacteria by
MALDI-TOF mass spectrometry. *Microbiology* **151**, 3403–3416 (2005).

65. Stavrum, R. *et al.* Modulation of transcriptional and inflammatory responses in
murine macrophages by the Mycobacterium tuberculosis Mammalian cell entry
(Mce) 1 complex. *PLoS One* **6**, (2011).

66. Katoch, V. M. Infections due to non-tuberculous mycobacteria (NTM). *Indian
Journal of Medical Research* **120**, 290–304 (2004).

67. Griffith, D. E. *et al.* An Official ATS/IDSA Statement: Diagnosis, Treatment, and
Prevention of Nontuberculous Mycobacterial Diseases. *Am. J. Respir. Crit. Care
Med.* **175**, 367–416 (2007).

68. Sapriel, G. & Brosch, R. Shared pathogenomic patterns characterize a new
phylotype, revealing transition towards host-adaptation long before speciation of
Mycobacterium tuberculosis. *Genome Biol. Evol.* **11**, 2420–2438 (2019).

69. Becq, J. *et al.* Contribution of horizontally acquired genomic islands to the
evolution of the tubercle bacilli. *Mol. Biol. Evol.* **24**, 1861–1871 (2007).

70. Jamwal, S. V. *et al.* Mycobacterial escape from macrophage phagosomes to the
cytoplasm represents an alternate adaptation mechanism. *Sci. Rep.* **6**, (2016).

71. Belisle, J. T. & Sonnenberg, M. G. Isolation of genomic DNA from mycobacteria.
Methods Mol. Biol. **101**, 31–44 (1998).

72. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: A flexible trimmer for
Illumina sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

73. Koren, S. *et al.* Canu: Scalable and accurate long-read assembly via adaptive
κ-mer weighting and repeat separation. *Genome Res.* **27**, 722–736 (2017).

74. Walker, B. J. *et al.* Pilon: An integrated tool for comprehensive microbial variant
detection and genome assembly improvement. *PLoS One* **9**, (2014).

75. Seemann, T. Prokka: Rapid prokaryotic genome annotation. *Bioinformatics* **30**,
2068–2069 (2014).

76.   Alikhan, N. F., Petty, N. K., Ben Zakour, N. L. & Beatson, S. A. BLAST Ring
Image Generator (BRIG): Simple prokaryote genome comparisons. *BMC*
Genomics **12**, (2011).

77.   Li, L., Stoeckert, C. J. & Roos, D. S. OrthoMCL: Identification of ortholog groups
for eukaryotic genomes. *Genome Res.* **13**, 2178–2189 (2003).

78.   Ihaka, R. & Gentleman, R. R: A Language for Data Analysis and Graphics. *J.*
Comput. Graph. Stat. **5**, 299–314 (1996).

79.   Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler
transform. *Bioinformatics* **25**, 1754–1760 (2009).

80.   Broad Institute. Picard tools. *https://broadinstitute.github.io/picard/* (2016).
Available at:
https://broadinstitute.github.io/picard/%5Cnhttp://broadinstitute.github.io/picard/.

81.   Alkan, C., Coe, B. P. & Eichler, E. E. GATK toolkit. *Nat. Rev. Genet.* **12**, 363–76
(2011).

82.   Naeem, R., Hidayah, L., Preston, M. D., Clark, T. G. & Pain, A. SVAMP:
Sequence variation analysis, maps and phylogeny. *Bioinformatics* **30**, 2227–2229
(2014).

83.   Stamatakis, A., Ludwig, T. & Meier, H. RAxML-III: A fast program for maximum
likelihood-based inference of large phylogenetic trees. *Bioinformatics* **21**, 456–463
(2005).

84.   Wu, M. & Scott, A. J. Phylogenomic analysis of bacterial and archaeal sequences
with AMPHORA2. *Bioinformatics* **28**, 1033–1034 (2012).

85.   Edgar, R. C. MUSCLE: a multiple sequence alignment method with reduced time
and space complexity. *BMC Bioinformatics* **5**, 113 (2004).

86.   Capella-Gutiérrez, S., Silla-Martínez, J. M. & Gabaldón, T. trimAl: A tool for
automated alignment trimming in large-scale phylogenetic analyses.
Bioinformatics **25**, 1972–1973 (2009).

87.   Haubold, B., Pierstorff, N., Möller, F. & Wiehe, T. Genome comparison without
alignment using shortest unique substrings. *BMC Bioinformatics* **6**, (2005).

88.   Camacho, C. *et al.* BLAST+: Architecture and applications. *BMC Bioinformatics*
902   **10**, (2009).
903   89.   Park, H. *et al.* Detection and identification of mycobacteria by amplification of the
904   internal transcribed spacer regions with genus- and species-specific PCR primers.
905   *J. Clin. Microbiol.* **38**, 4080–4085 (2000).

DATA AVAILABILITY

The *M. riyadhense* dataset is available at European Nucleotide Archive (ENA) under the

study accession no. PRJEB32162. The assemblies are available with DOI

10.5281/zenodo.2873972.

FUNDING

Work in AP's laboratory is supported by the KAUST faculty baseline fund

(BAS/1/1020-01- 01). Work in SG and AB's laboratory was supported in part by BBSRC

award BB/N004574/1.

CONFLICTS OF INTEREST

The authors have no conflicts of interest to declare.

ACKNOWLEDGEMENT

We would like to thank members of the Biological Core Lab (BCL) of King Abdullah

University of Science and Technology for sequencing the bacteria on the Illumina HiSeq

4000 and PacBio RSII platforms. The research protocol was approved by the Institutional

Review Board of King Fahad Medical City (Riyadh, Saudi Arabia; #16-345) and the

Institutional Biosafety and Bioethics Committee of King Abdullah University of Science

and Technology (Jeddah, Saudi Arabia; #18IBEC23).

AUTHOR CONTRIBUTION STATEMENT

AP conceived the comparative genomics part of the study, obtained the funding and

supervised the work; MG and AH conceived the clinical part of the study; MG and FA

wrote the clinical part of the text; FA, MG, TA, SF, AH, MAR, AR, and TS collected the

microbiological and clinical information; and AP, QG, SG, AB, and FA designed the

experiments. QG performed the data analysis and prepared the initial draft of the

manuscript, followed by edits from AP, SG, MG, SF and AB. QG, SM, ASm and JB

performed the transcriptome experiment. ASi and AB performed the TLC analysis. CN

and YS provided materials for the diagnostic markers and intellectual advice. All authors

have commented on various sections of the manuscript, which were finally curated and

incorporated in the final version by QG and AP.

FIGURES

Fig. 1: A phylogenetic tree of 148 species of mycobacteria showing close relationship of *M. riyadhense* and MTBC.

Twelve marker genes from the chromosome were used for generating the diagram and were selected by Amphora2, Dayhoff model selected by Modelgenerator applied to generate this phylogeny using RaxML. For details please see in the Materials and Methods. Circles on each branch indicates the bootstrap values (above 80%, 1,000 replicates). The relevant section of the phylogenetic tree has been blown-up for better illustration. The complete high definition version of the phylogenetic tree is available as Supplementary Fig. 7.

953 **Fig. 2: Comparison of the shared and unique gene orthologs amongst selected**

954 **mycobacteria.**

955 Venn diagrams represent the overlap of gene orthologs between the (A) *M. riyadhense*, *M. tuberculosis*

956 H37Rv, *M. marinum* M, *M. kansasii* 12478 and *M. szulgai* and (B) *M. riyadhense* MR226 and five species

957 within the MTBC.

Fig. 3: A hierarchical clustering of the presence (black) and absence (white) of _M. tuberculosis_ H37Rv toxin/antitoxin orthologs in _M. riyadhense_, _M. marinum_, _M. kansasii_, _M. shinjukuense_, _M. lacus_, _M. decipiens_ and MTBC species.

The orange colored blocks represent if only one of the toxin or antitoxin gene ortholog is present in a given pair of the toxin/antitoxin system. The overall presence and absence toxin/antitoxin gene systems in _M. riyadhense_ are highlighted by the red outline. The name of the toxin/antitoxin system are shown for each row on the right.

Fig. 4: LOS systems in *M. riyadhense* and other related mycobacteria and 2D-TLC

analysis of polar lipids extracted from selected *M. riyadhense* strains.

(A) Genetic locus map of the LOS biosynthesis gene cluster from *M. riyadhense*, *M. canettii* A and *M.*

tuberculosis H37Rv (drawn to scale). The arrows showing the direction of transcription and the genes are

colored according to the orthologous relationships. (B) Rough-dry colony morphology (MR193) and (C)

smooth morphology (MR226) of *M. riyadhense*. Polar lipids from two known LOS producers, *M. marinum*

(D) and *M. kansasii* (E) are included to show the migration pattern of LOS species in System E. (F)

2D-TLC analysis of polar lipids extracted from select *M. riyadhense* rough strain or smooth (G) strain. A

separate staining with alpha-napthol also confirmed that this was a glycolipid species from the same (H)

rough and (I) smooth strain. (D), (E), (F) and (G) were charred after staining with MPA, while (H) and (I)

were charred after staining with alpha napthol. LOS III from *M. riyadhense* is indicated by a solid arrow.

Fig. 5 A circular heatmap demonstrating the host gene expression values from RAW264.7 infections with *M. riyadhense* (MR), *M. kansasii* (MK) and *M. bovis* BCG Denmark (Denm) over 3, 24 and 48 hours as determined by the Myeloid Innate Immunity Gene Expression Panel (NanoString Technologies, USA).

Data represent Log2FC expression values compared to the uninfected control from each time point. The gene sets of each clusters are: A: Angiogenesis; B: Antigen Presentation; C: Cell Cycle and Apoptosis; D: Cell Migration and Adhesion; E: Chemokine signaling; F: Complement Activation; G: Cytokine Signaling; H: Differentiation and Maintenance of Myeloid Cells; I: ECM remodeling; J: Fc Receptor Signaling; K: Growth Factor Signaling; L: Interferon Signaling; M: Lymphocyte activation; N: Metabolism; O: Pathogen Response; P: T-cell Activation and Checkpoint Signaling; Q: TH1 Activation; R: TH2 Activation; S: TLR signaling. The circles 1-9 represent RAW264.7 transcriptome response upon infections of 1: Denm-3h; 2: Denm-24h; 3: Denm-48h; 4: MR-3h; 5: MK-3h; 6: MR-24h; 7: MK-24h; 8: MR-48h; 9: MK-48h. The cluster dendrogram in the middle show the hierarchical clustering of the individual gene expression datasets.

Fig. 6: Development of a rapid PCR-based diagnostic test for detection of *M.*

***riyadhense*.**

Agarose gel (2%) electrophoresis patterns of the PCR products are shown as part of the diagnostic test for

M. riyadhense. Lane M: DNA Marker, Lane 1-8: Varies *M. riyadhense* strains (From left to right: MR193,

MR206, MR210, MR222, MR226, MR244, MR246, MR1023), Lane 9-16: Varies *Mycobacterium* species

(From left to right: *M. tuberculosis*, *M. bovis*, *M. africanum*, *M. microti*, *M. oryis*, *M. kansasii*, *M. szulgai*

and *M. angelicum*) template, Template cocktail, mycobacterium species (*M. tuberculosis*, *M. bovis*, *M.*

kansasii, *M. marinum*, *M. szulgai*, *M. avium* and *M. angelicum*) with (Lane 17, +) and without (Lane 18,-)

M. riyadhense MR226 gDNA template. Lane 19: Non-template control (NTC). Upper Panel: MRDP (*M.*

 riyadhense diagnostic marker) set. Lower Panel B: *Mycobacterium* genus specific primer ITS-T and

 mycom-R amplified with mycobacterial gDNA.

Table 1 Clinical characteristics of the studied population

	Age (Years)	Gender	HIV status	Other co-morbid conditions	Site(s) of the infection	Smooth/Rough	Auramine-rhodamine stain	Ziehl–Neelsen stain	Therapy Initial medication/duration	Therapy Continuous medication/duration	Radiographic Location of lesion (Chest CT)	Baseline CD4 count (X10^9/L)	Baseline Viral load (Copies/mL)	Treatment Outcome
MR193	82	M	Unknown	Smoking; DM; HTN; heart disease	Pulmonary	R	4+	2+	NA	NA	Left upper lobe consolidation	NA	N/A	Died
MR206	32	M	Negative	Eisenmenger's syndrome; pulmonary HTN	Pulmonary	R	NA	1+	IMP/2M CLR/2M EMB/2M INH/1M PYZ/1M RIF/1M	CLR/4M EMB/4M	Bilateral air space consolidation and nodular opacity	NA	N/A	Survived
MR210	66	M	Positive	PCP co-infection	Pulmonary	S	NA	Negative	EMB/18M CLR/18M	NA	Bilateral ground glass opacities	0.7	415,927	Survived
MR222	37	M	Positive	Lymphoblastic lymphoma	Pulmonary; LN, Abdominal	S	NA	Negtive	CLR/1M EMB/1M IMP/1M	EMB/2M IMP/2M	Multiple lung lesions with central cavitation	0	838	Died
MR226	8	M	Negative	Free	Pulmonary; LA	S	NA	Negtive	NA	NA	Multiple bilateral nodular infiltrates, more in the left upper lobe	NA	N/A	Survived
MR244	28	M	Positive	Pulmonary disease; hematologic disease	Pulmonary; LN, Abdominal; Peritoneal;	S	NA	2+	CLR/1M RIF+INH/1M EMB/1M	MXF/4M	Multiple bilateral lung nodules; cavitation left lower lobe	0.02	399,652	Survived
MR246	17	M	Negative	Pulmonary disease; chronic diarrhea	Pulmonary; Mediastinal LN	S	NA	Negative	RIF/17M CLR/21M EMB/17M	NA	Left upper lobe consolidation	NA	N/A	Survived
MR1023	47	M	Negative	DM; subclinical hypothyrodism	Pulmonary	S	NA	Negative	NA	NA	Bilateral nodular infiltrate; cavity,left upper lobe	NA	N/A	Survived

1025

1026

Table.2 Comparison of general features of genome in Mycobacteria

Features	M. tuberculosis	M. africanum	M. bovis	M. canettii	M. riyadhense	M. kansasii	M. marinum
Genome Size (M)	4.41	4.39	4.26	4.48	Chromosome: 6.25 Plasmid pMRLP01: 0.55 Plasmid pMR01: 0.09	Chromosome: 6.43 Plasmid pMK12478: 0.14	Chromosome: 6.34 Plasmid pRAW: 0.11
GC content (%)	65.6	65.6	65.6	65.6	Chromosome: 65.35 Plasmid pMRLP01: 67.36 Plasmid pMR01: 65.69	Chromosome: 66.2 Plasmid pMK12478: 65.8	Chromosome: 65.8 Plasmid pRAW: 64.1
CDS	3,906	4,045	4,042	4,064	Chromosome: 5,490 Plasmid pMRLP01: 443 Plasmid pMR01: 88	Chromosome: 5,631 Plasmid pMK12478: 152	Chromosome: 5,149 Plasmid pRAW: 94
Gene Density(bp)	1101	1085	1032	1044	Chromosome: 1126 Plasmid pMRLP01: 1239 Plasmid pMR01: 1072	Chromosome: 1142 Plasmid pMK12478: 954	Chromosome: 1176 Plasmid pRAW: 1166
Conserved protein with assigned function	2855 (73%)	3078 (76%)	3072 (76%)	3125 (77%)	Chromosome: 4079 (74%) Plasmid pMRLP01: 217 (49%) Plasmid pMR01: 22 (25%)	Chromosome: 4468 (79%) Plasmid pMK12478:74 (49%)	Chromosome: 4091 (79%) Plasmid pRAW: 38 (40%)
Unique CDS	62 (1.59%)	25 (0.62%)	40 (0.99%)	83 (2.04%)	Chromosome:1059(19.29%) Plasmid pMRLP01:293 (66.14%) Plasmid pMR01:32 (36.36%)	Chromosome: 1,080 (19.18%) Plasmid pMK12478: 61 (40.13%)	Chromosome: 993 (19.29%) Plasmid pMK12478: 42 (44.68%)
tRNA	45	45	47	45	Chromosome: 51 Plasmid pMRLP01:1	Chromosome: 46 Plasmid: NA	Chromosome: 47 Plasmid: NA
rRNA	3	3	3	3	Chromosome: 3 Plasmid: NA	Chromosome: 3 Plasmid: NA	Chromosome: 3 Plasmid: NA
Transposase /integrase	53	68	56	98	Chromosome:110 Plasmid pMR01:1 Plasmid pMRLP01: 72	Chromosome:79 Plasmid pMK12478:3	Chromosome:41 Plasmid pRAW:4
Reference	1	2	3	4	MR226 This Study	5	6

1027

1028

1029

Table 3 Presence and absence of the *M. marinum* orthologs related to LOS synthesis in *M. riyadhense* and other species.

	M. smegmatis	M. kansasii	M. canettii	M. riyadhense	M. tuberculosis H37Rv
MMAR_1008	P	P	P	P	P
MMAR_2307	A	P	A	P	A
MMAR_2313	A	P	P	P	P
MMAR_2319	A	A	A	A	A
MMAR_2320 (wecE)	A	P	P	A	A
MMAR_2327	A	P	P	P	P
MMAR_2332	A	A	A	A	A
MMAR_2336 (galE)	A	P	A	A	A
MMAR_2340 (pks)	P	P	P	P	T
MMAR_2341	P	P	P	P	P
MMAR_2343 (pap)	P	P	P	P	A
MMAR_2353	P	P	A	P	A
MMAR_5170 (whiB4)	P	P	P	P	P

P: Present; A: Absent; T: Truncated