**A Sense of Place: Transcriptomics Identifies Environmental Signatures**

**in Cabernet Sauvignon Berry Skins in the Late Stages of Ripening.**

**Grant R. Cramer[1][§], Noé Cochetel[1], Ryan Ghan[1], Agnès**
**Destrac-Irvine[2], and Serge Delrot[2]**

[1]Department of Biochemistry and Molecular Biology, University of Nevada,

Reno, Reno, NV 89557, USA

[2]UMR Ecophysiology and Grape Functional Genomics, Institut des Sciences

de la Vigne et du Vin, University of Bordeaux, Villenave d'Ornon, France

[§]Corresponding author

Tel: +1 (775) 784-4204, Fax: +1 (775) 784-1419

Email addresses:

GRC: cramer@unr.edu

RG: rghan@unr.nevada.edu

NC: noe.cochetel@gmail.com

AD-I: agnes.destrac-irvine@inra.fr

SD: serge.delrot@inra.fr

**Abstract**

**Background**

Grape berry ripening is influenced by climate, the main component of the

"terroir" of a place. Light and temperature are major factors in the vineyard

that affect berry development and fruit metabolite composition.

**Results**

To better understand the effect of "place" on the transcript abundance during

the late stages of berry ripening, Cabernet Sauvignon berries grown in

Bordeaux, France and Reno, NV USA were compared at similar sugar levels

(19 to 26 °Brix, total soluble solids (TSS)). Day temperatures were warmer

and night temperatures were cooler in Reno. TSS was lower in Bordeaux

berries compared to Reno at maturity levels considered optimum for harvest.

RNA-seq analysis identified 4,455 differentially expressed genes (DEGs)

between Bordeaux and Reno grape skins at 22°Brix. Weighted Gene

Coexpression Network Analysis for all expressed transcripts for all four °Brix

levels measured indicated that the majority (80%) of the gene expression

differed significantly between the two locations. Top gene ontology

categories for the common gene sets were translation, photosynthesis, DNA

metabolism and catabolism. Top DEG gene ontology categories at the

22°Brix level involved response to stimulus, biosynthesis and response to

stress. Some DEGs included genes encoding terpene synthases, cell wall

enzymes, kinases, transporters, transcription factors and photoreceptors.

Most circadian clock genes had higher transcript abundance in Bordeaux.

Bordeaux berries had higher transcript abundance with DEGs associated

with seed dormancy, light, auxin, ethylene signaling, powdery mildew

infection, phenylpropanoid, carotenoid and terpenoid metabolism, whereas Reno berries were enriched with DEGs involved in water deprivation, cold response, ABA signaling and iron homeostasis.

**Conclusions**

Transcript abundance profiles in the berry skins at maturity were highly dynamic. RNA-seq analysis identified a smaller (20% of total) common core set of ripening genes that appear not to depend on rootstock, vineyard management, plant age, soil and climatic conditions. Much of the gene expression differed between the two locations and could be associated with multiple differences in environmental conditions that may have affected the berries in the two locations; some of these genes may be potentially controlled in different ways by the vinegrower to adjust final berry composition and reach a desired result.

**Keywords**

abiotic stress, biotic stress, grape berry development, RNA-seq, transcriptomics, *Vitis vinifera* L.

**Background**

*Vitis vinifera* grapevines originated approximately 65 million years ago from Eurasia and have been cultivated for at least the last 8000 years for its fruit that are crushed to make wine [1]. Grapevines are now grown throughout the world in many kinds of environments.

Grape berry development is a complex process involving three

developmental phases and multiple hormones [2, 3]. It is in the latter ripening

phase that many compounds involved in flavor and aromas are synthesized,

conjugated or catabolized [2, 3]. Most of these compounds reside in the skin

of the berry and seem to develop in the very last stages of berry

development [4-6].

Aroma and flavor are important sensory components of wine. They are

derived from multiple classes of compounds in grapes including important

volatile compounds from the grape and from yeast metabolism during grape

fermentation [5, 6]. Each grape cultivar produces a unique set of volatile and

flavor compounds at varying concentration that represents its wine typicity or

typical cultivar characteristics [6]. Esters and terpenes are volatile compound

chemical classes largely responsible for the fruity and floral aromas in wines

[5, 6]. Esters are largely produced during yeast fermentation from grape-

derived products such as aliphatic alcohols and aldehydes [7, 8]. Grape

lipoxygenases are thought to provide the six carbon precursors from fatty

acids for the synthesis of the fruity aroma, hexyl acetate [8], in yeast during

wine fermentation. Terpenes mostly originate from the grapes and are found

in both the free and bound (glycosylated) forms. Both plant fatty acid and

terpenoid metabolism pathways are very sensitive to the environment [9-13].

Climate has large effects on berry development and composition [14-16].

Besides grape genetics other factors may influence metabolite composition

including the local grape berry microbiome [17], the soil type [15] and the

rootstock [18-22]. While there is evidence that rootstock can affect fruit

composition and transcript abundance, this effect appears to be minor

relative to other environmental factors [18, 19, 21]. Many cultural practices used by the grape grower may directly or indirectly affect the environment sensed by the grapevine (e.g. row orientation, planting density, pruning, leaf removal, etc.) [23, 24]. Temperature and light are major contributors to "terroir". The terroir term is used in the wine industry to acknowledge the influence of the environment on the grapevine and the distinctive characteristics that are contributed to the typicity of a wine [2, 14, 15, 25]. It includes the biotic, abiotic and soil environments as well as the viticultural practices. In the present work, we will use the term "place" to address all of the above except for the viticultural practices.

Recently, a transcriptomic approach was used to elucidate the common gene subnetworks of the late stages of berry development when grapes are normally harvested at their peak maturity [4]. One of the major subnetworks associated with ripening involved autophagy, catabolism, RNA splicing, proteolysis, chromosome organization and the circadian clock. An integrated model was constructed to link light sensing with the circadian clock highlighting the importance of the light environment on berry development. In this report, in order to get a better understanding of how much of the gene expression in Cabernet Sauvignon berry skin could be attributed to environmental influences, we tested the hypothesis that there would be significant differences in gene expression during the late stages of Cabernet Sauvignon berry ripening between two widely different locations: one in Reno, NV, USA (RNO) and the other in Bordeaux, France (BOD). The analysis revealed a core set of genes that did not depend on location, climate, vineyard management, grafting and soil properties. Also, the

analysis revealed keys genes that are differentially expressed between the

two locations and links some of these differences to the effects of

temperature and other environmental factors on aromatic and other quality-

trait-associated pathways. Many gene families were differentially expressed

and may provide useful levers for the vinegrower to adjust berry composition.

Among others, these families encompassed genes involved in amino acid

and phenylpropanoid metabolism, as well as aroma and flavor synthesis.

**Results**

*Background Data for a "Sense of Place"*

To test the hypothesis that the transcript abundance of grape berries during

the late stages of ripening differed in two locations with widely different

environmental conditions, we compared the transcript abundance of grape

berry skins in BOD and RNO. The vineyards were planted in 2004 and 2009

in RNO and BOD, respectively. BOD vines were grafted on to SO4 rootstock

and RNO vines were grown on their own roots. Both BOD and RNO used a

vertical shoot positioning trellis design. The environmental variables between

the two vineyard sites had a number of differences. BOD has a slightly more

northern latitude than RNO; consequently, day lengths were slightly longer in

BOD at the beginning of harvest and slightly shorter at the end of harvest

(Table 1). On the final harvest dates, the day length differed between the two

locations by about 30 min.

Average monthly maximum temperatures were warmer in RNO than BOD,

but minimum September temperatures were cooler (Table 1). This made for

an average daily day/night temperature differential of 10°C and 20°C during

the harvest periods for BOD and RNO, respectively. The day temperatures

were warmer by about 6°C and the night time temperatures were about 4°C

cooler in RNO.

The Reno site was much drier than the Bordeaux site (Table 1). September

monthly precipitation totals were 65.5 and 2.03 mm and had average relative

humidities of 74 and 34% for BOD and RNO, respectively. The soil at the

BOD site was a gravelly soil with a pH of 6.2 and the soil at the RNO site

was a deep sandy loam with a pH of 6.7. No pathogen, nutrient deficiency or

toxicity symptoms were evident in the vines.

*Transcriptomics*

The analysis of transcript profiles of Cabernet Sauvignon grapes harvested

in RNO in September of 2012 was previously described [4]. Individual berry

skins were separated immediately from the whole berry and the individual

sugar (°Brix) level of the berry was determined. Similarly, Cabernet

Sauvignon berry skins from BOD were harvested from the middle of

September in 2013 until the end of the first week of October (Table 1). Berry

skins were separated and analyzed in the same manner as in RNO with the

same Illumina technology. Grapes were harvested at a lower °Brix range in

BOD (19.5 to 22.5°Brix) than in RNO (20 to 26°Brix) because fruit maturity

for making wine is typically reached in the BOD region at a lower sugar level.

Transcript abundance of the RNA-seq reads from both RNO and BOD was

estimated using Salmon software (0.9.1) tools [26] with the V2 assembly and

the V3 gene model annotation of the PN40024 reference genome for

grapevine [27]. The TPM (transcripts per million) were computed for each

gene from each experimental replicate (n = 3) from berry skins at different

sugar levels ranging from 19 to 26°Brix (Additional File 1).

Functional annotation of the genes from the V2 assembly/V3 annotation of the

grapevine genome [27] was not yet performed at the time of our analyses, so

we performed our own functional annotation (Additional file 2) using different

versions of the Arabidopsis protein database, the V1 PN40024 *Vitis vinifera*

reference genome annotation and the refseq plant protein database (see

Materials and Methods). Among the 42,413 genes, approximately 91% of

them were annotated (*i.e.* matched to one of the database accessions).

Two approaches were used to analyze the transcriptomic data, one focused

on expression at one similar sugar level and one more comprehensive, using

all of the sugar levels. Since sugar levels were not exactly the same between

the two locations, differentially expressed genes (4445 DEGs at an FDR p-

value < 0.05) were identified between the two locations at the sugar level

closest to the 22°Brix level (21.5°Brix in BOD vs 22°Brix in RNO) using

DESeq2 [28] (Additional file 3). DEGs will refer to this set of genes throughout

this manuscript. Gene set enrichment analysis with topGO was determined for

these 4445 genes (Additional file 4) and the top gene ontology (GO) categories

for biological processes based on number of genes identified were response

to stimulus (1464 genes, corrected p-value = 3.13E-28), biological regulation

(1271 genes, corrected p-value = 5.00E-07) and biosynthetic process (1260

genes, corrected p-value = 3.50E-09). Another important and highly significant

category was response to stress (834 genes, corrected p-value = 1.03E-24).

There were 625 GO categories in total that were significantly enriched

(Additional file 4). The relationship between the top 25 GO categories can be

seen in Additional file 5. We use the term "significantly" throughout this text to mean statistically significant at or below a p-value of 0.05. Amongst the top stimulus subcategories with the largest number of genes were hormone stimulus (513 genes; 287 of which were related to ABA stimulus), abiotic stimulus (518 genes) and biotic stimulus (283 genes). There were many other environmental stimuli categories with significant gene set enrichment including response to light stimulus (216 genes), response to osmotic stress (154 genes), and response to temperature stimulus (153 genes).

Using a more powerful approach to finely distinguish the expression data for all sugar levels, Weighted Gene Coexpression Network Analysis (WGCNA) identified gene sets common to and differentially expressed between BOD and RNO. All transcripts for all °Brix levels (Additional file 1) were used in this analysis. Additional details of the analysis are described in the Materials and Methods section. Twenty-two modules or gene subnetworks were defined (Additional file 6) and a heat map was generated displaying the module-trait relationships (Additional file 7); the grey module is not a real module but a place to put all genes not fitting into a real module. Nine modules had similar profiles for BOD and RNO (p-value > 0.05); these included darkred, lightgreen, cyan, lightyellow, grey60, royalblue, magenta, midnightblue and tan. The expression of approximately 19.5% (5845 out of 29,976) of the genes were similar (not significantly different) in both locations. Gene set enrichment analysis was performed with genes that had a kME > 0.80 for all modules (Additional file 8). The similar gene sets in common with both locations with decreasing transcript abundance as sugar levels increased were enriched with the GO categories involving translation

(lightgreen module) and photosynthesis (cyan module). The common gene

sets with increasing transcript abundance as sugar levels increased were

enriched with the GO categories involving DNA metabolism (magenta

module) and catabolism (tan module).

Most modules were positively or negatively correlated with BOD and RNO

berries (e.g. blue, black, yellow, red, turquoise, etc.). The blue module was

one of the largest modules and consisted of 4018 genes; it had the most

positive and negative correlations for BOD and RNO, respectively (Additional

file 7). This gene set was similar to the DEGs defined by DESeq2 with the

largest differences between BOD and RNO at 22°Brix. Gene set enrichment

analysis of genes within the blue module having a kME of 0.80 or higher

(1137 genes) revealed many common GO categories with the DEGs

(Additional file 8); 71% (265 of 374) of the GO categories from the blue

module subset were also found in the 625 GO categories of the DEGs (42%

of total). Many GO categories were more highly enriched in the blue module

(e.g. organic acid metabolic process was 1.31E-28 and 1.46E-18 for the blue

module and DEGs, respectively).

Other enriched GO categories in the DEGs were enriched in the red

subnetwork (e.g. phenylpropanoid metabolism, response to stress and

defense response; Additional file 8). The WRKY75 (Vitvi01g01680) gene

was the top hub gene (kME = 0.97) in this subnetwork (Additional file 6).

WRKY75 is a transcription factor that positively regulates leaf senescence. It

is induced by ethylene, ROS and SA and is a direct target of EIN3 [29].

Thus, the WGCNA results, which utilized all of the transcripts from all °Brix

levels were consistent with the DESeq2 results that only compared transcript

abundances at 22°Brix. The WGCNA results complemented the DESeq2

results by identifying hub genes and gene subnetworks, and additional

transcripts and GO categories associated with the differences in transcript

abundance in berry skins between BOD and RNO.

**Transcriptomic profiles dynamically changing with sugar levels**

*DEGs with largest increases in transcript abundance between sugar levels*

As a first approach to examining the 4445 DEG dataset, differences between

the transcript abundance in berries with the lowest and highest °Brix levels at

the two locations were determined (Additional file 3). Eight examples of the

many DEGs with the largest transcript abundance differences from BOD

(MLP43, EXL2, JMT and MYB24) and RNO (EXL2, BSMT1, HB-7, UPI, JMT,

NAC073 and MYB24) were selected and are presented in Figure 1.

They include several transcription factors, methyl transferases, a lipid

transfer protein and a protease inhibitor. The major point of showing this plot

is to highlight the general trends of continuously increasing transcript

abundance with sugar levels for these genes; half of these genes start at

similar transcript abundance levels around 20°Brix for both BOD and RNO

berries and increase in transcript abundance at a higher rate in BOD grapes

as sugar levels increase. The other half increased at approximately the same

rate for both locations but had higher amounts at the BOD location at the

same sugar levels. These data were fitted by linear regression to compare

the slopes of the lines. The slopes were significantly higher for MLP43,

EXL2, JMT, and MYB24 for BOD as compared to RNO berries, but not for

the other four genes (data not shown).

Figure 1. Plots of the top genes from berry skins from Bordeaux and Reno with the highest

increase in transcript abundance (TPM) between the lowest and highest sugar levels (°Brix).

Values are the means ± SE (n = 3). Error bars not shown are smaller than the symbol. The

symbol legend is displayed in the figure. MLP43 is major lipid transfer protein 43; ERG-1 is a

squalene epoxidase involved in ERGgosterol biosynthesis; JMT is a jasmonate o-methyltransferase; MYB24 is a protein with a MYB domain; BSMT1 is a salicylate/benzoate carboxyl methyltransferase; HB-7 is a homeobox protein with a HB domain; NAC073 is a NAC protein with a NAC domain; and UPI is unusual serine protease inhibitor.

The significantly increased rate of change for transcript abundance in the BOD berries indicated that the berries in BOD may have ripened at a faster rate relative to sugar level. The DEGs in Figure 1 are involved in sterol and jasmonate biosynthesis, defense responses and transcription. The gene set enrichment analysis of the top 400 DEGs with the greatest increase in transcript abundance from the lowest sugar level to the highest sugar level provided some crucial insights about the differences between berries from the two locations. The top 400 DEGs for BOD berries were highly enriched in biosynthetic processes involving amino acid and phenylpropanoid metabolism (Additional File 9); defense responses, response to fungus and response to ethylene stimulus were other highly enriched categories. The top 400 DEGs for RNO berries were enriched in response to water deprivation and response to hormone stimulus (Additional File 10).

*DEGs with largest decreases in transcript abundance between sugar levels*
Eight examples of DEGs with the greatest decrease in transcript abundance with increasing sugar levels are presented in Figure 2. They include lipid and cell wall proteins (e.g. expansin, protease inhibitor, wax synthase and lipid transferase proteins) and an aquaporin (GAMMA-TIP). The data were fitted

2    **Figure 2.** Plots of the top genes from berry skins from Bordeaux and Reno with the highest

3    decrease in transcript abundance (TPM) between the lowest and highest sugar levels

4    (°Brix). Values are the means ± SE (n = 3). Error bars not shown are smaller than the

5    symbol. The symbol legend is displayed in the figure. GDSL is a sequence motif; GBD is a

galactose binding domain; CPI is cysteine protease inhibitor; GAMMA-TIP is a tonoplast

intrinsic protein; LTP is a lipid transfer protein; MLP28 is a major latex protein; FOP1 is a

bifunctional wax ester synthase/diacylglycerol acyltransferase; and EXPA6 is an expansin

protein.

to linear regressions and the slopes statistically compared between BOD and

RNO berries. In some cases, the slopes of the linear regression lines of the

DEGs were not statistically different (LTP, MLP28, GBD-like and FOP1); in

other cases, there are similar amounts of transcript abundance around

20°Brix, but there are significantly different slopes (GDSL Lipase, CPI,

GAMMA-TIP, and EXPA6). Again, there is a trend for the transcript

abundance of many of these genes to change more significantly in BOD

berries than RNO berries relative to sugar level.

*Differences in autophagy genes between BOD and RNO*

Berry ripening is a degradative process involving autophagy. As a general

rule, the transcript abundance of genes involved in autophagy increased with

sugar level and had a higher transcript abundance in BOD berries relative to

RNO berries at the same sugar level (Fig. 3), supporting the hypothesis that

BOD berries ripened faster than RNO berries relative to sugar level. In

addition, the transcript abundance of a NAC6 transcription factor

(Vitvi14g01985), whose ortholog in arabidopsis positively regulates aging-

induced cell death and senescence in leaves [30, 31], was twice as high in

BOD berries at 22°Brix (log2 value of 4.5 vs 3.5 in RNO berries).

**Figure 3.** The transcript abundance of some autophagy-related (ATG) genes. Data shown

are the means ± SE; n = 3. Error bars not shown are smaller than the symbol. The symbol

legend is displayed in the figure. APG9 is Autophagy 9.

***Aroma- and flavor-associated DEGs***

Many aroma and flavor-associated compounds are synthesized in the late

stages of berry ripening and sensitive to the environment. The major

metabolic pathways affecting flavor and aromas in grapes and wines include

the terpenoid, carotenoid, amino acid, and phenylpropanoid pathways [6].

These pathways were identified by topGO to be highly enriched in the DEGs

and/or the blue module. Some of the DEGs are involved in these pathways

and will be presented in the following subsections.

*Terpene synthase genes with the greatest transcript abundance differences*

*between BOD and RNO*

Cultivar differences in berries are often ascribed to differences in aroma

compounds. One of the main classes of cultivar specific aroma compounds

is the terpene group [32]. The transcript abundance of a number of terpene

synthases were higher in BOD berries as compared to RNO berries (Fig. 4).

All but one of these (terpene synthase 55; TPS55) increased in transcript

abundance with increasing sugar levels. TPS35 is a β-ocimine synthase (β-

ocimine is a main component of snapdragon flower aroma [33]). TPS 26 is a

Cubebol/δ-Cadinene synthase, TPS 4 and 10 are (E)-α-bergamotene

synthases, and TPS 07, 14 and 28 are germacrene-D synthases; these

enzymes produce sesquiterpenes found in essential plant oils (see [32] and

references therein for the function of all of these terpenoid genes). TPS55 is

a linalool/nerolidol synthase which synthesizes acyclic terpene alcohols;

linalool contributes significantly to the floral aromas of grape berries and

wines. TPS68 is a copalyl diphosphate synthase involved in diterpenoid

biosynthesis and TPS69 is an ent-kaurene synthase. Both TPS68 and 69 are

diterpene synthases and are part of the ent-kaurene biosynthesis pathway.

*Other terpenoid and carotenoid metabolism-related genes*

Carotenoid metabolism is another biosynthetic pathway that contributes to

flavor and aroma in grapes [34]. There are a number of key genes that

contribute to terpenoid and carotenoid metabolism that have a higher

transcript abundance (Additional file 3) in BOD berries as compared to RNO

berries. For example, DXR (1-deoxy-D-xylulose 5-phosphate

**Figure 4.** Expression profiles of some terpene synthase (TPS) genes. Data shown are the

means ± SE; n = 3. Error bars not shown are smaller than the symbol. The symbol legend is

displayed in the figure.

reductoisomerase, Vitvi17g00816) catalyzes the first committed step and

HDS (4-hydroxy-3-methylbut-2-en-1-yl diphosphate synthase;

Vitvi06g00286) enzyme controls the penultimate steps of the biosynthesis of

isopentenyl diphosphate (IPP) and dimethylallyl diphosphate (DMAPP) via

the methylerythritol 4-phosphate (MEP) pathway. Another example is PSY

(Vitvi04g02011) that encodes phytoene synthase; PSY is the rate-limiting

enzyme in the carotenoid biosynthetic pathway.

*Amino acid and phenylpropanoid metabolism genes*

Amino acids contribute to the aroma and flavor of grapes and wines [7]. The

amino acid metabolism functional GO category is highly enriched in the

group of DEGs between BOD and RNO (Additional file 4) and more

specifically in the top 400 BOD DEGs (Additional file 9). Some examples of

genes involved in amino acid metabolism that have a higher transcript

abundance in BOD berries (see Fig. 5) are phenylalanine ammonia lyase 4

(*PAL4*, Vitvi16g00054 and 9 other paralogs), which catalyzes the first step in

phenylpropanoid biosynthesis, branched-chain-amino-acid aminotransferase

3 (*BCAT3*, Vitvi00g00789), which is involved in isoleucine, leucine and valine

biosynthesis, 3-deoxy-D-arabino-heptulosonate 7-phosphate synthase 1

(*DHS1*, Vitvi07g03049), which catalyzes the first committed step in aromatic

amino acid biosynthesis), and tyrosine aminotransferase 7 (*TAT7*;

**Figure 5.** The transcript abundance of some DEGs that are involved in the amino acid

metabolism. Data shown are the means ± SE; n = 3. Error bars not shown are smaller than

the symbol. The symbol legend is displayed in the figure. PAL4 is a phenylalanine ammonia

lyase; DHS1 is a 3-deoxy-D-arabino-heptulosonate 7-phosphate synthase; BCAT3 is a

branched-chain-amino-acid aminotransferase; and TAT7 is a tyrosine aminotransferase.

Vitvi19g00186), which is involved in tyrosine and phenylalanine metabolism.

Included in this group was a large number of stilbene synthases (STS),

which are part of the phenylpropanoid pathway; these STSs had a higher

transcript abundance in BOD berries as compared to RNO berries, with very

similar transcript abundance profiles to PAL4 (see Additional file 11 for

examples).

***DEGs associated with abiotic stimuli***

*Light-responsive genes*

In a previous analysis, WGCNA defined a circadian clock subnetwork that was highly connected to transcript abundance profiles in late ripening grapevine berries [4]. To compare the response of the circadian clock in the two different locations, we plotted all of the genes of the model made earlier [4]. Most core clock genes (Additional file 12) and light sensing and peripheral clock genes (Additional file 13) had significantly different transcript abundance in BOD berries than that in RNO berries at the same sugar level (profiles bracketed in a red box). All but one of these (PHYC) had higher transcript abundance in BOD berries relative to RNO berries. The transcript abundance of other genes had nearly identical profiles (not bracketed in a red box). These data are summarized in a simplified clock model (Fig. 6), which integrates PHYB as a key photoreceptor and temperature sensor [35, 36] that can regulate the entrainment and rhythmicity of the core circadian clock, although to be clear it is the protein activity of PHYB, not the transcript abundance that is regulating the clock.

*Chilling-responsive genes*

Temperatures were colder in Reno than BOD, reaching chilling temperatures in the early morning hours. A number of previously identified chilling-responsive genes [37] in Cabernet Sauvignon had a higher transcript abundance in RNO berries as compared to BOD berries (Fig. 7). These genes included *CBF1* (previously named CBF4, but renamed to be consistent with the ortholog of arabidopsis), a transcription factor that

**Figure 6.** A simplified model of the core circadian clock genes. Black lines and bars

represent known inhibitory reactions, red arrows indicate known stimulatory reactions, and

blue lines represent known physical interactions. PHYB is phytochrome B; PIF7 is

Phytochrome Interacting Factor 7; COP1 is Constitutive Photomorphogenic 1; FAR1 is Far-

Red Impaired Response 1; FHY3 is Far Red Elongated Hypocotyl 3; ELF4 is Early

Flowering 4; ELF3 is Early Flowering 3; CCA1 is Circadian Clock Associated 1; PRR9 is

Psuedo-Response Related 9; PRR7 is Psuedo-Response Related 7; PRR5 is Psuedo-

Response Related 5; TOC1 is Timing of CAB expression 1; and RVE1 is Reveille 1.

regulates the cold stress regulon [38], *ERF6L1* (named *ERF4* in earlier grape

annotations), an AP2/ERF domain transcription factor, *CML41*, that encodes

a calmodulin-like protein, *XTH23*, that encodes a xyloglucan

endotransglucosylase/hydrolase that is a cell wall loosening enzyme,

*SULTR3*;4, a sulfate transporter, and *TPPD*, a trehalose phosphate

**Figure 7.** Transcript profiles of some cold responsive genes. Data shown are the means ±

SE; n = 3. Error bars not shown are smaller than the symbol. The symbol legend is

displayed in the figure. CBF1 is C-Repeat Binding Factor 1; ERF6L1 is Ethylene Response

Factor 6-like 1; CML41 is Calmodulin 41; TPPD is Trehalose 6-Phosphate Phosphatase D;

XTH23 is Xyloglucan Endotransglucosylase/Hydrolase 23 and SUFTR3;4 is Sulfate

Transporter 3;4.

phosphatase, identified as a chilling-sensitive gene in grapevine involved in

trehalose metabolism [39]; trehalose 6-phosphate is a signaling molecule in

plants that has diverse roles in developmental programs, sugar regulation

and abiotic stress tolerance, including temperature stress [40].

***DEGs associated with biotic stimuli***

The top DEGs in BOD berries were highly enriched in biotic stimuli genes.

Some of these genes included pathogenesis proteins (PR); these genes had

higher transcript abundance in BOD berry skins relative to RNO berries (Fig.

8).

**Figure 8.** Expression profiles of pathogenesis proteins (PR) involved with powdery mildew.

Data shown are the means ± SE; n = 3. Error bars not shown are smaller than the symbol.

The symbol legend is displayed in the figure. MLA10 is Intracellular Mildew A 10.

The transcript abundance of PR10 increased with the sugar level. PR10

transcript abundance increases in response to powdery mildew (*Erysiphe*

*necator*) inoculation in Cabernet Sauvignon leaves [41]; Vitvi05g01760

corresponds to the 1610704_at Affymetrix probe set in the VV12 experiment

in PlexDB (plexdb.org). Furthermore, many other genes induced by powdery

mildew were also at much higher transcript abundance levels in BOD

berries. These included a PR3 protein that is a class IV chitinase, a

thaumatin-like protein (PR5) and a number of the stilbene synthases

mentioned in the phenylpropanoid metabolism section above (Additional file

11). MLA10 (Vitvi12g01802; Affymetrix probe set 1615715_at in [41])

matches to a fungal protein from *E. necator* and was used as a control probe

set for the presence of powdery mildew [41]. Vitvi12g01802 had a much

higher transcript abundance in BOD berries than that in RNO berries. These

data indicate that powdery mildew infection may have been higher in BOD

berries and that this infection may induce some of the phenylpropanoid

pathway as well.

***DEGs associated with hormonal stimuli***

*Auxin signaling genes*

Auxin transport (31 genes; 1.96E-04) and cellular response to auxin stimulus

(47 genes, 3.09E-04) were highly enriched GO categories for the DEGs

(Additional file 4). Auxin is known to have multiple effects on grape berry

ripening [42, 43]. Auxin can delay berry ripening at the veraison stage, which

is at the beginning of berry ripening. Some auxin metabolism (*GH3.1*) and

signaling genes such as *ABF5, IAA13, IAA15, IAA27*, and *ARF5* had a

higher transcript abundance in RNO berries (Additional file 3). Other auxin

metabolism (*GH3*.6, *JAR1*) and auxin signaling genes had a higher transcript

abundance in BOD including *ARF2*, *ARF8*, *ARF10*, *ARF11*, *IAA1*, *IAA10*,

*IAA16*, *ARAC1* (auxin-induced proteolysis), and *GID1B*, a gibberellin

receptor (Additional file 3).

*ABA metabolism and signaling genes*

ABA is a stress hormone that responds to water deficits in grapevine [44]. A

number of ABA-related genes are differentially expressed in berry skins

between the two locations (Additional file 14). *NCED3.1* and *NCED3.2*

(previously described as *NCED3* and *NCED5*, respectively), which are

responsive to water deficit [45, 46], had higher transcript abundance in RNO

and *NCED6*, which is highly expressed in embryos [45], had higher transcript

abundance in BOD. NCED6, but not NCED3 is involved in seed ABA and

seed dormancy [46]. Additionally a number of other genes involved in the

ABA signaling pathway had higher transcript abundance including some

ABF, bZIP and MYB transcription factors and kinases identified to be in the

ABA signaling pathway [47, 48]. Interestingly, BAM1 (Barley Any Meristem 1)

was identified to be the receptor to a root signaling peptide hormone

(CLE25) that responds to water deficit and upregulates *NCED3* transcript

abundance in arabidopsis leaves [49]. The transcript abundance of BAM1

was higher in RNO berries than that of BOD berries (Additional File 14).

There were no differences in the transcript abundance of CLE25 (data not

shown).

*Ethylene signaling genes higher in BOD*

There were 72 DEGs that were enriched in the response to ethylene GO

category (Additional file 4). Ethylene is a stress hormone that responds to

many types of biotic [50] and abiotic [51] stresses in addition to its role in fruit

development and ripening [52]. Many ethylene-related genes had a higher

transcript abundance in BOD berries. These included ethylene biosynthesis,

ethylene receptors and ERF transcription factors (Fig. 9). ERF1 and ERF2

are at the beginning of the ethylene signaling pathway and are direct targets

of EIN3 [48, 53]. Other ERF transcription factors (e.g. ERF98 and ERF59)

identified as hubs in the ethylene signaling pathway in arabidopsis leaves

[54] were also differentially expressed in a similar manner as ERF1 and

ERF2 between the two locations (data not shown).

### DEGs associated with mineral nutrients

*Iron homeostasis genes*

Nineteen DEGs were associated with genes enriched in iron homeostasis

(Additional file 4); five of these DEGs, *NRAMP2* (Vitvi18g00695), *SIA1*

(Vitvi12g02500), *VIT1* (Vitvi01g00512), *FER4* (Vitvi08g00763) and *ABS4*

(Vitvi18g00583) were in the top 400 DEGs in RNO berries. Only *VIT1* was in

the top 400 DEGs for BOD berries. Nine examples of DEGs involved in iron

homeostasis are shown in Figure 10. Iron homeostasis genes *VIT1*, *IREG3*,

and *ATH13* have higher transcript abundance in BOD berries than in RNO

berries. Iron homeostasis genes *SIA1*, *YSL3*, *FER4*, *FER2*, and *ABS4* had

higher transcript abundance in RNO berries compared to BOD berries.

Average available iron soil concentrations were about 5 times higher in the

BOD vineyard soil compared to the RNO vineyard soil (Table 1).

2  **Figure 9.** The transcript abundance of DEGs involved with ethylene metabolism and

3  signaling. Data shown are the means ± SE; n = 3. Error bars not shown are smaller than the

4  symbol. The symbol legend is displayed in the figure. ACS6 is 1-Aminocyclopropane-1-

5  Carboxylic Acid (ACC) Synthase 6; ACO1 is ACc Oxidase 1; ACOL is ACc Oxidase-Like;

6  ERS1 is Ethylene Response Sensor 1; CTR1 is Constitutive Triple Response 1, EIN3 is

Ethylene-Insenstive 3; and ERF1 and ERF2 are Ethylene Response Factors 1 and 2.

**Figure 10.** Expression profiles of DEGs involved in iron homeostasis. Data shown are the

means ± SE; n = 3. Error bars not shown are smaller than the symbol. The symbol legend is

displayed in the figure. SIA1 is Salt-Induced ABC1 Kinase 1; VIT1 is Vacuolar Iron

Transporter 1; FER2 and 4 are Ferritin 2 and 4; YSL3 is Yellow Stripe-like 3; IREG3 is Iron-

Regulated Protein 3; NRAMP3 is Natural Resistance-Associated Macrophage Protein 3;

ATH13 is ABC2 homolog 13; and ABS4 is Abnormal Shoot 4.

**Discussion**

The common gene sets for both locations represented approximately 20% of

the genes expressed. The processes involved included the increase of

catabolism and the decline of translation and photosynthesis. It is clear that

these processes play important roles in berry ripening. Most transcript

abundances varied with increasing sugar levels and berry maturation and most of these varied with the vineyard site. Many of the DEGs were enriched with gene ontologies associated with environmental or hormonal stimuli.

*DEG expression profiles of grape berry skins were associated with environmental factors and seed development*

Plants are exposed to a multitude of factors that influence their physiology even in controlled agricultural fields such as vineyards. The vineyards in BOD and RNO are exposed to very different environments; these environmental influences were reflected in some of the DEG gene sets with enriched gene ontologies. Our data support the hypothesis that transcriptomic dynamics in the late stages of berry ripening are sensitive to local environmental influences on the grapevine.

While most transcript abundances in berries are largely influenced by genetics or genotype, environment also plays a large role [2]. It is impossible with the experimental design of this study to determine the amount that each of the environmental factors contributed to the amount of differential expression in these two locations. There are too many variables and too many potential interactions to determine anything conclusively. All we can say is that these genes are differentially expressed between the two locations, which are likely due to known and unknown factors (a sense of place or terroir).

In this study we compared two different clones of Cabernet Sauvignon, one on rootstock and the other on its own roots. There are likely to be some effects on the transcript abundance in the berries between these grapevines

as a result of the genetic differences between their roots and their clonal

shoots/scions. Not knowing what these genes might be from previous

studies prevents us from drawing any clues. These and other factors most

certainly affect the berries to some degree. However, our data indicate that

grape berry skins also were responsive to a multitude of potential

environmental factors in the two vineyard locations and possibly from signals

coming from the maturing seed. We say potential environmental factors

because we did not control for these factors; we associated transcript

abundance with the factors that were different in the two locations. The

transcript abundance profiles along with functional annotation of the genes

gave us clues to factors that were influencing the berries and then

associations were made with the known environmental variables. Further

experiments are required to follow up on these observations.

The environmental factors that we were able to associate with differences in

transcript abundance (DEGs) between the two locations included air

temperatures, light, moisture (rainfall and relative humidity) and biotic stress.

These factors in turn were associated with transcript abundance involved

with physiological responses and berry traits such as seed and embryo

development, hormone signaling (ABA, ethylene and auxin),

phenylpropanoid metabolism, and the circadian clock. In the following

sections we discuss in more detail some of the possible environmental

factors that were reflected in the enriched gene ontologies found in the gene

sets from this study.

*Transcript abundance of light-responsive genes*

Light regulates the transcript abundance of many genes in plants. It has

been estimated that 20% of the plant transcriptome is regulated by white

light and this includes genes from most metabolic pathways [55]. Light is

sensed by a variety of photoreceptors in plants [36]; there are red/far red,

blue and UV light receptors. PHYB is a key light sensor, regulating most of

the light sensitive genes [56] and sensing the environment through red light

to far-red light ratios and temperature [36, 57]. PHYB entrains the circadian

clock affecting the rate of the daily cycle [58] and the expression of many the

circadian clock genes [56]; PHYB induces morning phase genes and

represses evening phase genes. Other photoreceptors can entrain the

circadian clock as well [36].

PHYB and the circadian clock are central regulators of many aspects of plant

development including seed germination, seedling growth, and flowering [36,

58, 59]. The circadian clock influences the daily transcript abundance of

genes involved in photosynthesis, sugar transport and metabolism, biotic

and abiotic stress, even iron homeostasis [58].

Light signaling was very dynamic in the berry skin transcriptome in the late

stages of berry ripening with a higher transcript abundance of many light

signaling genes in BOD berries. Many photoreceptors that interact with the

circadian clock had a higher gene expression in BOD berries. In the

circadian clock model, Circadian Clock Associated 1 (*CCA1*) is an early

morning gene and has its highest expression at the beginning of the day. It is

at the start of the circadian core clock progression through the day, whereas

the transcript abundance of Timing Of CAB Expression 1 (*TOC1*) is highest

at the end of the day and finishes the core clock progression (Fig. 6). In both

of these cases, there is a higher transcript abundance of these genes in

BOD than in RNO.

The evening complex is a multi-protein complex composed of Early

Flowering 3 (ELF3), Early Flowering 4 (ELF4) and Phytoclock 1 (PCL1 also

known as LUX) that peaks at dusk. Of the three of these proteins, ELF4 had

a small difference in transcript abundance between locations around the

20°Brix level (Fig. 6), whereas there was no distinction between locations or

sugar levels for ELF3 (Fig. 6) or PCL1 (Additional file 12). The transcript

abundance of ELF3 increased with sugar level and shortening of the day

length (the higher sugar level comes later in the season and thus is at a

shorter day length). ELF3, as part of the evening complex (EC), has direct

physical interactions with PHYB, COP1 (Constitutive Photomorphogenic 1)

and TOC1 [60] linking light and temperature signaling pathways directly with

the circadian clock. It is interesting that most of the components of the clock

showed significant differences in transcript abundance between BOD and

RNO, except for the three proteins that make up the evening clock.

The transcript abundance profile of PHYB was similar in both BOD and RNO

berries (Fig. 6), however the changes in transcript abundance with sugar

level occurred in BOD berries at a lower sugar level. There was a gradual

decline of PHYB transcript abundance with increasing sugar level until the

last measurement at the fully mature stage, where there was a large

increase in transcript abundance. A very similar profile is observed for

Reveille 1 (RVE1).  RVE1 promotes seed dormancy in arabidopsis and

PHYB interacts with RVE1 by inhibiting its expression [61]. PIF7

(Phytochrome Interacting Factor 7), interacts directly with PHYB to suppress

PHYB protein levels [62]. Likewise, PIF7 activity is regulated by the circadian

clock [63]. PIF7 had different transcript abundance profiles in the BOD and

RNO berries. It was somewhat similar to the profiles in PHYB in RNO berries

and the opposite in BOD berries, indicating a more complicated role of the

transcript abundance for this protein in the two different locations. Note that

transcript abundance does not necessarily infer protein abundance and so

making functional associations here is not necessarily accurate. The

transcript abundance of two of the other grape phytochromes (PHYA and

PHYE) did not vary significantly between the two locations or at different

sugar levels. PHYC had a higher transcript abundance in RNO berries and

did not change much with different sugar levels.  Many other light receptors

(e.g. CRY3, FAR1, FRS5, etc.) had higher transcript abundance in BOD

berries (Additional file 13). Thus, light sensing through the circadian clock is

a complicated process with multiple inputs.

RVE1 follows a circadian rhythm [64]. It behaves like a morning-phased

transcription factor and binds to the EE element, but it is not clear if it is

affected directly from the core clock (e.g. TOC1 or EC which repress other

morning gene paralogs like *CCA1* and *LHY*) or through effects of PHYB or

both. PHYB downregulates RVE1; RVE1 promotes auxin concentrations and

decreases gibberellin (GA) concentrations [61]. Warmer night temperatures

(as in BOD) cause more rapid reversion of the active form of PHYB to the

inactive form [36] and thus may promote a higher expression/activity of

RVE1. $P_r$ (phytochrome in the red form, which is the physiologically inactive

form) appears to accelerate the pace of the clock [58]. It is unclear what role

phytochromes might have in seed and fruit development in grapes.

Very little is known about the effect of PHY on fruit development in general.

In one tomato study, the fruit development of *phy* mutants was accelerated

[65], suggesting that PHYB as a temperature/light sensor and a regulator of

the circadian clock may influence fruit development. Carotenoid

concentrations, but not sugar concentrations, also were affected in these

mutants.

Photoperiod affects the transcript abundance of PHYA and PHYB in grape

leaves [66]. In the present study, the transcript abundance of the majority of

the photoreceptor genes in berry skins, including red, blue and UV light

photoreceptors, had a higher transcript abundance in BOD berries

(Additional file 13). It is unclear what the effect of PHYB and the circadian

clock have on grape berry development.  However, there are clear

differences between the two locations and it seems likely that PHYB and the

circadian clock are key grape berry sensors of the environment, affecting

fruit development and composition.

*Transcript abundance of temperature-responsive genes*

The berry transcriptome is sensitive to temperature [2, 3]. Temperature

related genes were differentially expressed at the two locations in our study.

The RNO berries were exposed to a much larger temperature differential

between day and night than BOD berries and were also exposed to chilling

temperatures in the early morning hours during the late stages of berry

ripening (Table 1). The transcript abundance of some cold-responsive genes

was higher in RNO berry skins than in BOD berry skins, including *CBF1* and

*ERF6L1*.

CBF1 transcript abundance is very sensitive to chilling temperatures; it is a master regulator of the cold regulon and improves plant cold tolerance [38, 67, 68]. PIF7 binds to the promoter of CBF1, inhibiting CBF1 transcript abundance, linking phytochrome, the circadian clock and CBF1 expression [63]. Our data are consistent with this model; transcript abundance of PIF7 was higher and CBF1 transcript abundance was lower in BOD berry skins than RNO berry skins (Fig. 6 and 7).

The transcript abundance of *ERF6L1* was identified in another study to peak at a TSS where optimum berry flavor and ripeness occur [69]. It was hypothesized that this gene may be associated with berry ripening and the transcript abundance involved with aroma-related compounds. However, in a later study, the transcript abundance of *ERF6L1* did not peak, but steadily declined with increasing sugar levels [4]. A major difference between these two studies is that in the earlier study the berries were shipped on ice 24 h before freezing in liquid nitrogen, whereas in the latter study, berries were harvested and flash frozen immediately. Thus, the increased *ERF6L1* transcript abundance may have been responding to the chilling temperatures rather than to the sugar concentration or there was an interaction between the two factors. This chilling response may also occur during some commercial harvests, when berries are chilled down before starting fermentation. The cold-responsive transcripts identified in this study (Fig. 7) may be good markers for chilling stress since their abundance was increased by chilling temperatures in shoot tips [37], leaves [39] and berry skins (this study). We do not have an explanation for the large decrease in

transcript abundance for many of these genes at 26°Brix for the RNO berries

(Fig. 7).

*Transcript abundance of dehydration and seed dormancy genes*

ABA concentrations in plants increase in response to dehydration and ABA

triggers a major signaling pathway involved in osmotic stress responses and

seed development [70]. ABA concentrations only increase in the seed

embryo near the end of seed development when the embryo dehydrates and

goes into dormancy. ABA concentrations remain high to inhibit seed

germination. The transcript abundance of ABA signaling genes such as

ABF2 and SnRK2 kinases increase after application of ABA to cell culture

[71] and in response to dehydration [48] in leaves of Cabernet Sauvignon.

The data in this study are consistent with the hypothesis that BOD berries

are riper at lower sugar levels.  The ABA signaling genes in the berry skins

had higher transcript abundance in BOD berries indicating that ABA

concentrations were higher in BOD than RNO berries even though RNO

berries were exposed to drier conditions (Table 1). ABA concentrations may

be higher in the BOD berry skins based upon the higher transcript

abundance of important ABA signaling and biosynthesis genes encoding for

ABF2, SnRK2 kinases and NCED6. We hypothesize that this is seed derived

ABA since water deficits were not apparent in BOD with the recent rainfall

and high humidity. In contrast, *NCED3* genes had higher transcript

abundance in RNO berry skins, which might occur as the result of the very

low humidity and large vapor pressure deficit (the vines were irrigated). The

lower expression of *NCED6* in RNO berry skins may indicate that the seeds

in the berry were more immature than the BOD berries. The higher

expression of other seed development and dormancy genes (e.g. *RVE1*,

*ARF2*, *ARF10*, etc.) in the berry skins support the argument that BOD berries

(and seeds) matured at a lower sugar level than the RNO berries.

The ABA concentrations in the berry skins are a function of biosynthesis,

catabolism, conjugation and transport. ABA in seeds increase as the seed

matures and some of this ABA may be transported to the skin. In fact,

*ABCG40* genes, which encode ABA transporters, had higher transcript

abundance in BOD berry skins than that in RNO (Additional file 3). Part of

the ABA in skins may be transported from the seed and part of it might be

derived from biosynthesis in the skins. *NCED6* transcript abundance in the

skins was higher in BOD berries. Perhaps the transcript abundance of

*NCED6* in the skin is regulated by the same signals as the embryo and

reflects an increase in seed maturity. *AtNCED6* transcript abundance is not

responsive to water deficit in arabidopsis, but *AtNCED3* and *AtNCED5* are

[46]. This is consistent with the higher *NCED3* and *BAM1* transcript

abundance in RNO berries. Thus, there are complex responses of ABA

metabolism and signaling. It would appear that there may be two different

ABA pathways affecting ABA concentrations and signaling: one involved with

embryo development and one involved with the water status in the skins.

Auxin is also involved with ABA signaling during the late stages of embryo

development in the seeds. Auxin signaling responses are complex. ABF5 is

an auxin receptor that degrades Aux/IAA proteins, which are repressors of

ARF transcriptional activity [72]. Thus, a rise in auxin concentration releases

Aux/IAA repression of ARF transcription factors, activating auxin signaling. In

the berry skins, there was a diversity of transcriptional responses of *Aux/IAA*

and *ARF* genes in the two locations, some with increased transcript

abundance and others with decreased transcript abundance. As with ABA

signaling, there may be multiple auxin signaling pathways operating

simultaneously.

One pathway appears to involve seed dormancy. *ARF2* and *ARF10* genes

had a higher transcript abundance in BOD berries. ARF10 upregulates *ABI3*

expression and promotes seed dormancy [73]; ARF2 also promotes

dormancy through the ABA signaling pathway [74]. This is consistent with

the hypothesis that BOD berries reach maturity at a lower sugar level than

RNO berries. Auxin may play a role in promoting seed dormancy through

ARF2 and ARF10 effects on ABA signaling in the late stages of berry

ripening.

*Transcript abundance of biotic stress genes*

Grapevines have very dynamic gene expression responses to pathogens

[75, 76]. The top 150 DEGs for BOD berries were highly enriched with biotic

stress genes. The higher rainfall and high relative humidity in BOD would

make moist conditions suitable for pathogenic fungi to grow. We detected a

much higher transcript abundance of powdery mildew-responsive genes in

BOD berries and this may be connected to a higher transcript abundance of

ethylene and phenylpropanoid genes as part of a defense response. The

transcript abundance profiles of some of these genes (e.g. PR10, PAL4,

STS10, ACS6, and ERF2; see Figs. 4, 5, 8 and 9) are remarkably similar.

Increased ethylene signaling in grapevines has been associated with powdery mildew infection and phenylpropanoid metabolism and appears to provide plant protection against the fungus [77, 78]. Genes involved with phenylpropanoid metabolism, especially *PAL* and *STS* genes, appear to be quite sensitive to multiple stresses in the environment [79]. In arabidopsis there are four *PAL* genes [80]. *PAL1* and *PAL2* appear to be involved with flavonoid biosynthesis; *PAL4* provides pathogen resistance. Ten different *PAL4* orthologs had a higher transcript abundance in BOD berry skins; many *STS* genes also had a higher transcript abundance in BOD berry skins (Additional file 11). Stilbenes are phytoalexins and provide pathogen resistance in grapes and *STS* genes are strongly induced by pathogens [75]. Thus, the higher transcript abundance of powdery mildew genes may be associated with the higher transcript abundance of genes in the ethylene and phenylpropanoid pathways.

*Transcript abundance of iron homeostasis genes*

The transcript abundance of a number of iron homeostasis genes were significantly different in the two locations (Fig. 10) and there was a difference in soil available iron concentrations in the two locations. However, iron uptake and transport in plants is complicated depending on multiple factors, such as pH, soil redox state, organic matter composition, solubility in the phloem, etc. Thus, it is impossible to predict iron concentrations in the berry without direct measurements. The roles of these genes in iron homeostasis and plant physiological functions are diverse. Iron supply can affect anthocyanin concentrations and the transcript abundance of genes in the

phenylpropanoid pathway in Cabernet Sauvignon berry skins [81]. One of

the DEGs, SIA1, is located in the chloroplast in arabidopsis and appears to

function in plastoglobule formation and iron homeostasis signaling in concert

with ATH13 (also known as OSA1) [82]. Another DEG, YSL3, is involved in

Fe transport [83]. It acts in the SA signaling pathway and appears to be

involved in defense responses to pathogens. It also functions in iron

transport into seeds [84]. FER4 is one of a family of ferritins (iron-binding

proteins) found in arabidopsis [85]. FER2 is a specific ferritin that is

associated with the dry seed during fruit development [86]. It is a storage

form of iron and also is involved in protection against oxidative stresses. The

transcript abundance of FER2, FER3 and FER4 is induced by chilling

temperatures in the light [87], but not FER1. VIT1 and NRAMP3 are vacuolar

iron transporters [86] and are also involved in iron storage in seeds.

Interestingly, increased *VIT1* transcript abundance leads to decreased *FER2*

transcript abundance [86]. This response is consistent within the studied

berries as *VIT1* transcript abundance increased with sugar level and had a

higher transcript abundance in BOD berries than that of RNO berries and the

transcript abundance of *FER2* declined with sugar levels in BOD berries to a

lower amount than that of RNO berries.

Other DEGs are also responsive to iron supply. IREG3 (also known as

MAR1) appears to be involved in iron transport in plastids; its transcript

abundance increases with increasing iron concentrations [88]. ABS4 is a

MATE (multidrug and toxic compound extrusion) efflux transporter that is

located in the Golgi complex and its transcript abundance is induced by iron

and osmotic stress [89].

It is unclear what specific roles these iron homeostasis genes are playing in grape berry skins, but they appear to be involved in iron storage in seeds and protection against oxidative stress responses [85, 86]. One possible explanation for the transcript abundance profiles in the BOD and RNO berry skins is that ferritins are known to bind iron and are thought to reduce the free iron concentrations in the chloroplast, thus, reducing ROS production that is caused by the Fenton reaction [85]. As chloroplasts senesce during berry ripening, iron concentrations may rise as a result of the catabolism of iron-containing proteins in the thylakoid membranes; thus, berry skins may need higher concentrations of ferritins to keep free iron concentrations low. This might explain the increase in ferritin transcript abundance with increasing sugar levels. The transcript abundance differences between BOD and RNO in iron homeostasis genes also may be related to an earlier berry ripening (seed development and chloroplast degradation) in BOD berries (earlier decline at lower sugar levels) of transcript abundance of *FER2* and higher transcript abundance of *VIT1*, and the chilling temperatures (higher transcript abundance of *FER2* and *FER4*) in RNO. Most soils contain 2 to 5% iron including available and unavailable iron; soils with 15 and 25 µg g$^{-1}$ of available iron are considered moderate for grapevines [90], but soils with higher concentrations are not considered toxic. Therefore, for both soils in this study, iron concentrations can be considered to be very high but not toxic. The higher available iron concentrations in the BOD vineyard may be associated with the wetter conditions (more reductive conditions) and the lower soil pH.

*Environmental influences on transcript abundance in other studies*

Other researchers using Omics approaches have identified environmental factors that influence grape berry transcript abundance and metabolites. One study investigated the differences in transcript abundance in berries of Corvina (a black-skinned grape cultivar that makes red wine) in 11 different vineyards over three years [91]. They determined that approximately 18% of the berry transcript abundance was affected by the environment. Climate had an overwhelming effect but viticultural practices were also significant. Phenylpropanoid metabolism was very sensitive to the environment and *PAL* transcript abundance was associated with *STS* transcript abundance.

In another study of a white grape cultivar, Garganega, berries were analyzed by transcriptomic and metabolomic approaches [92]. Berries were selected from vineyards at different altitudes and soil types. Again, phenylpropanoid metabolism was strongly influenced by the environment. Carotenoid and terpenoid metabolism were influenced as well.

Two studies investigated the grape berry transcriptomes during the ripening phase in two different regions of China, a dry region in Western China and a wet region in Eastern China [93, 94]. These two locations mirror some of the differences in our conditions in our study, namely moisture, light and elevation, although the China western region has higher night temperatures and more rainfall than the RNO location. In the Cabernet Sauvignon study [93], they compared the berry transcriptomes (with seeds removed) from the two regions at three different stages: pea size, veraison and maturity. The TSS at maturity was slightly below 20°Brix. Similar to our study, the response to stimulus, phenylpropanoid and diterpenoid metabolism GO

categories were highly enriched in mature berries between the two locations.

Differences in the transcript abundance of NCED and PR proteins were also noted. Like in our study, the authors associated the transcript abundance of these proteins to the dry (drought response) and wet (pathogen defense) locations, respectively.

In the second study comparing these two regions in China [94], the effects of the environment on the metabolome and transcriptome of Muscat Blanc à Petits Grains berries were investigated over two seasons; specifically, terpenoid metabolism was targeted. Like in our study, the transcripts in terpenoid were in higher abundance in the wetter location. The transcript abundances were correlated with terpenoid concentrations and a coexpression network was constructed. A specific set of candidate regulatory genes were identified including some terpene synthases (TPS14), glycosyl transferases and 1-hydroxy-2-methyl-2-butenyl 4-diphosphate reductase (HDR). We examined the transcript abundance of some of these candidate genes in our own data but did not find significant differences between our two locations. The contrasting results between our study and Wen et al. (2015) could be for a variety of reasons such as different cultivar responses, berry versus skin samples, or different environmental conditions that affect terpenoid production.

Terpenoid metabolism is influenced by the microclimate [24] and is involved in plant defense responses to pathogens and insects [32, 95]. Light exposure to Sauvignon Blanc grapes (a white grape cultivar) was manipulated by removing adjacent leaves without any detectable differences in berry temperatures [24]. Increased light exposure increased specific carotenoid

and terpene concentrations in the berry. The responses of carotenoid and

terpenoid production to temperature are less clear. Some effect of

temperature was associated with carotenoid and terpenoid production, but to

a lesser extent than light [24]. Higher concentrations of rotundone, a

sesquiterpene, have been associated with cooler temperatures [96]. Water

deficit can also alter carotenoid and terpenoid metabolism in grapes [11, 97].

Terpenes can act as signals for insect attacks and attract insect predators

[95]. Thus, terpenoid metabolism is highly sensitive to the environment and

influenced by many factors.

In contrast to these studies, excess light and heat can affect transcript

abundance and damage berry quality. In addition to a higher rate of malate

catabolism, anthocyanin concentrations and some of the transcript

abundances associated with them are decreased as well [98, 99].

*Temperature effects on berry maturity and total soluble solids*

BOD berries reached maturity at a lower °Brix level than RNO berries; the

cause is likely to be the warmer days and cooler nights in RNO. Higher day

temperature may increase photosynthesis and sugar transport and cooler

night temperatures may reduce fruit respiration. °Brix or TSS approximates

the % sugar in a berry and is a reliable marker of berry maturity in any given

location [100]; however, TSS is an unreliable marker of berry maturity when

comparing grapes from very different climates. The differences in TSS

between BOD and RNO are consistent with other studies on the temperature

effects on berry development. Indirect studies have associated gradual

warming over the last century to accelerated phenology and increased sugar

concentrations in the grape berries [101-104]. Increasing temperature can accelerate metabolism, including sugar biosynthesis and transport, but the increase in metabolism is not uniform. For example, the increase in anthocyanin concentration during the ripening phase is not affected as much as the increase in sugar concentration [105]. These responses vary with the cultivar [102], complicating this kind of analysis even further.

Direct studies of temperature effects on Cabernet Sauvignon berry composition also are consistent with our data. In one study, the composition of Cabernet Sauvignon berries was altered substantially for vines grown in phytotrons at 20 or 30°C temperatures (temperatures that are very similar to the BOD and RNO temperatures occurring in the present study) [106]. Cooler temperatures promoted anthocyanin development and malate concentrations (inhibited malate catabolism) and higher temperatures promoted TSS (°Brix) and proline concentrations [106]. In a second study, vines were grown at 20 or 30°C day temperatures with night temperatures 5°C cooler than the day [107]. In this study, higher temperatures increased berry volume and veraison started earlier by about 3 to 4 weeks [107]. The authors concluded that warmer temperatures hastened berry development. In a third study, Cabernet Sauvignon berry composition was affected in a similar manner by soil temperatures that differed by 13°C [108].

TSS concentrations are also affected by light and the vine water status. Light is generally not a factor because there is usually a large enough leaf area and sufficient light levels to saturate this source to sink relationship [109, 110]. Sun-exposed Cabernet Sauvignon berries in the vineyard had higher TSS than shaded berries [109]. This sunlight effect was attributed largely to

an increase in berry temperature rather than an increase in the fluence rate

per se. A higher grapevine water status results in larger berry size and lower

sugar concentrations [111] and water deficit is known to increase sugar

concentrations in Cabernet Sauvignon [11]. However, temperature is thought

to have the largest effect on sugar concentrations [16].

Other transcriptomic data in the present study indicated that BOD berries

were more mature at a lower sugar level than RNO berries. These included

the transcript abundance profiles of genes involved in autophagy, auxin and

ABA signaling, iron homeostasis and seed development. Many of these

DEGs had an accelerated rate of change in BOD berries. While these

transcripts are in the skins, they may be influenced by signals coming from

the seed. In addition, there was a higher transcript abundance for most

genes involved with the circadian clock in BOD berries. PHYB can regulate

the circadian clock [58] and PHYB activity is very sensitive to night

temperatures (BOD had higher night temperatures); PHYB reversion is

accelerated to the inactive form at warmer temperatures [36]. The inactivity

of phytochrome promotes the expression of RVE1, which promotes auxin

concentrations and seed dormancy [61]. Thus, all things considered, it is

likely that temperature and/or the temperature differentials between day and

night significantly contributed to the differences in the rate of berry

development and sugar accumulation in the two locations.

*Are there reliable markers to harvest berries at maturity?*

Determining maturity of grapes is a difficult and error prone process. Reliable

markers could aid in the decision of when to harvest the grapes. "Optimum"

maturity is a judgement call and will ultimately depend on the winemaker's or

grower's specific goals or preferences. A combination of empirical factors

can be utilized including °Brix, total acidity, berry tasting in the mouth for

aroma and tannins, seed color, etc. TSS by itself may not be the best marker

for berry ripening as it appears to be uncoupled from berry maturity by

temperature. Phenylpropanoid metabolism, including anthocyanin

metabolism, is also highly sensitive to both abiotic and biotic stresses and

may not be a good indicator of full maturity. Thus, color may not be a good

indicator either. Specific developmental signals from the seed or embryo,

such as those involved with auxin and ABA signaling, may provide more

reliable markers for berry ripening in diverse environments, but will not be

useful in seedless grapes. Aromatic compounds may also be reliable

markers but they will need to be generic, developmental markers that are not

influenced by the environment. This study revealed many genes that are not

reliable markers because they were expressed differently in different

environments. One candidate marker that is noteworthy is ATG18g

(Vitvi07g00210). Its transcript abundance increased and was relatively linear

with increasing °Brix and these trends were offset at the two locations

relative to their level of putative fruit maturity (Fig. 3). ATG18g is required for

the autophagy process [112] and maybe important during the fruit ripening

phase. It was found to be a hub gene in a gene subnetwork associated with

fruit ripening and chloroplast degradation [4]. Further testing will be required

to know if it is essential for fruit ripening and whether its transcript

abundance is influenced by abiotic and biotic stresses in grape berry skins.

*Conclusions*

The ultimate function of a fruit is to produce fully mature seeds in order to reproduce another generation of plants. Berry ripening involves complex signaling that tells other organisms when the fruit is ready for consumption and seed dispersal. In this study, we tested and confirmed the hypothesis that the transcript abundance in grape skins differed in two different locations with different environments. Another goal was to distinguish transcripts with common profiles from differentially expressed genes. The observations made in this study provide lists of such genes and generated a large number of hypotheses to be tested in the future. WGCNA was particularly powerful and enhanced our analyses. It is clear that berry ripening may respond to a multitude of factors both before and during the late stages of berry ripening. The transcriptomic analysis in the late stages of berry ripening in this study indicated that transcript abundance was very dynamic and possibly influenced by a number of environmental and internal (e.g. seed signals) factors. We could discern these influences using functional analysis of the genes and GO enrichment analysis of the transcriptomic datasets. Temperature, light, moisture and the local microbiome were major factors that may have contributed to the transcript abundance profiles of the berry skins. While earlier fruit development stages clearly have an effect on fruit composition, the dynamic changes in transcript abundance in the late stages of berry ripening indicated that berries still have a "sense of place"; it appears that the environment continues to affect berry metabolite composition, which may impact berry flavor, color and other quality-related traits. Future studies are required to follow up on these observations. It

appears that fruit ripening is very malleable. Manipulation of the canopy (time

and intensity of leaf removal at different locations on the plant) may offer a

powerful lever to adjust gene expression and berry composition, since these

parameters are strongly affected by light and temperature.

**Methods**

*Plant materials*

Grapevines in Reno (*Vitis vinifera* L. cv. Cabernet Sauvignon clone 8) were

grown on their own roots at the Valley Road Experimental Vineyard at the

Nevada Agricultural Experiment Station on the campus of the University of

Nevada, Reno. Research approval was obtained by Grant R. Cramer from

the Nevada Agricultural Experiment Station and the University of Nevada,

Reno. The grapevines were originally obtained as certified material from

Inland Desert Nursery, Benton, City, Washington, USA. The grapes were

harvested between September 10 and October 2012 depending on maturity.

Maturity was assessed using a digital refractometer (HI 96811, Hanna

Instruments, Woonsocket, RI, USA) to measure soluble solids (°Brix) that are

mostly made up of sugars. Berry clusters were collected between 11.00 h

and 13.00 h (near solar noon) in an attempt to minimize temporal

transcriptional response variations caused by the circadian clock. At harvest,

individual berry °Brix levels were determined with a digital refractometer.

Separated berry skins were placed into 50 mL centrifuge tubes in liquid

nitrogen according to sugar level (1 ± 0.5 °Brix increments; 19 to 27 °Brix). In

this way berries were collected over many days from multiple clusters from

multiple vines from 3 different individually drip-irrigated blocks in the

vineyard. Stem-water potentials were measured weekly to determine

irrigation levels that would maintain the water status of the vines. The vines

were regularly sprayed for fungal prevention. Each block in the vineyard was

considered an experimental replicate. Soil testing was performed by A & L

Western Agricultural Laboratories (Modesto, CA, USA).

Cabernet Sauvignon clone CA33 412 grapevines in Bordeaux were grown

on SO4 rootstock at the VitAdapt parcel on the Institut National de

Recherche Agronomique (INRA) research station in Villenave d'Ornon, in the

Aquitaine region of France. There were 5 replicate blocks within this site to

mitigate soil variation. Additional details of the VitAdapt project can be found

in [113]. Berries for the three replicates were collected at the 14:00 h of the

day. Environmental conditions and variables of the Reno and BOD vineyards

are listed in Table 1. The vines were regularly sprayed for fungal prevention.

Soil testing was performed by Aurea Agroscience (Blanquefort, France).

*RNA extraction*

Three experimental replicates at 20, 22, 24 and 26 °Brix were used for

sequencing. In RNO, total RNA from the three experimental replicates was

extracted from approximately 250 mg of frozen, finely ground, skin tissue (a

mixture of frozen berry skins from multiple vines and clusters from three

different blocks) using a modified CTAB extraction protocol followed by an

additional on-column DNase digestion using a Qiagen RNeasy Mini Kit

(Qiagen, Valencia, CA, USA) as in [114]. RNA quality and quantity were

assessed with a Nanodrop ND-1000 spectrophotometer (ThermoFisher

Scientific, Waltham, MA, USA) and an Agilent 2100 Bioanalyzer (Agilent

Technologies, Santa Clara, CA, USA).

In BOD, total RNA was extracted according to Reid et al. [115] from 1 g of

frozen, finely ground berry skins. Traces of genomic DNA were removed by

a DNAse I treatment according to the manufacturer's instructions (Ambion

TURBO DNA-free DNase, Life Technologies). The RNA was quantified using

a Nanodrop 2000c spectrophotometer (Thermo Scientific) and their integrity

was checked on an 1.8% agarose gel.

*RNA-seq library preparation and sequencing*

Cabernet Sauvignon 50 bp single-end, barcoded libraries were constructed

and sequenced by the Neuroscience Genomics Core at the University of

California, Los Angeles for the RNO samples and by the Genome Center at

the University of California, Davis for BOD samples using Illumina TruSeq

RNA library prep kits (Illumina Inc., San Diego, CA, USA) according to the

manufacturer's instructions. The barcoded libraries were pooled, multiplexed,

and were sequenced using Illumina TruSeq chemistry (version 3.0) on a

HiSeq2000 sequencer (Illumina Inc., San Diego, CA, USA).

*Transcript abundance and statistical analysis*

The single-end sequence fragments (reads) generated by Illumina

sequencing were base-called and demultiplexed. The fastq files generated

were inspected for sequence quality and contamination using FastQC [116].

Illumina adapters were removed from sequences with Trimommatic [117]

version 0.36.  The transcript abundance was estimated from the trimmed

fastq files using Salmon [26] version 0.9.1 using the gcBias, seqBias,

fldMean and fldSD flags. The index file utilized by Salmon for read

alignments was built from the V2 PN40024 *Vitis vinifera* reference genome

transcriptome annotation [27] after filtering out genes with "No start", "No

stop", "Pseudogene", "The sequence of the model RefSeq protein was

modified", "Appear only in secondary splicing variant" annotations and using

a k-mer size of 17. The salmon output (quant files) was imported into

DESeq2 [28] version 1.18.1 using the tximport R package [118] for

determination of differentially expressed genes (DEGs).

*Coexpression network analysis*

A coexpression gene network analysis was performed using the WGCNA

software package (v1.63) in R [119, 120] using all the libraries for each

location. Prior to this analysis, low-expressed genes were removed with a

minimum threshold of 10 counts in all the libraries. Counts data were

transformed using the function varianceStabilizingTransformation of the

package DESeq2. The resulting set of counts was used for network

construction and module detection using the function blockwiseModules. An

adjacency matrix was created by calculating the biweight mid-correlation

raised to a power β of 5 and the maxPoutliers parameter set to 0.05. The

subsequent Topological Overlap Matrix (TOM) was used for module detection

using the DynamicTreecut algorithm with a minimal module size of 30 and a

branch merge cut height of 0.25. The module eigengenes were used to

evaluate the association between the resulting modules (22) and the

experimental traits (Brix degrees and locations).

*Gene functional annotation*

Gene models of the V3 annotation were searched against different protein

databases with the blastx function of the DIAMOND version 0.9.19 software

[121] using default parameters and reporting alignments in the 1% range of

the top alignment score. For each V3 gene model, the best blast hit was kept

(1-to-1). For multiple hits with the same score (1-to-many), the first hit was

kept as the representative result but the other hits are still accessible. The

databases used were Araport11 (release 06.17.16, https://www.araport.org/),

the *Arabidopsis thaliana* TAIR10 (http://www.arabidopsis.org/index.jsp) and

the *Vitis vinifera* IGGP 12X from Ensembl plants 38

(https://plants.ensembl.org/index.html) and the refseq plant protein database

(release 03.12.18, https://www.ncbi.nlm.nih.gov/refseq/). The corresponding

gene annotations were obtained from the Araport11 gff file (release

06.22.16), TAIR10 functional descriptions (release 01.16.13), a manually

curated and actualized grapevine V1 annotation and from the refseq

database.

*Functional enrichment of GO categories*

Gene Ontology (GO) enrichment was performed using the R software

package topGO [122]. Enriched functional categories with an FDR adjusted

p-value > 0.01 after the Fisher's test were filtered for further analysis. For

gene ontology (GO) categories assignments, interproscan version 5.32-71.0

was performed on the full genome with the options -iprlookup and -

goterms [123]. These new annotations were combined with the previously

manually curated GO annotations of the V1 gene models.

**Abbreviations**

**ABA:** ABscisic Acid **ABCG:** Adenosine triphosphate Binding Casette G

**ABF:** Abscisic acid responsive elements Binding Factor **ABS**: Abnormal

Shoot **ACO1**: ACc Oxidase 1 **ACOL**: ACc Oxidase-Like **APG9:**

AutoPhaGy 9 **ARF:** Auxin Response Factor **ATG:** AuTophaGy **ATH**:

Arabidopsis Thaliana abc2 Homolog **BAM:** Barley Any Meristem **BCAT:**

Branched-Chain-amino-acid AminoTransferase **BOD:** BOrDeaux **bp:**

**base pair CCA1:** Circadian Clock Associated 1 **CLE**: CLavata3/Esr-

related **COP1:** COnstitutive Photomorphogenic 1 **CML**: CalModuLin

**CRY3:** CRYptochrome 3 **cv**: cultivar **DEG(s):** Differentially Expressed

Gene(s) **DHS:** 3-deoxy-D-arabino-heptulosonate 7-phosphate synthase

**DMAPP:** dimethylallyl diphosphate **DXR**: 1-deoxy-D-Xylulose 5-

phosphate Reductoisomerase  **EC:** Evening Complex **EE:** Evening

Element **EIN:** Ethylene INsensitive **ELF:** EarLy Flowering **ERF:** Ethylene

Response Factor **FAR1:** Far-Red impaired Response 1 **FDR:** False

Discovery Rate **FER:** FERrittin **FOP1**: FOlded Petal 1 **FRS:** FAR1-

Related Sequence **GO:** Gene Ontology **HDS:** 4-Hydroxy-3-methylbut-2-

en-1-yl Diphosphate Synthase **IAA:** Indole Acetic Acid **IPP:** IsoPentenyl

Pyrophosphate **IREG**: Iron REGulated **JAR**: JAsmonate Resistant **MEP:**

MethylErythritol 4-Phosphate **MLA:** intracellular MiLdew A **NCBI:**

National Center for Biotechnology Information **NCED:** Nine-Cis

Epoxycarotenoid Dioxygenase **PAL:** PhenylAlanine Lyase **PCL1:**

PhytoCLock 1 **PHY:** PHYtochrome **PIF:** Phytochrome Interacting Factor

**PR**: Pathogenesis Related **PRR:** Psuedo-Response Regulator **PSY:**

Phytoene SYnthase **RNO**: ReNO **RVE1:** ReVeillE 1 **SA**: Salicylic Acid

**SIA**: Salt Induced Abc kinase **SnRK:** SNF1 Related protein Kinase **STS**: STilbene Synthase **SULTR**: SULfur TRansporter **TAIR:** The Arabidopsis Information Resource **TAT**: TyrosineAmino Transferase **TOC:** Timing Of Cab expression **TPPD:** Trehalose Phosphate Phosphatase D **TPL:** ToPLess **TPM:** Transcripts Per Million **TPS:** TerPene Synthase **TSS:** Total Soluble Solids **VIT**: Vacuolar Iron Transporter **WGCNA:** Weighted Gene Co-expression Network Analysis **XTH**: Xyloglucan endoTransglucosylase/Hydrolase **YSL**: Yellow Stripe-Like

**Declarations**

**Ethics**

Not applicable.

**Consent to publish**

Not applicable.

**Competing interests**

The authors declare that they have no competing interests.

**Authors' contributions**

GRC, SG, and AD-I designed the experiments. RG and AH collected berries, analyzed the °Brix levels and extracted the RNA. NC functionally annotated the V3 gene models, performed the WGCNA and assisted with estimation of the TPMs. GRC processed and analyzed the data and wrote the body of the paper. All authors reviewed, edited and approved the final version of the manuscript.

## Availability of supporting data

## Acknowledgements

We would like to thank Aude Habran for her assistance in collecting berry skins and extraction RNA in BOD berries and thank the many UNR Biochemistry senior thesis undergraduate students that assisted in berry separation in the RNO berries.

### Funding

This research was supported by the University of Nevada, Reno, Agriculture Experimental Station with the Multi-state USDA-Hatch Grant NEV00383A to GRC.

**Tables**

**Table 1.** Environmental variables for the harvest times in BOD and RNO. Grapes were harvested in RNO from September 10 to 26, 2012 and in BOD from September 17 to October 8, 2013.

| Environmental Variable | BOD (2013) | RNO (2012) |
|---|---|---|
| Elevation (m) | 25 | 1,373 |
| Average Daily Solar Radiation (kW-hr m$^{-2}$) | 1.17 | 5.86 |
| Day length Starting Harvest Date | 12:25:36 | 12:38:34 |
| Day length Ending Harvest Date | 11:20:57 | 11:57:37 |
| Maximum Temperature Starting Harvest Date | 19 | 30.5 |
| Minimum Temperature Starting Harvest Date | 13 | 13.9 |
| Maximum Temperature Ending Harvest Date | 18 | 27.8 |
| Minimum Temperature Ending Harvest Date (°C) | 11 | 6.7 |
| Ave September Maximum Temperature (°C) | 23.9 | 30.2 |
| Ave September Minimum Temperature (°C) | 13.9 | 10.2 |
| Latitude | 44°47'23.83'' N | 39°52'96'' N |
| Longitude | 0°34'39.3'' W | 119°81'38'' W |
| September Precipitation (mm) | 65.5 | 2.03 |
| Average Monthly Relative Humidity (%) | 74 | 34 |
| Soil Type | Gravelly soil | Sandy Loam |
| Soil pH | 6.2 | 6.7 |
| Soil Fe (µg g-1) | 193 | 38 |
| Root stock | SO4 | Own-rooted |

**Supplemental Information**

**Additional file 1.** Log$_2$ transcripts per million (TPM) to genes uniquely mapped to the V2 PN40024 grape genome.

**Additional File 2.** Functional annotation of all V3 structurally annotated genes.

**Additional file 3.** Differentially expresses genes (DEGs) determined by DESeq2 for berry skin samples at 22°Brix from BOD and RNO.

**Additional file 4.** Statistical results from gene set enrichment analysis of the DEGs using topGO.

**Additional file 5.** Image of the top 25 connected GO categories in the topGO network of the DEGs in Additional file 4.

**Additional file 6.** Module membership (MM) of all filtered transcripts as defined by WGCNA. Values are the kME (module eigengene connectivity).

**Additional file 7.** Heatmap correlation of berry traits (°Brix level, BOD, RNO) of each of 19 gene modules. Gene modules were identified by a color name (MMcolornumber) as assigned by the WGCNA R package. Values in each heatmap block are the correlation (left value) and p-value (in parentheses) of the module with the berry trait.

**Additional file 8.** topGO analysis of all modules with genes having a kME >0.80. The results for each module are on a separate tab within the file.

**Additional file 9.** topGO analysis of the top 400 genes with the greatest differential expression between high and low °Brix samples in BOD berry skins.

**Additional file 10.** topGO analysis of the top 400 genes with the greatest differential expression between high and low °Brix samples in RNO berry skins.

**Additional file 11.** Transcript profiles of stilbene synthase (STS) genes that were differentially expressed.

**Additional file 12**. Transcript abundance of circadian clock genes from BOD and RNO berry skins. The data are placed on a circadian clock model derived from [4]. Lines in the model represent known interactions between genes; red arrows are positive interactions, black lines are negative interactions, and blue lines indicate direct physical interactions but the direction, positive or negative, is unknown. No lines indicate that there are no known interactions at this time. Transcript profiles outlined in red highlight significantly higher transcript abundance for the BOD berries.

**Additional file 13**. A model of the peripheral genes including light sensing genes that interact with core circadian clock genes in BOD and RNO berry skins. Lines represent gene interactions as described in Additional file 12. Red and blue lightning bolts represent the reception of their respective light wavelengths for each gene symbol. Transcript profiles outlined in red highlight significantly higher transcript abundance for the BOD berries.

**Additional file 14**. Transcript profiles of ABA biosynthesis and signaling genes that are differentially expressed between BOD and RNO berry skins.

**References**

1.      This P, Lacombe T, Thomas MR: **Historical origins and genetic diversity of wine grapes**. *Trends in Genetics* 2006, **22**(9):511-519.

2.      Kuhn N, Guan L, Dai ZW, Wu BH, Lauvergeat V, Gomes E, Li SH, Godoy F, Arce-Johnson P, Delrot S: **Berry ripening: recently heard through the grapevine**. *Journal of Experimental Botany* 2014, **65**(16):4543-4559.

3.      Fortes AM, Teixeira RT, Agudelo-Romero P: **Complex Interplay of Hormonal Signals during Grape Berry Ripening**. *Molecules* 2015, **20**(5):9326-9343.

4.      Ghan R, Petereit J, Tillett RL, Schlauch KA, Toubiana D, Fait A, Cramer GR: **The common transcriptional subnetworks of the grape berry skin in the late stages of ripening**. *BMC Plant Biol* 2017, **17**(1):94.

5.      Gonzalez-Barreiro C, Rial-Otero R, Cancho-Grande B, Simal-Gandara J: **Wine aroma compounds in grapes: a critical review**. *Crit Rev Food Sci Nutr* 2015, **55**(2):202-218.

6.      Boss PK, Kalua CM, Nicholson EL, Maffei SM, Böttcher C, Davies C: **Fermentation of grapes throughout development identifies stages critical to the development of wine volatile composition**. *Australian Journal of Grape and Wine Research* 2018, **24**(1):24-37.

7.      Boss PK, Pearce AD, Zhao Y, Nicholson EL, Dennis EG, Jeffery DW: **Potential grape-derived contributions to volatile ester concentrations in wine**. *Molecules* 2015, **20**(5):7845-7873.

8.      Dennis EG, Keyzers RA, Kalua CM, Maffei SM, Nicholson EL, Boss PK: **Grape contribution to wine aroma. Production of hexyl acetate, octyl acetate and benzyl acetate during yeast fermentation is dependent upon precursors in the must**. *Journal of Agricultural and Food Chemistry* 2012.

9.      Sun P, Schuurink RC, Caissard JC, Hugueney P, Baudino S: **My Way: Noncanonical Biosynthesis Pathways for Plant Volatiles**. *Trends Plant Sci* 2016, **21**(10):884-894.

10.     Schwab W, Wust M: **Understanding the Constitutive and Induced Biosynthesis of Mono- and Sesquiterpenes in Grapes (Vitis vinifera): A Key to Unlocking the Biochemical Secrets of Unique Grape Aroma Profiles**. *J Agric Food Chem* 2015, **63**(49):10591-10603.

11.    Deluc LG, Quilici DR, Decendit A, Grimplet J, Wheatley MD, Schlauch KA, Merillon JM, Cushman JC, Cramer GR: **Water deficit alters differentially metabolic pathways affecting important flavor and quality traits in grape berries of Cabernet Sauvignon and Chardonnay**. *BMC Genomics* 2009, **10**:212.

12.    Cramer GR, Evans J, Ardelean R, Keady M, Quilici D, Schooley DA: **Impacts of regulated-deficit irrigation on the flavor components of grapes and wines.** In: *Macromolecules and Secondary Metabolites of Grapevine and Wine.* Edited by Jeandet P, Clément C, Conreaux A. Paris: Lavoisier; 2007: 53-59.

13.    Koundouras S, Marinos V, Gkoulioti A, Kotseridis Y, van Leeuwen C: **Influence of vineyard location and vine water status on fruit maturation of nonirrigated cv. Agiorgitiko (*Vitis vinifera* L.). Effects on wine phenolic and aroma components**. *J Agric Food Chem* 2006, **54**:5077-5086.

14.    Drappier J, Thibon C, Rabot A, Geny-Denis L: **Relationship between wine composition and temperature: Impact on Bordeaux wine typicity in the context of global warming— Review**. *Critical Reviews in Food Science and Nutrition* 2017:1-17.

15.    van Leeuwen C, Friant P, Choné X, Tregoat O, Koundouras S, Dubourdieu D: **Influence of Climate, Soil, and Cultivar on Terroir**. *American Journal of Enology and Viticulture* 2004, **55**(3):207.

16.    Martínez-Lüscher J, Kizildeniz T, Vučetić V, Dai Z, Luedeling E, van Leeuwen C, Gomès E, Pascual I, Irigoyen JJ, Morales F, Delrot S: **Sensitivity of Grapevine Phenology to Water Availability, Temperature and CO2 Concentration**. *Frontiers in Environmental Science* 2016, **4**:48.

17.    Bokulich NA, Collins TS, Masarweh C, Allen G, Heymann H, Ebeler SE, Mills DA: **Associations among Wine Grape Microbiome, Metabolome, and Fermentation Behavior Suggest Microbial Contribution to Regional Wine Characteristics**. *mBio* 2016, **7**(3).

18.    Berdeja M, Nicolas P, Kappel C, Dai ZW, Hilbert G, Peccoux A, Lafontaine M, Ollat N, Gomès E, Delrot S: **Water limitation and rootstock genotype interact to alter grape berry metabolism through transcriptome reprogramming**. *Horticulture Research* 2015, **2**:15012.

19.    Marè C, Aprile A, Roncaglia E, Tocci E, Corino LG, De Bellis L, Cattivelli L: **Rootstock and soil induce transcriptome modulation of phenylpropanoid pathway in grape leaves**. *Journal of Plant Interactions* 2013, **8**(4):334-349.

20.    Cookson SJ, Ollat N: **Grafting with rootstocks induces extensive transcriptional re-programming in the shoot apical meristem of grapevine**. *BMC Plant Biol* 2013, **13**:147.

21.    Corso M, Vannozzi A, Ziliotto F, Zouine M, Maza E, Nicolato T, Vitulo N, Meggio F, Valle G, Bouzayen M, Müller M, Munné-Bosch S, Lucchin M, Bonghi C: **Grapevine Rootstocks Differentially Affect the Rate of Ripening and Modulate Auxin-Related Genes in Cabernet Sauvignon Berries**. *Frontiers in plant science* 2016, **7**:69.

22.    Cochetel N, Escudie F, Cookson SJ, Dai Z, Vivin P, Bert PF, Munoz MS, Delrot S, Klopp C, Ollat N, Lauvergeat V: **Root transcriptomic responses of grafted grapevines to heterogeneous nitrogen availability depend on rootstock genotype**. *Journal of Experimental Botany* 2017, **68**(15):4339-4355.

23.    Downey MO, Dokoozlian NK, Krstic MP: **Cultural Practice and Environmental Impacts on the Flavonoid Composition of Grapes and Wine: A Review of Recent Research**. *American Journal of Enology and Viticulture* 2006, **57**(3):257-268.

24.    Young PR, Eyeghe-Bickong HA, du Plessis K, Alexandersson E, Jacobson DA, Coetzee Z, Deloire A, Vivier MA: **Grapevine Plasticity in Response to an Altered Microclimate: Sauvignon Blanc Modulates Specific Metabolites in Response to Increased Berry Exposure**. *Plant Physiology* 2016, **170**(3):1235-1254.

25.    Fabres PJ, Collins C, Cavagnaro TR, Rodriguez Lopez CM: **A Concise Review on Multi-Omics Data Integration for Terroir Analysis in Vitis vinifera**. *Frontiers in plant science* 2017, **8**:1065.

26.    Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C: **Salmon provides fast and bias-aware quantification of transcript expression**. *Nat Methods* 2017, **14**(4):417-419.

27.    Canaguier A, Grimplet J, Di Gaspero G, Scalabrin S, Duchene E, Choisne N, Mohellibi N, Guichard C, Rombauts S, Le Clainche I, Berard A, Chauveau A, Bounon R, Rustenholz C, Morgante M, Le Paslier MC, Brunel D, Adam-Blondon AF: **A new version of the grapevine reference genome assembly (12X.v2) and of its annotation (VCost.v3)**. *Genom Data* 2017, **14**:56-62.

28.    Love MI, Huber W, Anders S: **Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2**. *Genome Biology* 2014, **15**(12):1-21.

29.    Guo P, Li Z, Huang P, Li B, Fang S, Chu J, Guo H: **A Tripartite Amplification Loop Involving the Transcription Factor**

**WRKY75, Salicylic Acid, and Reactive Oxygen Species Accelerates Leaf Senescence**. *The Plant Cell* 2017, **29**(11):2854-2870.

30. Qiu K, Li Z, Yang Z, Chen J, Wu S, Zhu X, Gao S, Gao J, Ren G, Kuai B, Zhou X: **EIN3 and ORE1 Accelerate Degreening during Ethylene-Mediated Leaf Senescence by Directly Activating Chlorophyll Catabolic Genes in Arabidopsis**. *PLoS Genet* 2015, **11**(7):e1005399.

31. Li Z, Peng J, Wen X, Guo H: **Ethylene-insensitive3 is a senescence-associated gene that accelerates age-dependent leaf senescence by directly repressing miR164 transcription in Arabidopsis**. *The Plant Cell* 2013, **25**(9):3311-3328.

32. Martin DM, Aubourg S, Schouwey MB, Daviet L, Schalk M, Toub O, Lund ST, Bohlmann J: **Functional annotation, genome organization and phylogeny of the grapevine (Vitis vinifera) terpene synthase gene family based on genome assembly, FLcDNA cloning, and enzyme assays**. *BMC Plant Biology* 2010, **10**:226.

33. Dudareva N, Martin D, Kish CM, Kolosova N, Gorenstein N, Faldt J, Miller B, Bohlmann J: **(E)-beta-ocimene and myrcene synthase genes of floral scent biosynthesis in snapdragon: function and expression of three terpene synthase genes of a new terpene synthase subfamily**. *The Plant Cell* 2003, **15**(5):1227-1241.

34. Young PR, Lashbrooke JG, Alexandersson E, Jacobson D, Moser C, Velasco R, Vivier MA: **The genes and enzymes of the carotenoid metabolic pathway in *Vitis vinifera* L**. *BMC Genomics* 2012, **13**:243.

35. Jung J-H, Domijan M, Klose C, Biswas S, Ezer D, Gao M, Khattak AK, Box MS, Charoensawan V, Cortijo S, Kumar M, Grant A, Locke JCW, Schäfer E, Jaeger KE, Wigge PA: **Phytochromes function as thermosensors in Arabidopsis**. *Science* 2016, **354**(6314):886.

36. Legris M, Klose C, Burgie ES, Rojas CC, Neme M, Hiltbrunner A, Wigge PA, Schafer E, Vierstra RD, Casal JJ: **Phytochrome B integrates light and temperature signals in Arabidopsis**. *Science* 2016, **354**(6314):897-900.

37. Tattersall EA, Grimplet J, Deluc L, Wheatley MD, Vincent D, Osborne C, Ergul A, Lomen E, Blank RR, Schlauch KA, Cushman JC, Cramer GR: **Transcript abundance profiles reveal larger and more complex responses of grapevine to chilling compared to osmotic and salinity stress**. *Funct Integr Genomics* 2007, **7**(4):317-333.

38. Jaglo-Ottoson KR, Gilmour SJ, Zarka DG, Schabenberger O, Thomashow MF: *Arabidopsis* **CBF1 overexpression induces COR genes and enhances freezing tolerance**. *Science* 1998, **280**:104-106.

39. Fernandez O, Vandesteene L, Feil R, Baillieul F, Lunn JE, Clement C: **Trehalose metabolism is activated upon chilling in grapevine and might participate in Burkholderia phytofirmans induced chilling tolerance**. *Planta* 2012, **236**(2):355-369.

40. Lunn JE, Delorge I, Figueroa CM, Van Dijck P, Stitt M: **Trehalose metabolism in plants**. *Plant Journal* 2014, **79**(4):544-567.

41. Fung RW, Gonzalo M, Fekete C, Kovacs LG, He Y, Marsh E, McIntyre LM, Schachtman DP, Qiu W: **Powdery mildew induces defense-oriented reprogramming of the transcriptome in a susceptible but not in a resistant grapevine**. *Plant Physiology* 2008, **146**(1):236-249.

42. Davies C, Boss PK, Robinson SP: **Treatment of grape berries, a nonclimacteric fruit with a synthetic auxin, retards ripening and alters the expression of developmentally regulated genes**. *Plant Physiology* 1997, **115**:1155-1161.

43. Böttcher C, Boss PK, Davies C: **Delaying Riesling grape berry ripening with a synthetic auxin affects malic acid metabolism and sugar accumulation, and alters wine sensory characters**. *Functional Plant Biology* 2012, **39**(9):745.

44. Cramer GR: **Abiotic stress & plant responses from the whole vine to the genes**. *Australian Journal of Grape and Wine Research* 2010, **16**:86-93.

45. Tan BC, Joseph LM, Deng WT, Liu L, Li QB, Cline K, McCarty DR: **Molecular characterization of the Arabidopsis 9-cis epoxycarotenoid dioxygenase gene family**. *Plant Journal* 2003, **35**(1):44-56.

46. Frey A, Effroy D, Lefebvre V, Seo M, Perreau F, Berger A, Sechet J, To A, North HM, Marion-Poll A: **Epoxycarotenoid cleavage by NCED5 fine-tunes ABA accumulation and affects seed dormancy and drought tolerance with other NCED family members**. *Plant Journal* 2012, **70**(3):501-512.

47. Rattanakon S, Ghan R, Gambetta GA, Deluc LG, Schlauch KA, Cramer GR: **Abscisic acid transcriptomic signaling varies with grapevine organ**. *BMC Plant Biol* 2016, **16**(1):72.

48. Hopper DW, Ghan R, Schlauch KA, Cramer GR: **Transcriptomic network analyses of leaf dehydration responses identify highly connected ABA and ethylene signaling hubs in three**

grapevine species differing in drought tolerance. *BMC Plant Biol* 2016, **16**(1):118.

49. Takahashi F, Suzuki T, Osakabe Y, Betsuyaku S, Kondo Y, Dohmae N, Fukuda H, Yamaguchi-Shinozaki K, Shinozaki K: **A small peptide modulates stomatal control via abscisic acid in long-distance signalling**. *Nature* 2018.

50. Broekaert WF, Delaure SL, De Bolle MF, Cammue BP: **The role of ethylene in host-pathogen interactions**. *Annu Rev Phytopathol* 2006, **44**:393-416.

51. Dubois M, Van den Broeck L, Inze D: **The Pivotal Role of Ethylene in Plant Growth**. *Trends Plant Sci* 2018, **23**(4):311-323.

52. Liu M, Pirrello J, Chervin C, Roustan JP, Bouzayen M: **Ethylene Control of Fruit Ripening: Revisiting the Complex Network of Transcriptional Regulation**. *Plant Physiology* 2015, **169**(4):2380-2390.

53. Chang KN, Zhong S, Weirauch MT, Hon G, Pelizzola M, Li H, Huang SS, Schmitz RJ, Urich MA, Kuo D, Nery JR, Qiao H, Yang A, Jamali A, Chen H, Ideker T, Ren B, Bar-Joseph Z, Hughes TR, Ecker JR: **Temporal transcriptional response to ethylene gas drives growth hormone cross-regulation in Arabidopsis**. *Elife* 2013, **2**:e00675.

54. Van den Broeck L, Dubois M, Vermeersch M, Storme V, Matsui M, Inze D: **From network to phenotype: the dynamic wiring of an Arabidopsis transcriptional network induced by osmotic stress**. *Mol Syst Biol* 2017, **13**(12):961.

55. Jiao Y, Ma L, Strickland E, Deng XW: **Conservation and divergence of light-regulated genome expression patterns during seedling development in rice and Arabidopsis**. *The Plant Cell* 2005, **17**(12):3239-3256.

56. Hu W, Su YS, Lagarias JC: **A light-independent allele of phytochrome B faithfully recapitulates photomorphogenic transcriptional networks**. *Mol Plant* 2009, **2**(1):166-182.

57. Delker C, van Zanten M, Quint M: **Thermosensing Enlightened**. *Trends Plant Sci* 2017, **22**(3):185-187.

58. Hsu PY, Harmer SL: **Wheels within wheels: the plant circadian system**. *Trends Plant Sci* 2014, **19**(4):240-249.

59. Johansson M, Staiger D: **Time to flower: interplay between photoperiod and the circadian clock**. *Journal of Experimental Botany* 2015, **66**(3):719-730.

60. Huang H, Alvarez S, Bindbeutel R, Shen Z, Naldrett MJ, Evans BS, Briggs SP, Hicks LM, Kay SA, Nusinow DA: **Identification of Evening Complex Associated Proteins in Arabidopsis by Affinity Purification and Mass Spectrometry**. *Mol Cell Proteomics* 2016, **15**(1):201-217.

61. Jiang Z, Xu G, Jing Y, Tang W, Lin R: **Phytochrome B and REVEILLE1/2-mediated signalling controls seed dormancy and germination in Arabidopsis**. *Nat Commun* 2016, **7**:12377.

62. Leivar P, Monte E, Al-Sady B, Carle C, Storer A, Alonso JM, Ecker JR, Quail PH: **The Arabidopsis phytochrome-interacting factor PIF7, together with PIF3 and PIF4, regulates responses to prolonged red light by modulating phyB levels**. *The Plant Cell* 2008, **20**(2):337-352.

63. Kidokoro S, Maruyama K, Nakashima K, Imura Y, Narusaka Y, Shinwari ZK, Osakabe Y, Fujita Y, Mizoi J, Shinozaki K, Yamaguchi-Shinozaki K: **The phytochrome-interacting factor PIF7 negatively regulates DREB1 expression under circadian control in Arabidopsis**. *Plant Physiology* 2009, **151**(4):2046-2057.

64. Rawat R, Schwartz J, Jones MA, Sairanen I, Cheng Y, Andersson CR, Zhao Y, Ljung K, Harmer SL: **REVEILLE1, a Myb-like transcription factor, integrates the circadian clock and auxin pathways**. *Proc Natl Acad Sci U S A* 2009, **106**(39):16883-16888.

65. Gupta SK, Sharma S, Santisree P, Kilambi HV, Appenroth K, Sreelakshmi Y, Sharma R: **Complex and shifting interactions of phytochromes regulate fruit development in tomato**. *Plant Cell Environment* 2014, **37**(7):1688-1702.

66. Kuhn N, Ormeno-Nunez J, Jaque-Zamora G, Perez FJ: **Photoperiod modifies the diurnal expression profile of VvPHYA and VvPHYB transcripts in field-grown grapevine leaves**. *Journal of Plant Physiology* 2009, **166**(11):1172-1180.

67. Stockinger EJ, Gilmour SJ, Thomashow MF: ***Arabidopsis thaliana* CBF1 encodes an AP2 domain-containing transcriptional activator that binds to the C-repeat/DRE, a cis-acting DNA regulatory element that stimulates transcription in response to low temperature and water deficit**. *Proceedings of the National Academy of Sciences of the United States of America* 1997, **94**(February):1035-1040.

68. Tillett RL, Wheatley MD, Tattersall EA, Schlauch KA, Cramer GR, Cushman JC: **The Vitis vinifera C-repeat binding protein 4 (VvCBF4) transcriptional factor enhances freezing tolerance in wine grape**. *Plant Biotechnology Journal* 2012, **10**(1):105-124.

69. Cramer GR, Ghan R, Schlauch KA, Tillett RL, Heymann H, Ferrarini A, Delledonne M, Zenoni S, Fasoli M, Pezzotti M: **Transcriptomic analysis of the late stages of grapevine ( Vitis vinifera cv. Cabernet Sauvignon) berry ripening reveals significant induction of ethylene signaling and flavor pathways in the skin**. *BMC Plant Biol* 2014, **14**(1):370.

70. Nakashima K, Yamaguchi-Shinozaki K: **ABA signaling in stress-response and seed development**. *Plant Cell Rep* 2013, **32**(7):959-970.

71. Nicolas P, Lecourieux D, Kappel C, Cluzet S, Cramer G, Delrot S, Lecourieux F: **The basic leucine zipper franscription factor ABSCISIC ACID RESPONSE ELEMENT-BINDING FACTOR2 is an important transcriptional regulator of abscisic acid-dependent grape berry ripening processes**. *Plant Physiology* 2014, **164**(1):365-383.

72. Wang R, Estelle M: **Diversity and specificity: auxin perception and signaling through the TIR1/AFB pathway**. *Curr Opin Plant Biol* 2014, **21**:51-58.

73. Liu X, Zhang H, Zhao Y, Feng Z, Li Q, Yang HQ, Luan S, Li J, He ZH: **Auxin controls seed dormancy through stimulation of abscisic acid signaling by inducing ARF-mediated ABI3 activation in Arabidopsis**. *Proc Natl Acad Sci U S A* 2013, **110**(38):15485-15490.

74. Wang L, Hua D, He J, Duan Y, Chen Z, Hong X, Gong Z: **Auxin Response Factor2 (ARF2) and its regulated homeodomain gene HB33 mediate abscisic acid response in Arabidopsis**. *PLoS Genet* 2011, **7**(7):e1002172.

75. Jeandet P, Douillet-Breuil AC, Bessis R, Debord S, Sbaghi M, Adrian M: **Phytoalexins from the Vitaceae: biosynthesis, phytoalexin gene expression in transgenic plants, antifungal activity, and metabolism**. *J Agric Food Chem* 2002, **50**(10):2731-2741.

76. Dufour MC, Magnin N, Dumas B, Vergnes S, Corio-Costet MF: **High-throughput gene-expression quantification of grapevine defense responses in the field using microfluidic dynamic arrays**. *BMC Genomics* 2016, **17**(1):957.

77. Jacobs AK, Dry IB, Robinson SP: **Induction of different pathogenesis-related cDNAs in grapevine infected with powdery mildew and treated with ethephon**. *Plant Pathology* 1999, **48**(3):325-336.

78. Belhadj A, Telef N, Cluzet S, Bouscaut J, Corio-Costet MF, Merillon JM: **Ethephon elicits protection against Erysiphe**

**necator in grapevine**. *J Agric Food Chem* 2008, **56**(14):5781-5787.

79.   Dixon RA, Paiva NL: **Stress-Induced Phenylpropanoid Metabolism**. *The Plant Cell* 1995, **7**(7):1085-1097.

80.   Huang J, Gu M, Lai Z, Fan B, Shi K, Zhou YH, Yu JQ, Chen Z: **Functional analysis of the Arabidopsis PAL gene family in plant growth, development, and response to environmental stress**. *Plant Physiology* 2010, **153**(4):1526-1538.

81.   Shi P, Li B, Chen H, Song C, Meng J, Xi Z, Zhang Z: **Iron Supply Affects Anthocyanin Content and Related Gene Expression in Berries of Vitis vinifera cv. Cabernet Sauvignon**. *Molecules* 2017, **22**(2).

82.   Manara A, DalCorso G, Leister D, Jahns P, Baldan B, Furini A: **AtSIA1 AND AtOSA1: two Abc1 proteins involved in oxidative stress responses and iron distribution within chloroplasts**. *New Phytol* 2014, **201**(2):452-465.

83.   Chen CC, Chien WF, Lin NC, Yeh KC: **Alternative functions of Arabidopsis Yellow Stripe-Like3: from metal translocation to pathogen defense**. *PLoS One* 2014, **9**(5):e98008.

84.   Chu HH, Chiecko J, Punshon T, Lanzirotti A, Lahner B, Salt DE, Walker EL: **Successful reproduction requires the function of Arabidopsis Yellow Stripe-Like1 and Yellow Stripe-Like3 metal-nicotianamine transporters in both vegetative and reproductive structures**. *Plant Physiology* 2010, **154**(1):197-210.

85.   Ravet K, Touraine B, Boucherez J, Briat JF, Gaymard F, Cellier F: **Ferritins control interaction between iron homeostasis and oxidative stress in Arabidopsis**. *Plant Journal* 2009, **57**(3):400-412.

86.   Ravet K, Touraine B, Kim SA, Cellier F, Thomine S, Guerinot ML, Briat JF, Gaymard F: **Post-translational regulation of AtFER2 ferritin in response to intracellular iron trafficking during fruit development in Arabidopsis**. *Mol Plant* 2009, **2**(5):1095-1106.

87.   Tarantino D, Petit JM, Lobreaux S, Briat JF, Soave C, Murgia I: **Differential involvement of the IDRS cis-element in the developmental and environmental regulation of the AtFer1 ferritin gene from Arabidopsis**. *Planta* 2003, **217**(5):709-716.

88.   Conte S, Stevenson D, Furner I, Lloyd A: **Multiple antibiotic resistance in Arabidopsis is conferred by mutations in a chloroplast-localized transport protein**. *Plant Physiology* 2009, **151**(2):559-573.

89. Seo PJ, Park J, Park MJ, Kim YS, Kim SG, Jung JH, Park CM: **A Golgi-localized MATE transporter mediates iron homoeostasis under osmotic stress in Arabidopsis**. *Biochemistry Journal* 2012, **442**(3):551-561.

90. Moyer MM, Singer SD, Davenport JR, Hoheisel G-A: **Vineyard Nutrient Management in Washington State**. In.: Washington State University Extension; 2018: 1-45.

91. Dal Santo S, Tornielli GB, Zenoni S, Fasoli M, Farina L, Anesi A, Guzzo F, Delledonne M, Pezzotti M: **The plasticity of the grapevine berry transcriptome**. *Genome Biol* 2013, **14**(6):r54.

92. Dal Santo S, Fasoli M, Negri S, D'Inca E, Vicenzi N, Guzzo F, Tornielli GB, Pezzotti M, Zenoni S: **Plasticity of the Berry Ripening Program in a White Grape Variety**. *Frontiers in plant science* 2016, **7**:970.

93. Sun R, He F, Lan Y, Xing R, Liu R, Pan Q, Wang J, Duan C: **Transcriptome comparison of Cabernet Sauvignon grape berries from two regions with distinct climate**. *Journal of Plant Physiology* 2015, **178**:43-54.

94. Wen YQ, Zhong GY, Gao Y, Lan YB, Duan CQ, Pan QH: **Using the combined analysis of transcripts and metabolites to propose key genes for differential terpene accumulation across two regions**. *BMC Plant Biol* 2015, **15**:240.

95. Singh B, Sharma RA: **Plant terpenes: defense responses, phylogenetic analysis, regulation and clinical applications**. *3 Biotech* 2015, **5**(2):129-151.

96. Zhang P, Barlow S, Krstic M, Herderich M, Fuentes S, Howell K: **Within-Vineyard, Within-Vine, and Within-Bunch Variability of the Rotundone Concentration in Berries of Vitis vinifera L. cv. Shiraz**. *J Agric Food Chem* 2015, **63**(17):4276-4283.

97. Ou C, Du X, Shellie K, Ross C, Qian MC: **Volatile compounds and sensory attributes of wine from Cv. Merlot (Vitis vinifera L.) grown under differential levels of water deficit with or without a kaolin-based, foliar reflectant particle film**. *J Agric Food Chem* 2010, **58**(24):12890-12898.

98. Degu A, Ayenew B, Cramer GR, Fait A: **Polyphenolic responses of grapevine berries to light, temperature, oxidative stress, abscisic acid and jasmonic acid show specific developmental-dependent degrees of metabolic resilience to perturbation**. *Food Chemistry* 2016, **212**:828-836.

99. Rienth M, Torregrosa L, Luchaire N, Chatbanyong R, Lecourieux D, Kelly MT, Romieu C: **Day and night heat stress trigger**

**different transcriptomic responses in green and ripening grapevine (vitis vinifera) fruit**. *BMC Plant Biol* 2014, **14**:108.

100. Bonada M, Sadras VO: **Review: critical appraisal of methods to investigate the effect of temperature on grapevine berry composition**. *Australian Journal of Grape and Wine Research* 2015, **21**(1):1-17.

101. Bock A, Sparks TH, Estrella N, Menzel A: **Climate-Induced Changes in Grapevine Yield and Must Sugar Content in Franconia (Germany) between 1805 and 2010**. *PLOS ONE* 2013, **8**(7):e69015.

102. Petrie PR, Sadras VO: **Advancement of grapevine maturity in Australia between 1993 and 2006: putative causes, magnitude of trends and viticultural consequences**. *Australian Journal of Grape and Wine Research* 2008, **14**(1):33-45.

103. Jones GV, Davis RE: **Climate influences on grapevine phenology, grape composition, and wine production and quality for Bordeaux, France**. *American Journal of Enology and Viticulture [print]* 2000, **51**(3):249-261.

104. Duchêne E, Schneider C: **Grapevine and climatic changes: a glance at the situation in Alsace**. *Agron Sustain Dev* 2005, **25**(1):93-99.

105. Sadras VO, Moran MA: **Elevated temperature decouples anthocyanins and sugars in berries of Shiraz and Cabernet Franc**. *Australian Journal of Grape and Wine Research* 2012, **18**(2):115-122.

106. Buttrose MS, Hale CR, Kliewer WM: **Effect of Temperature on the Composition of 'Cabernet Sauvignon' Berries**. *American Journal of Enology and Viticulture* 1971, **22**(2):71.

107. Guillaumie S, Fouquet R, Kappel C, Camps C, Terrier N, Moncomble D, Dunlevy JD, Davies C, Boss PK, Delrot S: **Transcriptional analysis of late ripening stages of grapevine berry**. *BMC Plant Biol* 2011, **11**:165.

108. Zelleke A, Kliewer WM: **Influence of Root Temperature and Rootstock on Budbreak, Shoot Growth, and Fruit Composition of Cabernet Sauvignon Grapevines Grown under Controlled Conditions**. *American Journal of Enology and Viticulture* 1979, **30**(4):312.

109. Bergqvist J, Dokoozlian N, Ebisuda N: **Sunlight Exposure and Temperature Effects on Berry Growth and Composition of Cabernet Sauvignon and Grenache in the Central San Joaquin**

**Valley of California**. *American Journal of Enology and Viticulture* 2001, **52**(1):1.

110. Kliewer WM, Dokoozlian NK: **Leaf Area/Crop Weight Ratios of Grapevines: Influence on Fruit Composition and Wine Quality**. *American Journal of Enology and Viticulture* 2005, **56**(2):170.

111. Triolo R, Roby JP, Plaia A, Hilbert G, Buscemi S, Di Lorenzo R, van Leeuwen C: **Hierarchy of Factors Impacting Grape Berry Mass: Separation of Direct and Indirect Effects on Major Berry Metabolites**. *American Journal of Enology and Viticulture* 2018, **69**(2):103.

112. Liu Y, Bassham DC: **Autophagy: pathways for self-eating in plant cells**. *Annu Rev Plant Biol* 2012, **63**:215-237.

113. Destrac-Irvine A, Van Leeuwen C: **The VitAdapt project: extensive phenotyping of a wide range of varieties in order to optimize the use of genetic diversity within the *Vitis vinifera* species as a tool for adaptation to a changing environment.** Villnave D'Ornon, France: Vigne & Vin Publications Internationales; 2017.

114. Ghan R, Van Sluyter SC, Hochberg U, Degu A, Hopper DW, Tillet RL, Schlauch KA, Haynes PA, Fait A, Cramer GR: **Five omic technologies are concordant in differentiating the biochemical characteristics of the berries of five grapevine (Vitis vinifera L.) cultivars**. *BMC Genomics* 2015, **16**(1):946.

115. Reid KE, Olsson N, Schlosser J, Peng F, Lund ST: **An optimized grapevine RNA isolation procedure and statistical determination of reference genes for real-time RT-PCR during berry development**. *BMC Plant Biol* 2006, **6**:27.

116. Andrews S: **FastQC**. In*. https://www.bioinformatics.babraham.ac.uk/projects/fastqc/*; 2014.

117. Bolger AM, Lohse M, Usadel B: **Trimmomatic: a flexible trimmer for Illumina sequence data**. *Bioinformatics* 2014, **30**(15):2114-2120.

118. Soneson C, Love MI, Robinson MD: **Differential analyses for RNA-seq: transcript-level estimates improve gene-level inferences**. *F1000Res* 2015, **4**:1521.

119. Langfelder P, Horvath S: **WGCNA: an R package for weighted correlation network analysis**. *BMC Bioinformatics* 2008, **9**:559.

120. Langfelder P, Horvath S: **Fast R Functions for Robust Correlations and Hierarchical Clustering**. *J Stat Softw* 2012, **46**(11).

121. Buchfink B, Xie C, Huson DH: **Fast and sensitive protein alignment using DIAMOND**. *Nature Methods* 2014, **12**:59.

122. **topGO: Enrichment Analysis for Gene Ontology. R package version 2.36.0.** [http://www.mpi-sb.mpg.de/~alexa]

123. Jones P, Binns D, Chang HY, Fraser M, Li W, McAnulla C, McWilliam H, Maslen J, Mitchell A, Nuka G, Pesseat S, Quinn AF, Sangrador-Vegas A, Scheremetjew M, Yong SY, Lopez R, Hunter S: **InterProScan 5: genome-scale protein function classification**. *Bioinformatics* 2014, **30**(9):1236-1240.

124. Leinonen R, Sugawara H, Shumway M, International Nucleotide Sequence Database C: **The sequence read archive**. *Nucleic Acids Res* 2011, **39**(Database issue):D19-21.