Dopamine waves as a mechanism for spatiotemporal credit assignment

Arif A. Hamid[1], Michael J. Frank[2,3]*, Christopher I. Moore[1,3]*

[1]Department of Neuroscience, [2]Department of Cognitive Linguistics & Psychological Sciences,
[3]Carney Institute for Brain Science, Brown University, Providence, RI

* These authors contributed equally to this work; alphabetical order

Abstract

Significant evidence supports the view that dopamine shapes reward-learning by
encoding prediction errors. However, it is unknown whether dopamine decision-signals are
tailored to the functional specialization of target regions. Here, we report a novel set of wave-like
spatiotemporal activity-patterns in dopamine axons across the dorsal striatum. These waves
switch between different activational motifs and organize dopamine transients into localized
clusters within functionally related striatal subregions. These specific motifs are associated with
distinct task contexts: At reward delivery, dopamine signals rapidly resynchronize into
propagating waves with opponent directions depending on instrumental task contingencies.
Moreover, dopamine dynamics during reward pursuit signal the extent to which mice have
instrumental control, and interact with reward waves to predict future behavioral adjustments.
Our results are consistent with a computational architecture in which striatal dopamine signals
are sculpted by inference about instrumental controllability, and provide evidence for a
spatiotemporally "vectorized" role of dopamine in credit assignment.

Main text

Dopamine supports reward learning and motivational activation, but details about what
decision variables are encoded, and how they are delivered to postsynaptic targets, continue to
be refined[1-3]. The dopamine-reward prediction error (RPE) hypothesis emphasizes that
dopamine conveys deviations from reward expectation in reinforcement learning (RL) theory[4].
This formulation generally treats dopamine as a "global" spatio-temporally uniform signal, a view
based on two key findings. First, extensively divergent dopamine axons[5,6] provide an
architecture for broadcast-like communication. Second, dopamine cell spikes measured in the
midbrain are highly synchronized[7,8], putatively implementing a redundant population code[9-11] for
RPEs[12]. These observations form the basis for an influential view[13,14] of what dopamine
communicates and how it is delivered: scalar RPEs that are uniformly delivered to all recipient
subregions. The notion of uniform encoding also extends to alternative accounts for dopamine's
role in motivation[2] by relaying scalar value signals[15] .

A key limitation of this global view is that scalar (or, spatio-temporally uniform) decision
variables are neither computationally advantageous, nor reflected in forebrain dopamine
dynamics. Postsynaptic striatal subregions are functionally specialized[16,17], receiving distinct
cortical and thalamic afferents[18,19], and express unique compliments of biomarkers[20].
Accordingly, rewards[21], their motivated pursuit[15,22] and predictive stimuli[23] produce vastly
different dopamine time courses across the dorsal-ventral and medial-lateral axes of the
striatum. While these observations indicate regional heterogeneity, the extent to which
dopamine inputs reflect the computational requirements of postsynaptic areas remains elusive.
For example, there is some theoretical motivation[24,25] and empirical support[26,27] for delivery of
vector-valued RPEs that depend on a target region's computational specialty. Nonetheless, we
currently lack a clear understanding of organizing principles for striatal dopamine activity, and
what normative computational functions may be served by such heterogeneity.

Related striatal subregions get correlated dopamine input

We set out to characterize the spatio-temporal organizational rules of dopamine activity
across the dorsal striatum. Standard methods for dopamine measurement typically survey small
territories (10s – 100s of micrometers) and are ill-suited to probing large-scale organization. To
overcome these limitations, we injected *cre*-dependent GCAMP6f into the midbrain of DAT-cre
mice, and imaged dopamine axons through a 7mm^2 chronic imaging window over the dorsal
striatum (DS)[28] (**Fig. 1a**). This approach provided optical access to 60-80% of the dorsal surface
of the mouse striatum, with a view of dorsomedial (DMS), dorsolateral (DLS) and partial access
to the posterior-tail (TS) region of the striatum (**Fig. 1b**). We imaged the activity of dopamine
axons at multiple levels of resolution with one or two photon microscopy.

Figure 1

Figure 1: Dorsal striatal dopamine activity is spatiotemporally asynchronous and clusters into contiguous territories.

a, Schematic of methods for imaging dopamine axons over dorsal striatum. GCaMP6 was injected into midbrain of DAT-cre mice. Cortex overlying dorsal striatum was aspirated, together with insertion of 3 mm diameter imaging cannula, and activity was imaged in head-restrained mice. **b**, Top-down field of view. **c**, Average delta f/F from two regions (top) that exhibit decorrelated activity (bottom). **d**, Activity of several ROIs from the same session as **c**, time series are sorted such that medial areas are top ROIs, and lateral regions are represented at the bottom. **e**, Correlation matrix across ROIs for different 5sec epochs (highlighted in bottom of **d**), showing patterns of correlations that evolve in time. For example, middle shows global correlation, whereas left and right panels show instances of antagonistic activity patterns in top and bottom set of ROIs. **f**, Results from spatial correlation from seed pixels, evaluating the Pearson's correlation of with all other regions. *Top* panel shows medial seed pixels that are highly correlated with nearby regions, and show graded decrease in correlation for distant regions. Same analysis was repeated for a set of pixels in central striatum (*middle*) and lateral seed pixels (*bottom*). **g**, Quantification of sessions-wide correlation between each pair of pixels as a function of distance, separated by medio-lateral (orange) and anterio-posterior distances. (n= 8 mice, p<0.001 wilcoxon signed-rank test for difference of ML vs AP slopes)

h, Paiwise correlation matrix using hierarchical clustering summarizes similarity of dopamine activity. **i**, *Top*, anatomical projection of pixels that share similarity at the highest cluster limit of two outlining medial and lateral subregions of the dorsal striatum. Increasing the cluster threshold to 20 (*bottom*) revealed smaller, but anatomically contiguous regions of the striatum.

Using a head-fixed preparation, we began by focusing on spontaneous activation of
dopamine axons in mice resting on a wheel in a dark chamber without external stimuli. To first
test if dopamine axons were globally activated, we compared fluorescence signals in DS
regions-of-interest (ROIs) (**Fig. 1c**). While ROIs were sometimes globally synchronized[28], we
observed decorrelated patterns across striatal subregions that evolved across time (**Fig.**
1c,d,e). These patterns of activation were observed across multiple anatomical scales (see
Extended Data Fig. 1 for micron-scale organization), indicating that dopamine afferents can
become recruited asynchronously.

To examine how activity is spatially coordinated, we computed the Pearson's correlation
between pixels' fluorescence as a function of anatomical distance. Dopamine axons showed
strong local correlations that gradually decreased with distance (**Fig. 1f,g**), comparable to the
organization of striatal spiny-neuron activity[29]. Strikingly however, this distance-dependent falloff
was selective to the medio-lateral axis, and was not present on the anterio-posterior axis (**Fig.**
1g), suggesting an organization rule that promoted selective mediolateral decorrelation.

To further examine the topographical organization of dopamine signals, we leveraged
standard cluster analyses (**Fig. 1h**). In every dataset (n = 31 sessions, 8 mice), the highest
cluster threshold identified two contiguous subregions outlining well-established[30,31] DS
subregions; medial (DMS) and lateral (DLS) striatum (**Fig. 1i top**). Further increasing cluster
limits progressively (**Extended Data Fig. 2**) revealed smaller subdomains of DS (**Fig. 1i**
bottom), resembling striatal sub-clusters previously identified based on glutamatergic input
patterns[18]. These areas had similar clustering patterns across days and animals (**Extended**
Data Fig. 3), with 25-30 optimal clusters identified in our field of view (**Extended Data Fig. 4**).
Shuffling the pixelwise temporal or spatial indices produced random clusters (**Extended Data**
Fig. 4), indicating a critical dependence on the underlying spatio-temporal activity pattern.
Together, these results provide evidence for regional coordination of dopamine transmission
and provided an initial basis for evaluating whether these signals are modulated by the
underlying subregion's computational specialty.

Wave-like patterns coordinate dopamine activity

What spatiotemporal patterns produce systematically decorrelated dopamine signals?
We noticed that full-field fluorescence exhibited complex but spatially and temporally continuous
trajectories throughout the striatum, similar to travelling waves described in other cortical and
subcortical brain regions[32-35] (**Fig. 2a,b, Supplementary Video 1**). To quantify these complex
trajectories, we used optic flow algorithms[36] to compute frame-by-frame flow fields (see
methods for details; **Supplementary Video 2**).

Figure 2

Figure 2: Wave-like spatiotemporally continuous sequences of dopamine-axon activation switch between motifs.

a, *Top* row shows individual frames for different epochs of a transient as dopamine axon activity emerges and extinguishes in DS. *Bottom* row displays the corresponding flow vector fields computed for each pixel. Notice the divergence of vector fields during the rise phase of fluorescence, and convergent vectors during fall phase. **b**, Average fluorescence (green) across the entire field of view lasting ~300ms sampled at 40Hz, and corresponding, flow magnitude in the fluorescence signal (red). **c**, Flow trajectory of fluorescence for 5 frames during the onset (*left,* red lines), or offset (*left,* blue lines) phase of the wave from **b**. Each line shows the pattern of flow from individually seeded pixels. **d**, Heatmap quantifying how divergent the vector fields are at each pixel during onset or offset of activity (left and right respectively). A positive value indicates that diverging pattern of flow at each pixel indicating that fluorescence is entering the striatum from those locations. Conversely, negative values are sink regions with converging flow vectors. **e**, Peak-normalized projection of the flow vector divergence for the onset (*top*) or offset (*bottom*) of all transients in one session (n=1516 events). Note that a repeated configuration of pixels serve as sinks and sources. **f**, Distribution of propagation velocity (n=8 mice, 1625 +/- 213 events per mouse). Error bar denotes SEM.**g**, Distribution of interwave intervals for the same data **f**. **h**, *Top,* quiver plot summarizing the direction and magnitude of waves in a single session, and distribution of angle of wave propagation for each animal, *bottom* (n=8 mice, all p < 0.001, Omnibus test for angular uniformity). **i**, *Left,* vector fields (yellow) superimposed onto source pixels (white) for waves that are sourced at the midline and propagate bidirectionally outward. *Right*, corresponding fluorescence time course in ROIs on a medio-lateral gradient of the striatum (*inset*). **j**, Same format as **i**, for lateral source and medial flow or, **k**, medial source and laterally flowing wave.

The onset of activity in GCaMP fluorescence originated from clustered "source"
locations, and rapidly migrated to other regions (**Fig. 2c,d, left**). By contrast, activity terminated
as a result of flow toward "sink" locations (**Fig. 2c,d, right**). A repeated configuration of pixels
had a high probability of serving as sinks and sources (**Fig. 2e, Extended Data Fig. 5**),
indicating that local rules may dictate the initiation and termination of dopamine activity.

Dopamine waves entered the dorsal striatum with exponentially decaying inter-wave-
intervals (**Fig. 2g**) and propagate with a range of velocities (median = 3.8 mm/s, interquartile
range = 2.5, **Fig. 2f**). The overall direction of flow is bimodally distributed, with a biased medial-
lateral propagation axis (**Fig. 2h**, all p < 0.001, Omnibus test for angular uniformity).

We next sought to determine if the collection of complex trajectories were made up of
simpler, repeated sequences that may influence the time course of dopamine arriving at
different parts of the striatum. Indeed, the combination of initiation loci and flow direction gave
rise to motif waves that were scaled by propagation velocity and extent of striatum covered. We
focused our attention on three motifs that produced most of the dopamine transients (**Extended
Data Fig. 5**).

First, source pixels clustered at the juncture of DMS and DLS (**Fig. 2i**) initiated
dopamine activity that rapidly spread bilaterally outward (Type-1*, "Center-Out" or CO wave,* **Fig.
2i, left**). These waves radiate across the stratum with the fastest velocities, arriving at all
subregions with almost zero lag (**Fig. 2i, right***).* Second, source pixels in lateral DLS initiated a
wave that propagates medially (Type-2*, "latero-medial" or LM wave*). LM waves advanced
across the striatum relatively slowly and delivered dopamine transients to DMS that were
delayed relative to DLS in proportion to propagation speed (**Fig. 2j right**). Third, a medially
sourced wave propagated laterally (Type-3*, "medio-lateral" or ML wave*, **Fig. 2k, left**),
terminating in DLS. ML waves activate dopamine axons in the medial striatum first and recruit
lateral regions with substantial delay (**Fig. 2k, right**). Together, these results demonstrate that
wave-like patterns are a fundamental organizational principle of dopamine axonal activity,
prescribing how activity initiates, propagates and terminates across DS.

Rewards evoke directional dopamine waves

What is the functional role of dopamine waves in adaptive behavior? We set out to
determine the computational significance of wave-like trajectories in the context of the well-
studied role of striatal dopamine in instrumental behavior. The dorsal striatum exhibits graded
behavioral specialty, with the DMS orchestrating goal-directed behaviors involving action-
outcome contingencies, and the DLS implicated in stimulus-response behaviors[30,31,37].
Inactivation or manipulation of dopamine in DMS degrades goal-directed planning and action
due to an inability to learn whether rewards are under instrumental control[38,39].

We thus designed two operant tasks intended to manipulate action-outcome
contingencies, and asked whether dopamine dynamics carry information about the degree of
instrumental controllability (**Fig. 3a,b,c**). First, in an 'instrumental' task, rewards were contingent

on mice running on a wheel to traverse linearized distance, with the progress to reward
indicated by an auditory tone that escalated in frequency (**Fig. 3b,d**). On each trial, the distance
that was needed to run for tone transitions (and ultimately, reward) was randomly selected from
a uniform distribution (50-150 cm, **Fig. 3c, left**). Thus, while the mouse was in control of tone
transitions, the specific contingencies varied across trials. A second 'pavlovian' task was
administered in separate sessions. The task structure was identical except tone-transitions
occurred after fixed durations within a trial (randomly drawn, 4-8 sec, **Fig. 3c right**), and
progress to reward was unrelated to running. Trained mice exhibited anticipatory lick trajectories
that increased with ascending tone frequency in both tasks (**Fig. 3e,f**), indicating that mice used
these tones to update their online judgment of progress to reward. Analysis of run bouts
(**Extended Data Fig. 6**) revealed that mice invested goal-directed effort to receive rewards
selectively in the instrumental task.

As in spontaneous conditions reported above, dopamine waves were ubiquitous during
task-performance. Notably, reward delivery immediately resynchronized irregular patterns into
smooth waves (**Fig. 3g,h**) that had opponent directions depending on task conditions.
Specifically, completion of a trial in the instrumental task triggered ML waves (**Fig. 3i,k bottom,**
Supplementary video 3), whereas rewards in the pavlovian task promoted LM waves (**Fig. 3j,k**
top, Supplementary video 4, p<0.001 Watson-Wilson test for equality of mean directions in
two tasks, n=6 mice for instrumental task, n=8 mice for pavlovian task).

These patterns evolved dynamically with learning: Reward-waves were initially irregular
in naive animals but became progressively smooth and directional with experience in task (**Fig.**
3l, Supplementary video 5). The dynamic sculpting of the spatiotemporal patterns by training
and task demands ruled out explanations related to the intrinsic anatomy or biophysics of
dopamine axons that would constrain the array of observed activation patterns. Thus, we
conclude that dopamine waves carry behaviorally relevant decision signals and set out to
formalize their precise contribution. In particular, the continuous propagation of dopamine
across the striatum both in space and time motivated a revision of standard "temporal-
difference" models wherein a single reward-value influences learning about earlier events that
are predictive of rewards. We reasoned that these views could be expanded to include
"spatiotemporal differences" in which waves carry additional, graded information about structural
sub-circuits that are most likely to be responsible for rewards.

Figure 3: Reward delivery promotes directional waves that depend on instrumental requirement of task.

a, Schematic of test chamber. **b**, Changes in tone frequency for short, medium and long trials tiling fraction of trial complete. **c**, In the instrumental task (*left*), tone transitions are linked to rotation of the wheel, and change in tone frequency. The total distance to travel on each trial is drawn from a uniform distribution of 50-150 centimeters. In pavlovian task (*right*), the passage of time escalated tones, and the duration to wait was also drawn form a uniform distribution of 4-8 seconds. **d**, Example trials in the instrumental task. When the mouse traverses linearized distance rapidly, the tones also escalate quickly, but if the mouse pauses running, auditory tones signal the completed fraction of distance. **e**, Example licking behavior in pavlovian, sorted by delay to reward. Mice increase lickrate in anticipation of reward. **f**, These anticipatory licks were not influenced by distance to run, or duration to wait, but increased in proportion to progress to reward signaled by tones (two-way ANOVA effect of tones $F_{(8,683)} = 3.32$), $p = 0.001$ and effect of 4 duration bins $F_{(3,683)}=0.48$, $p = 0.7$ in pavlovian sessions. For instrumental sessions, effect of tones $F_{(8,359)} = 8.41$, $p<0.001$ and influence of four distance bins $F_{(3,359)} = 0.13$, $p=0.9$). **g**, Alignment of trial-by-trial wave velocity across the striatum. Rewards consistently resynchronized dopamine axons into wave in the instrumental task (n=123 trials), but also in the pavlovian task **h** (111 trials). **i,** *Top,* example flow vectors (black arrows) and source locations (contour plot representing source regions) across pixels for a single trial. *Bottom,* peak-normalized fluorescence time course across trials produced by mediolateral waves on the medio-lateral gradient of the striatum. **j,** Same format as **i** for pavlovian session. **k,** Mean flow vectors for reward epoch (0-1s post reward) for each trial in pavlovian and instrumental sessions shown in **h** and **i**. **l,** Flow trajectory of fluorescence in response to reward as mice gained experience with the task. *Top,* Naive mice had irregular responses in the first two days of reward exposure, and at *bottom*, the same mice exhibit smooth waves after 3 weeks of learning reward contingency. See supplementary video 5 for responses plotted.

Dopamine waves implement spatio-temporal credit assignment

Our functional interpretation of dopamine dynamics is that the opponent wave trajectories at reward are relevant for spatiotemporal credit assignment. The key inference the animal must make is whether it is in control of the reward-predictive tone transitions, and moreover, which specific contingency applies in the current trial (i.e., distance to run to advance tones). Thus, for mice to preferentially run in the instrumental task (and persist running for long-distance trials), the extent of instrumental controllability should guide reward-evoked dopamine to favor the DMS (i.e. strengthen action-outcome learning). Trial by-trial controllability is partly ambiguous in the task because contingencies were stochastic (drawn from uniform distributions), and mice natively run to varying levels. Nonetheless, we reasoned that task contingencies could still be inferred within trials based on the extent that tone transitions are congruent with locomotion, and dopamine signals can be informed by such congruency.

To formalize this notion, we constructed a multi-agent mixture of experts (MoE) model, extending earlier hierarchically nested corticostriatal circuit models of learning and decision making [24,25] (**Fig. 4a, Extended Data Fig. 7**, see *Methods* for details). At the highest layer (*level 1*) is an expert, putatively corresponding to DMS, that computes the online evidence for action-outcome contingencies and thus task controllability (**Fig. 4a**). Sub-experts within that area (*level 2*) represent specific contingencies (e.g., distance needed to run is short, medium or long) based on previous exposure to the tone transition distributions, learned as a semi-markov decision process via temporal difference learning[40]. Sub-expert prediction errors (PEs; level 3) occur at tone transitions and are used to compute evidence for (or against) the accuracy of each sub-expert's prediction. This formulation allows an agent to flexibly adapt behavior based on task contingencies (**Extended Data Fig. 7**) [24,25] and expands the RL account of dopamine to allow both RPE and value signals to be informed by their inferred causal contributions[41-44].

This architecture makes novel predictions at multiple levels which can potentially tie together the separable roles of dopamine during reward pursuit (performance) and learning. First, when reward waves initiate in the DMS (i.e. ML waves in the instrumental task), that region will receive the most credit, and hence mice will be faster to initiate running on the next trial and will persist in doing so until rewards are obtained. Second, the reward wave dynamics should be informed by a trace of which circuit ("expert") was most responsible for the reward (i.e., which circuit's predictions were most valuable). We posited that DA dynamics during the tone transitions (anticipatory epoch) could provide such a responsibility signal; that is, the sub-circuit that best predicts the action-outcome contingencies will exhibit increases in dopamine. These levels of dopamine can facilitate mice's motoric output to be guided most-strongly by that subexpert, while also signaling the degree to which it is responsible for future rewards. We thus hypothesized that DA dynamics during anticipation would impact how reward-waves circulate among striatal subregions and the behavioral expression of running in future trials. Finally, at the most fine-grained level, our model predicts that RPEs should occur at tone transitions to inform the extent of instrumental controllability. In the remaining sections we unpack and test each of these predictions.

The first prediction is that dopamine waves experienced at reward outcome reflects a
measure of credit assignment across the striatal experts. ML waves deliver dopamine first to
medial subregions (**Fig. 2i,j, 3i,j**), and these DMS-biased signals would selectively strengthen
corticostriatal representations for action-outcome contingencies that compete for instrumental
control in future trials. As such, we predicted that stronger ML waves at reward would enhance
instrumental learning that will drive future running. Indeed, we found a significant correlation
between the trial-by-trial magnitude of reward wave and latency to start running on the next trial
(n=6 mice, mean r = -0.32, p = 0.0019 two-sided t-test on correlation coefficients). Furthermore,
these wave magnitudes predicted the velocity even late in the next trial, 10.2 ±1.4 seconds after
the reward response (**Fig. 4c**). The influence of these waves in future-trial behavioral
adjustments indicated that they are used for learning functions. Further, these effects were
selective to instrumental sessions, indicating that DMS sourced ML waves promote learning
about instrumental contingency that is employed for future reward pursuit.

Anticipatory dopamine ramps provide eligibility for credit assignment

If reward-waves reflect credit assignment, what determines which subregion should
receive the credit? Canonical accounts in RL invoke dopamine RPEs that have graded effects
on learning depending on "eligibility" signaled by recent MSN activity[45,46]. As noted above, we
considered the possibility that local dopamine dynamics during the anticipation epoch
themselves signal a coarser measure of eligibility in terms of which subregion was responsible.
This trace would be in proportion to the value of the underlying subregion's predictions,
providing a tag for a subregion's credit.

We thus focused on the activity of dopamine axons during the anticipatory period as
mice drew closer to reward. In the instrumental task, we observed a buildup of activity in the
DMS (**Fig. 4d,f**), ramping in proportion to the progress to reward[15,47]. Strikingly, the opposite
profile was observed in the pavlovian condition (**Fig. 4e,f**), with decreasing ramps even as the
mouse continued to increase licking in anticipation of rewards. These findings do not support
extant models of DA ramps in accumbens or midbrain, where they have been linked to value
functions or RPEs[15,48-50], none of which predict opposite profiles across the two tasks.

Instead, we posited that anticipatory dopamine dynamics in the DMS reflects the
evidence of agency or controllability, and that subregions within might differentially represent
distinct controllable transition functions (which vary from trial to trial). Escalating tones in our
tasks provide information about online action-outcome contingency. For example, if tone
transitions consistently follow locomotion (as in **Fig. 3d**), they signal evidence for control. The
opposite inference can be made in the pavlovian task when tone transitions diverge from
locomotion. Respectively increasing or decreasing ramps in the instrumental and pavlovian
tasks accumulate in MoE 'distance' expert-weights as controllability is confirmed (or
contradicted) with each tone transition (**Fig. 4a, right, Extended Data Fig. 7**).

Thus, according to our model, DA ramps do not reflect a monolithic value function, but
rather the value of the underlying sub-region's agentic predictions for reward pursuit, as a
marker of that region's responsibility. Consequently, we argue here that the computational
function of dopamine waves at reward-outcome is to assign spatio-temporal credit by delivering

dopamine to striatal subregions with different latencies as a function of their graded
responsibility signals. This proposal is also motivated by theory and observations that
dopamine-mediated plasticity at striatal synapses is strongly attenuated with delayed dopamine
release[45,46].

This credit assignment interpretation makes additional testable predictions at both
physiological and behavioral levels. If dopamine ramps during reward anticipation hold
persistent information about a sub-region's prediction accuracy, they should modify the impact
of dopamine bursts at reward to focus preferentially on the sub-region with the highest
accuracy. As such, striatal areas that ramp with the steepest slopes during anticipation (highest
eligibility) should receive a reward response soonest (largest credit, **Fig. 4g**). Indeed,
anticipatory ramp slopes across pixels were significantly correlated with the fastest latency to
peak fluorescence following reward for both tasks (**Fig. 4h,i**).

Second, if DMS ramps signal responsibility for learning about instrumental control, then
trial-by-trial DMS ramp slopes should also modulate the impact of reward waves on next-trial
velocity. Indeed, we found that the impact of ML waves on future velocity in the instrumental
task (**Fig. 4c**) were dependent on the level of DMS ramps in the previous trial. When DMS
ramps were steep, reward waves strongly predicted speeded velocity in subsequent trials; this
effect was absent when ramps were weak (**Fig. 4k**, p = 0.016 Wilcoxon signed-rank test, n=6
mice). Together these results suggest that anticipatory dopamine ramps provide a tag for how
midbrain driven reward-credit circulate across the striatum to deliver a reinforcement signal for
future performance.

Figure 4

346
347
Figure 4: Anticipatory epoch dopamine dynamics reflect inferred controllability, trial-
specific task statistics, and modulate reward responses in line with a mixture of striatal
experts.
a, Schematic of hierarchical mixture of experts model as a framework for interpreting functional
relevance of dopamine dynamics applied to escalating tone tasks. At the lowest level, individual
states (representing auditory tones) induce reward prediction errors if they misalign with learned
contingencies. These prediction errors are accumulated within trials to provide evidence for or

against "sub-experts" specialized to represent local task contingencies (e.g. short, medium or long
durations/distances). At the highest level, instrumental or Pavlovian "experts" computes the
overall (weighted across sub-experts) evidence for instrumental task requirements, used to infer
controllability of the value function. These distance expert responsibility weights accumulate
within and across trials in the Instrumental task and decline in the Pavlovian task, and are used
to adjust model velocities. **b**, Proposed implementation of hierarchical task signals in striatal
dopamine activity. Widefield and 2-photon imaging at the micron level was used to test sub-
region-specific computations in dopamine terminals. **c**, Multiple regression predicting future
running speed of mice in the late phase of the next trial as a function of trial-by-trial wave
propagation-velocity (in 1 second bins) surrounding the reward from the previous trial. Reward-
induced wave velocity predicted future running speed in instrumental (blue) but not Pavlovian
(pink) sessions. Regression coefficients significantly different from zero (blue, asterix p=0.005,
two-tailed t-test). Error bars are S.E.M. **d,** Anticipatory and reward response in the medial and
lateral DS in a representative instrumental session. White points indicate start of trial. **e,** same
format as **d** for pavlovian session. **f,** Aggregate ramping profile during anticipatory epoch for the
DMS. Mean activity for each session was z-scored and averaged across mice. Activity in DMS
but not DLS showed task-dependent ramping profiles in line with inferred controllability. Shaded
regions represent S.E.M. **g,** Schematic for testing whether anticipatory epochs ramp slope is
related to the latency to peak dopamine in the outcome epoch. **h**, Results of the relationship from
two representative sessions, each from instrumental and pavlovian condition. For both tasks,
ramp slope was inversely related to subsequent latency to peak reward response. **i, j** summarize
the distribution of correlation coefficients across sessions. **k**, Multiple regression as in **c** for
instrumental sessions. In each session, trial-by-trial ramp slope was median split into low-ramp
or high-ramp trials. Across trials, steeper DMS ramps magnified the impact of reward waves on
subsequent trial running speed, in line with credit assignment. **l,** Anticipatory epoch ramps in a
sample instrumental (*top*) and pavlovian (*bottom*) sessions were expressed to varying extent on
the medio-lateral axis. ROIs were drawn with fixed distance from the edge of the field of view and
illustrated by inset. **m,** Quantification of ramp slope across sessions. Error bars represent S.E.M.
n, Local subregions within the dorsal striatum respond to distinct distance contingencies during
reward pursuit, reminiscent of sub-expert dynamics. Contingency specialization map in an
example session with color indicating the average distance that produces the steepest ramp
slopes for each pixel. **o,** Example time course of anticipatory ramps in two example subregions
for their respective preferred (high ramp trials), and non-preferred (low ramp trials). See Extended
Data Fig. 7c for similar pattern of activation in model simulations.

Thus far, we have focused on the coarsest division of labor related to the highest level in
our model (controllability, level 1), but the agent's ability to infer control depends on underlying
sub-experts that learn distinct action-outcome contingencies (level 2, **Fig. 4a,b**). Such a
hierarchical scheme implies that striatal subregions should also differentially ramp for different
distance contingencies (**Fig. 4a**). Overall, we observed that DS dopamine ramps are expressed
in a gradient across both tasks, with the strongest ramps in the most medial portions (**Fig. 4l,m**).
These results are in line with previous work on progressive instrumental specialization of DS on
the mediolateral axis[51]. Moreover, contiguous territories within the DS exhibit varying ramp
profiles for different distance conditions (**Fig. 4n, Extended Data Fig. 8**), with each area
expressing the steepest dopamine ramps in preferred set of trials with related distance
requirements (**Fig. 4o**). On a trial-by-trial basis, we further observed a significant rank
correlation between each pixel's ramp slope and latency to peak response during reward (mean
$r = -0.13$, spearman's correlation $p<0.001$ for all instrumental sessions). These results indicate
that the heterogeneously expressed anticipatory ramp gradients across the striatum modulate
the spread of reward waves, further strengthening the relationship between eligibility and credit
assignment. These findings further support our interpretation that is motivated by MoE account
by demonstrating that DMS consists of smaller sub-regions that learn, and express predictions
for a variety of potential instrumental contingencies.

These findings led us to ask whether waves organized the response of dopamine axons
on the micron scale, and functionally, how evidence for instrumental controllability accumulated
in single axon segments. The ramp-like responses we observed at the coarser scale using
widefield, one-photon imaging may emerge from trial-by-trial ramps within individual axons, or
from a weighted distribution of sharply tuned activation patterns. To directly address these
questions, we used 2-photon imaging in two mice to examine the behavior of individual axons in
the DMS (**Fig. 5a**).

Figure 5

416
417
Figure 5: Single dopamine axons show wave-like reward dynamics, tone-specific
transients, and distance-dependent ramping during instrumental anticipation.
a, Schematic of imaged region (*top*), and example field of view of dopamine axons in DMS. **b**,
Sequence of frames showing how individual dopamine axons respond to reward. Time relative to
solenoid click is shown below each frame. Note the activation of leftmost fibers first, then

progressive activation of more lateral axons. **c**, Average time course of reward response from
rectangular ROIs equally distributed along the ML axis. **d**, Activity of dopamine segments that
respond to tone transitions during anticipation. The activity in each trial is shown as percent trial
complete instead of alignment in time. *Top,* heatmap shows the trial-by-trial responses (106 total
trials) of groups of pixels in the 2-photon field of view that respond transiently at specific fraction
of trial completed. The responses of nine different types are concatenated. *Bottom,* average delta
f/F for each type. **E**, Transient response-peaks are not tuned to time or distance run within trial.
Each time series is aligned to trial-start and binned into 3 distances: short (50-80cm, dark blue),
medium (80-120cm), and long (120-150cm, light blue). Note that the peak location of response
arrives at different distances in each trial. By contrast, when aligned to % trial complete, in **f**, the
peak response arrives with fixed delay from tone transitions (illustrated at the top for both panels)
for all distance contingencies. The transient responses for each tone had larger amplitudes for
shorter trials (ie when rewards are predicted to occur sooner, in line with state-dependent reward
prediction errors within the lowest level of MoE). **g**, Individual axon segments highlighted to
demonstrate example ramp-like trajectories. **h**, Some axon segments ramped downward across
all distance trajectories when aligned to distance travelled or fraction of trial completed as in **i**. **j,**
An axon segment that progressively ramps upward only in short distance trials. **k**, same alignment
as **i**. **l,** pixelwise map of ramp slope during anticipation.

Similar to our observations at the macro scale, reward delivery recruited dopamine
axons in a spatial sequence that was directional (**Fig. 5b,c**), demonstrating that wave-like
activation patterns also organize individual axon lattices on the micron scale.

The activity in individual dopamine axons were also modulated during the anticipation
epoch. Strikingly, segments of axons transiently responded to auditory tone transition, tiling the
full sequence of escalating tones (**Fig. 5d**). The timing of these responses was not affected by
distance travelled (**Fig. 5e**), but reliably responded to changes in tone frequency across a
variety of distance contingencies (**Fig. 5f**). The systematic tuning of these axons to tone-
transitions are consistent with PEs at the lowest level of our model (**Fig. 4a, Extended Data
Fig. 7**) that are used to update the online evidence of predictions within each sub-expert. Each
tone is represented as a unique state within a sub-expert's semi-markov process, and PEs arise
at tone transitions when they misalign with the predicted distance (or dwell time) until state
change. Furthermore, the model predicts larger PEs for state transitions indicative of rewards
that will arrive when the distance to run is shorter, due to temporal discounting (**Extended Data
Fig. 7**). Supporting this prediction, we observed that tone responses were largest in trials that
had shorter distance contingencies, and progressively decreased in amplitude for longer trials
(**Fig.5e,f,** mean r = -0.33, and -0.14 n=2 mice; see **Extended Data Fig. 9**). These PE-like
responses were distributed throughout the 2-photon field of view, with equivalent fractions of
pixels selectively tuned to each tone transition (**Extended Data Fig. 9**).

We also noted that contiguous segments of axon lattices had single trial ramps that were
either upward (**Fig. 5h,i**) or downward (**Fig. 5j,k**) as mice get closer to reward. Instead of tuning
to tone-transitions reported above, these dopamine ramps were selectively expressed for
different contingencies, ramping to varying extents depending on the required distance in
separate trials. Together these results provide evidence for two, simultaneous classes of nearby
dopamine axon segments (**Extended Data Fig. 9**) used for sub-expert computations: transient
PE signals that respond to state transitions, and ramping segments that accumulate evidence
for controllability as predicted by a sub-expert.

Discussion

Our report of dopamine waves provides the earliest evidence for a foundational
organizational principle of dopamine axons that correlate activity within functionally related
striatal boundaries. In the cortex, travelling waves have been described to facilitate (or
constrain) computations that are topographically organized[52-54]. Similarly, we interpret the
computational significance of dopamine waves as orchestrating dopamine release to striatal
subregions that exhibit a graded functional specialization on the medio-lateral axis[29,51,55]. Thus,
waves are a natural candidate for solving the spatiotemporal credit assignment problem when
multiple, topographically organized striatal actors/sub-experts compete to guide action selection
across multiple levels of abstraction[24,25,56] . We used a very simple task to manipulate reward
and sensory statistics, requiring mice to resolve ambiguity about instrumental contingency by
comparing predicted and actual tone transitions. Consistent with the MoE account, wave
directions during reward were sensitive to controllability of task structure, and -- only in the

controllable task -- dopamine waves were related to future behavioral adjustment on a trial-by-
trial basis.

We also describe anticipatory epoch ramping dynamics that appear to signal the value of
a subregion's prediction about reward contingency. These dynamics may serve a dual purpose.
First, they could promote online behavioral flexibility (e.g., optimize reward-rate and minimize
energetic costs) according to the predictions of the most accurate subregions during reward
pursuit. Second, these activity patterns would also signal which subregion was most responsible
for behavioral output and hence provide a low dimensional tag for responsibility (akin to an
eligibility trace in RL[57]), which would then allow for reward-driven RPEs to preferentially credit
the appropriate subregion and the eligible MSNs within it. While the two functions are not
mutually exclusive, our data provide strong support for the second interpretation: On a trial-by-
trial basis, the degree of ramping across regions was related to the latency to reward peak
elicited by the wave, and the combination of ramp slope and wave magnitude was predictive of
subsequent-trial behavioral adjustments. These findings accord with views that dopamine
signals can have different functions during reward pursuit and outcome, which can be gated by
local microcircuit elements that regulate plasticity windows [3,58-61]. Moreover, we also interpret
transient and localized RPEs during reward pursuit as facilitating inference about the current
task state (i.e., determining credit), whereas RPEs during reward itself facilitates reinforcement
learning; a dual operation that can also be gated[43,61-63]. Put together, the synthesis of our data
and computational simulations imply that dopamine signals are spatio-temporally vectorized
during both epochs, tailored to underlying region's computational specialty.

Although the dorsal striatal dopamine dynamics support the computations of the
Distance expert in the MoE, one limitation of our study is that we did not identify or assess the
dopamine dynamics with properties of the 'Time' expert. Many studies investigating RPEs
involve classical conditioning in which temporal representations are evident in the midbrain[64-66].
Models and data have suggested that ramping signals related to timing may be present in other
regions upstream of the DA system[67-69]. Another limitation is that we did not deduce the origin
of dopamine waves, which may be inherited from sequential firing of midbrain dopamine cells
that have a topographical projection pattern[70]. To date, such dynamics have not been reported
in the literature, potentially because limited studies investigated the activity of a large population
dopamine neurons simultaneously[71]. Another likely mechanism may involve local sculpting of
dopamine release within the striatum. Wave-like patterns have been reported in neocortex[34,72]
and striatal cholinergic interneurons[73], both of which can potently regulate dopamine axon
activity[74-76]. Moreover, dopamine waves at reward outcome may also be a consequence of the
interaction between primed excitability of dopamine axons during the anticipatory epoch and
midbrain-sourced synchronous reward bursts. The combination of these two patterns may
produce sequential activation that propagates across the striatum in proportion to expressed
ramp during anticipation.

Methods

Animals and Surgery. All procedures were conducted in accordance with the guidelines of the NIH and approved by Brown University Institutional Animal Care and Use Committee. We used 17 DAT-cre mice (9 females, 8 males; *Jax Labs # 020080*) that were maintained on reversed 12hr cycle and all behavioral training and testing was performed during the dark phase. To achieve selective expression of GCaMP6f in dopamine cells, we followed standard surgical procedures for stereotaxic injection of cre-dependent virus. Briefly, mice were anesthetized with isoflurane (2% induction and maintained at 0.75-1.25% in 1 liter/min oxygen). To attain widespread infection of dopamine cells throughout the midbrain, we drilled two burr holes above the midbrain (-3.2mm AP, 0.4mm and 1.0mm ML relative to bregma) and injected 0.1-0.2μL of AAV-syn-Flex-GCaMP6f at two depths per burr hole (3.8 and 4.2 mm relative to brain surface). We next secured a metal head post to the skull and implanted an imaging cannula over the ipsilateral dorsal striatum. The cannula is a custom fabricated stainless-steel cylinder (Microgroup; 3mm diameter and 2.5-3mm height) with a 3mm coverslip (CS-3R, Warner Instruments) glued at the bottom with optical adhesive (Norad Optical #71). To insert the cannula into the brain, a 3mm diameter craniotomy was first drilled over the striatum (at bregma, centered on 2.0mm ML), and then gently removed the dura and slowly aspirated the overlying cortex until white colossal fibers were clearly visible (~0.8-1.2mm from brain surface). These fibers were also gently aspirated layer by layer until the underlying dorsal striatal tissue was uniformly exposed. A sterile imaging cannula was progressively lowered until the coverslip contacted striatal tissue uniformly. Dental cement was applied to secure implant to the skull and mice received a single dose of slow-release buprenorphine and allowed to recover for 1-2 weeks with post-operative care.

Behavioral Training. After full recovery from surgery, mice underwent 2-3 days of habituation in operant chambers outfitted with a 3D printed wheel (15 cm diameter), audio speakers and a solenoid-gated liquid reward dispenser. Following acclamation, mice were water-restricted, receiving 1ml/day in addition to water earned during task performance. We used custom LabVIEW scripts to control operant boxes during training and testing in behavioral tasks. In the first stage of training, mice received non-contingent rewards that were delivered randomly (3-15 second apart, uniform distribution) for 3 consecutive days. Next, training in a "pavlovian" task began, wherein rewards were delivered after a variable delay from trial start. The start of each trial is signaled by the onset of a 4.3kHz tone that continued to escalate in frequency in proportion to fraction of trial completed. We used nine different frequencies that were selected to minimize harmonic overlap; 4.3kHz, 6.2kHz, 8.3kHz, 10kHz, 12.4kHz, 14.1kHz, 16kHz ,8.4kHz, 20kHz. Across trials, the duration to wait for reward is randomly drawn from a uniform distribution (4-8 seconds). At the end of a trial, the auditory sound is turned off, and the solenoid delivered 3μL of water reward to a spout in front of the mouse. Licking behavior is detected using capacitive touch sensors (AT42QT1010, Sparkfun). In some catch trials, the initial 4.3kHz tone turned off after 0.5s and the mouse did not have continuous information of progress to reward. For clarity, we only focused on escalating-tone trials. The next trial started after a variable inter-trial-interval of 3-8 seconds. After 2-3 weeks of the pavlovian task, activity of dopamine axons in the striatum was imaged in a test chamber with a widefield and 2-photon

imaging system. The same animals were then further trained on a distance-variant of the same
task, where reward delivery is now contingent on mice running on the wheel. Mice were
exposed to the "instrumental" task in training chambers requiring them to run on the wheel to
traverse linearized distances, also randomly selected from a uniform distribution (50-150cm).
Progress to reward was indicated by the same tone frequencies, and the angular position of the
wheel recorded using a miniature rotary encoder (MA3A10250N, US Digital). All behavioral data
is digitized and stored to disc at 50Hz.
571
Widefield and two-photon imaging. All imaging was performed using a multi-photon
microscope with modular laser-scanning and light microscopy designed by Bruker/Prairie
Technologies. For widefield imaging, we used a full-spectrum LED illumination with FITC filter
cassette for illumination at 470nm and detection centered at 530nm. Images were acquired
using a CoolSnap ES2 CCD camera (global shutter, Photometrics) and synchronized with
behavioral events through TTL triggers. All widefield images during behavioral tasks were
acquired with a 4X objective (Olympus), 100ms exposure (10Hz) and 8X on-camera binning to
achieve a sample resolution of 40µm/pixel (unless indicated otherwise). Two-photon microscopy
was performed using a 20X air objective (Olympus) on the same imaging platform with a
femtosecond pulsed TiSapphire laser source (MaiTai DeepSee, 980nm power measured at
objective was 20-50mW) that was scanned across the sample using a resonant (x-axis) and
non-resonant (y-axis) galvanometer scanning mirrors. Returning photons were collected through
an imaging path onto milti-alkali PMTs (R3896, Hamamatsu), and recorded frames were online-
averaged to achieve a sampling rate of 10-15Hz.
586
Data Analysis and statistics. All images were processed with custom routines in MATLAB.
Each session is preprocessed for image registration, and alignment to behavioral events based
on triggers. Movement artifacts and image drift in the XY plane were corrected using rigid-body
registration using a DFT-based method[77]. To cluster the activity of dopamine axons, we used
the K-means algorithm in MATLAB. To compute robustness of clustering results, we used the
adjusted rand Index measure which computes the similarity of two clusters based on the
probability of member overlap (corrected for chance; 0=random clusters, 1=exact same
membership). To examine how robust the clustering results were, we re-clustered the same
dataset 100 times in K-means using random initialization and varied cluster limits. We compared
the extent that pixels were re-clustered into the same group using the adjusted rand index as an
indicator of robustness of underlying structure of the data that produced clusters (see Extended
Data Fig. 4. To additionally test the extent spatial relationship between the pixels, or the how
similarity in temporal activation influenced the identified clusters, we repeated the same analysis
but shuffled the spatial or temporal relationships between the pixels. To estimate the optimal
number of clusters within each dataset, we computed the Bayesian information criterion (BIC)
on the K-means algorithm.
We characterized flow patterns in dopamine waves by adapting standard optical flow
algorithms in machine vision that are adapted for imaging of fluorescence signals[34,36,78]. Briefly,
flow trajectories were computed for any two successive frames as a displacement of intensity
across the pixels in time. This method allows us to evaluate a pixel-by-pixel velocity vector fields
that summarizes the direction and strength of flow at each pixel. While there are multiple

methods to achieve this calculation[36,78], we adapted a combined Global-Local (CGL)
algorithm[79,80] that combines the Lucas-Kanade and Horn-Schunck methods. The frame-by-
frame vector fields calculated using the CGL method was further processed to extract sink and
source locations and also flow trajectories across multiple frames (**Fig. 2a, bottom**). The frame-
by-frame flow magnitude for each frame (or flow-velocity, with units of mm/second) is computed
by averaging the length of vectors at each pixel (e.g. **Fig. 2b, red**). The locations of sinks or
sources were estimated based on local vector orientations: i.e sinks are points of inward flow,
whereas sources are points of outward flow. We estimated the pixel-wise likelihood of sinks and
sources by simply computing the divergence of the vector field in each frame ("*divergence*"
function in MATLAB, **Supplementary Video 2**). The flow trajectory across frames were
calculated from vector fields using the "*stream3*" function in MATLAB from seeded pixels (e.g.
Fig3l).
For alignment of fluorescence time series, DMS and DLS masks were defined using one
of three methods: i) manual drawing, ii) boundaries using cluster boundaries (as in Fig. 1i, top)
and iii) uniformly spaced ROIs on the mediolateral axis (as in Fig. 4l, inset). Each animal
performed multiple behavioral sessions, and we used one session per animal (n=6 mice in
instrumental task, and n=8 mice in pavlovian task) that had the largest Δf/F deviations to avoid
results from being dominated by a few animals.
To determine the influence of reward-wave on behavioral performance on the next trial,
we performed a multiple regression predicting the running velocity of mice late (i.e. 75-100% of
trial complete) in the next trial based on reward aligned wave magnitude (1-sec bins, **Fig. 4c**).
To determine whether DMS ramp slopes influenced how last-trial wave outcome, on the next
trial, we conditioned this analysis on the ramping profile in the DMS, median split into low, and
high ramp conditions (**Fig 4k**). We evaluated the correlation between the ramp slope and
latency to peak by first peak-normalizing the reward response in 2-sec window and finding the
time index (after reward) for which the fluorescence signal reached peak levels. To examine
whether anticipatory dopamine ramps had a preference for different distance conditions (**Fig.
4n,** also see **Extended Data Fig. 7**), we sorted the trials based on the expressed ramps in each
pixel and averaged the distance contingency in trials with top 90% ramping.
TIFF stacks of 2-photon images of dopamine axon segments were also pre-processed
for registration and alignment with behavioral data. To draw ROIs of these segments for
assessing organization of responses (**Extended Data Fig. 1**), we followed the Howe and
Dombeck [28]. Otherwise, we generally used pixel-wise analyses.
641
Computational model. We modeled mouse behavior using a mixture of experts / multi-agent
RL architecture[25], extended here to accommodate the sequential tone structure with semi-
markov dynamics[40]. We modeled the two task structures as separate "experts" that learned a
value function *V* as a function of either elapsed time as in classical temporal difference learning
applied to Pavlovian condition, or as a function of distance travelled. Because mice were trained
on both time and distance tasks, multiple sub-experts (representing clusters in mediolateral
coordinates of striatum) were pre-trained for 2000 trials to span a range of contingencies (e.g.,
400ms, 600ms, or 800ms per tone transition; or 5, 10 or 15cm). For simplicity, we modeled the
task with discrete sub-experts that specialized on (had been preferentially exposed to) particular
times/distances. However, one can easily generalize the framework to the continuous case

(e.g., using basis functions[81]) and the discrete space can be modeled with arbitrary resolution
by simply increasing the number of sub-experts. Moreover, various models have shown that
prediction errors can be used to segregate learning of different latent task states[43,56].
Subexpert and expert learning. The value function for each time sub-expert s estimates
the discounted future reward $V^s(X_{i,t}) = r(t) + \gamma V^s(X_{i,t+1})$ and was trained via temporal
differences[82] based on reward prediction errors $\delta(X_{i,t}) = r(t) + \gamma V(X_{i,t+1}) - V(X_{i,t})$. Each auditory
tone was modeled as a distinct state $X_{i,t}$ or $X_{i,d}$ with semi-markov dynamics. That is, the onset of
each tone i would advance the state vector to the corresponding position even if the tone
occurred earlier or later in absolute time/distance. Thus the value function learned for each sub-
expert was tied to the current state (tone) and the (discretized) dwell time (t) or distance (d)
since it has been entered, and not to the absolute time or distance that passed from the onset of
the first state. This semi-markov process was based on the assumption that the tone stimuli
induce a neural state representation upon which TD is computed[40,81] and evidence that rodents
are endowed with such a rich state representation [83.] The value function was learned by
adjusting weights in response to the X state vector, with $V(X_{i,t}) = w_t X_{i,t}$ and $w_t \leftarrow w_t + \alpha \delta(t)$,
where α is a learning rate. The distance experts were trained analogously but with the X vector
advancing with discretized distance steps taken rather than passive time. Thus if the agent
stopped moving, the $X_{i,d}$ vector remained constant until it moved again, and if it moved faster
than usual, the $X_{i,d}$ vector would advance to later states accordingly. We fixed α=0.25 and γ
=.95 for all experts but verified that the patterns were robust to other settings.
Performance and inference. After learning, the on-line evidence (responsibilities, fig 4B,
modeling the ramps) for each sub-expert was computed as an approximation to the likelihood of
the trial-wise tone transitions for that sub-expert given the dwelling time or distance since the
last tone. We adopted a hybrid Bayesian-RL formulation[25]. Each expert learns a value function
associated with the tones, and inference about which expert is responsible is computed in
proportion to the log likelihood ratio of the observations (tone transitions at particular instances
following elapsed time or distance) given their predictions relative to the other experts.
From a Bayesian perspective, the attentional weights for each expert can be
evaluated by computing the posterior probability that each expert encompasses the best
account of the observed data x: $P(s|x) = P(x|s) P(s) / \Sigma P(x|s_i) P(s_i)$. Thus the evidence for each
expert is computed by considering its prior evidence and the likelihood that the observed tone
transitions or rewards (positive or negative) would have been observed under the expert's
model, relative to all other experts. For example, if there was a low probability for a tone
transition at a particular moment under a given expert, then the likelihood of that observation
given the expert's model is low. Once the posterior evidence for each expert is computed, one
can then apply Bayesian model averaging to allocate attentional weights to each expert in
proportion to their log evidence.
Rather than a fully Bayesian realization, we instead implemented an RL approximation
that we posited would more directly relate to corticostriatal DA mechanisms[25]. Instead of
computing the likelihood directly, each expert was penalized as a function of its reward
prediction errors. In particular we updated the evidence for each sub-expert's predictions in
terms of a responsibility weight $\omega`_s$ which was decremented when the corresponding sub-expert
experienced a reward prediction error: $\omega`_s \leftarrow \omega`_s - \delta_s$, where δ_s is the positive reward
prediction error according to the corresponding sub-expert's value function at the state vector X.

(Similar results hold if using $|\delta_s|$ instead of only positive RPEs to decrement expert weights, but
positive RPEs dominate). Intuitively, experts with more prediction errors are less likely to have
been responsible for the outcome (tone transition or reward). These responsibility weights were
then normalized relative to all sub-experts as an approximation to the log evidence for a given
subexpert: $\omega_{si} = exp(\beta\omega^c si) / \sum_j exp(\beta\omega^c sj)$, where β is an inverse temperature parameter.
Thus, in contrast to standard RL in which RPEs reinforce actions that yield rewards, during
inference, more frequent RPEs for a given subexpert are indicative that it is less responsible for
observations compared to those that predict these observations. Such a scheme is compatible
with extant models that use reward prediction errors for state creation and inference separate
from reinforcement per se [25,43,56,63]. We posited that these RPEs correspond to the phasic
events observed at tone transitions in the 2p imaging data. The accumulation of these
responsibility weights were posited to relate to the 1p imaging data in discrete sub-regions of
DMS.
709   Finally, a second-level task selection process was implemented to arbitrate responsibility
between the overall distance expert and overall time expert (each of which constituted a
weighted combination of their subordinate experts). This inference process was identical to that
for the sub-experts, with responsibility updated based on their experienced prediction errors: $\omega`_D$
$\leftarrow \omega`_D - \delta_D$, where $\omega`_D$ is the accumulated responsibility of the distance expert based on its
reward prediction errors, $\delta_D = r(t) + \gamma V_D(t+1) - V_D(t)$. The value function for the distance and
time experts V_D and V_T are in turn weighted averages according to the inferred responsibilities of
the subordinate experts within each structure: $V_D(t) = \Sigma\omega_{sD}V_{SD}(t)$ and $V_T = \Sigma\omega_{sT}V_{ST}(t)$. Similarly,
the value function of the agent as a whole is the weighted average value function across the two
experts $V(t) = \omega_D V_D(t) + \omega_T V_T(t)$. These responsibility weights for each task structure were
again normalized across tasks, $\omega_D = e^{\beta w`D} / e^{\beta w`D} + e^{\beta w`T}$.
720   For each distance or time, 100 test trials were run with 10 tones each and an inter trial
interval was randomly drawn from 5-15s. The agent as a whole selects actions in terms of
speeds to run for a period of time at each tone transition or after it has completed it's previous
running. Speeds were selected in proportion to the inferred responsibility of the DMS expert,
together with some stochasticity: speed(t) = $5*(\omega_D(t)-0.5)) + \epsilon$, where ϵ was drawn from a
uniform distribution with a mean of 3. Stochasticity facilitates the agent ability to disambiguate
distance from time tasks within a trial (a constant speed would equate the prediction errors for
the two tasks given appropriate sub-experts). Increasing speed with inferred DMS expert
responsibility ω_D allows the model to capture the increased running with instrumental task
structure (extended Fig 7). More detailed investigation of how speeds may be optimized
according to reward/effort/delay tradeoffs will be examined in future work.

References

1   Schultz, W. Dopamine reward prediction-error signalling: a two-component response.
Nat Rev Neurosci **17**, 183-195, doi:10.1038/nrn.2015.26 (2016).
2   Berridge, K. C. The debate over dopamine's role in reward: the case for incentive
salience. *Psychopharmacology (Berl)* **191**, 391-431, doi:10.1007/s00213-006-0578-x
(2007).
3   Berke, J. D. What does dopamine mean? *Nat Neurosci* **21**, 787-793,
doi:10.1038/s41593-018-0152-y (2018).
4   Schultz, W., Dayan, P. & Montague, P. R. A neural substrate of prediction and reward.
Science **275**, 1593-1599, doi:10.1126/science.275.5306.1593 (1997).
5   Prensa, L. & Parent, A. The nigrostriatal pathway in the rat: a single-axon study of the
relationship between dorsal and ventral tier nigral neurons and the striosome/matrix
striatal compartments. *Journal of Neuroscience* **21**, 7247-7260 (2001).
6   Matsuda, W. *et al.* Single nigrostriatal dopaminergic neurons form widely spread and
highly dense axonal arborizations in the neostriatum. *J Neurosci* **29**, 444-453,
doi:10.1523/JNEUROSCI.4029-08.2009 (2009).
7   Hyland, B. I., Reynolds, J. N., Hay, J., Perk, C. G. & Miller, R. Firing modes of midbrain
dopamine cells in the freely moving rat. *Neuroscience* **114**, 475-492 (2002).
8   Li, W., Doyon, W. M. & Dani, J. A. Acute in vivo nicotine administration enhances
synchrony among dopamine neurons. *Biochemical pharmacology* **82**, 977-983 (2011).
9   Joshua, M. *et al.* Synchronization of midbrain dopaminergic neurons is enhanced by
rewarding events. *Neuron* **62**, 695-704, doi:10.1016/j.neuron.2009.04.026 (2009).
10   Kim, Y., Wood, J. & Moghaddam, B. Coordinated activity of ventral tegmental neurons
adapts to appetitive and aversive learning. *PLoS One* **7**, e29766,
doi:10.1371/journal.pone.0029766 (2012).
11   Mohebi, A. *et al.* Dissociable dopamine dynamics for learning and motivation. *Nature*
570, 65-70, doi:10.1038/s41586-019-1235-y (2019).
12   Eshel, N., Tian, J., Bukwich, M. & Uchida, N. Dopamine neurons share common
response function for reward prediction error. *Nat Neurosci* **19**, 479-486,
doi:10.1038/nn.4239 (2016).
13   Schultz, W. Predictive reward signal of dopamine neurons. *J Neurophysiol* **80**, 1-27,
doi:10.1152/jn.1998.80.1.1 (1998).
14   Glimcher, P. W. Understanding dopamine and reinforcement learning: the dopamine
reward prediction error hypothesis. *Proc Natl Acad Sci U S A* **108 Suppl 3**, 15647-
15654, doi:10.1073/pnas.1014269108 (2011).
15   Hamid, A. A. *et al.* Mesolimbic dopamine signals the value of work. *Nat Neurosci* **19**,
117-126, doi:10.1038/nn.4173 (2016).
16   Haber, S. N. The primate basal ganglia: parallel and integrative networks. *J Chem*
Neuroanat **26**, 317-330, doi:10.1016/j.jchemneu.2003.10.003 (2003).
17   Graybiel, A. M. Habits, rituals, and the evaluative brain. *Annu Rev Neurosci* **31**, 359-387,
doi:10.1146/annurev.neuro.29.051605.112851 (2008).
18   Hunnicutt, B. J. *et al.* A comprehensive excitatory input map of the striatum reveals novel
functional organization. *Elife* **5**, e19103 (2016).
19   Hintiryan, H. *et al.* The mouse cortico-striatal projectome. *Nature neuroscience* **19**, 1100
(2016).
20   Riedel, A. *et al.* Principles of rat subcortical forebrain organization: a study using
histological techniques and multiple fluorescence labeling. *Journal of chemical*
neuroanatomy **23**, 75-104 (2002).
21   Brown, H. D., McCutcheon, J. E., Cone, J. J., Ragozzino, M. E. & Roitman, M. F.
Primary food reward and reward-predictive stimuli evoke different patterns of phasic

dopamine signaling throughout the striatum. *Eur J Neurosci* **34**, 1997-2006,
doi:10.1111/j.1460-9568.2011.07914.x (2011).
22 Shnitko, T. A. & Robinson, D. L. Regional variation in phasic dopamine release during
alcohol and sucrose self-administration in rats. *ACS Chem Neurosci* **6**, 147-154,
doi:10.1021/cn500251j (2015).
23 Menegas, W., Babayan, B. M., Uchida, N. & Watabe-Uchida, M. Opposite initialization to
novel cues in dopamine signaling in ventral and posterior striatum in mice. *Elife* **6**,
e21886 (2017).
24 Doya, K., Samejima, K., Katagiri, K.-i. & Kawato, M. Multiple model-based reinforcement
learning. *Neural computation* **14**, 1347-1369 (2002).
25 Frank, M. J. & Badre, D. Mechanisms of hierarchical reinforcement learning in
corticostriatal circuits 1: computational analysis. *Cerebral cortex* **22**, 509-526 (2011).
26 Gershman, S. J., Pesaran, B. & Daw, N. D. Human reinforcement learning subdivides
structured action spaces by learning effector-specific values. *J Neurosci* **29**, 13524-
13531, doi:10.1523/JNEUROSCI.2469-09.2009 (2009).
27 Badre, D. & Frank, M. J. Mechanisms of hierarchical reinforcement learning in cortico–
striatal circuits 2: Evidence from fMRI. *Cerebral cortex* **22**, 527-536 (2011).
28 Howe, M. & Dombeck, D. A. Rapid signalling in distinct dopaminergic axons during
locomotion and reward. *Nature* **535**, 505 (2016).
29 Klaus, A. *et al.* The spatiotemporal organization of the striatum encodes action space.
Neuron **95**, 1171-1180. e1177 (2017).
30 Balleine, B. W., Delgado, M. R. & Hikosaka, O. The role of the dorsal striatum in reward
and decision-making. *Journal of Neuroscience* **27**, 8161-8165 (2007).
31 Yin, H. H. & Knowlton, B. J. The role of the basal ganglia in habit formation. *Nature
Reviews Neuroscience* **7**, 464 (2006).
32 Grinvald, A., Lieke, E. E., Frostig, R. D. & Hildesheim, R. Cortical point-spread function
and long-range lateral interactions revealed by real-time optical imaging of macaque
monkey primary visual cortex. *Journal of Neuroscience* **14**, 2545-2568 (1994).
33 Lubenov, E. V. & Siapas, A. G. Hippocampal theta oscillations are travelling waves.
Nature **459**, 534 (2009).
34 Mohajerani, M. H. *et al.* Spontaneous cortical activity alternates between motifs defined
by regional axonal projections. *Nature neuroscience* **16**, 1426 (2013).
35 Muller, L., Chavane, F., Reynolds, J. & Sejnowski, T. J. Cortical travelling waves:
mechanisms and computational principles. *Nature Reviews Neuroscience* **19**, 255
(2018).
36 Afrashteh, N., Inayat, S., Mohsenvand, M. & Mohajerani, M. H. Optical-flow analysis
toolbox for characterization of spatiotemporal dynamics in mesoscale optical imaging of
brain activity. *NeuroImage* **153**, 58-74 (2017).
37 Corbit, L. H. & Janak, P. H. Posterior dorsomedial striatum is critical for both selective
instrumental and Pavlovian reward learning. *Eur J Neurosci* **31**, 1312-1321,
doi:10.1111/j.1460-9568.2010.07153.x (2010).
38 Balleine, B. W. & O'doherty, J. P. Human and rodent homologies in action control:
corticostriatal determinants of goal-directed and habitual action.
Neuropsychopharmacology **35**, 48 (2010).
39 Wunderlich, K., Smittenaar, P. & Dolan, R. J. Dopamine enhances model-based over
model-free choice behavior. *Neuron* **75**, 418-424 (2012).
40 Daw, N. D., Courville, A. C. & Touretzky, D. S. Representation and timing in theories of
the dopamine system. *Neural computation* **18**, 1637-1677 (2006).
41 Chang, Y.-H., Ho, T. & Kaelbling, L. P. in *Advances in neural information processing
systems.* 807-814.

42   O'Reilly, R. C. & Frank, M. J. Making working memory work: a computational model of
learning in the prefrontal cortex and basal ganglia. *Neural computation* **18**, 283-328
(2006).
43   Gershman, S. J., Norman, K. A. & Niv, Y. Discovering latent causes in reinforcement
learning. *Current Opinion in Behavioral Sciences* **5**, 43-50 (2015).
44   Russell, S. J. & Zimdars, A. in *Proceedings of the 20th International Conference on*
Machine Learning (ICML-03).   656-663.
45   Yagishita, S. *et al.* A critical time window for dopamine actions on the structural plasticity
of dendritic spines. *Science* **345**, 1616-1620, doi:10.1126/science.1255514 (2014).
46   Shindou, T., Shindou, M., Watanabe, S. & Wickens, J. A silent eligibility trace enables
dopamine-dependent synaptic plasticity for reinforcement learning in the mouse
striatum. *European Journal of Neuroscience* **49**, 726-736 (2019).
47   Howe, M. W., Tierney, P. L., Sandberg, S. G., Phillips, P. E. & Graybiel, A. M. Prolonged
dopamine signalling in striatum signals proximity and value of distant rewards. *Nature*
500, 575-579, doi:10.1038/nature12475 (2013).
48   Lloyd, K. & Dayan, P. Tamping Ramping: Algorithmic, Implementational, and
Computational Explanations of Phasic Dopamine Signals in the Accumbens. *PLoS*
Comput Biol **11**, e1004622, doi:10.1371/journal.pcbi.1004622 (2015).
49   Gershman, S. J. Dopamine ramps are a consequence of reward prediction errors.
Neural Comput **26**, 467-471, doi:10.1162/NECO_a_00559 (2014).
50   Morita, K. & Kato, A. Striatal dopamine ramping may indicate flexible reinforcement
learning with forgetting in the cortico-basal ganglia circuits. *Frontiers in neural circuits* **8**,
36 (2014).
51   Thorn, C. A., Atallah, H., Howe, M. & Graybiel, A. M. Differential dynamics of activity
changes in dorsolateral and dorsomedial striatal loops during learning. *Neuron* **66**, 781-
795 (2010).
52   Muller, L., Reynaud, A., Chavane, F. & Destexhe, A. The stimulus-evoked population
response in visual cortex of awake monkey is a propagating wave. *Nature*
communications **5**, 3675 (2014).
53   Nauhaus, I., Busse, L., Carandini, M. & Ringach, D. L. Stimulus contrast modulates
functional connectivity in visual cortex. *Nature neuroscience* **12**, 70 (2009).
54   Bringuier, V., Chavane, F., Glaeser, L. & Frégnac, Y. Horizontal propagation of visual
activity in the synaptic integration field of area 17 neurons. *Science* **283**, 695-699 (1999).
55   Barbera, G. *et al.* Spatially compact neural clusters in the dorsal striatum encode
locomotion relevant information. *Neuron* **92**, 202-213 (2016).
56   Collins, A. G. & Frank, M. J. Cognitive control over learning: Creating, clustering, and
generalizing task-set structure. *Psychological review* **120**, 190 (2013).
57   Singh, S. P. & Sutton, R. S. Reinforcement learning with replacing eligibility traces.
Machine learning **22**, 123-158 (1996).
58   Morris, G., Arkadir, D., Nevet, A., Vaadia, E. & Bergman, H. Coincident but distinct
messages of midbrain dopamine and striatal tonically active neurons. *Neuron* **43**, 133-
143, doi:10.1016/j.neuron.2004.06.012 (2004).
59   Threlfell, S. & Cragg, S. J. Dopamine signaling in dorsal versus ventral striatum: the
dynamic role of cholinergic interneurons. *Front Syst Neurosci* **5**, 11,
doi:10.3389/fnsys.2011.00011 (2011).
60   Bradfield, L. A., Bertran-Gonzalez, J., Chieng, B. & Balleine, B. W. The thalamostriatal
pathway and cholinergic control of goal-directed action: interlacing new with existing
learning in the striatum. *Neuron* **79**, 153-166 (2013).
61   Franklin, N. T. & Frank, M. J. A cholinergic feedback circuit to regulate striatal population
uncertainty and optimize reinforcement learning. *Elife* **4**, e12029 (2015).

62   Schoenbaum, G., Stalnaker, T. A. & Niv, Y. How did the chicken cross the road? With her striatal cholinergic interneurons, of course. *Neuron* **79**, 3-6 (2013).

63   Redish, A. D., Jensen, S., Johnson, A. & Kurth-Nelson, Z. Reconciling reinforcement learning models with behavioral extinction and renewal: implications for addiction, relapse, and problem gambling. *Psychological review* **114**, 784 (2007).

64   Pan, W.-X., Schmidt, R., Wickens, J. R. & Hyland, B. I. Dopamine cells respond to predicted events during classical conditioning: evidence for eligibility traces in the reward-learning network. *Journal of Neuroscience* **25**, 6235-6242 (2005).

65   Hollerman, J. R. & Schultz, W. Dopamine neurons report an error in the temporal prediction of reward during learning. *Nature neuroscience* **1**, 304 (1998).

66   Soares, S., Atallah, B. V. & Paton, J. J. Midbrain dopamine neurons control judgment of time. *Science* **354**, 1273-1277 (2016).

67   Brown, J., Bullock, D. & Grossberg, S. How the basal ganglia use parallel excitatory and inhibitory learning pathways to selectively respond to unexpected rewarding cues. *Journal of Neuroscience* **19**, 10502-10511 (1999).

68   Hazy, T. E., Frank, M. J. & O'Reilly, R. C. Neural mechanisms of acquired phasic dopamine responses in learning. *Neuroscience & Biobehavioral Reviews* **34**, 701-720 (2010).

69   Mello, G. B., Soares, S. & Paton, J. J. A scalable population code for time in the striatum. *Current Biology* **25**, 1113-1122 (2015).

70   Lerner, T. N. *et al.* Intact-Brain Analyses Reveal Distinct Information Carried by SNc Dopamine Subcircuits. *Cell* **162**, 635-647, doi:10.1016/j.cell.2015.07.014 (2015).

71   Engelhard, B. *et al.* Specialized coding of sensory, motor and cognitive variables in VTA dopamine neurons. *Nature*, 1 (2019).

72   Kasanetz, F., Riquelme, L. A., Della-Maggiore, V., O'Donnell, P. & Murer, M. G. Functional integration across a gradient of corticostriatal channels controls UP state transitions in the dorsal striatum. *Proceedings of the National Academy of Sciences* **105**, 8124-8129 (2008).

73   Rehani, R. *et al.* Activity patterns in the neuropil of striatal cholinergic interneurons in freely moving mice represent their collective spiking dynamics. *Eneuro* **6** (2019).

74   Krebs, M. *et al.* Glutamatergic control of dopamine release in the rat striatum: evidence for presynaptic N-methyl-D-aspartate receptors on dopaminergic nerve terminals. *Journal of neurochemistry* **56**, 81-85 (1991).

75   Threlfell, S. *et al.* Striatal dopamine release is triggered by synchronized activity in cholinergic interneurons. *Neuron* **75**, 58-64, doi:10.1016/j.neuron.2012.04.038 (2012).

76   Cachope, R. *et al.* Selective activation of cholinergic interneurons enhances accumbal phasic dopamine release: setting the tone for reward processing. *Cell Rep* **2**, 33-41, doi:10.1016/j.celrep.2012.05.011 (2012).

77   Guizar-Sicairos, M., Thurman, S. T. & Fienup, J. R. Efficient subpixel image registration algorithms. *Optics letters* **33**, 156-158 (2008).

78   Townsend, R. G. & Gong, P. Detection and analysis of spatiotemporal patterns in brain activity. *PLoS computational biology* **14**, e1006643 (2018).

79   Bruhn, A., Weickert, J. & Schnörr, C. in *Joint Pattern Recognition Symposium.* 454-462 (Springer).

80   Liu, C. *Beyond pixels: exploring new representations and applications for motion analysis*, Massachusetts Institute of Technology, (2009).

81   Ludvig, E. A., Sutton, R. S. & Kehoe, E. J. Stimulus representation and the timing of reward-prediction errors in models of the dopamine system. *Neural computation* **20**, 3034-3054 (2008).

82   Sutton, R. S. & Barto, A. G. *Reinforcement learning : an introduction*. illustrated edn, (MIT Press, 1998).

83   Gardner, M. P., Schoenbaum, G. & Gershman, S. J. Rethinking dopamine as
933   generalized prediction error. *Proceedings of the Royal Society B* **285**, 20181645 (2018).

Acknowledgements. We thank Matthew Nassar, Joshua Berke, Peter Dayan, Theresa Desrochers, for valuable discussion of the project and feedback on an earlier version of the manuscript, and members of the Frank and Moore laboratories for feedback at various stages of the project. We also thank Ines Belghiti and Aneri Soni for help with adapting the MoE model implementation to the tone task. GCaMP6 was developed and generously made available by the HHMI Janelia Farm Research Campus GENIE Project. This work was supported by HHMI Hanna Gray Fellowship to AAH, R01MH080066 and NSF grant 1460604 to MJF, and awards from Carney Institute for Brain Sciences and Dean's office to CIM.

Contributions. AAH designed and performed the experiments, analyzed the data, and applied computational model and wrote the paper. MJF designed and supervised the study, developed computational model, and contributed to writing and revising of the paper. CIM designed and supervised the study and revised the paper.

Competing Interests: The authors declare no competing interests.

Data and code availability. All data and code is available from corresponding author(s) upon reasonable request.

Materials and Correspondence: Correspondence and material request is to be addressed to Arif Hamid (Arif_Hamid@brown.edu), Christopher Moore (Christopher_Moore@brown.edu) or Michael Frank (Michael_Frank@brown.edu).

Extended Data Figure 1

<u>Extended Data Fig. 1:</u> Individual dopamine axons also exhibit decorrelated activity patterns.

a, Example frames illustrating that different portions of axon laticies are activated asynchronously. **b,** Representative timeseries of fluorescence from two axon segments outlined in blue and red at **a**. **c,** Additional examples of activity in dopamine axon segments. Data is organized such that rearby axons are plotted closer. **d,** Quantification of correlation between the sessionwide timesereries of axons based anatomical distances. Note that nearby axons are highly correlated, but they exhibit a distance dependent falloff as reported in Fig. 1g, although a different anatomical scale.

Extended Data Figure 2

Extended Data Fig. 2: Dopamine axon activity clusters into hierarchical domains.

Data from one session, mean projection of fluorescence is displayed at the top left.
Progressively increasing cluster limits identifies contiguous striatal subregions that decompose
into sub-clusters.

Extended Data Figure 3

969 **Extended Data Fig. 3: Clustering patterns in all 8 mice.**

970 We provide the K-means cluster of each of the animals examined. Plotting format follows panels
971 in Fig 1.

Extended Data Figure 4

Extended Data Fig. 4: **Cluster patterns are robust.**

a, Adjusted rand-index score of cluster patterns determined using the K-means for re-clustering (black) or shuffling the temporal (red) or spatial(blue) indices of pixels. Results are shown for 100 reclustering iternation with or without shuffling. Note that randomizing the temporal or spatial relationships of fluorescence activity results in random clusters. **b,** BIC score for K-means results all sessions examined (n=31 sessions, 8 mice; gray), and average (purple). **c,** Distribution of optimal number of clusters identified using the BIC metric. **d,** Distribution of areas of identified clusters across all mice.

Extended Data Fig. 5: **Local sources and sinks initiate and terminate dopamine activity,**
delivering temporally delayed dopamine to striatal subregions.

 a, Flow pattern (*top*) amd divergence map (*bottom*) for sinks that are clustered in medial, lateral
or central striatal regions. **b,** same format as **a**, for source locations. **c,** Time Course of activity
across the mediolateral gradient for a one minute recording epoch. Blue boxes focus on
transient events that were produced by ML, LM, or CO waves that deliver dopamine to different
parts of the dorsal striatum with relative lags.

Extended Data Fig. 6: The running behavior of mice is more structured and goal-directed in the instrumental task.

a, Example velocity profile for an instrumental session. *Top* shows the trial by trial velocity aligned to the end of trial (reward receipt) White dots indicate the start of the trial. *Bottom* illustrates mean velocity trajectory for different distance contingencies. **b,** Same format as **a** but for pavlovian session. Note that the running behavior of the mouse is disorganized relative to task events (quantified in later sessions). **c,** Example session showing variability the latency to start running on the next trial. **d,** We quantified this latency for training sessions across all mice and observed a significantly shorter latency to initiate next trial running. X-axis is displayed in log scale. **e,** Single trial trajectories of position from trial start during instrumental sessions, and pavlovian sessions in **f**. Circles denote mouse position at the end of a trial. **g,** Overall, mice ran less distance that the requirement in instrumental sessions (i.e. 50-150cm) , and **h**, mice chose not to run at all in a significant fraction of trials during the pavlovian task (note that running is required for rewards in instrumental sessions).

1007

Extended Data Fig. 7: Within-trial dynamics of model variables at all three levels under different task conditions.

a,b Positive and negative accumulation of distance expert (level-1, equivalent to DMS) weights under (a) instrumental and (b) Pavlovian task condition, for short, medium and long trial types. Each trace is the average dynamics on the very first trial, averaged for 10 simulation sessions. Similar dynamics accumulate across trials within a session when the task is repeated (not

shown). Note that the ramp shape is convex in the first trial but concave for later trials. **c,** Within
the distance expert, sub-experts (level-2) specialize on distinct contingencies and the weights
ramp accordingly depending on task conditions. **d,** RPEs within a sub-expert in which tone
transitions occur at unexpected times/distances (RPEs are zero for sub-experts that perfectly
predict the current contingency; not shown). Note the larger magnitude RPEs for short
compared to longer trials, as seen empirically (Fig. 5). Escalation of RPEs across the trial is due
to temporal discounting. Similar to the empirical data, the impact of larger RPEs on short
distances is more evident later in the trial. **e)** Model velocities (averaged across simulations)
recapitulate increase in running in instrumental compared to pavlovian sessiona. The model
selects speeds in proportion to inferred responsibility of the distance ("DMS") expert, which
accumulates across a session.

Extended Data Figure 8

1025

1026 **Extended Data Fig. 8:** **Additional example session of striatal subregions that have**
1027 **preferred distance contingency.**

1028 The trial-by-trial ramp slope during anticipation epoch was distinctly modulated for different
1029 striatal subregions. Color maps are in same format as Fig. 4n.

Extended Data Fig. 9: Discrete and ramp-like responses in dopamine axon segments.

a, Discrete, tone responses in two mice. Format is as in Fig 5. **b,** Example ramping patterns in the two animals. **c,** Examination of the anatomical distribution of pixels that exhibit tone-transition tuning. *Top* row summarizes data from mouse-1, and *bottom* is for the second mouse. Leftmost panels show the mean projection of field of view, and the next three panels show the individual pixels that display discrete responses early (first transition), mid-trial (5th transition) and late-trial (last-tone transition) responses. Moreover, the anatomical organization of these pixels are intermingled. Fifth panel shows the anatomical position of all phasically responding pixels color coded for which transition they respond to. Finally, Last panel on the right shows the anatomical distribution and trial-averaged ramp slopes of pixels within the 2-photon field of view that exhibit sustained upward or downward activity during the anticipatory epoch. Same format for mouse-2 at *bottom* **d,** Quantification of how peak response at tone-transition is affected by distance to needed to run on current trial. Our simulations predict (see Extended Data Fig. 7) that shorter trials will elicit larger PEs. We found a significantly negative correlation overall in both mice (p<0.001, *left*), but the influence of distance was more prominent for later tones (*middle*, filled bar are have p<0.05) as in the model (Extended Data fig 7). Right panel shows that similar fraction of pixels that were responsive to each tone transition.

Supplementary Video 1: Example recording session demonstrating the activity pattern of
dopamine axons in the dorsal striatum. Video playback is 1X.

Supplementary Video 2: Video illustrating extraction of flow trajectories on a frame by frame
basis. Video playback is slowed down 0.25X.

Supplementary Video 3: Reward response in Instrumental task. Clock at top left displays time
relative to reward. Video playback is 1X.

Supplementary Video 4: Reward response in Pavlovian task. Clock at top left displays time
relative to reward. Video playback is 1X.

Supplementary Video 5: Progressively organized and continuous reward response to
unpredicted reward deliver in naive mice (*top*), or animals that have received training in
pavlovian sessions for 3 weeks. Clock at top left displays time relative to reward. Video
playback is slowed down 0.5X.