Quantifying genetic effects on disease mediated by assayed gene expression levels

Douglas W. Yao[1,*], Luke J. O'Connor[1,2], Alkes L. Price[2,3,4], and Alexander Gusev[5,6,*]

[1]Program in Bioinformatics and Integrative Genomics, Harvard Medical School, Boston, MA
[2]Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, MA
[3]Department of Biostatistics, Harvard T.H. Chan School of Public Health, Boston, MA
[4]Program in Medical and Population Genetics, Broad Institute, Cambridge, MA
[5]Department of Medical Oncology, Dana-Farber Cancer Institute and Harvard Medical School, Boston, MA
[6]Division of Genetics, Brigham and Women's Hospital, Boston, MA
[*]Correspondence should be addressed to D.W.Y. (douglasyao@g.harvard.edu) or A.G. (Alexander_Gusev@dfci.harvard.edu)

August 8, 2019

Abstract

Disease variants identified by genome-wide association studies (GWAS) tend to overlap with expression quantitative trait loci (eQTLs). However, it remains unclear whether this overlap is driven by mediation of genetic effects on disease by expression levels, or whether it primarily reflects pleiotropic relationships instead. Here we introduce a new method, mediated expression score regression (MESC), to estimate disease heritability mediated by the cis-genetic component of assayed steady-state gene expression levels, using summary association statistics from GWAS and eQTL studies. We show that MESC produces robust estimates of expression-mediated heritability across a wide range of simulations. We applied MESC to GWAS summary statistics for 42 diseases and complex traits (average $N = 323K$) and cis-eQTL data across 48 tissues from the GTEx consortium. We determined that a statistically significant but low proportion of disease heritability (mean estimate 11% with S.E. 2%) is mediated by the cis-genetic component of assayed gene expression levels, with substantial variation across diseases (point estimates from 0% to 38%). We further partitioned expression-mediated heritability across various gene sets. We observed an inverse relationship between cis-heritability of expression and disease heritability mediated by expression, suggesting that genes with weaker eQTLs have larger causal effects on disease. Moreover, we observed broad patterns of expression-mediated heritability enrichment across functional gene sets that implicate specific gene sets in disease, including loss-of-function intolerant genes and FDA-approved drug targets. Our results demonstrate that steady-state expression levels in bulk tissues are informative of regulatory disease mechanisms, while also motivating additional assays to more fully capture the regulatory effects of GWAS variants.

Introduction

In the past decade, genome-wide association studies (GWAS) have shown that most disease-associated variants lie in noncoding regions of the genome[1–3], leading to the hypothesis that regulation of gene expression levels is the primary biological mechanism through which genetic variants affect complex traits, and motivating large scale expression quantitative trait loci (eQTL) studies[4,5]. Many statistical methods have been developed to integrate eQTL data with GWAS data to gain functional insight into the genetic architecture of disease. These methods fall into two classes. The first class operates on individual genes and includes colocalization tests[6–10], which identify genes with shared causal variants for their expression levels and disease, and transcriptome-wide association studies[11–16], which identify genes with significant cis-genetic correlations between their expression and disease. The second class of methods operates on the entire genome and partitions disease heritability by SNP categories defined by eQTL status (i.e. whether or not SNPs are eQTLs)[17–20]. Application of these methods to eQTL and GWAS data has shown that many genes have

eQTLs that colocalize with GWAS loci[6–10] and/or exhibit significant cis-genetic correlations between their expression and trait[11–16, 21–28], while eQTLs as a whole are significantly enriched for disease heritability[17–20].

However, the true causal relationships between SNPs, gene expression levels, and disease remain uncertain for the vast majority of genes[9, 29–31]. This uncertainty arises from the fact that several different causal scenarios can result in similar patterns of enrichment/overlap between GWAS loci and eQTLs, summarized as follows: (1) mediation, in which a SNP affects gene expression levels, which then affect disease risk; (2) pleiotropy, in which a SNP affects gene expression levels and independently affects disease risk through an alternative mechanism; and (3) linkage, in which a SNP that affects gene expression levels is in linkage disequilibrium (LD) with a different, independent disease SNP. These scenarios carry very different biological interpretations. In particular, only scenario (1) is informative of the SNP's mechanism of action on disease, so it is vital to distinguish scenarios (2) and (3) from scenario (1). However, existing methods are unable to do so consistently. Among the individual-gene methods mentioned above, colocalization tests can sometimes rule out linkage as an explanation for overlap between eQTLs and disease SNPs, but cannot rule out pleiotropy[13, 32], while transcriptome-wide association studies cannot rule out either pleiotropy or linkage[13, 29], especially for genes with only one strong eQTL[15]. Among the genome-wide methods, some aim to rule out linkage through fine-mapping of eQTLs[19], but none aim to rule out pleiotropy. Thus, it remains unclear whether enrichment/overlap between eQTLs and disease SNPs usually reflects mediation, or whether it more commonly reflects pleiotropy and/or linkage[9, 29]. For example, in the case of autoimmune diseases, most instances of overlap between significant disease loci and immune cell eQTLs are driven by linkage[9], suggesting that linkage may be more prevalent than mediation[31].

In this study, we aim to quantify the proportion of disease heritability that is specifically mediated in cis by the assayed expression levels (scenario 1 from above) of the set of all genes, and of genes in specific functional categories. We first define expression-mediated heritability under a generative model featuring both mediated and non-mediated (including pleiotropic and linkage) effects of SNPs on the trait. We introduce a method, mediated expression score regression (MESC), to estimate expression-mediated heritability from GWAS summary statistics, ancestry-matched LD scores, and eQTL effect sizes obtained from external expression panels. Intuitively, MESC distinguishes mediated from non-mediated effects in a set of genes via the idea that mediation (unlike pleiotropy and linkage) induces a linear relationship between the magnitude of eQTL effect sizes and disease effect sizes.

Results

Definition of expression-mediated heritability

We first define heritability mediated by the cis-genetic component of gene expression levels (h^2_{med}), which is the quantity that our method aims to estimate. We model individual SNP effect sizes on a disease as the sum of a mediated component, defined as causal cis-eQTL effect sizes (where "cis" includes SNPs located within 1 Mb of the gene) multiplied by gene-trait effect sizes, and a non-mediated component, which includes pleiotropic, linkage, and trans-eQTL-mediated effects (Figure 1a; Methods). Under our model, h^2_{med} is defined as the squared mediated component of each SNP effect size summed across all SNPs (Figure 1b; Methods). We consider additional causality scenarios, such as reverse mediation, cis-by-trans mediation, and mediation by unobserved intermediaries, and we justify that these scenarios do not compromise our definition of h^2_{med} (see "Modes of expression causality" in Supplementary Note).

In this initial definition of h^2_{med}, which we call $h^2_{med;causal}$, cis-eQTL effect sizes are taken in the causal cell types/contexts for the disease, which are likely unobserved in practice. We extend this definition of h^2_{med} to allow for "tagging" effects of *assayed* expression in a set of tissues T, whereby assayed expression levels that are genetically correlated with unmeasured expression levels in causal cell types/contexts can contribute to h^2_{med}. We define $h^2_{med;assayed}(T)$ in terms of $h^2_{med;causal}$ as follows:

$$h^2_{med;assayed}(T) = r^2_g(T)h^2_{med;causal} \tag{1}$$

where $r^2_g(T)$ is the average squared genetic correlation between expression in T and expression in the causal cell types/contexts for the trait. $h^2_{med;assayed}(T)$ represents the final quantity that we estimate in practice,

and can be conceptualized as the heritability mediated/tagged by the cis-genetic component of assayed gene expression levels in a given set of tissues T.

Estimating expression-mediated heritability using MESC

In order to estimate $h^2_{med;assayed}(T)$, we first decompose $h^2_{med;causal}$ into the product of three quantities: the number of genes, the average cis-heritability of expression across all genes, and the average squared gene-trait effect size across all genes (Fig 1c; Methods). The average cis-heritability of expression across genes can be readily estimated with existing methods[33]. In order to estimate the average squared gene-trait effect size across all genes, we propose an approach that involves regressing causal squared SNP-trait effect sizes on causal squared cis-eQTL effect sizes summed across genes (Figure 1d; Methods). We show that this approach produces an appropriate estimate of $h^2_{med;assayed}(T)$ when using cis-eQTL effect sizes in assayed tissues T (Methods). For brevity, we refer to $h^2_{med;assayed}(T)$ as h^2_{med} for the remainder of the manuscript (where the set of tissues T is implicit).

We make two extensions to this simplified regression model. First, we model tagging effects due to LD between SNPs, which allows us to perform this regression using summary statistics from GWAS and eQTL studies (Figure 1d; Methods). Second, to avoid bias (see below), we stratify the regression across gene categories D, which allows us to estimate the heritability mediated by the assayed expression of genes in D, $h^2_{med}(D)$ (Methods). The regression equation used to estimate $h^2_{med}(D)$ is

$$E[\chi^2_k] = N \sum_c \tau_c \ell_{k;c} + N \sum_d \pi_d \mathcal{L}_{k;d} + 1 \tag{2}$$

where χ^2_k is the GWAS χ^2-statistic of SNP k, N is the number of samples, τ_c is the per-SNP contribution to non-mediated heritability of SNPs in SNP category C, $\ell_{k;c}$ is the *LD score*[2,34] of SNP k with respect to SNP category C (defined as $\ell_{k;c} = \sum_{j \in C} r^2_{jk}$), π_d is the per-gene contribution to $h^2_{med}(D)$, and $\mathcal{L}_{k;d}$ is the *expression score* of SNP k with respect to gene category D (defined as $\mathcal{L}_{k;d} = \sum_{i \in D} \sum_j^M r^2_{jk} \beta^2_{ij}$). $\mathcal{L}_{k;d}$ can be conceptualized as the total expression cis-heritability of genes in D that is tagged by SNP k.

Equation (2) allows us to estimate π_d and τ_c via computationally efficient multiple regression of GWAS chi-square statistics against LD scores and expression scores. In order for equation (2) to provide unbiased estimates of $h^2_{med}(D)$, the following key assumptions must be met:

- Within each gene category D, the magnitude of eQTL effect sizes is uncorrelated with the magnitude of gene-trait effect sizes

- Within each SNP category C, the magnitude of eQTL effect sizes is uncorrelated with the magnitude of non-mediated effect sizes

Note that if we estimate total h^2_{med} using a single gene category containing all genes and a single SNP category containing all SNPs (i.e. if we perform this regression without stratifying genes and SNPs), these two assumptions are likely violated in practice, leading to biased estimates of h^2_{med}. The first assumption is violated in the scenario that genes under selection systematically have low expression heritability and large effect on disease[5,35], causing h^2_{med} to be underestimated. We can ameliorate this bias by stratifying genes in a manner that captures the dependence between eQTL effect sizes and gene-trait effect sizes across the genome (e.g. stratifying genes by the magnitude of their expression cis-heritability). The second assumption from above is violated in the scenario that the magnitude of non-mediated SNP effect sizes and eQTL effect sizes tend to both be larger in biologically active regions of the genome[36–39] (e.g. promoters, enhancers), causing h^2_{med} to be overestimated. We can ameliorate this bias by stratifying SNPs in a manner that captures the dependence between eQTL effect sizes and non-mediated effect sizes across the genome (e.g. stratifying SNPs according to whether they lie in promoters, enhancers, etc.). In summary, stratifying the regression over strategically defined SNP and gene categories is necessary for us to obtain unbiased estimates of h^2_{med} in practice due to violations to the two above assumptions (see below for simulation results and "Model assumptions" in Methods for additional discussion).

In practice, we estimate LD scores from an external reference panel. We estimate expression scores in individual tissues from raw gene expression data and genotypes from external expression reference panels[11].

We do not exclude genes from our analyses based on any significance threshold, in contrast to previous approaches[9–13,19]. When applicable, we meta-analyze expression scores across tissues (Methods).

Throughout this study, we present estimates of three quantities that are a function of h^2_{med} and/or $h^2_{med}(D)$. These quantities include the proportion of heritability mediated by expression (defined as h^2_{med}/h^2_g), the proportion of expression-mediated heritability for gene category D (defined as $h^2_{med}(D)/h^2_{med}$), and the enrichment of expression-mediated heritability for D (defined as the proportion of expression-mediated heritability in D divided by the proportion of genes in D). We estimate standard errors and p-values for all quantities by block jackknife[2,34]. When applicable, we perform random effects meta-analysis of these quantities across diseases and complex traits (Methods). We have released open source software implementing our method (URLs).

Figure 1: **Schematic of MESC** (**a**) SNP effect sizes are modeled as the sum of a mediated component (defined as cis-eQTL effect sizes β multiplied by gene-trait effect sizes α) and a non-mediated component γ. (**b**) Heritability mediated by the cis-genetic component of gene expression levels (h^2_{med}) is defined as the squared mediated component of SNP effect sizes summed across all SNPs (assuming that genotypes are standardized). h^2_{med} can be rewritten as the product of the number of genes G, the average expression cis-heritability $E[h^2_{cis}]$, and the average gene-trait effect size $E[\alpha^2]$ (**c**) The basic premise behind MESC is to regress squared SNP effect sizes on squared eQTL effect sizes, in which case the slope will equal $E[\alpha^2]$ given appropriate effect size independence assumptions. (**d**) MESC can be extended to account for tagging effects due to LD. Summary MESC essentially involves regressing squared GWAS summary statistics on squared eQTL summary statistics with an LD score covariate, in which case the coefficient for the eQTL summary statistics will be $E[\alpha^2]$. In the figure, we show a simplified LD architecture with two discrete levels of LD.

Simulations assessing calibration and bias

We performed simulations to assess the calibration and bias of MESC in estimating h^2_{med}/h^2_g and its standard error from simulated complex trait and expression data under a variety of genetic architectures (Methods). We performed all simulations using real genotypes from UK Biobank[40] ($N_{GWAS} = 10,000$ samples; $M = 98,499$ SNPs from chromosome 1). We simulated causal cis-eQTL effect sizes for $G = 1000$ genes, gene

effect sizes on a complex trait, and non-mediated effect sizes of each SNP on the complex trait from normal or point-normal distributions corresponding to various levels of h_{med}^2, with total h_g^2 fixed at 0.5. We then simulated complex trait phenotypes from these effect sizes, with normally distributed environmental effects. To emulate an external expression panel, we simulated expression phenotypes from the eQTL effect sizes using a separate set of genotypes, with normally distributed environmental effects (N_{eQTL} = 100-1000 samples). For all simulations other than Figure 2c, eQTL effect sizes used to generate expression panel phenotypes and complex trait phenotypes were identical (i.e. $r_g^2(T) = 1$).

First, we evaluated the impact of expression panel sample size and different methods of estimating expression scores on the bias of MESC. We used five different methods to estimate expression scores from simulated expression panels with sample sizes ranging from 100 to 1000. These methods include: (1) eQTL summary statistics, (2), LASSO[41], (3) LASSO with REML correction, (4) BLUP[42], and (5) BLUP with REML correction (see "Comparing different methods of estimating expression scores" in Supplementary Note for description of these methods). We then used these expression scores to estimate h_{med}^2/h_g^2 from the simulated complex trait phenotypes. We observed that LASSO with REML correction gave us approximately unbiased estimates of h_{med}^2/h_g^2 with around 500 or more expression panel samples (Figure 2a), whereas all other methods of estimating expression scores produced biased results at all sample sizes tested (Supplementary Figure 1) due to large sampling noise and/or systematic bias in their estimates of expression scores (Supplementary Note and Supplementary Figure 2). Notably, all methods (including LASSO with REML correction) produced biased estimates of h_{med}^2/h_g^2 at N_{eQTL} comparable to individual tissue expression panels ($N_{eQTL} < 200$). We performed additional simulations in which we estimated h_{med}^2/h_g^2 using expression scores meta-analyzed across multiple simulated tissues (Supplementary Note). We obtained approximately unbiased estimates of h_{med}^2/h_g^2 with 5 tissues × 200 samples per tissue (Supplementary Figure 3), demonstrating that we were able to ameliorate low-N_{eQTL} dependent downward bias in h_{med}^2/h_g^2 via meta-analysis of expression scores across tissues.

To test the bias of MESC in sparse genetic/gene architectures, we performed simulations in which we varied the number of eQTLs per gene, the proportion of SNPs with non-mediated effects, and the proportion of genes with gene-trait effects. We observed that MESC was robust to the number of eQTLs per gene and sparsity of gene effect sizes or non-mediated effect sizes (Figure 2b).

Next, we tested the calibration of jackknife standard errors computed by MESC for the h_{med}^2 enrichment of a gene category D, defined as (proportion of h_{med}^2 in D) / (proportion of genes in D). This is the main quantity that we aim to estimate for analyses involving functional gene sets. We simulated a gene category containing 200 genes. We then performed two sets of simulations in which this gene category was either enriched or not enriched for h_{med}^2 (Supplementary Note). We observed well-calibrated h_{med}^2 enrichment standard errors for the gene category under both the null and causal enrichment scenarios (Supplementary Figure 4).

Finally, we sought to verify that MESC estimates the quantity $r_g^2(T)h_{med;causal}^2$ when estimating expression scores from a tissue with squared genetic correlation $r_g^2(T)$ with the causal tissue. We simulated expression phenotypes for assayed tissues that were genetically correlated with expression in the causal tissue ($r_g^2(T) = 0.5$-1), then predicted expression scores from the assayed tissues and estimated h_{med}^2/h_g^2 from these expression scores (Methods). We obtained unbiased estimates of $r_g^2(T)h_{med;causal}^2/h_g^2$ for each value of $r_g^2(T)$ (Figure 2c).

In summary, we show that MESC produces approximately unbiased estimates of h_{med}^2/h_g^2 and well-calibrated standard errors under a variety of genetic architectures at $N_{eQTL} > 500$. Given that many individual tissue expression panels have N_{eQTL} of 200 or fewer[5], this suggests that meta-analysis of expression scores across tissues is necessary to obtain unbiased estimates of h_{med}^2/h_g^2 in practice. Note that all simulations in this section were conducted assuming that the two main effect size independence assumptions were satisfied across the genome; see below for simulations under violations of these assumptions.

Figure 2: **Simulation results assessing bias of MESC.** We simulated expression and complex trait architectures corresponding to various levels of h^2_{med}/h^2_g. GWAS sample size was fixed at 10,000 and h^2_g was fixed at 0.5. Error bars represent mean standard errors across 300 simulations. (**a**) Impact of expression panel sample size on estimates of h^2_{med}/h^2_g. Expression scores were estimated from simulated expression panel samples using LASSO with REML correction. See Supplementary Figure 1 for estimates of h^2_{med}/h^2_g using other methods of estimating expression scores. (**b**) Impact of sparse genetic/eQTL architectures on estimates of h^2_{med}/h^2_g. Expression panel sample size was set to 1,000. (**c**) Estimates of h^2_{med}/h^2_g with $r^2_g(T) < 1$. Expression panel sample size was set to 1,000.

Simulations under violations of effect size independence assumptions

We performed simulations to assess the bias of MESC in the presence of violations to the two main effect size independence assumptions, and to assess how well partitioning genes and SNPs ameliorated this bias (see "Model assumptions" in Methods for discussion of these assumptions).

In order to simulate violations to gene-eQTL effect size independence, we simulated eQTL effect sizes and gene effect sizes so that the per-gene h^2_{med} was constant across genes, but the magnitude of eQTL effect sizes and gene effect sizes were strongly inversely proportional across genes (Methods). This emulates the scenario in which negative selection causes low-heritability genes to have larger gene-trait effects. Non-mediated effect sizes were simulated in the same manner as previous simulations. As expected, when estimating h^2_{med}/h^2_g in this scenario using expression scores corresponding to a single gene category, we obtained downwardly biased estimates of h^2_{med}/h^2_g. When we estimated h^2_{med}/h^2_g with genes stratified by their estimated expression cis-heritability into 5 gene categories D, we obtained approximately unbiased estimates of h^2_{med}/h^2_g (Figure 3a), as well as approximately unbiased estimates of $h^2_{med}(D)/h^2_g$ for each D (Supplementary Figure 5). This result demonstrates that partitioning genes into expression cis-heritability bins can capture a continuous underlying relationship between eQTL effect sizes and gene-trait effect sizes across the genome, enabling us to obtain unbiased estimates of h^2_{med}/h^2_g in the presence of violations to gene-eQTL effect size independence.

In order to simulate violations to pleiotropy-eQTL effect size independence, we performed simulations in which specific SNP categories were enriched for both eQTL effects and non-mediated effects. This emulates the presence of regulatory hotspots in the genome with a high concentration of both eQTL effects and non-mediated effects, and it induces a genome-wide positive correlation between eQTL effect size magnitude and non-mediated effect size magnitude. We selected three functional SNP categories from the baselineLD model[2,43]—transcription start sites, coding regions, and conserved regions—that have been shown to be highly enriched for both expression cis-heritability[44] and complex trait heritability[2]. We then simulated 10x expression cis-heritability enrichment and 10x total heritability enrichment for SNPs in any of these three categories, which is close to empirical estimates[2,44]. Gene effect sizes were simulated in the same manner as previous simulations, and we estimated h^2_{med}/h^2_g using a single SNP category containing all SNPs. However, we observed no discernible upward bias in these estimates of h^2_{med}/h^2_g (Supplementary Figure 6), demonstrating that stronger violations to pleiotropy-eQTL effect size independence are necessary to induce bias in estimates of h^2_{med}/h^2_g.

In order to simulate more extreme patterns of colocalization between eQTL effects and non-mediated effects, we performed three sets of simulations in which *all* eQTL effect sizes and non-mediated effect sizes were entirely localized in each of the three SNP categories. When we estimated h^2_{med}/h^2_g for each simulation using a single SNP category containing all SNPs, we obtained upwardly biased estimates of h^2_{med}/h^2_g. We then estimated h^2_{med}/h^2_g for each simulation in two additional ways: (1) with SNPs stratified by the full baselineLD model, and (2) with SNPs stratified by a misspecified baselineLD model missing the causal category and window around the causal category, emulating a realistic scenario in which the SNP model does not fully capture all sources of non-mediated effects. When we estimated h^2_{med}/h^2_g with SNPs stratified by the full baselineLD model, we obtained unbiased estimates of h^2_{med}/h^2_g. When we estimated h^2_{med}/h^2_g with SNPs stratified by the misspecified baselineLD model, we obtained estimates of h^2_{med}/h^2_g that were nearly unbiased (Figure 3b, Supplementary Figure 7), demonstrating that the remaining SNP categories in the baselineLD model were able act as a reasonable proxy to the missing causal category.

In summary, we show that partitioning SNPs by the baselineLD model can capture diverse patterns of non-independence between eQTL effect sizes and non-mediated effect sizes across the genome, allowing us to obtain unbiased estimates of h^2_{med}/h^2_g even in the presence of extreme violations to pleiotropy-eQTL effect size independence.

Figure 3: **Simulation results for violations of effect size independence assumptions.** Error bars represent mean standard errors across 300 simulations. (**a**) We simulated eQTL effect sizes and gene effect sizes so that the per-gene h^2_{med} was constant across genes, but the magnitude of eQTL effect sizes and gene effect sizes were strongly inversely proportional across genes. (**b**) We simulated all eQTL effects and non-mediated effects to lie within coding regions. Results in which all eQTL effects and non-mediated effects lie in conserved regions or transcription start sites can be found in Supplementary Figure 6.

Comparison to other methods in simulations

To our knowledge, no published methods specifically aim to estimate heritability mediated by expression levels. The closest analogues are approaches that measure the genome-wide heritability enrichment of eQTLs[17–20] using GCTA[33] or stratified LD score regression (S-LDSC)[2,43]. If pleiotropy-eQTL effect size independence holds *across the genome*, then the heritability enrichment of a SNP category corresponding to the set of all eQTLs will accurately reflect h^2_{med} (i.e. the heritability enrichment will be 1x when $h^2_{med} = 0$ and will increase when $h^2_{med} > 0$). Given pleiotropy-eQTL effect size independence, we observed that S-LDSC has a well-calibrated false positive rate for detecting heritability enrichment of the eQTL category when $h^2_{med} = 0$, with increasing power to detect heritability enrichment of the eQTL category as h^2_{med} increased

(Supplementary Figure 8).

However, in the presence of violations to pleiotropy-eQTL effect size independence, S-LDSC will detect significant heritability enrichment of the eQTL category even in the absence of mediation. Like the simulation performed in Supplementary Figure 6, we simulated 10x enrichment of eQTL effect sizes in three SNP categories (coding regions, transcription start sites, and conserved regions). With h^2_{med} fixed at 0, we then varied the heritability enrichment of the three SNP categories from 2.5x to 10x. We observed that S-LDSC detected significant heritability enrichment of the eQTL category in the absence of mediation when total heritability enrichment was 5x or greater, whereas MESC had a well-calibrated false positive rate at all levels of enrichment (Figure 4). Note that this result does not imply that S-LDSC or other heritability partitioning methods are flawed, but rather that they cannot specifically distinguish mediated effects from non-mediated effects when they are applied to annotations generated from eQTL data. We did not compare MESC to the colocalization methods of Chun et al.[9] or Ongen et al.[10], since these methods only operate on genome-wide significant GWAS loci and also do not attempt to distinguish pleiotropy from mediation.

Figure 4: **Comparison of MESC to heritability partitioning.** We fixed h^2_{med}/h^2_g at 0 for all simulations. We simulated 10x enrichment of eQTL effect sizes for SNPs within three causal SNP categories (coding regions, transcription start sites, and conserved regions). We then simulated heritability enrichment in the same three SNP categories ranging from 2.5x to 10x. In the figure, we show the proportion of simulations in which the null hypothesis that $h^2_{med}/h^2_g = 0$ is rejected by MESC, and the proportion of simulations in which the null hypothesis of no h^2_g enrichment for the set of all eQTLs is rejected by stratified LD-score regression (S-LDSC). 300 simulations were performed.

Estimation of expression-mediated heritability for 42 diseases and complex traits

We applied MESC to estimate the proportion of heritability mediated by the cis-genetic component of assayed expression levels (h^2_{med}/h^2_g) for 42 independent diseases and complex traits from the UK Biobank[40] and other publicly available datasets (average $N = 323$K; see Supplementary Table 1 for list of traits). We obtained estimates of expression cis-heritability for each gene in each of 48 tissues from the GTEx consortium[5] using GCTA[33]. We estimated individual-tissue expression scores from raw expression data (with appropriate quality control steps) and individual-level genotypes in 48 tissues from GTEx. (Methods; see Supplementary Table 2 for list of tissues). In contrast to approaches that restrict to genes with significantly heritable expression[11, 12], we included in our analysis all gene-tissue pairs for which LASSO converged when estimating eQTL effect sizes, resulting in an average of 13,674 genes analyzed per tissue (see Supplementary Table 2 for number of genes per tissue). To capture dependencies between eQTL effect sizes and non-

mediated effect sizes and avoid upward bias in h^2_{med}/h^2_g estimates, we partitioned SNPs by 72 functional categories specified by the baselineLD v2.0 model[2,43] (Methods). To capture dependencies between eQTL effect sizes and gene-trait effect sizes and avoid downward bias in h^2_{med}/h^2_g estimates, we partitioned genes into 5 bins based on the magnitude of their estimated expression cis-heritability. Finally, we meta-analyzed expression scores in two ways: within groups of GTEx tissues with common biological origin, and across all 48 GTEx tissues (Methods). For each trait, we used MESC to estimate three versions of h^2_{med}/h^2_g using three different types of expression scores: (1) individual-tissue expression scores, (2) tissue-group meta-analyzed expression scores, and (3) all-tissue meta-analyzed expression scores.

Across all traits, we observed an average h^2_{med}/h^2_g of 0.11 (S.E. 0.02) from all-tissue meta-analyzed expression scores. Of the 42 traits, 26 had h^2_{med}/h^2_g estimates greater than 0 at nominal significance (p-value < 0.05). In Figure 5a, we report h^2_{med}/h^2_g estimates from all-tissue and tissue-group meta-analyzed expression scores for 12 significant traits (after pruning closely related traits). h^2_{med}/h^2_g estimates for all 42 traits from all three types of expression scores can be found in Supplementary Figure 9 and Supplementary Table 3. Between the three types of expression scores, we observed the lowest estimates of h^2_{med}/h^2_g from individual-tissue expression scores (Figure 5b, Supplementary Figure 9). Moreover, we observed a positive correlation between the sample size of the tissue and magnitude of the estimated h^2_{med}/h^2_g ($R^2 = 0.71$) (Supplementary Figure 10). These results demonstrate that tissue sample size affected our ability to obtain unbiased estimates of h^2_{med}/h^2_g from individual-tissue expression scores, consistent with our simulations that suggest downward bias in h^2_{med}/h^2_g estimates at $N_{eQTL} < 200$ (Figure 2a).

When we estimated h^2_{med}/h^2_g using tissue-group expression scores and all-tissue expression scores, we obtained consistently higher estimates of h^2_{med}/h^2_g than individual-tissue expression scores (Figure 5b). Moreover, the tissue-group resulting in the highest estimate of h^2_{med}/h^2_g for each trait was generally consistent with the known causal tissue of the trait (Figure 5a), unlike the highest individual tissue in many instances (Supplementary Figure 9). These results suggest we were able to account for low-N_{eQTL} dependent downward bias in h^2_{med}/h^2_g by meta-analysis of expression scores across tissues, consistent with our simulation result in Supplementary Figure 3.

Overall, we did not observe a relationship between h^2_{med}/h^2_g and h^2_g across all traits ($R^2 = 0.004$) (Supplementary Figure 11). The traits that did not have a h^2_{med}/h^2_g estimate significantly greater than 0 comprised traits with a known low amount of polygenicity (balding and various pigmentation traits including skin color, tanning, sunburn occasion, and hair color) and various complex neurological traits (autism, number children ever born, depressive symptoms, morning person, neuroticism, smoking status, anorexia).

To explore the nature of dependencies between eQTL effect sizes, gene-trait effects, and non-mediated effects across the genome, we estimated h^2_{med}/h^2_g for all traits as before, but without stratifying genes and/or SNPs (Supplementary Figure 12). When we did not stratify genes by expression cis-heritability bins, we obtained much lower estimates of h^2_{med}/h^2_g (mean estimate 0.011 with S.E. 0.003). This result is consistent with our simulation results for Figure 3a and suggests a genome-wide negative correlation between the magnitude of eQTL effect sizes and gene-trait effect sizes in practice. Moreover, when we did not stratify SNPs by the baselineLD model, we obtained much higher estimates of h^2_{med}/h^2_g (mean estimate 0.45 with S.E 0.03). This result is consistent with our simulation results for Figure 3b and suggests a genome-wide positive correlation between the magnitude of eQTL effect sizes and non-mediated effect sizes in practice. Together, these results demonstrate that both gene-eQTL independence and pleiotropy-eQTL independence are strongly violated in practice, justifying the necessity of stratifying genes by expression cis-heritability bins and stratifying SNPs by the baselineLD model to obtain unbiased estimates of h^2_{med}/h^2_g.

Finally, to investigate whether the number of expression cis-heritability bins influenced our estimates of h^2_{med}/h^2_g, we estimated h^2_{med}/h^2_g for all traits as before, but with genes stratified into 10 bins rather than 5 based on their estimated expression cis-heritability. We obtained very similar estimates of h^2_{med}/h^2_g (Supplementary Figure 13), demonstrating that 5 bins were adequate to capture genome-wide dependencies between the eQTL effect sizes and gene-trait effect sizes.

Genes with low expression heritability explain more expression-mediated disease heritability

To investigate the relationship between magnitude of expression cis-heritability (h^2_{cis}) and amount of complex trait heritability mediated by those genes, we looked at the proportion of h^2_{med} (defined as $h^2_{med}(D)/h^2_{med}$

Figure 5: **Estimates of proportion of heritability mediated by expression.** (**a**) Estimated proportion of heritability mediated by the cis-genetic component of assayed gene expression levels (h^2_{med}/h^2_g) for 12 traits. See Supplementary Figure 9 for estimates of h^2_{med}/h^2_g for all 42 traits. Error bars represent jackknife standard errors. For each trait, we report the h^2_{med}/h^2_g estimate for "All tissues" (expression scores meta-analyzed across all 48 GTEx tissues) and "Best tissue group" (expression scores meta-analyzed within 7 tissue groups). Here, "best" refers to the tissue group resulting in the highest estimates of h^2_{med}/h^2_g compared to all other tissue groups. (**b**) h^2_{med}/h^2_g estimates meta-analyzed across all 42 traits. Here, "Best tissue" refers to the individual tissue resulting in the highest estimates of h^2_{med}/h^2_g compared to all other tissues.

for gene category D) mediated by genes stratified into 10 bins by their h^2_{cis}. We performed all analyses using all-tissue meta-analyzed expression scores. Across 26 traits with h^2_{med} significantly greater than 0, we observed an inverse relationship between meta-tissue h^2_{cis} and proportion of h^2_{med} across gene bins (Figure 6, Supplementary Table 4), with 32% of h^2_{med} explained by the lowest 2 bins (mean meta-tissue $h^2_{cis} = 0.014$) and only 3% of h^2_{med} explained by the highest 2 bins (mean meta-tissue $h^2_{cis} = 0.30$). This result suggests that genes with less heritable expression have larger causal effect sizes on the complex trait.

There are several reasons why genes with less heritable expression might have larger causal effects on the complex trait. One explanation is that negative selection purifies out strong eQTLs for genes with large effect on complex traits[5, 35]. Alternatively, genes with low meta-tissue h^2_{cis} may comprise genes with tissue-specific eQTLs, which have been shown to be enriched for disease heritability[5, 19, 20].

In support of the first explanation, we observed that the probability of being loss-of-function intolerant[45] (i.e. pLI) and the level of selection against protein-truncating variants[46] (i.e. s_{het}) were inversely proportional to meta-tissue h^2_{cis} (Supplementary Figure 14). Moreover, consistent with the first explanation, we observed that the proportion of meta-tissue h^2_{cis} explained by the top eQTL was proportional to overall meta-tissue h^2_{cis} (i.e. genes with low meta-tissue h^2_{cis} tended to be affected by multiple weaker eQTLs rather than a single strong eQTL) (Supplementary Figure 15). To investigate the second explanation, we considered the possibility that most low meta-tissue h^2_{cis} genes were primarily genes with nonzero individual-tissue h^2_{cis} in only one or a small number of tissues, which is consistent with the genes having tissue-specific eQTLs. However, we did not observe strong evidence for this hypothesis (see "Role of tissue specificity in explaining low heritability genes" in Supplementary Note). Moreover, we did not observe a strong relationship between meta-tissue h^2_{cis} and the mean expression level of the gene across tissues, nor the number of tissues in which the gene was expressed (Supplementary Figure 17).

In summary, our results support the hypothesis that negative selection contributes to the inverse relationship between expression cis-heritability and proportion of h^2_{med}.

Figure 6: **Low heritability genes explain more expression-mediated disease heritability.** Estimated proportion of expression-mediated heritability $(h^2_{med}(D)/h^2_{med})$ for 10 gene bins stratified by magnitude of expression cis-heritability. Results are meta-analyzed across 26 traits with nominally significant h^2_{med}. Results for individual traits can be found in Supplementary Table 4. In the legend, the number in parentheses next to the bin number indicates the average expression cis-heritability of genes in the gene bin.

Expression-mediated heritability enrichment in functional gene sets

To gain insight into the distribution of expression-mediated effect sizes across various functional gene sets, we estimated h^2_{med} enrichment, defined as (proportion of h^2_{med}) / (proportion of genes), for these gene sets. We gathered gene sets relevant for disease and specific biological pathways from publicly available sources (see Supplementary Table 5 for list of gene sets). We analyzed 827 gene sets from three main sources: (1) 10 gene sets reflecting various broad metrics of gene essentiality (Methods; URLs); (2) 780 gene sets reflecting specific biological pathways, including gene sets from the KEGG[47], Reactome[48], and Gene Ontology (GO)[49] pathway databases (Methods; URLs); and (3) 37 gene sets comprising of genes specifically expressed in 37 different GTEx tissues[50] (Methods). We restricted our analyses to large gene sets with at least 200 genes (Methods), since we observed large standard errors in h^2_{med} enrichment estimates for gene sets with 200 or fewer genes (Supplementary Figure 18). We applied MESC to estimate the h^2_{med} enrichment for each of the gene sets in 26 significant traits. As before, we stratified SNPs by the baselineLD model and genes by 5 cis-heritability bins. We performed all analyses using all-tissue meta-analyzed expression scores (with the exception the tissue-specific expression analyses in Figure 7c, in which case we used tissue-group meta-analyzed expression scores).

Out of 21,502 gene set-complex trait pairs (827 gene sets × 26 complex traits), we observed 226 gene set-complex trait pairs (comprising of 117 unique gene sets) with FDR-significant h^2_{med} enrichment (FDR < 0.05 accounting for 21,502 hypotheses tested). Significant h^2_{med} enrichment estimates ranged from 1.5x to 51x across gene-set complex trait pairs. The full list of h^2_{med} enrichment estimates for all 21,502 gene set-complex trait pairs is reported in Supplementary Table 6.

In Figure 7a, we show h^2_{med} enrichment estimates for all 10 broadly essential gene sets meta-analyzed across 26 complex traits. h^2_{med} enrichment estimates for individual traits can be found in Supplementary Figure 19. We observed significant h^2_{med} enrichment (p < 0.05 after Bonferroni correction for 10 hypotheses) for 8 gene sets, including ExAC loss-of-function intolerant genes[45] (3.9x enrichment; p = 2.3×10^{-25}), FDA-approved drug targets[51] (5.2x enrichment; p = 2.0×10^{-5}), genes essential in mice[52–54] (4.0x enrichment; p

$= 1.1 \times 10^{-10}$), and genes nearest to GWAS peaks[55] (3.9x enrichment; p $= 5.0 \times 10^{-46}$).

Of the 780 pathway gene sets, we observed that 97 had a significant h^2_{med} enrichment (FDR < 0.05) in at least one of the 26 complex traits. In Figure 7b, we show the h^2_{med} enrichment estimates of a representative set of 156 gene set-complex traits pairs, comprising of 13 pathway-specific gene sets $\times$ the same 12 traits presented in Figure 5a. More comprehensive results, comprising of h^2_{med} enrichment estimates for all 97 significant pathway gene sets $\times$ 26 complex traits, can be found in Supplementary Figure 20. Most gene sets exhibited highly trait-specific patterns of enrichment that were consistent with the known biology of the trait. In particular, we observed significant h^2_{med} enrichment of:

- Various brain-related pathways (including fragile X mental retardation protein (FMRP)-interacting genes and neuronal/synapse genes) for brain traits (schizophrenia, years of education, BMI)

- Matrisome genes for height

- Wnt signaling pathway for bone density

- Various developmental pathways (including genes involved in tissue/gland development, ossification, and embryonic morphogenesis) for growth-related traits (height, bone density, forced vital capacity, FEV1-FVC ratio, systolic blood pressure, waist-hip ratio)

- Various immune pathways (including genes involved in immune response, lymphocyte activation, and cytokine-mediated signaling pathway) for immune traits (eczema, respiratory/ear-nose-throat diseases, ulcerative colitis, Crohn's disease, hypothyroidism, white blood cell count, eosinophil count)

- Hemostasis/wound response for platelet count

- Lipid metabolism/binding/localization genes for lipid traits (high-density lipoproteins, high cholesterol)

- Very general regulatory pathways (including genes involved in regulation of cell differentiation, proliferation, transcription, response to stimulus, and communication) for many traits

Finally, we investigated whether genes specifically expressed in 37 different GTEx tissues[50] were enriched for h^2_{med} (Methods). It has previously been shown that SNPs near the top 10% of genes most differentially expressed in putative causal tissues/cell types for specific traits are highly enriched for complex trait heritability[50]. However, it is unknown whether the effects of these SNPs are mediated by the expressed genes, or whether they impact the trait through alternate mechanisms in the same loci. We examined the same set of specifically expressed genes in GTEx tissues investigated in ref.[50]. In additional to stratifying SNPs by the baselineLD model, we included a SNP category corresponding to 100Kb windows around specifically expressed genes, which is the same SNP category used to identify complex trait heritability enrichment in ref.[50]. We found significant h^2_{med} enrichment (FDR < 0.05) in genes specifically expressed in brain tissues for brain-related traits (schizophrenia, years of education, BMI) (Figure 7c), suggesting that the complex trait heritability of SNPs near genes specifically expressed in causal tissues (at least for brain traits) is mediated by the expression of those genes.

Figure 7: **Mediated heritability enrichment estimates for functional gene sets.** For all plots, x axis represents complex traits and y axis represents gene sets. The number in parentheses next to each gene set label indicates the number of genes in that gene set. (**a**) h^2_{med} enrichment estimates for 10 broadly essential gene sets meta-analyzed across 26 complex traits. h^2_{med} enrichment estimates for individual traits can be found in Supplementary Figure 19. (**b**) For ease of display, we report h^2_{med} enrichment estimates for a representative set of 13 pathway-specific gene sets across 11 complex traits. h^2_{med} enrichment estimates for additional complex traits and gene sets can be found in Supplementary Figure 20 and Supplementary Table 6. (**c**) h^2_{med} enrichment estimates for 37 gene sets corresponding to the top 10% most differentially expressed genes in 37 GTEx tissues. Brain tissues (13 total) are indicated as so in the figure. h^2_{med} enrichment estimates for additional complex traits, with individual GTEx tissues labelled, can be found in Supplementary Figure 21.

Discussion

We have developed a new method, mediated expression score regression (MESC), to estimate complex trait heritability mediated by the cis-genetic component of assayed expression levels (h^2_{med}) from GWAS summary statistics and eQTL effect sizes estimated from an external expression panel. Our method is distinct from existing methods that identify and quantify overlap between eQTLs and GWAS hits (including colocalization tests[6–10], transcriptome-wide association studies[11–15], and heritability partitioning by eQTL status[17–20]) in that it specifically aims to distinguish directional mediated effects from non-directional pleiotropic and linkage effects. Moreover, our polygenic approach does not require individual eQTLs or GWAS loci to be significant and is not impacted by the sparsity of eQTL effect sizes, so unlike other approaches[9–13, 19] we do not exclude genes or SNPs from our analyses based on any significance thresholds. We applied our method to summary statistics for 42 traits and eQTL effect sizes estimated from 48 GTEx tissues. We show that across traits, a

significant but modest proportion of complex trait heritability (on average 0.11 with S.E. 0.02) is mediated by the cis-genetic component of expression levels. Though many previous approaches have hypothesized that SNPs impact complex traits by directly modulating gene expression levels, our results provide the first concrete genome-wide evidence for this hypothesis. On the other hand, the fact that our h^2_{med}/h^2_g estimates are low for most traits suggests that assayed steady-state expression levels in bulk postmortem tissues from studies such as GTEx do not capture most of the mediated effect of complex trait heritability, motivating additional assays to better identify molecular mechanisms impacted by regulatory GWAS variants.

There are two explanations for our low h^2_{med}/h^2_g estimates. The first explanation is that the complex trait heritability mediated by the cis-genetic component of expression levels in causal cell types/contexts for the trait (i.e. $h^2_{med;causal}$) is high in reality, but assayed expression from the GTEx consortium poorly tags expression in causal cell types/contexts (i.e. $r^2_g(T)$ is low). The second explanation is that assayed expression adequately tags expression in causal cell types/contexts, but $h^2_{med;causal}$ is low in reality. In particular, complex trait heritability can be mediated in ways other than through gene expression in cis, including mediation through protein-coding changes or splicing, or mediation through expression levels purely in trans (note that our model does capture cis-by-trans effects, in which a SNP acts as a trans-eQTL for a gene via its effects in cis on the expression of another gene (Supplementary Note)). Both the above explanations are consistent with motivating additional assays. If the first explanation is predominant, then context-specific expression[56] and single-cell expression[57] assays are desirable to capture expression levels in highly cell type/context specific scenarios. If the second explanation is predominant, then additional assays such as splicing[58], histone mark[4], chromosome conformation[59], and trans-eQTL[60] assays can be informative for probing other molecular mechanisms impacted by GWAS variants. We anticipate that these additional assays will allow us to explain a greater proportion of mediated complex trait heritability, and that MESC will be used to benchmark the progress made by future QTL studies.

We observed that expression scores meta-analyzed across tissues gave us higher estimates of h^2_{med}/h^2_g than individual-tissue expression scores, suggesting that disease-associated cis-eQTLs often affect expression in multiple tissues. This result is consistent with previous studies that reported higher heritability enrichment of cis-eQTLs meta-analyzed across all GTEx tissues compared to individual tissues[19], higher prediction accuracy for imputed expression using joint prediction from multiple tissues compared to individual tissues[61], and high cis-genetic correlations of expression between tissues overall[44,62]. Nevertheless, we did obtain higher estimates of h^2_{med}/h^2_g using expression scores from the presumed causal tissue group for a given disease compared to other tissue groups, suggesting some tissue-specific effects.

Despite the fact that assayed expression levels only explain a small proportion of overall disease heritability, we were still able to extract useful biological insight from partitioning expression-mediated heritability across gene categories. In particular, we observed a strong inverse relationship between proportion of h^2_{med} and expression cis-heritability across genes, suggesting that genes with low expression cis-heritability have large effects on complex traits. This result may be explained by negative selection purifying out strong eQTLs for genes with large effect on disease[35], or by genetic compensatory mechanisms that buffer against the phenotypic effects of genes with strong genetic effects on their expression[63,64]. Alternatively, genes with low expression cis-heritability may have highly cell type-specific eQTLs whose effect sizes are diluted by the inclusion of other cell types in a bulk assayed tissue sample. Regardless of the underlying nature of low-heritability genes, this result suggests that integrative association tests that prioritize genes based on probability of colocalization between eQTLs and GWAS hits[6,8,9] and/or significance of genetic correlation between expression and trait[11–13] may not detect the most mechanistically important genes, since these methods have lower power for genes with weaker eQTLs. Instead, our result suggests that genes with weaker eQTLs should be prioritized, and it motivates the implementation of larger eQTL studies and/or cell-type specific assays to more accurately detect these weak eQTLs.

We did not observe significantly nonzero h^2_{med} for low polygenicity traits (pigmentation traits and balding) and many neurological traits (autism, number children ever born, depressive symptoms, morning person, neuroticism, smoking status, anorexia). There is evidence that pigmentation traits are driven in large part by coding changes in MC1R[65] and a small number of other genes[66–68]. Moreover, there is evidence that eQTLs important for neurological traits can be unique to fetal brain expression and thus highly context-specific[69,70]. Neither the protein-coding changes in pigmentation genes nor the context-specific eQTLs in brain genes will be captured by steady-state expression level measurements in assayed postmortem tissues, and thus any SNP effects mediated by them will also not be observed in our analysis.

There are several limitations to our method. First, our method makes the assumption that the magnitude of eQTL effect sizes is uncorrelated with the magnitude of both gene-trait effect sizes (gene-eQTL independence) and non-mediated effect sizes (pleiotropy-eQTL independence) within each SNP/gene category included in the model. Violations to genome-wide gene-eQTL independence can be resolved by stratifying genes by the magnitude of their expression cis-heritability; given that expression cis-heritability is directly estimable for each gene in practice, violations to genome-wide gene-eQTL independence are readily resolvable (see Figure 3a and Supplementary Figure 13). Meanwhile, violations to genome-wide pleiotropy-eQTL independence can be resolved by stratifying SNPs by the magnitude of their eQTL effect sizes; however, due to LD, we cannot obtain precise estimates of eQTL effect sizes for individual SNPs. Instead, our model assumes that the SNP categories in the baselineLD model act as a reasonable proxy for eQTL effect size magnitude. If the magnitude of eQTL effect sizes and non-mediated effect sizes are positively correlated within the baselineLD categories included in our analyses, then our estimates of h^2_{med}/h^2_g would be upwardly biased. However, given that the baselineLD model was able to account for extreme violations to pleiotropy-eQTL independence in simulations (Figure 3b), and that the average h^2_{med}/h^2_g across traits was only 0.11, this upward bias (if present) is likely to be small. An alternative approach to estimating h^2_{med}/h^2_g involves regressing pairwise products of GWAS summary statistics on pairwise products of eQTL summary statistics[71]. Unlike MESC, this approach yields unbiased estimates h^2_{med}/h^2_g even when pleiotropy-eQTL independence is violated within SNP categories. However, the approach requires aggressive LD pruning in order to retain SNP pairs in linkage equilibrium, causing the approach to be highly underpowered in practice (L.J.O, unpublished data). The second limitation of our method is that it relies on the accurate estimation of expression scores from external expression panel samples. In order for our method to be well-powered, it requires large expression panel sample sizes that can only be obtained through meta-analysis across individual tissues at current sample sizes. Third, the quantity that our method estimates in practice (i.e. heritability mediated by *assayed* gene expression levels) can potentially be much smaller than the theoretical quantity of heritability mediated by expression levels in causal cell types/contexts if assayed gene expression levels do not adequately capture expression levels in causal cell types/contexts. Fourth, our method can only provide reliable h^2_{med} enrichment estimates for large gene sets on the order of 200 or more genes, so smaller gene pathways or individual genes cannot be prioritized using our method. Fifth, our method does not capture non-additive effects of SNPs on gene expression or gene expression on trait.

Despite these limitations, our method provides a novel framework to distinguish mediated effects from pleiotropic and linkage effects and will be useful for quantifying the improvement of new molecular QTL studies over existing assays in capturing regulatory disease mechanisms. Moreover, partitioning mediated heritability can provide insight into regulatory effects mediated by specific gene sets or pathways.

Acknowledgments

We thank B. Pasaniuc, R. Ophoff, H. Shi, and S. Groha for helpful discussions. This research was funded by NIH grants T32HG002295, R01 MH115676, R01 MH107649, R01 MH101244, and U01 HG009379.

Author contributions

D.W.Y, L.J.O, A.L.P, and A.G conceived of the project. D.W.Y and A.G. designed experiments. D.W.Y performed the experiments and analyzed the data. D.W.Y. and A.G. wrote the manuscript with input from L.J.O. and A.L.P.

URLs

MESC software, https://github.com/douglasyao/mesc; 1000G Phase 3 data, ftp://ftp.1000genomes.ebi.ac.uk/vol1/ftp/release/20130502, baselineLD annotations, https://data.broadinstitute.org/alkesgroup/LDSCORE/; S-LDSC software, https://github.com/bulik/ldsc, GTEx v7 data, https://www.gtexportal.org/home/datasets; Macarthur lab gene sets, https://github.com/macarthur-lab/gene_lists; Molecular Signatures Database gene sets, http://software.broadinstitute.org/gsea/msigdb/collections.jsp

Online Methods

Definition of h^2_{med}

We model trait $\boldsymbol{y}$ for N individuals as follows:

$$\boldsymbol{y} = \boldsymbol{X\gamma} + \boldsymbol{XB\alpha} + \boldsymbol{\epsilon} \tag{3}$$

where $\boldsymbol{y}$ is an N-vector of phenotypes (standardized to mean 0 and variance 1), $\boldsymbol{X}$ is an $N \times M$ genotype matrix for M SNPs (standardized to mean 0 variance 1), $\boldsymbol{\gamma}$ is an M-vector of non-mediated SNP effect sizes on the trait (including pleiotropic, linkage, and trans-eQTL-mediated effects), $\boldsymbol{B}$ is an $M \times G$ matrix of cis-eQTL effect sizes *in the causal cell types/contexts* for G genes, $\boldsymbol{\alpha}$ is a G-vector of causal gene expression effect sizes on the trait, and $\boldsymbol{\epsilon}$ is an N-vector of environmental effects. We treat all variables as random. We define $h^2_{med;causal}$ as follows:

$$h^2_{med;causal} = Var[\boldsymbol{XB\alpha}] \tag{4}$$

Under the assumption that α and β are independent of each other, we can rewrite this as follows:

$$h^2_{med;causal} = E_{B,\alpha}[Var[\boldsymbol{XB\alpha} \mid \boldsymbol{B}, \boldsymbol{\alpha}]] + Var_{B,\alpha}[E[\boldsymbol{XB\alpha} \mid \boldsymbol{B}, \boldsymbol{\alpha}]] \tag{5}$$

$$= E_{B,\alpha}\left[\sum_i^G \sum_j^M \beta_{ij}^2 \alpha_i^2 \right] \tag{6}$$

$$= GE[\alpha^2]E[h^2_{cis}] \tag{7}$$

where $E[\alpha^2]$ is the average squared per-gene effect of expression on trait and $E[h^2_{cis}]$ is the average cis-heritability of expression across all genes. (6) follows because $E[\boldsymbol{XB\alpha} \mid \boldsymbol{B}, \boldsymbol{\alpha}] = 0$. We define $h^2_{nonmed;causal}$ in a similar fashion:

$$h^2_{nonmed;causal} = Var[\boldsymbol{X\gamma}] \tag{8}$$

$$= ME[\gamma^2] \tag{9}$$

where $E[\gamma^2]$ is the average squared per-SNP effect on trait that is not mediated by gene expression.

In practice, expression levels in causal cell types/contexts for the complex trait are likely not assayed. Given a set of assayed tissues T (which may or may not be causal for the complex trait), we define $h^2_{med;assayed}(T)$ as follows:

$$h^2_{med;assayed}(T) = r_g^2(T)h^2_{med;causal} \tag{10}$$

while we define $h^2_{nonmed;assayed}(T)$ as $h^2_{nonmed;causal} + (1 - r_g^2(T))h^2_{med;causal}$. Here, $r_g^2(T) = \frac{\sum_i^G \sum_j^M Cov(\beta_{ij}^2, \beta_{ij}'^2)}{\sum_i^G \sum_j^M \sqrt{Var(\beta_{ij}^2)Var(\beta_{ij}'^2)}}$ and denotes the average squared genetic correlation between expression in assayed tissues T vs. in causal cell types/contexts, where β_{ij}' is the cis-eQTL effect size of SNP j on gene i in T. Note that β_{ij}' can refer to either single tissue or meta-tissue cis-eQTL effect sizes, depending on whether T contains one or multiple tissues.

MESC with SNP effect sizes known

For illustrative purposes, we walk through a derivation for MESC in the idealized scenario that we know (1) the true eQTL effect sizes, β, of each SNP on each gene (in the causal cell types/contexts for the trait) and (2) the true phenotypic effect sizes, ω, of each SNP on y. In practice, due to LD between SNPs, these quantities are not directly estimable.

Under the generative model (3), the total effect of SNP k on the complex trait is

$$\omega_k = \sum_i^G \beta_{ik} \alpha_i + \gamma_k$$

Upon squaring ω_k we have

$$E[\omega_k^2 \mid \beta_{1k} \ldots \beta_{ik}] = \sum_i^G E[\alpha_i^2 \mid \beta_{1k} \ldots \beta_{ik}]\beta_{ik}^2 + E[\gamma_k^2 \mid \beta_{1k} \ldots \beta_{ik}] \tag{11}$$

Under certain effect size independence assumptions (see below), this simplifies to

$$E[\omega_k^2 \mid \beta_{1k} \ldots \beta_{ik}] = E[\alpha^2] \sum_i^G \beta_{ik}^2 + E[\gamma^2] \tag{12}$$

We use equation (12) to estimate $E[\alpha^2]$ by regressing ω^2 for all SNPs on $\sum_i^G \beta_i^2$ and taking the slope, while we estimate $E[\gamma^2]$ by taking the intercept. See Figure 1c for a plot illustrating this approach.

$E[\alpha^2]$ can be multiplied by $GE[h_{cis}^2]$ to obtain $h_{med;causal}^2$, while $E[\gamma^2]$ can be multiplied by M to obtain $h_{nonmed;causal}^2$. Note that violations to effect-size independence assumptions will lead to biased estimates of $h_{med;causal}^2$ and $h_{nonmed;causal}^2$, but can be remedied by stratifying SNPs and genes (see below).

Model assumptions

In this section, we describe the two main effect size independence assumptions that are needed to derive equation (12), we discuss how they might be violated, and we show that conditioning on SNP- and gene-level annotations can ameliorate any resulting bias. These assumptions are:

- Across all genes (indexed by i), the magnitude of gene effect sizes is uncorrelated with the magnitude of eQTL effect sizes (i.e. $Cov(\alpha^2, \beta^2) = 0$). We refer to this assumption as gene-eQTL effect size independence.

- Across all SNPs (indexed by k), the magnitude of non-mediated SNP effect sizes is uncorrelated with the magnitude of eQTL effect sizes (i.e. $Cov(\gamma^2, \beta^2) = 0$). We refer to this assumption as pleiotropy-eQTL effect size independence.

Violation of gene-eQTL effect size independence

Gene-eQTL effect size independence is violated in the scenario that less heritable genes have larger causal effect sizes on the trait, which is supported by evidence that evolutionarily constrained genes tend to have fewer eQTLs[5]. A negative correlation between the magnitude of gene effect sizes and eQTL effect sizes across the genome will result in downwardly biased estimates of $E[\alpha^2]$ and upwardly biased estimates of $E[\gamma^2]$, as illustrated in Supplementary Figure 23a. The downward bias in h_{med}^2 arises due to the fact that during regression, SNPs with larger eQTL effect sizes are implicitly weighted more than SNPs with smaller eQTL effect sizes, so the average slope will be biased toward the value of α^2 for high heritability genes.

In order to account for violations to gene-eQTL effect size independence, we can stratify genes by the magnitude of their cis-heritability so that within each gene category, gene-eQTL effect size independence approximately holds. We can then obtain unbiased estimates of $E[\alpha_D^2]$ for each gene category D as illustrated in Supplementary Figure 23b. In practice, we stratify genes by 5 bins according to their cis-heritability, which we show adequately captures genome-wide dependence of gene effect sizes on eQTL effect sizes in simulations (Figure 3a).

Violation of pleiotropy-eQTL effect size independence

Pleiotropy-eQTL effect size independence is violated in the presence of regulatory hotspots with high biological activity in the genome[2, 36–39], resulting in an increased number and/or magnitude of both eQTLs and pleiotropic/linkage effects in these hotspots. A positive correlation between the magnitude of non-mediated effect sizes and eQTL effect sizes across the genome will result in upwardly biased estimates of $E[\alpha^2]$ and downwardly biased estimates of $E[\gamma^2]$, as illustrated in Supplementary Figure 24a.

In order to account for violations to pleiotropy-eQTL effect size independence, we can stratify SNPs by the magnitude of their eQTL effect sizes so that within each SNP category, pleiotropy-eQTL effect size independence approximately holds. We can then obtain unbiased estimates of overall $E[\alpha^2]$ and $E[\gamma_C^2]$ for each SNP category C as illustrated in Supplementary Figure 24b. Note that in practice, LD between SNPs makes it is difficult or impossible to identify the exact SNPs that act as eQTLs. This in turn makes it impractical to stratify SNPs according to eQTL effect size. Therefore, we instead stratify SNPs by a set of comprehensive functional SNP annotations, the baselineLD model[2, 43], which should capture most known regulatory hotspots in the genome and act as a reasonable proxy to eQTL effect sizes (see Figure 3b for simulation results).

MESC with eQTL effect sizes assayed in non-causal tissues

In this section, we show that when we carry out the regression procedure described in "MESC with SNP effect sizes known" using eQTL effect sizes assayed in non-causal tissues T, we obtain an estimate of the quantity $h^2_{med;assayed}(T)$, as defined above.

Let β represent cis-eQTL effect sizes in causal cell types/contexts for the trait, and β' represent cis-eQTL effect sizes in assayed tissues T. We start with regression equation (12):

$$E[\omega_k^2] = E[\alpha^2] \sum_i^G \beta_{ik}^2 + E[\gamma^2]$$

The ordinary least squares estimate of the coefficient from regressing ω^2 on $\sum_i^G \beta'^2_i$ is

$$
\begin{aligned}
\alpha'^2 &= \frac{\sum_i^G \sum_j^M Cov(\omega_j^2, \beta'^2_{ij})}{\sum_i^G \sum_j^M Var(\beta'^2_{ij})} \\
&= \frac{\sum_i^G \sum_j^M Cov(\alpha_i^2 \beta_{ij}^2 + \gamma_j^2, \beta'^2_{ij})}{\sum_i^G \sum_j^M Var(\beta'^2_{ij})} \\
&= E[\alpha^2] \frac{\sum_i^G \sum_j^M Cov(\beta_{ij}^2, \beta'^2_{ij})}{\sum_i^G \sum_j^M Var(\beta'^2_{ij})}
\end{aligned}
$$

The third line follows given the $Cov(\gamma^2, \beta'^2) = 0$ and $Cov(\alpha^2, \beta'^2) = 0$. Let $r_g^2(T) = \frac{\sum_i^G \sum_j^M Cov(\beta_{ij}^2, \beta'^2_{ij})}{\sum_i^G \sum_j^M \sqrt{Var(\beta_{ij}^2) Var(\beta'^2_{ij})}} =$ the average squared genetic correlation between expression in T vs. in causal cell types. Given this definition, we have

$$\alpha'^2 = r_g^2(T) E[\alpha^2] \sum_i^G \sqrt{\frac{\sum_j^M Var(\beta_{ij}^2)}{\sum_j^M Var(\beta'^2_{ij})}} \tag{13}$$

For simplicity, we make the assumption that the magnitude of eQTL effect sizes is not systematically higher or lower in T vs. in causal cell types/contexts. Note that violations to this assumption can realistically occur in practice but will not bias our estimate of $h^2_{med;assayed}(T)$ (Supplementary Note). Given this assumption, we have

$$\alpha'^2 = r_g^2(T) E[\alpha^2]$$

We can then multiply α'^2 by $E[h'^2_{cis}]$ to obtain $h^2_{med;assayed}(T)$, where $E[h'^2_{cis}]$ is the average expression cis-heritability of genes in T.

MESC using summary statistics

We have previously derived an estimator for h^2_{med} in the idealized scenario that SNP effect sizes are given. In practice, we use GWAS summary statistics, which are affected by sampling noise and by LD. It has previously been shown that LD and sampling noise can be accounted for by regressing GWAS χ^2 statistics on LD scores, which measure the total of LD for each SNP[2,34]. Under our generative model (3), the marginal OLS estimate of the total effect size of a SNP k on the trait is given by

$$
\begin{aligned}
\hat{\omega}_k &= \frac{1}{N} \boldsymbol{X}_k^T \boldsymbol{y} \\
&= \frac{1}{N} (\boldsymbol{X}_k^T (\boldsymbol{X}\boldsymbol{\gamma} + \boldsymbol{X}\boldsymbol{B}\boldsymbol{\alpha} + \boldsymbol{\epsilon})) \\
&= \frac{1}{N} \boldsymbol{X}_k^T \boldsymbol{X}\boldsymbol{\gamma} + \frac{1}{N} \boldsymbol{X}_k^T \boldsymbol{X}\boldsymbol{B}\boldsymbol{\alpha} + \frac{1}{N} \boldsymbol{X}_k^T \boldsymbol{\epsilon} \\
&= \sum_j^M \gamma_j \hat{r}_{jk} + \sum_i^G \alpha_i \sum_j^M \hat{r}_{jk} \beta_{ij} + \boldsymbol{\epsilon}'
\end{aligned}
$$

Let $\hat{R} = \frac{1}{N} \boldsymbol{X}^T \boldsymbol{X}$ denote the in-sample LD matrix, and let $\boldsymbol{\epsilon}' = \frac{1}{N} \boldsymbol{X}^T \boldsymbol{\epsilon}$ denote the noise term in the summary statistics. The χ^2 statistic for SNP k (defined as $N\hat{\omega}_k^2$) is:

$$
E[\chi_k^2 \mid \hat{R}, \boldsymbol{B}] = N \sum_j^M E[\gamma_j^2 \mid \hat{R}, \boldsymbol{B}] \hat{r}_{jk}^2 + N \sum_i^G E[\alpha_i^2 \mid \hat{R}, \boldsymbol{B}] \sum_j^M \hat{r}_{jk}^2 \beta_{ij}^2 + N E[(\boldsymbol{\epsilon}')^2] \tag{14}
$$

$$
= N E[\gamma^2] \sum_j^M \hat{r}_{jk}^2 + N E[\alpha^2] \sum_i^G \sum_j^M \hat{r}_{jk}^2 \beta_{ij}^2 + N E[(\boldsymbol{\epsilon}')^2] \tag{15}
$$

$$
= \frac{N h^2_{nonmed;causal}}{M} \sum_j^M \hat{r}_{jk}^2 + \frac{N h^2_{med;causal}}{G E[h^2_{cis}]} \sum_i^G \sum_j^M \hat{r}_{jk}^2 \beta_{ij}^2 + 1 - h^2_{med;causal} - h^2_{nonmed;causal} \tag{16}
$$

In order for (15) to hold true, we must make two independence assumptions involving LD-dependent genetic architecture, in addition to the independence assumptions described in "Model assumptions" (see above). LD-dependent architecture, if not accounted for, is known to produce bias in heritability estimates[72,73]. These assumptions are:

- Across all genes (indexed by i), the magnitude of α_i is uncorrelated with the LD scores of eQTLs for gene i

- Across all SNPs (indexed by k), the magnitude of γ_k is uncorrelated with the LD score of SNP k

(16) follows (15) from our definitions of $h^2_{med;causal}$ and $h^2_{nonmed;causal}$. Since $E[\hat{r}_{jk}^2] \approx r_{jk}^2 + \frac{1}{N}$, we have

$$
E\left[\sum_j^M \hat{r}_{jk}^2 \right] \approx \sum_j^M r_{jk}^2 + \frac{M}{N}
$$

and

$$
\begin{aligned}
E\left[\sum_i^G \sum_j^M \hat{r}_{jk}^2 \beta_{ij}^2 \right] &\approx \sum_i^G \sum_j^M \left(r_{jk}^2 \beta_{ij}^2 + \frac{\beta_{ij}^2}{N} \right) \\
&\approx \sum_i^G \sum_j^M r_{jk}^2 \beta_{ij}^2 + \frac{G E[h^2_{cis}]}{N}
\end{aligned}
$$

Thus,

$$E[\chi_k^2] \approx \frac{Nh_{nonmed;causal}^2}{M}\left(\sum_j^M r_{jk}^2 + \frac{M}{N}\right) + \frac{Nh_{med;causal}^2}{GE[h_{cis}^2]}\left(\sum_i^G \sum_j^M r_{jk}^2\beta_{ij}^2 + \frac{GE[h_{cis}^2]}{N}\right) + 1 - h_{nonmed;causal}^2 - h_{med;causal}^2$$

$$\approx \frac{Nh_{nonmed;causal}^2}{M}\sum_j^M r_{jk}^2 + \frac{Nh_{med;causal}^2}{GE[h_{cis}^2]}\sum_i^G \sum_j^M r_{jk}^2\beta_{ij}^2 + 1$$

Defining LD scores $\ell_k = \sum_j^M r_{jk}^2$ and expression scores $\mathcal{L}_k = \sum_i^G \sum_j^M r_{jk}^2\beta_{ij}^2$, we arrive at our main equation for summary MESC regression:

$$E[\chi_k^2] \approx \frac{Nh_{nonmed;causal}^2}{M}\ell_k + \frac{Nh_{med;causal}^2}{GE[h_{cis}^2]}\mathcal{L}_k + 1 \tag{17}$$

Analogous to the derivation in "MESC with eQTL effect sizes assayed in non-causal tissues" (see above), we show that if we perform this regression using expression scores in assayed tissues T rather than expression scores in causal cell types/contexts, we will estimate $h_{med;assayed}^2(T)$ rather than $h_{med;causal}^2$ (see "MESC with assayed expression scores" in Supplementary Note).

Stratified MESC

In this section, we extend summary-based MESC to estimate $h_{med;assayed}^2(T)$ partitioned over groups of genes. Note that stratified MESC can be viewed as a special form of stratified LD score regression[2] (Supplementary Note). Given D potentially overlapping gene categories $\mathcal{D}_1, \ldots, \mathcal{D}_D$, we define $h_{med;causal}^2$ partitioned over gene categories as follows:

$$h_{med;causal}^2(\mathcal{D}_d) = \sum_{i \in \mathcal{D}_d} \alpha_i^2 \sum_j^M \beta_{ij}^2 \tag{18}$$

$$= |\mathcal{D}_d| \cdot E[\alpha_i^2 | i \in \mathcal{D}_d] \cdot E[h_{i;cis}^2 | i \in \mathcal{D}_d] \tag{19}$$

where $h_{med;causal}^2(\mathcal{D}_d)$ is the heritability mediated in cis through the expression of genes in category $\mathcal{D}_d$, $|\mathcal{D}_d|$ is the number of genes in $\mathcal{D}_d$, $E[\alpha_i^2 | i \in \mathcal{D}_d]$ is the average squared causal effect of expression on trait for genes in $\mathcal{D}_d$, and $E[h_{i;cis}^2 | i \in \mathcal{D}_d]$ is the average cis-heritability of expression of genes in $\mathcal{D}_d$. Similar to equation (12), equation (19) relies on an independence assumption between α and β, namely that $\alpha_i \perp \beta_i \mid i \in \mathcal{D}_d$.

For gene i, we model the variance of gene effect size α_i as

$$Var(\alpha_i) = \sum_{d:i \in \mathcal{D}_d} \pi_d$$

Assuming that gene categories $\mathcal{D}_d$ form a disjoint partition of the set of all genes, we have

$$\pi_d = \frac{E[h_{med;causal}^2(\mathcal{D}_d)]}{|\mathcal{D}_d|E[h_{i;cis}^2 | i \in \mathcal{D}_d]}$$

If gene categories are overlapping, then π_d can be conceptualized as the contribution of annotation $\mathcal{D}_d$ to $h_{med;causal}^2$ conditional on contributions from all other gene categories included in the model.

Given C potentially overlapping SNP categories $\mathcal{C}_1, \ldots, \mathcal{C}_C$, we define $h_{nonmed;causal}^2$ partitioned over SNP categories as follows:

$$h_{nonmed;causal}^2(\mathcal{C}_c) = \sum_{j \in \mathcal{C}_c} \gamma_j^2$$

$$= |\mathcal{C}_c| \cdot E[\gamma_j^2 | j \in \mathcal{C}_c]$$

where $h^2_{nonmed;causal}(\mathcal{C}_c)$ is the non-mediated heritability of SNPs in category $\mathcal{C}_c$, $|\mathcal{C}_c|$ is the number of SNPs in $\mathcal{C}_c$, and $E[\gamma_j^2|j \in \mathcal{C}_c]$ is the average squared non-mediated effect size of SNPs in $\mathcal{C}_c$.

For SNP j, we model the variance of non-mediated effect size γ_j as follows:

$$Var(\gamma_j) = \sum_{c:j\in\mathcal{C}_c} \tau_c$$

Assuming that SNP categories $\mathcal{C}_c$ form a disjoint partition of the set of all SNPs, we have

$$\tau_c = \frac{E[h^2_{nonmed;causal}(\mathcal{C}_c)]}{|\mathcal{C}_c|}$$

If SNP categories are overlapping, then τ_c can be conceptualized of as the contribution of annotation $\mathcal{C}_c$ to $h^2_{nonmed;causal}$ conditional on contributions from all other SNP categories included in the model.

The equation for stratified MESC is

$$E[\chi_k^2] = N \sum_c \tau_c \ell_{k;c} + N \sum_d \pi_d \mathcal{L}_{k;d} + 1 \tag{20}$$

where χ_k^2 is the GWAS χ^2-statistic of SNP k, N is the number of samples, $\ell_{k;c}$ is the LD score of SNP k with respect to SNP category $\mathcal{C}_c$ (defined as $\ell_{k;c} = \sum_{j\in\mathcal{C}_c} r_{jk}^2$), and $\mathcal{L}_{k;d}$ is the expression score of SNP k with respect to gene category $\mathcal{D}_d$ (defined as $\mathcal{L}_{k;d} = \sum_{i\in\mathcal{D}_d}\sum_j^M r_{jk}^2\beta_{ij}^2$). See Supplementary Note for a derivation of this equation. Analogous to unstratified MESC, when we perform this regression using expression scores in assayed tissues T rather than expression scores in causal cell types/contexts, we will estimate $h^2_{med;assayed}(T, \mathcal{D}_d) = r^2(T, \mathcal{D}_d)h^2_{med;causal}(\mathcal{D}_d)$, where $r^2(T, \mathcal{D}_d)$ is the average squared genetic correlation of expression between T and causal cell types/contexts for genes in $\mathcal{D}_d$.

Estimation of expression scores

In order to carry out the regression described in equation (20), we must first estimate expression scores $\mathcal{L}_{k;d}$ (where $\mathcal{L}_{k;d} = \sum_{i\in D}\sum_j^M r_{jk}^2\beta_{ij}^2$) from an external expression panel. $\mathcal{L}_{k;d}$ can be conceptualized as the total expression cis-heritability of genes in D that is tagged by SNP k, and is equivalent in expectation to squared eQTL summary statistics for SNP k summed across genes in D ($\sum_{i\in D} \hat{\beta}_{ik(sumstat)}^2$) modulo a constant error term that arises from squaring the sampling noise in the eQTL summary statistics. Thus, we can in theory use eQTL summary statistics to compute $\mathcal{L}_{k;d}$ and estimate h^2_{med} (see Supplementary Note).

However, because gene expression tends to have low cis-heritability across genes[5] and most individual tissue expression panels have small sample sizes (less than 500 samples per GTEx tissue), expression scores estimated using eQTL summary statistics can be very noisy and can lead to downwardly biased estimates of $h^2_{med;assayed}(T)$ due to regression attenuation[74] (Supplementary Figure 1). Therefore, we estimate individual-tissue expression scores as follows in order to reduce sampling noise. Assuming that individual level genotypes for expression panel samples are available, we leverage prior knowledge regarding the sparse nature of eQTL effect sizes and use LASSO[41] to obtain regularized estimates of causal eQTL effect sizes ($\hat{\boldsymbol{\beta}}_{LASSO}$), then multiply $\hat{\boldsymbol{\beta}}_{LASSO}^2$ by the element-wise squared LD matrix $\boldsymbol{R}^2$ to obtain $\hat{\mathcal{L}}_{k;d}$:

$$\hat{\mathcal{L}}_{k;d} = \sum_{i\in D} c_i \sum_j^M r_{jk}^2 \hat{\beta}_{ij(LASSO)}^2 \tag{21}$$

Here, c_i refers to a correction factor we apply to $\hat{\boldsymbol{\beta}}_{LASSO}$ to both reduce bias and improve prediction accuracy compared to uncorrected estimates. In particular, we observed that LASSO without correction produces downwardly-biased estimates of eQTL effect sizes in simulations (Supplementary Figure 2), leading to upwardly-biased estimates of h^2_{med} (Supplementary Figure 1). We calculate c_i as follows. First, we use REML to obtain an unbiased estimate of h^2_{cis} for each gene i ($\hat{h}_{i;cis}^2$), which in expectation is equivalent to $\sum_j^M \beta_{ij}^2$. For each gene, we then scale $\hat{\boldsymbol{\beta}}_{LASSO}^2$ by a factor c_i so that $c_i \sum_j^M \hat{\beta}_{ij(LASSO)}^2 = \hat{h}_{i;cis}^2$. We can obtain approximately unbiased estimates of the squared LD between two SNPs using the formula $r_{adj}^2 = \hat{r}^2 - \frac{1-\hat{r}^2}{N-2}$, where $\hat{r}^2$ denotes the standard biased estimator of r^2.

Other methods to estimate expression scores were also explored (Supplementary Note); we observed that the approach described in this section provided the best performance in simulations (Supplementary Figures 1,2).

Meta-analysis of expression scores

Given our method of computing expression scores in individual tissues outlined above, we meta-analyze expression scores across tissues as follows. We first obtain meta-tissue expression cis-heritability (h_{cis}^2) estimates for each gene by averaging individual-tissue h_{cis}^2 estimates across tissues. We scale individual-tissue LASSO-predicted causal eQTL effect sizes to the meta-tissue h_{cis}^2 (see above), then average the scaled causal eQTL effect sizes across tissues. Finally, we multiply the averaged causal eQTL effect sizes by the element-wise squared LD matrix to obtain expression scores. In simulations, we show that this method of meta-analyzing expression scores produces nearly unbiased estimates of h_{med}^2/h_g^2 at 5 tissues $\times$ 200 samples per tissue (Supplementary Figure 3), which is comparable to the number expression panel samples in given tissue group (Supplementary Table 2).

There are several reasons why meta-analyzing expression scores across tissues is desirable. We would like to select a set of tissues T in which eQTL effect sizes are as close as possible to those in causal cell types/contexts for the complex trait, which would maximize $r_g^2(T)$. $r_g^2(T)$ in practice refers to the squared correlation between *estimated* eQTL effect sizes in assayed tissues vs. true eQTL effect sizes in causal cell types/contexts, so sampling noise in eQTL effect size estimates will decrease $r_g^2(T)$. In simulations, we showed that 200 expression panel samples (comparable to the size of individual-tissue expression panels from GTEx) was inadequate to obtain unbiased estimates of h_{med}^2/h_g^2 (Figure 2a, Supplementary Figures 1,2). Thus, meta-analysis of expression scores across tissues is necessary in order to eliminate downward bias in $h_{med;assayed}^2(T)$ due to sampling noise in expression scores (Supplementary Figure 3). Moreover, agnostic of expression panel sample size, if meta-tissue eQTLs better tag eQTLs in causal cell types/contexts for the trait than individual tissue eQTLs, we will also increase $r_g^2(T)$ by meta-analyzing expression scores across tissues.

Simulations

All simulations were conducted using genotypes from UK Biobank[40] restricted to HapMap 3 SNPs[75] on chromosome 1 ($M = 98,499$ SNPs). All simulations followed the same overall procedure, outlined in chronological order:

1. *Simulation of expression data.* We simulated eQTL effect sizes for $G = 1,000$ genes, with eQTLs for each gene randomly selected in a 1 Mb window. We then simulated expression phenotypes for 1,000 expression panel samples (randomly selected from UK Biobank) using an additive generative model with appropriate environmental noise added, representing an expression panel. The exceptions were Figure 2a and Supplementary Figure 1, in which we varied the number of expression panel samples from 100 to 1,000, and Supplementary Figure 3, which had 200 samples per tissue.

2. *Simulation of GWAS data.* We simulated non-mediated SNP effect sizes and gene-trait effect sizes for all SNPs and genes corresponding to various levels of h_{med}^2/h_g^2. Total h_g^2 was fixed at 0.5 for all simulations. Together with the eQTL effect sizes simulated in the previous step, we used these effect sizes to simulate trait phenotypes using an additive generative model with appropriate environmental noise added for 10,000 GWAS samples (randomly selected from UK Biobank and distinct from the expression panel samples). We then produced GWAS summary statistics from this simulated data set using ordinary least squares.

3. *Estimation of expression scores.* We estimated expression scores from the expression panel samples using LASSO with REML correction (see "Estimation of expression scores" for description). For computational ease, we did not actually use REML to predict expression cis-heritability for each gene in each simulation; we instead took the true expression cis-heritability of the gene and added noise drawn from $N(0, 0.01^2)$ to simulate REML prediction error. This prediction error is consistent with empirical estimates (see Supplementary Figure 26).

4. *Estimation of h^2_{med}/h^2_g.* We estimated h^2_{med}/h^2_g using MESC with the previously estimated expression scores, in-sample LD scores (computed from the 10,000 GWAS samples), and GWAS summary statistics.

The main parameter being changed across simulations was the distribution of eQTL effect sizes, non-mediated SNP effect sizes, and gene-trait effect sizes across the genome (including various levels of sparsity and correlations between effect sizes).

Figure 2a: Impact of expression panel sample size on h^2_{med}/h^2_g estimates

For this simulation, we varied h^2_{med}/h^2_g from 0 to 1. For each gene, we selected 5 SNPs to act as cis-eQTLs with locations randomly selected within a random 1 Mb window. The average cis-heritability across all genes was set at $\overline{h^2_{cis}} = 0.05$. One eQTL for each gene was randomly selected to explain 80% of the total cis-heritability of the gene and had an effect size drawn from $\mathcal{N}(0, 0.8\overline{h^2_{cis}})$. The remaining eQTLs had effect sizes drawn from $\mathcal{N}(0, 0.2\overline{h^2_{cis}}/4)$. Expression phenotypes were simulated for each gene with environmental noise drawn from $\mathcal{N}(0, 1 - h^2_{cis})$. Non-mediated effect sizes were simulated for each SNP from $\mathcal{N}(0, (h^2_g - h^2_{med})/M)$. Gene-trait effect sizes simulated for each gene from $\mathcal{N}(0, h^2_{med}/(G\overline{h^2_{cis}}))$. Complex trait phenotypes were simulated with environmental noise drawn from $\mathcal{N}(0, 1 - h^2_g)$.

Figure 2b: Impact of sparse genetic/eQTL architectures on h^2_{med}/h^2_g estimates

We set $\overline{h^2_{cis}} = 0.05$. For the simulation involving 1 eQTL per gene, we drew its effect size from $\mathcal{N}(0, \overline{h^2_{cis}})$. For the simulation involving 10% of genes being causal, we simulated gene-trait effect sizes from a point-normal distribution, with nonzero effects drawn from $\mathcal{N}(0, h^2_{med}/(0.1G\overline{h^2_{cis}}))$. For the simulation involving 10% of SNPs being causal, we simulation non-mediated SNP effect sizes from a point-normal distribution, with nonzero effects drawn from $\mathcal{N}(0, (h^2_g - h^2_{med})/0.1M)$. The remainder of simulation parameters were the same as in Figure 2a.

Figure 2c: h^2_{med}/h^2_g estimates with $r^2_g(T) < 1$

We set $\overline{h^2_{cis}} = 0.05$. We simulated one eQTL per gene, with its effect size in the assayed tissue and causal tissue drawn from a multivariate normal distribution with mean $\mathbf{0}$ and 2×2 covariance matrix Σ with diagonal values $\Sigma_{ii} = \overline{h^2_{cis}}$ and off-diagonal values $\Sigma_{ij} = \overline{h^2_{cis}}\sqrt{\frac{1}{2}(r^2_g(T)(3 - p) - (1 - p))}$, where p represents the proportion of SNPs with non-zero eQTL effect sizes on any gene (i.e. $\frac{G}{M}$). Simulating eQTL effect sizes in this fashion results in an average squared genetic correlation of $r^2_g(T)$ across all genes. We simulated non-mediated effect sizes, expression phenotypes for the assayed tissues, and complex trait phenotypes in the same manner as in Figure 2a. eQTL effect sizes used to generate complex trait phenotypes were taken in the causal tissue. Expression scores were estimated from only the assayed expression phenotypes.

Figure 3a: Impact of violations to gene-eQTL independence on h^2_{med}/h^2_g estimates

We simulated 1 eQTL for each gene with effect size drawn from a normal distribution with mean 0 and variance $100 \cdot 2^{k/200}/(\sum_k 2^{k/200})$, where k randomly indexes the genes from 1 to 1000. Simulating eQTL effect sizes in this fashion results in a realistic continuous distribution of eQTL effect sizes, where the quintiles for expression cis-heritability across genes are 0.016, 0.032, 0.064, 0.13, and 0.26. Next, we simulated gene-trait effect sizes from a normal distribution with mean 0 and variance $h^2_{med}/(10000 \cdot 2^{k/200}/(\sum_k 2^{k/200}))$. This causes the magnitude of gene-trait effects to be strongly inversely correlated with the magnitude of eQTL effect sizes across genes, but the per-gene h^2_{med} remains constant. We simulated non-mediated effect sizes, expression phenotypes, and complex trait phenotypes in the same manner as in Figure 2a.

Figure 3b: Impact of violations to pleiotropy-eQTL independence on h^2_{med}/h^2_g estimates

We set $\overline{h^2_{cis}} = 0.05$. We simulated 1 eQTL per gene with effect size drawn from $\mathcal{N}(0, \overline{h^2_{cis}})$. All eQTLs were selected to fall in coding regions. Next, for all SNPs in coding regions, we simulated non-mediated

SNP effect sizes from $\mathcal{N}(0, (h_g^2 - h_{med}^2)/M_{coding})$, where M_{coding} is the number of SNPs that fall into coding regions. We simulated gene-trait effect sizes, expression phenotypes, and complex trait phenotypes in the same manner as in Fig 2a.

Figure 4: Comparison of MESC to heritability partitioning

We set $\overline{h_{cis}^2} = 0.05$. We simulated 1 eQTL per gene with effect size drawn from $\mathcal{N}(0, \overline{h_{cis}^2})$. To simulate 10x enrichment of eQTLs in coding, TSS, and conserved regions, we selected eQTL locations so that 10x more eQTLs per SNP were located in the three SNP categories than the remainder of the genome. We simulated non-mediated effect sizes so that the heritability enrichment of the three SNP categories was 2.5x, 5x, or 10x. We simulated expression phenotypes and complex trait phenotypes in the same manner as Figure 2a. For stratified LD-score regression, we defined the eQTL category as the set of all true eQTLs. For MESC, we stratified SNPs by the baselineLD model with the three SNP categories and windows around the SNP categories removed from the model.

Data and quality control

Genotypes

For MESC, we used European samples in 1000G[76] as reference SNPs to compute LD scores. Regression SNPs were obtained from HapMap 3[75]. SNPs with GWAS χ^2 statistics $> \max\{80, 0.001N\}$ (where N is the number of GWAS samples) and in the major histocompatibility complex (MHC) region were excluded. See Supplementary Note of ref.[2] for justification of these procedures.

For computing expression scores, we downloaded genotypes derived from sequencing data for GTEx v7 from the GTEx Portal (URLs) as described in ref.[5]. We retained SNPs that were from HapMap 3[75].

Expression data

We obtained processed and quantile normalized gene expression data for GTEx v7 from the GTEx Portal (URLs) as described in ref.[5]. For each tissue, the following covariates were included in all analyses: 3 genetic principal components, sex, platform, and 14-35 expression factors[77] as selected by the main GTEx analysis.

Estimation of expression scores from GTEx data

We used REML as implemented in GCTA[33] to estimate the expression cis-heritability for each gene in each individual GTEx tissue. We then used LASSO as implemented in PLINK[78] (with the LASSO tuning parameter set as the estimated expression cis-heritability of the gene) to estimate eQTL effect sizes for each gene in each individual GTEx tissue. In all procedures, we excluded gene-tissue pairs for which LASSO did not converge when predicting effect sizes. For Figure 5a and Supplementary Figure 8, we obtained causal eQTL effect size estimates in three different ways:

- *Meta-analysis across all tissues.* For each gene, we averaged the expression cis-heritability estimates across all 48 tissues. Within each tissue, we scaled the LASSO-predicted eQTL effect sizes to the averaged cis-heritability value. We then averaged the scaled eQTL effect sizes for each gene across all tissues. Genes were retained if they had a LASSO-converged eQTL effect size in at least one tissue.

- *Meta-analysis in tissue groups.* Of the 48 tissues, we grouped together 37 of them into 7 broad tissue groups: adipose, blood/immune, cardiovascular, CNS, digestive, endocrine, and skin (Supplementary Table 2). Within each tissue group, we averaged the expression cis-heritability estimates for each gene and scaled the LASSO-predicted eQTL effect sizes to the averaged cis-heritability value. We then averaged the scaled eQTL effect sizes for each gene across the tissues for each tissue group. Genes were retained in each tissue group if they had a LASSO-converged eQTL effect size in at least one tissue within that tissue group.

- *Individual tissues.* For each individual tissue, we scaled the LASSO-predicted eQTL effect sizes to the within-tissue-group averaged cis-heritability estimates.

The final eQTL effect sizes were then multiplied by the element-wise squared LD matrix in order to obtain expression scores (see "Estimation of expression scores").

All remaining results in the manuscript, including Figures 6,7a and Supplementary Figures 11,12,13,18,19,20, were obtained from expression scores meta-analyzed across all 48 GTEx tissues. The only exceptions were the tissue-specific expression analyses in Figure 7c and Supplementary Figure 21, in which we meta-analyzed expression scores within the tissue group of the focal tissue.

Set of 42 independent traits

Analogous to previous studies[19,79], we initially considered a set of 34 traits from publicly available sources and 55 traits from UK Biobank for which GWAS summary statistics had been computed using BOLT-LMM v2.3[80,81]. We restricted our analysis to 47 traits with z-scores of total SNP heritability above 6 (computed using stratified LD-score regression). The 47 traits included 5 traits that were duplicated across two datasets (genetic correlation of at least 0.9). For duplicated traits, we retained the data set with the larger sample size, leaving us with a total of 42 independent traits. When meta-analyzing estimates across traits, we performed random effects meta-analysis using the R package rmeta.

BaselineLD categories

In all our analyses, we stratified SNPs by 72 functional categories specified by the baselineLD model v2.0[2,43] (URLs). These annotations include coding, conserved, regulatory (e.g., promoter, enhancer, histone marks, transcription factor binding sites), and LD-related annotations. The original baselineLD model v2.0 contains 76 total categories; we removed 4 categories corresponding to QTL MaxCPP annotations[19] because the information contained in these annotations is redundant with the eQTL effect size information contained in expression scores.

Gene set analyses

In order for us to obtain to unbiased estimates of h^2_{med} enrichment for the gene sets in our analysis, we must ensure that the gene-eQTL effect size independence assumption holds within each gene set (see "Model assumptions" above). Thus, in order to capture potential correlations between the magnitude of eQTL effect sizes and gene-trait effect sizes within gene sets, we partitioned each gene set into three separate bins based on the magnitude of their expression cis-heritability relative to other genes in the gene set. We then estimated $h^2_{med}(D)$ for each individual bin and aggregated these values together to estimate the overall h^2_{med} enrichment of the gene set.

Broad gene sets

We obtained gene sets corresponding to all coding genes, genes near significant GWAS hits in the NHGRI GWAS catalog[55], genes essential in mice[52–54], genes essential in cultured cell lines[82], genes with any disease association in ClinVar[83], and genes that are FDA-approved drug targets[51] from the Macarthur lab GitHub page (URLs). We obtained an additional gene set for genes essential in cell lines[84], genes depleted for protein-truncating mutations[45,46], and genes depleted for missense mutations[85] from the supplementary data of the respective papers.

Pathway gene sets

We initially considered a set of 7,246 gene sets from the "canonical pathways" and "GO gene sets" collections from the Molecular Signatures Database[86] (URLs), consisting of gene sets from BioCarta, Reactome, KEGG, GO, PID, and other sources. We restricted our analysis to 780 gene sets for which the number of genes with LASSO estimates of eQTL effect sizes that converged in individual GTEx tissues was at least 100 when averaged across all individual tissues. (Note that this roughly corresponds to gene sets with greater than 200 total genes; see Supplementary Table 5).

Tissue specific expression gene sets

We initially considered the full set of 48 GTEx tissues. We restricted our analysis to 37 gene sets for which the focal tissue belonged to one of the 7 main tissue groups we defined in our previous analyses (Supplementary Table 2). From ref.[50], we obtained the set of 10% most specifically expressed genes in each of the 37 tissues.

References

[1] Maurano, M. T. *et al.* Systematic Localization of Common Disease-Associated Variation in Regulatory DNA. *Science* **337**, 1190–1195, DOI: 10.1126/science.1222794 (2012).

[2] Finucane, H. K. *et al.* Partitioning heritability by functional annotation using genome-wide association summary statistics. *Nat. Genet.* **47**, 1228–1235, DOI: 10.1038/ng.3404 (2015).

[3] Visscher, P. M. *et al.* 10 Years of GWAS Discovery: Biology, Function, and Translation. *The Am. J. Hum. Genet.* **101**, 5–22, DOI: 10.1016/j.ajhg.2017.06.005 (2017).

[4] Stunnenberg, H. G. *et al.* The International Human Epigenome Consortium: A Blueprint for Scientific Collaboration and Discovery. *Cell* **167**, 1145–1149, DOI: 10.1016/j.cell.2016.11.007 (2016).

[5] GTEx Consortium. Genetic effects on gene expression across human tissues. *Nature* **550**, 204–213, DOI: 10.1038/nature24277 (2017).

[6] Giambartolomei, C. *et al.* Bayesian Test for Colocalisation between Pairs of Genetic Association Studies Using Summary Statistics. *PLOS Genet.* **10**, e1004383, DOI: 10.1371/journal.pgen.1004383 (2014).

[7] Farh, K. K.-H. *et al.* Genetic and epigenetic fine mapping of causal autoimmune disease variants. *Nature* **518**, 337–343, DOI: 10.1038/nature13835 (2015).

[8] Hormozdiari, F. *et al.* Colocalization of GWAS and eQTL Signals Detects Target Genes. *The Am. J. Hum. Genet.* **99**, 1245–1260, DOI: 10.1016/j.ajhg.2016.10.003 (2016).

[9] Chun, S. *et al.* Limited statistical evidence for shared genetic effects of eQTLs and autoimmune-disease-associated loci in three major immune-cell types. *Nat. Genet.* **49**, 600, DOI: 10.1038/ng.3795 (2017).

[10] Ongen, H. *et al.* Estimating the causal tissues for complex traits and diseases. *Nat. Genet.* **49**, 1676–1683, DOI: 10.1038/ng.3981 (2017).

[11] Gamazon, E. R. *et al.* A gene-based association method for mapping traits using reference transcriptome data. *Nat. Genet.* **47**, 1091–1098, DOI: 10.1038/ng.3367 (2015).

[12] Gusev, A. *et al.* Integrative approaches for large-scale transcriptome-wide association studies. *Nat. Genet.* **48**, 245–252, DOI: 10.1038/ng.3506 (2016).

[13] Zhu, Z. *et al.* Integration of summary data from GWAS and eQTL studies predicts complex trait gene targets. *Nat. Genet.* **48**, 481–487, DOI: 10.1038/ng.3538 (2016).

[14] Barbeira, A. N. *et al.* Exploring the phenotypic consequences of tissue specific gene expression variation inferred from GWAS summary statistics. *Nat. Commun.* **9**, 1825, DOI: 10.1038/s41467-018-03621-1 (2018).

[15] Barfield, R. *et al.* Transcriptome-wide association studies accounting for colocalization using Egger regression. *Genet. Epidemiol.* **42**, 418–433, DOI: 10.1002/gepi.22131 (2018).

[16] Porcu, E. *et al.* Mendelian randomization integrating GWAS and eQTL data reveals genetic determinants of complex and clinical traits. *Nat. Commun.* **10**, 1–12, DOI: 10.1038/s41467-019-10936-0 (2019).

[17] Davis, L. K. *et al.* Partitioning the Heritability of Tourette Syndrome and Obsessive Compulsive Disorder Reveals Differences in Genetic Architecture. *PLOS Genet.* **9**, e1003864, DOI: 10.1371/journal.pgen.1003864 (2013).

Torres, J. *et al.* Cross-Tissue and Tissue-Specific eQTLs: Partitioning the Heritability of a Complex Trait. *The Am. J. Hum. Genet.* **95**, 521–534, DOI: 10.1016/j.ajhg.2014.10.001 (2014).

Hormozdiari, F. *et al.* Leveraging molecular quantitative trait loci to understand the genetic architecture of diseases and complex traits. *Nat. Genet.* **50**, 1041, DOI: 10.1038/s41588-018-0148-2 (2018).

Gamazon, E. R. *et al.* Using an atlas of gene regulation across 44 human tissues to inform complex disease- and trait-associated variation. *Nat. Genet.* **50**, 956, DOI: 10.1038/s41588-018-0154-4 (2018).

Hemani, G. *et al.* The MR-Base platform supports systematic causal inference across the human phenome. *eLife* **7**, e34408, DOI: 10.7554/eLife.34408 (2018).

Mancuso, N. *et al.* Large-scale transcriptome-wide association study identifies new prostate cancer risk regions. *Nat. Commun.* **9**, 4079, DOI: 10.1038/s41467-018-06302-1 (2018).

Wu, L. *et al.* A transcriptome-wide association study of 229,000 women identifies new candidate susceptibility genes for breast cancer. *Nat. Genet.* **50**, 968, DOI: 10.1038/s41588-018-0132-x (2018).

Wray, N. R. *et al.* Genome-wide association analyses identify 44 risk variants and refine the genetic architecture of major depression. *Nat. Genet.* **50**, 668, DOI: 10.1038/s41588-018-0090-3 (2018).

Gandal, M. J. *et al.* Transcriptome-wide isoform-level dysregulation in ASD, schizophrenia, and bipolar disorder. *Science* **362**, eaat8127, DOI: 10.1126/science.aat8127 (2018).

Gusev, A. *et al.* A transcriptome-wide association study of high-grade serous epithelial ovarian cancer identifies new susceptibility genes and splice variants. *Nat. Genet.* **51**, 815, DOI: 10.1038/s41588-019-0395-x (2019).

Huckins, L. M. *et al.* Gene expression imputation across multiple brain regions provides insights into schizophrenia risk. *Nat. Genet.* **51**, 659, DOI: 10.1038/s41588-019-0364-4 (2019).

Gamazon, E. R., Zwinderman, A. H., Cox, N. J., Denys, D. & Derks, E. M. Multi-tissue transcriptome analyses identify genetic mechanisms underlying neuropsychiatric traits. *Nat. Genet.* **51**, 933, DOI: 10.1038/s41588-019-0409-8 (2019).

Wainberg, M. *et al.* Opportunities and challenges for transcriptome-wide association studies. *Nat. Genet.* **51**, 592, DOI: 10.1038/s41588-019-0385-z (2019).

Mancuso, N. *et al.* Probabilistic fine-mapping of transcriptome-wide association studies. *Nat. Genet.* **51**, 675, DOI: 10.1038/s41588-019-0367-1 (2019).

Liu, B., Gloudemans, M. J., Rao, A. S., Ingelsson, E. & Montgomery, S. B. Abundant associations with gene expression complicate GWAS follow-up. *Nat. Genet.* **51**, 768, DOI: 10.1038/s41588-019-0404-0 (2019).

Hemani, G., Bowden, J. & Davey Smith, G. Evaluating the potential role of pleiotropy in Mendelian randomization studies. *Hum. Mol. Genet.* **27**, R195–R208, DOI: 10.1093/hmg/ddy163 (2018).

Yang, J., Lee, S. H., Goddard, M. E. & Visscher, P. M. GCTA: A Tool for Genome-wide Complex Trait Analysis. *Am. J. Hum. Genet.* **88**, 76–82, DOI: 10.1016/j.ajhg.2010.11.011 (2011).

Bulik-Sullivan, B. K. *et al.* LD Score regression distinguishes confounding from polygenicity in genome-wide association studies. *Nat. Genet.* **47**, 291–295, DOI: 10.1038/ng.3211 (2015).

O'Connor, L. J. *et al.* Polygenicity of complex traits is explained by negative selection. *bioRxiv* 420497, DOI: 10.1101/420497 (2018).

Albert, F. W. & Kruglyak, L. The role of regulatory variation in complex traits and disease. *Nat. Rev. Genet.* **16**, 197–212, DOI: 10.1038/nrg3891 (2015).

[37] Albert, F. W., Bloom, J. S., Siegel, J., Day, L. & Kruglyak, L. Genetics of trans-regulatory variation in gene expression. *eLife* **7**, e35471, DOI: 10.7554/eLife.35471 (2018).

[38] Yao, C. *et al.* Dynamic Role of trans Regulation of Gene Expression in Relation to Complex Traits. *The Am. J. Hum. Genet.* **100**, 571–580, DOI: 10.1016/j.ajhg.2017.02.003 (2017).

[39] Boyle, E. A., Li, Y. I. & Pritchard, J. K. An Expanded View of Complex Traits: From Polygenic to Omnigenic. *Cell* **169**, 1177–1186, DOI: 10.1016/j.cell.2017.05.038 (2017).

[40] Bycroft, C. *et al.* The UK Biobank resource with deep phenotyping and genomic data. *Nature* **562**, 203, DOI: 10.1038/s41586-018-0579-z (2018).

[41] Tibshirani, R. Regression Shrinkage and Selection via the Lasso. *J. Royal Stat. Soc. Ser. B (Methodological)* **58**, 267–288 (1996).

[42] Robinson, G. K. That BLUP is a Good Thing: The Estimation of Random Effects. *Stat. Sci.* **6**, 15–32, DOI: 10.1214/ss/1177011926 (1991).

[43] Gazal, S. *et al.* Linkage disequilibrium–dependent architecture of human complex traits shows action of negative selection. *Nat. Genet.* **49**, 1421–1427, DOI: 10.1038/ng.3954 (2017).

[44] Liu, X. *et al.* Functional Architectures of Local and Distal Regulation of Gene Expression in Multiple Human Tissues. *The Am. J. Hum. Genet.* **100**, 605–616, DOI: 10.1016/j.ajhg.2017.03.002 (2017).

[45] Lek, M. *et al.* Analysis of protein-coding genetic variation in 60,706 humans. *Nature* **536**, 285–291, DOI: 10.1038/nature19057 (2016).

[46] Cassa, C. A. *et al.* Estimating the selective effects of heterozygous protein-truncating variants from human exome data. *Nat. Genet.* **49**, 806–810, DOI: 10.1038/ng.3831 (2017).

[47] Kanehisa, M., Furumichi, M., Tanabe, M., Sato, Y. & Morishima, K. KEGG: new perspectives on genomes, pathways, diseases and drugs. *Nucleic Acids Res.* **45**, D353–D361, DOI: 10.1093/nar/gkw1092 (2017).

[48] Fabregat, A. *et al.* The Reactome Pathway Knowledgebase. *Nucleic Acids Res.* **46**, D649–D655, DOI: 10.1093/nar/gkx1132 (2018).

[49] Ashburner, M. *et al.* Gene Ontology: tool for the unification of biology. *Nat. Genet.* **25**, 25, DOI: 10.1038/75556 (2000).

[50] Finucane, H. K. *et al.* Heritability enrichment of specifically expressed genes identifies disease-relevant tissues and cell types. *Nat. Genet.* **50**, 621, DOI: 10.1038/s41588-018-0081-4 (2018).

[51] Wishart, D. S. *et al.* DrugBank 5.0: a major update to the DrugBank database for 2018. *Nucleic Acids Res.* **46**, D1074–D1082, DOI: 10.1093/nar/gkx1037 (2018).

[52] Blake, J. A. *et al.* The Mouse Genome Database (MGD): premier model organism resource for mammalian genomics and genetics. *Nucleic Acids Res.* **39**, D842–848, DOI: 10.1093/nar/gkq1008 (2011).

[53] Georgi, B., Voight, B. F. & Bućan, M. From Mouse to Human: Evolutionary Genomics Analysis of Human Orthologs of Essential Genes. *PLOS Genet.* **9**, e1003484, DOI: 10.1371/journal.pgen.1003484 (2013).

[54] Liu, X., Jian, X. & Boerwinkle, E. dbNSFP v2.0: A Database of Human Non-synonymous SNVs and Their Functional Predictions and Annotations. *Hum. Mutat.* **34**, E2393–E2402, DOI: 10.1002/humu.22376 (2013).

[55] MacArthur, J. *et al.* The new NHGRI-EBI Catalog of published genome-wide association studies (GWAS Catalog). *Nucleic Acids Res.* **45**, D896–D901, DOI: 10.1093/nar/gkw1133 (2017).

[56] Strober, B. J. *et al.* Dynamic genetic regulation of gene expression during cellular differentiation. *Science* **364**, 1287–1290, DOI: 10.1126/science.aaw0040 (2019).

Wijst, M. G. P. v. d. *et al.* Single-cell RNA sequencing identifies celltype-specific cis-eQTLs and co-expression QTLs. *Nat. Genet.* **50**, 493, DOI: 10.1038/s41588-018-0089-9 (2018).

Li, Y. I. *et al.* RNA splicing is a primary link between genetic variation and disease. *Science* **352**, 600–604, DOI: 10.1126/science.aad9417 (2016).

Delaneau, O. *et al.* Chromatin three-dimensional interactions mediate genetic effects on gene expression. *Science* **364**, eaat8266, DOI: 10.1126/science.aat8266 (2019).

Võsa, U. *et al.* Unraveling the polygenic architecture of complex traits using blood eQTL metaanalysis. *bioRxiv* 447367, DOI: 10.1101/447367 (2018).

Hu, Y. *et al.* A statistical framework for cross-tissue transcriptome-wide association analysis. *Nat. Genet.* **51**, 568, DOI: 10.1038/s41588-019-0345-7 (2019).

Price, A. L. *et al.* Single-Tissue and Cross-Tissue Heritability of Gene Expression Via Identity-by-Descent in Related or Unrelated Individuals. *PLOS Genet.* **7**, e1001317, DOI: 10.1371/journal.pgen.1001317 (2011).

Rossi, A. *et al.* Genetic compensation induced by deleterious mutations but not gene knockdowns. *Nature* **524**, 230–233, DOI: 10.1038/nature14580 (2015).

Wang, X. & Goldstein, D. B. Enhancer redundancy predicts gene pathogenicity and informs complex disease gene discovery. *bioRxiv* 459123, DOI: 10.1101/459123 (2018).

Valverde, P., Healy, E., Jackson, I., Rees, J. L. & Thody, A. J. Variants of the melanocyte–stimulating hormone receptor gene are associated with red hair and fair skin in humans. *Nat. Genet.* **11**, 328, DOI: 10.1038/ng1195-328 (1995).

Sulem, P. *et al.* Genetic determinants of hair, eye and skin pigmentation in Europeans. *Nat. Genet.* **39**, 1443–1452, DOI: 10.1038/ng.2007.13 (2007).

Sulem, P. *et al.* Two newly identified genetic determinants of pigmentation in Europeans. *Nat. Genet.* **40**, 835–837, DOI: 10.1038/ng.160 (2008).

Kenny, E. E. *et al.* Melanesian Blond Hair Is Caused by an Amino Acid Change in TYRP1. *Science* **336**, 554–554, DOI: 10.1126/science.1217849 (2012).

O'Brien, H. E. *et al.* Expression quantitative trait loci in the developing human brain and their enrichment in neuropsychiatric disorders. *Genome Biol.* **19**, 194, DOI: 10.1186/s13059-018-1567-1 (2018).

Walker, R. L. *et al.* Genetic control of gene expression and splicing in the developing human brain. *bioRxiv* 471193, DOI: 10.1101/471193 (2018).

O'Connor, L. J. *et al.* Estimating the proportion of disease heritability mediated by gene expression levels. *bioRxiv* 118018, DOI: 10.1101/118018 (2017).

Speed, D. *et al.* Reevaluation of SNP heritability in complex human traits. *Nat. Genet.* **49**, 986–992, DOI: 10.1038/ng.3865 (2017).

Gazal, S., Marquez-Luna, C., Finucane, H. K. & Price, A. L. Reconciling S-LDSC and LDAK functional enrichment estimates. *Nat. Genet.* **51**, 1202–1204, DOI: 10.1038/s41588-019-0464-1 (2019).

Spearman, C. The Proof and Measurement of Association between Two Things. *The Am. J. Psychol.* **15**, 72–101, DOI: 10.2307/1412159 (1904).

The International HapMap 3 Consortium. Integrating common and rare genetic variation in diverse human populations. *Nature* **467**, 52–58, DOI: 10.1038/nature09298 (2010).

The 1000 Genomes Project Consortium. A global reference for human genetic variation. *Nature* **526**, 68–74, DOI: 10.1038/nature15393 (2015).

[77] Stegle, O., Parts, L., Piipari, M., Winn, J. & Durbin, R. Using probabilistic estimation of expression residuals (PEER) to obtain increased power and interpretability of gene expression analyses. *Nat. Protoc.* **7**, 500–507, DOI: 10.1038/nprot.2011.457 (2012).

[78] Purcell, S. *et al.* PLINK: A Tool Set for Whole-Genome Association and Population-Based Linkage Analyses. *Am. J. Hum. Genet.* **81**, 559–575 (2007).

[79] Kim, S. S. *et al.* Genes with High Network Connectivity Are Enriched for Disease Heritability. *The Am. J. Hum. Genet.* **104**, 896–913, DOI: 10.1016/j.ajhg.2019.03.020 (2019).

[80] Loh, P.-R. *et al.* Efficient Bayesian mixed-model analysis increases association power in large cohorts. *Nat. Genet.* **47**, 284–290, DOI: 10.1038/ng.3190 (2015).

[81] Loh, P.-R., Kichaev, G., Gazal, S., Schoech, A. P. & Price, A. L. Mixed-model association for biobank-scale datasets. *Nat. Genet.* **50**, 906, DOI: 10.1038/s41588-018-0144-6 (2018).

[82] Hart, T. *et al.* Evaluation and Design of Genome-Wide CRISPR/SpCas9 Knockout Screens. *G3: Genes, Genomes, Genet.* **7**, 2719–2727, DOI: 10.1534/g3.117.041277 (2017).

[83] Landrum, M. J. *et al.* ClinVar: public archive of relationships among sequence variation and human phenotype. *Nucleic Acids Res.* **42**, D980–D985, DOI: 10.1093/nar/gkt1113 (2014).

[84] Wang, T. *et al.* Identification and characterization of essential genes in the human genome. *Science* **350**, 1096–1101, DOI: 10.1126/science.aac7041 (2015).

[85] Samocha, K. E. *et al.* A framework for the interpretation of *de novo* mutation in human disease. *Nat. Genet.* **46**, 944–950, DOI: 10.1038/ng.3050 (2014).

[86] Liberzon, A. *et al.* Molecular signatures database (MSigDB) 3.0. *Bioinformatics* **27**, 1739–1740, DOI: 10.1093/bioinformatics/btr260 (2011).