High-throughput analysis of adaptation using barcoded strains of *Saccharomyces cerevisiae*

Vincent J. Fasanello*, Ping Liu^, Carlos A. Botero†[a], Justin C. Fay‡^[a]

* Division of Biology and Biomedical Sciences; Washington University School of Medicine in St.

Louis; St. Louis, Missouri; 63110

^Department of Genetics; Washington University in St. Louis; St. Louis, Missouri; 63110

† Department of Biology; Washington University in St. Louis; St. Louis, Missouri; 63130

‡ University of Rochester; Rochester, New York; 14627

[a]These authors share senior authorship

20   **Running Title:** Barcoded Experimental Evolution

21   **Keywords:** Experimental Evolution, *Saccharomyces cerevisiae*, Genetic Barcode, Fitness Assay

22   **Corresponding Author:** Vincent J. Fasanello

23   ABSTRACT

Background: Experimental evolution of microbes can be used to empirically address a wide

range of questions about evolution and is increasingly employed to study complex phenomena

ranging from genetic evolution to evolutionary rescue. Regardless of experimental aims, fitness

assays are a central component of this type of research, and low-throughput often limits the

scope and complexity of experimental evolution studies. We created an experimental evolution

system in *Saccharomyces cerevisiae* that utilizes genetic barcoding to overcome this challenge.

Results: We first confirm that barcode insertions do not alter fitness in this system and can be

used to efficiently detect fitness differences via pooled competition-based fitness assays. Next,

we examine the effects of ploidy, chemical stress, and population bottleneck size on the

evolutionary dynamics and fitness gains (adaptation) in a total of 76 experimentally evolving

populations by conducting 2,136 fitness assays and analyzing 532 longitudinal-evolutionary

samples collected from the evolving populations. In our analysis of these data we describe the

strengths of this experimental evolution system and explore sources or error in our

measurements of fitness and evolutionary dynamics.

Conclusions: Our experimental treatments generated distinct fitness effects and evolutionary

dynamics, respectively quantified via multiplexed fitness assays and barcode lineage tracking.

These findings demonstrate the utility of this new resource for designing and improving high-

throughput studies of experimental evolution. The approach described here provides a

framework for future studies employing experimental designs that require high-throughput

multiplexed fitness measurements.

                                          BACKGROUND

Experimental evolution in microorganisms such as yeast, bacteria, and viruses has been used to

answer evolutionary questions that are experimentally intractable in organisms with longer

generation times [1–4]. A central benefit of most experimental evolution systems is the ability

to replicate and repeat evolution as well as the ability to store and compete evolved,

intermediate, and ancestral strains [4]. Indeed, if there is a single unifying theme to what we

have learned from experimental evolution it is that adaptation is universal and often repeatable

due to parallel changes down to the molecular level [5–9]. The power of this approach has led

to numerous investigations of the effects of population size [6,10,11] and structure [12–14],

mutation rate [15–18], and various environmental treatments [19–24].

Measuring adaptation by changes in fitness is a core requirement of experimental evolution

that often limits its implementation. Fitness assays involve direct competition of individuals or

populations against one another or a common reference, and traditionally have been calculated

using neutral markers scored by plating assays. For example, the long-term experimental

evolution conducted by Lenski and colleagues is scored by counting colonies that can ferment

arabinose based on colony color [25]. Many recent studies use fluorescently marked strains

such that the fitness of co-cultured strains can be assayed by flow-cytometry [e.g., 10,26,27].

However, the number of fitness assays, and thereby the resolution of those assays, remain

limited, which poses a challenge for large-scale projects. The primary constraint limiting the

scale of these studies is that measuring fitness requires replicate assays, often under a variety

of conditions. For example, an experiment with 100 strains evolved in a single environment

would require 800 fitness assays if one were to measure, in quadruplicate, the fitness of each

ancestral strain and each evolved strain in relation to a common reference. Furthermore, each

of the fitness assays would require the frequency of the two strains to be quantified at the

beginning and end of the competition in order to calculate fitness, resulting in 1600 individual

samples for a comprehensive look at fitness change in the 100 strains in question. Equally

important is the ability to detect small changes in fitness, which is directly related to the

number of replicate assays per strain and the noise of the assay itself.

Genetic barcodes can vastly increase the throughput of experimental evolution analyses

through pooled-fitness assay designs. In a barcoding approach, each strain is marked by

insertion of a unique neutral DNA sequence (i.e., a barcode) into its genome, enabling the easy

quantification of the relative abundance of multiple strains when simultaneously competed

against a common reference. Microarrays [28] and more recently, direct sequencing of

barcodes [29], have been used to measure the effects of thousands of single gene deletions on

fitness using this approach. Barcodes have also been used for lineage tracking in experimental

evolution. For example, Blundell and co-workers utilized a pool of half a million barcoded yeast

strains to detect and track the fate of lineages in large populations [30,31]. A further advantage

of this approach is that the barcodes can be used to measure the fitness of strains that have

been isolated from an evolved population in a traditional competition assay [32,33].

In this study we describe a set of barcoded yeast strains and characterize their utility for

assessing adaptation during experimental evolution. The system is composed of a collection of

Saccharomyces cerevisiae strains, individually barcoded with unique 20 bp sequences inserted

upstream of the *HO* locus; this barcoding strategy offers a number of advantages for

experimental evolution. First, cross-contamination between populations with different

barcodes can be detected and monitored. Second, populations can be initiated with mixtures of

multiple barcodes for tracking adaptive dynamics that occur during the experimental evolution

period [27,34]. Finally, fitness can be measured from the entire set of strains in a single pooled

fitness assay, which dramatically increases efficiency. To demonstrate the capabilities of this

system we began by conducting a series of proof-of-concept fitness assays and subsequently

applied what we learned in a short, 25-day (~250-generation) experimental evolution. These

analyses confirm that barcode insertions (1) do not alter the fitness of our source strains, and

(2) provide a high-throughput means of measuring fitness differences and evolutionary

dynamics for individual lineages from pooled samples obtained at different stages of

experimental evolution. We highlight the advantages of multiplexing samples with indexing as

well as the importance of limiting molecular contamination between initial sampling and library

construction. Taken together, this system represents a valuable resource for designing and

improving high-throughput studies in experimental evolution.

107   METHODS

Strains, media and culture methods

Barcoded yeast strains were constructed using two haploid derivatives of a diploid strain

collected from an Oak tree in Pennsylvania (YPS163) [35]: YJF153 (MATa, HO::dsdAMX4) and

YJF154 (MATalpha, HO::dsdAMX4). Barcoded kanMX deletion cassettes were amplified from

the MoBY plasmid collection [36] with primers containing homology to the promoter region (-

1,129 to -1,959) of HO. A set of 184 barcoded cassettes were selected based on confirmation of

correct barcodes by sequencing (Additional file 1 contains entries for barcodes used in this

project), then transformed into YJF153 and confirmed by PCR. Barcoded diploid strains were

made by mating barcoded haploids (YJF153) to YJF154, and confirming diploids by mating-type

PCR [37]. The final barcoded yeast strain library consisted of 184 haploid and 184 diploid

strains; 92 strains from this set were utilized in the proof-of-concept fitness assays and 77

strains from the set were utilized in the 250-generation experiment. Strains were stored at -

80°C as 15% glycerol stocks.

All evolution and fitness assays were conducted in complete medium (CM; 2 g/l dextrose, 0.17

g/l yeast nitrogen base without amino acid and ammonium sulfate, 0.5 g/l ammonium sulfate,

0.13 g/l dropout mix complete without yeast nitrogen base) with or without additional stresses

in 96-deep well plates (2.2-ml poly-propylene plates, square well, v-conical bottom; Abgene AB-

0932) covered with rayon acrylate breathable membranes (Thermo Scientific, 1256705).

Growth plates were incubated at 30°C for 24 hours inside an incubator (VWR, Forced Air

Incubator, basic, 120v, 7 cu. ft.) with agitation using a horizontal electromagnetic microplate

shaker (Union Scientific LLC, 9779-TC). Saturated 24-hour culture was diluted (1:1000) into

fresh medium at the same time each day to initialize the next round of growth for all evolution

and fitness assays.

Starting material for all evolution and fitness assays originated from -80°C freezer stocks of the

constructed barcoded yeast strains. Yeast were revived from -80°C freezer stocks via a single

round of growth (1 day, 10 generations) under standard culture conditions for these assays.

Samples collected during our experiments were stored both as (1) 15% glycerol stocks at -80°C

to maintain viable freezer stocks of yeast populations [i.e., a "frozen time vault", sensu 4], and

(2) pelleted samples at -20°C for DNA extraction.

Experimental design

Proof-of-concept fitness assays: The design of this new system for experimental evolution

began with a proof-of-concept analysis that allowed us to optimize our methods and to confirm

that the fitness of multiple strains in pooled samples could be simultaneously and accurately

measured with sequencing-based competition assay methods. This initial step involved

measuring the fitness of 91 barcoded yeast strains relative to an ancestral reference strain

simultaneously using a sequencing-based fitness assay (Figure 1, A.). Ten replicate fitness assays

were conducted under standard culture conditions (see *Strains, media and culture methods*).

Briefly, 92 yeast strains were revived from -80°C freezer stocks, mixed in equal proportions (i.e.,

such that each strain comprised 1/92 of the pooled population), and diluted (1:1000) into fresh

medium to initialize the two-day proof-of-concept fitness assays. Samples were obtained from

the undiluted initial mixtures and, 20 generations later, from the final overnight population

cultures. Fitness was measured by the change in barcode abundance relative to a 'reference'

strain (d1H10) from 10 replicates over the two-day period of approximately 20 generations, for

a total of 920 fitness assays (92 focal barcodes x 10 replicates). See *Fitness calculations*, below,

for a full description of the competition-based fitness assay methodology and for calculations of

fitness from barcode abundance data. DNA was isolated separately for each fitness assay

sample using a ZR Fungal/Bacterial DNA Kit (Zymo Research D6005) in individual 2.0 mL screw-

cap tubes following the manufacturer's instructions. Physical cell disruption by bead-beating

was carried out in a mixer mill (Retsch, MM 300) at 30 Hz (1800 min^{-1}) for ten minutes (1-

minute on, 1-minute off, times ten cycles). Following extraction, DNA was amplified with

forward/reverse primers containing a 9-12 bp index for multiplex sequencing. PCR products

were quantified, pooled and purified to form a single multiplexed library for sequencing.

Additional control samples were also included in the library to track barcode cross-

contamination. See *Library construction and sequencing*, below, for a detailed description of

the library preparation protocol used for these and all other samples.

250-Generation evolution experiment: After establishing the feasibility of the analysis strategy

and optimizing culture, assay, and processing methods, 152 yeast strains were evolved for 25

days (i.e., ca. 250 generations at 9.97 generations per day) under different scenarios of

selection in a second set of experiments. Specifically, these yeast strains were evolved by serial

dilution in one of six different treatments (Figure 1, B1.; See Additional file 2 for treatment

descriptions). Evolutionary treatments involved growth in either complete media (CM), CM

with ethanol (8% by volume) or CM with NaCl (0.342 M). Serial transfers were achieved either

through standard dilution (1:1000), reduced dilution (1:250), or increased dilution (1:4000).

Note that the 1:250 dilution treatment and the 1:4000 dilution treatment are expected to have

undergone 200 and 300 generations of evolution, respectively, rather than the 250 generations

of evolution expected in the treatments with a standard 1:1000 dilution. This difference in

number of generations is caused by variation in the number of possible doubling events when

different numbers of cells are added to media with the same (limiting) amount of glucose.

Additionally, haploid and diploid yeast were evolved under standard culture conditions (see

Strains, media and culture methods) to assess the effects of ploidy. To initialize this evolution

experiment, barcoded yeast strains were revived from -80°C freezer stocks. Barcoded yeast

pairs slated to evolve in competition were then mixed in equal proportions and diluted in fresh

medium, according to the experimental design, and allowed to evolve for 25 days.

Samples were obtained from the initial undiluted mixtures and on day 25 (ca. 250 generations

later) to use, respectively, as the starting material for the Generation-0 and the Generation-250

fitness assays . These end-point assays of relative fitness were developed to characterize fitness

outcomes in the 250-generation experiment (See *End-point assay of relative fitness change*).

Additional samples were obtained at regular intervals to characterize evolutionary dynamics

(See *Evolutionary dynamics*). These two sets of assays are described in detail in the two

following sections.

Evolutionary Dynamics: Relative proportions of barcoded yeast strain pairs evolved in

competition within the same well were quantified at generations 0, 100, 150, 200, 220, 240,

and 250 (Days 0, 10, 15, 20, 22, 24 and 25) for a total of 532 evolutionary dynamics samples

across 76 evolving two-barcode populations. These samples were pooled for DNA extraction

such that there was no barcode overlap within pools. DNA was subsequently extracted (see the

Proof-of-concept fitness assays section for details). Libraries were constructed for sequencing as

described in the *Library construction and sequencing* section. From these data, the relative

proportions of barcoded yeast strains within each pair were measured at each time-point.

Using this information, the time-point (t-max) and magnitude (m-max) of the maximum change

in relative abundance in comparison to the starting conditions, the time-point (t-max-rate) and

magnitude (m-max-rate) of the maximum rate of change between adjacent time-points, the

time-point (t-max-diff) and maximum difference (m-max-diff) in barcode proportions, and the

total cumulative change in barcode relative abundance were calculated for each two-barcode

population. Barcodes approaching fixation (hereafter referred to as "fixed") were also noted

and were defined as cases in which a single barcode in the pair obtained (and maintained) a

proportion of 0.95 or greater by (through) generation 250 of the evolution experiment.

End-point assays of relative fitness: This second type of assay was designed to assess the fitness

of a given focal strain relative to a reference using a barcoded competition-based assay.

Because they involve taking samples (and quantifying fitness) at the start and end of an

experimental evolution, end-point fitness assays provides a means of measuring fitness change

across the entire experiment in the design utilized here. In this case, the assays involved

quantifying barcode fitness using pooled samples from generation 0 (Figure 1, B2.) and

generation 250 (Figure 1, B3.) from the 250-generation evolution experiment (Figure 1, B1.).

Briefly, Generation-0 and Generation-250 yeast strains were revived and samples from each

time point were pooled such that there was no overlap in barcoded yeast strain identity within

each pool. Pools of revived generation-0 and generation-250 yeast were independently

combined in equal proportion (50 pooled-yeast : 50 ancestral reference) with an ancestral

reference strain (re: an 'unevolved' barcoded yeast strain); these pooled samples were then

diluted into fresh medium to initialize the fitness assays. Four replicate Fitness assays were

conducted for each pool of generation-0 and generation-250 yeast. Fitness assays employed a

standard 1:1000 transfer dilution across all samples (see *Strains, media and culture methods*).

Samples evolved in CM plus additional stresses were assayed in the same media type that they

were evolved in. Diploid yeast were competed against a diploid reference strain (strain ID:

d1H10), while haploid yeast were competed against the haploid version of this same reference

(strain ID: h1H10). In these assays, yeast samples for DNA extraction were obtained from the

undiluted initial mixtures (fitness assay starting material), and, 20 generations later, from the

final overnight population cultures (fitness assay end) to assess fitness of generation-0 and

generation-250 yeast strains. See *Fitness Calculations*, below, for a full description of fitness

assay methodology and for calculations of fitness from fitness assay barcode abundance data.

Libraries were constructed for sequencing as described in the *Library construction and*

sequencing section, below.

Library construction and sequencing: Barcode sequencing libraries for the proof-of-concept

fitness assays and all samples from both components of the 250-generation evolution

experiment were constructed by amplification of MoBY barcodes [36] with primers containing

241 Ion Torrent adaptors with indexes assigned to distinguish samples from one another (Additional

242 file 3).

244 PCR products for library construction were generated using 25 cycles and were subsequently

245 quantified with a Qubit 3.0 Flourometer, high sensitivity assay kit. These products were then

246 combined at equimolar concentrations and purified using a Zymo DNA Clean & Concentrator kit

247 (Zymo Research D4014) to create a single library for sequencing. The DNA extraction and PCR

248 steps were repeated for samples that did not attain sufficient DNA concentrations (no samples

249 were exposed to > 25 rounds of PCR). We note that PCR jackpotting can be an issue in DNA

250 library construction and that it can add noise to measurements of barcode frequency. To

251 minimize the influence of this phenomenon on our results, we used a large amount of template

252 DNA [38], and conducted all fitness assays in four replicates. Additionally, we avoided

253 misclassification of barcodes by using high fidelity Taq Plus DNA Polymerase (Lambda Biotech,

254 Taq Plus Master mix Red), and MOBY barcode sequences that differ from one another in at

255 least six substitutions.

257 DNA libraries were sequenced using an Ion Torrent sequencer (Ion Proton System, Ion Torrent)

258 at the Genomics Core Facility at Saint Louis University with a customized parameter to assess

259 polyclonality after 31bp (the start position of the forward Ion Torrent adapter index sequence).

260 A single sequencing run was used for each pooled library (library 1 – proof-of-concept fitness

261 assays, library 2 – evolution experiment: evolutionary dynamics samples, Generation-0 fitness

assays, and Generation-250 fitness assays). An additional library was constructed and

sequenced for a set of samples from library 2 with elevated barcode cross-contamination rates.

Sequence data processing & calculations

Sequence datasets: Sequence data in FASTQ format were parsed and demultiplexed using

custom scripts in R (see the *Availability of data and materials* statement, below). A total of

142,243,245 raw reads that matched the forward Ion Torrent adapter indices included in our

experiment (omitting reads that matched no forward adapter, polyclonal reads, low quality

reads, and adapter dimer reads) were recovered across the three sequenced libraries.

104,365,740 reads (73.4%) were retained for analysis that perfectly matched a forward

sequencing adapter index (9-12 bp), reverse sequencing adapter index (9-12 bp) pair, and a

MOBY genetic barcode (20 bp) included in the full experimental design.

Barcode cross-contamination rate: The barcoded yeast experimental evolution system has an

innate ability to detect and track barcode cross-contamination that could arise during

evolution, over the course of short-term fitness assays, or during DNA library preparation.

Barcode cross-contamination rate was defined as the mean number of counts mapping to any

given barcoded yeast strain included in the full experimental design (sequenced library), but not

expected to be present in the given sample (pair of forward and reverse IonTorrent adapter

Indices). Barcode cross-contamination rates were calculated separately for each unique

forward-reverse index pair included in each sequencing library and represent the amount of

noise an average contaminating barcode strain contributes to each sample. The barcode cross-

contamination rate of sample (primer pair) j, B_j, was measured as,

Equation 1.

$$B_j = \frac{\frac{1}{m}\sum_{i=1}^{m} C_i}{T_j}$$

where m is the number of barcoded strains that could potentially contribute to barcode cross-

contamination in experiment j (i.e., the number of unique strain IDs included in the full library,

but not expected in sample j), C_i is the number of barcoded yeast strain counts recovered for

contaminating barcode i, and T_j is the total number of counts recovered in sample j across all

barcoded yeast strains included in the full experimental design.

Contamination rate summaries are reported separately for the proof-of-concept fitness assays

and the evolution experiment (the latter containing evolutionary dynamics, generation-0 fitness

assay and generation-250 fitness assays samples). For the subset of evolution experiment

samples that were sequenced in two separate libraries, only the replicate with a lower

contamination rate was retained for contamination rate summary reporting and all

downstream analyses. In all statistical analyses reported below, contamination rate is initially

included as a potential predictor and subsequently removed from the model if its effect was

found to be non-significant.

Fitness calculations: Fitness was evaluated via sequencing-based assays that involved

competing barcoded yeast strains against a common ancestral reference strain for 2 days (20

generations). Reported fitness values for the yeast strains in all fitness assays are relative to the

same barcoded ancestral reference strain (strain ID: d1H10 for diploids; strain ID: h1H10 for

haploids). The relative fitness of barcoded yeast strain i at generation gn, $w_{i\,gn}$, was measured

as,

Equation 2.

$$w_{i_{gn}} = \left(\left(ln\frac{Ci_{20}}{R_{20}} - ln\frac{Ci_0}{R_0}\right)/20\right)^e$$

where Ci and R refer to barcode counts for the focal barcode and reference barcode at fitness

assay generation 0 (initial mixtures; fitness assay initial measurement) and fitness assay

generation 20 (final overnight cultures; fitness assay final measurement), and 20 is the number

of generations between measurement at fitness assay generation 0 and fitness assay

generation 20 [39]. The change in fitness of strain i between Day 0 and Day 25 in for our 250-

generation evolution experiment, Δw_i, we therefore computed as,

Equation 3.

$$\Delta w_i = w_{i_{g250}} - w_{i_{g0}}$$

where $w_{i\,g0}$ and $w_{i\,g250}$ are the strain's fitness at generation 0 and 250 as measured from

Equation 2.

Statistical analysis

Analysis & visualization tools: All Analyses and statistical test were conducted in R version 3.5.2

[40]. Data processing uses primarily base-R functionality; the plyr package [41] is used in some

cases for data frame manipulation. All Statistical models with linear mixed effects were

generated using the lme4 [42], and lmerTest [43] packages. Weighted t-tests were assessed

with the weights package [44]. Power analyses were conducted using the pwr package [45].

Finally, figures and tables were generated with the ggplot2 [46] and sjPlot [47] packages,

respectively; multi-panel figures were built using the gridExtra package [48]. Raw p-values are

reported unless otherwise noted; some tables included in the supplement report raw and

corrected p-values (i.e., false discovery rate) for cases where only corrected values are

presented in the main text.

Reads: Analysis of multiplex barcode sequencing data requires careful consideration of sample

size. Because count data are ultimately handled as relative frequencies (proportions), it was

necessary to consider underlying sample size or "confidence" in each piece of data within the

full dataset for all calculations and analyses. That is, entries with more reads were explicitly

assumed to contribute more to summary calculations and statistical models. Variation in

sample size was thus controlled for by weighting all calculations by the read sample size and by

including such weights in downstream statistical models. This sample size metric considers both

the total counts recovered for a multiplexed sample (unique forward and reverse index

sequence adapter pair) and the number of counts recovered for the focal barcoded yeast

strain(s) for that entry. In the analyses presented here, read calculations for constructed

variables utilize harmonic means rather than arithmetic means; the harmonic mean is sensitive

to small values and therefore does not overinflate the read values for any variables constructed

from a combination of high- and low-read data sources.

Proof-of-concept fitness assays: Fitness differences among 92 constructed barcoded yeast

strains used in the proof-of-concept fitness assays were assessed using a weighted linear model

with mean-corrected fitness as the response variable and yeast strain ID as the predictor

variable. Barcode cross-contamination is not included in this model as all fitness assay samples

contained all 92 focal barcodes; barcode cross-contamination is reported from a separate set of

1-barcode and 2-barcode samples for the Proof-of-concept fitness assays.

Power Analysis: Power analyses were conducted to calculate the sensitivity of the system in

measuring treatment-level fitness effects, fitness changes for individual barcoded yeast strains,

and initial fitness variation among individual strains. Power was calculated using sample size

and fitness effect as variables of interest with a fixed value for error in fitness measurements.

Error in fitness was estimated from the root mean squared error among replicates from the

Proof-of-Concept fitness assays as well as from replicate measures of fitness change for

individual barcodes from the 250-generation experiment. Power analyses were conducted

using the pwr package in R; the pwr.f2.test function was used to calculate power to discern

treatment-level differences in fitness and the pwr.t.test function was used to calculate power

to detect individual barcode fitness changes and initial fitness differences among individual

barcodes.

Multiplex barcode sequencing and cross-contamination: Barcode cross-contamination rate

was assessed (Equation 2) and summarized, separately for the proof-of-concept fitness assays

and the 250-generation experiment. The effects of barcode cross-contamination rate on the

change in fitness and the error among replicate fitness measures for each barcode were

assessed by including barcode cross-contamination as a predictor in two sets of linear mixed-

effects models with either change in fitness over 250 generations of experimental evolution or

standard error in fitness change over 250 generations (across four replicate measures for each

barcode) as the response variable. Additionally, a weighted 1-tailed t-test was utilized to assess

whether barcode cross-contamination rates for samples collected in the 250-generation

experiment decreased after re-extracting DNA and resequencing.

Fitness change in 250 generations of experimental evolution: Individual barcoded yeast strains

that exhibited any change in fitness over the 250-generation fitness experiment were identified

using a weighted linear model with change in fitness as the response variable and a strain

identifier (treatment plus yeast strain Barcode ID) as the predictor variable.

Next, the effects of evolutionary treatment (medium type, ploidy, and transfer dilution) on

change in fitness over 250 generations of experimental evolution was assessed using a

weighted linear mixed-effects model with cross-contamination rates as a covariate. A random

effect of strain identifier (treatment plus yeast strain barcode ID) was placed on the model

intercept. A second weighted linear model was fit to assess potential sources of error in the

measurement of fitness in this system. Here, a model was constructed using a comprehensive

set of factors that could have impacted the standard error in fitness measurement among

replicates. These included: treatment, cross-contamination rate, magnitude fitness change,

median count (median number of counts returned for a focal barcode across all sampling points

that contribute to the fitness change calculation), and median proportion (median proportion

of the focal barcode in its mixed 2-barcode well across all sampling points).

Evolutionary dynamics: For all analyses of evolutionary dynamics, only the first barcoded yeast

strain from each well was included to ensure that data were not double-counted. A set of

weighted linear mixed effects models, with treatment as the predictor variable and either t-

max, m-max, t-max-rate, m-max-rate, t-max-diff, m-max-diff, or total cumulative change in

barcoded relative abundance across all time-points as the dependent variable, were employed

to assess treatment differences in evolutionary dynamics. Full models also included initial

barcode abundance as a predictor term because initial barcode abundance could impact

subsequent dynamics. Initial barcode abundance was subsequently removed from models

when deemed nonsignificant, resulting in removal from all but one model (t-max-diff). Barcode

fixation rate was not assessed statistically due to the small number of fixation events observed

(n=7/76 populations).

410   RESULTS

Proof-of-concept fitness assays

We constructed 92 diploid yeast strains, each with a unique 20 bp barcode inserted upstream

of the deleted *HO* gene. Proof-of-Concept fitness assays were then used to measure any fitness

differences among these constructed "barcoded" strains, to estimate our power to detect small

fitness differences, and to assess our ability to measure fitness using multiplexed barcode

sequencing. We assayed fitness simultaneously for our pool of 92 barcodes by measuring

changes in barcode abundance relative to a 'reference' strain (barcode ID: d1H10) in 10

replicates over a two-day period of approximately 20 generations. A total of 920 fitness assays

were performed in this section of this study (92 barcodes x 10 replicates).

A few of the barcoded strains showed significant differences in fitness (3/92 at a 5% false

discovery rate, FDR; 1/92 at a 1% FDR) (Figure 2, Additional file 4). The root mean squared error

(rMSE) among replicated measures of fitness was 1.76e-2, indicating good power (80%) to

detect fitness differences of 1.2% between strains with four replicates at a nominal significance

level of P = 0.05 (Figure 3a, Additional file 5). We subsequently used these data to estimate

power to detect treatment effects in our proposed experimental design for the 250-generation

experiment. With a modest number of barcoded yeast strains per treatment (n=22), we

estimate high power (99.8%) to detect 1% fitness differences between treatments at a nominal

significance level of P = 0.05 with four replicate fitness assays (Figure 3b, Additional file 6). With

these promising results, we proceeded to conduct a more comprehensive test of the utility of

barcoded strains in a practical experimental evolution context (i.e., the utility to evaluate the

effects of treatment on fitness outcomes and evolutionary dynamics).

Experimental evolution

To empirically evaluate the strengths of this barcoded-strain system for studies of experimental

evolution, we conducted a 25-day evolution experiment, which is approximately equal to 250

generations of evolution. Our design included 76 populations, each initiated with two barcodes

per well. Samples spanned six treatments, which varied in yeast strain ploidy, growth medium,

and daily transfer dilution (Additional file 2). To measure any changes in fitness we competed

the evolved generation-250 strains and their generation-0 ancestors against a common

reference strain over a two-day period of approximately 20 generations. With four replicate

fitness assays for each barcode, this amounted to 1,216 fitness assays (2 barcodes * 2 time-

points * 76 populations * 4 replicates). We also measured the change in barcode frequency of

barcoded strains within each evolving population (barcode pair) from 532 longitudinal-

evolutionary dynamics samples (76 barcoded pairs * 7 timepoints). From these samples we

quantified the magnitude and timing of changes in barcode frequency, which should be

influenced by changes in the fitness of the evolving barcoded strains present within each

population.

Multiplex barcode sequencing: We utilized a two-step multiplexed design to obtain high

throughput estimates of fitness based on barcode sequencing in our fitness assays. In the first

step we leveraged the strain-identifying barcodes by pooling multiple strains together and

simultaneously competing them against an ancestral reference strain to estimate relative

fitness. In our second multiplexing step we PCR-amplified each fitness assay sample with unique

forward and reverse indexed sequencing adaptors. This latter step enabled us to assign

sequencing reads to the appropriate fitness assay after sequencing many samples in concert as

a single library. We constructed one library for each experiment (Library 1 – Proof-of-Concept

fitness assays; Library 2 – 250-generation evolution experiment evolutionary dynamics and

fitness assays samples).

From 84,776,627 demultiplexed reads, we obtained a median of 2,961 reads per barcode in

each sample. However, we also detected barcode cross-contamination, defined as barcodes

with sample indexes that shouldn't exist in our sequenced libraries. The average rate of

barcode cross-contamination per sample was low, 0.04%, and consistently present in nearly all

of our samples (Additional file 7 A, B). Contamination could occur during culturing of yeast

strains, liquid handling during preparation of libraries [4,25], or potentially via index switching

during sequencing procedures [49–51]. However, the low but uniform rate of cross

contamination we observe is more consistent with library preparation and sequencing than

yeast contamination. This is further evidenced by the fact that no growth was observed in

culture blanks during any fitness assays nor during the 25-day experimental evolution

experiment.

To test whether the barcode cross-contamination rate depends on liquid handling during library

preparation, we reprocessed a subset of samples with high cross contamination rates using

identical starting material (multiple yeast sample aliquots were created at the time of sample

collection). Cross contamination rates decreased significantly in these reprocessed samples (t =

22.3, df = 65, p < 10e-6), but a low level of background contamination remained (Original

Samples: mean = 0.75%; reprocessed samples: mean = 0.05%) (Additional file 8).

In our analyses, the presence of low abundance barcode cross-contamination was removed

from all samples in which a barcode is not expected to occur. However, this doesn't eliminate

contamination in samples where a barcode is expected to occur. In such cases, error in

estimates of barcode frequency is highest when a barcode's frequency is low and approaches

the cross-contamination rate. The effects of contamination rate and other system-specific

covariates on experimental outcomes are presented in detail in the next section.

End-point fitness assays: We assessed fitness in the 152 evolved strains (evolution experiment

generation-250 strains) as well as their ancestors (evolution experiment generation-0 strains).

For each strain and timepoint, fitness was measured in comparison to a common ancestral

reference strain via a two-day competition-based fitness assay in the same media type under

which the strain had been evolved. The root mean squared error (rMSE) among replicated

measures of fitness was 4.31e-2 for these assays. This is slightly higher than our proof-of-

concept fitness assays but is still expected to provide reasonable power to detect fitness

changes for individual barcodes (80% power to detect fitness changes of 2.99%) and fitness

differences among treatments (80% power to detect treatment differences of 0.81%) (Figure 3c

& Figure 3d, Additional file 9, Additional file 10). Among a number of covariates tested we

found that only the cross-contamination rate (P = 0.01) and magnitude fitness change (P =

6.03e-4) were associated with error among replicates (Additional File 11).

Twenty three percent (35/152) of the strains showed significant increases in fitness between

generation 0 and generation 250 at a 1% false discovery rate, FDR (Figure 4, Additional File 12).

Within this subset, the average fitness increase was 6.80% with a range of 2.75% to 23.5%.

Relatively few strains with significant increases in fitness were found in the CM treatment

(3/42; 7.14%) and ethanol stress treatment (2/22; 9.09%). A higher proportion of strains

exhibited significant increases in fitness in the lower 1:250 dilution treatment (9/22; 40.9%), the

salt stress treatment (11/22; 50.0%), and the haploid treatment (10/22; 45.5%). No strains in

the 1:4000 dilution treatment (0/22) exhibited increases in fitness. No strains (0/152) exhibited

a significant decrease in fitness.

Next, we evaluated the effect of evolutionary treatment on fitness change. We found

significant variation in fitness among treatments (P < 10e-6, Figure 5, Additional file 13).

Relative to our standard culture conditions, diploids in CM ("no stress") with 1:1000 transfer

dilution, we found greater (more positive) fitness change in diploid strains evolved in salt stress

(P < 10e-6), in haploid strains evolved in standard culture conditions (P = 0.01), and in diploid

strains evolved with a less extreme (1:250) daily transfer dilution (P = 9.49e-4). Relative to the

CM ("no stress") diploid treatment (our standard culture conditions), we found less (less

positive) fitness change in diploid strains evolved in ethanol stress (p = 3.99e-3). We found no

significant effect of a more extreme (1:4000) daily transfer dilution on fitness change (p=0.39).

In addition to the treatment effects, there was also a negative effect of barcode cross

contamination rate on fitness change (P < 10e-6).

Evolutionary dynamics: In experimental evolution, adaptation can influence the relative

abundance of barcodes evolving in competition [27,34]. In addition to characterizing fitness

change for the evolved barcodes, we tracked the relative proportions of the 76 barcoded-yeast

pairs at seven timepoints during our 250-generation evolution experiment (days 0, 10, 15, 20,

22, 24, and 25) to examine the evolutionary dynamics generated by adaptation and other

processes (e.g., drift). We collected 8 measurements from these 532 evolutionary dynamics

samples (76 populations * 7 timepoints): (1) barcode fixation, (2) the time-point and (3)

magnitude of the maximum change in relative abundance in comparison to the starting

conditions, (4) the time-point and (5) magnitude of the maximum rate of change between

adjacent time-points, (6) the time-point and (7) magnitude of the maximum difference in BC

proportions, and (8) the total cumulative change in barcoded relative abundance summed

across all time-points.

Out of the seven measures of adaptive dynamics that were amenable to statistical testing (i.e.,

all adaptive dynamics measures other than fixation rate, which had sparse imbalanced binary

data), six significantly differed by treatment (Table 1; see additional files 14:27 for individual

adaptive dynamics plots and results tables). Relative to the CM-diploid treatment (our standard

culture conditions) both the salt and haploid treatments were associated with a larger

maximum change in relative abundance in comparison to the start (p = 5.24e-5, p = 5.85e-3,

respectively) and a more extreme maximum difference in barcode proportion (p = 2.77e-4, p =

6.01e-6, respectively). The salt treatment was also associated with an earlier time-point of the

maximum change in abundance relative to the start (p = 0.01), and the haploid treatment was

associated with an earlier maximum rate of change (p = 1.35e-5). The total cumulative change

in barcode abundance was greater for the 1:4000 dilution treatment than the 1:1000 dilution

treatment (p = 1.77e-3). Total cumulative change in barcode abundance was significantly less

for haploids than diploids (p = 4.9e-2), and significantly less in the ethanol stress treatment than

in the CM with no stress treatment (p = 2.98e-2). Finally, we observed barcodes that

approached fixation in eleven percent (9/76) of our populations. Most fixation events were

observed in the salt (3/11; 27.3%) and haploid (5/11; 45.5%) treatments, one instance was

found in the CM-diploid treatment and no instances of fixation were observed in the ethanol,

1:4000 dilution, nor 1:250 dilution treatments.

555   DISCUSSION

Experimental evolution has proven to be a valuable approach for studying a range of

evolutionary questions. In this study we implemented a genetic barcoding system in *S.*

cerevisiae to increase the efficiency and throughput of these types of studies. Through

barcoding of ancestral strains we were able to increase the complexity of our experimental

design, increase the throughput of fitness assays by multiplexing of independent samples, and

provide a relatively simple means to track the evolutionary dynamics of barcodes evolving in

competition during the experimental evolution period. Accordingly, we have demonstrated the

potential for detecting fitness differences in six experimental evolution treatments and have

shown that, while the typically low levels of barcode cross-contamination we observe cannot

be completely eliminated, their effects on inference can be minimized through simple statistical

procedures. Below we discuss the merits and drawbacks of the barcoding system, and its

capabilities to increase efficiency and throughput in measuring fitness changes that occur

during experimental evolution.

One important consideration of a barcoding system like the one we examine here, is that

barcoded strains are not necessarily identical to one another at the beginning of an experiment

(i.e., have equal fitness) even if all barcoded variants are produced from a single ancestral

clone. Although we found no significant fitness differences among the majority of barcoded

strains, we note that we did indeed observe a few strains with significant deviations from the

population mean fitness. Given the location of our barcode insertions (i.e., a non-functional

region of the genome) it is unlikely that the barcodes themselves generated these fitness

differences. A perhaps more likely explanation is that these differences arose from mutations

that occurred during transformation [29] or shortly thereafter. Regardless, this limitation could

be mitigated by (1) removing strains with unequal generation-0 fitness entirely from their

analyses, (2) quantifying initial fitness differentials and including them as a covariate in

downstream analyses, or (3) looking explicitly at change in fitness for each individual barcode as

we do here. We note that initial fitness differentials may be of interest themselves, given that

they can potentially impact evolutionary outcomes [52–54] and the types of mutations that

successfully spread through an experimental population [55].

In addition to throughput, the power to detect small changes in fitness in evolved strains is a

critical parameter of any experimental evolution system. We estimated 80% power to detect a

fitness difference of 1.2% in the proof-of-concept assays and 80% power to detect a fitness

change of 3.0% in the 250-generation evolution experiment for any individual barcode. While a

formal power analysis is not often reported, a review of the literature suggests that prior

studies have detected fitness differences between 1 and 5% [26,32,56,57]. For example, one

previous barcoded fitness assay found between replicate deviations of 1-2% and between batch

deviations of up to 5% [32]. Some studies report confidence intervals significantly lower than

1%, but these studies tend to employ a large number of replicates [12,55,58]. We also note that

our power to detect treatment effects was much higher than our ability to detect changes in

fitness for individual barcoded strains. Specifically, we estimated 99.8% power to detect fitness

differences of 1% between treatments using the error in our proof-of-concept fitness assays,

and 87.7% power to detect fitness differences of 1% between treatments using the actual 250-

generation experiment data.

While the error measured in our fitness assays is consistent with most prior studies, pooled

fitness assays do have additional sources of error. Prior studies using pooled fitness assays

estimated that pools containing 1000 cells per strain are expected to have an error rate of 4.4%

[59] after two rounds of dilution due to sampling error. The expected number of cells per

barcoded strain at inoculation in our system is 9,000 (3.0×10^8 cells/ml at saturation in 0.6ml of

media, diluted 1:1000 and divided by up to 20 strains per assay). However, the number of cells

could be much lower due to sampling or to low frequency of the barcode of interest in a well at

the end of the 250 generations of evolution. Lineages with large changes in fitness are expected

to deviate the most from equal numbers of cells in a well. Consistent with this possibility, the

RMSE of fitness in generation 250 (4.34) is higher than the RMSE of fitness in generation 0

(2.92), where barcode frequencies are much more uniform. This provides one explanation for

our finding that error among replicates is associated with the magnitude of the fitness change.

Thus, while there are some limitations to pooled fitness assays, these limitations can be

compensated by examination a large number of strains per treatment –as done in this study–

or by increasing the number of replicate assays.

Another important consideration when designing an experimental evolution system is that it

must be fairly robust to contamination. While culture contamination is rare in experimental

evolution [25,60], barcode cross-contamination is possible. Our results are consistent with prior

work on this topic. Specifically, while we found no evidence of culture contamination in any

growth blanks, we detected a uniformly low level of diffuse barcode cross-contamination in

multiplexed fitness assays and evolutionary dynamics samples. Because there was no

conspicuous pattern of cross-contamination, we suggest that the observed contamination was

likely introduced during sample preparation and/or DNA sequencing. DNA extraction is a likely

source of cross-contamination in samples processed in strip-tubes or 96-well plate formats that

prioritize throughput. However, we minimized the chance for contamination in the DNA

isolation step in our experiments by isolating DNA with individual reaction tubes for each

sample. Furthermore, contamination in the DNA extraction step would not be consistent with

the low-level diffuse contamination observed, a pattern that tends to be more indicative of

contamination occurring downstream of DNA extraction. Several other sources of

contamination are possible, including primer contamination during the index addition for PCR

[61] and index switching during library construction or sequencing [49–51]. The latter possibility

seems nevertheless less likely because all PCR steps were performed separately prior to pooling

of libraries and because no previous studies report index switching for IonTorrent sequence

data to our knowledge.

The strength and efficiency of barcoded experimental evolution is further evidenced by this

study. Barcoding enabled us to evolve a large number of strains (152) for 250 generations

across six treatments, and to conduct a total of 532 evolutionary dynamics measurements and

1,216 fitness assays related to these manipulations in a relatively short amount of time and

with limited resources. Reassuringly, our results are largely consistent with prior work. We find

greater fitness increases, i.e., a greater rate of adaptation, in haploids than diploids. This finding

agrees with another study that found faster rates of adaptation and larger effective population

sizes in haploids relative to diploids [10]. In a related study, Selmecki et al., [27] found faster

adaptation in tetraploids than diploids, but no difference in rate of adaptation between

haploids and diploids, potentially suggesting a trend opposite to ours of increasing adaptive

rate with increasing ploidy. However, differences between haploids and diploids must be

treated with caution because there is mounting evidence that haploid yeast evolves towards

diploidy under experimental evolution conditions [20,27,62]. We also find greater fitness

increase in complete medium (CM) plus NaCl stress than in CM alone, which was not surprising

given what is known about adaptation to NaCl stress in *S. cerevisiae* [20,63–65]. In contrast, we

were surprised to detect less fitness increase in CM plus EtOH stress than in CM alone. There

are several non-mutually exclusive explanations for this result. It is possible that ethanol did not

present a significant stress (selective pressure) to the cells once they had attained physiological

adaptation to the medium, i.e., acclimation [19]. It is also possible that adaptations to CM and

adaptations to ethanol exhibit antagonistic pleiotropy, similar to what has been found in

experiments contrasting rich and poor media [66] or exploring adaptation to other chemical

stressors [67]. Pleiotropy could also shed light on the marked adaptation observed in the NaCl

treatment given that adaptation to CM and NaCl stress may exhibit complementarity via

synergistic or positive pleiotropy [21,68–70]. We find no difference in fitness change between

our standard (1:1000) dilution treatment and a treatment with a more extreme (1:4000) daily

transfer dilution. We do, however, find a greater increase in fitness when a less extreme (1:250)

daily transfer dilution was used instead of the standard (1:1000) dilution. While these transfer

dilution findings are not necessarily expected [12,71–73], they nevertheless support earlier

findings that less extreme bottlenecks favor the maintenance of adaptive mutants [74] and that

large populations are less adaptively constrained than small ones in simple environments [73].

They are also consistent with previous evidence for greater mean fitness increase in wide

versus narrow bottleneck populations [11].

In addition to high-throughput fitness assays, barcoding enabled us to track the relative

proportions of barcoded lineage pairs evolving in competition during the experimental

evolution period [30,31,75,76]. As expected, we found general agreement between the

evolutionary dynamics results and endpoint fitness assay results in our six experimental

treatments. For example, the haploid and NaCl stress treatments both displayed dynamics

consistent with more extreme increases in fitness, including greater change in barcode

abundance relative to the starting conditions and a greater maximum difference in barcode

abundance than diploids in CM. Interestingly, haploids also showed signs of earlier adaptation

than diploid strains in similar conditions, as evidenced by an earlier generation of maximum

rate of change in barcode abundance (also noted in [27,30,31]) and a lower total change in

barcode abundance over 250 generations. Although this latter result may seem paradoxical, it is

consistent with the observation that haploids adapt earlier than diploid strains [10], and is

expected if a greater proportion of the total change in barcode abundance of haploids in our

experiments happened in the first 100 generations (before we collected any evolutionary

dynamics data).

Despite significant differences in barcode dynamics, there are some limitations to interpreting

these results. Because we mostly assessed abundance every 50 generations, it is possible that

we missed some of the adaptive dynamics; dense temporal sampling is ideal for a full picture of

evolutionary dynamics in populations subjected to experimental evolution [76]. Furthermore,

due to projected read depth constraints, barcode frequencies over time were not measured in

replicate for the 250-generation evolution project. Finally, changes in barcode abundance over

long time scales may not be directly related to final fitness as these measures are calculated

over 250 generations of evolution, during which relative fitness relationships between evolving

barcodes could change significantly. For example, the 1:4000 transfer dilution bottleneck

treatment had elevated rates of change in barcode abundance and a high amount of total

change in barcode abundance without a concomitant increase in fitness, possibly as a result of

the potentially strong effects of drift when severe bottlenecks are frequent. We suggest that

future studies employ a denser and more even longitudinal-evolutionary dynamics sampling

scheme, with replication, to maximize the value and precision of this type of lineage tracking or

evolutionary dynamics data.

703   CONCLUSIONS

In summary, we conclude that the barcoded yeast system that we describe here greatly

increases the throughput of fitness measurements and provides a relatively simple means for

lineage tracking. These two characteristics greatly increase our ability to design more complex

and potentially more useful evolutionary experiments. We note that although barcode cross-

contamination imposes some limitations on the implementation of this system, it is possible to

track the origin and rates of such contamination and, therefore, to statistically consider its

effects on experimental outcomes. Thus, we conclude that this system represents a significant

step forward toward the design and implementation of experimental evolution systems capable

of high throughput fitness measurements.

715   REFERENCES

1.   de Varigny H. Experimental Evolution. London, United Kingdom: MacMillan and Co; 1892.

2.   Garland T, Rose MR. Experimental Evolution concepts, methods, and applications of

selection experiments. University of California Press; 2009.

3.   Kassen R. Experimental Evolution and the Nature of Biodiversity. Greenwood Village, CO

USA: Roberts and Company publishers, Inc.; 2014.

4.   Van den Bergh B, Swings T, Fauvart M, Michiels J. Experimental Design, Population

Dynamics, and Diversity in Microbial Experimental Evolution. Microbiol Mol Biol Rev

[Internet]. 2018;82(3):e00008-18. Available from:

http://mmbr.asm.org/lookup/doi/10.1128/MMBR.00008-18

5.   Bailey SF, Guo Q, Bataillon T. Identifying drivers of parallel evolution: A regression model

approach. Genome Biol Evol. 2018;10(10):2801–12.

6.   Bailey SF, Blanquart F, Bataillon T, Kassen R. What drives parallel evolution?: How

population size and mutational variation contribute to repeated evolution. BioEssays.

2017;39(1):1–9.

7.   Graves JL, Hertweck KL, Phillips MA, Han M V., Cabral LG, Barter TT, et al. Genomics of

parallel experimental evolution in drosophila. Mol Biol Evol. 2017;34(4):831–42.

8.   Burke MK, Liti G, Long AD. Standing genetic variation drives repeatable experimental

evolution in outcrossing populations of saccharomyces cerevisiae. Mol Biol Evol.

2014;31(12):3228–39.

9.   Kohn LM, Anderson JB. The underlying structure of adaptation under strong selection in

12 experimental yeast populations. Eukaryot Cell. 2014;13(9):1200–6.

10.   Gerstein AC, Cleathero LA, Mandegar MA, Otto SP. Haploids adapt faster than diploids

across a range of environments. J Evol Biol. 2011;24(3):531–40.

740 11. Schoustra SE, Bataillon T, Gifford DR, Kassen R. The properties of adaptive walks in

741   evolving populations of fungus. PLoS Biol. 2009;7(11).

742 12. Kryazhimskiy S, Rice DP, Desai MM. Population subdivision and adaptation in asexual

743   populations of saccharomyces cerevisiae. Evolution (N Y). 2012;66(6):1931–41.

744 13. Low-Décarie E, Kolber M, Homme P, Lofano A, Dumbrell A, Gonzalez A, et al. Community

745   rescue in experimental metacommunities. Proc Natl Acad Sci. 2015;112(46):14307–12.

746 14. Bell G, Gonzalez A. Adaptation and Evolutionary Rescue Environmental Deterioration.

747   Science (80-) [Internet]. 2011;332(June):1327–30. Available from:

748   http://www.ncbi.nlm.nih.gov/pubmed/21659606

749 15. Loewe L, Textor V, Scherer S. High Deleterious Genomic Mutation Rate in Stationary

750   Phase of Escherichia coli. Science (80-). 2003;302(5650):1558–60.

751 16. Swings T, van Den Bergh B, Wuyts S, Oeyen E, Voordeckers K, Verstrepen KJ, et al.

752   Adaptive tuning of mutation rates allows fast response to lethal stress in escherichia coli.

753   Elife. 2017;6:1–24.

754 17. Lenski RE, Sniegowski P, Gerrish P. Evolution of high mutation rates in experimental

755   populations of E. coli. Nature. 1997;387(June):703–5.

756 18. Perfeito L, Fernandes L, Mota C, Gordo I. Adaptive mutations in bacteria: High rate and

757   small effects. Science (80-). 2007;317(5839):813–5.

758 19. Huang CJ, Lu MY, Chang YW, Li WH. Experimental Evolution of Yeast for High-

759   Temperature Tolerance. Mol Biol Evol. 2018;35(8):1823–39.

760 20. Dhar R, Sägesser R, Weikert C, Yuan J, Wagner A. Adaptation of Saccharomyces

761   cerevisiae to saline stress through laboratory evolution. J Evol Biol. 2011;24(5):1135–53.

21.   Dhar R, Sägesser R, Weikert C, Wagner A. Yeast adapts to a changing stressful

environment by evolving cross-protection and anticipatory gene regulation. Mol Biol

Evol. 2013;30(3):573–88.

22.   Hughes BS, Cullum AJ, Bennett AF. Evolutionary adaptation to environmental pH in

experimental lineages of Escherichia coli. Evolution (N Y). 2007;61(7):1725–34.

23.   Horinouchi T, Suzuki S, Hirasawa T, Ono N, Yomo T, Shimizu H, et al. Phenotypic

convergence in bacterial adaptive evolution to ethanol stress. BMC Evol Biol [Internet].

2015;15(1):1–14. Available from: http://dx.doi.org/10.1186/s12862-015-0454-6

24.   Zhou A, Baidoo E, He Z, Mukhopadhyay A, Baumohl JK, Benke P, et al. Characterization of

NaCl tolerance in Desulfovibrio vulgaris Hildenborough through experimental evolution.

ISME J [Internet]. 2013;7(9):1790–802. Available from:

http://dx.doi.org/10.1038/ismej.2013.60

25.   Lenski RE, Rose MR, Simpson SC, Tadler SC. Long-Term Experimental Evolution in

Escherichia coli. I. Adaptation and Divergence During 2,000 Generations. Am Nat

[Internet]. 1991;138(6):1315–41. Available from:

http://www.journals.uchicago.edu/doi/10.1086/285289

26.   Gresham D, Desai MM, Tucker CM, Jenq HT, Pai DA, Ward A, et al. The repertoire and

dynamics of evolutionary adaptations to controlled nutrient-limited environments in

yeast. PLoS Genet. 2008;4(12).

27.   Selmecki AM, Maruvka YE, Richmond PA, Guillet M, Shoresh N, Sorenson AL, et al.

Polyploidy can drive rapid adaptation in yeast. Nature [Internet]. 2015;519(7543):349–

51. Available from: http://dx.doi.org/10.1038/nature14187

28.   Roth FP, Smith AM, Kaper F, Nislow C, Giaever G, Thompson MJ, et al. Quantitative

785   phenotyping via deep barcode sequencing. Genome Res. 2009;19(10):1836–42.

786   29.   Giaever G, Nislow C. The yeast deletion collection: A decade of functional genomics.

787   Genetics. 2014;197(2):451–65.

788   30.   Levy SF, Blundell JR, Venkataram S, Petrov DA, Fisher DS, Sherlock G. Quantitative

789   evolutionary dynamics using high-resolution lineage tracking. Nature.

790   2015;519(7542):181–6.

791   31.   Blundell JR, Levy SF. Beyond genome sequencing: Lineage tracking with barcodes to

792   study the dynamics of evolution, infection, and cancer. Genomics [Internet].

793   2014;104(6):417–30. Available from: http://dx.doi.org/10.1016/j.ygeno.2014.09.005

794   32.   Venkataram S, Dunn B, Li Y, Agarwala A, Chang J, Ebel ER, et al. Development of a

795   Comprehensive Genotype-to-Fitness Map of Adaptation-Driving Mutations in Yeast. Cell.

796   2016;166(6):1585-1596.e22.

797   33.   Li Y, Venkataram S, Agarwala A, Dunn B, Petrov DA, Sherlock G, et al. Hidden Complexity

798   of Yeast Adaptation under Simple Evolutionary Conditions. Curr Biol [Internet].

799   2018;28(4):515-525.e6. Available from: https://doi.org/10.1016/j.cub.2018.01.009

800   34.   Kao KC, Sherlock G. Molecular characterization of clonal interference during adaptive

801   evolution in asexual populations of Saccharomyces cerevisiae. Nat Genet.

802   2008;40(12):1499–504.

803   35.   Sniegowski PD, Dombrowski PG, Fingerman E. Saccharomyces cerevisiae and

804   Saccharomyces paradoxus coexist in a natural woodland site in North America and

805   display different levels of reproductive isolation from European conspecifics. FEMS Yeast

806   Res. 2002;1(4):299–306.

807   36.   Magtanong L, Andersen RJ, Porter J, Barker SL, Nislow C, Boone C, et al. A molecular

808   barcoded yeast ORF library enables mode-of-action analysis of bioactive compounds. Nat

809   Biotechnol. 2009;27(4):369–77.

810   37.   Huxley C, Green ED, Dunbam I. Rapid assessment of S. cerevisiae mating type by PCR.

811   Trends Genet. 1990;6(C):236.

812   38.   Cha RS, Thilly WG. PCR Methods and Applications: Specificity, Efficiency, and Fidelity of

813   PCR. Genome Res [Internet]. 1993;S18–30. Available from:

814   https://genome.cshlp.org/content/3/2/S8.full.pdf

815   39.   Hartl DL, Clark AG. Principles of Population genetics. Vol. 116. Sunderland, MA: Sinauer

816   associates; 1997.

817   40.   R Core Team. R: A language and environment for statistical computing [Internet]. Vienna,

818   Austria; 2012. Available from: http://www.r-project.org/

819   41.   Hadley Wickham. The Split-Apply-Combine Strategy for Data Analysis. J Stat Softw

820   [Internet]. 2011;40(1):1–29. Available from:

821   http://www.ncbi.nlm.nih.gov/pubmed/12345678

822   42.   Bates D, Mächler M, Bolker B, Walker S. Fitting Linear Mixed-Effects Models using lme4. J

823   Stat Softw [Internet]. 2015;67(1). Available from: http://arxiv.org/abs/1406.5823

824   43.   Kuznetsova A, Brockhoff PB, Christensen RHB. lmerTest Package: Tests in Linear Mixed

825   Effects Models . J Stat Softw. 2017;82(13).

44.   Pasek J. Weights: Weighting and Weighted Statistics. 2018.

45.   Champely S. pwr: Basic Functions for Power Analysis. 2018.

46.   Wikham H. ggplot2: Elegant Graphics for Data Analysis. 1st ed. New York: Springer-Verlag

New York; 2009.

47.   Ludecke D. sjPlot: Data Visualization for Statistics in Social Science [Internet]. 2019.

Available from: https://cran.r-project.org/package=sjPlot

48.   Auguie B. gridExtra: Miscellaneous Finctions for "Grid" Graphics. 2017.

49.   Sinha R, Stanley G, Gulati GSG, Ezran C, Travaglini KJK, Wei E, et al. Index Switching

Causes "Spreading-Of-Signal" Among Multiplexed Samples In Illumina HiSeq 4000 DNA

Sequencing. bioRxiv [Internet]. 2017;125724. Available from:

https://www.biorxiv.org/content/early/2017/04/09/125724

50.   Costello M, Fleharty M, Abreu J, Farjoun Y, Ferriera S, Holmes L, et al. Characterization

and remediation of sample index swaps by non-redundant dual indexing on massively

parallel sequencing platforms. BMC Genomics. 2018;19(1):1–10.

51.   Illumina. Effects of Index Misassignment on Multiplexing and Downstream Analysis.

Illumina. 2017;1–4.

52.   Kryazhimskiy S, Rice DP, Jerison ER, Desai MM. Global Epistasis Makes Adaptation

Predictable Despite sequence-level stochasticity. Science (80-). 2014;344(6191):1519–

22.

53.   Barrick JE, Kauth MR, Strelioff CC, Lenski RE. Escherichia coli rpoB mutants have

increased evolvability in proportion to their fitness defects. Mol Biol Evol.

2010;27(6):1338–47.

54.   Jerison ER, Kryazhimskiy S, Mitchell JK, Bloom JS, Kruglyak L, Desai MM. Genetic variation

in adaptability and pleiotropy in budding yeast. Elife. 2017;6.

55. MacLean RC, Perron GG, Gardner A. Diminishing returns from beneficial mutations and

pervasive epistasis shape the fitness landscape for rifampicin resistance in Pseudomonas

aeruginosa. Genetics. 2010;186(4):1345–54.

56. Hughes BS, Cullum AJ, Bennett AF. An Experimental Evolutionary Study on Adaptation to

Temporally Fluctuating pH in *Escherichia coli*. Physiol Biochem Zool [Internet].

2007;80(4):406–21. Available from:

http://www.journals.uchicago.edu/doi/10.1086/518353

57. Barrick JE, Lenski RE. NIH Public Access. Spring. 2010;(517):119–29.

58. Gallet R, Cooper TF, Elena SF, Lenormand T. Measuring selection coefficients below 10 -3:

Method, Questions, and Prospects. Genetics. 2012;190(1):175–86.

59. Pierce SE, Davis RW, Nislow C, Giaever G. Genome-wide analysis of barcoded

saccharomyces cerevisiae gene-deletion mutants in pooled cultures. Nat Protoc.

2007;2(11):2958–74.

60. Cooper TF, Lenski RE. Experimental evolution with E. coli in diverse resource

environments. I. Fluctuating environments promote divergence of replicate populations.

BMC Evol Biol. 2010;10(11).

61. Lo YMD, Mehal WZ, Fleming KA. False-Positive Results and the Polymerase Chain

Reaction. Lancet. 1988;332(8612):679.

62. Gerstein AC, Otto SP. Cryptic fitness advantage: Diploids invade haploid populations

despite lacking any apparent advantage as measured by standard fitness assays. PLoS

One. 2011;6(12).

63. Park WK, Yang JW, Kim HS. Identification of novel genes responsible for salt tolerance by

transposon mutagenesis in Saccharomyces cerevisiae. J Ind Microbiol Biotechnol.

2015;42(4):567–75.

64. Tekarslan-Sahin SH, Alkim C, Sezgin T. Physiological and transcriptomic analysis of a salt-

resistant saccharomyces cerevisiae mutant obtained by evolutionary engineering. Bosn J

Basic Med Sci. 2018;18(1):55–65.

65. Blomberg A. Global changes in protein synthesis during adaptation of the yeast

Saccharomyces cerevisiae to 0.7 M NaCl. J Bacteriol. 1995;177(12):3563–72.

66. Minty JJ, Lesnefsky AA, Lin F, Chen Y, Zaroff TA, Veloso AB, et al. Evolution combined with

genomic study elucidates genetic bases of isobutanol tolerance in Escherichia coli.

Microb Cell Fact. 2011;10:1–38.

67. Reyes LH, Abdelaal AS, Kao KC. Genetic Determinants for n -Butanol Tolerance in Evolved

Escherichia coli Mutants: Cross Adaptation and Antagonistic Pleiotropy between n -

Butanol and Other Stressors. Appl Environ Microbiol. 2013;79(17):5313–20.

68. Ostman B, Hintze A, Adami C. Impact of epistasis and pleiotropy on evolutionary

adaptation. Proc R Soc B Biol Sci. 2012;279(1727):247–56.

69. McGee LW, Sackman AM, Morrison AJ, Pierce J, Anisman J, Rokyta DR. Synergistic

pleiotropy overrides the costs of complexity in viral adaptation. Genetics.

2016;202(1):285–95.

70. Hughes KA, Leips J. Pleiotropy, constraint, and modularity in the evolution of life

histories: insights from genomic analyses. Ann N Y Acad Sci. 2017;1389(1):76–91.

71. Gerrish PJ, Lenski RE. The fate of competing beneficial mutations in an asexual

population. Genetica. 1998;102(107):127–44.

72. De Visser JAGM, Zeyl CW, Gerrish PJ, Blanchard JL, Lenski RE. Diminishing returns from

mutation supply rate in asexual populations. Science (80-). 1999;283(5400):404–6.

73. De Visser JAGM, Rozen DE. Clonal interference and the periodic selection of new

beneficial mutations in Escherichia coli. Genetics. 2006;172(4):2093–100.

74. Wahl LM, Gerrish PJ, Saika-Voivod I. Evaluating the impact of population bottlenecks in

experimental evolution. Genetics. 2002;162(2):961–71.

75. Cooper VS. Experimental Evolution as a High-Throughput Screen for Genetic Adaptations.

mSphere. 2018;3(3):1–7.

76. Hegreness M, Shoresh N, Hartl D, Kishony R. An Equivalence Principle for the

Incorporation of Favorable Mutations in Asexual Populations. 2006;311(March):1615–8.

77. Fasanello VJ, Liu P, Botero CA, Fay JC. Data From: High-throughput analysis of adaptation

using barcoded strains of Saccharomyces cerevisiae [Internet]. 2019. Available from:

https://www.ncbi.nlm.nih.gov/bioproject/PRJNA555990/

78. Fasanello VJ. VinceFasanello/MM_Code_Supplement: Code Supplement v1.0.0: High

throughput analysis of adaptation using barcoded strains of Saccharomyces cerevisiae.

2020 Jan 24 [cited 2020 Jan 24]; Available from:

https://doi.org/10.5281/zenodo.3627318

912   DECLARATIONS

914 **Ethics approval and consent to participate**

915 Not applicable

Consent for publication

Not applicable

Availability of data and materials

The dataset supporting the conclusions of this article is available in the NCBI Sequence Read

Archive (SRA) repository, BioProject Number PRJNA555990,

https://www.ncbi.nlm.nih.gob.bioproject/PRJNA555990 [77]. Data formatted for analysis,

intermediate data frames, as well as the custom R scripts utilized for all data processing,

statistical analysis, and figure generation are available from GitHub

(github.com/VinceFasanello/MM_Code_Supplement); a static version of the repository is

available from Zenodo [78]. A readme file is available in the repository with the instructions

necessary to reproduce the analyses and to confirm the results presented in this article.

Supplementary figures, tables, and files referenced throughout the main text are available as

"additional files"; detailed descriptions of all such files can be found in Additional File S1. Strains

are available from the Fay Lab upon request.

Competing interests

The authors have no competing interests.

Funding

V.F. was supported by funding provided by the National Science Foundation Graduate Research

Fellowship Program (NSF-GRFP) (DGE-1745038) and the Botero and Fay labs. J.C.F. was

supported by the National Institutes of Health (GM080669).

Authors' contributions

J.C.F., C.A.B., P.L., and V.J.F conceived of and designed the study. P.L., and V.J.F. conducted the

lab work. V.J.F. processed the data, performed the statistical analysis., and created

visualizations. V.J.F., J.C.F., and C.A.B. drafted and edited the manuscript.

Acknowledgements

The authors thank the members of the Botero and Fay labs, specifically Angela Chira, Trevor

Fristoe, Emery Longan, and Dilys Vela for ongoing feedback and helpful advice.

950   FIGURES

Figure 1

Overview of experimental design. Barcoded yeast strains were used for proof-of-concept fitness

assays (A, 2 days) and experimental evolution (B, 25 days) with gray boxes indicating one day

(10 generations). During the 25-day, 250-generation, experimental evolution (B1), evolutionary

dynamics samples were collected at generations 0, 100, 150, 200, 220, 240, and 250 (cyan

arrows). Fitness of the ancestral (B2) and evolved (B3) strains were quantified via (2-day, 20-

generation) fitness assays (red arrows).

Figure 2

Histogram of fitness from 92 barcoded yeast strains. Fitness is the deviation from the

population mean and was quantified by competition against a common reference strain via

Proof-of-concept fitness assay. Orange and green bars indicate yeast strains with fitness values

significantly and not significantly different from the population mean at a false discovery rate of

5%, respectively.

Figure 3

Power surfaces for POC fitness assays data (green) and fitness change data from the 250-

generation experiment (orange). Power to detect individual barcode fitness deviation from

population mean calculated using error in the POC fitness assay data (A, green) and power to

detect individual barcode fitness change using error in the 250-generation experiment (A,

orange); both employ four replicate measures of fitness. Power to detect treatment-level

fitness differences for two treatments with 22 barcodes in each as estimated using error in the

POC fitness assay data (B, green) and the 250-generation experiment (B, orange).

Figure 4

Change in fitness over 250 generations of experimental evolution. Histogram depicts fitness

increase quantified by competition against a common reference strain. Fitness increase is

generation-250 fitness minus generation-0 fitness. Orange and green bars indicate strains with

fitness values significantly greater or not significantly greater than their own fitness at

generation 0 at a false discovery rate of 1%, respectively. No strains significantly decreased in

fitness. 16/152 entries with large standard error (> 0.029) among replicates are not displayed,

but are included for all statistical testing. Dashed lines indicate the means of each group, with

black indicating the overall mean.

Figure 5

Change in fitness across six treatments in 250 generations of experimental evolution. Violin

plots show the density of 152 yeast strains' change in fitness with treatment labels indicating

the medium, ploidy and dilution rate. Points indicate mean fitness change across 4 replicates

for individual barcodes; point sizes reflect the number of reads underlying each datapoint and

colors indicate evolutionary treatments. Treatment mean fitness changes are depicted as heavy

black crossbars. 16/152 entries with large standard error (> 0.029) among replicates are not

displayed, but are included for all statistical testing. Treatments significantly different from the

reference, diploid yeast evolved under a standard 1:1000 transfer dilution in CM, are marked

with an asterisk.

1000

1001

1002

1003 TABLES

1004

1005 **Table 1**

1006 Associations between evolutionary dynamics and evolutionary treatments.

Measure	Treatment[1]				
	CM + ETOH	CM + NaCl	1:4000 dil.	1:250 dil.	Haploid
t-max	N.S.	-48.702 **	N.S.	N.S.	N.S.
m-max	N.S.	0.292 ***	N.S.	N.S.	0.178 **
t-max-rate	N.S.	N.S.	N.S.	N.S.	-52.717 ***
m-max-rate	N.S.	N.S.	0.006 *	N.S.	N.S.
t-max-diff	N.S.	N.S.	N.S.	N.S.	N.S.
m-max-diff	N.S.	0.175 ***	N.S.	N.S.	0.204 ***
total change	-0.007 *	N.S.	0.018 *	N.S.	-0.007 *

1007

1008 [1]Estimates for significant differences between each evolutionary treatment and the control

1009 (CM-diploid, 1:1000 transfer) are shown for seven different measures of barcoded dynamics (t-

1010 max, m-max, t-max-rate, m-max-rate, t-max-diff, m-max-diff, and total change).

1011

1012 ADDITIONAL FILES

1013

File Name: Additional file 1

File Format: Word Document, .docx

Title of Data: Yeast strains and inserted MOBY barcode sequences.

Description of Data: Columns 1 and 2, resp., contain diploid and haploid yeast strain ID's. Only

yeast strains used in the current project are included (diploids, n = 92; haploids, n = 13).

Columns 3 and 4, resp., contain the MOBY barcode uptag (5'-3') and MOBY barcode downtag

(3'-5-) sequences that uniquely identify the haploid and diploid yeast strains in the same row.

1021

File Name: Additional file 2

File Format: Word Document, .docx

Title of Data: Treatments for 250-generation experimental evolutions.

Description of Data: Treatment ID is a numerical ID for each of six treatments; Evo. Wells is the

number of populations assigned to each treatment; Barcodes per Evo. Well is the number of

sympatric barcodes per evolutionary population; Ploidy, Evo. Medium, and Evo. Transfer

Dilution indicate the yeast strain ploidy, evolutionary medium type, and daily passage dilution

for each treatment, respectively.

1030

File Name: Additional file 3

File Format: Word Document, .docx

Title of Data: Ion Proton sequencing primers.

Description of Data: Oligo Name indicates the name of the forward or reverse primer; Oligo

Names that contain numbers preceded by an "R" are reverse primers. Full Sequence notes the

full Oligo sequence in the 5'-3' direction. Barcode sequence shows the unique genetic barcode

within each primer, in the 5'-3' direction, used in multiplexing and demultiplexing libraries.

1038

File Name: Additional file 4

File Format: Excel document, .xlsx

Title of Data: Estimated fitness change in 250 generations evolution and significance of 92

barcoded yeast strains.

Description of Data: Summary of results for a fixed-0-intercept linear model of mean-corrected

barcoded yeast initial fitness. Predictors are individual barcoded yeast strain ID's. FDR adjusted

p-values for a 2-tailed test are reported.

1046

File Name: Additional file 5

File Format: html document, .html

Title of Data: Data table of power calculations using error in the POC fitness assays data to

support figure 3 panel A.

Description of Data: Data used to generate figure 3 panel A. Columns, in order, are: fitness

change (fitchange), population standard deviation in the POC fitness assays data among

replicate measures of fitness (psd), number of replicates per timepoint (n), power, and

significance level (sig.level). fitchange and n are varied to calculate power; psd and sig.level are

kept constant.

1056

File Name: Additional file 6

File Format: html document, .html

Title of Data: Data table of power calculations using error in the POC fitness assays data to

support figure 3 panel B.

Description of Data: Data used to generate figure 3 panel B. Columns, in order, are: degrees of

freedom in the numerator (u), degrees of freedom in the denominator (v), power, effect size

(f2), and significance level (sig.level). v and f2 are varied to calculate power; u and sig.level are

kept constant.

1065

File Name: Additional file 7

File Format: pdf document, .pdf

Title of Data: barcode cross-contamination rates for sequenced libraries.

Description of Data: A. Histogram of mean barcode contamination rate (as a percentage of total

count) for the cross-contamination diagnostic samples included in the Proof of Concept Fitness

Assays sequenced library. One datapoint is reported for each unique forward-reverse index pair

(sample) in the sequenced library. B. Histogram of mean barcode contamination rate (as a

percentage of total count) for samples included in the 250-generation experimental evolution

project. For resequenced samples, only the less contaminated sample is retained. One

datapoint is reported for each unique forward-reverse index pair (sample) in this consensus

library.

1077

File Name: Additional file 8

File Format: pdf document, .pdf

Title of Data: Decreased contamination after reprocessing samples.

Description of Data: Violin plots showing decrease in barcode contamination rate for a set of

samples from the 250-generation evolution experiment that were DNA extracted, PCR

amplified, and sequenced two separate times. Colors depict sequencing runs: cyan for run 1.,

and orange for run 2. Point sizes reflect the number of reads underlying each datapoint. Mean

contamination rate for run 1., and run 2., are depicted as heavy black crossbars.

1086

File Name: Additional file 9

File Format: html document, .html

Title of Data: Data table of power calculations using error in the change in fitness data for the

250-generation experiment to support figure 3 panel C.

Description of Data: Data used to generate figure 3 panel C. Columns, in order, are: fitness

change (fitchange), population standard deviation in the POC fitness assays data among

replicate measures of fitness (psd), number of replicates per timepoint (n), power, and

significance level (sig.level). fitchange and n are varied to calculate power; psd and sig.level are

kept constant.

1096

File Name: Additional file 10

File Format: html document, .html

Title of Data: Data table of power calculations using error in the change in fitness data for the

250-generation experiment to support figure 3 panel D.

Description of Data: Data used to generate figure 3 panel D. Columns, in order, are: degrees of

freedom in the numerator (u), degrees of freedom in the denominator (v), power, effect size

(f2), and significance level (sig.level). v and f2 are varied to calculate power; u and sig.level are

kept constant.

1105

File Name: Additional file 11

File Format: html document, .html

Title of Data: Change in fitness over 250 generations of experimental evolution.

Description of Data: Summary of results for a fixed-0-intercept linear model of barcoded yeast

fitness change over 250-generations of experimental evolution. Predictors are generation-250

barcoded yeast strains. FDR adjusted p-values for a 1-tailed (greater) test are reported.

1112

File Name: Additional file 12

File Format: Excel document, .xlsx

Title of Data: Standard error in change in fitness across siz treatments in 250 generations of

experimental evolution.

Description of data: summary of results for linear model of standard error in change in

barcoded yeast strain fitness over 250-generations of experimental evolution.

1119

File Name: Additional file 13

File Format: html document, .html

Title of Data: Change in fitness across six treatments in 250 generations of experimental

evolution.

Description of Data: summary of results for linear mixed effects model of change in barcoded

yeast strain fitness over 250-generations of experimental evolution.

1126

File Name: Additional file 14

File Format: html document, .html

Title of Data: t-max by treatment model results.

Description of Data: summary of results for linear mixed effects model of generation of

maximum deviation in barcode abundance from initial conditions over 250-generations of

experimental evolution by treatment.

1133

File Name: Additional file 15

File Format: pdf document, .pdf

Title of Data: t-max by treatment visualization.

Description of Data: Violin plot of generation of maximum deviation in barcode abundance

from initial conditions for 152 yeast strains evolved across six evolutionary treatments for 250

generations. Point sizes reflect the number of reads underlying each datapoint and colors

indicate evolutionary treatments. Treatment means are depicted as heavy black crossbars.

Treatments significantly different from the control treatment are marked with an asterisk. The

treatment with diploid yeast evolved under a standard 1:1000 transfer dilution in CM is

selected as the reference level in this model.

1144

File Name: Additional file 16

File Format: html document, .html

Title of Data: m-max by treatment model results.

Description of Data: summary of results for linear mixed effects model of maximum deviation in

barcode abundance from initial conditions over 250-generations of experimental evolution by

treatment.

1151

File Name: Additional file 17

File Format: pdf document, .pdf

Title of Data: m-max by treatment visualization.

Description of Data: Violin plot of magnitude of maximum deviation in barcode abundance

from initial conditions for 152 yeast strains evolved across six evolutionary treatments for 250

generations. Point sizes reflect the number of reads underlying each datapoint and colors

indicate evolutionary treatments. Treatment means are depicted as heavy black crossbars.

Treatments significantly different from the control treatment are marked with an asterisk. The

treatment with diploid yeast evolved under a standard 1:1000 transfer dilution in CM is

selected as the reference level in this model.

1162

File Name: Additional file 18

File Format: html document, .html

Title of Data: t-max-rate by treatment model results.

Description of Data: summary of results for linear mixed effects model of generation of

maximum rate of change in barcode abundance over 250-generations of experimental

evolution by treatment.

1169

File Name: Additional file 19

File Format: pdf document, .pdf

Title of Data: t-max-rate by treatment visualization.

Description of Data: Violin plot of generation of maximum rate of change in barcode abundance

for 152 yeast strains evolved across six evolutionary treatments for 250 generations. Point sizes

reflect the number of reads underlying each datapoint and colors indicate evolutionary

treatments. Treatment means are depicted as heavy black crossbars. Treatments significantly

different from the control treatment are marked with an asterisk. The treatment with diploid

yeast evolved under a standard 1:1000 transfer dilution in CM is selected as the reference level

in this model.

1180

File Name: Additional file 20

File Format: html document, .html

Title of Data: m-max-rate by treatment model results.

Description of Data: Summary of results for linear mixed effects model of magnitude of

maximum rate of change in barcode abundance over 250-generations of experimental

evolution by treatment.

1187

File Name: Additional file 21

File Format: pdf document, .pdf

Title of Data: m-max-rate by treatment visualization.

Description of Data: Violin plot of magnitude of maximum rate of change in barcode abundance

for 152 yeast strains evolved across six evolutionary treatments for 250 generations. Point sizes

reflect the number of reads underlying each datapoint and colors indicate evolutionary

treatments. Treatment means are depicted as heavy black crossbars. Treatments significantly

different from the control treatment are marked with an asterisk. The treatment with diploid

yeast evolved under a standard 1:1000 transfer dilution in CM is selected as the reference level

in this model.

1198

File Name: Additional file 22

File Format: html document, .html

Title of Data: t-max-diff by treatment model results.

Description of Data: Summary of results for linear mixed effects model of generation of

maximum difference in sympatric barcode abundance over 250-generations of experimental

evolution by treatment.

1205

File Name: Additional file 23

File Format: pdf document, .pdf

Title of Data: t-max-diff by treatment visualization.

Description of Data: Violin plot of generation of maximum difference in sympatric barcode

abundance for 152 yeast strains evolved across six evolutionary treatments for 250

generations. Point sizes reflect the number of reads underlying each datapoint and colors

indicate evolutionary treatments. Treatment means are depicted as heavy black crossbars.

Treatments significantly different from the control treatment are marked with an asterisk. The

treatment with diploid yeast evolved under a standard 1:1000 transfer dilution in CM is

selected as the reference level in this model.

1216

File Name: Additional file 24

File Format: html document, .html

Title of Data: m-max-diff by treatment model results.

Description of Data: Summary of results for linear mixed effects model of magnitude of

maximum difference in sympatric barcode abundance over 250-generations of experimental

evolution by treatment.

1223

File Name: Additional file 25

File Format: pdf document, .pdf

Title of Data: m-max-diff by treatment visualization.

Description of Data: Violin plot of magnitude of maximum difference in sympatric barcode

abundance for 152 yeast strains evolved across six evolutionary treatments for 250

generations. Point sizes reflect the number of reads underlying each datapoint and colors

indicate evolutionary treatments. Treatment means are depicted as heavy black crossbars.

Treatments significantly different from the control treatment are marked with an asterisk. The

treatment with diploid yeast evolved under a standard 1:1000 transfer dilution in CM is

selected as the reference level in this model.

1234

File Name: Additional file 26

File Format: html document, .html

Title of Data: Total change in barcode abundance by treatment model results.

Description of Data: summary of results for linear mixed effects model of total (cumulative)

change in sympatric barcode abundance over 250-generations of experimental evolution by

treatment.

1241

File Name: Additional file 27

File Format: pdf document, .pdf

Title of Data: Total change in barcode abundance by treatment visualization.

Description of Data: Violin plot of total (cumulative) change in sympatric barcode abundance

for 152 yeast strains evolved across six evolutionary treatments for 250 generations. Point sizes

reflect the number of reads underlying each datapoint and colors indicate evolutionary

treatments. Treatment means are depicted as heavy black crossbars. Treatments significantly

different from the control treatment are marked with an asterisk. The treatment with diploid

yeast evolved under a standard 1:1000 transfer dilution in CM is selected as the reference level

in this model.

A

B

Experiment

● POC Fitness Assays

● 250 Generation Evolution Fitness Assays