Quantifying and engineering probiotic bacterial adhesion to mucus

Zachary J. S. Mays[1], Todd C. Chappell[1], and Nikhil U. Nair[1*]

[1]Department of Chemical and Biological Engineering, Tufts University, Medford, MA, 02155

[*]Correspondence: nikhil.nair@tufts.edu or @nair_lab

KEYWORDS

Lactic acid bacteria; probiotics; microbiome; cell adhesion; mucus; mucoadhesion

ABSTRACT

Mucus in the gastrointestinal (GI) tract is the primary point-of-interaction between humans and their gut

microbiota. This not only intimates that mucus ensures protection against endogenous and exogenous opportun-

ists but provision for the human microbiota to reside and flourish. With the emergence of living therapeutics, en-

gineered microbes can deliver and produce increasingly complex medicine, and controlling the mucoadhesive

properties of different microbial chassis can dictate dose-response in a patient. Here we present a redesigned, *in*

vitro, plate-based assay to measure the mucus adhesion of various probiotics. Cell-mucus interactions were isolat-

ed by immobilizing mucus to the plate surface. Binding parameters were derived for each probiotic strain by

measuring cell adhesion over a wide range of cell concentrations, providing dose-dependent adhesion metrics.

Surface proteins and cell components known to influence mucoadhesion were then heterologously expressed or

altered in *Lactococcus lactis* MG1363 and *E. coli* Nissle 1917 to control mucus-binding capacity, avidity, and

cooperativity.

INTRODUCTION

As a new era of precision medicine and personalized therapeutics becomes increasingly mainstream, there too

is heightened interest in our ability to leverage the human gut microbiota, a complex and rich community of ap-

proximately 1000 species of microbes living in the gastrointestinal (GI) tract. Although causality remains uncertain,

recent evidence suggests a strong relationship between many disease states and a perturbed, dysbiotic, gut

microbiota (Cho and Blaser, 2012; Shreiner, Kao and Young, 2015). One strategy to restore microbial balance, and

the basis of 282 clinical trials in the last decade (US National Library of Medicine, 2018), is to introduce probiotics

– microbes shown to confer health benefits. Probiotics have been shown to inhibit or exclude enteric pathogens

(Ollech et al., 2016; Shafaghi et al., 2016), modulate or improve mucosal barrier function (Mennigen et al., 2009),

reduce inflammatory responses (Gionchetti et al., 2003), ameliorate metabolic or toxin imbalances (Krishnan et al.,

2018), maintain intestinal pH, and contribute to neurohumoral signaling (Hyland and Stanton, 2016). Furthermore,

engineered probiotics as therapeutic delivery vehicles (Waller et al., 2017; Bober, Beisel and Nair, 2018; Mays and

Nair, 2018) are being explored for pathogen eradication (Hwang et al., 2017), vaccines (Rosales-Mendoza, Angulo

and Meza, 2016), metabolic disorders, enzyme-replacement therapies (Isabella et al., 2018; West et al., 2019), can-

cer treatments (Zhang et al., 2012; Chowdhury et al., 2019), and diagnostics (Danino et al., 2015; Riglar et al.,

2017).

Somewhat counterintuitively, the stability of a healthy gut microbiota likely arises from its diversity and com-

petitiveness rather than established cooperative metabolic networks (Coyte, Schluter and Foster, 2015). Thus, an

argument can be made that when developing probiotic therapeutics, one design focus should be to enhance fac-

tors conferring competitive opportunity (of course, while safeguarding against overgrowth). This dichotomy is fur-

ther exemplified by the fact that lactic acid bacteria (LAB), which encompass the majority of common probiotics

studied for disease amelioration, make up less than 1% of the human microbiota (Douillard and de Vos, 2014).

Furthermore, orally-administered probiotics are rarely detected past a week after ingestion, and persistence is

highly strain- and patient-specific (Derrien and van Hylckama Vlieg, 2015), making dose-response relationships

unreliable (Ouwehand, 2017). From an engineering perspective, directing a microbe toward a synthetic application

inherently places a burden that conciliates competitive advantage against its natural counterpart in the body; es-

sentially, current strategies cannot compete with the autochthonous members of the microbiota.

The most prominent point-of-interaction between microbes and humans is at mucus membranes, revealing

mucus adhesion as a prime target for controlling probiotic occupancy. Indeed, the cell-surface structures of com-

mensal bacteria are hypothesized to have coevolved in the presence of mucus glycans, creating preferential

mucoadhesive mechanisms (Boekhorst *et al.*, 2006). Additionally, the extracellular polysaccharides (EPS) secreted

for LAB biofilm formation can influence colonization through mucus associations (Lebeer *et al.*, 2009; Salas-Jara *et*

al., 2016), while pathogenic biofilm formation is impeded by LAB extracellular polysaccharides and mucus itself

(Caldara *et al.*, 2012). Mucin has also been observed to increase surface-protease expression in LAB, suggesting

the utilization of mucus as a nutrient source (Wickström *et al.*, 2013). Despite these interactions and the mucus

matrix of the GI tract providing a topography for microbes to find spatial niches, orally-administered natural pro-

biotics and engineered microbial therapeutics do not currently provide any means of controlling biogeographical

targeting, residence time, or an ability to colonize the gut. The ability to accurately measure and engineer mucus

adhesion could address these limitations and provide a new avenue to increase efficacy of natural and engineered

probiotic therapies.

Methodologies to quantify bacterial adhesion to mucus have been adapted from traditional cell adhesion as-

says and inherit similar advantages and disadvantages (Mackie *et al.*, 2017). The general procedure behind these

adhesion assays is to quantify the fraction of residual bacteria bound to mucus-functionalized surfaces after

washes (Khalili and Ahmad, 2015). Surfaces are often modified by incubating with mucus or by culturing gut epi-

thelial cell/organ tissue. However, recent studies showed that bacteria adhere to polystyrene, the support of mu-

cus functionalization, as well or better than mucin- or cell culture-coated microplate wells (Laparra and Sanz,

2009). Additionally, there are several examples of bacterial strains showing different adhesion characteristics de-

pending on the mucus source (Li *et al.*, 2008; MacKenzie *et al.*, 2010; Le *et al.*, 2013), and different mucus composi-

tional profiles can be observed in co-culture models simply by differentiating the HT29 cell line with either metho-

trexate (MTX) or 5-fluorouracil (FU) (Leteurtre *et al.*, 2004). The question arises then, whether these adhesion data

are the result of the mucus composition affecting cell-cell associations, cell-mucus associations, or mucus-plastic

associations. Measuring more holistic binding parameters when quantifying mucus adhesion would allow for bet-

ter comparability across conditions.

In order to address these disconnects, we report an improved *in vitro* method for quantifying bacterial adhe-

sion to mucus. Rather than relying on natural nonspecific interactions to coat a microplate with mucus, we cova-

lently functionalized surfaces with mucus, providing robust and full coverage. Our assay is able to isolate the cell

adhesion events associated directly with mucus. We then measure adhesion over a range of cell concentrations to

develop strain-specific mucus-binding curves. By more clearly describing mucus adhesion using global quantifiers,

we can better compare various binding characteristics related to binding capacity, binding avidity, and

cooperativity for common probiotic strains as well as engineered strains. We also demonstrate how the expression

of recombinant characterized and putative mucus-binding proteins, or deletion thereof, can be used to alter mu-

cus-binding characteristics of probiotics and how these changes are reflected in the binding curves. Such altered

mucus-binding has never been engineered previously. Overall, our methods and results provide a novel and ro-

bust approach to quantifying bacterial adhesion to mucus and advances design of microbial therapeutics with

tunable pharmacological release factors.

RESULTS AND DISCUSSION

Silanization and EDC-coupling covalently binds mucus to the plate surface

A comparison of traditional *in vitro* models used to study bacterial adhesion to the intestinal epithelium re-

vealed that differences between bacterial strains binding to mucus are significantly confounded by their ability to

strongly stick to the abiotic surface of the microplate (Laparra and Sanz, 2009). To remedy this, we sought to cova-

lently link mucus to the plate, providing full mucus coverage and removing cell-plate interactions. To perform this,

the microplate surface was silanized with (3-aminopropyl)triethoxysilane (APTES) to provide a surface of available

primary amine groups. The most abundant mucin protein in the small intestine is Muc2, which is dominated by

sialylated and sulfated glycans, providing freely accessible carboxyl groups (Larsson *et al.*, 2009; Koropatkin, Cam-

eron and Martens, 2012). Using the primary amine on the plate and available carboxyl groups of terminal glycans

of PIM, porcine intestinal mucus (PIM) was then crosslinked to the silanized microplate using a common *N*-ethyl-

N′-(3-dimethyl-aminopropyl) carbodiimide and *N*-hydroxysuccinimide (EDC/NHS) reaction mixture (Figure 1).

Previous reports indicate that it is possible to immobilize mucus using EDC-coupling, wherein mucus provides

primary amine groups for the reaction (Uchida *et al.*, 2004). Because our proposed reaction scheme requires mu-

cin to donate carboxyl groups, we wanted to confirm that EDC-coupling was indeed possible with the plate sur-

face and that the reaction was not hindered by a self-reaction involving amines within the mucins themselves. To

accomplish this, we performed two EDC reactions in tandem. First, between sulfo-cyanine5 amine (cya5) and PIM

in which mucus provides carboxyl groups and a subsequent filtration removes any unbound fluorophore. Second,

between PIM and the microplate in which vigorous wash steps can separate non-specifically bound mucus from

that which is covalently attached to the plate. Fluorescence was observed before and after washing, and the pres-

ence of mucus was confirmed by a follow-up Muc2-specific enzyme-linked immunosorbent assay (ELISA) (Figure

2a). As shown in Figure 2b, when both EDC-coupling reactions are performed, not only is fluorescence observed

initially, suggesting covalent linking to the mucus, but it remains after multiple washes, suggesting the mucus is

stably linked to the microplate surface. Absence of the EDC reagent at either stage results in no fluorescence or

plate binding, as expected. Similar results were observed from the subsequent Muc2-specific ELISA (Figure 2c).

These results confirm the directionality of the EDC-coupling reaction and that mucus was covalently bound to the

microtiter plate. Covalently bound mucus provided further opportunities for optimizing the cell adhesion assay,

including automated wash steps.

Modified microtiter plate assay reveals dependence of cell adhesion on mucus type

Bacterial strains isolated from mammalian intestines have been reported to preferentially bind mucus over

other general proteins such as bovine serum albumin (BSA) (MacKenzie *et al.*, 2010). Unfortunately, when using

the traditional cell adhesion assay to test the mucoadhesive ability of *Lactobacillus reuteri* ATCC PTA-6475 at

OD_{600} 0.5, we were not able replicate preferential binding to mucus over BSA (Figure 2d). Only after implementing

our newly modified covalent immobilization, did we observe significant differences between binding to polysty-

rene plate, BSA, porcine gastric mucus (PGM), and PIM by *L. reuteri* ATCC PTA-6475. Specifically, there was a larger

percent of adhered *L. reuteri* ATCC PTA-6475 cells to PIM than the plate, BSA, and PGM (Figure 2d), potentially

recapitulating this strain's co-evolution with, and preferential binding to, intestinal niches. It has been previously

reported that mucus gel reconstituted from commercially available PGM does not accurately replicate the pH-dependent rheological characteristics of native mucus (Celli *et al.*, 2007) and has an altered structure, likely from industrial processing (Wagner, Wheeler and Ribbeck, 2018). Sialic acid content is an important determinant of the viscosity and elasticity of mucus (Zanin *et al.*, 2016), which may explain the diminished cell adhesion capacity observed with PGM (Figure 2d).

We obtained similar preferential binding to covalently immobilized PIM at a high cell density (OD_{600} 10.0) of *L. reuteri* ATCC PTA-6475 while the traditional assay demonstrated equivalent and non-specific maximum binding capacity of 35% under all conditions (Figure 2d). This was likely observed because the more vigorous washing required at higher cell loadings removes non-covalently attached proteins from the plastic surface. Additionally, while *L. reuteri* ATCC PTA-6475 did not appear to bind any better to PGM than BSA or the plate at the low cell density, the binding capacity significantly increased at the higher cell density, becoming significantly higher than the 35% maximum threshold of the uncoated plate (Figure 2d). It can then be inferred that covalently immobilized mucus enables detection of specific cell-mucus interactions rather than non-specific cell-plastic binding events that dominate with non-covalently functionalized plates. A final important aspect of the new assay is a higher throughput and automated wash procedure, adding consistency between technician operability without sacrificing replicate variability.

Different probiotics display vastly different mucoadhesive characteristics

Cell mucoadhesion is typically reported as a percent or fractional residual bound cells at a single cell concentration, often at OD_{600} 0.5 (Jonsson, Ström and Roos, 2001; Roos and Jonsson, 2002; Li *et al.*, 2008; MacKenzie *et al.*, 2010; Etzold *et al.*, 2014; Jensen *et al.*, 2014). To assess binding characteristics in greater resolution, we measured adhesion of several probiotic strains and species across a wide range of cell concentrations (OD_{600} 0.01 to 100). While there are some previous reports measuring dose-dependent adhesion of probiotics, these were either performed using a Caco2 monolayer (Lee *et al.*, 2000), therein not directly measuring mucus adhesion, or performed at very low cell concentrations (OD_{600} < 0.1), likely due to the limited dynamic range of the traditional assay (Mukherjee and Ramesh, 2015). With our more robust immobilized mucus plate, we aimed to develop a more

quantitative and descriptive mucoadhesive measurement by expanding the dose range, introducing shear stress in

the form of orbital plate shaking, and fitting a sigmoidal curve to the adhesion data to make direct comparisons

without the need to normalize between strains. With many cell-mucus interactions determined to be lectin-like

recognition of mucus glycans by surface proteins (Tallon *et al.*, 2007; Petrova *et al.*, 2016; Valeriano *et al.*, 2016;

Latousakis *et al.*, 2019), a sigmoidal Hill-type shape (Hill, 1910) was chosen with the hypothesis that cell-mucus

binding mirrors the energetics of ligand-receptor binding.

Using this method, distinct strain-specific differences between the mucoadhesive abilities of common probiot-

ics become apparent (Figure 3). This was particularly interesting for *L. reuteri* human isolates. *L. reuteri* ATCC PTA-

6475 and *L. reuteri* ATCC PTA-5289, which are nearly identical at the genome level (Walter, Britton and Roos, 2011)

clustered in the MLSA (multilocus sequence analysis) lineage II phylogenic group (Oh *et al.*, 2010). Notably, these

strains also contain variations of cell and mucus-binding protein A (CmbA) (Jensen *et al.*, 2014) and have been

previously reported to have strong adhesion to mucus and intestinal epithelial cells (IEC) (Jensen *et al.*, 2012). In

contrast, *L. reuteri* DSM 17938 is from a different phylogenic branch (lineage VI) and has a much lower reported

adherence to IECs (Jonsson, Ström and Roos, 2001), which concurs with our observation that it attains a low max-

imum adherence capacity (A_{max} = 4.3×10^7 cells/cm^2).

Escherichia coli Nissle 1917 (EcN) is an extensively studied probiotic sold over-the-counter as Mutaflor® to

treat chronic inflammatory and infectious intestinal diseases and is significantly used in the development of mi-

crobial therapeutics (Lasaro *et al.*, 2009; Hancock, Dahl and Klemm, 2010; Joeres-Nguyen-Xuan *et al.*, 2010; Maltby

et al., 2013; Ou *et al.*, 2016; Scaldaferri *et al.*, 2016; West *et al.*, 2019). Though there are conflicting reports whether

EcN can persist in human intestines (Anderson *et al.*, 2003; Hancock, Dahl and Klemm, 2010), it has been shown to

colonize the porcine intestine al tract. Surprisingly, therefore, our results inicate a surprisingly poor ability to ad-

here to PIM, as determined by this assay (A_{max} 12.9×10^7 cells/cm^2). Like *L. reuteri* DSM 17938, its binding plateaus

to its maximum relatively easily (Table 1). It has been suggested that biofilm-like aggregation is necessary for EcN

colonization (Kankainen *et al.*, 2009) and that mucoadhesion is influential in initial binding before other competi-

tive mechanisms take over. Additionally, compared to pathogenic strains, commensal *E. coli* have more flexibility

in their ability to perform metabolic shifts depending on the mono- and disaccharide carbon sources available

from mucin glycans (Fabich *et al.*, 2008). This suggests commensal *E. coli*, such as EcN, may utilize mucus by find-

ing a nutritional niche rather than adhering as a persistence mechanism.

Another commonly used clinical probiotic is *Lactobacillus rhamnosus* GG (ATCC 53103), which has been report-

ed to have strong mucoadhesion primarily through SpaCBA pilus-mediated interactions (Nishiyama, Sugiyama

and Mukai, 2016). Notably, both EcN and *L. rhamnosus* GG show moderate mucus-binding ability and use moon-

lighting proteins with the secondary function of mucus adhesion but have quite different binding characteristics.

While the probiotic nature of *Weissella confusa* is in question (Fairfax, Lephart and Salimnia, 2014), it was observed

to have a similar binding capacity (A_{max} 45 $\times 10^7$ cells/cm^2) to *L. rhamnosus* GG, which corroborates a report of the-

se strains binding strongly to HT29-MTX, a mucus-producing differentiated epithelial cell line (Ayeni *et al.*, 2011).

Finally, *Lactococcus lactis* MG1363 is undoubtedly the most well-studied LAB and the basis of centuries of food

engineering. Importantly, *L. lactis* MG1363 showed poor binding ability to PIM compared to other probiotic strains

(Figure 3g). Despite similar curve parameters to *L. reuteri* DSM 17938, *L. lactis* MG1363 has a higher sigmoidal co-

efficient, creating a tighter window of concentrations in which mucus binding is influential compared to other

strains.

Expression or removal of surface proteins can be used to alter adhesion characteristics

There are several surface proteins characterized on LAB that either moonlight or directly mediate

mucoadhesion (Jensen *et al.*, 2014). A database search revealed a number of known and putative mucus-

associated proteins found on the *L. reuteri* strains being evaluated. Constitutively expressing three of these pro-

teins in *L. lactis* MG1363 highly, moderately, and minimally altered the stain mucoadhesion characteristics.

Cell and mucus-binding protein (CmbA) has been reported to significantly impact the ability of *L. reuteri* ATCC

PTA-6475 to bind to Caco-2 cells and PIM (Boekhorst *et al.*, 2006). The protein also contains a 288 base pair tan-

dem repeat region, a characteristic found in surface proteins associated with mucus binding in various LAB (Bâth

et al., 2005). When expressing CmbA on the surface of *L. lactis* MG1363, the mucus-binding ability of the strain

drastically improved, creating a binding curve with similar parameters to that of *L. reuteri* ATCC PTA-6475 (Figure

4a and Table 2).

Despite the observed poor mucus-binding ability of *L. reuteri* DSM 17938, the cell surface of *L. reuteri* SD2112,

the genetically equivalent parental strain, has been characterized with multiple identified putative mucus binding

proteins, including Lr0793 and Lr1064 (Aleljung *et al.*, 1994; Roos *et al.*, 1996). The protein Lr0793 is an ATP-

binding cassette (ABC)-transporter component of an amino acid transporter, homologous to collagen-binding

protein (CnBP) and mucus adhesion promoting protein A (MapA) with 97% sequence identity. These homologs

have been described as binding to collagen type I (Rojas, Ascencio and Conway, 2002), gastric mucus (Saraiva *et*

al., 2015), and Caco-2 cells (Båth *et al.*, 2005). However, expressing Lr0793 on the surface of *L. lactis* MG1363 only

slightly improved maximum adherence capacity to PIM compared to wild-type while OD_{A50} remained unchanged

(Figure 4c and Table 2). Similarly, Lr1064 is an ABC-transporter component of a metal ion transporter and member

of the LraI family (Jenkinson, 1994). These proteins have been reported to be adhesins of oral streptococci (Båth *et*

al., 2005). When expressed on *L. lactis* MG1363, Lr1064 increased the maximum adherence capacity (27.8×10^7

cells/cm^2) while keeping the OD_{A50} similar to wild-type, thus moderately improving mucoadhesion (Figure 4b and

Table 2). Additionally, when testing the binding ability of *L. lactis* expressing Lr1064 on BSA rather than PIM, there

was no observed improved adhesion, reinforcing the observed differences are in fact mucus-specific cell adhesion

caused from surface protein expression (Figure S1). These studies demonstrate binding curved of recombinant *L.*

lactis recapitulates those of the strain from which the mucus binding proteins were sourced.

Expression of each mucus-associated protein was confirmed by Western Blot (Figure S2), and surface display

was confirmed by flow cytometry (Figure S3) using anti-His-tag antibodies. The pSIP expression vectors have pre-

viously shown successful secretion of the reporter nuclease A (NucA) in LAB (Troge *et al.*, 2012). We displayed an

inactive mutant (NucA^{E41Q}) on *L. lactis,* as a display control, and confirmed that this strain demonstrated no im-

provement in mucus binding (Figure 4d), again reiterating that the *L. reuteri* proteins enhance binding in a mucus-

specific manner.

The major mucus adhesin on EcN is considered to be the flagellum, formed with the structural subunit FliC

(Monteiro *et al.*, 2009). Additionally, EcN expresses a cellulose synthase (BscA) necessary for biofilm formation and

adhesion to epithelial cells (Friedlander, Vogel and Aizenberg, 2015). We sought to assess the role of FliC and

BcsA in EcN mucus binding. Somewhat unexpectedly, and counter to previous reports, deleting both genes

(EcN$^{\Delta fliC, \Delta bscA}$) increase the mucoadhesive ability of EcN (Figure 4g), shown by a curve shift up and to the left. This

relationship is reversed, however, when the adhesion of each strain was subsequently tested on polystyrene alone

(Figure S4a). Recently, the *E. coli* flagellum was shown to adhere more tightly to hydrophobic abiotic surfaces than

hydrophilic substrates and may in fact impede other surface adhesins as a tradeoff for mobility (Karlskås *et al.*,

2014). This behavior is consistent with our results in Figure 4g, suggesting the adhesion of EcN to the hydrophilic

mucus surface was improved by removing the flagellum, and it is possible that previous measurements of EcN

mucoadhesion were either confounded by the cells binding other available abiotic surfaces like plastic. These re-

sults emphasize the importance of covalently immobilizing mucus and measuring global binding parameters

across a wide range of cell concentrations. Beyond the initial observation that more EcN$^{\Delta fliC, \Delta bscA}$ cells remain

bound than EcN, the binding parameters show a stark difference in curve shape between the wild-type and

knockout strains, perhaps suggesting different mechanisms of adhesion (Table 2). This is further emphasized when

observing binding on polystyrene alone and with the observation that the removal of the flagellum improves

binding to PIM and BSA indiscriminately (Figure S4). Despite a 46% increase in adhesion at OD$_{600}$ 0.5, EcN$^{\Delta fliC, \Delta bscA}$

retains a similar curve shape to EcN because nonspecific interactions dominate the adhesion of both cell types to

polystyrene (Figure S4a). Curve shapes are altered when protein-protein interactions (mucus-specific or otherwise)

at surfaces dominate (Figure 4h).

CONCLUSIONS

Our improved mucus adhesion assay isolates mucus-specific interactions by immobilizing a confluent layer of

mucus to a plate surface. This removes previously observed variability caused through non-specific adhesion

mechanisms to the polystyrene while retaining the benefits of a minimalistic adhesion assay. The assay is further

improved by implementing automated wash steps for consistency, and, measuring adhesion over a wide range of

cell concentrations facilitates the interpolation of a mucus-binding curve. Curve parameters offer a more global

metric for strain-specific mucoadhesion without the need to normalize to a reference strain. We observe clear

dose-dependent differences in the binding ability of a number of probiotic strains with the curve parameters ex-

tending these relationships beyond the traditional measurement of percent adhesion at one cell density. Im-

portantly, known and putative mucus-associated factors were expressed on the surface of a poor mucus binder, *L.*

lactis MG1363, and mucoadhesion was significantly altered, reinstating the mucus-specific interactions of these

cell-surface elements. Furthermore, a new understanding of EcN mucoadhesion was observed with the polar fla-

gellum, originally considered a major mucoadhesin, actually impeding mucus adhesion and more general hydro-

philic surface adhesion. This emphasizes the advantage of our updated assay over a traditional approach in which

nonspecific interactions with the abiotic plate surface likely confounded mucoadhesion measurements. These re-

sults offer promise in the delivery and control of drug-release in the emerging field of living microbial therapeu-

tics.

SIGNIFICANCE

This work presents an improved *in vitro* adhesion assay to determine mucus-specific adhesion mechanisms of

bacteria. The development of mucus-binding curves directly quantifies strain-specific dose-depending

mucoadhesion with curve parameters acting as a more universal metric for adhesion than the traditional percent

adhesion normalized to a reference strain. This work is a significant advancement for the delivery and control of

therapeutics through various probiotic chassis.

ACKNOWLEDGEMENTS

The authors would like to thank Dr. Huan-Hsuan Hsu and Prof. Xiaocheng Jiang for their advisement and re-

sources when developing the surface immobilization procedure. A special thanks to Josef R. Bober for his advise-

ment with surface-display in LAB. This work was financially supported by grant numbers R03HD090444 and

DP2HD091798 of the National Institutes of Health.

AUTHOR CONTRIBUTIONS

Z.J.S.M. performed all experiments. Z.J.S.M. and N.U.N. designed all experiments, analyzed the data, and wrote the

manuscript. T.C.C. performed gene deletions in EcN.

DECLARATION OF INTERESTS

The authors declare no competing interests

Figure 1. Schematic of mucus immobilization strategy

The structure of mucus presents an opportunity for covalent immobilization chemistry. Gel-forming mucins such

as Muc2 form (a.) macromolecular sheets of trimers of dimers that crosslink or interact to form a mesh. Each mu-

cin monomer has a highly glycosylated proline-threonine/serine (PTS) domain with the (b.) highly sialylated core 3

O-glycan dominating in the intestines. Sialic acid, or N-acetylneuraminic acid, provides a freely available carboxylic

acid that can be (c.) coupled to a primary amine presented on a glass or plastic surface using EDC/NHS crosslink-

ing chemistry.

Figure 2. Validating the modified cell mucoadhesion assay and comparing to the traditional assay

(a.) Two EDC/NHS reactions were performed in tandem followed by a Muc2-specific ELISA to determine the suc-

cess of the mucus immobilization. (b.) Fluorescence microscopy was then implemented to measure of the robust-

ness and directionality of the coupling between mucus and sulfo-cyanine5 and between mucus and the plate sur-

face. Each condition was imaged before and after vigorous washing and after the ELISA. (c.) Luminescence counts

were quantified to determine mucus coverage, with the final optimized condition (1% PIM) measuring 10-fold

higher than other conditions, as expected. (d.) With these final conditions, the traditional adhesion assay was

compared to the newly optimized assay using the strong binder *L. reuteri* ATCC PTA-6475 as a metric for success

on different protein surfaces.

Figure 3. Dose-dependent mucus-binding curves for common probiotics

(a.-g.) Using our optimized plate assay, the number of cells adhered to immobilized mucus was measured over a large range of concentrations for common probiotics. Data points are reported as the mean values of quadruplicate wells across three independent experiments. Hill-type sigmoidal curves were fit, with curve parameters a metric for strain-specific mucoadhesive characteristics (Table 1). (h.) With the traditional assay typically performed at OD_{600} 0.5, comparative analysis across strains was visualized at this density.

Figure 4. Altered dose-dependent mucoadhesion by heterologous protein expression

Known and putative mucus-associating surface proteins from *L. reuteri* were heterologously expressed in *L. lactis*

MG1363. The (e.) wild-type *L. lactis* MG1363 binding curve (orange) was (a.-c.) compared to the engineered coun-

terpart (black) as well as the *L. reuteri* strain from which the protein was sourced (green or blue). (d.) As a display

control, an inactivated NucA[E41Q] did not alter mucus binding compared to the wild-type strain. Alternatively, the

flagellum of EcN, considered a known mucus-binding factor, and cellulose synthase, a contributor to biofilm for-

mation, were (g.) knocked out but surprisingly improved mucoadhesion compared to the (f.) wild-type (pink). (h.)

With the traditional assay typically performed at OD_{600} 0.5, comparative analysis across strains was visualized at

this density.

314 Table 1. Summary of mucus-binding curve parameters for common probiotics

Strain	[a]A_{max} ($\times 10^7$ cells/cm^2)	[b]OD_{A50}	n
L. reuteri ATCC PTA-6475	77.4 ± 3.1	14.3 ± 1.5	0.98 ± 0.05
L. reuteri ATCC PTA-5289	133.5 ± 7.5	30.0 ± 4.1	0.97 ± 0.05
L. reuteri DSM 17938	4.3 ± 0.3	6.2 ± 1.1	1.11 ± 0.11
Weissella confusa	44.7 ± 0.0	[a]203.7 ± 0.1	0.95 ± 0.001
L. rhamnosus GG	44.5 ± 0.3	[a]99.0 ± 0.8	0.80 ± 0.02
E. coli Nissle 1917	12.9 ± 0.2	17.7 ± 0.2	0.98 ± 0.04
L. lactis MG1363	6.8 ± 0.4	3.7 ± 0.4	1.52 ± 0.14

315
316 Parameters determined from sigmoidal curve fitting of experimental adhesion data, shown with 95% confidence
317 intervals

318 [a]Extrapolated value from curve fitting

319 [b]Standard curves to correlate OD_{600} to CFU/mL are reported for each strain in Table S1.

322 Table 2. Summary of mucus-binding curve parameters when altering mucoadhesive protein expression

Strain	[a]A_{max} ($\times 10^7$ cells/cm^2)	OD_{A50}	n
L. lactis MG1363	6.8 ± 0.4	3.7 ± 0.4	1.52 ± 0.14
+ NucA^{E41Q}	5.5 ± 0.4	4.0 ± 0.5	1.5 ± 0.13
+ CmbA	104.3 ± 4.6	11.5 ± 0.8	1.37 ± 0.12
+ Lr1064	27.8 ± 1.7	9.4 ± 0.9	1.64 ± 0.12
+ Lr0793	13.0 ± 0.5	4.1 ± 0.5	1.48 ± 0.22
E. coli Nissle 1917	12.9 ± 0.2	17.7 ± 0.2	0.98 ± 0.04
ΔfliC,ΔbscA	368.3 ± 5.5	[a]230.5 ± 3.6	0.78 ± 0.02

323
324 Parameters determined from sigmoidal curve fitting of experimental adhesion data, shown with 95% confidence
325 intervals

326 [a]Extrapolated value from curve fitting

STAR ⬤ METHODS

⬤ EXPERIMENTAL MODEL AND SUBJECT DETAILS

Microbial strains, plasmids, and growth conditions

All lactobacilli and *Weissella confusa* were cultured in de Man-Rogosa-Sharpe (MRS) medium (RPI Corp, Mt.

Prospect, IL) at 37 °C in static microaerobic conditions. *Lactococcus lactis* subsp. cremoris MG1363 was cultured in

GM17 medium (M17 broth (BD Difco, Franklin Lakes, NJ), 0.5 % (w/v) glucose) at 30 °C in static microaerobic con-

ditions. *E. coli* strains were cultured in lysogeny broth (LB) (VWR International, Randor, PA) at 37 °C with rotary

shaking at 250 rpm. All media was solidified using 1.5 % (w/v) agar (Teknova Inc, Hollister, CA). *E. coli* TG1$\Delta endA$

was used as a host for the construction of the pSIP-based expression vectors listed in Table S2 and cultured as

above only supplemented with erythromycin (200 µg/mL) (RPI Corp). When *L. lactis* subsp. cremoris MG1363 was

used as hosts for these vectors, the strains were cultured as above only supplemented with erythromycin (5

µg/mL).

All cloning was performed in *E. coli* TG1$^{\Delta endA}$. Beginning with the plasmid pSIP411, the inducible promoter

PsppQ (Rud *et al.*, 2006) was replaced with the constitutive synthetic promoter P11 (Mathiesen *et al.*, 2009) using

the surrounding sites for restriction endonucleases BglII and NcoI. The genomic DNA of *L. reuteri* ATCC PTA-6475

and *L. reuteri* DSM 17938 were used as templates to amplify the known and putative mucus-associating factors in

this study using colony PCR. The pSIP411-P11 backbone, *lr1064*, and *lr0793* were digested with NcoI and XhoI

while *cmbA* was digested with PciI and XhoI. A site mutation was introduced into *nucA* using overlap PCR and the

inactive *nucA*^{E41Q} was inserted into a surface-display pSIP411-P11 variant (previously unpublished) by digestion

with NcoI and XmaI. This created a fusion construct of secretion signal peptide lp3050 (Fredriksen *et al.*, 2012), the

N-terminal LysM anchoring motif lp3014 (Fredriksen *et al.*, 2012), and *nucA*^{E41Q}. All cloning enzymes were pur-

chased from New England Biolabs (Ipswich, MA).

 METHOD DETAILS

 Transformation conditions

L. lactis MG1363 transformations require GSGM17 medium (GM17, 0.5 M sucrose, 2.5% (w/v) glycine).

Electrocompetent cells were prepared by first growing in 5 mL GM17 overnight from a frozen stock optimized for

GSGM17 growth. The resulting overnight was subcultured 1:9 in 10 mL GSGM17 and grown overnight. This final

overnight culture was subcultured 1:7 in 40 mL GSGM17 and grown 6-7 h (OD_{600} 0.17). Cells were centrifuged at

3000 × g at 4 °C for 25 min and washed one in 40 mL ice-cold EP buffer (0.5 M sucrose, 10 % (w/v) glycerol). The

cells were then suspended in 20 mL ice-cold EP buffer and incubated for 15 min on ice. Finally, cells were sus-

pended in 400 µL ice-cold EP buffer and stored at −80 °C in 50 µL aliquots. To transform, 500 ng DNA, always di-

luted in dH2O to a final volume of 5 µL, was added to 50 µL electrocompetent cells and incubated on ice for 2

min. This mix was transferred to a pre-chilled 2 mm electroporation cuvette and electroporation was performed

(2.0 kV, 25 µF, 200 Ω) using a Gene Pulser Xcell (Bio-Rad Laboratories, Hercules, CA) electroporation system. Cells

were recovered by immediately adding 950 µL pre-warmed GM17 supplemented with recovery salts (20 mM

$MgCl_2$, 2 mM $CaCl_2$) and incubating statically at 30 °C for 1.5 h. Cells were plated on selective GM17 with erythro-

mycin (5 µg/mL) and grown for at least 24 h at 30 °C.

Isolation of primary porcine intestinal mucus

The intestines of 8-10-week-old pigs were obtained from a local supplier (Research 87 Inc, Boylston, MA). The

small intestine was sectioned into 60 cm segments, squeezed of chyme, and placed in ice-cold phosphate buff-

ered saline (PBS) (137 mM NaCl, 2.7 mM KCl, 10 mM Na_2HPO_4, 2 mM KH_2PO_4, pH 7.4). Each section was then

splayed longitudinally, washed by dipping into a tray of ice-cold PBS, and moved to a second tray of ice-cold PBS.

Pinning one end of an intestinal segment, a silicone spatula was used to scrape the mucus layer, placing the mu-

cus in a beaker of ice-cold PBS. The crude PIM was centrifuged at 11000 × g at 4 °C for 10 min, and the superna-

tant was centrifuged again at 26000 × g at 4 °C for 15 min using an Optima L-90K (Beckman Coulter, Brea, CA)

ultracentrifuge with a 70-Ti fixed angle rotor (Beckman Coulter). This clarified PIM supernatant was then lyophi-

lized and stored at 4 °C.

Plate surface functionalization

The wells of 96-well clear polystyrene U-bottom (Greiner Bio-One, Kremsmünster, Austria) microtiter plates

were activated by oxygen plasma etching using a PE-200 (Plasma Etch, Inc. Carson City, NV) industrial benchtop

plasma processing system with a R301 RF Generator (Seren IPS, Inc., Vineland, NJ) power supply (105 W, 1 min

plasma time, 50 psi vacuum pressure, 20 cc/min O_2 flow). Immediately, 200 µL of APTES (5% v/v in ethanol) was

added to each well, and the plate was sealed with parafilm and incubated at room temperature for 15 min. The

plate was then washed twice with ethanol and dried with an air stream, always minimizing exposure to humidity or

water. A 1% PIM solution was prepared in PBS. Separately, a 10x solution of EDC (Thermo Fisher Scientific, Wal-

tham, MA) was made in PBS. The EDC was added to the 1% PIM to a final concentration of 5 mM and placed on a

rocker for 5 min. During this time, NHS (TCI Chemicals, Portland, OR) was weighed out and added to the 1% PIM

to a final concentration of 10 mM, again using a rocker to mix for 5 min. Finally, 200 µL of the PIM/EDC/NHS was

added to each well of the APTES functionalized plate. The plate was then sealed with parafilm and incubated at

room temperature for 2 h before being left at 4 °C overnight.

Fluorescence microscopy, ELISA, and Western blot analysis

A coupling reaction was prepared as described in the "Plate surface functionalization" section above with a 1%

PIM solution combined with EDC/NHS and mixed for 5 min. Sulfo-cyanine5 amine (cya5) (Lumiprobe, Hunt Valley,

MA) was added to a final concentration of 1 mg/mL, and the reaction was incubated for 2 h at room temperature,

wrapped in foil to prevent light exposure. A Zeba spin desalting column (Thermo Fisher Scientific) was then used

to remove excess fluorophore. The mucus-cya5 conjugate was then used in a second coupling reaction, conjugat-

ing the mucus to a 96-well clear polystyrene U-bottom (Greiner Bio-One) microtiter plate. After a 2 h incubation,

excess volume was aspirated from each well condition. Microscope images were captured using a DMi8 (Leica Mi-

crosystems, Inc., Buffalo Grove, IL) microscope with 100x objective. Phase contrast images were taken with 1 ms

exposure. Fluorescent images were taken by exciting with a Y5 filter cube with 400 ms exposure and gain set to 4 on LAS X (Leica Microsystems, Inc.) software. Wells were then washed three times with a plate washer as described in the "Mucus adhesion assay" section below.

Microscope images were again taken before an ELISA assay was performed. Each well was blocked for 1 h with cold blocking buffer (5% (w/v) skim milk powder in TBST (20 mM Tris, 150 mM NaCl, 0.1% (v/v) Tween, pH 7.6)). Wells were then washed three times with ice cold TBST, shaking at 150 rpm for 5 min each time. Wells were resuspended in 200 µL blocking buffer containing 0.1 µg rabbit anti-Muc2 (MyBioSource, Inc., San Diego, CA) primary antibody and incubated on ice with shaking at 150 rpm for 1 h. After another three washes, the wells were resuspended in 200 µL blocking buffer containing 0.04 µg goat anti-rabbit IgG-HRP (Abcam, Cambridge, MA) secondary antibody and incubated on ice with shaking at 150 rpm for 30 min. Using a SuperSignal West Femto (Thermo Fischer Scientific) chemiluminescence detection kit, images and luminescence values were recorded after 30 s exposure on a UVP ChemiDoc-It2 810 (Analytik Jena US, LLC., Upland, CA) imager on VisionWorks LS (Analytik Jena US, LLC) analysis software.

Flow cytometry

Five milliliter cultures of *L. lactis* MG1363 wild-type or carrying expression vectors were grown overnight as described in the "Microbial strains, plasmids, and growth conditions" section above. Cells were centrifuged at 3000 × g for 5 min and washed twice with PBS before being suspended at OD_{600} 1.0. A 200 µL aliquot was resuspended in 200 µL PBSA (2% (w/v) BSA in PBS) and incubated on ice for 30 min. The cells were then resuspended in 50 µL PBSA containing 1 µg mouse anti-6xHis (Thermo Fisher Scientific) primary antibody and incubated for 1 h at room temperature. The cells were washed three times with PBSA before being suspended in 50 µL PBSA containing 0.4 µg goat anti-mouse IgG-AlexaFluor488 (Thermo Fisher Scientific) secondary antibody and incubated for 15 min on ice, wrapped in foil to prevent light exposure. Finally, the cells were washed three times with PBSA and suspended in 500 µL PBSA. Samples were run on an Attune NxT (Thermo Fisher Scientific) flow cytometer, using a 50 mW blue laser to collect at least 10,000 events. Histograms were analyzed on FCS Express 6 (De Novo Software, Glendale, CA) software.

Gene knockouts in *E. coli* Nissle 1917

Antibiotic cassette insertion and gene knockout (KO) were performed using lambda red recombineering (Datsenko and Wanner, 2000). Kanamycin (kan), chloramphenicol (cat), and ampicillin (amp) were used at 50, 25, and 100 μg/mL, respectively, to maintain plasmids or select for integration of resistance cassettes. Amplification of the cat cassette from pKD3 was used for *bcsA* KO, while amplification of the kan cassette from pKD4 was used for *fliC* KO. Verification of antibiotic resistance cassette integration and removal was verified by Taq colony PCR. Plasmid curing and cassette removal was verified by absence of growth on LB agar plates. All primers for cassette amplification and removal verification are listed in the ★ Key Resources Table.

Fluorescent labelling of bacteria

All strains were grown to early stationary phase in 200 mL of their respective growth media (approximately 10 h for LAB strains). To promote flagellum expression *E. coli* was grown for 24 h at 30 °C at 85 rpm. Cells were then centrifuged at 3000 × g at 4 °C for 10 min and washed three times with 50 mL PBS. Cells were then diluted in PBS to OD_{600} 1.0 and labelled by the addition of carboxyfluorescein diacetate (cFDA) (Thermo Fisher Scientific) to a final concentration of 10 μM by incubating at 37 °C for 40 min with rotary shaking at 85 rpm. Again, carboxyfluorescein (cF)-labelled cells were washed three times as above and suspended in 10 mL PBS.

Mucus adhesion assay

Cells were diluted using PBS from the 10 mL stock to OD_{600} values approximately evenly-spaced across three orders of magnitude. The mucus solution left after plate functionalization was removed from the microtiter plate by aspiration. Two hundred microliters of each dilution were plated in four technical replicates. A separate 225 μL of each OD_{600} dilution was placed in microfuge tubes, and both the plate and tubes were then incubated at 37 °C for 4 h with rotary shaking at 500 rpm. Using an AquaMax4000 (Molecular Devices, San Jose, CA) plate washer with a cell wash head, the plate was washed three times with PBS (center aspiration 50 μL/well/s; wall dispense 7.2 mL/plate/s; probe height 3 mm) ending with a final aspiration step. The adhered cells were lysed by adding 200 μL lysis buffer (1% (w/v) SDS, 0.1 M NaOH) and incubating at 37 °C for 1 h with rotary shaking at 750 rpm. Addition-

ally, the 225 μL cell dilutions were centrifuged at 5000 × g for 5 min and suspended in lysis buffer as fluorescence

controls. All lysates were transferred to a 96-well F-bottom black FluoTrac (Greiner Bio-One) microtiter plate. The

plate was passed over with a flame to remove bubbles and fluorescence was measured (Em/485 nm, Ex/520 nm)

using a SpectraMax M3 (Molecular Devices) plate reader. Blank wells containing only lysis buffer were used for

background subtraction.

▮ QUANTIFICATION AND STATISTICAL ANALYSIS

Adhesion curve fitting and statistical analysis

Plate data was background corrected by subtracting the relative fluorescence units (RFU) of blank wells con-

taining only lysis buffer. The means of experimental replicates were used for further data analysis. By plotting the

standards as RFU vs OD_{600}, the slope of a linear trend (C_1) was used to associate RFU per unit OD_{600}. This relation-

ship, along with the volume-to-surface-area relationship (V/A) derived from a microtiter plate schematic (Figure

S5) and the CFU/mL per OD_{600} (C_2), can be used to convert values to adhered cells per unit area (A_y):

$$A_y = (RFU_y)(C_1{}^{-1})(C_2)(\frac{V}{A})$$

459   Values for C1 and C2 are summarized in Table S1. A typical sigmoidal (Hill-type) curve was adapted to model fit

460   cell adhesion:

$$A_y = \frac{A_{max}OD_y^n}{OD_{A50}^n + OD_y^n}$$

Here, A_y is the adhered cells per unit area at cell concentration y, A_{max} is the maximum possible adhered cells per

unit area, OD is the OD_{600} for cell concentration y, OD_{A50} is the OD_{600} required for half the maximum number of

cells to adhere to the well, and n is the sigmoid factor. The curve parameters were obtained by using the nonlinear

curve fitting function in MATLAB (fitnlm) with robust bi-square least squares regression analysis, which applies a

weighting factor to discount outliers. Plots of cells per unit area versus OD_{600} were created to visualize the adhe-

sion differences between strains.

⬛ KEY RESOURCES TABLE

SUPPLEMENTAL INFORMATION

Figure S1. Adhesion of *L. lactis* MG1363 to different substrates

The observed increase in adhesion of *L. lactis* MG1363 when expressing Lr1064 is shown to be specific to mucus;

adhesion does not improve over wild-type *L. lactis* MG1363 when both are adhered to BSA. Only the curve fit is

shown for clarity.

475

477 His-tag fused (a.) empty vector, (b.) CmbA, (c.) Lr0793, and (d.) Lr1064 were expressed in *L. lactis* MG1363. Bands

478 show the surface-stripped, soluble, and insoluble fractions. Proteins expressed at the correct size (ladder not

479 shown).

Histogram #	Parameter	Median
1	BL1-A	11.00
2	BL1-A	132.00
3	BL1-A	47.00
4	BL1-A	57.00
5	BL1-A	86.00

Figure S3. Flow-cytometry confirms surface display of recombinant mucus-associated factors

His-tag fused mucus-associated factors were labelled with AlexaFluor488 conjugated anti-His-tag antibody. Wild-

type *L. lactis* MG1363 was used as a negative control and inactivated NucA^{E41Q} was used as a positive control for

surface display. A shift to the right on each histogram is an increase in fluorescence, correlated to protein display.

Median fluorescence units are shown in the table for each corresponding histogram.

Figure S4. Adhesion of *E. coli* Nissle 1917 to different substrates

While the deletion of FliC and BscA increased the ability of EcN to bind to mucus, (a.) the relationship was the re-

verse when adherence of wild-type EcN and EcN$^{\Delta fliC, \Delta bscA}$ was quantified on uncoated polystyrene. (b.) Further in-

vestigation shows that EcN$^{\Delta fliC, \Delta bscA}$ seems to indiscriminately bind hydrophilic surfaces as adhesion was increased

on mucus and BSA.

Figure S5. Schematic of Greiner U-bottom plate geometry

Sourced from Greiner 96-Well U-bottom product document and used to calculate well surface area

502 Table S1. Linear relationships between RFU and OD_{600} and OD_{600} and CFU/mL for each strain

	RFU/OD_{600}	CFU/mL ($\times 10^8$) per OD_{600}
L. reuteri ATCC PTA-5289	3109	3.7
L. reuteri ATCC PTA-6475	2794	3.3
L. reuteri DSM 17938	2334	6.7
W. confusa	397	3.8
L. rhamnosus GG	1002	1.4
E. coli Nissle 1917	51	8.0
L. lactis MG1363	7005	6.8

503
504 Values for CFU/mL acquired by growing cells overnight, checking the OD_{600} of three dilutions, and plating a
505 10,000x dilution of those concentrations to count CFUs.

ABBREVIATIONS

ABC, ATP-binding cassette

APTES, (3-aminopropyl)triethoxysilane

BSA, bovine serum albumin

BscA, cellulose synthase catalytic subunit A

cFDA, carboxyfluorescein diacetate

CmbA, cell and mucus binding protein A

CnBP, collagen binding protein

EcN, *E. coli* Nissle 1917

EDC, *N*-ethyl-*N'*-(3-dimethyl-aminopropyl) carbodiimide

ELISA, enzyme-linked immunosorbent assay

EPS, extracellular polysaccharide

FU. 5-fluorouracil

GI, gastrointestinal

IEC, intestinal epithelial cell

LAB, lactic acid bacteria

MapA, mucus adhesion promoting protein A

MTX, methotrexate

NHS, *N*-hydroxysuccinimide

NucA, nuclease A

PGM, porcine gastric mucus

PIM, porcine intestinal mucus

PTS, proline-threonine/serine

RFU, relative fluorescence unit

REFERENCES

Aleljung, P. *et al.* (1994) "Purification of collagen-binding proteins of Lactobacillus reuteri NCIB 11951," *Current Microbiology*, 28(4). doi: 10.1007/BF01575966.

Anderson, G. G. *et al.* (2003) "Intracellular bacterial biofilm-like pods in urinary tract infections," *Science*, 301(5629). doi: 10.1126/science.1084550.

Ayeni, F. A. *et al.* (2011) "Evaluation of the functional potential of Weissella and Lactobacillus isolates obtained from Nigerian traditional fermented foods and cow's intestine," *International Journal of Food Microbiology*. Elsevier B.V., 147(2), pp. 97–104. doi: 10.1016/j.ijfoodmicro.2011.03.014.

Båth, K. *et al.* (2005) "The cell surface of Lactobacillus reuteri ATCC 55730 highlighted by identification of 126 extracellular proteins from the genome sequence," *FEMS Microbiology Letters*. Oxford, UK: Elsevier B.V, 253(1). doi: 10.1016/j.femsle.2005.09.042.

Bober, J. R., Beisel, C. L. and Nair, N. U. (2018) "Synthetic Biology Approaches to Engineer Probiotics and Members of the Human Microbiota for Biomedical Applications," (March), pp. 277–300.

Boekhorst, J. *et al.* (2006) "Comparative analysis of proteins with a mucus-binding domain found exclusively in lactic acid bacteria," *Microbiology*. England: Soc General Microbiol, 152(1). doi: 10.1099/mic.0.28415-0.

Caldara, M. *et al.* (2012) "Mucin biopolymers prevent bacterial aggregation by retaining cells in the free-swimming state," *Current Biology*. Elsevier Ltd, 22(24). doi: 10.1016/j.cub.2012.10.028.

Celli, J. P. *et al.* (2007) "Rheology of gastric mucin exhibits a pH-dependent sol-gel transition," *Biomacromolecules*, 8(5). doi: 10.1021/bm0609691.

Cho, I. and Blaser, M. J. (2012) "The human microbiome: at the interface of health and disease.," *Nature Reviews Genetics*, 13(4). doi: 10.1038/nrg3182.

Chowdhury, S. *et al.* (2019) "Programmable bacteria induce durable tumor regression and systemic antitumor immunity," *Nature Medicine*. Springer US, 25(July). doi: 10.1038/s41591-019-0498-z.

Coyte, K. Z., Schluter, J. and Foster, K. R. (2015) "The ecology of the microbiome: Networks, competition, and stability.," *Science*, 350(6261). doi: 10.1126/science.aad2602.

Danino, T. *et al.* (2015) "Programmable probiotics for detection of cancer in urine," *Science Translational Medicine*, 7(289), p. 289ra84. doi: 10.1126/scitranslmed.aaa3519.

Datsenko, K. A. and Wanner, B. L. (2000) "One-step inactivation of chromosomal genes in Escherichia coli K-12 using PCR products," *Proceedings of the National Academy of Sciences*, 97(12), pp. 6640–6645. doi: 10.1073/pnas.120163297.

Derrien, M. and van Hylckama Vlieg, J. E. T. (2015) "Fate, activity, and impact of ingested bacteria within the human gut microbiota," *Trends in Microbiology*. Elsevier Ltd, 23(6). doi: 10.1016/j.tim.2015.03.002.

Douillard, F. P. and de Vos, W. M. (2014) "Functional genomics of lactic acid bacteria: From food to health," *Microbial Cell Factories*. BioMed Central Ltd, 13(Suppl 1). doi: 10.1186/1475-2859-13-S1-S8.

Etzold, S. *et al.* (2014) "Structural and molecular insights into novel surface exposed mucus adhesins from Lactobacillus reuteri human strains," *Molecular Microbiology*. HOBOKEN: WILEY-BLACKWELL, 92(3). doi: 10.1111/mmi.12574.

Fabich, A. J. *et al.* (2008) "Comparison of carbon nutrition for pathogenic and commensal Escherichia coli strains in
the mouse intestine," *Infection and Immunity*, 76(3), pp. 1143–1152. doi: 10.1128/IAI.01386-07.

Fairfax, M. R., Lephart, P. R. and Salimnia, H. (2014) "Weissella confusa: Problems with identification of an oppor-
tunistic pathogen that has been found in fermented foods and proposed as a probiotic," *Frontiers in Microbiology*,
5(JUN), pp. 1–5. doi: 10.3389/fmicb.2014.00254.

Fredriksen, L. *et al.* (2012) "Surface display of N-terminally anchored invasin by Lactobacillus plantarum activates
NF-κB in monocytes," *Applied and Environmental Microbiology*, 78(16). doi: 10.1128/aem.01227-12.

Friedlander, R. S., Vogel, N. and Aizenberg, J. (2015) "Role of flagella in adhesion of Escherichia coli to abiotic sur-
faces," *Langmuir*, 31(22). doi: 10.1021/acs.langmuir.5b00815.

Gionchetti, P. *et al.* (2003) "Prophylaxis of pouchitis onset with probiotic therapy: A double-blind, placebo-
controlled trial," *European Journal of Gastroenterology and Hepatology*, 15(7). doi: 10.1097/00042737-200307000-
00020.

Hancock, V., Dahl, M. and Klemm, P. (2010) "Probiotic Escherichia coli strain Nissle 1917 outcompetes intestinal
pathogens during biofilm formation," *Journal of Medical Microbiology*, 59(4). doi: 10.1099/jmm.0.008672-0.

Hill, A. v (1910) "The possible effects of aggregation of haemoglobin on its dissociation curves," *J. Physiol. London*,
40.

Hwang, I. Y. *et al.* (2017) "Engineered probiotic Escherichia coli can eliminate and prevent Pseudomonas
aeruginosa gut infection in animal models," *Nature Communications*. Nature Publishing Group, 8. doi:
10.1038/ncomms15028.

Hyland, N. and Stanton, C. (2016) *The gut-brain axis: Dietary, probiotic, and prebiotic interventions on the*
microbiota. London, UK⬜: Academic Press is an imprint of Elsevier.

Isabella, V. M. *et al.* (2018) "Development of a synthetic live bacterial therapeutic for the human metabolic disease
phenylketonuria," *Nature Biotechnology*. Nature Publishing Group, 36(August), pp. 857–867. doi: 10.1038/nbt.4222.

Jenkinson, H. F. (1994) "Cell surface protein receptors in oral streptococci," *FEMS Microbiology Letters*, 121(2). doi:
10.1111/j.1574-6968.1994.tb07089.x.

Jensen, H. *et al.* (2012) "In vitro testing of commercial and potential probiotic lactic acid bacteria," *International*
Journal of Food Microbiology, 153(1–2). doi: 10.1016/j.ijfoodmicro.2011.11.020.

Jensen, H. *et al.* (2014) "Role of Lactobacillus reuteri cell and mucus binding protein A (CmbA) in adhesion to in-
testinal epithelial cells and mucus in vitro," *Microbiology (United Kingdom)*, 160(PART 4). doi:
10.1099/mic.0.073551-0.

Joeres-Nguyen-Xuan, T. H. *et al.* (2010) "Survival of the probiotic Escherichia coli Nissle 1917 (EcN) in the gastroin-
testinal tract given in combination with oral mesalamine to healthy volunteers," *Inflammatory Bowel Diseases*,
16(2). doi: 10.1002/ibd.21042.

Jonsson, H., Ström, E. and Roos, S. (2001) "Addition of mucin to the growth medium triggers mucus-binding activi-
ty in different strains of Lactobacillus reuteri in vitro," *FEMS Microbiology Letters*, 204(1). doi: 10.1016/S0378-
1097(01)00369-X.

Kankainen, M. *et al.* (2009) "Comparative genomic analysis of Lactobacillus rhamnosus GG reveals pili containing a
human-mucus binding protein," *Proc Natl Acad Sci U S A*, 106(40). doi: 10.1073/pnas.0908876106.

Karlskås, I. L. *et al.* (2014) "Heterologous protein secretion in lactobacilli with modified pSIP vectors," *PLoS One*. San Francisco: Public Library of Science, 9(3). doi: 10.1371/journal.pone.0091125.

Khalili, A. A. and Ahmad, M. R. (2015) "A review of cell adhesion studies for biomedical and biological applications," *International Journal of Molecular Sciences*, 16(8). doi: 10.3390/ijms160818149.

Koropatkin, N. M., Cameron, E. A. and Martens, E. C. (2012) "How glycan metabolism shapes the human gut microbiota," *Nature Reviews Microbiology*, 10(5). doi: 10.1038/nrmicro2746.

Krishnan, S. *et al.* (2018) "Gut Microbiota-Derived Tryptophan Metabolites Modulate Inflammatory Response in Hepatocytes and Macrophages," *Cell Reports*. ElsevierCompany., 23(4). doi: 10.1016/j.celrep.2018.03.109.

Laparra, J. M. and Sanz, Y. (2009) "Comparison of in vitro models to study bacterial adhesion to the intestinal epithelium," *Letters in Applied Microbiology*, 49(6). doi: 10.1111/j.1472-765X.2009.02729.x.

Larsson, J. M. H. *et al.* (2009) "A complex, but uniform O-glycosylation of the human MUC2 mucin from colonic biopsies analyzed by nanoLC/MSn," *Glycobiology*, 19(7). doi: 10.1093/glycob/cwp048.

Lasaro, M. A. *et al.* (2009) "F1C fimbriae play an important role in biofilm formation and intestinal colonization by the escherichia coli commensal strain nissle 1917," *Applied and Environmental Microbiology*, 75(1). doi: 10.1128/AEM.01144-08.

Latousakis, D. *et al.* (2019) "Serine-rich repeat protein adhesins from Lactobacillus reuteri display strain specific glycosylation profiles," *Glycobiology*, 29(1). doi: 10.1093/glycob/cwy100.

Le, D. T. L. *et al.* (2013) "Unraveling the role of surface mucus-binding protein and pili in muco-adhesion of Lactococcus lactis," *PLoS ONE*, 8(11). doi: 10.1371/journal.pone.0079850.

Lebeer, S. *et al.* (2009) "Identification of a gene cluster for the biosynthesis of a long, galactose-rich exopolysaccharide in Lactobacillus rhamnosus GG and functional analysis of the priming glycosyltransferase," *Applied and Environmental Microbiology*, 75(11). doi: 10.1128/AEM.02919-08.

Lee, Y. K. *et al.* (2000) "Quantitative approach in the study of adhesion of lactic acid bacteria to intestinal cells and their competition with enterobacteria," *Applied and Environmental Microbiology*, 66(9). doi: 10.1128/AEM.66.9.3692-3697.2000.

Leteurtre, E. *et al.* (2004) "Differential mucin expression in colon carcinoma HT-29 clones with variable resistance to 5-fluorouracil and methotrexate," *Biology of the Cell*, 96(2). doi: 10.1016/j.biolcel.2003.12.005.

Li, X. J. *et al.* (2008) "The adhesion of putative probiotic lactobacilli to cultured epithelial cells and porcine intestinal mucus," *Journal of Applied Microbiology*. Blackwell Publishing Ltd, 104(4). doi: 10.1111/j.1365-2672.2007.03636.x.

MacKenzie, D. A. *et al.* (2010) "Strain-specific diversity of mucus-binding proteins in the adhesion and aggregation properties of Lactobacillus reuteri," *Microbiology*. England: Society for General Microbiology, 156(11). doi: 10.1099/mic.0.043265-0.

Mackie, A. R. *et al.* (2017) "Innovative Methods and Applications in Mucoadhesion Research," *Macromolecular Bioscience*, 17(8). doi: 10.1002/mabi.201600534.

Maltby, R. *et al.* (2013) "Nutritional Basis for Colonization Resistance by Human Commensal Escherichia coli Strains HS and Nissle 1917 against E. coli O157:H7 in the Mouse Intestine," *PLoS ONE*, 8(1). doi: 10.1371/journal.pone.0053957.

Mathiesen, G. *et al.* (2009) "Genome-wide analysis of signal peptide functionality in Lactobacillus plantarum
WCFS1," *BMC Genomics*, 10. doi: 10.1186/1471-2164-10-425.

Mays, Z. J. and Nair, N. U. (2018) "Synthetic biology in probiotic lactic acid bacteria: At the frontier of living thera-
peutics," *Current Opinion in Biotechnology*. Elsevier Ltd, 53, pp. 224–231. doi: 10.1016/j.copbio.2018.01.028.

Mennigen, R. *et al.* (2009) "Probiotic mixture VSL#3 protects the epithelial barrier by maintaining tight junction
protein expression and preventing apoptosis in a murine model of colitis," *American Journal of Physiology-*
Gastrointestinal and Liver Physiology, 296(5). doi: 10.1152/ajpgi.90534.2008.

Monteiro, C. *et al.* (2009) "Characterization of cellulose production in Escherichia coli Nissle 1917 and its biological
consequences," *Environmental Microbiology*, 11(5). doi: 10.1111/j.1462-2920.2008.01840.x.

Mukherjee, S. and Ramesh, A. (2015) "Bacteriocin-producing strains of Lactobacillus plantarum inhibit adhesion of
Staphylococcus aureus to extracellular matrix: Quantitative insight and implications in antibacterial therapy," *Jour-*
nal of Medical Microbiology, 64(12). doi: 10.1099/jmm.0.000181.

Nishiyama, K., Sugiyama, M. and Mukai, T. (2016) "Adhesion Properties of Lactic Acid Bacteria on Intestinal Mucin,"
Microorganisms, 4(4). doi: 10.3390/microorganisms4030034.

Oh, P. L. *et al.* (2010) "Diversification of the gut symbiont lactobacillus reuteri as a result of host-driven evolution,"
ISME Journal. England: Nature Publishing Group, 4(3). doi: 10.1038/ismej.2009.123.

Ollech, J. E. *et al.* (2016) "Use of probiotics in prevention and treatment of patients with Clostridium difficile infec-
tion," *Best Practice and Research: Clinical Gastroenterology*. Baillière Tindall, 30(1). doi: 10.1016/j.bpg.2016.01.002.

Ou, B. *et al.* (2016) "Genetic engineering of probiotic Escherichia coli Nissle 1917 for clinical application," *Applied*
Microbiology and Biotechnology. Applied Microbiology and Biotechnology, 100(20). doi: 10.1007/s00253-016-
7829-5.

Ouwehand, A. C. (2017) "A review of dose-responses of probiotics in human studies," *Beneficial Microbes*, 8(2). doi:
10.3920/BM2016.0140.

Petrova, M. I. *et al.* (2016) "The lectin-like protein 1 in Lactobacillus rhamnosus GR-1 mediates tissue-specific ad-
herence to vaginal epithelium and inhibits urogenital pathogens," *Scientific Reports*. Nature Publishing Group,
6(October). doi: 10.1038/srep37437.

Riglar, D. T. *et al.* (2017) "Engineered bacteria can function in the mammalian gut long-term as live diagnostics of
inflammation," *Nature Biotechnology*, 35(7), pp. 653–658. doi: 10.1038/nbt.3879.

Rojas, M., Ascencio, F. and Conway, P. L. (2002) "Purification and Characterization of a Surface Protein from Lacto-
bacillus fermentum 104R That Binds to Porcine Small Intestinal Mucus and Gastric Mucin Purification and Charac-
terization of a Surface Protein from Lactobacillus fermentum 104R That Binds to ," 68(5). doi:
10.1128/AEM.68.5.2330.

Roos, S. *et al.* (1996) "A collagen binding protein from Lactobacillus reuteri is part of an ABC transporter system?,"
FEMS Microbiology Letters, 144(1). doi: 10.1016/0378-1097(96)00334-5.

Roos, S. and Jonsson, H. (2002) "A high-molecular-mass cell-surface protein from Lactobacillus reuteri 1063 ad-
heres to mucus components," *Microbiology*. England: Soc General Microbiol, 148(2). doi: 10.1099/00221287-148-2-
433.

Rosales-Mendoza, S., Angulo, C. and Meza, B. (2016) "Food-grade organisms as vaccine biofactories and oral de-
livery vehicles," *Trends in Biotechnology*. Elsevier Ltd, 34(2). doi: 10.1016/j.tibtech.2015.11.007.

Rud, I. *et al.* (2006) "A synthetic promoter library for constitutive gene expression in Lactobacillus plantarum," *Mi-*
crobiology, 152(4). doi: 10.1099/mic.0.28599-0.

Salas-Jara, M. *et al.* (2016) "Biofilm Forming Lactobacillus: New Challenges for the Development of Probiotics,"
Microorganisms, 4(4). doi: 10.3390/microorganisms4030035.

Saraiva, T. D. L. *et al.* (2015) "Milk fermented with a 15-lipoxygenase-1-producing Lactococcus lactis alleviates
symptoms of colitis in a murine model.," *Current pharmaceutical biotechnology*, 16(5). doi:
10.2174/1389201015666141113123502.

Scaldaferri, F. *et al.* (2016) "Role and mechanisms of action of Escherichia coli Nissle 1917 in the maintenance of
remission in ulcerative colitis patients: an update," *World Journal of Gastroenterology*, 22(24). doi:
10.3748/wjg.v22.i24.5505.

Shafaghi, A. *et al.* (2016) "The Effect of Probiotic Plus Prebiotic Supplementation on the Tolerance and Efficacy of
Helicobacter Pylori Eradication Quadruple Therapy: a Randomized Prospective Double Blind Controlled Trial,"
Middle East Journal of Digestive Diseases. Iranian Society of Gastroenterology and Hepatology, 8(3). doi:
10.15171/mejdd.2016.30.

Shreiner, A. B., Kao, J. Y. and Young, V. B. (2015) "The Gut Microbiome in Health and in Disease," *Curr Opin*
Gastroenterol. NIH Public Access, 31(1). doi: 10.1097/MOG.0000000000000139.The.

Tallon, R. *et al.* (2007) "Strain- and matrix-dependent adhesion of Lactobacillus plantarum is mediated by
proteinaceous bacterial compounds," *Journal of Applied Microbiology*, 102(2). doi: 10.1111/j.1365-
2672.2006.03086.x.

Troge, A. *et al.* (2012) "More than a marine propeller - the flagellum of the probiotic Escherichia coli strain Nissle
1917 is the major adhesin mediating binding to human mucus," *International Journal of Medical Microbiology.*
Elsevier GmbH., 302(7–8), pp. 304–314. doi: 10.1016/j.ijmm.2012.09.004.

Uchida, H. *et al.* (2004) "A New Assay Using Surface Plasmon Resonance (SPR) to Determine Binding of the Lacto-
bacillus acidophilus Group to Human Colonic Mucin," *Bioscience, Biotechnology, and Biochemistry*, 68(5). doi:
10.1271/bbb.68.1004.

US National Library of Medicine (2018) *Search: Probiotic; Lactic Acid Bacteria; Intestine | Exclude Unknown | Com-*
pleted | 01/012018 - 12/31/2018.

Valeriano, V. D. *et al.* (2016) "Carbohydrate-binding specificities of potential probiotic Lactobacillus strains in por-
cine jejunal (IPEC-J2) cells and porcine mucin," *Journal of Microbiology*, 54(7). doi: 10.1007/s12275-016-6168-7.

Wagner, C. E., Wheeler, K. M. and Ribbeck, K. (2018) "Mucins and Their Role in Shaping the Functions of Mucus
Barriers," *Annual Review of Cell and Developmental Biology*, 34(1). doi: 10.1146/annurev-cellbio-100617-062818.

Waller, M. C. *et al.* (2017) "Toward a genetic tool development pipeline for host-associated bacteria," *Current*
Opinion in Microbiology. Elsevier Ltd, 38, pp. 156–164. doi: 10.1016/j.mib.2017.05.006.

Walter, J., Britton, R. A. and Roos, S. (2011) "Host-microbial symbiosis in the vertebrate gastrointestinal tract and
the Lactobacillus reuteri paradigm," *Proc Natl Acad Sci U S A*, 108 Suppl. doi: 10.1073/pnas.1000099107.

West, K. A. *et al.* (2019) " An engineered E. coli Nissle improves hyperammonemia and survival in mice and shows
dose-dependent exposure in healthy humans ," *Science Translational Medicine*, 11(475). doi:
10.1126/scitranslmed.aau7975.

Wickström, C. *et al.* (2013) "Surface-associated MUC5B mucins promote protease activity in Lactobacillus
fermentum biofilms," *BMC Oral Health*, 13(1). doi: 10.1186/1472-6831-13-43.

Zanin, M. *et al.* (2016) "The Interaction between Respiratory Pathogens and Mucus," *Cell Host and Microbe*. Elsevier
Inc., 19(2). doi: 10.1016/j.chom.2016.01.001.

Zhang, Yunlei *et al.* (2012) "Escherichia coli Nissle 1917 targets and restrains mouse b16 melanoma and 4T1 breast
tumors through expression of azurin protein," *Applied and Environmental Microbiology*, 78(21), pp. 7603–7610. doi:
10.1128/AEM.01390-12.