Title:

Molecular responses to freshwater limitation in the mangrove tree *Avicennia germinans*

(Acanthaceae)

Short running title:

Black-Mangroves responses to water limitation

Authors:

Mariana Vargas Cruz[1], Gustavo Maruyama Mori[2], Dong-Ha Oh[3], Maheshi Dassanayake[3],

Maria Imaculada Zucchi[4], Rafael Silva Oliveira[1], Anete Pereira de Souza[1].

[1]Department of Plant Biology, Institute of Biology, University of Campinas (Unicamp),

Campinas, SP 13083-863, Brazil.

[2]Institute of Biosciences, São Paulo State University (Unesp), São Vicente, SP 11330-900,

Brazil.

[3]Department of Biological Sciences, Louisiana State University (LSU), Louisiana, LA 70803,

United States.

[4]São Paulo Agency for Agribusiness Technology (APTA), Piracicaba, SP 13400-790, Brazil.

*Author for correspondence. Tel: +55 19 3521 1132 Email: anete@unicamp.br

Abstract

Environmental variation along the geographical space can shape populations by natural selection. In the context of global warming, accompanied by substantial changes in precipitation regimes, it is crucial to understand the role of environmental heterogeneity in tropical trees adaptation, given their disproportional contribution to water and carbon biogeochemical cycles. Here we investigated how heterogeneity in freshwater availability along tropical wetlands has influenced molecular variations of the Black-Mangrove (*Avicennia germinans*). Fifty-seven trees were sampled in seven sites differing markedly on precipitation regime and riverine freshwater inputs. Using 2,297 genome-wide single nucleotide polymorphic markers, we found signatures of natural selection by the genotype association with the precipitation of the warmest quarter and the annual precipitation. We also found candidate loci for selection, based on statistical deviations from neutral expectations of interpopulation differentiation. Most candidate loci present within coding sequences were functionally associated with central aspects of drought-tolerance or plant response to drought. Complementarily, our results suggest the occurrence of rapid evolution of a population, likely in response to sudden and persistent limitation in plant access to soil water, following a road construction in 1974. Observations supporting rapid evolution included reduction in tree size and changes in allele frequencies and in transcripts expression levels associated with increased drought-tolerance, through accumulation of osmoprotectants and antioxidants, biosynthesis of plant cuticle, protection against stress-induced proteins degradation, stomatal closure, photorespiration and photosynthesis. We describe a major role of spatial heterogeneity in freshwater availability in the specialization of this typically tropical tree.

Keywords: *Avicennia germinans* (Black Mangrove), drought-tolerance, tropical tree, nextRAD, RNA-Seq, ecological genomics.

1.   Introduction

Natural ecosystems are characterized by wide environmental heterogeneity over the

geographical space. These spatial variations in several abiotic conditions can shape

populations specializations, particularly in widespread species, through changes in

frequencies of genotypes and phenotypes (Kawecki & Ebert, 2004). Environmental gradients

are, therefore, natural laboratories for the study of environmental selection (De Frenne et al.,

2013). As sessile organisms, plants are incapable of escaping from unfavorable conditions

and, therefore, are ideal models for investigating adaptation through phenotypic plasticity

and/or adaptive genetic variation. They are often subject to a wide range of environmental

factors, such as water, light, temperature and nutrients availability. Among the various

environmental factors that determine adaptive phenotypic and genotypic diversity in plants,

freshwater availability has a prominent role (Choat et al., 2018; Phillips et al., 2010).

Accordingly, population differentiation in drought-tolerance has been widely identified in

various studies, providing valuable insights on evolutionary consequences of spatial variation

in freshwater availability in plant species (Aranda et al., 2014; Donovan, Ludwig, Rosenthal,

Rieseberg, & Dudley, 2009; Etterson, 2004; Heschel & Riginos, 2005; Keller et al., 2011;

Ramírez-Valiente et al., 2018). Reduced freshwater availability resulting from the

combination of high atmospheric temperature with low rainfall and air humidity represent a

major threat to plants, especially trees and forest ecosystems they form (Asner et al., 2016;

Bennett, McDowell, Allen, & Anderson-Teixeira, 2015; McDowell & Allen, 2015). These

changes influence major components of resource-use: it reduces the soil water potential,

which limits the water and nutrients supply to leaves, and raises the air vapor pressure deficit

(VPD), increasing water loss through transpiration (McRae, 1980; Novick et al., 2016). In

response to these conditions, plants close their stomata (McAdam & Brodribb, 2015; Tyree &

Sperry, 1989), decreasing the chance of death from hydraulic failure (Rowland et al., 2015),

despite causing negative impacts on photosynthesis and productivity (Lawlor, 2002).

Currently, there is a great interest in understanding genotypic and phenotypic basis of

trees resistance to drought, as these mechanisms are key to improve predictions of

environmental consequences of extreme events and to elaborate plans to mitigate forest loss

(Corlett, 2016; da Costa et al., 2010; Phillips et al., 2009). Substantial advances in the

understanding of phenotypic characteristics that enhance drought-tolerance in trees have been

achieved recently (Bartlett, Scoffoni, & Sack, 2012; Bennett et al., 2015; Hacke, Sperry,

Wheeler, & Castro, 2006; Powell et al., 2017), however little is known about the molecular

basis of water-stress tolerance, particularly in tropical species (Holliday et al., 2017), which

contribute disproportionally to global carbon cycle (Corlett, 2016).

In this study, we investigated the role of environmental selection along a gradient of

freshwater availability in shaping the molecular variation of a typically tropical and abundant

tree, *Avicennia germinans* (L.) L. (Acanthaceae). The species is particularly suitable for the

study of mechanisms involved in trees adaptation to drought, since it is the most widespread

mangrove in the Atlantic East-Pacific biogeographic region (Ellison, Farnsworth, & Merkt,

1999), belonging to the most tolerant mangrove genus to limiting conditions for water

acquisition, as drought, freezing and salinity (Pranchai et al., 2017; Reef & Lovelock, 2015;

Stuart, Choat, Martin, Holbrook, & Ball, 2007). These trees occur naturally under daily and

seasonal fluctuations in temperature, air humidity, freshwater inputs and soil salinity

(Tomlinson, 1986). In tropical arid zones or during dry seasons, VPD, soil salinity and water-

deficits can reach extreme levels, hindering the maintenance of water and ion homeostasis and

limiting carbon gain and plant growth (Clough, Sim, Inlet, Bay, & Rivers, 1989; Lin &

Sternberg, 1992).

We hypothesized that spatial heterogeneity in freshwater availability shapes adaptive

variation in allele frequencies and in expression profiles of transcripts associated with the

response to and tolerance of limiting freshwater in tropical trees. We sampled *A. germinans*

individuals along a wide longitudinal range in an equatorial region (from 0°43'12" S to

8°31'48" S of latitude), showing narrow spatial heterogeneity in temperature and solar

radiation, but encompassing high variation in the intensity and duration of the dry season, in

annual precipitation levels and in riverine freshwater input, which influence levels of soil

salinity (Figure 1). We used the Nextera-tagmented reductively-amplified DNA (nextRAD)

sequencing approach for the identification and genotyping of genome-wide single nucleotide

polymorphisms (SNPs). We analyzed the organization of the genetic structure and performed

statistical tests to identify candidate loci for selection. To minimize false positives in the

detection of candidate loci (Lotterhos & Whitlock, 2015), we used distinct approaches: (1)

based on the identification of loci deviating from neutral models of interpopulational genetic

variation (F_{ST} outlier tests) and (2) based on direct genetic-environment (G-E) association

tests. RNA sequencing (RNA-Seq) was used to assemble and characterize the transcriptome

of the species, providing a functional basis for the annotation of candidate loci for selection.

Additionally, we examined the role of molecular adaptation to extreme soil freshwater

limitation via differential gene expression analysis between samples from adjacent sites

differing in tidal inundation frequency (Lara & Cohen, 2006; Pranchai et al., 2017),

acclimated in pots under homogeneous, watered conditions. Our results provide converging

signs of adaptive responses of *A. germinans* to the environmental heterogeneity in freshwater

availability, including a case suggesting the rapid evolution of a population, with changes in

phenotype and in the genetic profile. We highlight the relevance of environmental

heterogeneity in freshwater availability as a key selective pressure in tropical tree species.

2.   Materials and methods

2.1   Study system

Avicennia germinans (L.) L. is one of the most widespread mangroves in the Atlantic-East Pacific biogeographic region. It presents a generalist insect pollination system (Nadia, Menezes, & Machado, 2013), and is predominantly outcrossing, although it supports self-fertilization (Mori, Zucchi, & Souza, 2015; Nettel-Hernanz, S.Dodd, Ochoa-Zavala, Tovilla-Hernández, & Días-Gallegos, 2013). The species reproduces by producing cryptoviviparous, buoyant, and salt-tolerant propagules, allowing for long-distance sea dispersal (Mori, Zucchi, Sampaio, & Souza, 2015; Tomlinson, 1986).

2.2   Study area

Sampling sites were located over more than 1,800 km of the north-northeast Brazilian coastline, between 0.724° S and 8.526° S of latitude, along a spatial gradient in freshwater availability (Figure 1a, Table 1). We obtained environmental data for each sampling site, consisting of 21 bioclimatic and oceanographic variables from the public databases WorldClim (Hijmans, Cameron, Parra, Jones, & Jarvis, 2005) (15 precipitation, radiation and air vapor pressure layers) and Marspec (Sbrocco & Barber, 2013) (four sea surface salinity and temperature layers). To minimize the collinearity of variables, we used a correlation threshold of 0.8 (Supplemental Figure S1). We classified the study area into three distinguishable regions, based on a principal component analysis (Figure 1b): (1) the Amazon Macrotidal Mangrove Coast (AMMC), the world's largest continuous mangrove belt (Nascimento Jr., Souza-Filho, Proisy, Lucas, & Rosenqvist, 2013), which has a mean annual precipitation above 2,000 mm yr^{-1} and is influenced by the mouth of the Amazon River; (2)

mangroves of Northeast Brazil, which show limited forest development (Schaeffer-Novelli,

Cintrón-Molero, Adaime, & de Camargo, 1990) and is characterized by the lack of riverine

freshwater inputs and a mean annual precipitation below 2,000 mm yr^{-1}, with pronounced and

long dry seasons, and less than 30 mm of precipitation in the driest quarter; and (3) a region

influenced by the southward-flowing branch of the South Equatorial coastal current (South of

the SEC), characterized by reduced riverine freshwater inputs, mean annual precipitation

below 2,000 mm yr^{-1}, but less pronounced dry season, showing more than 100 mm of

precipitation in the driest quarter.

Table 1. Characterization and geographic location of *Avicennia germinans* sampling sites.

Sampling site ID	Climate classification†	Latitude; Longitude	Number of samples	
			nextRAD	RNA-Seq
MRJ	Am	00.72 S; 48.49 W	8	-
PA-arid‡	Am	00.90 S; 46.69 W	9	3
PA-humid	Am	00.94 S; 46.72 W	9	3
ALC	Aw	02.41 S; 44.41 W	5	-
PNB	Aw	02.78 S; 41.82 W	9	-
PRC	Aw	03.41 S; 39.06 W	7	-
TMD	As	08.53 S; 35.01 W	10	-

155 †Köppen-Geiger climate classification system (Alvares, Stape, Sentelhas, Gonçalves, &
156 Sparovek, 2013). *Am*: Tropical monsoon climate; *Aw*: Tropical wet savanna climate; *As*:
157 Tropical dry savanna climate.
158 ‡Samples from a forest flooded exclusively during spring tides (Lara & Cohen, 2006).
159

Figure 1. *Avicennia germinans* **sampling sites along a low-latitude salinity and precipitation gradient.** (a) *Top:* Geographical location of the study area and sampling sites. Black points represent sampling sites, green area represents the occurrence of the species, and blue areas represent ponds and rivers. *Bottom:* Environmental variation based on sea surface salinity and precipitation variables across sampling sites (source: WorldClim and Marspec). *AMMC*: Amazon Macrotidal Mangrove Coast. *SEC*: South Equatorial Current. (b) Bidimensional projection of sampling sites, based on a principal component analysis. PC1 and PC2 retained 79.7% of the variance of non-collinear environmental variables (eigenvectors). *AnnP*: Annual precipitation. *MinVap*: Minimum air water vapor pressure. *MinTCM*: Minimum temperature of the coldest month. *SSTCM*: Sea surface temperature of the coldest month. *MeanSST*: Mean sea surface temperature. *MeanSSS*: Mean sea surface salinity. *PWarmQ*: Precipitation of the warmest quarter. *MaxRad*: Maximum solar radiation. *SSTRange*: Sea surface temperature range. *PDM*: Precipitation of the driest month.

In the AMMC, we visited two adjacent sites, which require a more detailed

description, both located in the peninsula of Ajuruteua, state of Pará, between the Maiaú and

Caeté estuaries (Figure 2a). The peninsula was covered by a preserved mangrove forest

(Cohen & Lara, 2003), until it was divided by the construction of a road (Figure 2b), in 1974.

The hydrology part of the forest was dramatically changed, no longer being influenced by the

Caeté River. Instead, it started flooding exclusively during the highest spring tides of the

Maiaú River. These changes resulted in the forest dieback and subsequent recolonization of

the impacted area, mainly by *A. germinans*. Pore water salinity accumulated to extremely high

levels (100 ppt at 50-cm depth), and the air surface temperature frequently exceeds 40 °C

(Vogt et al., 2014). These environmental features contributed to the dwarfism of recolonizing

individuals (Cohen & Lara, 2003; Pranchai et al., 2017), whose shrub architecture (up to 2.0

m in height) contrasts to the former tall morphology (up to 30 m in height), still observed on

surrounding areas, where the hydrology remained unaltered (Menezes, Berger, & Mehlig,

2008) (Figure 2b). Throughout this work, we refer to the western side of the road as 'PA-arid'

and to the eastern side as 'PA-humid'.

190
Figure 2. PA-arid and PA-humid sampling sites and RNA sequencing (RNA-Seq)
experimental design. (a) Geographical location of sites, highlighted by a square (red colored
area is the PA-arid site). (b) Photograph of a section of the Bragança-Ajuruteua road (Pará,
Brazil) along which severe changes in hydrology altered the mangrove community and tree
morphology. (c) RNA-Seq experimental design. *Photographs authors:* G.M. Mori and M.V.
Cruz.
197

2.3 DNA extraction and sequencing

Leaves from 57 adult *A. germinans* trees were sampled in seven distinct sites (Table 1)

and stored in bags with silica gel. DNA extraction was performed using the DNeasy Plant

Mini Kit (QIAGEN) and NucleoSpin Plant II (Macherey Nagel). DNA quality and quantity

were assessed using 1% agarose gel electrophoresis and QuantiFluor dsDNA System in a

Quantus fluorometer (Promega). NextRAD libraries were constructed by SNPsaurus

(SNPsaurus, LLC) (Russello, Waterhouse, Etter, & Johnson, 2015). Genomic DNA

fragmentation and short-adapter ligation were performed with Nextera reagent (Illumina,

Inc.), followed by amplification, in which one of the primers matched the adapter and

extended by nine arbitrary nucleotides of DNA. Thus, amplicons were fixed at the selective

end, and their lengths depended on the initial fragmentation, leading to consistent genotyping

of amplified loci. Subsequently, nextRAD libraries were sequenced in a HiSeq 2500

(Illumina, Inc), with 100-bp single-end chemistry (Supplemental Figure S2).

2.4   SNP detection and genotyping

Genotyping-by-sequencing used custom scripts (SNPsaurus, LLC) to create a

reference catalog of abundant reads. Read mapping to the catalog allowed two mismatches.

Biallelic loci present in at least 10% of samples were selected for the following steps. Using

VCFtools 0.1.12b (Danecek et al., 2011), we selected high-quality sequences (Phred score >

30), allowing a maximum of 65% missing data and one SNP per sequence, requiring a

minimum coverage of 8x and a minor allele frequency ≥ 0.05 (Supplemental Figure S2). To

reduce false SNP detection rates due to paralogy or low-quality genotype calls, we used a

maximum read of 56, resulting from the product of the average read depth and 1.5 standard

deviation of the mean.

2.5   Genetic diversity analysis and identification of candidate SNP loci associated with

water-stress tolerance

For each sampling site, we estimated the genetic diversity using per-site nucleotide

diversity (π), calculated using VCFtools (Danecek et al., 2011); private alleles (pA), observed

(H_O) and expected (H_E) heterozygosities, using poppr 2.8.2 (Kamvar, Tabima, & Grünwald,

2014); and the percentage of polymorphic loci (%Poly), using adegenet 2.1.1 (Jombart &

Ahmed, 2011). We also estimated pairwise genetic differentiation (F_{ST}), the inbreeding

coefficient (F_{IS}) and its 95% confidence interval, through 1000 bootstrap resampling over all

SNP loci, using hierfstat 0.04-22 (Goudet, 2005). The genetic structure of *A. germinans* was

described by a multivariate model-free method, the discriminant analysis of principal

components (DAPC) (Jombart, Devillard, & Balloux, 2010), and ADMIXTURE 1.3.0

(Alexander, Novembre, & Lange, 2009). We considered numbers of groups from 1 to 50 and

the Bayesian information criteria to determine the number of groups (K). The optim.a.score

function was used to avoid overfitting during discrimination steps. For ADMIXTURE

analysis, we performed three separate runs for values of K varying from 1 to 15, using the

block-relaxation method for point estimation. Computing was terminated when estimates

increased by less than 0.0001. The lowest level of cross-validation error indicated the most

likely K-value.

To detect signatures of natural selection, we used three distinct methods. Loci that

were highly correlated with the environmental structure (false discovery rate (FDR) <0.05)

were detected using the R package LEA (Frichot & François, 2015), which is based on

analyses of genetic structure and ecological association (G-E association tests). In these

analyses, we used non-collinear environmental bioclimatic and oceanographic variables

(Supplemental Figure S1) retrieved from the public databases WorldClim (Hijmans et al.,

2005) (precipitation, radiation and air vapor pressure layers) and Marspec (Sbrocco & Barber,

2013) (sea surface salinity and temperature layers). Since the environmental divergence

between PA-arid and PA-humid is so evident (Figure 2), yet undistinguishable based on

highest-resolution environmental datasets, we excluded PA-arid samples from G-E

association tests.

To overcome limitations in the acquisition of environmental data for the PA-arid site,

two other methods were used to detect candidate loci for selection. These methods are solely

based on deviations from neutral expectations of allele frequency distributions, regardless of

environmental variation data across sampling sites (F_{ST} outlier tests). With Pcadapt 2.0 (Luu,

Bazin, & Blum, 2016), population structure was determined by a principal component

analysis. A disproportional relation to this structure (FDR < 0.05) was indicative of selection.

With Lositan (Antao, Lopes, Lopes, Beja-Pereira, & Luikart, 2008), which uses the FDIST2

method (Beaumont & Nichols, 1996), we assessed the relationship between F_{ST} and H_E to

describe the neutral distribution under an island migration model. Hence, we detected loci

with excessively high or low F_{ST}. Because Lositan may show partially divergent results

among independent simulations, we only considered candidate loci conservatively identified

in three independent Lositan simulations assuming an infinite allele model of mutation, with a

confidence interval of 0.99 and a FDR < 0.05, using the neutral mean F_{ST} and forcing the

mean F_{ST} options.

Demographic histories can affect the statistical power of tests of selection (Lotterhos

& Whitlock, 2015); thus, type I and type II errors are frequently associated with these

approaches (Narum & Hess, 2011). We minimized the potential for false discovery by

identifying consensus candidate loci for selection among distinct methods used for two

datasets: (1) the entire genotypic dataset, including PA-arid individuals, using Lositan and

Pcadapt consensus candidates and (2) a subset of genotypic data, which excluded the PA-arid

samples due to absence of environmental data, using Lositan, Pcadapt and LEA consensus

candidates (Supplemental Figure S2).

2.6   Functional annotation of candidate loci putatively under natural selection

Functional annotation of candidate loci was obtained by reciprocal nucleotide

alignment between nextRAD sequences and the reference transcriptome characterized in the

present study. Blast+ 2.2.31 (Camacho et al., 2009) was used with a threshold of at least 50

aligned nucleotides, a maximum of one mismatch and no gaps (Supplemental Figure S2).

2.7   Plant material for transcriptome assembly and differential expression analysis

On August 16th, 2013, three similarly sized seedlings of *A. germinans* with 5-9 stem

nodes were collected from the PA-arid and PA-humid sites, separated by the Bragança-

Ajuruteua road, and transplanted with the surrounding soil into 3.0-L pots (Figure 2c). Based

on previous observations of other *Avicennia* species (Almahasheer, Duarte, & Irigoien, 2016;

Duarte, Thampanya, Terrados, Geertz-Hansen, & Fortes, 1999), we estimate that all samples

were between at least 3.5 months to at most 11 months old. For acclimation in a

homogeneous environment, pots were placed in open-air and naturally shaded conditions and

were watered daily after the sunset, with 300 mL of tap water. Plants were harvested at noon,

after 72 hours of acclimation, washed with water and split into roots, stems and leaves with a

sterile blade to be stored in RNAlater (Ambion Inc., Austin, TX, USA) and transported to the

laboratory for RNA extraction.

2.8   RNA extraction, cDNA library preparation and RNA sequencing

RNA extraction was performed according to Oliveira, Viana, Reátegui, & Vincentz

(2015). To assess purity, integrity and concentration, we used 1% agarose gel electrophoresis

and a NanoVue spectrophotometer (GE Healthcare Life Sciences, Buckinghamshire, UK).

Subsequently, cDNA-enriched libraries were constructed using TruSeq RNA Sample

Preparation kits (Illumina Inc., California, USA). Libraries qualities were assessed using an

Agilent 2100 Bioanalyzer (Agilent Technologies, California, USA) and concentrations were

quantified using quantitative real-time PCR (qPCR), with a Sequencing Library qPCR

Quantification kit (Illumina Inc.). Sequencing was performed with two 36-cycle TruSeq SBS

paired-end kits (Illumina Inc.) on a Genome Analyzer IIx platform (Illumina Inc.).

2.9   *Transcriptome assembly and functional annotation of transcripts*

Raw data were filtered by quality, using Phred >20 for 70% of the read length, and

adapters were trimmed using NGS QC Toolkit 2.3 (Patel & Jain, 2012). Filtered reads were

de   novo   assembled   into   transcripts   using   the   CLC   Genomics   Workbench

(https://www.qiagenbioinformatics.com/). The distance between paired-end reads was set to

300-500 bp, the k-mer size was set to 45 bp, and the remaining default parameters were not

changed.

Reads were mapped to the transcriptome using Bowtie1 (Langmead, Trapnell, Pop, &

Salzberg, 2009), and contiguous sequences (contigs) without read-mapping support were

removed from the assembly. For transcript annotation, we used blast+ 2.2.31 (Camacho et al.,

2009), with an e-value <1e-10, using reference sequences from manually curated databases,

as the National Center for Biotechnology Information (NCBI) RefSeq protein and RefSeq

RNA (O'Leary et al., 2016) and representative proteins and cDNA from The Arabidopsis

Information Resource (TAIR) (Berardini et al., 2015). We removed putative contaminant

contigs from the assembly, which did not match plant sequences but showed high similarity to

nonplant sequences from the NCBI's RefSeq database. Sequences were assigned to the Kyoto

Encyclopedia of Genes and Genomes (KEGG) orthology (KO) identifiers using the KEGG

Automatic Annotation Server (KAAS) (Moriya, Itoh, Okuda, Yoshizawa, & Kanehisa, 2007).

Protein family domains were searched using the Pfam database and the HMMER3 alignment

tool (Finn et al., 2014).

We identified putative open reading frames (ORFs) using the program TransDecoder

(http://transdecoder.sf.net) with default parameters. Redundant transcripts were detected using

Cd-hit-est 4.6.1 (Li & Godzik, 2006) in the local alignment mode, with 95% identity and 70%

coverage of the shortest sequence thresholds. To minimize redundancy, in the final assembly,

we retained only sequences with the longest putative ORF and the longest putative non-

coding transcripts from each Cd-hit-est clusters. Functional categories were assigned to

putative coding sequences using the *Arabidopsis thaliana* association file from the Gene

Ontology Consortium website (Blake et al., 2015) (Supplemental Figure S2).

2.10   Analysis of differentially expressed transcripts (DET)

Counts of reads mapped to assembled transcripts per sequenced sample were used as

input files in DET analyses. Reads that mapped to multiple transcripts were excluded. The

count matrix was normalized and used to detect transcripts with significant differential

expression between PA-arid and PA-humid samples from the Bragança-Ajuruteua road

(Figure 2) with the EdgeR Bioconductor package (Robinson, McCarthy, & Smyth, 2010) at a

FDR <0.05. Gene Ontology (GO) term enrichment analyses were performed using the goseq

R Bioconductor package (Young, Wakefield, Smyth, & Oshlack, 2010), which takes the

length bias into account, with the default Wallenius approximation method and a p-value

cutoff set to <0.05 (Supplemental Figure S2).

3.   Results

3.1   Population genetics analyses of **A. germinans** *along the equatorial coast of Brazil*

Genetic diversity indices are shown in Table 2. The lowest levels of genetic variation

were observed in the TMD site. The remaining sites, influenced by the northern branch of the

SEC, presented considerably higher levels of diversity, including the PA-arid site, which was

one of the most genetically diverse across the study region. All sampling sites deviated from

Hardy-Weinberg equilibrium (HWE), with an excess of heterozygosity in all sites, besides

ALC, which showed a small heterozygosity deficit (F_{IS}=0.03). HWE deviation was highest in

TMD (F_{IS}=-0.43) but relatively low in all remaining sites, ranging from -0.15 (PNB) to +0.03

(ALC).

Table 2. Genetic diversity statistics, based on analyses of 2,297 genome-wide polymorphic
loci detected in 57 individuals sampled in seven sampling sites along the equatorial Atlantic
coastline of South America. π: Nucleotide diversity; H_E: expected heterozygosity; H_O:
observed heterozygosity; pA: private alleles; %Poly: percentage of polymorphic loci; F_{IS}:
inbreeding coefficient.

Sampling site	π (mean)	π (s.e.)	H_E	H_O	pA	%Poly	F_{IS}	Low Level (95% c.i.)	High Level (95% c.i.)
MRJ	0.264	0.206	0.244	0.292	1	72.70	-0.136	-0.154	-0.102
PA-arid	0.317	0.189	0.295	0.326	1	83.76	-0.050	-0.067	-0.022
PA-humid	0.301	0.186	0.280	0.328	0	84.46	-0.113	-0.132	-0.084
ALC	0.282	0.214	0.249	0.269	0	69.83	0.030	0.009	0.070
PNB	0.208	0.210	0.195	0.234	0	58.21	-0.154	-0.180	-0.111
PRC	0.323	0.193	0.296	0.327	1	82.15	-0.035	-0.053	-0.002
TMD	0.123	0.192	0.130	0.168	46	34.52	-0.426	-0.452	-0.376

A substantial genetic structure was observed over the Equatorial Brazilian coast

(Figure 3). The genetic diversity, based on 2,297 genome-wide SNPs genotyped in 57

individuals, was organized into four distinct genetic clusters (K=4) (Figure 3a, Supplemental

Figure S3a-b). The greatest genetic divergences (F_{ST}>0.46; Nei's distance>0.255) were

observed between individuals from TMD and those from all other sites (Figure 3b-c,

Supplemental Table S1). The TMD site also presented the highest number of private alleles

(pA=46). Most remarkable was the divergence between individuals from PA-arid and its

adjacent site, PA-humid (F_{ST}=0.25; Nei's distance=0.181). This divergence was even greater

than the observed divergence of PA-arid from PRC individuals (Supplemental Table S1),

located approximately 900 km distant from each other (F_{ST}=0.23; Nei's distance=0.177), on

the semi-arid coast of Brazil. The divergent gene pool of the PA-arid population was also

evident when these samples were excluded from structure analyses. The most likely number

of ancestral populations dropped from four to three (Figure 3c-d, Supplemental Figure S3c-d),

and the overall structure remained unchanged (Figure 3d). At a finer scale, individuals from

PA-humid and MRJ sites, on the Amazon Macrotidal Mangrove Coast (AMMC), seemed to

be derived from the same ancestral population. This population, in turn, diverged from the

population that may have given rise to individuals from ALC, PNB, and PRC sites, in

Northeast Brazil (Figure 3a-b).

Figure 3. Genetic structure inferred from genome-wide single nucleotide polymorphisms (SNPs) detected in *Avicennia germinans*. (a) Attribution of ancestry implemented in the program Admixture 1.3.0 for all sampled individuals; stacked bars represent individuals, and each color represents one ancestral cluster (K=4). (b) Scatterplot of the first two principal components of the multivariate discriminant analysis of principal components (DAPC) of total genetic variance; all sampled individuals are represented as points; distinct symbols indicate sampling sites. (c) Attribution of ancestry (using Admixture 1.3.0) for all sampled individuals, excluding PA-arid samples; stacked bars represent individuals, and each color represents one ancestral cluster (K=3). (d) Scatterplot of the first two principal components of the DAPC of total genetic variance; sampled individuals, except for individuals sampled in the PA-arid site, are represented as points; distinct symbols indicate sampling sites.

3.2 **De novo** *assembly and annotation of the* **Avicennia germinans** *reference*

transcriptome

We used RNA-Seq to sequence and *de novo* assemble a reference transcriptome from

leaves, stems and roots of *A. germinans* seedlings, providing a functional context for this

study. A total of 249,875,572 high-quality-filtered, paired-end, 72-bp reads, representing

78.25% of the raw data was used in the assembly. The reference transcriptome comprised

47,821 contigs, after removal of misassembled, redundant and contaminant sequences.

Putative ORFs were identified in 29,854 contigs, subsequently annotated as putative protein-

coding transcripts. The remaining 17,967 contigs were classified as putative non-coding

transcripts. A detailed characterization of contigs can be found in Supplemental Table S2.

Raw reads were mapped back to the reference transcriptome, with over 82% uniquely

mapped to a single transcript and only 1.31% mapped to more than one transcript

(Supplemental Table S3). We found 91.74% of the plant universal orthologs from the

BUSCO database (Simão, Waterhouse, Ioannidis, Kriventseva, & Zdobnov, 2015)

represented in the reference transcriptome (Supplemental Table S3). We also found from

20,529 to 31,348 putative orthologous sequences between the *A. germinans* transcriptome and

four other publicly available transcriptomes derived from the genus *Avicennia* L.

(Acanthaceae) (Supplemental Table S4).

As expected, most putative protein-coding transcripts of the reference transcriptome

could be annotated using relevant databases (92.6%), whereas only a few putative non-coding

transcripts could be annotated (33.6%) (Supplemental Figure S4). We found 2,207 putative

coding transcripts (7.4%) and 11,925 putative non-coding transcripts (66.4%) unique to *A.*

germinans, which may represent lineage specific sequences.

3.3   Detection of candidate loci responding to environmental selection

Genome-wide signatures of selection were detected from genotypic data retrieved

from samples from all seven sites of collection. Fifty-six putative outlier loci were

consistently identified by two F_{ST} outlier methods, solely based on deviations from neutral

expectations of the distribution of genetic diversity. Eleven of these loci aligned to sequences

in the reference transcriptome, of which eight were highly similar to proteins associated with

the response or tolerance to drought in *A. thaliana* or *Sesamum indicum* (Figure 4a,

Supplemental Figure S5).

425
Figure 4. Candidate loci for selection detected from *Avicennia germinans* sampled along
tropical mangrove forests of the north-northeastern Brazilian coast. (a) Schematic
representation of key biological processes associated with candidate loci. *Arabidopsis*
thaliana gene symbols listed below biological processes represent the functional annotation of
candidate loci present within putative protein-coding regions of the genome. *§ Labeled gene*
symbols: candidates obtained using F_{ST} outlier tests performed with all sampled individuals,
including PA-arid samples. *¶ Labeled gene symbols*: candidates obtained by combining
genetic-environmental association tests and F_{ST} outlier approaches, using a subset of samples,
without PA-arid ones, for which environmental characterization could not be retrieved from
public databases. (b) Number of candidate SNPs associated with non-collinear environmental
variables.

The exclusion of PA-arid samples was necessary for G-E association tests, due to

limitations in the resolution of Marspec and WorldClim environmental layers. From the

remaining subset of 48 individuals from six sampling sites, further 153 candidate loci for

selection were identified by G-E correlation and two F_{ST} outlier approaches (Figure 4b). Out

of these candidate loci, 24 aligned to the reference transcriptome, of which 20 were putative

protein-coding showing high similarity to gene models from *A. thaliana* or *S. indicum*

(Supplemental Figure S5).

Among all candidate loci for selection detected along the sampling region, we found

14 loci associated with plant growth and development, wood formation, cell wall metabolism,

biogenesis of the photosynthetic apparatus, abiotic stress perception and response and protein

protection from stress-induced aggregation, among other processes (Figure 4a, Table 3).

Table 3. Annotation of candidate SNP loci putatively under selection effect identified by
two (Lositan and Pcadapt) or three (Lositan, Pcadapt and LEA) distinct methods, from
two datasets. (1) All sampled individuals, including PA-arid samples, using F_{ST} outlier
approaches and (2) a subset of samples, excluding individuals from the PA-arid site,
combining genetic-environmental association tests and F_{ST} outlier tests.

Dataset	Transcript ID	Similar to†	Environmental association‡	Putative function
1	Ag_23357	AT4G28500, NAC073, NAC domain containing protein 73	NA	Regulation of the biosynthesis of cellulose and hemicellulose in wood fibers and the expression of lignin-polymerizing and signaling genes (Hussey et al., 2011; Zhong, Lee, Zhou, McCarthy, & Ye, 2008)
1	Ag_47094	AT4G17220, MAP70-5, microtubule-associated proteins 70-5	NA	Fundamental for the development of secondary cell wall band patterning in xylem tracheids and for wood formation, playing a role in the anisotropic expansion of cells (Pesquet, Korolev, Calder, & Lloyd, 2010)
1	Ag_26495	AT5G67200, Leucine-rich repeat protein kinase (LRR-RLK) family protein	NA	May play a role in hormones and abiotic stress sensing and signaling (ten Hove et al., 2011; Van der Does et al., 2017) and in regulating adaptation to salt stress (Lorenzo et al., 2009)
1	Ag_43440	Not annotated	NA	Not annotated
1 and 2	Ag_4919	AT1G09910, Rhamnogalacturonate lyase family protein	PWQ (8.31e-05)	Associated with the degradation of the cell wall polysaccharides, and pectin (McDonough, Kadirvelraj, Harris, Poulsen, & Larsen, 2004)

1 and 2	Ag_20543	AT5G18950, Tetratricopeptide repeat (TPR)-like superfamily protein	PWQ (0.00018)	Associated with photosynthetic machinery biogenesis, stabilization and repair (Bohne, Schwenkert, Grimm, & Nickelsen, 2016)
1 and 2	Ag_5627	PREDICTED: GTP-binding protein OBGC, chloroplastic [Sesamum indicum]	PWQ (8.31e-05)	Associated with thylakoid membrane biogenesis and chloroplast protein synthesis and essential for early embryogenesis in response to light stimulus (Bang et al., 2009)
1 and 2	Ag_29619	AT2G18500, OFP7, ovate family protein 7	PWQ (0.00019)	May be involved in various aspects of plant growth and participate in suppressing cell elongation (Wang, Chang, & Ellis, 2016; Wang, Chang, Guo, & Chen, 2007)
1 and 2	Ag_38780	AT2G24395, chaperone protein dnaJ-related	PWQ (8.31e-05)	Regulation of the activity of Hsp70 chaperones (P. Walsh, Bursać, Law, Cyr, & Lithgow, 2004) and protein protection against stress-induced denaturation (Frydman, 2001)
2	Ag_12760	AT3G54810, GATA8, Plant-specific GATA-type zinc finger transcription factor family protein	PWQ (0.00363); PDM (0.01916)	Associated with the control of plant growth and development (Manfield, Devlin, Jen, Westhead, & Gilmartin, 2006)
2	Ag_135	AT1G28440, HSL1, HAESA-like 1	PWQ (9.09e-05)	The Ag_135 transcript has a kinase domain and is associated with growth and abiotic stress signaling (ten Hove et al., 2011)
2	Ag_15249	AT2G34790, MEE23, FAD-binding Berberine family protein	PWQ (8.31e-05)	Involved in the lignification of the cell wall; mediates the oxidation of monolignols (Daniel et al., 2015) and is required for endosperm development (Pagnussat et al., 2005)
2	Ag_10980	AT5G13530, KEG, keep on going	PWQ (0.00041); PDM (0.02034)	Associated with the control of plant growth and development and critical for seedling regulation of abscisic acid perception and signaling (Stone, Williams, Farmer, Vierstra, & Callis, 2006)
2	Ag_26982	AT3G01640, GLCAK, glucuronokinase G	PWQ (0.00011)	Involved in the biosynthesis of UDP-glucuronic acid, providing nucleotide sugars for the polymerization of cell-wall compounds (Pieslinger, Hoepflinger, & Tenhaken, 2010)
2	Ag_11269	AT2G25730, unknown protein	PWQ (0.00016)	The Ag_11269 transcript has a TPR 2 domain and, thus, may be involved in thylakoid membrane biogenesis and with the photosynthetic machinery stabilization and repair (Bohne, Schwenkert, Grimm, & Nickelsen, 2016)

2	Ag_43282	AT1G04300, MUSE13, TRAF-like superfamily protein	PWQ (8.31e-05)	MUSE13 regulates the turnover of nucleotide-binding domain and leucine-rich repeat-containing immune receptors. Loss of MUSE13 and MUSE14 leads to enhanced pathogen resistance, NLR accumulation, and autoimmunity. May regulate ATG6 ubiquitination and formation of autophagosomes (Huang et al., 2016)
2	Ag_14879	AT2G42520, P-loop containing nucleoside triphosphate hydrolases superfamily protein	PWQ (0.00109)	AT2G42520 has important roles in RNA metabolism such as splicing, RNA transport, ribosome biogenesis, translation and decay and unwinds double-stranded RNA molecules through the hydrolysis of NTP
2	Ag_20289	AT1G48650, DEA(D/H)-box RNA helicase family protein	PDM (0.01222); PANNUAL (5.80E-05); SSS (0.01461); WVPMAX (0.03580); RADMIN (0.03813)	AT1G48650 has important roles in RNA metabolism such as splicing, transport, ribosome biogenesis, translation and decay; it unwinds double-stranded RNA molecules through the hydrolysis of NTP
2	Ag_24006	AT1G11050, Protein kinase superfamily protein	PWQ (0.00341); PDM (0.00671); PANNUAL (0.01962)	Ag_24006 has a protein kinase domain
2	Ag_22233	AT5G01020 Protein kinase superfamily protein	PWQ (0.00022)	Ag_22233 has a protein kinase domain
2	Ag_43463	AT5G43960, Nuclear transport factor 2 (NTF2) family protein with RNA binding (RRM-RBD-RNP motifs) domain	PWQ (0.00011)	mRNA-binding protein, controls the fate and expression of a transcript
2	Ag_36772	AT2G35680, Phosphotyrosine protein phosphatases superfamily protein	PWQ (0.00026)	AT2G35680 may possess phosphatase activity
2	Ag_13254	AT4G30100, P-loop containing nucleoside triphosphate hydrolases superfamily protein	PWQ (8.31e-05)	AT4G30100 has a hydrolase activity

2	Ag_29952	AT5G11970, Protein of unknown function (DUF3511)	PWQ (8.62e-05)	Unknown function
2	Ag_1345	AT5G55960, unknown protein	PWQ (0.00018)	Unknown function
2	Ag_43837	Not annotated	PWQ (8.62e-05)	Not annotated

†Blastx hit to *Arabidopsis thaliana* or *Sesamum indicum* [when indicated] gene models.
‡(*FDR*): False discovery rate values (Benjamini-Hochberg procedure) for genetic-
environment association tests; *NA*: genetic-environment association tests were not available
for dataset (1) due to unavailability of environmental data for the PA-arid sites in public
databases; *PWQ*: Precipitation of the warmest quarter; *PDM*: Precipitation of the driest
month; *PANNUAL*: Annual precipitation; SSS: Mean sea surface salinity; *WVPMAX*:
Maximum water vapor pressure; *RADMIN*: Minimum radiation.
462

3.4   *Differential transcript expression analysis*

Transcriptome sequencing of seedlings grown under contrasting field conditions,

revealed significant expression differences in 2,454 transcripts, despite previous

acclimatization under homogeneous shaded, well-watered conditions (Figure 2c). Most DETs

were detected in roots (2,337) and stems (1,383), followed by leaves (361) (Figure 5,

Supplemental Figure S6). We refer to DETs that showed higher expression levels in samples

from the PA-arid site than in samples from the PA-humid site as "DET-Arid" and to DET

showing a significantly higher expression in samples from the PA-humid than in samples

from the PA-arid site as "DET-Humid".

472
**Figure 5. Functional categories of differentially expressed transcripts (DETs) identified
between samples grown in the PA-arid and PA-humid sampling sites.** Categories were
selected based on biological processes previously identified to be involved in the response,
acclimation or resistance to water stress in model plants.

The functional annotation and subsequent assignment of most putative protein-coding

transcripts to GO terms (76.24%) (Supplemental Figure S4) enabled the assessment of DETs

that highlighted key aspects of a differential response of *A. germinans* to contrasting source

environments, differing markedly in hydrological regime, soil pore water salinity, solar

radiation and surface temperature (Lara & Cohen, 2006; Pranchai et al., 2017; Vogt et al.,

2014). We focused this analysis on enriched biological processes previously identified to be

involved in the tolerance, resistance or response to osmotic and drought stress in various

crops, model and non-model species, including mangroves (Figure 5). A detailed

characterization of the differentially transcripts belonging to such processes is shown in

Supplemental Table S5. In the following subsections we summarize these results.

3.4.1. Photosynthesis

Mangroves have aerial, photosynthesizing roots, which contribute to carbon gain and

enable root respiration in anaerobic soils, using both atmospheric and photosynthetically

regenerated oxygen (Kitaya et al., 2002). Therefore, although various photosynthesis genes

frequently express only in basal levels in roots of the model plant, *Arabidopsis thaliana*

(Klepikova, Kasianov, Gerasimov, Logacheva, & Penin, 2016), they may express in higher

levels in mangroves, even during the seedling stage, when aerial roots are generally absent, as

observed in *A. germinans*, and in the congeneric *A. officinalis* (Krishnamurthy et al., 2017).

Differential transcripts expression analyses revealed that photosynthesis-associated transcripts

were enriched, mainly in roots and stems from the DET-Arid set. These included putative

enzymes and genes required for chloroplast biogenesis or development, for chlorophyll

biosynthesis, for the assembly of the photosynthetic apparatus and light-harvesting complexes

and involved in acclimation to fluctuating light. The DET-Arid set of roots samples also

presented transcripts associated with the C3 carbon fixation pathway, including,

glyceraldehyde-3-phosphate dehydrogenase subunits, calvin cycle proteins, and the

RIBULOSE BISPHOSPHATE CARBOXYLASE SMALL CHAIN 1A (RBCS1A), which

showed a 7.8 Fold-Change relative to roots of PA-humid samples.

3.4.2. Response to light

The DET-Arid sets of roots and stems were enriched in transcripts associated with the

response to light but not directly associated with photosynthesis. These included transcripts

similar to genes and transcription factors involved with light acclimation and essential for

maintaining electron flow and photosynthetic efficiency under changing light conditions

(Privat, Hakimi, Buhot, Favory, & Lerbs-Mache, 2003). The DET-Arid sets of roots and

stems also included putative light-signaling genes and photoreceptors, as phototropins,

cryptochromes and phytochromes. These photoreceptors are sensitive to light intensity and

control complex light and stress responses, including photoinduced movements as well as

growth and development under limiting light (Correll et al., 2003; Ohgishi, Saji, Okada, &

Sakai, 2004; Pedmale et al., 2016). Complementarily, we identified transcripts similar to

proteins that interact with these photoreceptors in the mediation of shade avoidance and

phototropism under low light, and to low light-induced transcription factors, which regulate

developmental processes and growth in response to shade avoidance. Moreover, in the DET-

Arid sets of roots and stems, we found transcripts associated with chloroplast accumulation

upon low blue light and with the response to sugar starvation induced by dark.

3.4.3.   Response to water deprivation and response to salt

525   The DET-Arid sets of roots and stems were also enriched in transcripts associated

with the response to osmotic stress and water deprivation, including putative genes that play

relevant roles in drought and salt stress resistance. For instance, we found transcripts

associated with abscisic acid (ABA)-dependent stomatal closure, a well-known mechanism

for maintaining water status under drought, which severely compromises growth (Murata,

Mori, & Munemasa, 2015). Additionally, DET-Arid sets of roots and stems included

transcripts associated with drought-induced ion transporters, which protect chloroplasts from

deleterious Na^+ concentrations (Müller et al., 2014) and involved in Ca^{2+} homeostasis under

abiotic stress. Complementarily, we found in the DET-Arid sets of roots and stems transcripts

associated with biosynthesis and accumulation of oligosaccharides, which may act as

osmoprotectants, increasing tolerance to osmotic stress (ElSayed, Rafudeen, & Golldack,

2014; Nishizawa, Yabuta, & Shigeoka, 2008; Taji et al., 2002). Additionally, transcripts

associated with drought-induced epicuticular wax biosynthesis, transport and deposition in

cell walls were among DET-Arid sets of roots and stems. We also detected in these sets of

DETs several transcripts associated with reactive oxygen species (ROS) dissipation for the

control of cell damage caused by drought, high light or salt stress, including copper/zinc

superoxide dismutases, a chloroplastic drought-induced stress protein (CDSP32) and MAP

KINASE 6 (MAPK6).

3.4.4. Response to heat

Transcripts similar to genes that confer tolerance to heat were enriched among the

DET-Arid sets of roots and stems. We detected several putative co-chaperones and

chaperones, which protect proteins from structural degradation (Al-Whaibi, 2011), and

transcripts associated with putative components of thermomemory, required for long-term

maintenance of acquired thermotolerance (Charng et al., 2006).

3.4.5. Photorespiration

As a response to drought, plants close their stomata, reducing the CO_2:O_2 ratio in

mesophyll cells, increasing levels of photorespiration. Rates of photorespiration may also be

increased by high temperatures, which reduce both the solubility of CO_2 in the mesophyll

cells and the ability of the ribulose-1,5-bisphosphate carboxylase/oxygenase (RuBisCO) to

discriminate between O_2 and CO_2 (Kozaki & Takeba, 1996; Wingler, Lea, Quick, & Leegood,

2000). On the one hand, photorespiration decreases photosynthesis efficiency, on the other, it

plays an essential role in protecting the photosynthetic machinery from damage caused by

excessive photochemical energy (Kozaki & Takeba, 1996; Wingler et al., 2000). In the DET-

Arid set of roots, we detected enrichment of putative genes associated with photorespiration,

similar to enzymes participating in transamination, required for maintaining photosynthesis

under photorespiratory conditions and for carbon flow during photorespiration, and which

catalyze the concluding reaction of photorespiration.

4.   Discussion

Our study suggests that spatial variation in freshwater availability play an important

role in driving adaptation of the typically tropical and widespread tree, *Avicennia germinans*.

In the following subsections, we describe how variation in oceanographic and bioclimatic

variables along the equatorial Brazilian coast seem to influence the organization of genome-

wide genetic diversity of the species. Moreover, we show that drastic and persistent restriction

in soil freshwater availability likely caused the rapid evolution of phenotype, gene pool and

gene expression profile of a recently founded population of *A. germinans*, without depletion

of genetic diversity, and despite clear possibility of gene flow with an adjacent, unchanged

population.

4.1   Gradual environmental variation in freshwater availability may partly explain the

organization of non-neutral genetic variation in **A. germinans**

The genetic structure inferred by genome-wide SNPs (Figure 3) suggested the

importance of both neutral and non-neutral environmental drivers of variation. The greatest

divergence was observed between samples influenced by the northern and southern branches

of the South Equatorial sea current (SEC) (Figure 1), acting on the Atlantic coast of South

America, corroborating previous results found for other coastal trees, using putatively neutral

molecular markers (Francisco, Mori, Alves, Tambarussi, & Souza, 2018; Mori, Zucchi, &

Souza, 2015; Takayama, Tateishi, Murata, & Kajita, 2008). The flow direction of coastal

currents provides a neutral explanation to the structure of genome-wide diversity of sea

dispersed species, due to restrictions in the dispersal of buoyant propagules, likely facilitating

the accumulation of random genetic divergences (Francisco et al., 2018; Mori, Zucchi, &

Souza, 2015; Wee et al., 2014; Yan, Duke, & Sun, 2016). Our results also suggest that the

lower precipitation of the warmest quarter in northern sites (AMMC and Northeast Brazil

regions) (Figure 1) may play an additional, non-neutral role in shaping this north-south

divergence. We hypothesize that the more even distribution of rainfall throughout the year in

the TMD site likely alleviates water-stress in *A. germinans*, whereas limited rainfall in the

warmest quarter of northern regions increases soil aridity and salinity, reducing opportunities

for rehydration and potentially favoring increased drought-tolerance. It is plausible that this

environmental filter contributes to the genetic divergence observed between TMD and

remaining populations (Figure 3). This hypothesis is corroborated by the detection of 26 loci

candidate for selection correlated to the spatial variation in the precipitation of the warmest

quarter over the study area (Figure 4b). Although most of these loci were poorly

characterized, hampering inferences about their functional relevance in the environmental

context, we were able to associate 11 candidates with biological processes influenced by

drought, as photosynthesis, cell wall metabolism, cell elongation, plant growth, protein

protection from stress-induced degradation and regulation of abscisic acid signaling (Figure 4,

Table 3). The adaptive importance of freshwater limitation in tropical trees was also

suggested in the mangrove, *A. schaueriana* (M. V. Cruz et al., 2018), and in tropical forests

and savannas (Ciemer et al., 2019), for which it was similarly suggested that an

environmental filtering mechanism driven by rainfall variability likely favored the survival of

more drought resistant lineages.

Although *Avicennia* propagules can remain viable for long periods and present

transoceanic dispersal (Mori, Zucchi, Sampaio, et al., 2015), at a finer scale, we observed a

genetic divergence between sites on the AMMC region (MRJ and PA-humid) and on

Northeast Brazilian mangroves (ALC, PNB and PRC) (Figure 3). The AMMC region shows

higher annual precipitation and is more strongly influenced by riverine freshwater inputs than

the remaining sites, due to its closer proximity to the Amazon River Delta (Figure 1).

Conversely, the Northeast Brazilian coastline is characterized by reduced rainfall and the lack

of riverine freshwater inputs (Figure 1), which limit plants access to soil freshwater, due to

increased salinity, potentially contributing to the local specialization of individuals. In fact,

we were able to detect two loci associated with sea surface salinity data and total annual

precipitation, based on both F_{ST} outlier methods and G-E association tests (Figure 4b). These

loci showed similarity to a poorly characterized protein kinase and to an RNA hydrolase

(Table 3). Given the unclear functional relevance of these putative adaptive loci in the

environmental context, we recommend future efforts to analyze the role of demographic

history in *A. germinans*, to find additional explanations for the genetic structure observed

along the equatorial Brazilian coastline (Figure 3). Additionally, candidate loci detected in our

study may play a role in drought adaptations, but their molecular functions need to be further

characterized in plants adapted to physiological drought. The functional role of these proteins

in drought tolerance may not get highlighted in genetic screenings performed on drought-

sensitive model plants, distantly related to *A. germinans*.

4.2   Rapid evolution of **A. germinans** *in response to abrupt limitation in access to soil*

freshwater

The PA-arid population of *A. germinans* was originated after 1974, when the

construction of the Bragança-Ajuruteua road altered the hydrology of part of the mangrove

forest in the AMMC region (Figure 2a-b) (Cohen & Lara, 2003). This sudden environmental

change caused a large dieback of the mangrove vegetation, followed by a gradual

recolonization, mainly by *A. germinans*. We estimate that the time since this event occurred,

could represent at most 40 reproductive cycles of *A. germinans*, based on previous

observations of closely related *Avicennia* sp. (Almahasheer et al., 2016; M. V. Cruz et al.,

2018), although a conservative estimate could be four generations (Polidoro et al., 2010). The

modification of the architecture of recolonizing individuals of PA-arid, from previous

tall/arboreal, to now dwarf/shrub (Figure 2b) (Pranchai et al., 2017), suggests that limitation

in plant access to soil water favored smaller tree sizes, one of the most integrative

characteristic of drought resistance (Bennett et al., 2015; Corlett, 2016; Rowland et al., 2015).

On the one hand, we cannot determine how much of the observed difference in tree size is

inherited or determined by phenotypic plasticity. On the other hand, our results revealed a

substantial change in genome-wide allele frequencies of trees, which likely represent samples

of the first generations of the recolonizing population (Figure 3a, Table 2). Interestingly,

changes in allele frequencies at founding occurred without reductions in levels of diversity, as

estimated by various population genetics parameters (Table 2). Moreover, consistent with an

adaptive response to selection, likely driven by restrictions in plants access to soil freshwater,

F_{ST} outlier tests detected 56 candidate loci, of which eight were present within transcripts

associated with functions as suppression of cell elongation, wood and xylem tracheids

formation, photosynthetic machinery biogenesis and repair, regulation of adaptation to stress

and of protection from stress-induced protein degradation (Figure 4a, Table 3). The

simultaneous presence of high levels of diversity in the founding population (Table 2) and

strong selection on individual traits is common in natural populations, and is a signature of

substantial multivariate genetic constraints (B. Walsh & Blows, 2009), as expected in this

specific case, given the magnitude of environmental change.

A surprising aspect of the population genetic analyses, however, was that genome-

wide changes in allele frequencies occurred despite expectations of high levels of gene flow,

given the proximity (<10.0 m) of contrasting populations, and characteristics of the species,

which is preferably allogamous, with entomophyllic flowers (Mori, Zucchi, & Souza, 2015;

Nadia et al., 2013; Nettel-Hernanz et al., 2013). In the absence of a geographic barrier to gene

flow, recombination is expected to hinder adaptation, unless there is very strong selection or

reproductive isolation (Kawecki & Ebert, 2004; Mayr, 1963). As we expect selection to be

very strong in this case, due to extreme environmental change, it could be enough to

counteract the homogenizing effect of gene flow or even to cause shifts in the timing of

flowering (Cho, Yoon, & An, 2017), resulting in reproductive isolation.

Besides genetic changes, we also identified transcripts expression differences between

seedlings from PA-arid and PA-humid sites after acclimatization in pots under shaded, well-

watered conditions (Figure 2c, Figure 5), indicating an additional molecular basis for

phenotypic divergences. Because differential gene expression influence trait variation (Wolf,

Lindell, & Backstrom, 2010), it can be substantial between distinct locally adapted

populations (Akman, Carlson, Holsinger, & Latimer, 2016; Gould, Chen, & Lowry, 2018).

Differential expression was observed mainly in roots and stems of *A. germinans* seedlings,

which are the first organs exposed to increased salinity and water deficits in mangroves, thus

where osmotic stress sensing and signaling to the whole plant are triggered (Chaves, Maroco,

& Pereira, 2003; Janiak, Kwas□niewski, & Szarejko, 2016). Such transcriptome changes

were associated with various biological processes previously identified to be involved in

central aspects of osmotic-, heat- and UV-stress in model and non-model plants (Ding et al.,

2013; Fan et al., 2018; Zhang et al., 2015). Being sessile under extreme environmental

conditions, PA-arid plants should need to rapidly adjust photosynthesis gene expression to

intermittent freshwater availability (Chaves, Flexas, & Pinheiro, 2009; Urban, Aarrouf, &

Bidel, 2017). The increased freshwater supply and the restriction in solar irradiance under

experimental conditions, compared to source-site (Figure 2c), likely required PA-arid

individuals to broadly adjust their photosynthetic machinery. Conversely, although

environmental conditions were relaxed in the RNA-Seq experiment, we observed several

transcripts associated with high temperature, drought and salinity response among DET-Arid,

including epicuticular wax and cutin synthesis, export and deposition, accumulation of

osmoprotectants and antioxidants and ion homeostasis. We suggest that these findings

represent key regulatory mechanisms, by which PA-arid plants tolerate a predominantly

warm, highly saline and dry environment, with extreme UV levels. These strategies may have

enhanced the relative capacity of PA-arid seedlings to grow under a high leaf water-deficit

status and infrequent freshwater inputs.

We suggest that future efforts to understand the molecular mechanisms of adaptation

of the PA-arid population should focus on characterizing epigenetic differences between PA-

arid and PA-humid. Epigenetic changes can emerge faster than adaptive genetic changes, also

playing an important role in early adaptive process (Kenkel & Matz, 2016; Pavey, Nosil, &

Rogers, 2010). Although the mechanisms are not entirely clear, our results suggest, through

independent molecular approaches, that a rapid evolution of this population occurred, with

gene expression and genetic changes. The desertification of the PA-arid site represents a

drastic and persistent environmental change, analogous to a freshwater exclusion experiment

for mangroves of the tropical Brazilian coastline, in which this limitation likely caused the

rapid evolution of recolonizing *A. germinans* individuals. Although it is difficult to

demonstrate rapid evolution in nature, there is growing empirical evidence that when

environmental selection is very intense, evolutionary processes may occur on a very fast time

scale (Amorim et al., 2017; Donihue et al., 2018; Schoener, 2011), as observed in this case.

4.3   Implications for conservation

Wetlands of the Northeastern coast of Brazil and the ecosystem services they provide

are predicted to be particularly impacted by future reductions in precipitation and freshwater

availability (Osland et al., 2018). Despite these threats, the extant gene pool of northern

populations of *A. germinans* seems to harbor sufficient diversity to enable species persistence

through natural selection of drought-resistant plants, as observed in the case of PA-arid.

Recolonizing individuals in this location share alleles with individuals from distant areas, as

the PRC and ALC sites (Figure 3a), in Northeast Brazil, suggesting that these forests might

have contributed as source of adaptive variation through sea-dispersed propagules in the

recolonization of the PA-arid site, in addition to the selection on migrants from adjacent areas.

In the context of an increasingly drying climate, reforestation plans for populations of

A. germinans located south of the SEC should consider using mixed stocks of seedlings

(Moritz, 1999), to introduce genetic variation associated with increased drought-tolerance

present in Northeastern mangroves, while also maintaining local alleles, possibly associated

with site-specific environmental characteristics.

4.4   Concluding remarks

We provide novel insights into how limited access to soil freshwater can change allele

frequencies, gene expression and phenotypes of a dominant tropical tree. These findings are

consistent with the predictions of a hypothesis of natural selection, but we acknowledge that

other evolutionary processes could also play a role in driving the observed patterns, opening

new opportunities for future investigation. Research on the genomic basis of tree adaptation

are often limited by difficulties in implementing empirical tests, given their long generation

time and the scarcity of basic biological information. In tropical forests, mostly found in

developing countries, the lack of resources imposes additional challenges for adaptation

studies. Advances in the understanding of the genomic basis of drought-tolerance in tropical

trees can support effective protection plans and mitigating climate change. As shown in this

study, such knowledge can improve predictions of the persistence of the ecosystems they

form and services they provide and generate key insight for conservation and management

efforts (Holliday et al., 2017; Moran, Hartig, & Bell, 2016).

Acknowledgments

We acknowledge Ilmarina Menezes for assistance with fieldwork and HPC services at

Louisiana State University. M.V.C. received fellowships from the São Paulo Research

Foundation - FAPESP (PhD 2013/26793-7) and from the Coordination for the Improvement

of Higher Education Personnel - CAPES Computational Biology Program (PhD SWE

8084/2015-07, PhD 88887.177158/2018-00). G.M.M. received fellowships from FAPESP

(PD 2013/08086-1, PD BEPE 2014/22821-9) and a grant from the Brazilian National Council

for Scientific and Technological Development - CNPq (PD 448286/2014-9). M.D. and D-

H.O. received grants from the United States National Science Foundation award (MCB

1616827) and the Next-Generation BioGreen21 Program of Republic of Korea (PJ01317301).

R.S.O. received a CNPq productivity scholarship and a grant from Microsoft research-

FAPESP (2011/52072-0). A.P.S. received a grant from CAPES - Computational Biology

Conflicts of interest

754 The authors have no conflict of interest to disclose.

6. References

Akman, M., Carlson, J. E., Holsinger, K. E., & Latimer, A. M. (2016). Transcriptome
sequencing reveals population differentiation in gene expression linked to functional
traits and environmental gradients in the South African shrub Protea repens. *New
Phytologist*, *210*(1), 295–309. doi:10.1111/nph.13761
Al-Whaibi, M. H. (2011). Plant heat-shock proteins☐: A mini review. *Journal of King Saud
University - Science*, *23*(2), 139–150. doi:10.1016/j.jksus.2010.06.022
Alexander, D. H., Novembre, J., & Lange, K. (2009). Fast model-based estimation of ancestry
in unrelated individuals. *Genome Research*, *19*(9), 1655–1664.
doi:10.1101/gr.094052.109
Almahasheer, H., Duarte, C. M., & Irigoien, X. (2016). Phenology and Growth dynamics of
Avicennia marina in the Central Red Sea. *Scientific Reports*, *6*, 37785.
doi:10.1038/srep37785
Alvares, C. A., Stape, J. L., Sentelhas, P. C., Gonçalves, J. L. de M., & Sparovek, G. (2013).
Köppen's climate classification map for Brazil. *Meteorologische Zeitschrift*, *22*(6), 711–
728. doi:10.1127/0941-2948/2013/0507
Amorim, M. E., Schoener, T. W., Ramalho, G. C. C. S., Lins, A. C. R., Piovia-Scottd, J., &
Brandão, R. A. (2017). Lizards on newly created islands independently and rapidly adapt
in morphology and diet. *Proceedings of the National Academy of Sciences*, *114*(33),
8812–8816. doi:10.1073/pnas.1709080114
Antao, T., Lopes, A., Lopes, R. J., Beja-Pereira, A., & Luikart, G. (2008). LOSITAN: A
workbench to detect molecular adaptation based on a F st -outlier method. *BMC
Bioinformatics*, *9*(1), 1–5. doi:10.1186/1471-2105-9-323
Aranda, I., Cano, F. J., Gascó, A., Cochard, H., Nardini, A., Mancha, J. A., … Sánchez-
gómez, D. (2014). Variation in photosynthetic performance and hydraulic architecture
across European beech (Fagus sylvatica L .) populations supports the case for local
adaptation to water stress. *Tree Physiology*, *35*(1), 34–46. doi:10.1093/treephys/tpu101
Asner, G. P., Brodrick, P. G., Anderson, C. B., Vaughn, N., Knapp, D. E., & Martin, R. E.
(2016). Progressive forest canopy water loss during the 2012 – 2015 California drought.
Proceedings of the National Academy of Sciences, *113*(2), 249–255.
doi:10.1073/pnas.1523397113
Bang, W. Y., Hata, A., Jeong, I. S., Umeda, T., Masuda, T., Chen, J., … Bahk, J. D. (2009).
AtObgC, a plant ortholog of bacterial Obg, is a chloroplast-targeting GTPase essential
for early embryogenesis. *Plant Molecular Biology*, *71*(4–5), 379–390.
doi:10.1007/s11103-009-9529-3

Bartlett, M. K., Scoffoni, C., & Sack, L. (2012). The determinants of leaf turgor loss point
and prediction of drought tolerance of species and biomes: A global meta-analysis.
Ecology Letters, *15*, 393–405. doi:10.1111/j.1461-0248.2012.01751.x
Beaumont, M. A., & Nichols, R. A. (1996). Evaluating Loci for Use in the Genetic Analysis
of Population Structure. *Proceedings of the Royal Society B: Biological Sciences*,
263(1377), 1619–1626. doi:10.1098/rspb.1996.0237
Bennett, A. C., McDowell, N. G., Allen, C. D., & Anderson-Teixeira, K. J. (2015). Larger
trees suffer most during drought in forests worldwide. *Nature Plants*, *1*, 1–5.
doi:10.1038/NPLANTS.2015.139
Berardini, T. Z., Reiser, L., Li, D., Mezheritsky, Y., Muller, R., Strait, E., & Huala, E. (2015).
The Arabidopsis Information Resource: Making and Mining the "Gold Standard"
Annotated Reference Plant Genome. *Genesis*, *53*, 474–485. doi:10.1002/dvg.22877
Blake, J. A., Christie, K. R., Dolan, M. E., Drabkin, H. J., Hill, D. P., Ni, L., … Westerfeld,
M. (2015). Gene ontology consortium: Going forward. *Nucleic Acids Research*, *43*,
D1049–D1056. doi:10.1093/nar/gku1179
Bohne, A. V., Schwenkert, S., Grimm, B., & Nickelsen, J. (2016). Roles of Tetratricopeptide
Repeat Proteins in Biogenesis of the Photosynthetic Apparatus. *International Review of
Cell and Molecular Biology*, *324*, 187–227. doi:10.1016/bs.ircmb.2016.01.005
Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., & Madden,
T. L. (2009). BLAST+: architecture and applications. *BMC Bioinformatics*, *10*, 421.
doi:10.1186/1471-2105-10-421
Charng, Y. -y., Liu, H. -c., Liu, N. -y., Chi, W. -t., Wang, C. -n., Chang, S. -h., & Wang, T. -t.
(2006). A Heat-Inducible Transcription Factor, HsfA2, Is Required for Extension of
Acquired Thermotolerance in *Arabidopsis*. *Plant Physiology*, *143*(1), 251–262.
doi:10.1104/pp.106.091322
Chaves, M. M., Maroco, J. P., & Pereira, J. S. (2003). Understanding plant responses to
drought - from genes to the whole plant. *Functional Plant Biology*, *30*(3), 239–264.
doi:10.1071/FP02076
Cho, L. H., Yoon, J., & An, G. (2017). The control of flowering time by environmental
factors. *Plant Journal*, *90*(4), 708–719. doi:10.1111/tpj.13461
Choat, B., Brodribb, T. J., Brodersen, C. R., Duursma, R. A., López, R., & Medlyn, B. E.
(2018). Triggers of tree mortality under drought. *Nature*, *558*(7711), 531–539.
doi:10.1038/s41586-018-0240-x
Ciemer, C., Boers, N., Hirota, M., Kurths, J., Müller-hansen, F., Oliveira, R. S., &
Winkelmann, R. (2019). Higher resilience to climatic disturbances in tropical vegetation
exposed to more variable rainfall. *Nature Geoscience*, *12*, 174–179. doi:10.1038/s41561-
019-0312-z
Clough, B. F., Sim, R. G., Inlet, T., Bay, M., & Rivers, I. (1989). Oecologia of mangroves in
response to salinity and vapour pressure deficit. *Oecologia*, (459), 38–44.
Cohen, M. C. L., & Lara, R. J. (2003). Temporal changes of mangrove vegetation boundaries
in Amazonia: Application of GIS and remote sensing techniques. *Wetlands Ecology and
Management*, *11*(4), 223–231. doi:10.1023/A:1025007331075
Corlett, R. T. (2016). The Impacts of Droughts in Tropical Forests. *Trends in Plant Science*,
21(7), 584–593. doi:10.1016/j.tplants.2016.02.003
Correll, M. J., Coveney, K. M., Raines, S. V., Mullen, J. L., Hangarter, R. P., & Kiss, J. Z.
(2003). Phytochromes play a role in phototropism and gravitropism in *Arabidopsis* roots.
Advances in Space Research, *31*(10), 2203–2210. doi:10.1016/S0273-1177(03)00245-X
Cruz, M. V., Mori, G. M., Müller, C. S., Cristina, C., Oh, D., Dassanayake, M., … Silva, R.

(2018). Local adaptation of a dominant coastal tree to freshwater availability and solar
radiation suggested by genomic and ecophysiological approaches. *Biorxiv*, 1–43.
doi:10.1101/378612

Cruz, M. V., Mori, G. M., Oh, D., Dassanayake, M., Zucchi, M. I., Oliveira, R. S., & Souza,
A. P. (2019). Data from: Molecular responses to freshwater limitation in the mangrove
tree Avicennia germinans (Acanthaceae). *Dryad Digital Repository*.
doi:doi:10.5061/dryad.h11t255

da Costa, C. L., Galbraith, D., Almeida, S., Tanaka Portela, B. T., da Costa, M., de Athaydes
Silva Junior, J., … Meir, P. (2010). Effect of seven years of experimental drought on the
aboveground biomass storage of an eastern Amazonian rainforest. *New Phytologist*, *187*,
579–591. doi:10.1111/j.1469-8137.2010.03309.x

Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., … Durbin,
R. (2011). The variant call format and VCFtools. *Bioinformatics*, *27*(15), 2156–2158.
doi:10.1093/bioinformatics/btr330

Daniel, B., Pavkov-Keller, T., Steiner, B., Dordic, A., Gutmann, A., Nidetzky, B., …
Macheroux, P. (2015). Oxidation of monolignols by members of the berberine bridge
enzyme family suggests a role in plant cell wall metabolism. *Journal of Biological*
Chemistry, *290*(30), 18770–18781. doi:10.1074/jbc.M115.659631

De Frenne, P., Graae, B. J., Rodríguez-Sánchez, F., Kolb, A., Chabrerie, O., Decocq, G., …
Verheyen, K. (2013). Latitudinal gradients as natural laboratories to infer species'
responses to temperature. *Journal of Ecology*, *101*(3), 784–795. doi:10.1111/1365-
2745.12074

Ding, Y., Liu, N., Virlouvet, L., Riethoven, J. J., Fromm, M., & Avramova, Z. (2013). Four
distinct types of dehydration stress memory genes in *Arabidopsis thaliana*. *BMC Plant*
Biology, *13*(1). doi:10.1186/1471-2229-13-229

Donihue, C. M., Herrel, A., Fabre, A. C., Kamath, A., Geneva, A. J., Schoener, T. W., …
Losos, J. B. (2018). Hurricane-induced selection on the morphology of an island lizard.
Nature, *560*(7716), 88–91. doi:10.1038/s41586-018-0352-3

Donovan, L. A., Ludwig, F., Rosenthal, D. M., Rieseberg, L. H., & Dudley, S. A. (2009).
Phenotypic selection on leaf ecophysiological traits in Helianthus. *New Phytologist*,
183(3), 868–879. doi:10.1111/j.1469-8137.2009.02916.x.

Duarte, C. M., Thampanya, U., Terrados, J., Geertz-Hansen, O., & Fortes, M. D. (1999). The
determination of the age and growth of SE Asian mangrove seedlings from internodal
counts. *Mangroves and Salt Marshes*, *3*(4), 251–257. doi:10.1023/A:1009967401337

Ellison, A. M., Farnsworth, E. J., & Merkt, R. E. (1999). Origins of mangrove ecosystems and
the mangrove biodiversity anomaly. *Global Ecology and Biogeography*, *8*(2), 95–115.
doi:10.1046/j.1466-822X.1999.00126.x

ElSayed, A. I., Rafudeen, M. S., & Golldack, D. (2014). Physiological aspects of raffinose
family oligosaccharides in plants: Protection against abiotic stress. *Plant Biology*, *16*(1),
1–8. doi:10.1111/plb.12053

Etterson, J. R. (2004). Evolutionary potential of Chamaecrista fasciculata in relation to
climate change. I . Clinal patterns of selection along and environmental gradient in the
great plains. *Evolution*, *58*(7), 1446–1458. doi:10.1111/j.0014-3820.2004.tb01726.x

Fan, L., Wang, G., Hu, W., Pantha, P., Tran, K., Zhang, H., … Qiu, Q. (2018). Plant
Physiology and Biochemistry Transcriptomic view of survival during early seedling
growth of the extremophyte Haloxylon ammodendron. *Plant Physiology and*
Biochemistry, *132*(9), 475–489. doi:10.1016/j.plaphy.2018.09.024

Finn, R. D., Bateman, A., Clements, J., Coggill, P., Eberhardt, R. Y., Eddy, S. R., … Punta,

M. (2014). Pfam: The protein families database. *Nucleic Acids Research*, *42*(D1), 222–
230. doi:10.1093/nar/gkt1223
Francisco, P. M., Mori, G. M., Alves, F. M., Tambarussi, E., & Souza, A. P. De. (2018).
Population genetic structure, introgression, and hybridization in the genus Rhizophora
along the Brazilian coast. *Ecology and Evolution*, *8*(6), 3491–3504.
doi:10.1002/ece3.3900
Frichot, E., & François, O. (2015). LEA: An R package for landscape and ecological
association studies. *Methods in Ecology and Evolution*, *6*(8), 925–929.
doi:10.1111/2041-210X.12382
Frydman, J. (2001). Folding of newkly translated proteins in vivo: the role of molecular
chaperones. *Annual Review of Biochemistry*, *701*(1), 603–647.
Goudet, J. (2005). Hierfstat, a package for R to compute and test hierarchical F -statistics.
Molecular Ecology Notes, *5*(1), 184–186. doi:10.1111/j.1471-8278
Gould, B. A., Chen, Y., & Lowry, D. B. (2018). Gene Regulatory Divergence Between
Locally Adapted Ecotypes in Their Native Habitats. *Molecular Ecology*, *27*(21), 4174–
4188. doi:10.1111/mec.14852
Hacke, U. G., Sperry, J. S., Wheeler, J. K., & Castro, L. (2006). Scaling of angiosperm xylem
structure with safety and efficiency. *Tree Physiology*, *26*(6), 689–701.
doi:10.1093/treephys/26.6.689
Heschel, M. S., & Riginos, C. (2005). Mechanisms of selection for drought stress tolerance
and avoidance in *Impatiens capensis* (Balsaminaceae). *American Journal of Botany*,
92(1), 37–44. doi:10.3732/ajb.92.1.37
Hijmans, R. J., Cameron, S. E., Parra, J. L., Jones, P. G., & Jarvis, A. (2005). Very high
resolution interpolated climate surfaces for global land areas. *International Journal of*
Climatology, *25*(15), 1965–1978. doi:10.1002/joc.1276
Holliday, J. A., Aitken, S. N., Cooke, J. E. K., Fady, B., González-Martínez, S. C., Heuertz,
M., … Plomion, C. (2017). Advances in ecological genomics in forest trees and
applications to genetic resources conservation and breeding. *Molecular Ecology*, 706–
717. doi:10.1111/mec.13963
Huang, S., Chen, X., Li, M., Ao, K., Huang, S., Chen, X., … Li, X. (2016). Plant TRAF
Proteins Regulate NLR Immune Receptor Turnover. *Cell Host & Microbe*, *19*(2), 204–
215. doi:10.1016/j.chom.2016.01.005
Hussey, S. G., Mizrachi, E., Spokevicius, A. V., Bossinger, G., Berger, D. K., & Myburg, A.
A. (2011). SND2, a NAC transcription factor gene, regulates genes involved in
secondary cell wall development in *Arabidopsis* fibres and increases fibre cell area in
Eucalyptus. *BMC Plant Biology*, *11*, 1–17. doi:10.1186/1471-2229-11-173
Janiak, A., Kwaśniewski, M., & Szarejko, I. (2016). Gene expression regulation in roots
under drought. *Journal of Experimental Botany*, *67*(4), 1003–1014.
doi:10.1093/jxb/erv512
Jombart, T., & Ahmed, I. (2011). adegenet 1 .3-1: new tools for the analysis of genome-wide
SNP data. *Bioinformatics*, *27*(21), 3070–3071. doi:10.1093/bioinformatics/btr521
Jombart, T., Devillard, S., & Balloux, F. (2010). Discriminant analysis of principal
components: a new method for the analysis of genetically structured populations. *BMC*
Genetics, *11*(1), 94. doi:10.1186/1471-2156-11-94
Kamvar, Z. N., Tabima, J. F., & Grünwald, N. J. (2014). Poppr: an R package for genetic
analysis of populations with clonal , partially clonal , and / or sexual reproduction.
PeerJ, *2:e281*. doi:10.7717/peerj.281
Kawecki, T. J., & Ebert, D. (2004). Conceptual issues in local adaptation. *Ecology Letters*,

7(12), 1225–1241. doi:10.1111/j.1461-0248.2004.00684.x

Keller, S. R., Soolanayakanahally, R. Y., Guy, R. D., Silim, S. N., Olson, M. S., & Tiffin, P.
(2011). Climate-driven local adaptation of ecophysiology and phenology in balsam
poplar, Populus balsamifera L. (Salicaceae). *American Journal of Botany*, *98*(1), 99–108.
doi:10.3732/ajb.1000317

Kenkel, C. D., & Matz, M. V. (2016). Gene expression plasticity as a mechanism of coral
adaptation to a variable environment. *Nature Ecology & Evolution*, *1*, 059667.
doi:10.1038/s41559-016-0014 |

Kitaya, Y., Yabuki, K., Kiyota, M., Tani, A., Hirano, T., & Aiga, I. (2002). Gas exchange and
oxygen concentration in pneumatophores and prop roots of four mangrove species.
Trees, *16*, 155–158. doi:10.1007/s00468-002-0167-5

Klepikova, A. V., Kasianov, A. S., Gerasimov, E. S., Logacheva, M. D., & Penin, A. A.
(2016). A high resolution map of the Arabidopsis thaliana developmental transcriptome
based on RNA-seq profiling. *Plant Journal*, *88*(6), 1058–1070. doi:10.1111/tpj.13312

Kozaki, A., & Takeba, G. (1996). Photorespiration protects C3 plants from photooxidation.
Nature, *384*, 557–560. doi:10.1038/384557a0

Krishnamurthy, P., Mohanty, B., Wijaya, E., Lee, D., Lim, T., Lin, Q., … Kumar, P. P.
(2017). Transcriptomics analysis of salt stress tolerance in the roots of the mangrove
Avicennia officinalis. *Nature Scientific Reports*, *7*, 1–19. doi:10.1038/s41598-017-
10730-2

Langmead, B., Trapnell, C., Pop, M., & Salzberg, S. (2009). Ultrafast and memory-efficient
alignment of short DNA sequences to the human genome. *Genome Biol.*, *10*(3), R25.
doi:10.1186/gb-2009-10-3-r25

Lara, R. J., & Cohen, M. C. L. (2006). Sediment porewater salinity, inundation frequency and
mangrove vegetation height in Bragança, North Brazil: An ecohydrology-based
empirical model. *Wetlands Ecology and Management*, *14*(4), 349–358.
doi:10.1007/s11273-005-4991-4

Lawlor, D. W. (2002). Limitation to photosynthesis in water-stressed leaves: Stomata vs.
Metabolism and the role of ATP. *Annals of Botany*, *89*(SPEC. ISS.), 871–885.
doi:10.1093/aob/mcf110

Li, W., & Godzik, A. (2006). Cd-hit: A fast program for clustering and comparing large sets
of protein or nucleotide sequences. *Bioinformatics*, *22*(13), 1658–1659.
doi:10.1093/bioinformatics/btl158

Lin, G., & Sternberg, L. da S. L. (1992). Comparative study of water uptake and
photosynthetic gas exchange between scrub and fringe red mangroves, Rhizophora
mangle L. *Oecologia*, *90*(3), 399–403. doi:10.1007/BF00317697

Lorenzo, L. de, Merchan, F., Laporte, P., Thompson, R., Clarke, J., Sousa, C., & Crespi, M.
(2009). A Novel Plant Leucine-Rich Repeat Receptor Kinase Regulates the Response of
Medicago truncatula Roots to Salt Stress. *The Plant Cell*, *21*(1), 668–680.
doi:10.1105/tpc.108.059576

Lotterhos, K. E., & Whitlock, M. C. (2015). The relative power of genome scans to detect
local adaptation depends on sampling design and statistical method. *Molecular Ecology*,
24(5), 1031–1046. doi:10.1111/mec.13100

Luu, K., Bazin, E., & Blum, M. G. B. (2016). pcadapt: An R package to perform genome
scans for selection based on principal component analysis. *Molecular Ecology*
Resources, *33*, 67–77. doi:10.1111/1755-0998.12592

Manfield, I. W., Devlin, P. F., Jen, C.-H., Westhead, D. R., & Gilmartin, P. M. (2006).
Conservation, Convergence, and Divergence of Light-Responsive, Circadian-Regulated,

and Tissue-Specific Expression Patterns during Evolution of the *Arabidopsis* GATA
Gene Family. *Plant Physiology*, *143*(2), 941–958. doi:10.1104/pp.106.090761
Mayr, E. (1963). *Animal species and evolution* (Belknap). Cambridge.
McAdam, S. A. M., & Brodribb, T. J. (2015). The Evolution of Mechanisms Driving the
Stomatal Response to Vapor Pressure Deficit. *Plant Physiology*, *167*(3), 833–843.
doi:10.1104/pp.114.252940
McDonough, M. A., Kadirvelraj, R., Harris, P., Poulsen, J. C. N., & Larsen, S. (2004).
Rhamnogalacturonan lyase reveals a unique three-domain modular structure for
polysaccharide lyase family 4. *FEBS Letters*, *565*(1–3), 188–194.
doi:10.1016/j.febslet.2004.03.094
McDowell, N. G., & Allen, C. D. (2015). Darcy's law predicts widespread forest mortality
under climate warming. *Nature Climate Change*, *5*(7), 669–672.
doi:10.1038/nclimate2641
McRae, G. J. (1980). A Simple Procedure for Calculating Atmospheric Water Vapor
Concentration. *Journal of the Air Pollution Control Association*, *30*(4), 394–394.
doi:10.1080/00022470.1980.10464362
Menezes, M. P. M. de, Berger, U., & Mehlig, U. (2008). Mangrove vegetation in Amazonia: a
review of studies from the coast of Pará and Maranhão States, north Brazil. *Acta*
Amazonica, *38*(3), 403–420. doi:10.1590/S0044-59672008000300004
Moran, E. V., Hartig, F., & Bell, D. M. (2016). Intraspecific trait variation across scales:
Implications for understanding global change responses. *Global Change Biology*, *22*(1),
137–150. doi:10.1111/gcb.13000
Mori, G. M., Zucchi, M. I., Sampaio, I., & Souza, A. P. (2015). Species distribution and
introgressive hybridization of two Avicennia species from the Western Hemisphere
unveiled by phylogeographic patterns. *BMC Evolutionary Biology*, *15*(1), 1–15.
doi:10.1186/s12862-015-0343-z
Mori, G. M., Zucchi, M. I., & Souza, A. P. (2015). Multiple-geographic-scale genetic
structure of two mangrove tree species: the roles of mating system, hybridization, limited
dispersal and extrinsic factors. *Plos One*, *10*, e0118710.
doi:10.1371/journal.pone.0118710
Moritz, C. (1999). Conservation units and translocations: Strategies for conserving
evolutionary processes. *Hereditas*, *130*(3), 217–228. doi:10.1111/j.1601-
5223.1999.00217.x
Moriya, Y., Itoh, M., Okuda, S., Yoshizawa, A. C., & Kanehisa, M. (2007). KAAS: An
automatic genome annotation and pathway reconstruction server. *Nucleic Acids*
Research, *35*, 182–185. doi:10.1093/nar/gkm321
Müller, M., Kunz, H. H., Schroeder, J. I., Kemp, G., Young, H. S., & Ekkehard Neuhaus, H.
(2014). Decreased capacity for sodium export out of *Arabidopsis* chloroplasts impairs
salt tolerance, photosynthesis and plant performance. *Plant Journal*, *78*(4), 646–658.
doi:10.1111/tpj.12501
Murata, Y., Mori, I. C., & Munemasa, S. (2015). Diverse Stomatal Signaling and the Signal
Integration Mechanism. *Annual Review of Plant Biology*, *66*(1), 369–392.
doi:10.1146/annurev-arplant-043014-114707
Nadia, T. D. E. L., Menezes, N. L. D. E., & Machado, I. C. (2013). Floral traits and
reproduction of Avicennia schaueriana Moldenke (Acanthaceae): a generalist
pollination system in the Lamiales. *Plant Species Biology*, *28*(1), 70–80.
doi:10.1111/j.1442-1984.2011.00361.x
Narum, S. R., & Hess, J. E. (2011). Comparison of FST outlier tests for SNP loci under

selection. *Molecular Ecology Resources*, *11*(SUPPL. 1), 184–194. doi:10.1111/j.1755-
0998.2011.02987.x
Nascimento Jr., W. R., Souza-Filho, P. W. M., Proisy, C., Lucas, R. M., & Rosenqvist, A.
(2013). Mapping changes in the largest continuous Amazonian mangrove belt using
object-based classification of multisensor satellite imagery. *Estuarine, Coastal and Shelf*
Science, *117*, 83–93. doi:10.1016/j.ecss.2012.10.005
Nettel-Hernanz, A., S.Dodd, R., Ochoa-Zavala, M., Tovilla-Hernández, C., & Días-Gallegos,
J. R. (2013). Mating system analyses of tropical populations of the Black Mangrove,
Avicennia germinans (L.) L. (Avicenniaceae). *Botanical Sciences*, *91*(1), 115–117.
Nishizawa, A., Yabuta, Y., & Shigeoka, S. (2008). Galactinol and Raffinose Constitute a
Novel Function to Protect Plants from Oxidative Damage. *Plant Physiology*, *147*(3),
1251–1263. doi:10.1104/pp.108.122465
Novick, K. A., Ficklin, D. L., Stoy, P. C., Williams, C. A., Bohrer, G., Oishi, A. C., …
Phillips, R. P. (2016). The increasing importance of atmospheric demand for ecosystem
water and carbon fluxes. *Nature Climate Change*, *6*(11), 1023–1027.
doi:10.1038/nclimate3114
O'Leary, N. A., Wright, M. W., Brister, J. R., Ciufo, S., Haddad, D., McVeigh, R., … Pruitt,
K. D. (2016). Reference sequence (RefSeq) database at NCBI: Current status, taxonomic
expansion, and functional annotation. *Nucleic Acids Research*, *44*, D733–D745.
doi:10.1093/nar/gkv1189
Ohgishi, M., Saji, K., Okada, K., & Sakai, T. (2004). Functional analysis of each blue light
receptor, cry1, cry2, phot1, and phot2, by using combinatorial multiple mutants in
Arabidopsis. *Proceedings of the National Academy of Sciences*, *101*(8), 2223–2228.
doi:10.1073/pnas.0305984101
Oliveira, R. R., Viana, A. J. C., Reátegui, A. C. E., & Vincentz, M. G. A. (2015). An efficient
method for simultaneous extraction of high-quality RNA and DNA from various plant
tissues. *Genetics and Molecular Research*, *14*(4), 18828–18838.
doi:10.4238/2015.December.28.32
Osland, M. J., Gabler, C. A., Grace, J. B., Day, R. H., Mccoy, M. L., Mcleod, J. L., …
Hartley, S. B. (2018). Climate and plant controls on soil organic matter in coastal
wetlands. *Global Change Biology*, (February). doi:10.1111/gcb.14376
Pagnussat, G. C., Yu, H.-J., Ngo, Q. A., Rajani, S., Mayalagu, S., Johnson, C. S., …
Sundaresan, V. (2005). Genetic and molecular identification of genes required for female
gametophyte development and function in *Arabidopsis*. *Development*, *132*(3), 603–614.
doi:10.1242/dev.01595
Patel, R. K., & Jain, M. (2012). NGS QC toolkit: A toolkit for quality control of next
generation sequencing data. *PLoS ONE*, *7*(2), e30619.
doi:10.1371/journal.pone.0030619
Pavey, S. A., Nosil, P., & Rogers, S. M. (2010). The role of gene expression in ecological
speciation. *Annals of the New York Academy of Sciences*, *1206*, 110–129.
doi:10.1111/j.1749-6632.2010.05765.x
Pedmale, U. V., Huang, S. S. C., Zander, M., Cole, B. J., Hetzel, J., Ljung, K., … Chory, J.
(2016). Cryptochromes Interact Directly with PIFs to Control Plant Growth in Limiting
Blue Light. *Cell*, *164*(1–2), 233–245. doi:10.1016/j.cell.2015.12.018
Pesquet, E., Korolev, A. V., Calder, G., & Lloyd, C. W. (2010). The microtubule-associated
protein AtMAP70-5 regulates secondary wall patterning in *Arabidopsis* wood cells.
Current Biology, *20*(8), 744–749. doi:10.1016/j.cub.2010.02.057
Phillips, O. L., Aragão, L. E. O. C., Lewis, S. L., Fisher, J. B., Lloyd, J., López-González, G.,

… Vargas, P. N. (2009). Drought Sensitivity of the Amazon Rainforest. *Science*,
323(5919), 1344–1347. doi:10.1126/science.1164033
Phillips, O. L., Heijden, G. van der, Lewis, S. L., López-González, G., Aragão, L. E. O. C.,
Lloyd, J., … Vilanova, E. (2010). Drought–mortality relationships for tropical forests.
New Phytologist, *187*, 631–646. doi:10.1111/j.1469-8137.2010.03359.x Key
Pieslinger, A. M., Hoepflinger, M. C., & Tenhaken, R. (2010). Cloning of glucuronokinase
from *Arabidopsis thaliana*, the last missing enzyme of the myo-inositol oxygenase
pathway to nucleotide sugars. *Journal of Biological Chemistry*, *285*(5), 2902–2910.
doi:10.1074/jbc.M109.069369
Polidoro, B. A., Carpenter, K. E., Collins, L., Duke, N. C., Ellison, A. M., Ellison, J. C., …
Yong, J. W. H. (2010). The loss of species: Mangrove extinction risk and geographic
areas of global concern. *PLoS ONE*, *5*(4), e10095. doi:10.1371/journal.pone.0010095
Powell, T. L., Wheeler, J. K., de Oliveira, A. A. R., da Costa, A. C. L., Saleska, S. R., Meir,
P., & Moorcroft, P. R. (2017). Differences in xylem and leaf hydraulic traits explain
differences in drought tolerance among mature Amazon rainforest trees. *Global Change
Biology*, *23*(10), 4280–4293. doi:10.1111/gcb.13731
Pranchai, A., Jenke, M., Vogt, J., Grueters, U., Yue, L., Mehlig, U., … Berger, U. (2017).
Density-dependent shift from facilitation to competition in a dwarf Avicennia germinans
forest. *Wetlands Ecology and Management*, 1–12. doi:10.1007/s11273-017-9561-z
Privat, I., Hakimi, M. A., Buhot, L., Favory, J. J., & Lerbs-Mache, S. (2003). Characterization
of *Arabidopsis* plastid sigma-like transcription factors SIG1, SIG2 and SIG3. *Plant
Molecular Biology*, *51*(3), 385–399. doi:10.1023/A:1022095017355
Ramírez-Valiente, J. A., Deacon, N. J., Etterson, J., Center, A., Sparks, J. P., Sparks, K. L., …
Cavender-Bares, J. (2018). Natural selection and neutral evolutionary processes
contribute to genetic divergence in leaf traits across a precipitation gradient in the
tropical oak Quercus oleoides. *Molecular Ecology*, *27*, 2176–2192.
doi:10.1111/mec.14566
Reef, R., & Lovelock, C. E. (2015). Regulation of water balance in Mangroves. *Annals of
Botany*, *115*(3), 385–395. doi:10.1093/aob/mcu174
Robinson, M. D., McCarthy, D. J., & Smyth, G. K. (2010). edgeR: a Bioconductor package
for differential expression analysis of digital gene expression data. *Bioinformatics*, *26*(1),
139–140. doi:10.1093/bioinformatics/btp616
Rowland, L., Da Costa, A. C. L., Galbraith, D. R., Oliveira, R. S., Binks, O. J., Oliveira, A. A.
R., … Meir, P. (2015). Death from drought in tropical forests is triggered by hydraulics
not carbon starvation. *Nature*, *528*(7580), 119–122. doi:10.1038/nature15539
Russello, M. A., Waterhouse, M. D., Etter, P. D., & Johnson, E. A. (2015). From promise to
practice: pairing non-invasive sampling with genomics in conservation. *PeerJ*, *3*, e1106.
doi:10.7717/peerj.1106
Sbrocco, E. J., & Barber, P. H. (2013). MARSPEC: ocean climate layers for marine spatial
ecology. *Ecology*, *94*(4), 979–979. doi:10.1890/12-1358.1
Schaeffer-Novelli, Y., Cintrón-Molero, G., Adaime, R. R., & de Camargo, T. M. (1990).
Variability of mangrove ecosystems along the Brazilian coast. *Estuaries*, *13*(2), 204–
218. doi:10.2307/1351590
Schoener, T. W. (2011). The Newest Synthesis: Understanding the Interplay of Evolutionary
and Ecological Dynamics. *Science*, *331*(6016), 426–429. doi:10.1126/science.1193954
Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V., & Zdobnov, E. M. (2015).
BUSCO: Assessing genome assembly and annotation completeness with single-copy
orthologs. *Bioinformatics*, *31*(19), 3210–3212. doi:10.1093/bioinformatics/btv351

Stone, S. L., Williams, L. A., Farmer, L. M., Vierstra, R. D., & Callis, J. (2006). KEEP ON
GOING, a RING E3 Ligase Essential for *Arabidopsis* Growth and Development, Is
Involved in Abscisic Acid Signaling. *Plant Cell Online*, *18*(12), 3415–3428.
doi:10.1105/tpc.106.046532
Stuart, S. A., Choat, B., Martin, K. C., Holbrook, N. M., & Ball, M. C. (2007). The role of
freezing in setting the latitudinal limits of mangrove forests. *New Phytologist*, *173*(3),
576–583. doi:10.1111/j.1469-8137.2006.01938.x
Taji, T., Ohsumi, C., Iuchi, S., Seki, M., Kasuga, M., Kobayashi, M., … Shinozaki, K.
(2002). Important roles of drought- and cold-inducible genes for galactinol synthase in
stress tolerance in *Arabidopsis thaliana*. *Plant Journal*, *29*(4), 417–426.
doi:10.1046/j.0960-7412.2001.01227.x
Takayama, K., Tateishi, Y., Murata, J., & Kajita, T. (2008). Gene flow and population
subdivision in a pantropical plant with sea-drifted seeds *Hibiscus tiliaceus* and its allied
species: Evidence from microsatellite analyses. *Molecular Ecology*, *17*(11), 2730–2742.
doi:10.1111/j.1365-294X.2008.03799.x
ten Hove, C. A., Bochdanovits, Z., Jansweijer, V. M. A., Koning, F. G., Berke, L., Sanchez-
Perez, G. F., … Heidstra, R. (2011). Probing the roles of LRR RLK genes in *Arabidopsis*
thaliana roots using a custom T-DNA insertion set. *Plant Molecular Biology*, *76*(1–2),
69–83. doi:10.1007/s11103-011-9769-x
Tomlinson, P. B. (1986). *The Botany of Mangroves*. New York: Cambridge University Press.
Tyree, M. T., & Sperry, J. S. (1989). Vulnerability of Xylem to Cavitation and Embolism.
Annual Review of Plant Physiology and Molecular Biology, 19–38.
doi:10.1146/annurev.pp.40.060189.000315
Van der Does, D., Boutrot, F., Engelsdorf, T., Rhodes, J., McKenna, J. F., Vernhettes, S., …
Zipfel, C. (2017). The *Arabidopsis* leucine-rich repeat receptor kinase MIK2/LRR-KISS
connects cell wall integrity sensing, root growth and response to abiotic and biotic
stresses. *PLoS Genetics*, *13*(6), 1–27. doi:10.1371/journal.pgen.1006832
Vogt, J., Lin, Y., Pranchai, A., Frohberg, P., Mehlig, U., & Berger, U. (2014). The importance
of conspecific facilitation during recruitment and regeneration: A case study in degraded
mangroves. *Basic and Applied Ecology*, *15*(8), 651–660. doi:10.1016/j.baae.2014.09.005
Walsh, B., & Blows, M. W. (2009). Abundant Genetic Variation + Strong Selection =
Multivariate Genetic Constraints: A Geometric View of Adaptation. *Annual Review of*
Ecology, Evolution, and Systematics, *40*(1), 41–59.
doi:10.1146/annurev.ecolsys.110308.120242
Walsh, P., Bursać, D., Law, Y. C., Cyr, D., & Lithgow, T. (2004). The J-protein family:
Modulating protein assembly, disassembly and translocation. *EMBO Reports*, *5*(6), 567–
571. doi:10.1038/sj.embor.7400172
Wang, S., Chang, Y., & Ellis, B. (2016). Overview of OVATE FAMILY PROTEINS, A
Novel Class of Plant-Specific Growth Regulators. *Frontiers in Plant Science*, *7*(March),
1–8. doi:10.3389/fpls.2016.00417
Wang, S., Chang, Y., Guo, J., & Chen, J. G. (2007). *Arabidopsis* Ovate Family Protein 1 is a
transcriptional repressor that suppresses cell elongation. *Plant Journal*, *50*(5), 858–872.
doi:10.1111/j.1365-313X.2007.03096.x
Wee, A. K. S., Takayama, K., Asakawa, T., Thompson, B., Onrizal, Sungkaew, S., … Webb,
E. L. (2014). Oceanic currents, not land masses, maintain the genetic structure of the
mangrove Rhizophora mucronata Lam. (Rhizophoraceae) in Southeast Asia. *Journal of*
Biogeography, *41*(5), 954–964. doi:10.1111/jbi.12263
Wingler, A., Lea, P. J., Quick, W. P., & Leegood, R. C. (2000). Photorespiration: metabolic

pathways and their role in stress protection. *Philosophical Transactions of the Royal*
Society B: Biological Sciences, *355*, 1517–1529. doi:10.1098/rstb.2000.0712
Wolf, J. B. W., Lindell, J., & Backstrom, N. (2010). Speciation genetics: current status and
evolving approaches. *Philosophical Transactions of the Royal Society B: Biological*
Sciences, *365*(1547), 1717–1733. doi:10.1098/rstb.2010.0023
Yan, Y.-B., Duke, N., & Sun, M. (2016). Comparative Analysis of the Pattern of Population
Genetic Diversity in Three Indo-West Pacific *Rhizophora* Mangrove Species. *Frontiers*
in Plant Science, *7*, 1434. doi:10.3389/fpls.2016.01434
Young, M. D., Wakefield, M. J., Smyth, G. K., & Oshlack, A. (2010). Gene ontology analysis
for RNA-seq: accounting for selection bias. *Genome Biology*, *11*(2), R14.
doi:10.1186/gb-2010-11-2-r14
Zhang, C., Zhang, L., Zhang, S., Zhu, S., Wu, P., Chen, Y., … Wu, G. (2015). Global
analysis of gene expression profiles in physic nut (Jatropha curcas L.) seedlings exposed
to drought stress. *BMC Plant Biology*, *15*(1), 1–14. doi:10.1186/s12870-014-0397-x
Zhong, R., Lee, C., Zhou, J., McCarthy, R. L., & Ye, Z.-H. (2008). A Battery of Transcription
Factors Involved in the Regulation of Secondary Cell Wall Biosynthesis in *Arabidopsis*.
The Plant Cell Online, *20*(10), 2763–2782. doi:10.1105/tpc.108.061325

1193
1194
Data accessibility

• Gene expression data and transcriptome sequences that support conclusions have been

deposited in GenBank with the accession code GSE123659;

• SNP genotypes are available at Dryad doi:10.5061/dryad.h11t255 (M. V. Cruz et al.,

2019).

Author contributions

A.P.S., G.M.M. and M.V.C. designed the study. M.V.C. and G.M.M. conducted

fieldwork, prepared samples for sequencing and wrote the manuscript. M.V.C., G.M.M.,

R.S.O., M.D. and D.H.O. analyzed the RNA-Seq and nextRAD results. A.P.S, M.I.Z.,

G.M.M. contributed material/reagents/analytical tools. All authors discussed the results and

contributed to the manuscript.

photosynthetic apparatus

MAP70-5[§]
NAC073[§]

development
KEG¶
GATA8¶
OPF7[§¶]
OBGC[§¶]
MAP70-5[§]