DUX4 regulates oocyte to embryo transition in human

Sanna Vuoristo[1,2#*], Christel Hydén-Granskog[3], Masahito Yoshihara[1], Lisa

Gawriyski[4], Anastassius Damdimopoulos[5], Shruti Bhagat[1,6], Kosuke Hashimoto[6],

Kaarel Krjutškov[1,7,8], Sini Ezer[8,9], Priit Paluoja[7], Karolina Lundin[2], Pauliina

Paloviita[2], Gaëlle Recher[10], Vipin Ranga[11], Tomi Airenne[11], Mahlet Tamirat[11], Eeva-

Mari Jouhilahti[8], Timo Otonkoski[8], Juha S. Tapanainen[2,3,12], Hideya Kawaji[6,13,14],

Yasuhiro Murakawa[6,15], Thomas R. Bürglin[16], Markku Varjosalo[4], Mark S. Johnson[11],

Timo Tuuri[2, 3], Shintaro Katayama[1,9*] and Juha Kere[1,8,9 *]

[1]Karolinska Institutet, Department of Biosciences and Nutrition, Huddinge, Sweden

[2]University of Helsinki, Department of Obstetrics and Gynecology, Helsinki, Finland

[3]Helsinki University Hospital, Department of Obstetrics and Gynecology, Helsinki,

Finland

[4]University of Helsinki, Institute of Biotechnology, Helsinki, Finland

[5]Karolinska Institutet, Bioinformatics and Expression Analysis Core Facility,

Huddinge, Sweden

[6]RIKEN Center for Integrative Medical Sciences, Yokohama, Japan

[7]Competence Centre for Health Technologies, Tartu, Estonia

[8] Stem Cells and Metabolism Research Program, University of Helsinki, Finland

[9]Folkhälsan Research Center, Helsinki, Finland

[10]Institut d'Optique Graduate School, CNRS - Université de Bordeaux, Talence, France

[11]Structural Bioinformatics Laboratory, Biochemistry, Faculty of Science and Engineering, Åbo Akademi University, Turku, Finland

[12]Oulu University Hospital, Oulu, Finland

[13]RIKEN Preventive Medicine and Diagnosis Innovation Program, Wako, Japan

[14]Tokyo Metropolitan Institute of Medical Science, Tokyo, Japan

[15]IFOM, The FIRC Institute of Molecular Oncology, Milan, Italy

[16]University of Basel, Department of Biomedicine, Basel, Switzerland

[#]Current affiliation

[*]Corresponding author

Abstract

In oocyte to embryo transition, the fertilized oocyte undergoes final maturation and the embryo genome is gradually activated during the first three cell divisions. How this transition is coordinated and which factors drive the processes in humans is largely unknown. Here we studied the role of the double homeodomain transcription factor DUX4 in regulating the human oocyte to embryo transition. DUX4 knockdown zygotes show delayed transcriptome reprogramming during the first three days after fertilization. Our combined experimental approaches allowed integrated analysis on the transcriptome, chromatin, and proteome data in human embryos or a DUX4 expressing human embryonic stem cell model. We conclude that DUX4 is a pioneering factor that regulates human oocyte to embryo transition through regulating oocyte mRNA degradation, as well as direct binding and activation of minor genome activation genes, and genomic repeat elements.

Mammalian pre-implantation development involves a series of coordinated processes,

starting with oocyte to embryo transition (OET). OET is a major developmental

reprogramming event from the oocyte to a totipotent embryo, involving fundamental

changes in the epigenetic landscapes, degradation of maternal mRNAs, and

embryonic genome activation (EGA)[1]. In humans, the major EGA takes place by the

8-cell stage[2-4]. Minor EGA genes are upregulated in the human 4-cell embryos where

they subsequently induce genes upregulated at the major EGA[3]. Until now, most

studies focusing on human EGA have concentrated on the genes that are expressed in

the cleavage stage (2-cell, 4-cell, and 8-cell) embryos[3,5] and set the stage for the

forthcoming lineage commitment[6,7]. How OET and EGA are orchestrated in the

human embryos and which genes act as pioneers remain poorly understood.

The conserved double homeodomain transcription factor DUX4 is expressed in early

human embryos[8-10]. It represents a plausible candidate to regulate the OET in humans,

given its capacity to activate EGA-related genes and the genomic repeat elements[8,9].

In this work we have used a combination of methods to investigate DUX4 in the

course of early human development. Our data highlight DUX4 mRNA upregulation

already in the zygotes followed by down-regulation within the next cell division.

Abundant cytoplasmic and nuclear DUX4 protein was apparent only during the first

two days of development. Our in-depth characterisation of DUX4 in the human

embryos suggested that it is not required for survival of the human embryos during

the first three days of development but that DUX4 regulates OET in the human.

siRNA silencing of DUX4 in zygotes delayed oocyte mRNA degradation. Our

comprehensive analysis of transcriptome and chromatin data in human embryos or

DUX4 expressing cell lines suggested that DUX4 opens up chromatin through direct

activation of the retroelements and unannotated genomic regions. Our protein

interaction data demonstrated that DUX4 interacts with transcriptional mediators,

chromatin modifiers, and ubiquitinases expressed in the human oocytes and early

embryos. We conclude that DUX4 regulates different aspects of the OET in humans

by affecting mRNA degradation, transcriptional regulation and chromatin structure.

Results

DUX4 is transiently expressed immediately after fertilization

We first measured DUX4 mRNA levels in human MII oocytes (N=20), zygotes

(N=59), and cleavage embryos (2-cell, N=4; 4-cell, N=14; 8-cell, N=15). We found

significant DUX4 upregulation in the zygotes, while few transcripts were found in the

MII oocytes or the cleavage embryos[10-13] (Fig. 1a). The expression of the DUX is

evolutionary conserved as shown by stage-specific expression in mouse, bovine, and

non-human primates[14]. We stained embryos with a monoclonal antibody targeting the

DUX4 protein and detected cytoplasmic DUX4 in the zygotes and all cleavage

embryos although less in 8-cell embryos (Fig. 1b). DUX4 protein was abundantly

present in the nuclei of the 2- and 4-cell embryos whereas nuclei of the 8-cell

embryos were mostly negative (Fig. 1b). In a single early 8-cell stage embryo there

was high variability in the nuclear DUX4 staining, consistent with a snapshot of on-

going degradation. We quantified nuclear DUX4 intensities in 3D and normalized

them to the cytoplasmic DUX4 intensities and found variable but increasing nuclear

signal from the zygotes up to the 4-cell embryos, while nuclear DUX4 was low or

absent in the 8-cell embryos (Fig 1c). These results demonstrated that DUX4

transcription takes place immediately after fertilisation and is followed by

cytoplasmic and nuclear localisation of the DUX4 protein during the first two days of

human embryo development.

DUX4 protein may form a relatively stable structure even when not bound to DNA

Given predominant DUX4 protein presence in the cytoplasm of the embryos as well

as stage-specific nuclear localization, we analysed the structural features of DUX4

(Fig. 2). Human DUX4 comprises two homeodomains (HD1: residues 19-81; and

HD2: 94-153; UniProt numbering), an intrinsically disordered region (IDR: 154-370)

and a C-terminal domain (residues 371-424; Fig. 2a). A stretch of hydrophobic

residues (308-323) within IDR is conserved in primates and could have a regulatory

role by interacting with other transcription co-regulators or masking the C-terminal

domain from solvent when no interacting partners are present. The C-terminal domain

is predicted to have two structurally ordered and evolutionarily conserved regions:

residues 371-387 and residues 414-423 (Fig. 2a). As a secondary structure the C-

terminal domain is predicted to contain three alpha helices and may form a stable fold

similar to that revealed by NMR for the Pax8 paired box domain (PDB: 2K27;[15]). We

found a nine amino acid transactivation domain (9aaTAD) located at the C-terminal

domain (371-379), also recently reported by Mitsuhashi et al.[16]. This motif is also

present in the PRD-like homeoprotein LEUTX[17] and might be recognized by other

proteins involved in the regulation of transcription, similar to the 9aaTAD motif of

the MLL that interacts with the cAMP-response element binding protein, a

transcriptional co-activator[18].

In the crystal structure of DUX4 (PDB: 6E8C;[19]) HD1 and HD2 bind DNA in a

symmetric manner (Fig. 2b). Residue R21, located at the N-terminal loop of HD1,

interacts with the residues I121, E135 and Q139 of HD2 and, equivalently, R96

located at the N-terminal arm of the HD2 interacts with the residues I46, E60 and Q64

of HD1 (Fig. 2b, c, d). Moreover, the main-chain carbonyl groups of G19 of HD1 and

G94 of HD2 respectively form a hydrogen bond with the main-chain nitrogen atom of

I121 of HD2 and I46 of HD1. We next asked whether these residues are conserved

within the DUX family of homeoproteins and within primates (Fig. 2e): based on our

sequence alignment, residues equivalent to G19, R21, I46, E60, Q64, G94, R96, I121,

E135 and Q139 of DUX4 are highly conserved. Interestingly, the residues G19, R21,

I46, E60, Q64, G94, R96, I121, E135 and Q139 of DUX4 (Fig. 2c,d) are not directly

involved in DNA binding, which prompted us to speculate that these residues could

be important for locking HD1 and HD2 together as a unit before DNA binding, a

hypothesis, which we further tested using molecular dynamics (MD) simulations.

Based on MD simulations and the RMSD among snapshots made every 10 ps (Fig.

2f), the DUX4 HD2-DNA complex appeared the most stable complex (average

RMSD over backbone atoms of 1.73 Å), followed by DUX4-DNA (1.77 Å) and

DUX4 HD1-DNA complex (2.3 Å). The DNA-free DUX4 HD1-HD2 structure (4.0

Å) was the least stabile yet the interactions between HD1 and HD2 were maintained

over the 100 ns simulation. Indeed, ionic interactions between R96 of HD2 and E50

and E60 of HD1 seem to be fundamental for the stability of the double HD structure

of DUX4: the electrostatic interactions/hydrogen bonds between R96 and E60 were

present during 92% of the simulation time. Additional stabilizing interactions between

the two HDs also take place between R21 (HD1) and E135 (HD2), and R22 (HD1)

and E125 (HD2). While these charged interactions hold the two HDs together, the

intermediate linker loop imparts flexibility, which could be vital to accommodate

DNA once DUX4 enters the nucleus and locates its binding motif. Even with bound

DNA, the linker loop fluctuates more relative to HD1 and HD2, as observed from the

RMSF values for the CA atoms of residues of the DUX4-DNA structure (Fig. 2g).

 DUX4 protein interacts with transcriptional mediators and chromatin modifiers

Abundant nuclear and cytoplasmic DUX4 in the human embryos and the modelled

stable structure of the DUX4 when not bound to DNA suggested that DUX4 might

have functions beyond DNA binding. To study this, we applied a recently published

MAC-tag method to identify DUX4 protein-to-protein interactions[20,21]. We identified

altogether 158 BioID and 43 AP-MS high-confidence DUX4 interactions, out of

which 19 appeared in both datasets (FDR < 0.05, corresponding to > 0.73 SAINT

Score). Single BioID interactions and AP-MS interactions together with the

interactions that appeared in both data sets, based on the scored frequency of

interaction with DUX4, are shown in Fig 3a. We concentrated on the DUX4

interacting proteins that scored above the median value (Fig 3b). Overrepresentation

Enrichment Analysis (ORA) of protein pathway markers (Reactome, KEGG) showed

significant enrichment (p < 0.05, FDR < 0.01) of markers linked to generic

transcription and 'RNA Polymerase II Transcription', 'Chromatin organization' and

'Chromatin modifying enzymes'. Comparing our list of genes to protein complex

databases such as ComplexPortal and Corum using Fisher's Exact Test yielded

significant overrepresentation of several variants of the SWI/SNF ATP-dependent

chromatin remodelling complex, Core mediator complex, NSL histone

acetyltransferase complex, SRCAP histone exchanging complex and the NuA4

histone acetyltransferase complex (p < 0.05, FDR 0.01) (Fig. 3b). There were 28

DUX4 interacting proteins classified as RNA binding (GO:0003723) and 19 out of

these were linked to spliceosome and pre-mRNA-splicing. In the protein complex

database Corum, DUX4 interactors were significantly overrepresented in the

Spliceosome, with 10 interactors comprising 7% of the whole complex. In addition,

we found six DUX4 interacting proteins (ZSCAN4, ZSCAN5C, ZSCAN5D, RFPLA,

RFPLB, RNF8, PTOV1, and MB3LB) that have not appeared in the analyses of other

transcription factors. As the protein interaction assay was run in a non-embryonic cell

line (HEK), we next studied which of the identified DUX4 interacting proteins are

expressed by human oocytes or embryos[3,11]. Importantly, the vast majority of the

genes coding for the DUX4 interacting proteins were expressed in oocytes (maternal

genes), embryos, or both. These results suggested that DUX4 could potentially

regulate maternal and embryonic proteins in the cytoplasm and nucleus during the

OET.

Knock-down of DUX4 in human zygotes leads to dysregulation of the maternal

transcriptome

We next asked how DUX4 regulates the OET and early human embryo development.

We microinjected either DUX4 targeting siRNA (siDUX4) or control siRNA

(siCTRL) together with GAP-GFP mRNA to triploid human zygotes and followed

their development for 48 h after the microinjections, until the third day of

development (Fig 4a). 18 h after microinjection, GAP-GFP protein was expressed in

all embryos, confirming successful microinjections. Staining for the DUX4 protein

was very faint or absent in the siDUX4 embryos but strongly positive in the siCTRL

embryos 24 h after microinjection, showing that the DUX4 targeting siRNA

efficiently down-regulated DUX4 (Fig 4b). The cells from the embryos were

collected 48 h after microinjections for single-cell-tagged reverse transcription RNA

sequencing (STRT RNA-seq[22,23]), which detects the transcript far 5'-ends (TFEs).

siDUX4 embryos did not arrest during the experiment by live imaging follow-up, but

their transcriptome was dysregulated in comparison to that of the siCTRL embryos. A

number of transcripts downregulated in the siCTRL appeared enriched in the siDUX4

embryos (Fig. 4c). In order to study how the siDUX4 enriched transcripts typically

behave during the first three days of development, we next compared siDUX4 and

siCTRL embryos to our published gene expression data set[3,10] on human MII oocytes,

zygotes, and cleavage cells. These analyses further confirmed that a large number of

TFEs that remained enriched in the siDUX4 embryos were typically downregulated

between the oocytes and the 4-cell embryos (Fig 4d), the zygotes and the 4-cell

embryos (Fig 4e), and the 4-cell and 8-cell embryos (Fig 4f), indicating delayed

degradation of the maternal transcripts. Gene expression enrichment analysis using

TopAnat[24] for the altogether 91 genes enriched in siDUX4 embryos resulted in terms

such as 'female germ cell' and 'oocyte', in agreement with non-degraded maternal

transcripts. As shown for the siDUX4 enriched gene set, a number of well-known

maternal genes such as *GDF9*, *ZP1*, *ZP2*, *ZP3*, *KHDC3L*, *WEE2*, *NPM3*, *TUBB8*, and

RERE failed to downregulate (Fig. 4g), demonstrating that the OET remained

incomplete after DUX4 abolishment in human zygotes.

DUX4 directly activates the minor EGA genes and a number of unannotated regions

We next generated two TetOn-DUX4 human embryonic stem cell (hESC) lines

expressing DUX4-ires-EmGFP under doxycycline, and studied the effects of DUX4

on the activity and accessibility of the genome (Fig 5a). *DUX4* mRNA (Fig 5b) and

protein (Fig 5c) expression promptly followed doxycycline induction and the known

DUX4 target genes *ZSCAN4* and *TRIM48* followed the induction with slight delay

(Fig 5b). We performed bulk RNA-seq using STRT RNA-seq methods and the data

analysis of the sorted EmGFP+ DUX4 expressing cells showed roughly equal

numbers of up- and downregulated TFEs (Fig 5d). Notably, the majority of the

upregulated TFEs were mapped to unannotated genomic regions, whereas the

downregulated TFEs were mapped to protein coding regions (Fig 5d). The known

target genes of DUX4, i.e., *ZSCAN4*, *LEUTX* and *TRIM48* were significantly

upregulated in the EmGFP+ cells (Fig 5e). Downregulated protein-coding TFEs

included a number of ribosomal genes and genes maintaining pluripotency. This is in

agreement with previous findings showing that DUX4 downregulates some

pluripotency markers[25]. We integrated the data from our RNA-seq and published

DUX4 ChIP-seq analysis[26,27] and found that out of the 32 minor EGA genes induced

in 4-cell embryos[3], 23 were induced in the EmGFP+ cells and 17 out of these

overlapped with DUX4 binding sites. This suggested that DUX4 can induce the

majority of the minor EGA-related genes (Fig 5f). We also identified three previously

unannotated DUX4 targets *KHDC1* pseudogene 1 (FE463525; Fig 5e and 6a), RING-

finger type E3 ubiquitin ligase (FE533694; Fig 5e and 6b), and RING-finger domain

protein 4.2 (FE130507; Fig 5e and 6c) that were induced in 4-cell embryos[3], were

upregulated by DUX4, and overlapped with DUX4 binding sites. We cloned novel

DUX4 target transcripts from a cDNA pool of human day 4 embryos, confirming

their presence in early human cleavage embryos (Fig 6). On the other hand, out of the

129 major EGA genes upregulated in 8-cell human embryos[3], 14 were upregulated by

DUX4 and interestingly, 33 were downregulated (Fig 5f). These data suggested that

DUX4 could upregulate some major EGA genes, but that most of them are likely

activated by the minor EGA genes (such as LEUTX[28]), and that DUX4 might also

negatively regulate major EGA genes. DUX4-induced TFEs were highly enriched

with DUX4 binding sites[26,27] (Fig 5g), and the most highly enriched motif in the

DUX4-induced TFEs was similar to the known DUX4 motif[29] (Fig 5h). Furthermore,

these TFEs were remarkably overrepresented with the DUX4 binding sites among

hundreds of transcription factors (Fig 5i).

Ectopic DUX4 expression causes chromatin opening at ERVL-MaLR elements similar

to 2-cell human embryos

We next integrated STRT RNA-seq and ATAC-seq data from EmGFP +/- sorted

TetOn-Dux4 hESC. DUX4 expression lead to consistent modifications in chromatin

accessibility across all studied clones (Fig 7a). The transcriptionally upregulated TFEs

correlated with more accessible (gained) genomic regions in the DUX4 induced cells

while the correlation was less obvious for the downregulated TFEs and less accessible

(lost) chromatin (Fig 7a). We found that the gained chromatin regions correlated with

upregulated TFEs and lost chromatin regions correlated with downregulated TFEs

(Fig 7b), likely implying that the transcriptional downregulation induced by DUX4

expression is faster than the nucleosome-mediated closing of the chromatin (Fig 7a

and b). We then asked how DUX4 expression modified openness of the different

regions of the chromatin. In general, DUX4 rapidly caused chromatin remodelling,

especially chromatin opening, far from transcription start sites (TSS), demonstrating

that the TSSs seem to be less targeted by the DUX4 expression than other genomic

regions (Fig 7c). We next focused on the ATAC-gained chromatin sites. These

chromatin regions were remarkably enriched with DUX4 binding sites compared with

unchanged regions (P<2.2e-16). Out of the ATAC-gained chromatin sites, 48.9%

overlapped with ERVL-MaLR elements and they were significantly enriched for the

DUX4 binding sites compared with non ERVL-MaLR overlapping sites (55.8% P <

2.2e-16) (Fig. 7d). The ATAC-gained ERVL-MaLR regions remarkably overlapped

with the open chromatin regions found in 2-cell human embryos[30] (Fig. 7e). Out of

the DUX4 induced gained chromatin regions that overlapped with those of the 2-cell

embryos, upregulated by DUX4 induction, and overlapped with DUX4 binding sites,

76.7% were unannotated. Only few protein-coding genes, for instance *ZSCAN4* and

the transcriptional and chromatin regulators *KDM5B (JARID1B)* and *ZNF296* were

included. These results show that DUX4 directly binds ERVL-MaLR elements and

converts the chromatin landscape of the hESCs towards that of the human 2-cell

embryos. These data also suggest that DUX4 largely functions through yet

unannotated genomic regions.

Discussion

The OET, including fertilization and activation of the oocyte to totipotent blastomeres

and subsequent EGA, gradually sets the stage for embryo development[31-33]. How the

OET is orchestrated in human and which factors are the main drivers are still poorly

known. One of the potential candidates driving the OET in human embryos is DUX4.

Our data here show that DUX4 transcripts appear immediately after fertilisation and

are downregulated rapidly following the first and the second cleavage divisions. The

origin of *DUX4* transcripts is still unclear. It could be one of the maternal dormant

RNAs, supported by the fact that DUX4 was absent in the majority of the oocytes but

was significantly induced in the zygotes. Dormant maternal RNAs are stored in the

oocytes as deadenylated transcripts and they are polyadenylated and translated only

after resumption of meiosis or after fertilisation[34]. The increasing nuclear DUX4

protein intensity from zygotes to 4-cell embryos and its disappearance in the nuclei of

8-cell embryos suggested that DUX4 can modify transcriptome and chromatin of the

embryos already before the genome activation takes place. Detailed mechanisms of

DUX4 protein degradation in the 8-cell embryos remain to be further investigated;

however, DUX4 upregulated and interacted with several protein ubiquitinases, such

as TRIM48, a well-known DUX4 target gene. We also identified two previously

unannotated and possibly embryo-specific putative RING-finger type E3 ubiquitin

ligases that were expressed in early human embryos and induced by DUX4. The

putative ubiquitinases RFPLA (RFPL4A) and RFPLB (RFPL4B) regulate protein

degradation in germ cells[35,36]. Another ubiquitin-ligase, RNF114, was recently shown

to be essential for the OET in the mouse[37]. Taken together, our data suggested that

DUX4 induces expression of ubiquitin ligases and also interacts with ubiquitinases,

possibly regulating the presence of DUX4 itself as well as general proteome during

OET.

Knock-down of DUX4 in the human zygotes did not cause mitotic arrest during the 2-

day experiment, in agreement with recent findings on Dux in mouse embryos where a

minority of embryos may proceed until blastocyst stage[8,38,39]. In the mouse Dux-/-

embryos, around 30% of the EGA transcripts that should be upregulated were

downregulated[39], while in the human DUX4 knock-down embryos, many of the

maternal, normally downregulated genes remained unchanged, suggesting that

Dux/DUX4 alone is sufficient for neither the OET nor the EGA. Another candidate

gene regulating OET is *PLAG1*[40]. De novo PLAG1 binding site is found in the EGA

genes in the human embryos, and the phenotype of Plag1+/- mice lacking the

maternal *Plag1* allele show enriched expression of maternal transcripts at the 2-cell

stage, when major EGA occurs in the mouse. The question remains how DUX4,

together with other factors such as PLAG1, coordinates regulation of the maternal and

EGA transcripts in human and which yet unnamed genes might be involved in the

OET in human.

Ectopic expression of DUX4 in the hESC caused opening of the chromatin regions

outside of TSSs, largely at ERVL-MaLR elements. Dux binding at *Mervl* loci drives

chromatin reorganisation at *Mervl* loci in the mouse 2-cell embryo-like cells, and

chromatin organisation during early mouse development is a consequence of the

Mervl integration[41]. Human 2-cell-like cells have not been established by now, but

importantly, in our experiments binding of the DUX4 at ERVL-MaLR elements[14]

could modify chromatin towards embryo-like stage even in the hESCs. Long terminal

repeat elements abundantly present in the genome have been suggested as key

elements contributing to the OET, when major epigenetic and chromatin changes take

place[42]. Our integrated analysis on chromatin openness and transcriptional regulation,

together with the DUX4 ChIP-seq data[26,27] demonstrated that DUX4 regulates several

transcripts and the corresponding genomic loci coding for chromatin modifiers and

epigenetic regulators, as also suggested by Liu et al.[43]. Taken together, our data

indicate that DUX4 alone is not a sufficient inducer for the first three days of human

embryo development but that it regulates the OET by regulating maternal RNA

degradation, EGA genes, and repetitive elements, all of which have been shown to be

crucial for the successful OET in organisms other than human. In addition to

regulating genetic elements by DNA-binding, DUX4 may regulate the proteome by

inducing ubiquitination pathway genes during the OET in human.

 Online methods

Human pre-implantation embryos for single cell RNA-sequencing using STRT method

We analysed single cell RNA-sequencing data from Töhönen et al.[3] for MII oocytes

(N=20), zygotes (N=59), 2-cell (N=4), 4-cell (N=15) and 8-cell (N=14) embryos.

For the DUX4 knockdown experiment 18 siCTRL cells (N=2) and 18 siDUX4 cells

(N=3) were analysed. The embryos were incubated in Ca^{2+}/Mg^{2+} -free culture

medium (Biopsy Medium, Origio) at 37°C heated stage for separation of the cells.

Individuals cells were briefly rinsed in Ca^{2+}/Mg^{2+} -free PBS and placed directly in

lysis buffer (5mM Tris-HCl, pH 7.0 (LifeTechnologies); 5mM DTT (Thermo

Scientific), 0.02 % Tx100 (Fisher Scientific); 0.5 U/µl Ribolock RNAse inhibitor

(ThermoFisher)). The library was prepared according to the published protocol[3,22,44].

The amplified libraries were sequenced on the Illumina HiSeq 2000 instrument.

Bulk RNA-sequencing using STRT method

Total RNA was isolated from the FAC-sorted DUX4-TetOn hESCs using the

RNAqueous Total RNA Isolation Kit (AM1912; ThermoFisherScientific). 20ng of

total RNA from each sample was used for library preparations. The libraries were

prepared using the STRT method as above, with minor modifications. Briefly, RNA

samples were placed in a 48-well plate in which a universal primer, template-

switching oligos, and a well-specific 8-bp barcode sequence (for sample

identification) were added to each well[23,45]. The synthesized cDNAs from the samples

were then pooled into one library and amplified by single-primer PCR with the

universal primer sequence. The resulting amplified library was then sequenced using

the Illumina NextSeq 500 instrument.

Preprocess of raw STRT RNAseq reads

The sequenced STRT raw reads were processed by STRTprep[23], v3dev branch

91a62d2 commit at https://github.com/shka/STRTprep/tree/v3dev. The processed

nonredundant reads were aligned to hg19 human reference genome sequences, ERCC

spike-in sequences and human ribosomal DNA unit (GenBank: U13369) with RefSeq

transcript alignments as a guide of exon junctions. For gene-based statistics, uniquely

mapped reads within (i) the 5'-UTR or the proximal upstream (up to 500 bp) of the

RefSeq protein coding genes, and (ii) within the first 50 bp of spike-in sequences,

were counted. For TFE-based statistics, the mapped reads were assembled according

to the alignments, and uniquely mapped reads within the first exons of the assembled

transcripts were counted, as described in Töhönen et al 2015[3].

Downstream STRT RNA-sequencing data analysis

Differentially expressed genes and TFEs required significantly different tendency on

the expression levels between two groups (q-value < 0.05), and significantly larger

variation than the technical variation (adjusted p-value < 0.05 by BH correction). The

former tendency was tested by the R package SAMstrt v0.99.0[45], and the latter

variation (fluctuation) was estimated based on gene-to-spikein (or TFE-to-spikein)

ratios in the squared coefficient of variation, described in Supplementary Text 1 of

Krjutskov et al. 2016[23]. The minimum value but non-zero was added to all the

normalized read counts and then the counts were divided by the minimum value so

that the logarithm of zero counts become zero. Enrichment analysis of anatomical

terms for the list of upregulated genes by siDUX4 was performed using the TopAnat[24]

(https://bgee.org/?page=top_anat). All human genes in the Bgee database were used

as background. STRT data of human early embryo were obtained from Töhönen et al.

2015 and 2017[3,10] and were overlapped with TFEs using the intersectBed function

from BEDTools[46] (v2.27.1). DUX4 ChIP-seq data was obtained from GSE33838[26]

and scores around the FEs were calculated with computeMatrix and visualized with

plotProfile from deepTools[47] (v3.1.3). Motif enrichment was analyzed using the

command findMotifsGenome.pl from HOMER[48] (v4.10.3) with the option "-size -

300,100". Enrichment analysis with publicly available ChIP-seq datasets was

conducted with ChIP-Atlas[49] (http://chip-atlas.org). A total of 7,216 human

transcription factor ChIP-seq datasets which had more than 500 peaks were analyzed.

Fold enrichment was calculated as (the number of ChIP-seq peaks overlapping with

upregulated TFEs / the number of upregulated TFEs) / (the number of ChIP-seq peaks

overlapping with all TFEs / the number of all TFEs). *P*-values were calculated with

Fisher's exact test and *Q*-values were calculated with the Benjamini & Hochberg

method. After excluding the TFEs annotated on ribosomal DNA, 6,425 upregulated

TFEs were used as foreground and 109,624 all the detected TFEs were used as

background both in the motif and ChIP-seq enrichment analysis.

Human ESC culture

hESC lines H1 (WA01) and H9 (WA09) were purchased from WiCell. The hESCs

were maintained on Geltrex-coated tissue culture dishes in Essential 8 culture medium

and passaged every three to five days by incubation with 0.5 mM EDTA (all from

Thermo Fisher Scientific).

Plasmid construction

The full-length DUX4 (NM_001293798.2) was synthesized and cloned between the

SalI and BamHI sites of the pB-tight-hMAFA-ires-EmGFP-pA-PGK-Puro vector (a

kind gift from Diego Balboa, Biomedicum Stem Cell Centre) at GenScript (Genscript,

NJ, USA).

Doxycycline-inducible DUX4 expressing human ESCs

The hESCs were incubated with StemPro Accutase (Thermo Fisher Scientific) until

the edges of the colonies started to curl up. The Accutase was aspirated and the cells

were gently detached in cold 5% FBS (Thermo Fisher Scientific) -PBS (Corning) and

counted. One million cells were centrifuged at 800rpm for 5 min and the pellet was

transferred into 120 μl of R-buffer containing 1 μg of pB-tight-DUX4-ires-EmGFP-

pA-PGK-Puro, 0.5 μg of pBASE and 0.5 μg of rtTA-M2-IN plasmids. 100 μl of the

cell-plasmid suspension was electroporated with two pulses of 1100 V, 20 ms pulse

width, using Neon Transfection system. The electroporated cells were plated on

Geltrex-coated dishes in Essential 8 medium with 10 μM ROCK inhibitor Y27632

(Peprotech). Fresh Essential 8 medium without ROCK-inhibitor was changed to the

cells on the day following the electroporation. The cells were selected with

Puromycin at 0.3 μg/mL. The TetOn-DUX4 hESC clones were picked manually on

Geltrex-coated 96-well plates, expanded and selected again with Puromycin.

Appearance of the EmGFP reporter protein was tested using Doxycycline at

concentrations ranging from 0.2 μg/ml to 1.0 μg/ml and detected using EVOS FL Cell

imaging system (Thermo Fisher Scientific). When indicated for the experiments

presented in this paper, the EmGFP+ DUX4 expressing hESC clones had been treated

with 1μg/ml of Doxicycline for 1, 2, 3, (qPCR) or 4 hours prior to downstream

analyses.

cDNA cloning of unannotated DUX4 targets

Single human 4-cell embryo cDNA library was prepared according to the protocol by

Tang et al.[50] and used for cloning of the putative transcripts. The transcripts were

amplified using Phusion High-Fidelity DNA polymerase (New England Biolabs)

according to manufacturer's instructions. Predicted KHDC1 pseudo gene 1, putative

RING-finger type E3 ubiquitin ligase and putative RING-finger domain protein

encoding genes were amplified using touchdown PCR: 98˚C for 30 s; 24 cycles of

98˚C for 10 s, annealing for 30 s, temperature decreasing from 63˚C to 56˚C, 1 ˚C/3

cycles, 72˚C for 30 s; 16 cycles of 98 ˚C for 10 s, 55˚C for 30 s, 72˚C for 30 s; final

extension 72˚C for 10 min. All PCR products were cloned into pCR4Blunt-TOPO

vector using Zero Blunt TOPO PCR Cloning kit (Invitrogen) and sequences were

verified by Sanger sequencing (Eurofins Genomics). Clone sequences are available

from the ENA browser at http://www.ebi.ac.uk/ena/data/view/LR694082-LR694089.

Bioinformatics analysis and molecular dynamics simulations of the DUX4 protein

The sequence for the human DUX4 (Q9UBX2) protein was obtained from the

UniProt database (The UniProt Consortium[30]), whereas other sequences were

retrieved from the non-redundant database of NCBI using blastp[51] and with human

DUX4 as the query sequence. Multiple sequence alignment was carried out using

MAFFT[52]. Secondary structures, solvent accessibility and disordered regions were

predicted using POLYVIEW-2D[53], SABLE[54], SCRATCH[55] and RaptorX-Property[56].

The 9aaTAD web server[57] was used to predict 9aaTAD motifs. The 2.12 Å resolution

crystal structure of the DUX4 HD1-linker-HD2 fragment bound to DNA[19] (PDB:

6E8C) was obtained from the Protein Data Bank[58]. PyMOL (version 1.6; Schrödinger

LLC) and Bodil[59] were used to analyze inter-HD interactions. Based on the DUX4

structure, molecular dynamics (MD) simulations, over all atoms, were used to explore

the dynamic states of the (1) double HD complex with (HD1-HD2 + DNA) and (2)

without (HD1-HD2) bound DNA and the individual HDs with bound DNA: (3) HD1

+ DNA and (4) HD2 + DNA. MD simulations of these four structures were carried

out with the AMBER package[60] (version 18) using the ff14SB[61] (for protein) and

OL15[62] (for DNA) force fields. The structures were solvated with explicit TIP3P

water molecules[63] within an octahedral box ensuring a 12 Å distance between the

boundaries of the simulation box and solute atoms. Sodium counter ions were added

to neutralize the system and additional Na^+/Cl^- ions were added to bring the salt

concentration to 150 mM. Periodic boundary conditions were employed and the

particle-mesh Ewald algorithm[64] was applied to electrostatic interactions with a

distance cutoff of 9 Å.

Prior to the production simulation, 5000 cycles of steepest descent and conjugate

gradient energy minimization were carried out on each system, initiated by

introducing a 25 kcal mol^{-1} $Å^{-2}$ restraint on solute atoms that was gradually reduced

to 0 kcal mol^{-1} $Å^{-2}$ over the total minimization. The systems were then heated from

100 K to 300 K during 100 ps with a 10 kcal mol^{-1} $Å^{-2}$ restraint on solute atoms,

followed by a 900 ps equilibration at constant pressure while systematically reducing

the restraint to 0.1 kcal mol^{-1} $Å^{-2}$. The equilibration protocol was finalized with a

restraint-free 5 ns simulation. The production simulation was performed for 100 ns at

constant temperature (300 K) and pressure (1 bar), which was maintained using the

Berendsen algorithm[65] with 5 ps coupling constant. Trajectories were saved every 10

ps and the resulting structural snapshots were analyzed further by calculating the root-

mean-square deviations (RMSD; over backbone atoms) and root-mean-square

fluctuations (RMSF; over Cα atoms), as well monitoring hydrogen bond interactions

using the programs CPPTRAJ[66] and VMD[67].

Affinity purification of protein complexes, mass spectrometry and data analysis

Cell Culture and Affinity Purification

Cloning of DUX4 to MAC-tag Gateway® destination vector

DUX4 was first amplified in two-step PCR reaction from pB-tight-DUX4-ires-

EmGFP-pA-PGK-Puro and cloned to Gateway compatible entry clone using Gateway

BP Clonase II (Invitrogen) according to manufacturer's instructions. The entry clone

was further cloned to Gateway compatible destination vectors containing the C-

terminal and N-terminal tags as described[21]. Transfection and selection of the T-Rex

293 cells (Invitrogen, Life Technologies, R78007, cultured in manufacturer's

recommended conditions) and affinity purification of the final product was done as

previously[21].

Liquid Chromatography-Mass Spectrometry

Analysis was performed on a Q-Exactive mass spectrometer with an EASY-nLC 1000

system via an electrospray ionization sprayer (Thermo Fisher Scientific), using

Xcalibur version 3.0.63. Peptides were eluted from the sample with a C18 precolumn

(Acclaim PepMap 100, 75 μm x 2 cm, 3 μm, 100 Å; Thermo Scientific) and

analytical column (Acclaim PepMap RSLC, 65 μm x 15 cm, 2 μm, 100 Å; Thermo

Scientific), using a 60 minute buffer gradient ranging from 5 to 35% Buffer B, then a

5 min gradient from 35 to 80% Buffer B and 10 minute gradient from 80 to 100%

Buffer B (0.1% formic acid in 98% acetonitrile and 2% HPLC grade water). 4 μl of

peptide sample was loaded by a cooled autosampler. Data-dependent FTMS

acquisition was in positive ion mode for 80 min. A full scan (200-2000 m/z) was

performed with a resolution of 70,000 followed by top10 CID-MS^2 ion trap scans with

a resolution of 17,500. Dynamic exclusion was set for 30 s. Database search was

performed with Proteome Discoverer 1.4 (Thermo Scientific) using the SEQUEST

search engine on the Reviewed human proteome in UniProtKB/SwissProt databases

(http://www.uniprot.org, downloaded Nov. 2018). Trypsin was selected as the

cleavage enzyme and maximum of 2 missed cleavages were permitted, precursor

mass tolerance at ±15 ppm and fragment mass tolerance at 0.05 Da.

Carbamidomethylation of cysteine was defined as a static modification. Oxidation of

methionine and biotinylation of lysine and N-termini were set as variable

modifications. All reported data were based on high-confidence peptides assigned in

Proteome Discoverer (FDR < 0.01).

Identification of statistical confidence of interactions

Significance Analysis of INTeractome (SAINT[68]) -express version 3.6.3 and

Contaminant Repository for Affinity Purification (CRAPome,

http://www.crapome.org) were used to discover statistically significant interactions

from the AP-MS data[69]. The DUX4 LC-MS data was ran alongside a large dataset of

other transcription factors, as well as a large GFP control set. Final results represent

proteins with a SAINT score higher than 0.73, and present in all four replicates.

Overrepresentation Analysis

Overrepresentation analysis of statistically significant interactions in Gene Ontology

and Reactome was done in WebGestalt[70], and overrepresentation of prey proteins in

ComplexPortal[71]   (https://www.ebi.ac.uk/complexportal)   and   CORUM[72]

(https://mips.helmholtz-muenchen.de/corum/) was done using Fisher's exact test and

multiple testing correction in an in-house R-script.

Interaction network

Protein interaction networks were constructed from filtered SAINT data that was

imported to Cytoscape 3.6.0. Known prey-prey interactions were obtained from the

iRef database (http://irefindex.org).

RNA isolation, reverse transcription and quantitative real-time quantitative PCR

Total RNA was isolated by NucleoSpin RNA kit (Macherey Nagel). 1μg of RNA was

reverse transcribed by MMLV-RTase with oligodT, dNTPs, and Ribolock in MMLV-

RTase buffer (Thermo Fisher Scientific). 5X HOT FirelPol qPCR Mix (Solis

Biodyne) was used to measure relative mRNA levels with Lightcycler (Roche). Δ Δ

CT method was followed to quantify relative gene expression where *CYCLOPHILIN*

G was used as endogenous control. Relative expression of each gene was normalized

to the expression without doxycycline treatment.

Fluorescence associated cell sorting

TetOn-DUX4 hESCs were treated with TrypLE for 5 min and suspended into cold

FACS buffer (5% FBS-PBS). Single cell suspension was filtered through 40μm Cell

strainers and centrifuged at 800 rpm for 5min. The cell pellets were suspended in cold

FACS buffer and placed on ice. EmGFP- and EmGFP+ cells were separated to FACS

buffer by Sony SH800Z Cell Sorter with blue laser (488) and 100 μm nozzle.

ATAC-sequencing library preparation and data analysis

In principle the ATAC-sequencing libraries were prepared as in[73]. 5x10^4 EmGFP-

negative and EmGFP-positive TetOn-hESCs for four biological samples; TetOn-

DUX4 in H1 clone 2, H1 clone 8, H9 clone 3 and H9 clone 4, were centrifuged at

500g for 5 min. The pellets were washed in cold 1X PBS by centrifugation at 500g for

5min. Each cell pellet was lysed in 50 μl of cold lysis buffer (10 mM Tris-HCl, pH

7.4; 10 mM NaCl, 3 mM MgCl$_2$, and 0.1% IGEPAL CA-630) and centrifuged at 500g

at 4°C for 10 min. The pellet was then resuspended in the transposase reaction mix

(2.5 μl of transposase in TD buffer) and incubated at 37°C for 30min. The reactions

were purified through columns and eluted in 20 μl. After addition of the barcode

oligos the DNA samples were amplified for 12 cycles (98°C for 10 seconds, 63°C for

30 seconds and 72°C for 60 seconds) in Phusion PCR master mix (Thermo Fisher

Scientific). The PCR products were purified through the columns and eluted in 20 μl.

ATAC-seq data analysis

Bcl files were converted and demultiplexed to fastq using the bcl2fastq program.

STAR[74] was used to index the human reference genome (hg19), obtained from

UCSC, and align the resulting fastq files. The resulting bam files with the mapped

reads were then converted to tag directories with subsequent peaks calling using the

HOMER suit of programs[48]. HOMER was also employed for counting the reads in the

identified peak regions. The raw tag counts from the peaks were then imported to

R/Bioconductor and differential peak analysis was performed using the edgeR

package and its general linear models pipeline. Peaks with an FDR adjusted p value

under 0.05 were termed significant. Plotting was done in R using packages Complex

heatmap, ggplot2 and ggbeeswarm. RepeatMasker table downloaded from UCSC

(http://hgdownload.soe.ucsc.edu/goldenPath/hg19/database/rmsk.txt.gz)   was

converted to BED format and then intersected with the ATAC-seq peaks using the

intersectBed from BEDTools[46] to determine the peaks overlapped with ERVL-MaLR

elements. ATAC-seq data of human early embryo were obtained from GSE101571[30],

and scores around the ATAC-seq peaks were calculated with computeMatrix and

visualized with plotHeatmap from deepTools[47]. All the scripts and command line

options can be provided upon request.

Immunocytochemistry of the human ESC

The cells were fixed with 3.8% PFA, washed three times, permeabilised in 0.5% (v/v)

Tx100-PBS for 7 min, and washed with washing buffer (0.1% (v/v) Tween20-PBS).

The samples were incubated with ProteinBlock (Thermo Fisher Scientifi) at room

temperature for 10 min to prevent unspecific binding of primary antibody. Primary

antibody (rabbit MAb anti DUX4, clone E5-5, 1:400; Abcam) was diluted as

indicated in washing buffer and incubated at 4°C overnight. After washings,

fluorescence-conjugated secondary antibody (anti rabbit 594, A-21207; Thermo

Fisher Scientific) was diluted 1:1000 in washing buffer and incubated at room

temperature for 20 min. Nuclei were counterstained with DAPI 1:1000 in washing

buffer. The images were captured with Evos FL Cell Imaging system with 10X and

20X Plan Achromatic objectives.

Immunocytochemistry of the human embryos

The embryos were fixed in 3.8 % PFA at room temperature for 15min, washed three

times in the washing buffer (above), and permeabilised in 0.5 % Tx100-PBS at room

temperature for 15 min. Unspecific primary antibody binding was blocked as above.

DUX4 (as above) was incubated at 4°C overnight. The embryos were washed and

incubated in the secondary antibody (anti-rabbit 488, A-21202; Thermo Fisher

Scientific) diluted 1:500 in washing buffer (as above) at room temperature for 2

hours. After washings, nuclei were counterstained with DAPI 1:500 in washing

buffer.

Imaging of the fixed human embryos

The embryos were imaged in washing buffer on Ibidi 8-well μ slides with Leica TCS

SP8 confocal laser scanning microscope (Leica Microsystems, Mannheim, Germany)

using Leica HC PL APO CS2 40X/1.10NA and Leica HC PL APO CS2 63X/1.20NA

water objectives.

Confocal microscopy image analysis

Confocal images were processed using Fiji (http://fiji.sc). For the data presented in

the Fig 1b, images were smoothened by Gaussian filter (radius=1 pixel kernel). For

the quantification of the DUX4 intensity in the nucleus (Fig 1c), the DAPI channel

was denoised using rolling ball (radius=100). The images were smoothened in 3D

using Gaussian filter (radius=2 pixel kernel) and cell nuclei were segmented. The

segmented regions were used to measure average pixel intensity per nucleus in each

cell in the DUX4 channel. DUX4 intensity in the nucleus was normalized to intensity

of the corresponding cytoplasmic DUX4 staining in the single representative plane.

Culture and microinjection of human embryos

Human triploid zygotes were warmed using Gems Warming Set (Genea Biomedx)

and cultured in G-TL medium (Vitrolife) in 6 %O_2 and 6 % CO_2 at 37°C. 12 μl of

either 20 μM scrambled control siRNA (AM4611, Thermo Fisher Scientific) or

DUX4-targeting siRNA (cat. 4457308, Thermo Fisher Scientific) diluted in

nucleotide-free H_2O were mixed with total of 500 ng of GAP-GFP mRNA and

centrifuged at maximum speed at 4°C for 10 min. The embryos were microinjected

using Eppendorf microinjector and placed in G-TL medium in Geri dish for 3D time-

lapse imaging (Geri incubator, Genea Biomedx, Australia).

Human embryo live imaging

Imaging of the human triploid embryos was initiated immediately after

microinjections (Geri incubator). Images were captured in 3D every 15 minutes until

the embryos were removed for fluorescence staining or termination of the experiment.

Ethical approvals

Collection and experiments on human oocytes and embryos were approved by the

Helsinki University Hospital ethical committee, diary numbers 308/13/03/03/2015

and HUS/1069/2016. Human surplus oocytes, zygotes, and embryos were donated by

couples that had undergone infertility treatments at Helsinki University Hospital

Reproduction Medicine Unit. The donations were done with an informed consent and

patients understood that donating oocytes, zygotes, or embryos is voluntary.

Acknowledgements

We are grateful for all the couples that donated their surplus oocytes, zygotes, or

embryos for this project. We thank the IVF nurses at Helsinki University Hospital

Reproductive Medicine Unit for recruiting the couples for oocyte and embryo

donation program, Dr. Diego Balboa-Alonso for the pB-ires-EmGFP-pA-PGK-Puro

plasmid and discussions of the data and Dr. Jere Weltner for helpful discussions. We

acknowledge Biomedicum Imaging Unit (Helsinki), Functional Genomics Unit

(Helsinki), Biomedicum Flow Cytometry Unit (Helsinki), Bioinformatics and

Expression Analysis Core Facility (Stockholm) for skilled technical assistance. We

thank Dr. Jukka Lehtonen for scientific IT support (Biocenter Finland) and CSC IT

Center for Science for supercomputing. We thank prof. Outi Hovatta for inspiration

and introducing the senior author to the field of early embryogenesis. This study is

based on collaboration between the groups of the last seven authors. This project was

supported by Jane and Aatos Erkko foundation, Sigrid Jusélius foundation, Finnish

Academy, and Helsinki University Hospital funds. SV was supported by Jane and

Aatos Erkko foundation. MY was supported by the Scandinavia-Japan Sasakawa

Foundation, the Japan Eye Bank Association, the Astellas Foundation for Research on

Metabolic Disorders, and the Japan Society for the Promotion of Science Overseas

Research Fellowships. MT was supported by Joe, Pentti and Tor Memorial

Foundation, Graduate School of Åbo Akademi University. VR was supported by

Foundation of Åbo Akademi University and Magnus Ehrnrooth Foundation.

Contributions

SV, SK and JK conceived and coordinated the study. YM, TRB, MV, MSJ, TT, SK

and JK supervised the work in each contributing laboratory. Every author participated

in either planning or conducting respective experiments and analyzing or interpreting

the data. SV, MY, LG, VR, TA, MT, MSJ, JK wrote the manuscript. All authors

approved the final version of the manuscript.

Competing interests

The authors declare no competing interests.

Figure Legends

Figure 1. DUX4 expression in human embryos.

(a) Log_2 RPM of DUX4 mRNA reads in human oocytes (N=20), zygotes (N=59), 2-
cell (N=4), 4-cell (N=14), and 8-cell (N=15) embryos.

(b) Representative confocal images of zygotes (N=3), 2-cell (N=3), 4-cell (N=4), and
8-cell (N=2) human embryos stained with monoclonal DUX4 antibody E5-5 (green).
Nuclei counterstained with DAPI (magenta).

(c) 3D quantification of DUX4 intensity in the nuclei of the human embryos,
normalised to cytoplasmic DUX4 staining (single plane) with standard deviation.

Figure 2. Structural features of DUX4.

(a) Domain structure of full-length DUX4: N- and C-terminal amino acid residues, as
well as boundary residues for the homeodomains HD1 and HD2, three predicted
ordered C-terminal regions (disorder value < 0.5; red curve) and 9aaTAD motif
(blue). Conservation of residues in primates versus the human sequence C-terminal to
residue G153 (green curve) and sequence alignment and predicted secondary
structures (alpha helices) of the ordered regions. Residue numbering from UniProt ID
Q9UBX2.

(b) Crystal structure (PDB:6E8C; Lee et al., 2018) of DUX4 HD1 (blue) and HD2
(gold) in complex with the consensus DNA motif "TAATCTAATCA" (grey).
Disordered/linker regions, magenta; residues forming inter-homeodomain contacts
drawn as sticks.

(c) Structure of DUX4 HD1 (blue) and 2 (gold) without DNA. The coordinates of the

bound DNA of the X-ray structure shown in (b) were deleted from the coordinate file

of DUX4 structure. Coloring as in (b).

(d) View focused on the inter-homeodomain interactions shown in (c). Hydrogen

bonds, yellow dash lines; coloring as in (b).

(e) Sequence comparison of residues forming inter-homeodomain contacts.

(f) Root-mean-squared fluctuations (RMSF) of the C α atoms of the X-ray structure

of DUX4 with (red curve) and without (blue curve) bound DNA during a 100 ns MD

simulation. HD1 (blue), linker (magenta) and HD2 (gold).

(g) Root-mean-squared deviation (RMSD) of the backbone atoms of the X-ray

structure of DUX4 with and without bound DNA during a 100 ns MD simulation.

Homedomains with (HD1-HD2 +DNA) and without (HD1-HD2) bound DNA and

single homedomains (HD1+DNA and HD2 +DNA) with bound DNA were used in

the simulations.

Figure 3. The protein – protein interaction network of DUX4.

(a) High confidence protein-protein interactions detected by AP-MS (n=24) and

BioID (n=139) -methods (SAINT score > 0.74). Average spectral count of the

interaction filtered to show interactions larger than the median (AP-MS=4.125,

BioID=7.5).

(b) DUX4 interactome, filtered to show spectral counts larger than median. BioID -

interactions shown in red lines and AP-MS -interactions in blue, if protein appeared in

both data sets it is outlined in bold black. Known prey-prey interactions shown in grey

(iREF).

Figure 4. Knockdown of DUX4 in human embryos.

(a) Schematic of the experimental set up.

(b) Human embryos immunostained with DUX4 antibody (green) 24 h after microinjection with either control siRNA (left panel) or DUX4 targeting siRNA (right panel). Nuclei counterstained with DAPI (blue). Overlay of a single representative z plane and the corresponding z-planes shown for DUX4 staining, nucleus and bright field channels on the right side of each overlay. Scale bar 50 μm. **(c)** Scatter plot of the expression levels of TFEs across the siCTRL and siDUX4 embryos. Red dots represent significantly upregulated TFEs by siDUX4 and grey crosses represent TFEs, which showed no significant change. **(d-f)** TFEs upregulated (red dots) or showing no significant change (grey crosses) by siDUX4 in the human OET transcriptome as in[3]. Comparisons in d: oocyte to 4-cell, e: zygote to 4-cell, and f: 4-cell to 8-cell. The dotted line marks the cell division effect on cellular RNA content. *P*-values were calculated with Fisher's exact test for the frequency of the siDUX4-upregulated TFEs of the TFEs normally downregulated during respective stages.

(g) Expression levels of the oocyte-specific genes in siCTRL and siDUX4 embryos. Asterisks represent statistical significance (*q*-value < 0.05). Horizontal lines represent the median values in each group.

Figure 5. Transcriptome and ChIP-seq analysis on the TetOn-DUX4 hESCs.
(a) Schematic of the experimental set up.

(b) Doxicycline induction of TetOn-DUX4 hESCs induces expression of DUX4 and

DUX4 target genes, ZSCAN4 and TRIM48. Shown for H1-TetOn-DUX4 clone 2. All

clones selected for experiments followed the same trend. X-axis indicates time (h)

incubated in doxycycline. Relative mRNA expression levels were normalized to the

non-induced cells.

(c) 4-hour doxycycline induction upregulates DUX4 protein expression in the

nucleus, shown for H1-TetOn-DUX4 clone 2.

(d) Proportion of the upregulated and downregulated TFEs based on the genome

annotation.

(e) Expression level of putative DUX4 target genes. Asterisks represent statistical

significance (q-value < 0.05). Horizontal lines represent the median values in each

group.

(f) Proportion of the upregulated (Up), downregulated (Down), and non-significantly

changed (NS) TFEs by DUX4 induction among the minor (Oocyte to 4-cell embryo)

and major (4- to 8-cell embryo) EGA genes. One TFE out of the 129 major EGA

genes annotated on unassigned chromosome (ChrUn) and was excluded from the

analysis.

(g) DUX4 ChIP-seq intensity[26] around the peaks of reads within the upregulated TFEs

(blue) and all the detected TFEs (green).

(h) De novo motif enrichment analysis of the DUX4-induced TFEs. Top: the most

significantly enriched motif (P = 1e-961). Bottom: the best-matched known motif

(DUX4 ChIP-seq of myoblasts: GSE75791[29]; matched score = 0.92).

(i) Enrichment analysis of the DUX4-induced TFEs with publicly available ChIP-seq

datasets. A total of 7,216 ChIP-seq data for transcription factors are shown. ChIP-seq

data for DUX4 are shown in red. Dots on the left side of the dashed line are

underrepresented, whereas dots on the right side are overrepresented.

Figure 6. Novel DUX4 targets.

(a) Predicted *KHDC1* pseudogene 1 (clone K5.2), at chromosome 6 (73,918,461-

73,920,115) was expressed by the human 4-cell embryos (FE463525) and upregulated

in the TetOn -DUX4- hESCs (TFE93242). TFEs overlapped with DUX4 binding sites

(DUX4 ChIP). cDNA clone K5.2 (thick orange regions indicate exons and grey thin

regions indicate introns) corresponded to the *KHDC1* pseudogene 1 transcript

assembly in TetOn-DUX4 cells. Transcript assemblies (mRNA Genbank and human

ESTs), including unspliced, are shown.

(b) Putative RING-finger type E3 ubiquitin ligase at chromosome 2 (108,273,771-

108,277,850) was expressed by the human 4-cell embryos (FE130507) and it was

upregulated in the TetOn-DUX4 hESCs (TFE25640). DUX4 ChIP-seq peak

overlapped with the TFEs. RET11.1 was cloned from human 4-cell embryo (clone

RET11.1). Thick blue regions indicate exons and thin grey regions indicate introns.

Transcript assemblies (mRNA Genbank and human ESTs), including unspliced, are

shown.

(c) Putative RING-finger domain protein at chromosome 8 (210,701-215,100) was

expressed by the human 4-cell embryos (TFE533694) and induced by TetOn-DUX4

hESCs (TFE102707). ChIP-seq overlapped with the TFEs. Two cDNA clones, Ring

4.2 and Ring 10.22, were expressed in the human 4-cell embyos. Thick blue regions

indicate exons and grey thin regions indicate introns. Transcript assemblies (mRNA

Genbank and human ESTs), including unspliced, are shown.

Figure 7. Integrated analysis of the DUX4 induced changes in the chromatin and transcriptome of the hESCs.

(a) Heatmap of the 4,686 ATAC-sequencing reads across all samples at ATAC-seq peaks that overlap with differentially regulated TFE reads. Counts for each peak were standardized across each sample (mean=0, sd=1). Samples and peaks were then clustered using hierarchical clustering. The Separate heatmap of the ATAC-seq shows if the changes in the heatmap are significant (red: ATAC-reg gained; blue: ATAC-reg lost; grey: ATAC-reg non-significant). TFE-Reg heatmap shows if the overlapping TFE site is upregulated or downregulated (red: upregulated; blue: downregulated).

(b) Quasi random plot showing the results of the differential peak analysis on the ATAC-seq and STRT-RNA-seq. Each point is an ATAC-seq peak. Analyses were carried out on peaks that were repeated at least three times. Red: ATAC-seq gained; blue: ATAC-seq lost; grey: ATAC-seq non-significant. Y-axis: The log fold change of the ATAC-seq reads in the DUX4 expressing versus the control samples. X-axis: The ATAC-seq peaks overlapping either with the down-regulated TFEs or upregulated TFEs.

(c) Density plot showing distribution of the ATAC-seq peaks relative to the TSS of genes separated by how the peak is regulated by the DUX4-expression. red: ATAC-reg gained; blue: ATAC-reg lost; grey: ATAC-reg non-significant.

(d) Proportion of the peaks overlapped with ERVL-MaLR elements in the gained, non-significant and lost ATAC-reg peaks (pink) by DUX4 induction. Inset pie charts inidicate the proportion of the ATAC-gained peaks overlapping with DUX4 binding sites (green).

(e) ATAC-reg intensity of human early embryo[30] around the gained, non-significant

(NS), and lost ATAC-reg peaks by DUX4 induction which overlap with ERVL-

MaLR elements.

References

1   Jukam, D., Shariati, S. A. M. & Skotheim, J. M. Zygotic Genome Activation
in Vertebrates. *Developmental cell* **42**, 316-332,
doi:10.1016/j.devcel.2017.07.026 (2017).

2   Braude, P., Bolton, V. & Moore, S. Human gene expression first occurs
between the four- and eight-cell stages of preimplantation development.
Nature **332**, 459-461, doi:10.1038/332459a0 (1988).

3   Tohonen, V. *et al.* Novel PRD-like homeodomain transcription factors and
retrotransposon elements in early human development. *Nature*
communications **6**, 8207, doi:10.1038/ncomms9207 (2015).

4   Blakeley, P. *et al.* Defining the three cell lineages of the human blastocyst by
single-cell RNA-seq. *Development* **142**, 3613, doi:10.1242/dev.131235
(2015).

5   Galan, A. *et al.* Functional genomics of 5- to 8-cell stage human embryos by
blastomere single-cell cDNA analysis. *PloS one* **5**, e13615,
doi:10.1371/journal.pone.0013615 (2010).

6   Stirparo, G. G. *et al.* Integrated analysis of single-cell embryo data yields a
unified transcriptome signature for the human pre-implantation epiblast.
Development **145**, doi:10.1242/dev.158501 (2018).

7   Boroviak, T. *et al.* Single cell transcriptome analysis of human, marmoset and
mouse embryos reveals common and divergent features of preimplantation
development. *Development* **145**, doi:10.1242/dev.167833 (2018).

8   De Iaco, A. *et al.* DUX-family transcription factors regulate zygotic genome
activation in placental mammals. *Nat Genet* **49**, 941-945, doi:10.1038/ng.3858
(2017).

9   Hendrickson, P. G. *et al.* Conserved roles of mouse DUX and human DUX4 in
activating cleavage-stage genes and MERVL/HERVL retrotransposons.
Nature genetics **49**, 925-934, doi:10.1038/ng.3844 (2017).

10   Tohonen, V. *et al.* Transcriptional activation of early human development
suggests DUX4 as an embryonic regulator. doi:https://doi.org/10.1101/123208
(2017).

11   Yan, L. *et al.* Single-cell RNA-Seq profiling of human preimplantation
embryos and embryonic stem cells. *Nature structural & molecular biology* **20**,
1131-1139, doi:10.1038/nsmb.2660 (2013).

12   Dang, Y. *et al.* Tracing the expression of circular RNAs in human pre-
implantation embryos. *Genome biology* **17**, 130, doi:10.1186/s13059-016-
0991-3 (2016).

13   Xue, Z. *et al.* Genetic programs in human and mouse early embryos revealed
by single-cell RNA sequencing. *Nature* **500**, 593-597,
doi:10.1038/nature12364 (2013).

14   Whiddon, J. L., Langford, A. T., Wong, C. J., Zhong, J. W. & Tapscott, S. J.
Conservation and innovation in the DUX4-family gene network. *Nat Genet*
49, 935-940, doi:10.1038/ng.3846 (2017).

15   Codutti, L. *et al.* The solution structure of DNA-free Pax-8 paired box domain
accounts for redox regulation of transcriptional activity in the pax protein
family. *The Journal of biological chemistry* **283**, 33321-33328,
doi:10.1074/jbc.M805717200 (2008).

16   Mitsuhashi, H. *et al.* Functional domains of the FSHD-associated DUX4
protein. *Biology open* **7**, doi:10.1242/bio.033977 (2018).

17   Katayama, S. *et al.* Phylogenetic and mutational analyses of human LEUTX, a
homeobox gene implicated in embryogenesis. *Scientific reports* **8**, 17421,
doi:10.1038/s41598-018-35547-5 (2018).

18   Goto, N. K., Zor, T., Martinez-Yamout, M., Dyson, H. J. & Wright, P. E.
Cooperativity in transcription factor binding to the coactivator CREB-binding
protein (CBP). The mixed lineage leukemia protein (MLL) activation domain
binds to an allosteric site on the KIX domain. *The Journal of biological*
chemistry **277**, 43168-43174, doi:10.1074/jbc.M207660200 (2002).

19   Lee, J. K. *et al.* Crystal Structure of the Double Homeodomain of DUX4 in
Complex with DNA. *Cell reports* **25**, 2955-2962 e2953,
doi:10.1016/j.celrep.2018.11.060 (2018).

20   Varjosalo, M. *et al.* Interlaboratory reproducibility of large-scale human
protein-complex analysis by standardized AP-MS. *Nature methods* **10**, 307-
314, doi:10.1038/nmeth.2400 (2013).

21   Liu, X. *et al.* An AP-MS- and BioID-compatible MAC-tag enables
comprehensive mapping of protein interactions and subcellular localizations.
Nature communications **9**, 1188, doi:10.1038/s41467-018-03523-2 (2018).

22   Islam, S. *et al.* Highly multiplexed and strand-specific single-cell RNA 5' end
sequencing. *Nature protocols* **7**, 813-828, doi:10.1038/nprot.2012.022 (2012).

23   Krjutskov, K. *et al.* Single-cell transcriptome analysis of endometrial tissue.
Human reproduction **31**, 844-853, doi:10.1093/humrep/dew008 (2016).

24   Komljenovic, A., Roux, J., Wollbrett, J., Robinson-Rechavi, M. & Bastian, F.
B. BgeeDB, an R package for retrieval of curated expression datasets and for
gene list expression localization enrichment tests. *F1000Research* **5**, 2748,
doi:10.12688/f1000research.9973.2 (2016).

25   Dandapat, A., Hartweck, L. M., Bosnakovski, D. & Kyba, M. Expression of
the human FSHD-linked DUX4 gene induces neurogenesis during
differentiation of murine embryonic stem cells. *Stem cells and development*
22, 2440-2448, doi:10.1089/scd.2012.0643 (2013).

26   Geng, L. N. *et al.* DUX4 activates germline genes, retroelements, and immune
mediators: implications for facioscapulohumeral dystrophy. *Developmental*
cell **22**, 38-51, doi:10.1016/j.devcel.2011.11.013 (2012).

27   Young, J. M. *et al.* DUX4 binding to retroelements creates promoters that are
active in FSHD muscle and testis. *PLoS genetics* **9**, e1003947,
doi:10.1371/journal.pgen.1003947 (2013).

28   Jouhilahti, E. M. *et al.* The human PRD-like homeobox gene LEUTX has a
central role in embryo genome activation. *Development* **143**, 3459-3469,
doi:10.1242/dev.134510 (2016).

29 Eidahl, J. O. *et al.* Mouse Dux is myotoxic and shares partial functional
homology with its human paralog DUX4. *Human molecular genetics* **25**,
4577-4589, doi:10.1093/hmg/ddw287 (2016).

30 Wu, J. *et al.* Chromatin analysis in human early development reveals
epigenetic transition during ZGA. *Nature* **557**, 256-260, doi:10.1038/s41586-
018-0080-8 (2018).

31 Tadros, W. & Lipshitz, H. D. The maternal-to-zygotic transition: a play in two
acts. *Development* **136**, 3033-3042, doi:10.1242/dev.033183 (2009).

32 Walser, C. B. & Lipshitz, H. D. Transcript clearance during the maternal-to-
zygotic transition. *Current opinion in genetics & development* **21**, 431-443,
doi:10.1016/j.gde.2011.03.003 (2011).

33 Lee, M. T., Bonneau, A. R. & Giraldez, A. J. Zygotic genome activation
during the maternal-to-zygotic transition. *Annual review of cell and
developmental biology* **30**, 581-613, doi:10.1146/annurev-cellbio-100913-
013027 (2014).

34 Winata, C. L. *et al.* Cytoplasmic polyadenylation-mediated translational
control of maternal mRNAs directs maternal-to-zygotic transition.
Development **145**, doi:10.1242/dev.159566 (2018).

35 Suzumori, N., Burns, K. H., Yan, W. & Matzuk, M. M. RFPL4 interacts with
oocyte proteins of the ubiquitin-proteasome degradation pathway.
*Proceedings of the National Academy of Sciences of the United States of
America* **100**, 550-555, doi:10.1073/pnas.0234474100 (2003).

36 Rajkovic, A., Lee, J. H., Yan, C. & Matzuk, M. M. The ret finger protein-like
4 gene, Rfpl4, encodes a putative E3 ubiquitin-protein ligase expressed in
adult germ cells. *Mechanisms of development* **112**, 173-177 (2002).

37 Yang, Y. *et al.* The E3 ubiquitin ligase RNF114 and TAB1 degradation are
required for maternal-to-zygotic transition. *EMBO reports* **18**, 205-216,
doi:10.15252/embr.201642573 (2017).

38 Chen, Z. & Zhang, Y. Loss of DUX causes minor defects in zygotic genome
activation and is compatible with mouse development. *Nat Genet* **51**, 947-951,
doi:10.1038/s41588-019-0418-7 (2019).

39 De Iaco, A., Verp S., Offner S., Trono D. DUX is a non-essential synchronizer
of zygotic genome activation. doi:https://doi.org/10.1101/569434 (2019).

40 Madissoon, E. *et al.* Pleomorphic Adenoma Gene 1 Is Needed For Timely
Zygotic Genome Activation and Early Embryo Development. *Scientific
reports* **9**, 8411, doi:10.1038/s41598-019-44882-0 (2019).

41 Kruse K., D. N., Enriquez-Gasca R., Gaume X., Torres-Padilla M-E.,
Vaquerizas JM. Transposable elements drive reorganisation of 3D chromatin
during early embryogenesis. doi:https://doi.org/10.1101/523712 (2019).

42 Franke, V. *et al.* Long terminal repeats power evolution of genes and gene
expression programs in mammalian oocytes and zygotes. *Genome research*
27, 1384-1394, doi:10.1101/gr.216150.116 (2017).

43   Liu, L. *et al.* An integrated chromatin accessibility and transcriptome
landscape of human pre-implantation embryos. *Nature communications* **10**,
364, doi:10.1038/s41467-018-08244-0 (2019).

44   Islam, S. *et al.* Quantitative single-cell RNA-seq with unique molecular
identifiers. *Nature methods* **11**, 163-166, doi:10.1038/nmeth.2772 (2014).

45   Katayama, S., Tohonen, V., Linnarsson, S. & Kere, J. SAMstrt: statistical test
for differential expression in single-cell transcriptome with spike-in
normalization. *Bioinformatics* **29**, 2943-2945,
doi:10.1093/bioinformatics/btt511 (2013).

46   Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for
comparing genomic features. *Bioinformatics* **26**, 841-842,
doi:10.1093/bioinformatics/btq033 (2010).

47   Ramirez, F., Dundar, F., Diehl, S., Gruning, B. A. & Manke, T. deepTools: a
flexible platform for exploring deep-sequencing data. *Nucleic acids research*
42, W187-191, doi:10.1093/nar/gku365 (2014).

48   Heinz, S. *et al.* Simple combinations of lineage-determining transcription
factors prime cis-regulatory elements required for macrophage and B cell
identities. *Molecular cell* **38**, 576-589, doi:10.1016/j.molcel.2010.05.004
(2010).

49   Oki, S. *et al.* ChIP-Atlas: a data-mining suite powered by full integration of
public ChIP-seq data. *EMBO reports* **19**, doi:10.15252/embr.201846255
(2018).

50   Tang, F. *et al.* RNA-Seq analysis to capture the transcriptome landscape of a
single cell. *Nature protocols* **5**, 516-535, doi:10.1038/nprot.2009.236 (2010).

51   Johnson, M. *et al.* NCBI BLAST: a better web interface. *Nucleic acids*
research **36**, W5-9, doi:10.1093/nar/gkn201 (2008).

52   Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software
version 7: improvements in performance and usability. *Molecular biology and*
evolution **30**, 772-780, doi:10.1093/molbev/mst010 (2013).

53   Porollo, A. A., Adamczak, R. & Meller, J. POLYVIEW: a flexible
visualization tool for structural and functional annotations of proteins.
Bioinformatics **20**, 2460-2462, doi:10.1093/bioinformatics/bth248 (2004).

54   Adamczak, R., Porollo, A. & Meller, J. Accurate prediction of solvent
accessibility using neural networks-based regression. *Proteins* **56**, 753-767,
doi:10.1002/prot.20176 (2004).

55   Magnan, C. N. & Baldi, P. SSpro/ACCpro 5: almost perfect prediction of
protein secondary structure and relative solvent accessibility using profiles,
machine learning and structural similarity. *Bioinformatics* **30**, 2592-2597,
doi:10.1093/bioinformatics/btu352 (2014).

56   Wang, S., Li, W., Liu, S. & Xu, J. RaptorX-Property: a web server for protein
structure property prediction. *Nucleic acids research* **44**, W430-435,
doi:10.1093/nar/gkw306 (2016).

57 Piskacek, S. *et al.* Nine-amino-acid transactivation domain: establishment and
prediction utilities. *Genomics* **89**, 756-768, doi:10.1016/j.ygeno.2007.02.003
(2007).

58 Berman, H. M. *et al.* The Protein Data Bank. *Nucleic acids research* **28**, 235-
242, doi:10.1093/nar/28.1.235 (2000).

59 Lehtonen, J. V. *et al.* BODIL: a molecular modeling environment for
structure-function analysis and drug design. *Journal of computer-aided*
molecular design **18**, 401-419 (2004).

60 Case, D. *et al.* AMBER 2018. (2018).

61 Maier, J. A. *et al.* ff14SB: Improving the Accuracy of Protein Side Chain and
Backbone Parameters from ff99SB. *Journal of chemical theory and*
computation **11**, 3696-3713, doi:10.1021/acs.jctc.5b00255 (2015).

62 Zgarbova, M. *et al.* Refinement of the Sugar-Phosphate Backbone Torsion
Beta for AMBER Force Fields Improves the Description of Z- and B-DNA.
Journal of chemical theory and computation **11**, 5723-5736,
doi:10.1021/acs.jctc.5b00716 (2015).

63 Jorgensen, W., Chandrasekhar, J. & Madura, J. Comparison of simple
potential functions for simulating liquid water. *The Journal of chemical*
physics **79**, doi: https://doi.org/10.1063/1.445869 (1983).

64 Essmann, U. *et al.* A smooth particle mesh Ewald method. *The Journal of*
chemical physics **103**, 8577-8593, doi:10.1063/1.470117 (1995).

65 Berendsen, H. J. C., Postma, J. P., van Gunsteren, A., DiNola, A. & J.R., H.
Molecular dynamics with coupling to an external bath. *The Journal of*
chemical physics **81**, doi:https://doi.org/10.1063/1.448118 (1984).

66 Roe, D. R. & Cheatham, T. E., 3rd. PTRAJ and CPPTRAJ: Software for
Processing and Analysis of Molecular Dynamics Trajectory Data. *Journal of*
chemical theory and computation **9**, 3084-3095, doi:10.1021/ct400341p
(2013).

67 Humphrey, W., Dalke, A. & Schulten, K. VMD: visual molecular dynamics.
Journal of molecular graphics **14**, 33-38, 27-38 (1996).

68 Choi, H. *et al.* SAINT: probabilistic scoring of affinity purification-mass
spectrometry data. *Nature methods* **8**, 70-73, doi:10.1038/nmeth.1541 (2011).

69 Mellacheruvu, D. *et al.* The CRAPome: a contaminant repository for affinity
purification-mass spectrometry data. *Nature methods* **10**, 730-736,
doi:10.1038/nmeth.2557 (2013).

70 Wang, J., Vasaikar, S., Shi, Z., Greer, M. & Zhang, B. WebGestalt 2017: a
more comprehensive, powerful, flexible and interactive gene set enrichment
analysis toolkit. *Nucleic acids research* **45**, W130-W137,
doi:10.1093/nar/gkx356 (2017).

71 Meldal, B. H. *et al.* The complex portal--an encyclopaedia of macromolecular
complexes. *Nucleic acids research* **43**, D479-484, doi:10.1093/nar/gku975
(2015).

72   Giurgiu, M. *et al.* CORUM: the comprehensive resource of mammalian
protein complexes-2019. *Nucleic acids research* **47**, D559-D563,
doi:10.1093/nar/gky973 (2019).

73   Buenrostro, J. D., Wu, B., Chang, H. Y. & Greenleaf, W. J. ATAC-seq: A
Method for Assaying Chromatin Accessibility Genome-Wide. *Current*
protocols in molecular biology **109**, 21 29 21-29,
doi:10.1002/0471142727.mb2129s109 (2015).

74   Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics*
29, 15-21, doi:10.1093/bioinformatics/bts635 (2013).

1093

Fig.1

a

DUX4

b

Zygote

2-cell

4-cell

8-cell

c

DUX4 intensity

Fig. 2

Fig.3

Fig. 4

Fig. 5

a TetOn-DUX4 H1 cl 2, cl 8 H9 cl 3, cl 4 +Dox → FACS EmGFP+ → STRT RNA-seq, ATAC-seq; -Dox → FACS EmGFP-

b Expression level relative to 0 hours — DUX4, ZSCAN4, TRIM48 over 0 1 2 3 (h)

c - Dox / + Dox; Bright field; DUX4

d Number of TFEs — UP 6,427; DOWN 5,547
- upstream and 5'-UTR of coding
- CDS and 3'-UTR of coding
- upstream and 1st-exon of noncoding
- other exons of noncoding
- intron
- unannotated

e FE275069 ZSCAN4; FE270433 LEUTX; FE94476 TRIM48; FE463525 Unannotated; FE533694 Unannotated; FE130507 Unannotated — log$_{10}$(normalized reads); EmGFP− EmGFP+

f Oocyte to 4C — 9, 23; 17/23 TFEs overlap with DUX4 binding. 4C to 8C — 14, 33, 81. Up / Down / NS

g DUX4 ChIP-seq intensity — Up-TFEs, All-TFEs; Distance from the peak within TFE (kbp)

h The most enriched de novo motif in Up-TFEs; DUX4 motif (Eidahl JO et al. 2016)

i Enrichment analysis with ChIP−seq datasets — under / over; SRX107284 DUX4 Myoblasts; SRX2529721 DUX4 hESCs; SRX2146136 EZH2 LNCAP; SRX2769830 OTX2 D283 Med; SRX2436059 OTX2 D283 Med; −log$_{10}$ (Q-value); Fold enrichment (Up−TFEs/All−TFEs)

Fig. 6

a
KHDC1 pseudo gene 1
ENA accession LR694084, LR694085

b
RING-finger type E3 ubiquitin ligase
ENA accession LR694082, LR694083

c
RING-finger domain protein
ENA accession:
LR694086, LR694087 (Ring10.2)
LR694088, LR694089 (Ring4.2)

Fig. 7