Comparing the utility of *in vivo* transposon mutagenesis approaches in yeast species to infer gene essentiality

Anton Levitan[1,2], Andrew N. Gale[3], Emma K. Dallon[3], Darby W. Kozan[3], Kyle W. Cunningham[3], Roded Sharan[4] and Judith Berman[1]

[1]School of Molecular Microbiology and Biotechnology, Faculty of Life Sciences, Tel Aviv University, Tel Aviv, Israel

[2]Edmond J. Safra Center for Bioinformatics, Tel Aviv University, Tel Aviv, Israel

[3]Department of Biology, Johns Hopkins University, Baltimore, MD, USA

[4]Blavatnik School of Computer Science, Tel Aviv University, Tel Aviv, Israel

27**ABSTRACT**

29**Background**:

30*In vivo* transposon mutagenesis coupled with deep sequencing enables large-
31scale genome-wide mutant screens for genes essential in different growth
32conditions. Six large scale studies have now been performed with three yeast
33species (*S. cerevisiae, S. pombe* and *C. albicans*), each mutagenized with two
34of three different heterologous transposons (*AcDs*, *Hermes,* and *PiggyBac*).

36**Results**: We analyzed predictions of gene essentiality for each of the six
37studies and evaluated the ability of the data to predict gene essentiality using
38a machine-learning approach. Important data features included a sufficient
39number of independent insertions and the degree of random insertion
40distribution. All transposons showed some bias in insertion site preference,
41both because of jackpot events, specific insertion sequence preferences and
42preferences for short-range vs long range insertions. For *PiggyBac*, a stringent
43target sequence limited the ability to predict essentiality in genes with few or
44no target sequences. Furthermore, the machine learning approach is robust for
45predicting gene function in less well-studied species by leveraging cross-
46species orthologs. Finally, comparisons of isogenic diploid vs haploid *S.*
47*cerevisiae* isolates identified several genes that are haplo-insufficient, while
48most essential genes, as expected, were recessive.

Conclusions: We provide recommendations for the choice of transposons and
the inference of gene essentiality in genome-wide studies of eukaryotic
microbes such as yeasts, including species thathat have been less amenable to
classical genetic studies. These include maximizing the number of unique
insertions, avoiding transposons with stringent target sequences and a method
for cross-species transfer learning.

INTRODUCTION

Work with model yeasts such as *Saccharomyces cerevisiae* and *S. pombe* has
pioneered the combination of genetype/phenotype comparisons at a genomic
scale.   For these yeasts, with genome sequences available for over 20 years[1,
2] and facile gene replacement protocols, have relied heavily on
comprehensive collections of deletion mutants[3, 4] for high throughput
dissection of specific genotypes as well as for genetic interactions with drugs
(reviewed in Lehár et al.[5]) and for gene-gene interactions through systematic
analysis of double and triple mutant analysis (e.g., Reguly et al.[6], Kuzmin et
al.[7]). In animals and plants that are less amenable to such directed molecular
manipulations, the use of heterologous transposons *in vivo* has facilitated
genetic analysis, within the limitations imposed by the transposon
excision/insertion process[8, 9]. With the advent of deep sequencing, such
studies have also become more facile and have been performed in the two
model yeasts as well[10, 11].

In vivo transposon mutagenesis generally involves the introduction of a
heterologous DNA transposon, along with the genes (e.g., the relevant
transposase) required to induce its active transposition into a clonal isolate of a
species of interest. Upon induction, the transposase excises the transposon
from its original location (excision site) and inserts it into a single new position
in the genome. Each cell harbors, at most, a single transposition mutation
because the frequency of transposon excision and reinsertion is quite low. The
transposon is usually engineered for facile selection of excision and/or
reinsertion events, allowing detection and enrichment of these rare events.
83
In vivo transposon insertion provides several advantages, as it rapidly yields
large numbers of mutants in a single step and easily can be performed in
parallel strains with different mutations or genetic backgrounds. Because it
does not require much prior knowledge, it can also be performed in non-model
species, where each experiment is likely to be highly informative.   The only
transformation steps required are those used to engineer the starting strain.
This bypassing the problem of low transformation efficiency in many species. It
also avoids the unintended genome alterations (e.g., aneuploidies) that often
accompany DNA transformation[12]. The sites of transposon insertion
throughout the genome can be identified *en masse* using very large collections
of independent insertion event clones, coupled with deep sequencing of the
DNA immediately adjacent to the new transposon locus. For example, a high
throughput genotype/phenotype analysis of 30 bacterial species grown under
>170 different nutrient and stress conditions recently assigned functions to

98thousands of genes including ~300-600 genes per bacterium that are essential

99for viability[13].

101Three different transposon systems have been used for *in vivo* mutagenesis in

102yeasts: *AcDs from Zea mays, Hermes from Musca domestica and PiggyBac,*

103*from Trichoplusia ni*. *AcDs* has been used primarily in plant species but was

104also engineered for increased efficiency in the model yeast *S. cerevisiae*[14]

105and, later, in *C. albicans*[15]. *AcDs* does not display any insertion sequence

106preference, although it has a higher frequency of insertions into intergenic

107regions than coding regions and has a bias for reinsertion near the initial site of

108excision. *Hermes* has been used for mutagenesis in *S. pombe* and *S.*

109*cerevisiae*[17, 18]; it prefers to insert at genomic positions with the target

110sequence TnnnnA. *PiggyBac* (PB) has been used in mammalian systems such

111as rat, mouse[19] and also in *S. pombe*[17]. *PB* has a strong preference for

112insertion at TTAA sequences, which are generally more frequent in A-T-rich

113intergenic regions than within coding sequences.

115Transposon insertion within an ORF is generally assumed to cause a loss-of-

116function mutation. Identifying the phenotypes associated with loss-of-function

117mutations in specific genes allows the prediction of genetic functions. Cells in

118which the transposon inserted into a gene essential for viability will fail to grow

119and thus be lost from the population. By contrast, cells with mutations in non-

120essential genes are expected to be well-represented in the cell population.

121Insertion of a transposon carrying a strong promoter into an ORF could activate

122expression inappropriately; can be useful for the study of gain-of-function

123mutations.

125*In vivo* transposon mutagenesis studies of yeasts include analysis of *S.*

126*cerevisiae* with *Hermes* (this study) or with the mini-Ds derivative of the *AcDs*

127system[10], in *S. pombe* with *Hermes*[11] and *PB*[20] and in *C. albicans* with

128*AcDs*[21] and with *PB*[22] (Fig. 1A). In earlier work, we applied a machine

129learning (ML) approach to infer gene essentiality from the *C. albicans AcDs*

130data. Here, we modified the ML approach to predict the likelihood of

131essentiality for the complete set of predicted open reading frames these sixth

132*in vivo* transposon datasets. We compared the strengths and challenges of the

133different transposons in each species, with the goal of reaching insights

134concerning the number of insertion events required for accurate predictions,

135the distribution of mutations, and the degree to which different transposons,

136with different sequence dependencies, provided similar or different

137conclusions. The goal was to provide metrics that assist in determining the

138advantages and disadvantages of different transposon systems so as to

139optimize the data produced in a given *in vivo* transposon system and to

140suggest approaches for generating whole genome data in understudied yeast

141species.

143**RESULTS AND DISCUSSION**

145**A comparative analysis of the transposon mutagenesis studies**

We compared six *in vivo* transposon insertion mutagenesis experiments,
produced using three different heterologous transposons (*AcDs, Hermes* and
PiggyBac) in three different species (*S. cerevisiae, S. pombe and C. albicans*).
Details of the datasets are provided in the methods section and relevant
parameters are highlighted in Table 1. The number of transposition events
detected in the different studies varied considerably, from over 500,000 unique
insertion sites (hits) and 84 M total reads for the *C. albicans AcDs* (*CaAcDs),* to
as few as 37,500 unique hits and 6.1 M reads in the *S. pombe PiggyBac* (*SpPB*)
data set (Table 1). The number of reads per hit also varied considerably, from
41 to 170.
157

Overview of ML approach for gene essentiality prediction

We first mapped the hit and read frequency of the transposons in each of the
three reference genomes (Fig. 1C). Sites of transposon insertion were identified
based upon targeted sequencing of regions adjacent to the inserted
transposon. Slightly different sequencing protocols were used in the different
studies, but all six essentially amplified Tn-adjacent sequences and mapped
them.
Theoretically, essential genes have no hits and non-essential genes have many
hits; however, distinguishing essential and non-essential genes from the data is
not entirely straight-forward (e.g., Fig. 1C, Gene X_n). To address this
ambiguity, we extended a previous approach for gene essentiality
prediction[21]. We first chose input features from transposon data that were
likely to be informative in the essential/nonessential decision: the number of

unique insertion sites (hits) per ORF, the degree to which those insertion sites

were enriched in the population (reads), as well as normalization factors that

consider the insertion frequency as a function of chromosome position.

Training sets were built using information from the two model species with

gene essentiality data available from classical genetic approaches (e.g., com-

prehensive ORF deletion analysis) (reviewed in Giaever and Nislow[23] and

Spirek et al.[24]) (Table S1 (training sets)). The specificity and sensitivity of the

approach was analyzed using the AUC (area under the receiver operating char-

acteristic curve (Fig. 1D). For *C. albicans,* which did not have extensive prior

knowledge of gene essentiality, we constructed a training set from a core set of

genes whose orthologs were known to be essential in both model yeasts. This

approach is likely to be useful for other species that lack sufficient prior knowl-

edge of gene essentiality to construct a within-species training set.

We assessed the performance of each classifier by producing training sets

using genes known or inferred to be essential and non-essential, as described

in the Methods. The classifiers were trained and their performance, assessed

using the AUC measure, was high across most examined studies (>0.94). The

one exception was the *SpPB* study, which had far fewer unique insertion sites

(Table 1) and had an AUC of 0.785. The highest AUC levels were seen with the

AcDs in both *C. albicans* and *S. cerevisiae*. Of note, these two studies also had

the largest number of total hits and reads. All the considered ML features for

each ORF in every study and the predicted verdicts of essentiality are provided

196in Tables S2-S7. Below, we describe the main insights gained from this

197comparison.

199**Insight #1: Optimize the number of independent insertion sites (hits)**

200**for highest quality predictions of gene essentiality**

202The total number of unique insertion sites (hits) and the performance (AUC

203values) were high highly correlated, and this correlation was statistically

204significant (Fig. 2a; Pearson's r = 0.892; p-value = 0.0169). By contrast, the

205total number of sequencing reads showed a weaker correlation with the AUC

206that was not statistically significant (Fig, 2b; Pearson's r = 0.636; p-value =

2070.1741). If we disregard the worst performing *SpPB*, the correlation of the AUCs

208with the total number of hits rises dramatically to Pearson's r = 0.995; p-value

209= 0.0003, and the correlation with the total number of sequencing reads

210remains similarly weak: Pearson's r = 0.652; p-value = 0.2327. Thus, it a

211library with many independent hits will improve performance and simply

212increasing the number of sequencing reads is not likely to be sufficient to

213obtain optimal results. Increasing the number of independent hits requires

214collection of sufficient numbers of independent colonies soon after transposase

215induction and the resulting transposon excision and reinsertion. An advantage

216of *Hermes* is that most insertions occur during stationary phase, so

217transposase-inducing conditions can be tolerated throughout the growth

218period. Experimental designs that optimize isolation of independent events are

219critical.

221 Insight #2: Avoid libraries with high levels of jackpot events

Jackpot events are the appearance of extraordinarily high numbers of reads in

a very small number of insertion sites. When the number of reads greatly

exceeds the theoretical number of cell divisions in the experiment, this is likely

due to a transposition event happened prior to the induction of transposon

excision in the experiment. Jackpot events are a major pitfall in that much

sequencing capacity is wasted on detection of a single insertion site. Jackpot

events with >1M sequencing reads were present in 4 of the 6 data sets; *SpPB*

and *ScHermes* had no major jackpot events (no hits with ≥1000-fold more

reads than the average read/hit) (Table 1). Both of these libraries also had far

fewer total sequencing reads than the other studies (6M and 18M vs 24-84M for

the other libraries).

Of note, within a data set, some individual experiments had jackpot events that

were far more than others (Table 1), which would be expected if jackpot events

occur stochastically. Importantly, jackpot events were not clearly associated

with one of the three species or with the transposon type. This suggests that

jackpots arise from technical, rather than biological issues.

Avoiding jackpot events is important because they reduce data quality

considerably: the higher the number of reads at a few jackpot sites, the lower

the number of informative hits and reads. Avoiding the selection of cells in

which a transposon was already mobilized is key to ensuring that the number

of hits and reads provide good genome coverage. Dividing the cultures into

246dozens of small cultures and then re-pooling these sub-cultures after

247transposase induction can effectively dilute out most jackpot events. Preparing

248several independent libraries and sampling a few sequences in each may also

249be worthwhile. For example, if a tested library shows one sequence twice in

250one hundred colonies, it is likely to be a >1M jackpot event.

252**Insight #3. Consider which features are most important in the analysis**

253**of a given transposon**

255In decision tree based algorithms, such as Random Forest[25], every node is a

256condition of a split of the data by a single feature. The splitting process

257continues until it reaches a stop condition such as: all the features have been

258used, the obtained subset is very small or the training labels are the same for

259the obtained subset. The goal is to reduce entropy (uncertainty) in the data.

260Entropy is zero when all the labels in the obtained subset are the same; and is

261maximum when half of the labels are the same in the obtained subset (in a

262binary classification). Each split of the data by a given feature (node in the

263tree) reduces the entropy. The importance of a given feature in the Random

264Forrest classifier is the calculated decrease in entropy contributed by that

265feature. Here we describe the features of the classifiers, and discuss their

266relative importance.

269For each ORF, we calculated predictive features including the number of hits,

270number of reads and the length of each ORF, a neighborhood index, which

271normalizes for insertion bias due to genomic position (e.g., proximity to the

272initial excision site in the genome) and a freedom index, which reports the

273proportion of an ORF that is hit-free. The freedom index is especially useful for

274identifying genes with essential domains, that are able to tolerate insertions

275outside of the essential domain. The number of the transposon hits per

276transposon target sequences in an ORF was an additional feature used in the

277analysis (Figure 3), where applicable (in *PB* and *Hermes* studies). Furthermore,

278we calculated the number of hits and the number of reads normalized by the

279length of each ORF. We compared the 'feature importance' for each library to

280ask whether specific features were more important for the different classifiers

281and whether feature importance was characteristic for a given transposon or

282yeast species.

284The number of hits per ORF played an important role in determining

285essentiality, with essential genes having far fewer hits than non-essential ORFs

286(~7 times less, on average, across the 6 datasets), consistent with the strong

287correlation between number of hits and the AUC (Fig. 2A). The number of reads

288per ORF played a lesser role in these classifications, also consistent with the

289correlation above (Fig. 2B). Gene length also affected the probability of

290transposon insertion in a gene, and thus was a crucial normalization parameter

291for the numbers of hits and reads for every ORF.

293The Neighborhood Index (NI) feature made important contributions in all of the

294classifications (except *SpPB*, which had far less data). Importantly, the NI did

295not differ considerably between the different transposons, consistent with the

296idea that chromatin accessibility, 3D chromosome organization and other

297factors that may bias the insertion site frequency in a given organism affect

298the frequency of insertion of different transposons in a similar manner.

300The Freedom Index (FI) was a major contributor to both *ScAcDs* and *CaAcDs*

301predictions while results with the *PB* and *Hermes* datasets were mixed (Fig. 3).

302This is consistent with the idea that *AcDs* does not have a specific target

303sequence and thus inserts throughout ORFs, while *PB* and *Hermes* have fewer

304target sequences within ORFs*.* Thus, the FI more important in *AcDs*

305experiments because hits occur more randomly throughout an ORF.

307The importance of the number of hits in the proximal regulatory sequences

308(100 bp upstream to the start codon) to the essential/non-essential predictions

309was only minor, but is highly variable. For example, the impact of *ScHermes*

310was nearly twice that of *ScAcDs* for this feature. As described further below,

311we posit that this difference is due to cryptic enhancer/promoter activity in the

312miniDs transposon in *S. cerevisiae* that not seen with the *Hermes* transposon.

314**Insight #4: Consider the effect of transposon-specific target sequence**

315**specificity**

317Some transposons have preferred sites of insertion: *Hermes* prefers TnnnnA

318and *PiggyBac* inserts primarily at TTAA sequences; *AcDs* does not have an

319insertion site preference. Theoretically, the length of the insertion site

320sequence necessarily scales inversely with the number of potential unique hit

321sites. However, it was not clear at what insertion sequence length the

322resolution of studies of gene essentiality becomes limiting.

324The feature importance of the number of hits per transposon target sequence

325in an ORF, which should be a measure of library saturation, showed a varying

326degree of importance in the *PB* and *Hermes* studies. Curiously, its importance

327wasn't dependent on the type of the transposon or the target sequence

328prevalence in the genome. This likely because target sequences are preferred

329sites of insertion, yet are not exclusive or absolute. For example, *PiggyBac* in

330*C. albicans* had 1.6-fold more unique insertion sites than the theoretical

331number of target sequences in the *C. albicans* genome. By contrast, for both

332*ScHermes* and *SpHermes*, the number of target sequences available far

333outnumbered the number of unique hits. The proportion of target sequences

334not hit ranged from 8.9% for *CaPB* to 85% for *SpHermes* (Table 1) and the

335proportion of hits not in target sequences ranged from 14% in *SpPB* to ~50% in

336*CaPB* as well as both *Hermes* data sets. Thus, we surmise that the preference

337for target sequences is only a minor limitation for both of these transposons,

338except when the total number of hits is very low as in *SpPB*.

340Another critical issue is the number of genes that lack any preferred target

341sequences within the ORF; there are 228 and 185 ORFs without a single TTAA

342sequence in *C. albicans* and *S. pombe,* respectively. These ORFS have a lower

343probability of acquiring insertions and, if the genes are non-essential, they are

344much more likely to give false positive information (be predicted essential for

345lack of insertions). Indeed, 155 ORFs without TTAA sequences were predicted

346essential in the *CaPB* data and yet are predicted non-essential in the *CaAcDs*
347study. Similarly, 118 of the 185 ORFs lacking TTAA sequences were predicted
348essential from the *SpPB*study, but were non-essential in the *SpHermes* study.
349We assume that many of these genes are false positives, especially given that
350127 of the 185 ORFs lacking TTAA, including 95 of the 118 aforementioned
351ORFs, were non-essential in classical genetics studies of *S. pombe.*

354Next, we asked if the number of target sequences within an ORF affected the
355*CaPB* classification performance for that ORF. To address this, we compared
356the performance (AUC) to sets of ORFs filtered to exclude ORFs with different
357numbers of target sequences (from 0 to 10) from the training set used to train
358the classifier (Fig. 4). The AUC increased from ~0.94 for the entire training set
359to >0.98 for the training set containing only genes with 10 or more target sites
360(~50% of the genes in the training set). This suggests that studies using the
361*PiggyBac* transposon may struggle to correctly infer gene essentiality for ORFs
362with low numbers of target sites.

364**Insight #5. Consider whether the transposon can activate as well as**
365**disrupt gene expression.**

367The prediction of essentiality was based upon the assumption that transposon
368insertion into an ORF disrupted gene expression and produced loss-of-function
369allele. However, this is not necessarily the case for all genes. For example, if
370an insertion allele removes a regulatory domain from a protein, the protein

may become hyperactive, and thus the result would be a gain-of-function

allele. Additionally, some transposons may introduce enhancer and promoter

activities that could increase gene expression in some species. The miniDs

transposon used in the *ScAcDs* data is likely to contain such activities.[10]

Consistent with this idea, the *ScAcDs* dataset contains an average of 1.89

insertions within the first 100 bp of essential genes whereas the other datasets

including *CaAcDs,* which has a transposon modified from the *miniDs,* contain

significantly fewer (0.82 insertions in the first 100 bp of Ess genes).

Additionally, many essential genes of *S. cerevisiae* appeared to tolerate miniDs

insertions, but not *Hermes* insertions, at sites in the 5' UTR that are very close

to the start codon. Thus, the *miniDs* transposon in *S. cerevisiae* may facilitate

inappropriate activation of gene expression when inserted upstream or within

certain genes.

Cross-study analysis

Knowing the full set of essential and non-essential genes in eukaryotic

microbes, including pathogens of humans, animals and plants, will improve our

understanding of common and species-specific properties of these

understudied organisms. Furthermore, once a transposon library has been

collected, it can be screened under many other growth conditions to reveal

genotype/phenotype relationships. *In vivo* transposon analysis of gene

essentiality is a practical and feasible approach, because the cost in time and

resources for obtaining libraries is far lower than that for producing engineered

deletion mutants, especially given that the amount of baseline information

396(other than the genome sequence) about the organisms may be minimal. The

397only technical hurdle is to introduce the heterologous transposon of interest,

398either on a plasmid (where feasible) or into a useful locus within the genome of

399the relevant organism.

401An additional challenge is that ML approaches require a high-quality training

402dataset (of gold-standard essential and non-essential genes). For many non-

403model organisms, such training data is too sparse to build a robust training set.

404For *C. albicans,* we circumvented the low numbers of genes already known to

405be essential by relying upon genes that had been determined to be essential

406from comprehensive classical genetic deletion studies in both model yeasts (*S.*

407*cerevisiae* and *S. pombe)* and that had orthologs in *C. albicans.* Training on *S.*

408*cerevisiae* or *S. pombe* orthologs with consistently essential orthologs yielded

409good performance predictions for *C. albicans* (AUC: 0.940 to 0.993, Table 1)*.*

410*CaAcDs* performance was lower when training only on the 66 genes known to

411be essential plus the set of presumed non-essential genes (those that had been

412successfully deleted in *C. albicans* studies, AUC of ~0.92)[21].

414Next, we considered the quality of the learning performance for each dataset, if

415we trained on orthologs from one species and predicted essentiality of genes in

416a different organism (Figure 5). The transfer learning performance of the

417classifications was of a comparable quality to the single study classifiers for

418most *AcDs* and *Hermes* cases (Figure 5a and Figure 2). Furthermore, it

419displayed a somewhat symmetrical property: in most cases, there were minor

420differences in performance between train/test and test/train pairs (reducing the

421quality by ~ 0.5% to 5.7%) when the tests were between or among *AcDs* and

422*Hermes* experiments. Conversely, when testing for predictions from *PB* data

423that were trained on either *AcDs* or *Hermes,* the AUCs dropped more

424dramatically (up to ~21.9%). Thus, *PB* data was less transferable than the

425*Hermes* and *AcDs* data.

427The low *PB* transferability between *SpPB* and *CaPB* is likely due to the sparser

428target sequence distribution relative to either *Hermes* or *AcDs*, which causes

429*PB* studies to produce false positives as noted above, and thus might

430contribute to reduced performance in cross-study analyses. The lower

431performance of the classifiers in the original *PB* single studies (Table 1), also

432may have contributed to the reduced ability to predict essentiality in pools of

433*PB* mutants using cross-species models.

435Reduced differences cross-study performance could also be due to differences

436between the importance of different features in the classifiers for the different

437datasets. To test this possibility, we correlated the vector of the relative

438feature importance for each study with the feature importance in all the other

439studies (Figure 5b). The analysis distinguished 3 groups within the 6 studies,

440based on the correlation coefficient values for feature importance between

441members of the group: *CaAcDs* and *ScAcDs* (Pearson's r = 0.902); *Sp Hermes*

442and *CaPB* (Pearson's r = 0.898); and *SpPB* and *ScHermes* (Pearson's r =

4430.929). Notably, the quality of the transfer learning predictions appears to be

444independent of both the transposon type and the organism studied, with the

445exception of the *AcDs* studies. We presume that this is due to the lack of a

446specific target sequence for the *AcDs* transposon system.

448**Insight #6: As necessary, construct training sets using genes with**

449**orthologs in models where essentiality is known and then validate the**

450**training set manually.**

452We suggest that an initial training set of orthologous genes known to be

453essential and non-essential in related model organisms can be used to

454facilitate analysis of a transposon insertion study in a non-model organism with

455sparse essentiality information. An important caveat is that differences

456between gene function in different species can alter gene essentiality of a

457small number of these orthologs; thus, it is important to visually inspect this

458orthologous training set before applying it. The goal is to remove any genes

459with insertion patterns that are highly contradictory to the 'essentiality label'

460that the orthologs provided. For example, for *C. albicans*, the entire

461orthologous training set was reviewed in an unprejudiced fashion by three

462independent inspectors, who visually reviewed the insertion patterns in the

463*CaAcDs* data and manually labeled each gene as essential, non-essential or

464ambiguous. When all three inspectors classified a gene as non-essential (e.g.,

465many insertions throughout an ORF within a genome region that had many

466insertions outside of that ORF) and the orthologs were labeled 'essential' in the

467two model yeasts, we removed that gene from the training set.

469Once a training set has been established, and the features for the ORFs have

470been calculated, the Random Forest classifier can be run in a cross-validation

471scheme and the AUC can be calculated using the essentiality labels. This

472provides an efficient approach to obtain information about all of the genes in a

473species that has been sequenced but not subjected to much molecular

474manipulation. Clearly, the same approach can be used to compare the

475essentiality of the same sets of genes grown in different conditions as well,

476potentially providing large amounts of phenotypic data across an entire set of

477ORFs. If applied to a species that had not been the subject of many genetic

478studies, such data would represent a treasure-trove of information about genes

479that had not been previously studied and the phenotypes associated with loss-

480of-function of those genes.

482**A Comparative Analysis of Gene Essentiality Predictions**

484An important issue is whether different transposon insertion studies in the

485same organism had similar or different predictions from one another and from

486the known essentiality status of deletion mutants, which are by definition 'loss-

487of-function' null alleles. For *S. cerevisiae,* the classifiers displayed a high

488degree of agreement on the final verdicts of gene essentiality (Figure 6a),

489while more discrepancies were evident for the *C. albicans* and *S. pombe*

490studies. Both *PiggyBac* studies predicted a much higher number of essential

491genes than the *AcDs* or *Hermes* studies (Figure 6b and 6c) as expected from

492the paucity of target sequences that are likely to give false positive predictions

493discussed above. For example, the *CaPB* study had an average 5.84 target

sites per kb in genes likely to be false positives vs. 10.62 target sites per kb in

all the genes (Mann Whitney U: p-value < $2.38*e^{-78}$). Importantly, when

compared to the set of essential genes for each species determined by deletion

analysis, the transposon studies also did quite well, with only 20 to 35% of the

genes in disagreement. In some cases, such discrepancies were found to be

due to issues with the deletion collection isolates. For example, ~8% of the

original *S. cerevisiae* deletion collection carried aneuploidies or gene

amplifications,[26] and ~10% of *S. pombe* deletion strains retained a wild-type

copy of the ORF that had been targeted for deletion. Extra copies of the

'deleted' gene reduces the apparent number of essential genes.

Gene essentiality in haploid versus diploid strains of *S. cerevisiae*

S. cerevisiae is readily grown in both the diploid and haploid states, which

allows identification of the haplo-insufficient subset of genes among the set of

essential genes. Based on gene knockout studies, only 2 genes (*NDC1*, *MLC1*)

were classified as haplo-insufficient,[27, 28] while all other essential genes

were haplo-proficient (i.e. heterozygous knockouts in diploids were viable). To

determine whether additional haplo-insufficient genes exist in *S. cerevisiae*, we

collected *ScHermes* insertions in diploid strain BY4743 and compared them to

haploid strains BY4741 and BY4742. *Sc Hermes* transposon mutagenesis

libraries were used with the classifier that had been trained on the *SpHermes*

haploid training set data, applying the same threshold for classification (Figure

8). The classifier identified 155 genes as "essential in both haploid and

diploid", a number far higher than expected. Upon closer analysis, 98

519contained regions of poor mapping due to duplications elsewhere in the

520genome, 50 were categorized as dubious in the Saccharomyces Genome

521Database (yeastgenome.org), one (*LEU2*) had been deleted in the strains

522studied, and the two known haploinsufficient genes (*NDC1, MLC1*) had been

523identified, providing support for this approach. Upon visual inspection of data

524of the remaining four genes, one essential gene (*BCY1*) appeared

525haploinsufficient, whereas another (*RPC10*) contained numerous insertions in

526its 5' UTR in diploids but not haploids, suggesting that it might not be essential

527(Fig. 9). The other two ORFs predicted to be haploinsufficient are very small

528(165-225 bp) and also are within regions of sparse insertion density. Thus, we

529have lower confidence in the data for these two genes. Thus, transposon

530mutagenesis of a diploid strain successfully revealed the two known

531haploinsufficient genes and one new one (*BCY1*), which is known to be

532essential in the conditions employed in the screen, but not essential in other

533culture conditions[29].

535The classifier also identified 29 genes as "haploinsufficient" in diploids and not

536essential in haploids. Of these, 20 could be dismissed based on their

537annotation as dubious or the presence of duplicated (unmappable) segments.

538 All of the remaining 9 genes were small (87-528 bp) and were found in regions

539of sparse insertion density. These genes are annotated in SGD as non-

540essential and are probably false positives.

542These observations raise an important issue about data quality control. It is

543important to filter dubious and uninformative ORFs from the data set (as was

544done in the analysis of *CaAcDs*[21]. This includes genes with repeated domains

545or duplicate copies in the genome that prevent unambiguous mapping of short

546Illumina reads. Furthermore, predicting the essential/non-essential status for

547short ORFs and especially those located in regions with sparse intergenic

548insertions is more likely to be problematic.

550**Insight #7: Prediction quality increases considerably when**

551**uninformative data such as mitochondrial genome sequences and**

552**duplicated genes that are difficult to map.**

554**In summary,** we suggest a number of metrics for the inference of gene

555essentiality using *in vivo* transposon mutagenesis studies in yeasts, including

556those with little available genetic data. Maximizing the total number of unique

557transposon insertions is the most critical factor in achieving optimal

558performance of the classification. It can be attained by collecting many

559independent insertion clones, striving to reduce the possible jackpot events in

560the study, maximizing the depth of the coverage and by utilizing a transposon

561with a fairly permissive target sequence or no preferred target sequence.

562Furthermore, while transposons with relatively stringent target sequences have

563some advantages for screens that identify individual mutants, for determining

564gene essentiality they are less robust, as the low number of potential target

565sequences, and especially the lack of any target sequences, in certain ORFs

566will increase the likelihood of falsely classifying non-essential genes as

567essential. Additionally, we think that transposon mutagenesis is an ideal

568approach to gain large amounts of useful genotype/phenotype data

569understudied organisms: the cross-species learning methodology allows

570inference of gene essentiality based on conserved orthologs, especially when

571coupled with visual screening of the data. Finally, *in vivo* transposon

572mutagenesis is an incredibly useful tool for high throughout genomic studies,

573not only of gene essentiality *per se,* but also of genes required under specific

574selective conditions. We hope that the recommendations provided here will

575facilitate future work to understand gene in a wide range of yeast species.

MATERIALS AND METHODS

Data Acquisition

Experimental

Sc Hermes data was obtained as follows: The haploid and diploid strains of *S. cerevisiae* were transformed with plasmid pSG36.[30] A single colony was suspended in 100 mL synthetic complete (SC) medium lacking uracil and containing 2% galactose, divided into twenty 16 x 150 mm glass culture tubes, and shaken for 3 days at 30°C. This protocol yielded ~5 x 10^6 cells bearing transposon insertions per mL (~3% of all cells). To enrich for cells bearing transposon insertions, the twenty cultures were pooled, centrifuged, and the cell pellet was resuspended in 600 mL SC medium containing 2% glucose, 0.1 mg/mL nourseothricin, and 1 mg/mL 5-fluoroorotic acid, and then shaken overnight at 30°C. The cells were pelleted, resuspended in 600 mL of the same medium, and cultured as before. Finally, 60 mL of these enriched cells were pelleted, resuspended in 600 mL of the same medium, and cultured as before. These highly enriched cells were pelleted, resuspended in 15% glycerol, and frozen in aliquots at -80°C. To extract genomic DNA, 100 mg of thawed cell pellets were washed three times in 1 mL deionized water and extracted using Quick-DNA Fungal/Bacterial Miniprep kit (Zymo Research). A total of 2.4 µg of purified gDNA was fragmented by sonication in four separate tubes using a Diagenode Picoruptor. The fragmented DNA was then end repaired, ligated to splinkerette adapters, size selected with AMPure xp beads, and PCR amplified in separate reactions using transposon-specific and adapter-specific primers as detailed previously.[31] Samples were then PCR amplified to attach Illumina P5 and P7 (indexed) adapters, purified with AMPure xp beads, mixed with phiX-

174, loaded into MiSeq instrument (Illumina) and 75 bp of each end was

sequenced using primers specific for Hermes right inverted repeat and P7.

Detailed protocols and primer sequences are available upon request. De-

multiplexed reads were mapped to the *S. cerevisiae* S288C reference genome

using Bowtie2, and any mapped reads with a quality score < 20 or a mismatch

at nucleotide +1 were removed. This process was repeated a total of 3 times in

diploid strain BY4743, 2 times in haploid strain BY4741, and 1 time in haploid

strain BY4742. The diploid and haploid datasets were combined prior to

analyses. The *S.cerevisiae Hermes* data (mapped reads and counts) are

available at http://genome-euro.ucsc.edu/s/CunninghamLab/Hermes%20Vs

%20AcDs . FastQ files are available from Sequence Read Archive (SRA) and

the ArrayExpress Experiment Archive (ArrayExpress), which are core

repositories of the European Nucleotide Archive (ENA) at accession number

[XXXX] https://www.ebi.ac.uk/ena/about/data-repositories (to be updated when

accession number is issued).

Publicly available databases

The rest of the datasets analyzed here were obtained from previously

published studies. *ScAcDs* was published by Michel et al, 2017[10], from which

both WT1 & WT2 were combined for the analysis. The data was downloaded:

https://www.ebi.ac.uk/arrayexpress/experiments/E-MTAB-4885/samples/. *SpPB*

was published by Li et al, 2011[20] and the data was obtained from the SRA

database: SRR089408. *Sp Hermes* was published by Guo et al, 2013[22] and

626the data was obtained from the SRA database: SRR327340. *CaAcDs* was

627published by Segal et al, 2018[21] and the data was obtained from the SRA

628database, where SRR7824843, SRR7824841 and SRR7824838 files, were

629combined for the analysis. *CaPB* was published by Gao et al, 2018[22] and the

630data was obtained from the SRA database, where all the following files were for

631the analysis: DMSO (untreatment): SRR7704188, SRR7704193, SRR7704196;

6325-FOA (untreatment: SRR7704189, SRR7704194, SRR7704200; No drug screen:

633SRR7704195. All the SRR files were obtained using fastq-dump, with the

634following bash command: fastq-dump --gzip --skip-technical --readids --

635dumpbase --split-files --clip <SRR*******>

637**Data Processing**

638The .fastq files downloaded with fastq-dump, were processed using cutadapt to

639filter out reads not containing partial transposon sequences. Reads with

640transposon sequences were trimmed to remove the transposon sequences for

641alignment purposes, as follows: cutadapt --cores=8 -m 2 -g <primer

642sequence> <input fastq filename> -o <output fastq filename> --discard-

643untrimmed --overlap <overlap length>. In the analysis of the *Sc Hermes* study

644all the sequencing reads contained the transposon and the reads start at the

645first genomic base, thus required no filtering. In the analysis of the *Sp PiggyBac*

646we filtered the reads containing the transposons from the rest, by identifying

647the ACGCAGACTATCTTTCTAGGG sequence, cutting it out and aligning only the

648remaining part of the relevant reads. In the analysis of the *Ca AcDs* we filtered

649the reads containing the transposons from the rest, by identifying the

650GTATTTTACCGACCGTTACCGACCGTTTTCATCCCTA sequence, cutting it out and

651aligning only the remaining part of the relevant reads, starting 37bp

652downstream (Segal et al, 2018[21]). In the analysis of the *Ca PiggyBac,* we

653filtered the reads containing the transposons from the rest, by identifying the

654TGCATGCGTCAATTTTACGCAGACTATCTTTCTA sequence, cutting it out and

655aligning only the remaining part of the relevant reads, starting 3bp

656downstream. In the analysis of the *Sc AcDs* study we used the published

657transposon hitmaps of WT1 and WT2. In the analysis of the *Sp Hermes* study

658we used the published transposon hitmaps (Segal et al, 2018[21]).

660**Alignment of reads and mapping the transposon hits**

661bowtie2 indices were created for each organism and gffutils databases were

662created for each organism's genetic features, using the latest versions of the

663reference genomes (fasta) and the genomic feature files (gff), which were

664downloaded from the respective official sources for the three organisms: *S.*

665*cerevisiae* from https://downloads.yeastgenome.org, *S. pombe* from

666ftp://ftp.pombase.org/pombe/ and *C. albicans* http://www.candidagenome.org/

667download/. Sequencing reads were aligned using bowtie2 with the default

668settings. The resulting sam files were converted to bam using samtools. bam

669files were sorted using samtools and indexed via pysam. Transposon hits and

670their corresponding reads were mapped to the respective genomes and

671counted in each genomic feature. Transposon target sites were found in every

672genome using Biopython and counted in each genetic feature.

674**Gene essentiality classification**

Table 2 summarizes the features for machine learning classification that were

engineered from the mapped transposon hits, reads and the transposon target

sequences in the genomes. Random Forrest classification was performed,

using Python's scikit-learn library with the default parameters, except the

n_estimators parameter that was increased to 200, and the random_state

parameter was fixed at 0, for reproducibility purposes. The results were

validated using a 5-fold cross-validation. Essentiality labels for the training set

of each organism were obtained previously (Shiftman et al, 2018) and are

provided in Table S1.

Thresholds for the essentiality predictions in each classification were chosen as

follows: Two metrics were evaluated (Figure 8): 1) Minimum of the Euclidean

distance between (0, 1) and the receiver operating characteristic (ROC) curve.

2) Maximum of the vertical distance between the line describing a random

choice (a straight line from (0, 0) to (1, 1)) and the ROC curve. The first method

was chosen, and we verified that the second metric is reasonably close, to

eliminate any possible artifacts. We predicted the essentiality of all the

available genes for each organism based on their respective features, and

using the aforementioned method to choose the threshold for each binary

classification.

Figures were generated using Python's matplotlib and seaborn libraries. The

schematics were drawn using Inkscape. Mann Whitney U p-values and

Pearson's correlation coefficients and p-values were calculated using Python's

Scipy.

700 FUNDING ACKNOWLEDGEMENTS

701 European Research Council Advanced Award 340087 (RAPLODAPT) to J.B., the

702 Israel Science Foundation (grants no. 715/18, 757/12) to R.S. by NIH R21-

703 AI130722 and T32-GM007231 to KWC.

704 A.L. is supported by a fellowship from the Edmond J. Safra Center for Bioinfor-

705 matics.

References

1. Goffeau A, Barrell BG, Bussey H, Davis RW, Dujon B, Feldmann H, Galibert F, Hoheisel JD, Jacq C, Johnston M, et al: **Life with 6000 genes.** *Science* 1996, **274:**546, 563-547.

2. Wood V, Gwilliam R, Rajandream MA, Lyne M, Lyne R, Stewart A, Sgouros J, Peat N, Hayles J, Baker S, et al: **The genome sequence of** *Schizosaccharomyces pombe.* *Nature* 2002, **415:**871-880.

3. Giaever G, Chu AM, Ni L, Connelly C, Riles L, Veronneau S, Dow S, Lucau-Danila A, Anderson K, Andre B, et al: **Functional profiling of the** *Saccharomyces cerevisiae* **genome.** *Nature* 2002, **418:**387-391.

4. Kim DU, Hayles J, Kim D, Wood V, Park HO, Won M, Yoo HS, Duhig T, Nam M, Palmer G, et al: **Analysis of a genome-wide set of gene deletions in the fission yeast** *Schizosaccharomyces pombe.* *Nat Biotechnol* 2010, **28:**617-623.

5. Lehar J, Stockwell BR, Giaever G, Nislow C: **Combination chemical genetics.** *Nat Chem Biol* 2008, **4:**674-681.

6. Reguly T, Breitkreutz A, Boucher L, Breitkreutz BJ, Hon GC, Myers CL, Parsons A, Friesen H, Oughtred R, Tong A, et al: **Comprehensive curation and analysis of global interaction networks in** *Saccharomyces cerevisiae.* *J Biol* 2006, **5:**11.

7. Kuzmin E, VanderSluis B, Wang W, Tan G, Deshpande R, Chen Y, Usaj M, Balint A, Mattiazzi Usaj M, van Leeuwen J, et al: **Systematic analysis of complex genetic interactions.** *Science* 2018, **360**.

8. Munoz-Lopez M, Garcia-Perez JL: **DNA transposons: nature and applications in genomics.** *Curr Genomics* 2010, **11:**115-128.

9.	Kawakami K, Largaespada DA, Ivics Z: **Transposons as tools for functional genomics in vertebrate models.** *Trends Genet* 2017, **33:**784-801.

10.	Michel AH, Hatakeyama R, Kimmig P, Arter M, Peter M, Matos J, De Virgilio C, Kornmann B: **Functional mapping of yeast genomes by saturated transposition.** *Elife* 2017, **6**.

11.	Guo Y, Park JM, Cui B, Humes E, Gangadharan S, Hung S, FitzGerald PC, Hoe KL, Grewal SI, Craig NL, Levin HL: **Integration profiling of gene function with dense maps of transposon integration.** *Genetics* 2013, **195:**599-609.

12.	Bouchonville K, Forche A, Tang KE, Selmecki A, Berman J: **Aneuploid chromosomes are highly unstable during DNA transformation of *Candida albicans*.** *Eukaryot Cell* 2009, **8:**1554-1566.

13.	Price MN, Wetmore KM, Waters RJ, Callaghan M, Ray J, Liu H, Kuehl JV, Melnyk RA, Lamson JS, Suh Y, et al: **Mutant phenotypes for thousands of bacterial genes of unknown function.** *Nature* 2018, **557:**503-509.

14.	Lazarow K, Du ML, Weimer R, Kunze R: **A hyperactive transposase of the maize transposable element activator (Ac).** *Genetics* 2012, **191:**747-756.

15.	Mielich K, Shtifman-Segal E, Golz JC, Zeng G, Wang Y, Berman J, Kunze R: **Maize transposable elements Ac/Ds as Insertion mutagenesis tools in *Candida albicans*.** *G3 (Bethesda)* 2018, **8:**1139-1145.

16. Betrán E, Long M: **Dntf-2r, a young Drosophila retroposed gene with specific male expression under positive Darwinian selection.** *Genetics* 2003, **164:**977-988.

17. Park JM, Evertts AG, Levin HL: **The Hermes transposon of *Musca domestica* and its use as a mutagen of *Schizosaccharomyces pombe*.** *Methods* 2009, **49:**243-247.

18. Edskes HK, Mukhamedova M, Edskes BK, Wickner RB: **Hermes transposon mutagenesis shows [URE3] prion pathology prevented by a Ubiquitin-targeting protein: Evidence for carbon/nitrogen assimilation cross talk and a second function for Ure2p in *Saccharomyces cerevisiae*.** *Genetics* 2018, **209:**789-800.

19. Zhao S, Jiang E, Chen S, Gu Y, Shangguan AJ, Lv T, Luo L, Yu Z: **PiggyBac transposon vectors: the tools of the human gene encoding.** *Transl Lung Cancer Res* 2016, **5:**120-125.

20. Li J, Zhang JM, Li X, Suo F, Zhang MJ, Hou W, Han J, Du LL: **A piggyBac transposon-based mutagenesis system for the fission yeast *Schizosaccharomyces pombe.*** *Nucleic Acids Res* 2011, **39:**e40.

21. Segal ES, Gritsenko V, Levitan A, Yadav B, Dror N, Steenwyk JL, Silberberg Y, Mielich K, Rokas A, Gow NAR, et al: **Gene essentiality analyzed by in vivo transposon mutagenesis and machine learning in a stable haploid isolate of *Candida albicans*.** *MBio* 2018, **9**.

22. Gao J, Wang H, Li Z, Wong AH, Wang YZ, Guo Y, Lin X, Zeng G, Liu H, Wang Y, Wang J: ***Candida albicans* gains azole resistance by altering sphingolipid composition.** *Nat Commun* 2018, **9:**4495.

23. Giaever G, Nislow C: **The yeast deletion collection: a decade of**
functional genomics. *Genetics* 2014, **197:**451-465.

24. Spirek M, Benko Z, Carnecka M, Rumpf C, Cipak L, Batova M, Marova I,
Nam M, Kim DU, Park HO, et al: **S. pombe genome deletion project:**
an update. *Cell Cycle* 2010, **9:**2399-2402.

25. Breiman L: **Random Forests.** *Machine Learning* 2001, **45:**5-32.

26. Hughes TR, Roberts CJ, Dai H, Jones AR, Meyer MR, Slade D, Burchard J,
Dow S, Ward TR, Kidd MJ, et al: **Widespread aneuploidy revealed by**
DNA microarray expression profiling. *Nat Genet* 2000, **25:**333-337.

27. Chial HJ, Giddings TH, Jr., Siewert EA, Hoyt MA, Winey M: **Altered**
dosage of the *Saccharomyces cerevisiae* spindle pole body
duplication gene, NDC1, leads to aneuploidy and polyploidy. *Proc*
Natl Acad Sci U S A 1999, **96:**10200-10205.

28. Stevens RC, Davis TN: **Mlc1p is a light chain for the unconventional**
myosin Myo2p in *Saccharomyces cerevisiae*. *J Cell Biol* 1998,
142:711-722.

29. Matsumoto K, Uno I, Oshima Y, Ishikawa T: **Isolation and**
characterization of yeast mutants deficient in adenylate cyclase
and cAMP-dependent protein kinase. *Proc Natl Acad Sci U S A* 1982,
79:2355-2359.

30. Gangadharan S, Mularoni L, Fain-Thornton J, Wheelan SJ, Craig NL: **DNA**
transposon Hermes inserts into DNA in nucleosome-free regions
in vivo. *Proc Natl Acad Sci U S A* 2010, **107:**21966-21972.

31. Bronner IF, Otto TD, Zhang M, Udenze K, Wang C, Quail MA, Jiang RH,
Adams JH, Rayner JC: **Quantitative insertion-site sequencing**

(QIseq) for high throughput phenotyping of transposon mutants.

Genome Res 2016, **26:**980-989.

32. **UCSC Genome Browser on *S. cerevisiae* Apr. 2011**

(SacCer_Apr2011/sacCer3) Assembly

[http://genome-euro.ucsc.edu/cgi-bin/hgTracks?

db=sacCer3&lastVirtModeType=default&lastVirtModeExtraState=&virtM

odeType=default&virtMode=0&nonVirtPosition=&position=chrIX

%3A183118%2D203117&hgsid=232744249_LxCF1Ofg7R9Jcsy2DHsyDa9

84DWn]

Figure Legends
816
Figure 1. Overview of data acquisition and analysis: transposition events to

gene essentiality.

a. Three yeast species analyzed (Sp, *S. pombe*; Sc, *S. cerevisiae*; and Ca, *C.*

albicans) by in vivo transposition in this study and which transposons

(PB, PiggyBac; AcDs and Hermes) were used to mutagenize which

species. Note that each species was analyzed with two different

transposon systems.

b. Comparison of genome insertion sites for transposition events that

initiate from an extrachromosomal plasmid (left, red region of plasmid

circle) or a specific chromosomal locus (right, red bar on a given

chromosome). Horizontal lines represent multiple copies of the same

genome, each of which underwent a single insertion event (green arrow)

per genome. While transposition is generally random, a bias for loci in

close proximity to the initial transposon insertion site demands

normalization of the final data.

c. Mapped Tnseq analysis of the pool of transposition events yields the

chromosomal insertion sites (brown vertical lines) in the reference

chromosomes relative to the ORFs (purple regions). A close up of a small

region of a single chromosome (olive horizontal line) including 5 ORFs is

illustrated.

d. A training set is constructed using known or inferred labels (non-

essential, blue; essential, red) together with extracted features

calculated from the data and its position relative to ORFs. Features are

defined in Table 2.

e. The training set features, as well as features for all ORFs are used as

input for Random Forest classification (black rectangle); output is a

prediction of essentiality (red or blue as in d), for which an optimal

threshold is determined and applied to designate all genes in one of the

two categories.

847*Figure 2*. Contribution of unique hits and total number of reads to the quality of

848ML predictions for gene essentiality/non-essentiality.

849Performance of the classifier vs (a) the total number of unique insertion sites

850(hits) and (b) the total number of sequencing reads, in each study (organism

851abbreviations as in Fig. 1a; Ac, AcDs; H, Hermes; PB, PiggyBac).

853*Figure 3.* Feature importances in the different classiffiers.

854Importance of each feature used in the Random Forest classifier of essentiality

855for each dataset. Features are described in Table 2; Neighborhood index

856generally normalizes for non-random insertion frequencies across the genome;

857Freedom index reports on the largest proportion of an ORF that has no hits,

858which is a measure of domains that may be essential.[21]

860*Figure 4.* Analysis of the ability of the CaPB classifier to infer gene essentiality

861in genes with increasing number of target sequences.

862When only ORFs with a specific number of target sites are considered (>= x-

863axis), AUC rises accordingly (red), but the number of ORFs that can be

864analyzed necessarily decreases (numbers above red dots). This demonstrates

865the importance of the prevalence of the transposon target sequences in ORFs,

866for the quality of gene essentiality inference, using in-vivo transposon

867mutagenesis studies. X axis: Minimum number of target sequences per ORF

868needed for inclusion in the classification. Y-axis (red): CaPB classifier AUC.

870*Figure 5.* Analysis of ROC AUC values for Random Forest classification trained

871on data from a different organism and/or transposon in all possible

872combinations.

a. For each ROC AUC value in the table, training was performed on 80% of

the original training set used in the training species/transposon described

in the rows. This training data was then used to predict the essentiality of

the remaining 20% of the training set in the species/transposon

described in the columns. The train/test split ratio was similar to the 5-

fold cross-validation performed in the single study analyses.

b. For each study, the vector of the relative feature importance was

correlated with the feature importance in all the other studies. Pearson r

correlation coefficient values are presented.

Figure 6: Comparison with known essentials genes.

a. Comparison of the essentiality verdicts in *S. cerevisiae*, based on the

known essential genes from the literature, ScAcDs and ScHermes

classifiers.

b. Comparison of the essentiality verdicts in *S. pombe*, based on the known

essential genes from the literature, SpPB and SpHermes classifiers.

c. Comparison of the essentiality verdicts in *C. albicans*, based on the

known essential Sp and Sc orthologs from the literature, CaAcDs and

CaPB classifiers.

Classification thresholds differ slightly from the previously published

analyses [21] based on threshold selection applied systematically to all 6

studies (described in detail in Methods).

Figure 7. Gene essentiality in haploid and diploid *S. cerevisiae*.

Comparison of essential genes in haploid and diploid *S. cerevisiae* analyzed

with Sc Hermes. RF classifier was trained on the haploid ScHermes study and

predicted gene essentiality in a diploid strain, using the same threshold for the

final verdict. Mitochondrial genes were not considered.

Figure 8. Threshold optimization.

903Two metrics were evaluated: 1) Minimum of the Euclidean distance between (0,

9041) and the receiver operating characteristic (ROC) curve. 2) Maximum of the

905vertical distance between the line describing a random choice (a straight line

906from (0, 0) to (1, 1)) and the ROC curve.

908*Figure 9.* Suspected haploinsufficient genes in *S. cerevisiae*.

909Four genes suspected to be haploinsufficient in *S. cerevisiae: NDC1, MLC1,*

910*RPC10 and BCY1*, as they appear in the UCSD genome browser [32]. *NDC1* and

911*MLC1* were known to be haploinsufficient. *BCY1* appears to be a previously

912unknown haploinsufficient gene. *RPC10* might not be essential as it sustained

913hits in the 5' UTR in diploids but not haploids.

Figure 1. Overview of data acquisition and analysis: transposition events to gene essentiality.

A. Three yeast species analyzed (Sp, *S. pombe*; Sc, *S. cerevisiae*; and Ca, *C. albicans*) by in vivo transposition in this study and which transposons (PB, PiggyBac; AcDs and Hermes) were used to mutagenize which species. Note that each species was analyzed with two different transposon systems.

B. Comparison of genome insertion sites for transposition events that initiate from an extrachromosomal plasmid (left, red region of plasmid circle) or a specific chromosomal locus (right, red bar on a given chromosome). Horizontal lines represent multiple copies of the same genome, each of which underwent a single insertion event (green arrow) per genome. While transposition is generally random, a bias for loci in close proximity to the initial transposon insertion site demands normalization of the final data.

C. Mapped Tnseq analysis of the pool of transposition events yields the chromosomal insertion sites (brown vertical lines) in the reference chromosomes

relative to the ORFs (purple regions). A close up of a small region of a single chromosome (olive horizontal line) including 5 ORFs is illustrated.

D. A training set is constructed using known or inferred labels (non-essential, blue; essential, red) together with extracted features calculated from the data and its position relative to ORFs. Features are defined in Table 2.

E. The training set features, as well as features for all ORFs are used as input for Random Forest classification (black rectangle); output is a prediction of essentiality (red or blue as in d), for which an optimal threshold is determined and applied to designate all genes in one of the two categories.

F.

Figure 2. Contribution of unique hits and total number of reads to the quality of ML predictions for gene essentiality/non-essentiality.

        ☐   Performance of the classifier vs (a) the total number of unique insertion sites (hits) and (b) the total number of sequencing reads, in each study (organism abbreviations as in Fig. 1a; Ac, AcDs; H, Hermes; PB, PiggyBac).

Figure 3. Feature importances in the different classifiers.

⬚ Importance of each feature used in the Random Forest classifier of essentiality for each dataset. Features are described in <mark>Table XXX</mark>; Neighborhood index generally normalizes for non-random insertion frequencies across the genome; Freedom index reports on the largest proportion of an ORF that has no hits, which is a measure of domains that may be essential.[21]

⬚

Figure 4. Analysis of the ability of the CaPB classifier to infer gene essentiality in genes with increasing number of target sequences.

- When only ORFs with a specific number of target sites are considered (>= x-axis), AUC rises accordingly (red), but the number of ORFs that can be analyzed necessarily decreases (numbers above red dots). This demonstrates the importance of the prevalence of the transposon target sequences in ORFs, for the quality of gene essentiality inference, using in-vivo transposon mutagenesis studies.
- X axis: Minimum number of target sequences per ORF needed for inclusion in the classification.
- Y-axis (red): CaPB classifier AUC.

Figure 5. Analysis of ROC AUC values for Random Forest classification trained on data from a different organism and/or transposon in all possible combinations.

A. For each ROC AUC value in the table, training was performed on 80% of the original training set used in the training species/transposon described in the rows. This training data was then used to predict the essentiality of the remaining 20% of the training set in the species/transposon described in the columns. The train/test split ratio was similar to the 5-fold cross-validation performed in the single study analyses.

B. For each study, the vector of the relative feature importance was correlated with the feature importance in all the other studies. Pearson r correlation coefficient values are presented.

a. Transfer learning AUCs

b. Correlation of feature importance vectors

C.

Figure 6: Comparison with known essentials genes.
  A. Comparison of the essentiality verdicts in *S. cerevisiae*, based on the known essential genes from the literature, ScAcDs and ScHermes classifiers.
  B. Comparison of the essentiality verdicts in *S. pombe*, based on the known essential genes from the literature, SpPB and SpHermes classifiers.
  C. Comparison of the essentiality verdicts in *C. albicans*, based on the known essential Sp and Sc orthologs from the literature, CaAcDs and CaPB classifiers.
  ● Classification thresholds differ slightly from the previously published analyses[21]based on threshold selection applied systematically to all 6 studies (described in detail in Methods).

a.

b.

c.

Figure 7. Gene essentiality in haploid and diploid *S. cerevisiae*.

    ☐   Comparison of essential genes in haploid and diploid *S. cerevisiae* analyzed with Sc Hermes. RF classifier was trained on the haploid ScHermes study and predicted gene essentiality in a diploid strain, using the same threshold for the final verdict. Mitochondrial genes were not considered.

Figure 8. Threshold optimization.

- Two metrics were evaluated: 1) Minimum of the Euclidean distance between (0, 1) and the receiver operating characteristic (ROC) curve. 2) Maximum of the vertical distance between the line describing a random choice (a straight line from (0, 0) to (1, 1)) and the ROC curve.

-

Figure 9. Suspected haploinsufficient genes in *S. cerevisiae*.

- Four genes suspected to be haploinsufficient in *S. cerevisiae*: NDC1, MLC1, RPC10 and BCY1, as they appear in the genome.[32]NDC1 and MLC1 were known to be haploinsufficient. BCY1 appears to be a previously unknown haploinsufficient gene. RPC10 might not be essential as it sustained hits in the 5' UTR in diploids but not haploids.

NDC1

MLC1

RPC10

BCY1

Table 1. Statistics of the transposon data sets

	Ca AcDs	Sc AcDs	Sc Hermes	Sp Hermes	Ca PiggyBac	Sp PiggyBac
Transposon target sequence	-	-	TnnnnA	TnnnnA	TTAA	TTAA
Initial transposon insertion	Genome	Plasmid	Plasmid	Plasmid	Genome	Genome
Number of target sequences (10^3)	-	-	1154.84	1302.41	120.27	111.37
Total number of unique hits (10^3)	588.97	514.89	444.41	382.82	191.49	37.5
Target Sequences without a hit (10^3)	-	-	924.56	1110.2	10.69	79.09
Percent of target sequences without a hit	-	-	80.06%	85.24%	8.89%	71.02%
Percent of hits in target sequences	-	-	51.82%	50.21%	56.74%	86.06%
Total number of reads (10^6)	84.16	47.1	18.22	23.92	32.58	6.14
Average number of reads per hit	143	91	41	62	170	164
Standard deviation reads per hit	9250	4357	137	5069	4584	481
Highest reads per hit (10^3)	3254.83	2210.55	11.72	2355.61	1301.82	48.31
Highest reads per hit / average reads per hit	22761	24292	286	37994	7658	295
Number of over 106 reads per hit	10	2	0	2	1	0
Average number of hits per gene	29.11	44.05	30.36	24.64	11.81	1.32
Number of genes with 0 hits	300	261	332	99	397	2580
Average number of target sequences per gene	-	-	123.64	131.44	10.62	8.39
Number of genes with 0 target sequences	-	-	0	0	228	186
ROC AUC	0.993	0.985	0.972	0.962	0.94	0.785

Table 2. Classification features

Feature	Description
Hits	Number of transposon hits within the ORF
Reads	Number of reads associated with the transposon hits within the ORF
Neighborhood Index (NI)	Number of transposon hits within the ORF, normalized by length of the ORF and the surrounding 10 kbp
Freedom Index (FI)	Length of the largest hit-free region in the ORF, divided by the ORF's length
Hits 100 upstream	Number of transposon hits within the upstream region of the ORF
Hits per Target Seqs	Number of transposon hits divided by the number of transposon target sequences within an ORF
Reads per Length	Number of transposon hits divided by the length of the ORF
Hits per Length	Number of reads associated with the transposon hits divided by the length of the ORF