1

Distinct microbial communities alter litter decomposition rates in a fertilized coastal plain wetland

Megan E. Koceja[1], Regina B. Bledsoe[1], Carol Goodwillie[1], and Ariane L. Peralta[1,†]

[1]Department of Biology, East Carolina University, Howell Science Complex, Mail Stop 551,

Greenville, NC 27858, USA

†Corresponding author: telephone: +1 252.328.2712, e-mail: peraltaa@ecu.edu

Author contributions: MEK, RBB, CG, and ALP conceived and designed the research; MEK,

RBB, and ALP collected and analyzed the data; MEK and ALP wrote the manuscript; all authors

performed field work and edited the manuscript.

ABSTRACT

Human activities have led to increased deposition of nitrogen (N) and phosphorus (P) into soils. Nutrient enrichment of soils is known to increase plant biomass and rates of microbial litter decomposition. However, interacting effects of hydrologic position and associated changes to soil moisture can constrain microbial activity and lead to unexpected nutrient feedbacks on microbial community structure-function relationships. Examining how feedbacks of nutrient enrichment on decomposition rates is essential for predicting microbial contributions to carbon (C) cycling as atmospheric deposition of nutrients persists. This study explores how long-term nutrient addition and contrasting litter chemical quality influence soil bacterial community structure and function. We hypothesize that long-term nutrient enrichment of low fertility soils alters bacterial community structure and leads to higher rates of litter decomposition with decreasing C:N ratio of litter; but low nutrient and dry conditions limit constrain microbial decomposition of high C:N ratio litter. We leverage a long-term fertilization experiment to test how nutrient enrichment and hydrologic manipulation (due to ditches) affects decomposition and soil bacterial community structure in a nutrient poor coastal plain wetland. We conducted a litter bag experiment and characterized litter-associated and bulk soil microbiomes using 16S rRNA bacterial sequencing and quantified litter mass losses and soil physicochemical properties. Results revealed that distinct bacterial communities were involved in decomposing higher C:N ratio litter more quickly in fertilized compared to unfertilized especially under drier soil conditions, while decomposition rates of green tea litter (lower C:N ratio) were similar between fertilized and unfertilized plots. Bacterial community structure in part explained litter decomposition rates, and long-term fertilization and drier hydrologic status affected bacterial diversity and increased decomposition rates. However, community composition associated with high C:N litter was similar in wetter plots with available

nitrate detected, regardless of fertilization treatment. This study provides insight into long-term

fertilization effects on soil bacterial diversity and composition, decomposition, and the increased

potential for soil C loss as nutrient enrichment and hydrology interact to affect historically low

nutrient ecosystems.

Keywords: biodiversity-ecosystem function; carbon cycling; decomposition; microbiomes; tea bag

index; wetlands.

INTRODUCTION

Humans modify their landscapes through fossil fuel burning, deforestation, and intense

agricultural activity (Vitousek et al. 2010, Fowler et al. 2013). These anthropogenic disturbances

have led to increased atmospheric deposition of nitrogen (N) and phosphorus (P), and can be

particularly disruptive to historically nutrient-limited ecosystems (Guignard et al. 2017). This

increased nutrient deposition can cause a fertilization effect on plant-microbial interactions which

results in increased biomass and shifts in community structure (Cherif and Loreau 2009, Leff et

al. 2015, Harpole et al. 2016). This fertilization effect can increase plant biomass C, which fuels

heterotrophic microbial growth and leads to increased respiration of CO_2 (Hoosbeek et al. 2004,

Kuzyakov 2010). The extent to which nutrient enrichment predicts decomposition rates is also

determined by the interaction of soil microorganisms with the plant inputs and the abiotic soil

environment. While human alteration of the environment and its effect on C storage and release is

relatively well-documented in terrestrial ecosystems, the mechanisms governing the interactive

effects of nutrient enrichment on plant-soil-microbial relationships at terrestrial-aquatic interfaces

are relatively understudied. The availability of terminal electron acceptors in hydrologically

dynamic ecosystems are known to shape the resident microbial community and to control

biogeochemical functions (Bernhardt et al. 2017). This makes it challenging to predict microbial

responses to environmental change in ecosystem models.

Changes in the nutrient stoichiometry of both plants and surrounding soils are expected to

affect microbial community composition and function, leading to changes in elemental cycling.

Prior studies have shown that long-term nutrient addition enhances grassland plant biomass and

increases C inputs into soils (e.g., Harpole et al. 2016). These additional plant inputs also increase

CO_2 outputs via microbial respiration (Hoosbeek et al. 2004, Lange et al. 2015). If microbes are

not limited by N and P, the increase in plant-derived organic C inputs are more quickly metabolized

and soil CO_2 fluxes increase (Peralta and Wander 2008, Cotrufo et al. 2013, Castellano et al. 2015).

Plant biomass could also vary in C:N or C:P ratios, which could also cause changes in microbial

nutrient use and respiration rates. Thus, it is important to understand how changes in available

nutrients within plant litter alter microbial communities and CO_2 respiration rates as surrounding

environmental conditions also shift (Hoosbeek et al. 2004, Lange et al. 2015).

Nuanced interactions between litter chemical quality, abiotic environment, and resident

microbial composition can manifest in similar decomposition patterns (i.e., increased rates due to

increased nutrient availability). However, the degree to which microbial structure contributes to

the magnitude of litter mass loss is challenging to pinpoint since interactions among abiotic

(temperature, nutrient availability, moisture) and biotic (community structure) factors determine

decomposition rates (Kuzyakov and Blagodatskaya 2015, Meier et al. 2017, Deveau et al. 2018).

In this study, we examine the extent that litter-associated microbiomes are unique depending on

historical nutrient enrichment and how this community composition contributes to decomposition

of litter of varying chemical quality (i.e., C:N ratio).

The nutrient availability in upland conditions makes priming (using standing litter

nutrients) more pronounced as aerobic respiration is more energetically favored. In this study, we

measured the extent that microbial community change (due to nutrient enrichment) contributes to

enhanced decomposition using a model litter comparison. The Tea Bag Index (TBI) compares the

decomposition rates of two different plant litters, green and rooibos Lipton™ tea bags (Keuskamp

et al. 2013). Green tea has a measured mean C:N ratio of 12.229 while rooibos tea is measured to

have a much higher mean C:N ratio of 42.870. We used the TBI protocol to examine how long-

term fertilization affects decomposition rates in a nutrient-limited coastal plain wetland ecosystem.

We also characterized the bacterial communities (using targeted amplicon sequencing of the V4

region of the 16S rRNA gene) associated with the litter decomposition of two differing litter types

(green and rooibos teas) occurring in wetland soils exposed to fertilization or not. The extent that

community composition matters to decomposition rate depends on biotic and abiotic factors. In

this study, litter type, nutrient enrichment and hydrologic status resulted in distinct bacterial

communities.

MATERIALS AND METHODS

Experimental design of a long-term ecological experiment

Initiated in 2003, a long-term ecological experiment started at East Carolina University's

West Research Campus (WRC) (35.6298N, -77.4836W) examines the effects of mowing and

fertilization on coastal plain wetland plant and microbial communities. This study site is classified

as a jurisdictional wetland, and the plant community has been described as a mosaic of wet pine

flatwood habitat, pine savanna, and hardwood communities (Chester 2004). The soils were

characterized as fine, kaolinitic, thermic Typic Paleaquults (Coxville series), fine-loamy, siliceous,

semiactive, thermic Aeric Paleaquults (Lynchburg series), and fine-loamy, siliceous, subactive,

thermic Aquic Paleudults (Goldsboro series). These soils are moderately to poorly drained ultisols

(USDA NRCS 2019). The annual mean temperature is 16.4 °C and annual precipitation is 126 cm

(U.S. Climate Data 2019). Fertilization and mowing treatments are replicated on eight 20×30 m

blocks in a full factorial design, and the N-P-K 10-10-10 pellet fertilizer is applied 3× per year

(February, June, and October) for a total annual supplementation of 45.4 kg/ha for each nutrient.

Plots are mowed by bush-hog annually to simulate fire disturbance (Goodwillie et al. In Review).

Within each plot, annual soil and plant sampling takes place at three randomly-placed quadrats

(Fig. 1). The plant community in plots that are mowed is dominated by perennial forbs in two

major genera of the Asteraceae—*Eupatorium* and *Solidago*—and graminoid species. The relative

abundance of major grass species differs between mowed and fertilized plots, with switchcane

(*Arundinaria tecta*) most dominant in fertilized plots and broomsedge (*Andropogon virginicus*)

found almost exclusively in unfertilized plots. Sedges (e.g., *Rhychospora* and *Carex* spp.) and

species of *Juncus* are common in (wet) plots away from, but not in (dry) plots adjacent to the

drainage ditch. This hydrologic gradient is caused by a roadside drainage ditch such that four

blocks near the ditch are drier and four blocks away from the ditch are wetter (Goodwillie et al. In

Review). In 2018, volumetric soil moisture content (measured by capacitance), was >2 times

wetter in blocks away from the ditch compared to blocks adjacent to the ditch. However, this

hydrologic gradient has not yet been characterized by modeling flow according to water levels

over time. The ditch was not intentionally manipulated as part of our experimental design, but it

contributes an important ecological variable to the study (Goodwillie et al. In Review). For this

study, tea litter decomposition experiments were focused in the mowed plots only.

Soil sampling

We collected composite soil samples from all mowed/unfertilized and mowed/fertilized plots, which represented two soil cores (12 cm depth, 3.1 cm diameter) adjacent to each of three permanently installed 1 m^2 quadrats where plant community data are annually collected. Each composite bulk soil sample was passed through a 4 mm sieve and homogenized before further analysis. This bulk soil sampling occurred on November 14, 2019, about three months after the tea litter bags were collected from the field.

Soil physico-chemical analyses

We measured gravimetric soil moisture by weighing 20-30 g of field-moist soil, drying at 105 °C overnight, and then re-weighing. We calculated percent moisture as difference between field-moist and dried soils divided by the oven-dried soil weight. In addition, we measured pH of oven-dried soil by mixing a 1:1 soil:water solution and using a pH probe (Genemate-Bioexpress; Kaysville, Utah; USA). We measured total soil organic carbon and total nitrogen on finely ground dried soil using an elemental analyzer (2400 CHNS Analyzer; Perkin Elmer; Waltham, Massachusetts, USA) at the Environmental and Agricultural Testing Service laboratory (Department of Crop and Soil Sciences at North Carolina State University). Extractable inorganic N for each soil sample was measured on approximately 5 g of field moist soil. We added 45 ml of 2 M KCl to soil, shook the sample for about 1 hour, and gravity filtered. Total phosphate (PO_4^{3-}) was extracted by combining 0.1 g dried soil (ground and passed through a 500 µm sieve) with 0.5 ml of 50% w/v $Mg(NO_3)$ and ashing for 2 hours at 550 °C. Samples were hydrated with 10 mL of 1 M HCl, shaken for 16 hours at 250 RPM, and the filtered (22 µm filter). Water extractable PO_4^{3-}

was determined by combining 1 g dried soil (ground and passed through a 500 µm sieve) with

deionized water, shaken for 1 hour, and filtered (22 µm filter). Ammonium (NH_4^+), nitrate (NO_3^-),

and PO_4^{3-} ions in soil extracts were colorimetrically measured using a SmartChem 200 auto

analyzer (Unity Scientific Milford, Massachusetts, USA) at the East Carolina University

Environmental Research Laboratory.

Field experimental methods

The field protocol for this experiment was adapted from the Tea Bag Index Protocol

(Keuskamp et al. 2013) and applied within the eight-replicate fertilized and unfertilized plots of

the WRC. The experimental setup involved three bags of Lipton™ green tea and three bags of

Lipton™ rooibos tea per replicate plot (3 bags × 8 blocks × 2 treatment plots = 48 bags per each

tea bag type). In each treatment plot, we prepared six holes about 20 cm apart and 8 cm deep using

a hand trowel. We weighed green and rooibos tea bags and buried them in separate holes. Soil was

lightly packed around the tea bags, while keeping the labels visible above the soil. Tea bags were

recovered after 111 days (May 21–September 09, 2018) using hand trowels to loosen soil adjacent

to tea bag locations. Two of three green tea bags at each replicate plot were measured for mass

loss. The third green tea bag was placed into a sterile Whirl-pak bag and stored at -20 °C for tea-

associated bacterial community analysis. The rooibos tea was processed following the same

procedure. Upon returning to the lab, the tea bags for decomposition analysis were separated from

adhered soil particles, placed into individual aluminum weighing dishes, and oven-dried at 70 °C

for 48 hours. To quantify mass loss, the tea bags were reweighed following drying and compared

to their initial weight prior to soil incubation.

 Soil and tea-associated bacterial 16S rRNA sequencing

Following freezing, tea litter was removed from tea bags, and DNA was extracted from

samples using the Qiagen DNeasy PowerSoil kit. We also extracted DNA from soils using the

Qiagen DNeasy PowerSoil Kit. For each unfertilized or fertilized treatment, DNA was extracted

from green tea (15 total), rooibos tea (14 total), and bulk soil samples (16 total). Three litter

samples could not be located and were not retrieved from the field. Following extraction, sample

DNA was used as template in PCR reactions with a bacterial 515F/806RB barcoded primer set

originally developed by the Earth Microbiome Project to target the V4-V5 region of the bacterial

16S subunit of the rRNA gene (Apprill et al., 2015; Caporaso et al., 2012; Parada et al., 2016). For

each sample, three 50 µL PCR libraries were prepared by combining 35.75 µL molecular grade

water, 5 µL Amplitaq Gold 360 10x buffer, 5 µL $MgCl_2$ (25 mM), 1 µL dNTPs (40mM total,

10mM each), 0.25 µL Amplitaq Gold polymerase, 1 µL 515 forward barcoded primer (10 µM), 1

µL 806 reverse primer (10 µM), and 1 µL DNA template (10 ng/µL). Thermocycler conditions for

PCR reactions were as follows: initial denaturation (94 °C, 3 minutes); 30 cycles of 94 °C for 45

seconds, 50 °C for 30 seconds, 72 °C for 90 seconds; final elongation (72 °C, 10 minutes).

Triplicate PCR reactions were combined for each sample and cleaned according to the Axygen®

AxyPrep Magnetic Bead Purification Kits protocol (Corning Life Sciences). Following cleaning,

PCR product were quantified using Quant-iT dsDNA BR (broad-range) assay (Thermo Scientific,

Waltham, MA, USA). Libraries were pooled in equimolar concentration of 5 ng/µL after being

diluted to a concentration of 10 ng/µL. The Indiana University Center for Genomics and

Bioinformatics sequenced the pooled libraries using the Illumina MiSeq platform using paired-end

reads (Illumina Reagent Kit v2, 500 reaction kit).

Following sequencing, we processed raw sequences using a standard mothur pipeline

(v1.40.1) (Schloss et al. 2009, Kozich et al. 2013). Contigs were assembled from paired end reads

and quality trimmed using a moving average quality score (minimum score of 35 bp). Sequences

were aligned to the SILVA rRNA gene database (v.128) (Quast et al. 2013) and chimeric

sequences were removed using the VSEARCH algorithm (Rognes et al. 2016). Formation of

operational taxonomic units (OTUs) involved dividing sequences based on taxonomic class and

then binnned into OTUs with a 97% sequence similarity level. OTUs were classified using the

SILVA database (Yilmaz et al. 2014, Glöckner et al. 2017).

Statistical analyses

All statistical calculations were completed in the R environment (R v3.6.3, R Core

Development Team 2020). Decomposition was measured through mass loss of buried tea bags.

Initial and final tea bag weights were used to determine percent of loss. We ran a linear mixed

effects model with 'source', 'treatment', and 'ditch proximity' as fixed effects and 'block' as a

random effect using the lmer() function in the lmerTest package (Kuznetsova et al. 2017). For the

mass loss response variable, a linear mixed model was fit by REML and produced type II analysis

of variances tables (ANOVA) tables based on the Kenward-Roger's denominator degrees of

freedom method using the lmerModLmerTest() function. Calculations of decomposition rate (k)

and stabilization factor (S) for each tea bag were accomplished according to formulas found within

the TBI protocol. Decomposition rate, k, is a parameter representing short-term dynamics of new

input decomposition, while stabilization factor, S, is a parameter characterizing long-term carbon

storage. These values were used to draw comparisons between k and S values found within the

WRC wetland ecosystem to those of global ecosystems submitted to the Tea Bag Index protocol

(Keuskamp et al. 2013). Outcomes of these calculations were plotted on a graph, and locations of

groupings were compared to those of the larger multi-ecosystem index (Keuskamp et al. 2013).

For the bulk soil factors measured and the computed diversity (OTU richness, Shannon

Diversity, Simpson's Evenness) metrics for soil and tea litter associated bacterial communities, we

ran a series of linear mixed models. We ran linear mixed effects models with 'source', 'treatment'

and 'ditch' as fixed effects and 'block' as a random effect and fitted by the REML criterion using

the lmer() function in the lmerTest package (Kuznetsova et al. 2017). Statistical inferences for

fixed effects were calculated from type II ANOVA tables and the Kenward-Roger's degrees of

freedom method using the anova() function.

To visualize the community responses to fertilization and litter type, we used principal

coordinates analysis (PCoA) of bacterial community composition based on the Bray-Curtis

dissimilarity. We used a permutational multivariate analysis of variance (PERMANOVA) to

examine among-treatment differences in bacterial communities. We also include a comparison of

the bulk soil and tea litter associated bacterial communities (PERMANOVA, PCoA ordination)

but focused on the specific tea litter associated microbiome patterns. To identify which bacterial

species were most representative of each litter type, we ran an indicator species analysis. For the

indicator species analysis, we only included bacterial taxa with a relative abundance greater than

0.05 when summed across all plots. We performed PERMANOVA using the adonis() function in

the vegan package (Oksanen et al. 2017) and the indval() function in the indicspecies package

(Caceres and Jansen 2016).

To examine structure-function relationships between the tea-associated bacterial

community and the mass loss of the tea litter, we ran a linear regression to measure the relationship

between mass loss as a function of bacterial diversity (mass loss ~ OTU richness + Shannon

Diversity + Simpson's Evenness). In addition, we conducted a distance-based partial least squares

regression (dbplsr) measure how much variation in mass loss of litter (function) was explained by

the tea-associated bacterial community composition (structure) (mass loss ~ bacterial community).

We used the Bray-Curtis dissimilarity matrix and the generalized cross-validation estimate of the

prediction error ('gcv' method). We performed a distance-based partial least squares regression

analysis using the dbplsr() function in the dbstats and pls packages (Boj Del Val et al. 2007, Boj

et al. 2017, Mevik et al. 2019).

RESULTS

Long-term fertilization strongly influenced abiotic and biotic factors within this nutrient-

limited wetland environment. Both fertilization and ditch effects influenced soil pH, soil C:N ratio,

extractable nitrate concentrations, and water extractable phosphorus (Table S1A, Table S2B). A

subset of soil factors was distinct between fertilized and unfertilized soils in the mowed plots. Total

and water extractable phosphorus concentrations, extractable nitrate concentrations, soil C and N

content, moisture, and pH were higher in fertilized compared to unfertilized soils (Table S1B). In

addition, soil moisture, C:N ratio, and water extractable phosphorus were higher in wetter soils

away from the ditch compared to drier soils adjacent to the ditch (Table S1C). Finally, nitrate

concentrations were detectable only in wetter soils away from the ditch with higher concentrations

in fertilized soils compared to unfertilized soils while nitrate was below detection limits in all soils

near the ditch (Table S1A). These differences in fertilized compared to unfertilized soils and soil

moisture due to ditch proximity related to litter decomposition rates and altered bacterial

community structure.

To examine how litter type and nutrient addition influenced decomposition rates, we

measured mass loss of green (low C:N litter) and rooibos (high C:N litter) tea bags. Following an

111-day incubation, there was a significant litter type (source) × fertilization treatment interaction

(P=0.05) (Fig. 2, Table S2). The mass loss of green tea litter was ~24% higher than the overall

mass loss of rooibos tea litter (Fig. 2, Table S2), averaged across fertilized and unfertilized soils.

Both fertilization and proximity to drainage ditch also increased the rate of mass loss (Fig. 2, Table

S2). Across all tea types and drainage ditch proximities, fertilized soils showed an average mass

loss of ~7.5% more than that of unfertilized soils. Tea bags buried in closer proximity to the

drainage ditch had ~6% more mass loss averaged across all litter and treatment types.

We also compared the decomposition rates (k) and stabilization factors (S) from the long-

term fertilization experiment to a broader context of ecosystems (Keuskamp et al., 2013). The most

variable S and k values were measured in the wetter, fertilized plots (Fig. 3). The unfertilized

compared to fertilized samples were generally higher along the stabilization factor (S) axis.

Samples that were buried in the wetter, fertilized plots clustered along the higher end of the

stabilization factor (S), while litter buried in the drier plots (fertilized and unfertilized) had lower

estimated stabilization factor (S) (Fig. 3). Upon comparison with other ecosystems, a large number

of fertilized treatment samples grouped near the grassland-ambient ecosystem in Iceland and the

forest ecosystem in the Netherlands (Keuskamp et al. 2013). However, the unfertilized treatment

samples grouped near the peat-disturbed and peat-undisturbed ecosystems of Iceland and the peat

ecosystem of the Netherlands (Keuskamp et al. 2013). Within this coastal plain wetland system,

long-term nutrient enrichment shifted decomposition rates that were representative of grassland

and forested ecosystems (fertilized plots) or peatlands (unfertilized plots) (Fig. 3).

Fertilization effects on soil physicochemical properties and decomposition rates were

related to shifts in bulk soil and litter-associated bacterial communities. Fertilization effect, tea

type, and proximity to ditch (i.e., wetter vs. drier plots) influenced bacterial diversity in various

ways. Specifically, tea type (R^2=0.130, P=0.001) influenced bacterial community composition the

most, while the interaction of tea type and proximity to ditch affected bacterial community patterns

to a lesser degree (R^2=0.060, =0.002) (Fig. 4, Table S3) (see circle vs triangle). The proximity to

ditch (proxy for hydrology) also altered bacterial community composition (R^2=0.110, P=0.001)

(Fig. 4, Table S3) (see opened vs filled). Within tea type, fertilization treatment influenced

bacterial community composition (R^2=0.070, P=0.003) (Fig. 4, Table S3) (see green vs gray). For

the drier, ditched plots, bacterial communities associated with high C:N ratio litter (rooibos)

overlapped despite fertilization treatment, while bacterial communities were distinct between

fertilization treatments in the wetter plots (Fig. 4). When the bulk soil and tea-associated bacterial

communities were analyzed together, bulk soil or tea most strongly influenced bacterial

community composition (source: R^2=0.490, P=0.001) (Fig. S1, Table S4). In addition to these

observed patterns, OTU diversity was generally higher in fertilized compared to unfertilized plots,

especially in the drier plots associated with the drainage ditch (Fig. 5, Table S5). Specifically, OTU

richness values were highest in bulk soil compared to tea types (Fig. 5A, Table S5A), and Shannon

diversity values were highest in bulk soil compared to tea type in fertilized plots (Fig. 5B, Table

S5B). Lastly, bacterial evenness was similar across fertilization treatment, source type, and

proximity to ditch (Fig. 5C, Table S5C).

To further examine bacterial associations with litter decomposition, we used indicator

species analysis to identify a subset of bacterial taxa that represented each of the bulk soil or tea

associated microbiomes. The unclassified OTU within the class Spartobacteria and another OTU

within the order Acidobacteria Gp1 represented the unfertilized bulk soil bacterial community,

while an unclassified OTU within the order Solirubrobacterales represented bulk soils in the

fertilized, dry plots. In wetter plots, an unclassified OTU within the order Rhodospirillales

represented unfertilized plots while unclassified taxa within the orders Acidobacteria Gp1, Gp2,

and Gp6 signified fertilized bulk soils (Tables S6, S7). In addition, the unclassified OTU within

the Acidobacteria Gp3 and *Dongia* spp. represented the green tea-associated (low C:N litter)

bacterial communities in the unfertilized, dry plots while *Conexibacter* spp. were represented in

the fertilized, dry plots. In the wetter plots, *Phenylobacterium* spp. represented green tea-

associated bacterial communities in unfertilized plots and *Legionella* spp. in fertilized plots. The

OTUs *Acidisoma* spp. in unfertilized plots and *Dyella* spp. in fertilized characterized rooibos tea

associated bacterial communities in the dry plots. Lastly, the OTU *Lacibacterium* spp. represented

mowed plots, while *Dokdonella* spp. and an unclassified OTU within the genus *Microbacteriaceae*

represented rooibos tea associated communities in the wet plots (Tables S6, S7).

When examined together, the link between bacterial community structure and

decomposition function was relatively weak. No relationship between litter mass loss and bacterial

diversity was detected (R^2_{adj} = -0.038, P = 0.574). However, when the total bacterial community

was considered, the tea-associated bacterial community accounted for ~64 % of the variation in

litter mass loss (dbplsr, Component 1 R^2_{adj} = 63.56, Component 2 R^2_{adj} = 87.09, Component 3 R^2_{adj}

= 93.96) (Table 1).

DISCUSSION

Litter composition, fertilization, and ditch effects on soil moisture influenced bacterial

community structure-function relationships in unexpected ways. In this study, increases in litter

mass loss were greater in fertilized compared to unfertilized soils, especially in drier versus wetter

plots. Surprisingly, distinct microbial communities associated with different litter chemical

qualities. Litter mass loss following the field incubation indicates that similar bacterial

communities are capable of decomposing higher C:N ratio litter (rooibos tea) more quickly in

fertilized compared to unfertilized plots, particularly in drier compared to wetter plots. However,

lower C:N ratio litter (green tea) decomposition rates were similar among fertilized and

unfertilized plots, although slightly higher in drier compared to wetter plots but were represented

by distinct bacterial communities. As expected, the low C:N ratio litter provides a N source to

microbes regardless of external soil nutrient conditions, whereas the high C:N ratio litter is reliant

on N from the fertilized soil in order to support increased rates of litter decomposition (Duddigan

et al. 2020). These results suggest that shifts in microbial community composition are partly due

to differences in litter chemical quality and nitrate availability in soils.

The comparative results from the TBI-based decomposition experiment indicate that the

tea litter bags buried in the unfertilized treatment had decomposition rates (k) and C stabilization

factor (S) values similar to that of peatland ecosystems (Keuskamp et al. 2013). Lower

decomposition rates occurred in unfertilized plots (i.e., the ambient state of this nutrient-poor

habitat). When looking specifically at high C:N ratio litter, less mass was lost to decomposition.

This provides some evidence that C storage potential within these plots resembles that of peatlands

(Hill et al. 2018). Wetland ecosystems that are disconnected and isolated from pulses of nutrients

due to natural processes or agricultural or urban runoff are more commonly nutrient limited by N

and P (Vitousek et al. 2010). The plant species that are adapted to these low-nutrient ecosystems

can maintain positive population growth and contribute to organic C additions to soils. Further,

flooded environments, such as wetlands, are also known to support anaerobic microbial processes,

which result in slower rates of decomposition (Collins et al. 2015). Taken together, low nutrient

environments can be sites of high plant biodiversity leading to organic C inputs and balanced with

slower decomposition rates, which contribute to long-term C storage in soils (Hooper et al. 2005,

Kleber et al. 2011). These results reflect the coastal plain wetland environment in this study, which

offers some evidence that wetland environments store a disproportionate amount of C compared

to other ecosystems (Sutfin et al. 2016, Nahlik and Fennessy 2016). However, this C storage

capacity can be disrupted by nutrient enrichment. The fertilized plots, however, had k and S values

that grouped closer towards grassland and forest ecosystems (Riggs et al. 2015), which have lower

C storage potential because more available nutrients and oxic environments support aerobic

respiration and higher decomposition rates. Results from this study provide support that nutrient

enrichment can have a lasting influence on plant-soil-microbe interactions that affect C storage

potential of wetland ecosystems (Lambers et al. 2009, Allison et al. 2014, Hartman et al. 2017).

Patterns in bacterial community composition (i.e., beta diversity) but not the combined

OTU richness, Shannon diversity, Simpson's evenness diversity metrics, significantly explained

decomposition rates. Further, litter composition followed by the fertilization and ditch effects

determined bacterial community composition. A trend of higher decomposition rates was

associated with high bacterial diversity, especially in the fertilized compared to unfertilized plots.

Higher diversity has been associated with higher decomposition rates across different ecosystems

(Lange et al. 2015, Trivedi et al. 2016). Shifts in plant and microbial communities can fuel

decomposition rates and ultimately C losses from historically low nutrient ecosystems that

experience atmospheric nutrient deposition (Sardans and Peñuelas 2012, Wieder et al. 2015).

When bulk and tea-associated microbiomes were considered together, the bulk soil

bacterial communities were very distinct from the tea-associated bacterial communities. Within

the bulk soil, bacterial communities were distinct across fertilized and unfertilized dry plots, and

indicator taxa were phylogenetically distinct at the phylum level (Proteobacteria vs.

Acidobacteria). While results revealed similarity in bulk soil bacterial composition in the fertilized

and unfertilized wet plots, the indicator taxa also represented distinct phyla (unfertilized:

Verrucomicrobia + Acidobacteria, fertilized: Actinobacteria). In contrast, the indicator taxa for

tea-associated bacterial communities that overlapped in composition were similar at the phylum

level. Specifically, similar bacterial assemblages of rooibos tea-associated (high C:N litter)

microbiomes in the fertilized and unfertilized, dry plots were associated with Proteobacteria (Table

S6, unfertilized: *Lacibacterium spp.*, fertilized: *Dokdonella spp.*). Indicator bacterial taxa

representing high C:N ratio litter-associated microbiomes in fertilized, dry plots were *Dokdonella*

spp. (Proteobacteria phylum) and unclassified Microbacteriaceae (Actinobacteria phylum). The

genus *Dokdonella* has been involved in heterotrophic denitrification (Figueroa-González et al.

2016, Palma et al. 2018), while members of the family Microbacteriaceae have been

predominantly found in terrestrial and aquatic ecosystems with putative functions related to plant

pathogenicity (Glöckner et al. 2000, Young 2008, Lory 2014). Indicator taxa within the order

Alphaproteobacteria represented the similar bacterial communities of the high C:N litter-

associated microbiomes in unfertilized (indicator OTU *Acidisoma spp.*), wet plots and low C:N

litter-associated microbiomes in unfertilized, dry plots (indicator OTU *Phenylobacterium spp.*). In

addition, the community similarity between low C:N litter-associated microbiomes in unfertilized,

wet plots and high C:N litter-associated microbiomes in fertilized, wet plots were represented by

taxa within different Proteobacterial classes. Specifically, *Dongia spp.* (within

Alphaproteobacteria class) and *Dyella spp.* (within Gammaproteobacteria class) are similar in at

least some life history traits. Representative isolates were cultured from soils (*Dyella*) and wetland

sediments (*Dongia*) and were found to be aerobic, Gram-negative, and motile (Xie and Yokota

2005, Baik et al. 2013). Further, these assemblages also overlapped with green tea-associated

microbiomes in the fertilized, dry plots, which were represented by *Legionella spp.* (within class

Gammaproteobacteria). Legionellae are found in aquatic and other moist environments with their

free-living protozoan hosts (Barker and Brown 1994). In the wetland soil environment, they can

persist but are unlikely existing under optimal conditions.

419   The interaction between source (litter type) and fertilization treatment influenced mass loss

of litter but hydrologic setting and soil nitrate altered litter-associated microbiomes to varying

degrees. The ditch-derived hydrologic differences at this wetland resulted in different structure-

function relationships for high C:N ratio litter-associated microbiomes only. Under the relatively

dry plots (i.e., near drainage ditch), microbiomes associated with high C:N ratio litter were similar,

but mass loss of litter was higher in fertilized compared to unfertilized plots. In all other instances,

distinct, fertilized microbiomes were associated with higher decomposition rates compared to

unfertilized plots. In the case that community structure is the same and decomposition rates

increase, this pattern provides some evidence of more severe N limitation in drier compared to

wetter plots, where soil nitrate concentrations are below detection limits (Table S1B). This result

was only observed under high C:N ratio litter context since the low C:N litter provided much

needed N to litter-associated microbes. While soil ammonium levels are similar in this case, it is

possible that N mineralization is occurring without limitation, but nitrification processes could be

slowed due to low soil pH, resulting in differences in extractable soil nitrate but not ammonium

concentrations (Hinckley et al. 2019).

434   In this study, nutrient enrichment and hydrologic status resulted in distinct bacterial

communities associated with litter and bulk soils. Drier conditions and nutrient enrichment

increased decomposition rates until N limitation constrains microbial community structure. This

study provided the opportunity to compare the extent that community composition matters to

decomposition rate – using model litter. In these nutrient poor wetland soils undergoing nutrient

enrichment, the fertilization increased bacterial diversity the potential for increased C losses

through decomposition is observed and is expected to increase as wetlands are drained and

fertilized.

ACKNOWLEDGMENTS

We thank Allison Fisk, Emma Richards, Tom Vogel, John Stiller, Suelen Tullio, and Adam Gold

for laboratory and field assistance. We thank John Gill and the East Carolina University grounds

crew for their tireless efforts in maintaining the long-term ecological experiment. This work was

supported by the East Carolina University Undergraduate Research and Creative Activity Award

to MEK and the National Science Foundation (GRFP to RBB, DEB 1845845 to ALP). All code

and data used in this study can be found in a public GitHub repository

(https://github.com/PeraltaLab/WRC18_RhizoTeaDecomp) and the NCBI SRA (BioProject).

LITERATURE CITED

Allison, S. D., S. S. Chacon, and D. P. German. 2014. Substrate concentration constraints on

microbial decomposition. Soil Biology and Biochemistry 79:43–49.

Apprill A, McNally S, Parsons R, and Weber L. 2015. Minor revision to V4 region SSU rRNA

806R gene primer greatly increases detection of SAR11 bacterioplankton. Aquatic

Microbial Ecology 75:129–137.

Baik, K. S., Y. M. Hwang, J.-S. Choi, J. Kwon, and C. N. Seong. 2013. Dongia rigui sp. nov.,

isolated from freshwater of a large wetland in Korea. Antonie van Leeuwenhoek

104:1143–1150.

Barker, J., and M. R. W. Brown. 1994. Trojan Horses of the microbial world: protozoa and the

survival of bacterial pathogens in the environment. Microbiology 140:1253–1259.

Bernhardt, E. S., J. R. Blaszczak, C. D. Ficken, M. L. Fork, K. E. Kaiser, and E. C. Seybold.

2017. Control Points in Ecosystems: Moving Beyond the Hot Spot Hot Moment Concept.

Ecosystems 20:665–682.

Boj Del Val, E., M. M. C. Bielsa, and J. Fortiana. 2007. Selection of Predictors in Distance-

Based Regression. Communications in Statistics - Simulation and Computation 36:87–

98.

Boj, E., A. Caballe, P. Delicado, and J. Fortiana. 2017. dbstats: Distance-Based Statistics.

Caceres, M. D., and F. Jansen. 2016. indicspecies: Relationship Between Species and Groups of

Sites.

Caporaso, J. G., C. L. Lauber, W. A. Walters, D. Berg-Lyons, J. Huntley, N. Fierer, S. M.

Owens, J. Betley, L. Fraser, M. Bauer, N. Gormley, J. A. Gilbert, G. Smith, and R.

Knight. 2012. Ultra-high-throughput microbial community analysis on the Illumina

HiSeq and MiSeq platforms. The ISME Journal 6:1621–1624.

Castellano, M. J., K. E. Mueller, D. C. Olk, J. E. Sawyer, and J. Six. 2015. Integrating plant litter

quality, soil organic matter stabilization, and the carbon saturation concept. Global

Change Biology 21:3200–3209.

Cherif, M., and M. Loreau. 2009. When microbes and consumers determine the limiting nutrient

of autotrophs: a theoretical analysis. Proceedings of the Royal Society B: Biological

Sciences 276:487–497.

Chester, R.E. 2004. Floristic assessment of a wet mineral flat at the East Carolina University

West Research Campus and investigation of influential, human-mediated factors on the

plant community. Master's Thesis, East Carolina University, Greenville, NC.

Collins, D. P., W. C. Conway, C. D. Mason, and J. W. Gunnels. 2015. Decomposition of three

common moist-soil managed wetland plant species. Journal of Fish and Wildlife

Management 6:102–111.

Cotrufo, M. F., M. D. Wallenstein, C. M. Boot, K. Denef, and E. Paul. 2013. The Microbial

Efficiency-Matrix Stabilization (MEMS) framework integrates plant litter decomposition

with soil organic matter stabilization: do labile plant inputs form stable soil organic

matter? Global Change Biology 19:988–995.

Deveau, A., G. Bonito, J. Uehling, M. Paoletti, M. Becker, S. Bindschedler, S. Hacquard, V.

Hervé, J. Labbé, O. A. Lastovetsky, S. Mieszkin, L. J. Millet, B. Vajna, P. Junier, P.

Bonfante, B. P. Krom, S. Olsson, J. D. van Elsas, and L. Y. Wick. 2018. Bacterial–fungal

interactions: ecology, mechanisms and challenges. FEMS Microbiology Reviews

42:335–352.

Duddigan, S., L. J. Shaw, P. D. Alexander, and C. D. Collins. 2020. Chemical underpinning of

the Tea Bag Index: an examination of the decomposition of tea leaves. Research Article,

Hindawi. https://www.hindawi.com/journals/aess/2020/6085180/.

Figueroa-González, I., G. Quijano, I. Laguna, R. Muñoz, and P. A. García-Encina. 2016. A

fundamental study on biological removal of N2O in the presence of oxygen.

Chemosphere 158:9–16.

Fowler, D., M. Coyle, U. Skiba, M. A. Sutton, J. N. Cape, S. Reis, L. J. Sheppard, A. Jenkins, B.

Grizzetti, J. N. Galloway, P. Vitousek, A. Leach, A. F. Bouwman, K. Butterbach-Bahl, F.

Dentener, D. Stevenson, M. Amann, and M. Voss. 2013. The global nitrogen cycle in the

twenty-first century. Phil. Trans. R. Soc. B 368:20130164.

Glöckner, F. O., P. Yilmaz, C. Quast, J. Gerken, A. Beccati, A. Ciuprina, G. Bruns, P. Yarza, J.

Peplies, R. Westram, and W. Ludwig. 2017. 25 years of serving the community with

ribosomal RNA gene reference databases and tools. Journal of Biotechnology 261:169–

176.

Glöckner, F. O., E. Zaichikov, N. Belkova, L. Denissova, J. Pernthaler, A. Pernthaler, and R.

Amann. 2000. Comparative 16S rRNA analysis of lake bacterioplankton reveals globally

distributed phylogenetic clusters including an abundant group of Actinobacteria. Applied

and Environmental Microbiology 66:5053–5065.

Goodwillie, C., M. W. McCoy, and A. L. Peralta. In Review. Long-term fertilization, mowing,

and ditch drainage interact in wetland plant communities.

Guignard, M. S., A. R. Leitch, C. Acquisti, C. Eizaguirre, J. J. Elser, D. O. Hessen, P. D.

Jeyasingh, M. Neiman, A. E. Richardson, P. S. Soltis, D. E. Soltis, C. J. Stevens, M.

Trimmer, L. J. Weider, G. Woodward, and I. J. Leitch. 2017. Impacts of Nitrogen and

Phosphorus: From Genomes to Natural Ecosystems and Agriculture. Frontiers in Ecology

and Evolution 5.

Harpole, W. S., L. L. Sullivan, E. M. Lind, J. Firn, P. B. Adler, E. T. Borer, J. Chase, P. A. Fay,

Y. Hautier, H. Hillebrand, A. S. MacDougall, E. W. Seabloom, R. Williams, J. D.

Bakker, M. W. Cadotte, E. J. Chaneton, C. Chu, E. E. Cleland, C. D'Antonio, K. F.

Davies, D. S. Gruner, N. Hagenah, K. Kirkman, J. M. H. Knops, K. J. L. Pierre, R. L.

McCulley, J. L. Moore, J. W. Morgan, S. M. Prober, A. C. Risch, M. Schuetz, C. J.

Stevens, and P. D. Wragg. 2016. Addition of multiple limiting resources reduces

grassland diversity. Nature 537:93–96.

Hartman, W. H., R. Ye, W. R. Horwath, and S. G. Tringe. 2017. A genomic perspective on

stoichiometric regulation of soil carbon cycling. The ISME Journal 11:2652–2665.

Hill, B. H., C. M. Elonen, A. T. Herlihy, T. M. Jicha, and G. Serenbetz. 2018. Microbial

ecoenzyme stoichiometry, nutrient limitation, and organic matter decomposition in

wetlands of the conterminous United States. Wetlands Ecology and Management 26:425–

439.

Hinckley, B. R., J. R. Etheridge, and A. L. Peralta. 2019. Wetland Conditions Differentially

Influence Nitrogen Processing within Waterfowl Impoundments. Wetlands.

Hooper, D. U., F. S. Chapin, J. J. Ewel, A. Hector, P. Inchausti, S. Lavorel, J. H. Lawton, D. M.

Lodge, M. Loreau, S. Naeem, B. Schmid, H. Setälä, A. J. Symstad, J. Vandermeer, and

D. A. Wardle. 2005. Effects of biodiversity on ecosystem functioning: a consensus of

current knowledge. Ecological Monographs 75:3–35.

Hoosbeek, M. R., M. Lukac, D. van Dam, D. L. Godbold, E. J. Velthorst, F. A. Biondi, A.

Peressotti, M. F. Cotrufo, P. de Angelis, and G. Scarascia-Mugnozza. 2004. More new

carbon in the mineral soil of a poplar plantation under Free Air Carbon Enrichment

(POPFACE): Cause of increased priming effect? Global Biogeochemical Cycles 18.

Keuskamp, J. A., B. J. Dingemans, T. Lehtinen, J. M. Sarneel, and M. M. Hefting. 2013. Tea

Bag Index: a novel approach to collect uniform decomposition data across ecosystems.

Methods in Ecology and Evolution 4:1070–1075.

Kleber, M., P. S. Nico, A. Plante, T. Filley, M. Kramer, C. Swanston, and P. Sollins. 2011. Old

and stable soil organic matter is not necessarily chemically recalcitrant: implications for

modeling concepts and temperature sensitivity. Global Change Biology 17:1097–1107.

Kozich, J. J., S. L. Westcott, N. T. Baxter, S. K. Highlander, and P. D. Schloss. 2013.

Development of a dual-index sequencing strategy and curation pipeline for analyzing

amplicon sequence data on the MiSeq Illumina sequencing platform. Applied and

Environmental Microbiology 79:5112–20.

Kuzyakov, Y. 2010. Priming effects: Interactions between living and dead organic matter. Soil

Biology and Biochemistry 42:1363–1371.

Kuzyakov, Y., and E. Blagodatskaya. 2015. Microbial hotspots and hot moments in soil:

Concept & review. Soil Biology and Biochemistry 83:184–199.

Lambers, H., C. Mougel, B. Jaillard, and P. Hinsinger. 2009. Plant-microbe-soil interactions in

the rhizosphere: an evolutionary perspective. Plant and Soil 321:83–115.

Lange, M., N. Eisenhauer, C. A. Sierra, H. Bessler, C. Engels, R. I. Griffiths, P. G. Mellado-

Vázquez, A. A. Malik, J. Roy, S. Scheu, S. Steinbeiss, B. C. Thomson, S. E. Trumbore,

and G. Gleixner. 2015. Plant diversity increases soil microbial activity and soil carbon

storage. Nature Communications 6:6707.

Leff, J. W., S. E. Jones, S. M. Prober, A. Barberán, E. T. Borer, J. L. Firn, W. S. Harpole, S. E.

Hobbie, K. S. Hofmockel, J. M. H. Knops, R. L. McCulley, K. L. Pierre, A. C. Risch, E.

W. Seabloom, M. Schütz, C. Steenbock, C. J. Stevens, and N. Fierer. 2015. Consistent

responses of soil microbial communities to elevated nutrient inputs in grasslands across

the globe. Proceedings of the National Academy of Sciences 112:10967–10972.

Lory, S. 2014. The Family Mycobacteriaceae. Pages 571–575 *in* E. Rosenberg, E. F. DeLong, S.

Lory, E. Stackebrandt, and F. Thompson, editors. The Prokaryotes: Actinobacteria.

Springer Berlin Heidelberg, Berlin, Heidelberg.

Meier, I. C., A. C. Finzi, and R. P. Phillips. 2017. Root exudates increase N availability by

stimulating microbial turnover of fast-cycling N pools. Soil Biology and Biochemistry

106:119–128.

Mevik, B.-H., R. Wehrens, K. H. Liland, and P. Hiemstra. 2019. pls: Partial Least Squares and

Principal Component Regression.

Nahlik, A. M., and M. S. Fennessy. 2016. Carbon storage in US wetlands. Nature

Communications 7:13835.

Oksanen, J., F. G. Blanchet, M. Friendly, R. Kindt, P. Legendre, D. McGlinn, P. R. Minchin, R.

B. O'Hara, G. L. Simpson, P. Solymos, M. H. H. Stevens, E. Szoecs, and H. Wagner.

2017. vegan: Community Ecology Package.

Palma, T. L., M. N. Donaldben, M. C. Costa, and J. D. Carlier. 2018. Putative Role of

Flavobacterium, Dokdonella and Methylophilus Strains in Paracetamol Biodegradation.

Water, Air, & Soil Pollution 229:200.

Parada, A. E., D. M. Needham, and J. A. Fuhrman. 2016. Every base matters: assessing small

subunit rRNA primers for marine microbiomes with mock communities, time series and

global field samples. Environmental Microbiology 18:1403–1414.

Peralta, A. L., and M. M. Wander. 2008. Soil organic matter dynamics under soybean exposed to

elevated [CO2]. Plant and Soil 303:69–81.

Riggs, C. E., S. E. Hobbie, E. M. Bach, K. S. Hofmockel, and C. E. Kazanski. 2015. Nitrogen

addition changes grassland soil organic matter decomposition. Biogeochemistry

125:203–219.

Rognes, T., T. Flouri, B. Nichols, C. Quince, and F. Mahé. 2016. VSEARCH: a versatile open

source tool for metagenomics. PeerJ 4:e2584.

Sardans, J., and J. Peñuelas. 2012. The Role of Plants in the Effects of Global Change on

Nutrient Availability and Stoichiometry in the Plant-Soil System. Plant Physiology

160:1741–1761.

Schloss, P. D., S. L. Westcott, T. Ryabin, J. R. Hall, M. Hartmann, E. B. Hollister, R. A.

Lesniewski, B. B. Oakley, D. H. Parks, C. J. Robinson, J. W. Sahl, B. Stres, G. G.

Thallinger, D. J. V. Horn, and C. F. Weber. 2009. Introducing Mothur: open-source,

platform-independent, community-supported software for describing and comparing

microbial communities. Applied and Environmental Microbiology 75:7537–7541.

Sutfin, N. A., E. E. Wohl, and K. A. Dwire. 2016. Banking carbon: a review of organic carbon

storage and physical factors influencing retention in floodplains and riparian ecosystems.

Earth Surface Processes and Landforms 41:38–60.

Trivedi, P., M. Delgado-Baquerizo, C. Trivedi, H. Hu, I. C. Anderson, T. C. Jeffries, J. Zhou,

and B. K. Singh. 2016. Microbial regulation of the soil carbon cycle: evidence from

gene–enzyme relationships. The ISME Journal 10:2593–2604.

U.S. Climate Data. 2019. Climate data for Greenville, North Carolina (-77.3984, 35.64) Average

weather Greenville, NC - 27834 - 1981-2010 normals.

https://www.usclimatedata.com/climate/greenville/north-carolina/united-states/usnc0281

Vitousek, P. M., S. Porder, B. Z. Houlton, and O. A. Chadwick. 2010. Terrestrial phosphorus

limitation: mechanisms, implications, and nitrogen–phosphorus interactions. Ecological

Applications 20:5–15.

Wieder, W. R., C. C. Cleveland, W. K. Smith, and K. Todd-Brown. 2015. Future productivity

and carbon storage limited by terrestrial nutrient availability. Nature Geoscience 8:441–

444.

Xie, C.-H., and A. Yokota. 2005. Dyella japonica gen. nov., sp. nov., a γ-proteobacterium

isolated from soil. International Journal of Systematic and Evolutionary Microbiology

55:753–756.

Yilmaz, P., L. W. Parfrey, P. Yarza, J. Gerken, E. Pruesse, C. Quast, T. Schweer, J. Peplies, W.

Ludwig, and F. O. Glöckner. 2014. The SILVA and "All-species Living Tree Project

(LTP)" taxonomic frameworks. Nucleic Acids Research 42:D643–D648.

Young, J. M. 2008. An overview of bacterial nomenclature with special reference to plant

pathogens. Systematic and Applied Microbiology 31:405–424.

Tables

Table 1. Summary of distance-based partial least squares regression (dbplsr) representing how much variation in decomposition rate is explained (%) by each component (Comp) derived from a tea-associated bacterial community Bray-Curtis distance matrix.

	Comp 1	Comp 2	Comp 3	Comp 4	Comp 5	Comp 6
R^2	64.96	88.08	94.65	98.12	99.32	99.77
adjusted R^2	63.56	87.09	93.96	97.78	99.16	99.70
gvar	31.50	41.62	50.10	56.93	61.21	66.39
crit	2.47	0.91	0.44	0.17	0.07	0.03

gvar = total weighted geometric variability; crit = value of criterion defined in method

658   **Figure 1**

Figure 1. Experimental design of a long-term ecological experiment to test the effects of fertilizer and disturbance by mowing on plant and microbial communities at East Carolina University's West Research Campus (WRC), Greenville, NC, USA. The decomposition protocol for this experiment was adapted from the Tea Bag Index Protocol (Keuskamp et al. 2013)(Keuskamp et al., 2013) and applied within the eight-replicate fertilized and unfertilized plots of the WRC. In one quadrat per replicate plot, three bags of Lipton™ green tea and three bags of Lipton™ rooibos tea were buried and retrieved after 111 days. Bulk soil was sample as a composite sample representing two soil cores from three permanent quadrats per plot.

669   **Figure 2.**

671   **Figure 2**. Boxplots representing cumulative mass loss of green (low C:N ratio litter) and rooibos

672   (high C:N ratio litter) tea in fertilized (green) and unfertilized (gray) plots. Symbols represent

673   individual data points.

    Figure 3

683   **Figure 3.** Initial decomposition rate *k* and stabilization factor *S* for different tea bags within the

684   long-term fertilization experiment. Tea bags from unfertilized plots are indicated in grey, while

685   tea bags from fertilized plots are indicated in green. Open symbols represented drier plots that were

686   closer to the drainage ditch, while closed symbols represented wetter plots that were further from

687   the ditch.

Figure 4

Figure 4. Ordination based on a Principal Coordinates Analysis depicting bacterial community

composition according to tea type. Symbols are colored according to fertilization treatment (gray

= unfertilized, green = fertilized) and tea source (circles = low C:N ratio green tea, triangles = high

C:N ratio rooibos tea) at drier mowed plots situated close to the drainage ditch (open symbols)

compared to wetter mowed plots (closed symbols).

Figure 5

Figure 5. Boxplots depicting of bacterial diversity metrics (OTU richness, Shannon Diversity

Index H′, and Simpson's Evenness) associated with source (bulk soil, low C:N ratio green tea,

high C:N ratio rooibos tea). Boxplots and symbols are colored according to fertilization treatment

(gray = unfertilized, green = fertilized) at drier mowed plots situated close to the drainage ditch

compared to wetter mowed plots. Symbols represent data points for individual plot samples.