# Development and application of eDNA-based tools for the conservation of white-clawed crayfish

**Christopher R. Troth[1,2*], Alfred Burian[1], Quentin Mauvisseau[1,2], Mark Bulling[1], Jen Nightingale[3,4], Christophe Mauvisseau[5], Michael J. Sweet[1].**

[1] Aquatic Research Facility, Environmental Sustainability Research Centre, University of Derby, Derby, DE22 1GB, UK.

[2] SureScreen Scientifics Ltd, Morley Retreat, Church Lane, Morley, DE7 6DE, UK.

[3] Bristol Zoological Society, Clifton, Bristol, UK.

[4] School of Biological Sciences, University of Bristol, Bristol, UK.

[5] Fédération de Pêche et de Protection du Milieu Aquatique du Loir-et-Cher, 11 Rue Robert Nau, 41000 Blois, France.

*Corresponding authors: christopher.troth@sky.com; m.sweet@derby.ac.uk

**Key words:** environmental DNA, sampling methodology, species interactions, white-clawed crayfish.

**Abstract**

1. eDNA-based methods represent non-invasive and cost-effective approaches for species monitoring and

their application as a conservation tool has rapidly increased within the last decade. Currently, they are

primarily used to determine the presence/absence of invasive, endangered or commercially important

species, but they also hold potential to contribute to an improved understanding of the ecological

interactions that drive species distribution. However, this next step of eDNA-based applications requires

a thorough method development.

2. We developed an eDNA assay for the white-clawed crayfish (*Austropotamobius pallipes*), a flagship

species of conservation in the UK. Multiple subsequent *in-situ* and *ex-situ* validation tests aimed at

improving method performance allowed us to apply eDNA-based surveys to evaluate interactions

between white-clawed crayfish, crayfish plague and invasive signal crayfish.

3. The assay performed well in terms of specificity (no detection of non-target DNA) and sensitivity, which

was higher than more established traditional species survey methods. Quantification of species biomass

was, however, less reliable.

4. Comparison of eDNA sampling methods (precipitation vs. various filtration approaches) revealed that

optimal sampling method differed across environments and might depend on inhibitor concentrations.

5. Finally, we applied our methodology together with established assays for crayfish plague and the

invasive signal crayfish and demonstrated their significant interactions in a UK river system.

6. Our analysis highlights the importance of thorough methodological development of eDNA-based

assays. Only a critical evaluation of methodological strengths and weaknesses will allow us to capitalise

on the full potential of eDNA-based methods and use them as decision support tools in environmental

monitoring and conservation practices.

## Introduction

Since its initial conception as a method for aquatic ecological surveys (Ficetola et al., 2008), the use of environmental DNA (eDNA) based methods are rapidly increasing in popularity (Biggs et al., 2015; Jerde et al., 2013; . Advantages such as higher cost effectiveness compared to established traditional species survey methods and the non-invasive sampling approach have been emphasised (Goldberg et al., 2016; Huver et al., 2015). Nevertheless, the true potential of eDNA-based methods is just starting to be realized. Currently, eDNA-based tools are mostly used for simple presence/absence surveys, while they could also be used to study ecological interactions that determine species distribution and the conservation status of target species. However, such advances in application require careful method evaluations and the improvement of sampling approaches to increase reliability of detection and prevent false conclusions.

In the case of species-specific eDNA assays, the design and validation of the assay represents a critical first step (Geerts et al., 2018). During assay design, it is fundamental to ensure a high target specificity (Bylemans et al., 2018) by selecting suitable amplicon lengths, *in-silico* simulations and testing against amplification of non-target DNA. *In-vitro* laboratory validation should then ascertain that the assay complies with established guidelines (Bustin et al., 2009) and that limits of detection (LOD) and quantification (LOQ) are established. Further, field comparisons with established traditional survey methods are recommended to complement reliability assessments (Smart et al., 2015). However, both traditional survey approaches and eDNA-based methods are affected by various error sources potentially creating inconsistencies that require careful interpretation (Hinlo et al., 2017a).

Further, the reliability of eDNA-based tools is strongly influenced by sampling methodology (Hinlo et al., 2017b). Currently, precipitation and various filtration methods are applied to concentrate eDNA during field sampling. Filtration approaches have the advantage of collating eDNA from larger volumes of water

compared to precipitation-based methods (Mächler et al., 2016). However, they can also incorporate the

risk of missing particles below the filter pore size (Minamoto et al., 2016) and may lead to higher

concentrations of inhibitors preventing targeted eDNA amplification (Mauvisseau et al., 2019a). Previous

method comparisons have come to contrasting recommendations for difference species (Deiner *et al.*,

2015; Dickie *et al.*, 2018). Additionally, even for the same species the 'optimal' method for collecting eDNA

may vary between lentic (i.e. ponds or lakes) and lotic (i.e. rivers and canals) systems (Geerts et al., 2018)

and therefore careful method comparisons are recommended (Deiner et al., 2015).

In this study, we target the white-clawed crayfish, *Austropotamobius pallipes* (Lereboullet, 1858), an

endangered and important umbrella species in the UK and Western Europe (Füreder et al., 2010). Range

reduction of white-clawed crayfish began in the 1860s, with declines rapidly accelerating in the UK after

the introduction of invasive non-native signal crayfish (*Pacifastacus leniusculus*, Dana, 1852) from North

America in the 1970s (Holdich et al., 2009). Moreover, the spread of crayfish plague *Aphanomyces astaci*

(Schikora 1906), an oomycete pathogen carried by signal crayfish, has greatly exacerbated the negative

impact of invasive competitors, pollution and habitat degradation (Holdich et al., 2009). Despite its

legislative protection (EU Habitats Directive), white-clawed crayfish has continued to decline by as much

as 50-80% between 2000 and 2010 (Füreder et al., 2010). Due (at least in part) to the now rarity of the

native species, traditional survey methods are having unsatisfactory success in monitoring populations

(Gladman et al., 2010; Holdich and Reeve, 1991), highlighting the urgent need to develop new survey

tools.

Consequently, the aim of this study was to develop a highly reliable eDNA assay for the detection of white-

clawed crayfish*,* that allows the assessment of interactions with competing species and pathogens which

threaten their survival. We designed a primer and probe set for the amplification of white-clawed crayfish

DNA and critically evaluated the sensitivity and specificity of the assay through extensive *in-silico, in-vitro*

and *in-situ* tests. Moreover, we evaluated the impact of different sampling methodologies on the

reliability of the assay in mesocosm experiments and field tests implemented in different habitat types.

Finally, this allowed us to assess in a UK river system the relationship between white-clawed crayfish,

signal crayfish and crayfish plague, demonstrating the applicability of eDNA-based approaches for in-

depth ecological investigations and ecosystem management.

**Materials and Methods**

***Primer design and in-silico tests***

Primer/probe design and validation followed guidelines established by MacDonald and Sarre (2017) aimed

for assay development of species-specific eDNA methods. The primers and probe, targeting the

Cytochrome C Oxidase Subunit 1 (COI) mitochondrial gene of white-clawed crayfish, were designed *in-*

*silico* using Geneious Pro R10 (Kearse et al., 2012). The forward primer WC2302F 5' -

GCTGGGATAGTAGGGACTTCTTT - 3', reverse primer WC2302R 5' – CATGGGCGGTAACCACTAC - 3' and

probe WC2302P 5' - 6-FAM-CTGCCCGGCTGCCCTAATTC-BHQ-1 -3' amplified a 109bp fragment. To ensure

specificity, *in-silico* tests were run against published sequences of closely related and/or co-occurring

crayfish species.

***In-vitro validation***

The specificity of the assay was further tested *in-vitro* against extracted DNA of either taxonomically

similar, or co-occuring crayfish species. These included; *Faxonius limosus* (Rafinesque, 1817), *P.*

*leniusculus*, *Astacus astacus* (Linnaeus, 1758), *Astacus leptodactylus* (Eschscholtz, 1823), *Procambarus*

*clarkii* (Girard, 1852), and *Procambarus virginalis* (Lyko, 2017). DNA was extracted from crayfish tissues

using the Qiagen DNeasy® Blood & Tissue kit, following manufacturers' instructions. PCRs were performed

using the primers and methods from Folmer et al. (1994) and sequenced by Eurofins Genomics (Germany)

to confirm species identity of all specimens. Specificity of the newly designed assay was then assessed

using qPCR.

The reactions for both tissue and all eDNA samples contained; 12.5µL TaqMan® Environmental Master

Mix 2.0 (Life Technologies®), 6.5µL $DH_20$, 1µL (10µm) of each primer, 1µL (2.5µm) of probe with the

addition of 3µL template DNA. qPCR's were performed with 6 technical replicates (i.e. qPCR replicates) of

each sample on a StepOnePlus™ Real-Time PCR System (Applied Biosystems). qPCR conditions were as

follows: $50^{o}C$ for 5 min, denaturation at $95^{o}C$ for 8 min, followed by 50 cycles of $95^{o}C$ for 30 s and $55^{o}C$ for

1 min. Six no template controls (NTC's) were prepared using RT-PCR Grade Water (Ambion™) alongside a

duplicated serial dilution of control white-clawed crayfish DNA ($10^{-1}$-$10^{-3}$ ng $uL^{-1}$) for each qPCR plate that

was run. In each subsequent experiment, negative PCR controls were included in this fashion.

*Limits of detection (LOD) and quantification (LOQ)*

The reliability of our assay was also assessed, following the Minimum Information for Publication of

Quantitative Real-Time PCR Experiments (MIQE) Guidelines, which recommend the establishment of a

calibration curve to determine LOD and LOQ (Bustin et al., 2009). We prepared a serial dilution of DNA

extracted from white-clawed crayfish starting from 0.79ng $µL^{-1}$ to $7.9x10^{-8}$ ng $µL^{-1}$ with 10 qPCR replicates

per dilution analysed. The LOD was defined as the last standard dilution that resulted in a detection of

target DNA with at least one qPCR replicate at a threshold cycle (Ct) of <45. The LOQ was defined as the

last standard dilution in which targeted DNA was detected and quantified in a minimum of 90% of qPCR

replicates of the calibration curve under a Ct of 45 (Mauvisseau et al., 2019b).

*In-situ validation*

The reliability of the assay was further field tested by comparing eDNA-based and traditional capture-

mark-recapture sampling techniques at six sites of confirmed white-clawed crayfish presence (2017) in

the Centre-Val de Loire region, France. Each site was visited at least twice in subsequent nights between

22nd June and 1st of August 2018 (see supplementary information, Table S1 Individual white-clawed

crayfish were surveyed using a torching approach, counted and marked using a white waterproof marker

stain. In the second night the survey was repeated and marked, and non-marked crayfish were

differentiated. Population size was estimated using the Lincoln-Petersen Method $N = \left(\frac{Kn}{k}\right)$, where N is

the estimated population size, n is the number of crayfish marked on the 1st visit, K is the total number

captured on the 2nd visit and k is the number of those captured individual which were already marked on

the 2nd visit. (Petersen 1896, Lincoln 1930). Additionally, eDNA samples (two environmental replicates)

were collected at each site using the 0.22µm Sterivex filters (see below for detailed description). eDNA

samples were collected between the 22nd and 29th June 2018. The water volume filtered varied due to

cases of high turbidity (consistent minimum volume of 150mL, see Table S2 for list of all sample volumes).

Furthermore, additionally to the method below, eDNA filters were fixed with 2mL of ethanol to

accommodate for the longer storage and transport time between the field and the laboratory. All sampled

locations are part of an extensive monitoring programme for white-clawed crayfish population studies

and due to conservation reasons, locations of sites are not reported. The water temperature,

environmental variables and the volume of sample filtered for eDNA samples (varied due to variable

turbidity) were recorded at each site.

***Comparison of eDNA sampling methodologies in mesocosms***

Further, our aim was to assess the impact of eDNA sampling methodology on both the probability of eDNA

detection and the signal strength (i.e. Ct) of its detection. We tested differences between the most

common eDNA sampling methods utilised to date, including (i) ethanol precipitation (Biggs et al., 2015),

(ii) 2µm pump-based filtration (Strand et al., 2014), (iii) 0.45µm pressure filtration and (iv) 0.22µm

pressure filtration (Spens et al., 2017). All methods were assessed in two mesocosms, housed at Bristol

Zoological Gardens, Bristol, UK during autumn 2018. Both mesocosms were designed to the same

specifications but contained different water volumes and crayfish numbers. Mesocosm 1 had a volume of

3000L and contained 249 individual adult white-clawed crayfish and sub-adults (between 17 months and

four years old) with a total biomass of 1.3kg. Water parameters of mesocosm 1 were; pH 8, temperature

11°C, and under natural light conditions. Mesocosm 2 contained a volume of 1000L of water with the

same pH (8) but with higher water temperatures (16°C) and under artificial light conditions. This

mesocosm held a larger number of crayfish (379) but all were juvenile (five months old). The sensitivity of

juveniles to handling did not allow us to obtain the exact biomass of this mesocosm, but biomass was

estimated as 250g. Both mesocosms were set up as recirculating 'flow through filtration systems',

ensuring high water quality at all times. Six samples for each method and mesocosm combination were

collected from both mesocosms.

In brief; eDNA samples classified hereafter as 'precipitation' samples were collected following the protocol

outlined in Biggs et al. (2014). 1L of water (20 x 50mL subsamples) was collected from ~20cm below the

surface and after homogenization, a subsample of 90 mL (6x 15mL) was aliquoted into sterile tubes

containing a pre-mixed buffer solution (Biggs et al., 2014). Samples were stored at -20°C prior to extraction

and extracted following Tréguier et al. (2014). For batch of samples analysed, negative extraction controls

were included and processed analogously to other samples.

eDNA samples taken with a 2µm pump-based filtration consisted of 2L of water collected by the same

sub-sample method outlined above but were then filtered through a Millipore Glass fibre filter AP25,

47mm (2µm pore size) using a peristaltic pump (Masterflex E/S Portable Sampler, Cole-Parmer, USA). The

filter was housed in an In-Line Filter Holder 47mm (Merck) connected by silicone tubing. The combined

use of a peristaltic pump and a larger filter pore size allowed us to substantially increase the amount of

water filtered. The filter was then removed from the pump system and stored at -20°C before extraction.

Equipment was soaked and cleaned with 10% bleach between samples. Filters were extracted following

Spens et al. (2017). eDNA sample collections for 0.22µm and 0.45µm pressure filtration were undertaken

in the same manner. 20 sub-samples were collected and a 50mL syringe (BD Plastipak™, Ireland) was then

used to pressure filter 250mL of water through a sterile enclosed filter (Sterivex™, Merck®, Germany) with

either a pore size of 0.22µm (Polyethersulfone membrane) or 0.45µm (Polyvinylidene fluoride

membrane). All filters were stored at -20°C, and extracted following Spens et al. (2017).

*In-situ comparison of eDNA sampling methodologies*

Complementary to the tests in the mesocosm experiment, we also evaluated sampling methodologies

under natural conditions. However, we performed only pairwise method comparisons in order to contain

sampling effort in the field. As a test in a lentic system, eDNA samples were collected from a 1000m$^2$ pond

in the South West of England after the release of 40 white-clawed crayfish individuals (equal juvenile-

adult and male-female ratios, total biomass of 436g). Here, precipitation (sample volume: 90mL) was

compared against 0.22µm pressure filtration (sample volume: 250mL). Sampling started on the 20[th] April

2018 and was repeated two hours, seven days, 14 days and 35 days after crayfish release. At each

sampling time, three field replicates were taken from four sites around the pond for each method.

Additionally, 20 50mL sub-samples taken from the entire pond perimeter were pooled, homogenised and

sampled with three field replicates per method.

Our second field test was conducted in a lotic system. We sampled 10 sites (situated approx. 1km apart)

along a chalk stream river in Dorset (UK), during September 2017, and 4 sites along a river in Derbyshire

(UK). Here, ethanol precipitation was used in comparison to pump-based filtration (2µm, sample volume:

2L), using three field replicates at each site per method ($n$ = 42). Samples collected in the river system (20

pooled sub-samples as described above) were taken in an interval of 1-2m along a diagonal downstream-

to-upstream transect across the river. In this field test, we also assessed the ability to screen for crayfish

plague using both sampling methods. qPCRs in this instance were run using the primers and probe

developed by Strand et al. (2014). For all of the field method comparisons, bottled distilled water was

sampled on site and processed as field negative controls.

**Field test of white-clawed crayfish, signal crayfish and crayfish plague co-existence**

Finally, we assessed the distribution of white-clawed crayfish, signal crayfish and crayfish plague in a river-

system in Derbyshire (UK). Two field replicates were taken at each of eight sites along the river in

November 2017. Six of these sites were located in proximity to the inflow of tributaries and two field

replicates were taken upstream and downstream of their confluence to capture the influence of

populations potentially present within tributaries (supplementary information, Fig. S1). The other two

sites were sampled with two field replicates. Sampling was conducted using the precipitation method

outlined above and water samples were tested for the occurrence of all three species. Protocols of

Mauvisseau *et al.*, (2018) for signal crayfish and of Strand *et al.*, (2014) for crayfish plague were applied.

*Statistical Analysis*

Samples measured for the establishment of a standard curve were analysed using a linear regression to

evaluate the relationship between DNA dilution and Ct. A log-log data transformation decreased the

models Akaike Information Criterion (AIC; reduction by 23 units compared to second best model) and was

therefore used for downstream analyses. Residuals were tested for autocorrelation, normal distribution

and any remaining patterns (same procedure applied in all regression analyses). A logistic regression

analysis was also applied to test the relationship between DNA concentration and binomial detection data

assessing the change of detection probability with DNA concentrations. For the mesocosm and field

samples, the relationship between (i) the population density established by traditional sampling methods

and (ii) the Ct values and detection probability (calculated as the fraction of technical replicates that

resulted in positive detection) of eDNA measurements were examined in a linear regression model.

Differences in sample volumes between locations (due to turbidity) were accounted for by including

sample volume as a predictor in regression models, and log-log and untransformed models were

compared using AIC. Further, Ct and detection probability of different sampling methods were compared

using ANOVA analyses followed by Tukey's HSD post-hoc tests, and t-tests or nested ANOVA's (lotic and

lentic systems, where only two methods were compared). Prior to ANOVAs, heteroscedasticity was

evaluated, and data transformed if necessary. Finally, the co-existence of white-clawed crayfish with

signal crayfish and crayfish plague was tested in regression models using detection probability of all three

species. All described statistical analyses were performed using R version 3.4.1 (R Core Team (2017).

**Results**

***Assay development and* in-silico *and* in-vitro *validation***

Primers and probe were highly species-specific as *in-silico* and *in-vitro* tests did not reveal any matches

with non-target species (Table S3)*.* Analysis of the standard curve (Fig. 1A) revealed a strong dependency

of Ct values on DNA concentrations (y=-1.73x+20.8, *p*<0.001, $r^2$= 0.996). Likewise, the detection

probability was also positively related to DNA concentration in the sample (y=-0.18x+1.39, *p*=0.0016,

$r^2$=0.804; Fig. 1B), highlighting the possibility of a quantifiable assay being developed. Method sensitivity

analyses revealed a LOD of 7.9 x $10^{-5}$ng and a LOQ of 7.9 x $10^{-4}$ng crayfish DNA extract per µL$^{-1}$.

***In-situ validation***

Populations of white-clawed crayfish were found in five out of the six surveyed sites using traditional

survey methods. eDNA-based detection indicates the presence of white-clawed crayfish in all six sites,

though the site with no visual white-clawed crayfish sightings was characterised by a very low detection

probability. The Ct values from the six river sites were converted into DNA concentrations using the

calibration curve, which allowed us to compare the relationship between detection probability and DNA

concentration in laboratory and field samples (Fig. 1B). Four out of the six field sites lay outside of the 95%

confidence interval of the standard curve, indicating systematic differences between *in-vitro* validation

and field samples. As a direct result of limited recapture data, no recapture was observed in three out of

the six survey locations despite indication of the presence of a significantly sized population. Estimates of

population sites were therefore unable to be reliably made using the Lincoln-Petersen approach. The

relationship between the mean number of crayfish detected across all sampling sates was therefore

applied instead. The relationship of the mean number of crayfish detecting using conventional species

survey methods (torching) and detection probability of eDNA measurements (Fig. 1D) was significant, but

only when water temperature was included ($y=0.0118x_1-0.117x_2+1.77$; $x_1$=mean survey count,

$x_2$=temperature, $p=0.035$, $r^2=0.82$, best model resulted also in reduction of AIC by 10 units), with two out

of the six datapoints lying outside of the confidence interval. The relationship between Ct and the mean

number of crayfish detected using torching was marginally non-significant but showed a reasonable

model fit (Fig. 1C; $y=-0.00067\log(x)+3.76$, $p=0.079$, $r^2=0.47$, best model resulted in reduction of AIC by 5

units). Differences in filtered sample volume did not significantly influence results.

***Comparison of eDNA sampling methods***

In mesocosm experiments, sampling methodology had a significant impact on detection probability

(ANOVA $F_{(3,44)}=74.48$, $p<0.001$). Pairwise comparisons revealed that detection probabilities of all three

filtration-based methods (2μm, 0.22μm and 0.45μm) were comparable ($p>0.05$) but differed significantly

from the precipitation method ($p<0.001$, Fig. 2A). However, the *p*-value for the comparison between

0.45μm and 2μm was marginally non-significant ($p=0.051$). Similarly, methodologies also differed

significantly in Ct (ANOVA $F_{(3,178)}$=90.1, $p$<0.001). However, in contrast to detection probability, pairwise

tests indicated a difference between the 2μm filtration method and all the other approaches (p<0.001;

Fig. 2B; only samples with positive detection were included in the analysis).

*In-situ* comparisons of sampling methods in a lentic system were highly comparable to the mesocosm

experiment (Fig. 3 A-B). The precipitation method showed a significantly lower detection probability (T-

test, t=3.55, df=75.37, *p*<0.001) and a significantly higher Ct (t=-2.46, df=15.72, *p*<0.05) than the filtration-

based method (0.22μm). However, contrasting results were attained in lotic systems. Here, we assessed

the method for both, white-clawed crayfish and the crayfish plague (not present in mesocosms or ponds).

The detection probability of crayfish plague mirrored findings from other systems showing significantly

higher detection probabilities for the 2μm filtration method (nested ANOVA; $F_{(1,69)}$=4.92, p<0.05; Fig. 3E).

Ct values were not significantly different, but also indicated a better performance of the filtration-based

method (Fig. 3F). However, the results for white-clawed crayfish contrasted all other results. In lentic

systems, precipitation resulted in a higher detection probability (nested ANOVA $F_{(1,69)}$=13.77, *p*<0.001, Fig.

3C) and accordingly, lower Ct values (nested ANOVA; $F_{(1,34)}$=5.24, *p*=0.028; Fig. 3D). Consequently,

filtration-based methods performed consistently better except in lentic systems where eDNA from white-

clawed crayfish was more reliably assessed with the precipitation method.

**Field tests of species co-occurrence**

Finally, our joint assessment of white-clawed crayfish, signal crayfish and crayfish plague (Fig. 4)

demonstrated that white-clawed crayfish occurrence was related to both other species (Fig. 4, B,C). Whilst

univariate regressions were marginally non-significant (dependency of white-clawed crayfish on signal

crayfish: *p*=0.063; dependency of white-clawed crayfish on crayfish plague: *p*= 0.051), a multiple

regression analysis revealed significant relationships (y = -23.8$x_1$ + 13.1$x_2$ - 3.8, $r^2$=0.73 , *p*=0.016; y, $x_1$ and

$x_2$ represent detection probabilities of white-clawed crayfish, signal crayfish and crayfish plague,

respectively). Unsurprisingly, white-clawed crayfish was negatively impacted by the presence of signal

crayfish (Fig. 4B), yet contrary to expectation they were shown to be positively related with increase in

detection of crayfish plague (Fig. 4C). There was no apparent correlation between signal crayfish and

plague (Fig. 4D).

**Discussion**

Native crayfish species across Europe are threatened by invasive competitors and the crayfish plague,

resulting in a downward trajectory of native species' abundance and distribution (Holdich et al., 2009). In

this study, we present a novel assay for the detection of white-clawed crayfish, a flagship conservation

species in Western Europe. In rigorous *in-vitro* and *in-situ* tests, we evaluated the reliability of our assay

under various environmental conditions. Further, we applied our assay together with established eDNA-

based methods to assess the drivers of white-clawed crayfish occurrence. Overall, we were able to

demonstrate that our approach can not only be used for simple presence/absence surveys but also has

the potential to reveal species co-occurrence and interactions. However, our results also highlight that

such applications are only meaningful after thorough method testing and validation.

Field comparisons indicated a higher sensitivity of the eDNA assay compared to conventional surveys,

which only resulted in positive detection in five out of six sites. Such traditional surveys are reported to

vary in detection success (Gladman et al., 2010), therefore making comparisons between eDNA and these

methods challenging. Additionally, the two UK lotic river systems used to compare different sampling

approaches were since conventionally surveyed for the presence of crayfish, with no white-clawed

crayfish found, despite previous recorded presence, suggesting a higher sensitivity of the eDNA assay.

Whilst higher sensitivity is frequently reported for eDNA assays (Dejean et al., 2012; Smart et al., 2015),

and therefore likely represent a more reliable species detection approach, such results should be

interpreted with caution as eDNA-based approaches are associated with a risk of providing false positive

results (Furlan et al., 2016), for example via downstream transport of eDNA within river networks (Pont

et al., 2018). Moreover, false positives may result from historic eDNA, which is still present after the

extinction or emigration of the target species (Turner et al., 2015). In our case, this represents a valid

hypothesis as all field sites were populated by white-clawed crayfish a year before our field surveys (C.

Mauvisseau, personal communication). Consequently, it remains inconclusive whether the developed

eDNA approach truly has a higher sensitivity (i.e. false negative of torching method) or, in fact, white-

clawed crayfish was not present at the field site in question.

Further, an important component of our method validation was the comparison of different field sampling

approaches. Precipitation and filtration protocols to concentrate eDNA from the environment have

already been compared in a number of studies (Deiner et al., 2015; Spens et al., 2017). Most investigators

endorse filtration approaches (Hinlo et al., 2017b; Spens et al., 2017; Vörös et al., 2017), but optimal pore

size may differ between species (Spens et al., 2017). Moreover, method choice can also be environment-

dependant. For example, Eichmiller, et al. (2016) indicated filtration as the optimal eDNA-based method

for surveying the common carp, *Cyprinus carpio*. In contrast, Minamoto et al. (2016) highlighted

precipitation performed better – a result likely brought about by variations in the environment across

both studies. In our controlled mesocosm comparison, we found that a 2 μm filtration approach

outperformed precipitation and the other filtration methods tested. However, field comparisons revealed

contrasting results, again likely brought about by the different environments surveyed (Fig. 3). In this

scenario precipitation outperformed filtration (2 μm) in the lotic system.

One possible explanation for our divergent findings across different habitats is that target eDNA particles

differ in these environments. eDNA is exposed to continuous degradation through biotic (e.g. bacteria)

and abiotic (e.g. UV) factors (Strickler et al., 2015) and these degradation processes can affect eDNA

particle size distributions. Filtration has the advantage to collate eDNA from larger water volumes but is

linked to the risk of losing particles which are below filter pore sizes. Hence, the habitat-specific

differences in our method comparisons may be explained by the specific degradation processes within

the investigated river systems. A decrease of average eDNA particle size below the filter pore size would

substantially decrease detection probability of filtration approaches and explain our findings.

An alternative explanation for our results is linked to inhibition of eDNA amplification. Inhibitor

compounds (that interfere with qPCR processes), have been shown to affect target DNA amplification in

a non-linear way (Goldberg et al., 2016). If inhibitor concentration is low, amplification will not be strongly

impacted. However, if concentrations surpass a certain threshold, inhibitors may suppress the

amplification of even high concentrations of target eDNA (Mauvisseau et al., 2019a). Sampling methods

that differ in their water collection volumes and in the amount of concentrated target eDNA, will also

concentrate inhibitors to different degrees (Fig. 5). Consequently, sampling methods that reach higher

target eDNA concentrations may show a lower overall performance due to the non-linear relationship

between inhibitor concentrations and DNA amplification. This scenario will occur when inhibitors are

present in high concentrations and efficiently concentrated. Therefore, different ratios between target

eDNA and inhibitors in different environments can cause a shift in the relative performance of sampling

methods across habitats (Fig. 5). In our case, we did not include tests for inhibition, which include the

addition of synthetic DNA to qPCR reactions (i.e. failure to detect synthetic DNA indicates inhibition;

(Goldberg et al., 2016; Mauvisseau et al., 2019a). However, we recommend that such inhibition tests

should be included in future field method comparisons.

Both inhibition and different target-eDNA size distributions might also explain differences in method

comparisons between species in the same environment as observed for white-clawed crayfish and

crayfish plague in lotic habitats (Fig. 3). A fundamental distinction between the two species is that crayfish

plague depends for its proliferation on the frequent and abundant release of encapsulated spores (~8 µm

in diameter). It seems likely that these spores, which are designed for transport along large distances, will

show lower sensitivity to degradation than white-clawed crayfish DNA, which potentially could explain

our species-specific results.

Finally, we demonstrated that our approach can also be used for investigating ecological relationships

determining the distribution of endangered species. Our simultaneous assessments of white-clawed

crayfish, signal crayfish and crayfish plague revealed a negative impact of signal crayfish on white-clawed

crayfish (Fig. 4). Such negative impacts of invasive competitors on native crayfish species have been

frequently highlighted before (Holdich et al., 2009) and demonstrate the applicability of our approach.

Interestingly, however, we illustrated a positive relationship between white-clawed crayfish and crayfish

plague, which went against our expectation. Such co-occurrence might result from the one-time nature

of our sampling approach and reflect a disease outbreak within the crayfish population, which most

probably would result in local extinction (Strand et al., 2019). However, recent discoveries have also

indicated the potential of plague resistance in some white-clawed crayfish strains (Martín-Torrijos et al.,

2017). Such increased disease tolerance might facilitate a permanent co-existence of pathogen and host.

Consequently, further in-depth monitoring of species dynamics together with genetic profiling and

disease susceptibility tests should be a primary objective of future conservation planning.

**Conclusions**

Currently, many species-specific eDNA assays only cover *in-silico*, *in-vitro* and sometimes basic *in-situ*

validation steps (Dickie et al., 2018; Lacoursière-Roussel et al., 2016). Already published white-clawed

crayfish eDNA assays have shown some promising first results but yet need to go through the required

thorough level of *in-situ* evaluation (Atkinson et al., 2019; Robinson et al., 2018). Here we illustrate that

sampling methods can differ strongly in performance and recommend rigorous testing of eDNA assays to

optimise sampling strategies. However, our contrasting results of method comparisons across habitats

and species highlight that there might not be something like a universal 'optimal sampling method', but

that adjustments to account for local conditions are required. The resulting higher method reliability

increases the applicability of eDNA assays and paves the way for more detailed ecological studies to

improve species management and conservation.

**Acknowledgments**

For assistance with sample collection we thank Holly Thompson, Jacob Ball, Jack Greenhalgh, Isabelle

Parot and Antoine Monier. We would also like to thank Alex Lumsdon and the Environment Agency, Angus

Menzies and Dorset Wildlife Trust for funding and their assistance with sample collection. For providing

DNA samples: Catherine Souty-Grosset, Sina Tönges, Frank Lyko, Jenny Makkonen and Japo Jussila.

Funding was provided by SureScreen Scientifics, UK.

## References

**<u>References</u>**

Atkinson, S., Carlsson, J.E.L., Ball, B., Kelly-Quinn, M., Carlsson, J., 2019. eDNA assay for
Austropotamobius pallipes Field validation of an eDNA assay for the endangered white-clawed
crayfish Austropotamobius pallipes. bioRxiv. https://doi.org/10.1101/562710

Biggs, J., Ewald, N., Valentini, A., Gaboriaud, C., Dejean, T., Griffiths, R.A., Foster, J., Wilkinson, J.W.,
Arnell, A., Brotherton, P., Williams, P., Dunn, F., 2015. Using eDNA to develop a national citizen
science-based monitoring programme for the great crested newt (Triturus cristatus). Biol. Conserv.
183, 19–28. https://doi.org/10.1016/J.BIOCON.2014.11.029

Biggs, J., Ewald, N., Valentini, A., Gaboriaud, C., Griffiths, R., Foster, J., Wilkinson, J., Arnett, A., Williams,
P., Dunn, F., 2014. Analytical and methodological development for improved surveillance of the
Great Crested Newt Appendix 5 . Technical advice note for field and laboratory sampling of great
crested newt (Triturus cristatus) environmental DNA. Freshw. Habitats Trust. Oxford.

Bustin, S.A., Benes, V., Garson, J.A., Hellemans, J., Huggett, J., Kubista, M., Mueller, R., Nolan, T., Pfaffl,
M.W., Shipley, G.L., Vandesompele, J., Wittwer, C.T., 2009. The MIQE Guidelines: Minimum
Information for Publication of Quantitative Real-Time PCR Experiments. Clin. Chem. 55, 611–622.
https://doi.org/10.1373/clinchem.2008.112797

Bylemans, J., Furlan, E.M., Gleeson, D.M., Hardy, C.M., Duncan, R.P., 2018. Does Size Matter? An
Experimental Evaluation of the Relative Abundance and Decay Rates of Aquatic Environmental
DNA. Environ. Sci. Technol. 52, 6408–6416. https://doi.org/10.1021/acs.est.8b01071

Deiner, K., Walser, J.-C., Mächler, E., Altermatt, F., 2015. Choice of capture and extraction methods
affect detection of freshwater biodiversity from environmental DNA. Biol. Conserv. 183, 53–63.
https://doi.org/10.1016/J.BIOCON.2014.11.018

Dejean, T., Valentini, A., Miquel, C., Taberlet, P., Bellemain, E., Miaud, C., 2012. Improved detection of
an alien invasive species through environmental DNA barcoding: the example of the American
bullfrog *Lithobates catesbeianus*. J. Appl. Ecol. 49, 953–959. https://doi.org/10.1111/j.1365-
2664.2012.02171.x

Dickie, I.A., Boyer, S., Buckley, H.L., Duncan, R.P., Gardner, P.P., Hogg, I.D., Holdaway, R.J., Lear, G.,
Makiola, A., Morales, S.E., Powell, J.R., Weaver, L., 2018. Towards robust and repeatable sampling
methods in eDNA-based studies. Mol. Ecol. Resour. 18, 940–952. https://doi.org/10.1111/1755-
0998.12907

Ficetola, G.F., Miaud, C., Pompanon, F., Taberlet, P., 2008. Species detection using environmental DNA
from water samples. Biol. Lett. 4, 423–5. https://doi.org/10.1098/rsbl.2008.0118

Folmer, O., BLACK, M., HOEH, W., Lutz, R., Vrijenhoek, R., 1994. DNA primers for amplification of
mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. Mol. Mar. Biol.
Biotechnol. 3, 294–299. https://doi.org/10.1371/journal.pone.0013102

Füreder, L., Gherardi, F., Holdich, D., Reynolds, J., Sibley, P., Souty-Grosset, C., 2010. Austropotamobius
pallipes. The IUCN Red List of Threatened Species 2010: e.T2430A9438817 [WWW Document]. URL
http://www.iucnredlist.org/details/2430/0 (accessed 2.9.18).

Furlan, E.M., Gleeson, D., Hardy, C.M., Duncan, R.P., 2016. A framework for estimating the sensitivity of

eDNA surveys. Mol. Ecol. Resour. 16. https://doi.org/10.1111/1755-0998.12483

Geerts, A.N., Boets, P., Van den Heede, S., Goethals, P., Van der heyden, C., 2018. A search for
standardized protocols to detect alien invasive crayfish based on environmental DNA (eDNA): A lab
and field evaluation. Ecol. Indic. 84, 564–572. https://doi.org/10.1016/j.ecolind.2017.08.068

Gladman, Z.F., Yeomans, W.E., Adams, C.E., Bean, C.W., McColl, D., Olszewska, J.P., McGillivray, C.W.,
McCluskey, R., 2010. Detecting North American signal crayfish (Pacifastacus leniusculus) in riffles.
Aquat. Conserv. Mar. Freshw. Ecosyst. 20, 588–594. https://doi.org/10.1002/aqc.1130

Goldberg, C.S., Turner, C.R., Deiner, K., Klymus, K.E., Thomsen, P.F., Murphy, M.A., Spear, S.F., McKee,
A., Oyler-McCance, S.J., Cornman, R.S., Laramie, M.B., Mahon, A.R., Lance, R.F., Pilliod, D.S.,
Strickler, K.M., Waits, L.P., Fremier, A.K., Takahara, T., Herder, J.E., Taberlet, P., 2016. Critical
considerations for the application of environmental DNA methods to detect aquatic species.
Methods Ecol. Evol. 7, 1299–1307. https://doi.org/10.1111/2041-210X.12595

Hinlo, R., Furlan, E., Suitor, L., Gleeson, D., 2017a. Environmental DNA monitoring and management of
invasive fish: comparison of eDNA and fyke netting. Manag. Biol. Invasions 8, 89–100.
https://doi.org/10.3391/mbi.2017.8.1.09

Hinlo, R., Gleeson, D., Lintermans, M., Furlan, E., 2017b. Methods to maximise recovery of
environmental DNA from water samples. PLoS One 12, e0179251.
https://doi.org/10.1371/journal.pone.0179251

Holdich, D.M., Reeve, I.A.N.D., 1991. Distribution of freshwater crayfish in the British Isles, with
particular reference to crayfish plague, alien introductions and water quality. Aquat. Conserv. Mar.
Freshw. Ecosyst. 1, 139–158. https://doi.org/10.1002/aqc.3270010204

Holdich, D.M.M., Reynolds, J.D.D., Souty-Grosset, C., Sibley, P.J.J., 2009. A review of the ever increasing
threat to European crayfish from non-indigenous crayfish species. Knowl. Manag. Aquat. Ecosyst.
11. https://doi.org/10.1051/kmae/2009025

Huver, J.R., Koprivnikar, J., Johnson, P.T.J., Whyard, S., 2015. Development and application of an eDNA
method to detect and quantify a pathogenic parasite in aquatic ecosystems. Ecol. Appl.
https://doi.org/10.2307/24432104

Jerde, C.L., Chadderton, W.L., Mahon, A.R., Renshaw, M.A., Corush, J., Budny, M.L., Mysorekar, S.,
Lodge, D.M., 2013. Detection of Asian carp DNA as part of a Great Lakes basin-wide surveillance
program. Can. J. Fish. Aquat. Sci. 70, 522–526. https://doi.org/10.1139/cjfas-2012-0478

Kearse, M., Moir, R., Wilson, A., Stones-Havas, S., Cheung, M., Sturrock, S., Buxton, S., Cooper, A.,
Markowitz, S., Duran, C., Thierer, T., Ashton, B., Meintjes, P., Drummond, A., 2012. Geneious Basic:
an integrated and extendable desktop software platform for the organization and analysis of
sequence data. Bioinformatics 28, 1647–9. https://doi.org/10.1093/bioinformatics/bts199

Lacoursière-Roussel, A., Côté, G., Leclerc, V., Bernatchez, L., 2016. Quantifying relative fish abundance
with eDNA: a promising tool for fisheries management. J. Appl. Ecol. 53, 1148–1157.
https://doi.org/10.1111/1365-2664.12598

Lincoln, F. C. 1930. Calculating waterfowl abundances on the basis of banding returns. U.S. Department
of Agriculture, Circular 118.

MacDonald, A.J., Sarre, S.D., 2017. A framework for developing and validating taxon-specific primers for
specimen identification from environmental DNA. Mol. Ecol. Resour. 17, 708–720.
https://doi.org/10.1111/1755-0998.12618

Mächler, E., Deiner, K., Spahn, F., Altermatt, F., 2016. Fishing in the Water: Effect of Sampled Water
Volume on Environmental DNA-Based Detection of Macroinvertebrates. Environ. Sci. Technol. 50,
305–312. https://doi.org/10.1021/acs.est.5b04188

Martín-Torrijos, L., Campos Llach, M., Pou-Rovira, Q., Diéguez-Uribeondo, J., 2017. Resistance to the
crayfish plague, Aphanomyces astaci (Oomycota) in the endangered freshwater crayfish species,
Austropotamobius pallipes. PLoS One 12, e0181226.
https://doi.org/10.1371/journal.pone.0181226

Mauvisseau, Q., Coignet, A., Delaunay, C., Pinet, F., Bouchon, D., Souty-Grosset, C., 2018. Environmental
DNA as an efficient tool for detecting invasive crayfishes in freshwater ponds. Hydrobiologia 805,
163–175. https://doi.org/10.1007/s10750-017-3288-y

Mauvisseau, Q., Davy-Bowker, J., Bulling, M., Brys, R., Neyrinck, S., Troth, C., Sweet, M., 2019a.
Improving detection capabilities of a critically endangered freshwater invertebrate with
environmental DNA using digital droplet PCR. bioRxiv 661447. https://doi.org/10.1101/661447

Mauvisseau, Q., Tönges, S., Andriantsoa, R., Lyko, F., Sweet, M., 2019b. Early detection of an emerging
invasive species: eDNA monitoring of a parthenogenetic crayfish in freshwater systems,
Management of Biological Invasions.

Minamoto, T., Naka, T., Moji, K., Maruyama, A., 2016. Techniques for the practical collection of
environmental DNA: filter selection, preservation, and extraction. Limnology 17, 23–32.
https://doi.org/10.1007/s10201-015-0457-4

Pont, D., Rocle, M., Valentini, A., Civade, R., Jean, P., Maire, A., Roset, N., Schabuss, M., Zornig, H.,
Dejean, T., 2018. Environmental DNA reveals quantitative patterns of fish biodiversity in large
rivers despite its downstream transportation. Sci. Rep. 8, 10361. https://doi.org/10.1038/s41598-
018-28424-8

Petersen, C. G. J. 1896. The yearly immigration of young plaice into the Limfjord from the German Sea.
Report of the Danish Biological Station 1895 6:1–77.

R Core Team (2017). R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, A.U. https://www. R. org/.

Robinson, C.V., Uren Webster, T.M., Cable, J., James, J., Consuegra, S., 2018. Simultaneous detection of
invasive signal crayfish, endangered white-clawed crayfish and the crayfish plague pathogen using
environmental DNA. Biol. Conserv. 222, 241–252. https://doi.org/10.1016/j.biocon.2018.04.009

Smart, A.S., Tingley, R., Weeks, A.R., Van Rooyen, A.R., McCarthy, M.A., 2015. Environmental DNA
sampling is more sensitive than a traditional survey technique for detecting an aquatic invader.
Ecol. Appl. 25, 1944–1952. https://doi.org/10.1890/14-1751.1

Spens, J., Evans, A.R., Halfmaerten, D., Knudsen, S.W., Sengupta, M.E., Mak, S.S.T., Sigsgaard, E.E.,
Hellström, M., 2017. Comparison of capture and storage methods for aqueous macrobial eDNA
using an optimized extraction protocol: advantage of enclosed filter. Methods Ecol. Evol. 8, 635–

645. https://doi.org/10.1111/2041-210X.12683

Strand, D.A., Johnsen, S.I., Rusch, J.C., Agersnap, S., Larsen, W.B., Knudsen, S.W., Møller, P.R., Vrålstad,
T., 2019. Monitoring a Norwegian freshwater crayfish tragedy:  eDNA snapshots of invasion,
infection and extinction. J. Appl. Ecol. 56, 1365-2664.13404. https://doi.org/10.1111/1365-
2664.13404

Strand, D.A., Jussila, J., Johnsen, S.I., Viljamaa-Dirks, S., Edsman, L., Wiik-Nielsen, J., Viljugrein, H.,
Engdahl, F., Vrålstad, T., 2014. Detection of crayfish plague spores in large freshwater systems. J.
Appl. Ecol. 51, 544–553. https://doi.org/10.1111/1365-2664.12218

Strickler, K.M., Fremier, A.K., Goldberg, C.S., 2015. Quantifying effects of UV-B, temperature, and pH on
eDNA degradation in aquatic microcosms. Biol. Conserv. 183, 85–92.
https://doi.org/10.1016/J.BIOCON.2014.11.038

Tréguier, A., Paillisson, J.-M., Dejean, T., Valentini, A., Schlaepfer, M.A., Roussel, J.-M., 2014.
Environmental DNA surveillance for invertebrate species: advantages and technical limitations to
detect invasive crayfish Procambarus clarkii in freshwater ponds. J. Appl. Ecol. 51, 871–879.
https://doi.org/10.1111/1365-2664.12262

Turner, C.R., Uy, K.L., Everhart, R.C., 2015. Fish environmental DNA is more concentrated in aquatic
sediments than surface water. Biol. Conserv. 183, 93–102.
https://doi.org/10.1016/j.biocon.2014.11.017

Vörös, J., Márton, O., Schmidt, B.R., Gál, J.T., Jelić, D., 2017. Surveying Europe's Only Cave-Dwelling
Chordate Species (Proteus anguinus) Using Environmental DNA. PLoS One 12, e0170945.
https://doi.org/10.1371/journal.pone.0170945

## **Author Contributions**

527    C.T., M.S. and J.N. designed the experiment and methodology, C.T., Q.M., J.N. and C.M. collected field

528    samples, C.T., and Q.M. performed extraction and PCR, C.T., A.B. and M.B. analysed the data. The

529    manuscript was written by C.T., M.S. and A.B. and reviewed by all authors.

**Figure 1. (A)** Relationship between cycle threshold (Ct) and DNA concentration from White-clawed crayfish qPCR calibration curve. Limit of detection (LOD) and limit of quantification (LOQ) are illustrated by vertical lines (dashed-yellow and red respectively). **(B)** Change in detection probability with increasing DNA concentrations in calibration curve data (blue line). Red triangles represent field data (in-*situ* validation data) that is compared with the log-regression based on in-*vitro* calibration curve data. Three out of five field data points were outside the established confidence interval, indicating discrepancies between field and laboratory-based data sets. **(C)** Relationship between Ct values and white-clawed crayfish population monitored using traditional method. **(D)** Relationship between detection probability of eDNA and traditionally evaluated crayfish population sizes. The blue line and the light-blue shaded area reflect the results of a logit regression and its 95% confidence interval, respectively.

543
**Figure 2.** Comparison of the detection probability **(A)** and Ct values **(B)** of white-clawed crayfish using
different eDNA sampling methods (0.22μm filtration, 0.45μm filtration, 2μm filtration and precipitation)
in a controlled mesocosm experiment (* indicates statistical significance).

547
548 **Figure 3.** Comparison of the detection probability **(A, C, E)** and Ct values **(B, D, F)** of different eDNA
549 sampling methods (filtration and precipitation) for white-clawed crayfish in a lentic system **(Pond, A-B)**
550 (filter pore size 0.22 µm) and for both white-clawed crayfish **(River, C-D)** and crayfish plague **(River, E-F)**
551 in the same lotic system (filter pore size 2 µm) (* in panels signifies significant differences between
552 pairwise method).

553
**Figure 4. (A)** Detection of eDNA from white-clawed crayfish (green squares), signal crayfish (blue squares)
and crayfish plague. (purple squares) in a river catchment in Derbyshire. Eight locations were sampled and
are represented by red dots. The empty squares represent the negative qPCR replicates. **(B)** Indicates the
relationship between the detection probability of white-clawed crayfish and detection probability of
signal crayfish. **(C)** The relationship between the detection probability of white-clawed crayfish and
detection probability of crayfish plague. **(D)** The relationship between the detection probability of signal
crayfish and detection probability of crayfish plague.

eDNA concentration in the sample

**Figure 5.** Schematic of the co-dependency of detection probability on target eDNA and inhibitors concentrations in water samples. Detection probability increases with eDNA concentration and decreases with inhibitor concentrations but is low when both variables are high. Each water body is characterised by a certain ratio between inhibitor and target eDNA concentrations represented by black dotted lines (**1-3**). A change in sampling methods accompanied by a change in the sampled water volume will result in different concentrations of target eDNA and inhibitors in the sample and in shifts along dotted lines (**grey crosses and dots**). An increase in sampled water volume will therefore in some water bodies increase (Line 1) and in other decrease (Line 2) detection probability. The same is true when different eDNA assays in the same water body are considered. While eDNA concentrations of two targets may differ, inhibitor concentrations will be the same. Consequently, samples with the same water volume will have the same inhibitor concentrations (horizontal dashed lines). Nevertheless, changes in sampling volume and method can result in increased detection probability for one target (Line 3) but not for the other (Line 2).