The finger 2 tip loop of Activin A is required for the formation of its non-signaling complex with ACVR1 and type II Bone Morphogenetic Protein receptors

Vincent Idone[1*], Richard A. Corpina[1], Erich J. Goebel[2], Camille J. Cunanan[1], Alexandra Dimitriou[1], Hyonjong Kim[1], Qian Zhang[1], Ashique Rafique[1], Raymond Leidich[1], Xin Wang[1], Joyce McClain[1], Johanna Jimenez[1], Kalyan C. Nannuru[1], Nyanza J. Rothman[1], John B. Lees-Shepard[1], Andrew J. Murphy[1], Thomas B. Thompson [2], and Aris N. Economides[1*]

[1]Regeneron Pharmaceuticals, 777 Old Saw Mill River Road, Tarrytown, NY 10591; [2] University of Cincinnati, 231 Albert Sabin Way ML 0524, Cincinnati, OH 45267 USA

*to whom correspondence should be addressed: vincent.idone@regeneron.com, or aris@regeneron.com

Keywords: activin, activin A receptor, type I (ACVR1), fibrodysplasia ossificans progressive (FOP), structure-function, TGF-beta superfamily, non-signaling complex (NSC)

Abstract
Activin A functions in BMP signaling in two ways: it either engages ACVR1B to activate Smad2/3 signaling or binds ACVR1 to form a non-signaling complex (NSC). Although the former property has been studied extensively, the roles of the NSC remain unexplored. The genetic disorder fibrodysplasia ossificans progressiva (FOP) provides a unique window into ACVR1/Activin A signaling because in that disease Activin can either signal through FOP-mutant ACVR1 or form NSCs with wild type ACVR1. To explore the role of the NSC, we generated 'agonist-only' Activin A muteins that activate ACVR1B but cannot form the NSC with ACVR1. Using one of these muteins we demonstrate that failure to form the NSC in FOP results in more severe disease pathology. These results provide the first evidence for a biological role for the NSC in vivo and pave the way for further exploration of the NSC's physiological role in corresponding knock-in mice.

Impact Statement
The signaling complex formed by Activin A and ACVR1 is operant *in vivo* and is required to temper the degree of heterotopic ossification in the genetic disorder fibrodysplasia ossificans progressiva.

Introduction
Activin A is a protein that belongs to the TGFß/BMP family of ligands (Hinck, Mueller, & Springer, 2016). These ligands initiate signaling by driving the formation of heterotetrameric complexes of their cognate type I and type II receptors, and where a dimeric ligand brings together two type I and two type II receptors (Yadin, Knaus, & Mueller, 2016). There are seven type I and five type II receptors that mediate signaling for this family of ligands. The choice of type I receptor in the signaling complex is the main determinant of which of two signaling pathways is activated: Smad1/5/8, when the type I receptors are ACVRL1, ACVR1, BMPR1A,

or BMPR1B; and Smad2/3, when the type I receptors are ACVR1B, TGFBR1, or ACVR1C. Activin A utilizes ACVR1B as its type I receptor in conjunction with the type II receptors ACVR2A, ACVR2B, and, to a lesser extent, BMPR2 to signal through Smad2/3 (Aykul et al., 2017). The biological functions of Activin A have been investigated widely using multiple approaches (Makanji et al., 2014; Namwanje & Brown, 2016) including reverse genetics experiments wherein the gene encoding for Activin A – *Inhba* – has been 'knocked out' (Archambeault & Yao, 2010; Matzuk et al., 1995; Pangas et al., 2007). The results of those experiments have been largely interpreted in the context of Activin A acting as an agonist of ACVR1B to induce Smad2/3 or other signaling pathways (Archambeault & Yao, 2010; Makanji et al., 2014; Namwanje & Brown, 2016; Pangas et al., 2007).

Interestingly, ACVR1 was originally cloned as the type I receptor for Activin A (Attisano et al., 1993; Tsuchida, Mathews, & Vale, 1993), but the inability of Activin A to activate ACVR1, followed by the discovery that BMP7 activates it, led to a 'relabeling' of ACVR1 as a BMP receptor (Macías-Silva, Hoodless, Tang, Buchwald, & Wrana, 1998). However, recent studies pinpoint a previously unrecognized property of Activin A: that it can engage the type I receptor ACVR1, to form an ACVR1•Actvin A•type II receptor non-signaling complex (NSC) (Hatsell et al., 2015; Olsen et al., 2015). This NSC is unique in that its stoichiometry is identical to that of corresponding signaling complexes formed between the same receptors and BMPs, but also in that it is converted into a signaling complex in the ACVR1-driven genetic disorder fibrodysplasia ossificans progressiva (FOP). Excluding the special situation of FOP (see below), the NSC functions to tie down Activin A and render it unavailable for signaling, but also to tie down the type II receptors and ACVR1 and render them unavailable for engagement with BMPs, hence resulting in an apparent inhibition of ACVR1-mediated BMP signaling; in cells where ACVR1 is the main type I receptor, Activin A inhibits BMP6- (Hatsell et al., 2015) and BMP7-induced signaling (this work).

The picture is more complex in FOP. FOP is a rare autosomal-dominant genetic disorder that arises from missense mutations in the sequence encoding the intracellular domain of ACVR1 (Katagiri, Tsukamoto, & Kuratani, 2018). The most clinically important feature of FOP is the episodic, yet progressive and cumulative, formation of heterotopic bone in connective tissue, a process commonly referred to as heterotopic ossification (HO) (Huning & Gillessen-Kaesbach, 2014). FOP-mutant variants of ACVR1 display the neomorphic property of recognizing Activin A (as well as other Activins) as an agonistic ligand, much like a BMP (Hatsell et al., 2015; Hino et al., 2015). In mouse FOP, activation of FOP-mutant ACVR1 by Activin A is required for HO, as demonstrated by experiments where inhibition of Activin A, using highly specific monoclonal antibodies, halts both the occurrence and the progression of HO (Hatsell et al., 2015; Lees-Shepard, Nicholas, et al., 2018; Lees-Shepard, Yamamoto, et al., 2018; Upadhyay et al., 2017). Thus, in FOP, the ACVR1[FOP mutant]•Activin A•type II receptor complex, which is stoichiometrically identical to the NSC, acts as a signaling complex. However, since FOP is autosomal-dominant, one wild type copy of ACVR1 remains operational and hence capable of sequestering Activin A in NSCs. In mouse FOP, removal of the wild type copy of *Acvr1* exacerbates the degree of HO (Lees-Shepard, Yamamoto, et al., 2018). These data are consistent with a model where loss of wild type ACVR1 results in loss of the NSC and concomitant increase in the number of active complexes between Activin A and FOP-mutant ACVR1. Whereas these results indicate that the NSC is operant *in vivo*, they do not provide direct

evidence for the NSC, and also do not inform whether there are other biological roles for the NSC.

To date, there has been no systematic exploration of the biological roles of the NSC either within or outside of FOP. Outside of FOP, exploration of the NSC has been limited to myeloma cells in culture, wherein Activin A protects them from BMP9-induced apoptosis (Olsen et al., 2015); whether a similar phenomenon takes place *in vivo* is currently unknown. In order to begin to address such knowledge gaps, we engineered a set of 'agonist-only' Activin A muteins that retain their ability to activate ACVR1B but fail to form the NSC with ACVR1. Using these muteins, we show that the finger 2 tip loop (F2TL) region of Activin A is critical for the formation of the NSC, and we obtain conclusive *in vivo* evidence that the NSC is required for tempering the degree of HO in mouse FOP. Our successful engineering and *in vivo* application of agonist-only Activin A to explore the role of the NSC in FOP sets the stage for further exploration of the wider physiological roles of the NSC using the corresponding knock-in mice.

Results

A conceptual framework for engineering 'agonist-only' Activin A muteins
In order to set the stage for exploring the role of the ACVR1•Activin A•type II receptor NSC, we attempted to engineer Activin A muteins that:
1. Activate ACVR1B in a manner indistinguishable to wild type Activin A (Fig. 1A, left panels).
2. Preserve Activin A's type I receptor preferences (i.e. do not utilize other type I receptors such as TGFBR1 or ACVR1C).
3. Retain their interaction with Activin A's natural antagonists, Follistatin (FST) and Follistatin-like 3 (FSTL3) (Fig. 1A, right panels).
4. Cannot form a NSC with ACVR1 as evidenced by a reduction of inhibition of BMP7-induced activation down to the level of inhibition that is driven solely by binding of Activin A to its type II receptors (Fig. 1B). (Since binding of Activin A to type II receptors is a required property for signaling through ACVR1B (Greenwald et al., 2004), it follows that it is not possible to engineer an Activin A that signals through ACVR1B but does not compete with BMPs for type II receptor occupancy.)

As a first step to engineer Activin A muteins with such properties, we searched for BMP-TGFß family members that engage the same type II receptors as Activin A, utilize ACVR1B as their type I receptor, and do not engage ACVR1. Three ligands fulfill these requirements: Nodal, MSTN, and GDF11 (Yadin et al., 2016). Given that of these three, MSTN and GDF11 can also utilize TGFBR1 (Andersson, Reissmann, & Ibanez, 2006; Rebbapragada, Benchabane, Wrana, Celeste, & Attisano, 2003), we focused on Nodal as the main donor of type I receptor-binding regions to substitute into the corresponding regions of Activin A, while also exploring the outcomes of substituting the same regions from other ligands. To define the specific regions of Activin to replace with their Nodal-derived counterparts, we initially relied on previously engineered artificial ligands where one region of Activin A had been replaced by the corresponding region from other BMPs (Cash, Rejon, McPherron, Bernard, & Thompson, 2009; Harrison et al., 2004; Korupolu, Muenster, Read, Vale, & Fischer, 2008). Based on the premise

that the regions that are most likely to define receptor utilization preferences are also those that are more evolutionarily divergent, we chose regions where the sequence diversity of TGFß ligands was significant.

More specifically, several studies have demonstrated that the pre-helix and post-helix regions make critical contacts with type I receptors. Substitution of the region encompassed by the pre-helix and post-helix region (amino acids 355 to 391 of mature Activin A) with the corresponding region from BMP2, switched type I (but not type II) receptor utilization (Korupolu et al., 2008), whereas swapping the pre-helix region of Activin A with that of MSTN switched the preference of Activin A from ACVR1B to TGFBR1 (Cash et al., 2009). Moreover, the recently solved structure of GDF11 (an Activin family ligand) in complex with TGFBR1 and ACVR2B (PDB:6MAC), shows that the Finger 2 region is particularly important for determining type I receptor specificity for the Activin class (Goebel et al., 2019). Superimposition of Activin A into this ternary complex predicts that Activin A's finger 2 tip loop (F2TL; amino acids 406-409) also participates in binding to type I receptors (Fig. S1). To further explore potential interactions of Activin A with ACVR1, we generated predictions of specific molecular interactions through structural modeling. We started with the structure of Activin A bound to Fs288 (PDB:2B0U) (Thompson, Lerch, Cook, Woodruff, & Jardetzky, 2005), as Fs288 engages Activin A using the same interface as the type I receptor. This fixed the structure of Activin A as would be expected when bound to receptor. We then generated several models of ACVR1 based on closely related receptors and found BMPR1A to provide the best structural prediction. Superimposition of these models to the complex structure of TGFBR1•GDF11•ACVR2B followed by energy minimization of amino acid side chains suggested a key interaction at the fingertip. Specifically, D406 extends into the type I binding site (seen in several Activin A structures – PDB: 1S4Y,2B0U, 2ARV, 3B4V) and is flanked by two lysine residues from ACVR1 (Fig. S1) (Greenwald et al., 2004; Harrington et al., 2006; Stamler et al., 2008; Thompson et al., 2005). This observation supports a key role for D406 in binding ACVR1. Our model is further supported by the fact that there is a conserved hydrophobic residue (M79 in ACVR1) that rests in a hydrophobic pocket formed by the Activin A dimer, forming a "knob-in-hole" motif, which is also in the BMP2:BMPR1A (Allendorph, Vale, & Choe, 2006) and GDF11•TGFBR1 complexes (PDB:2GOO, 6MAC) (Goebel et al., 2019).

Along with the F2TL region, the pre- and post-helix regions also exhibit significant sequence diversity between ligands, hence making them attractive candidates for type I receptor selectivity (Fig. S2). In contrast, other regions that are known to be involved in engaging the type I receptors, such as finger 1 and H3 helix, were excluded as donors for substitution analysis; finger 1 was excluded because it displays a high level of sequence conservation between ligands, whereas the H3 helix was not utilized because it appears to play a structural role and not to be a major determinant of type I receptor choice (Allendorph, Isaacs, Kawakami, Izpisua Belmonte, & Choe, 2007). Hence, we chose to focus on these three regions – F2TL, pre-helix, and post-helix – as sources of sequences to transfer from Nodal (and other Activin class ligands) into Activin A.

Activin A with Nodal F2TL activates ACVR1B

Based on the design criteria and biochemical and structural information summarized above, we engineered Activin A muteins in which their pre-helix, post-helix, and F2TL regions were replaced with the corresponding regions of Nodal, generating **Activin A.Nod.pre**, **Activin A.Nod.post**, and **Activin A.Nod.F2TL**, respectively (Fig. 2). After expression and purification of their mature forms (Fig. S3), we first tested their ability to activate Smad2/3 signaling, a measure for utilization of ACVR1B as the type I receptor. Of these three muteins, the Activin A.Nod.F2TL exhibited activity very close to that of wild type Activin A (Fig. 3), whereas the other two were less active (Fig. S4).

The realization that the tip loop variant performed according to our specifications for activation of ACVR1B prompted closer examination of the finger 2 tip loop region sequences between Activin A and Nodal. This comparison revealed an intriguing difference between the fingertips of these ligands: that Nodal lacks an aspartic acid at position 406 (Fig. 2*B*). Moreover, an alignment of Activin A from different species reveals that this residue is highly conserved (Fig. S5), whereas it varies when the different Activins are compared (Fig. S2). Therefore, we tested whether a deletion of D406 would generate an Activin A mutein with properties similar to those of the Nodal tip loop variant (Activin A.Nod.F2TL). We compared the ability of Activin A.ΔD406, Activin A.Nod.F2TL, and wild type Activin A to activate ACVR1B using a CAGA-luciferase assay as a readout for Smad2/3 activation. As shown on Figure 3A, Activin A.Nod.F2TL is slightly less active than Activin A, whereas Activin A.ΔD406 exhibits identical activity to wild type Activin A, making Activin A.ΔD406 a second potential candidate for an agonist-only Activin A variant to study further.

Activin A.Nod.F2TL and Activin A.ΔD406 retain their type I receptor preferences
In order to exclude the possibility that by substituting the sequence of finger 2 we have inadvertently altered type I receptor preferences (precluding the deliberate reduction of binding to ACVR1), we employed an assay that measures dimerization of type I with type II receptors and investigated whether Activin A.Nod.F2TL and ΔD406 muteins displayed altered type I receptor preferences. In line with our bioassay results, both Activin A.Nod.F2TL and Activin A.ΔD406 are identical to Activin A in their ability to dimerize ACVR1B and TGFBR1 along with the corresponding type II receptors (Fig. 3B; Fig. S6).

Activin A.Nod.F2TL cannot form the NSC
In order to determine whether Activin A.Nod.F2TL and Activin A.ΔD406 have lost their ability to form the NSC with ACVR1 and type II receptors, we tested their capacity to inhibit BMP7-initiated activation of ACVR1 using a Smad1/5/8-responsive luciferase-based assay. Consistent with previous observations utilizing BMP6 (Hatsell et al., 2015) and BMP9 (Hatsell et al., 2015; Olsen et al., 2015), Activin A was able to inhibit activation of ACVR1 by BMP7 in dose-dependent manner. In contrast, both Activin A.Nod.F2TL and Activin A.ΔD406 were approximately 60-fold and 15-fold less effective at inhibiting BMP7, respectively (Fig. 4; Fig. S7). In addition, as would be expected for any Activin A variant that has reduced ability to form the NSC, both Activin A.Nod.F2TL (Fig. 3B) and Activin A.ΔD406 (Fig. S6) display greatly reduced ability to induce dimerization of ACVR1 with ACVR2A.

To exclude the possibility that this reduced ability to inhibit BMP7-initiated activation of ACVR1 is due to loss or reduction of binding of these muteins to type II receptors, we tested Activin A.Nod.F2TL and Activin A.ΔD406 for their ability to engage ACVR2B, one of the main type 2 receptors, using Biacore. In this binding assay, Activin A.Nod.F2TL and ActivinA.ΔD406 bound with high affinity, similar to wild type Activin A, confirming that the corresponding finger 2 tip loop substitutions in these two muteins do not affect type II receptor binding sites (Fig. S8). Furthermore, REGN2476, an anti-Activin A monoclonal antibody that neutralizes Activin A by blocking its binding to its type II receptors (Fig. S9), completely alleviates inhibition of BMP7 signaling by Activin A.Nod.F2TL further indicating that the remaining inhibition is a result of binding of this mutein to type II receptors (Fig. 4A). Similar results were obtained for ActivinA.ΔD406 in this assay (Fig. S7); however, its reduced effectiveness in inhibiting BMP7-initiated activation of ACVR1 indicates that ActivinA.ΔD406 can still interact with ACVR1 to form a functional NSC, albeit not as well as wild type Activin A.

To demonstrate that it is indeed the inability of Activin A.Nod.F2TL to form the NSC that accounts for its reduced ability to antagonize BMP7-initiated signaling by ACVR1, we used a neutralizing anti-Activin A antibody, H4H10442, that inhibits the interaction of Activin A with its type I receptors by binding to the finger 2 tip loop region (Fig. S10). We hypothesized that binding of H4H10442 to Activin A will mimic the effect of the Nod.F2TL substitution, and hence result in a BMP7 signaling inhibition profile identical to that of Activin A.Nod.F2TL. Indeed, H4H10442 shifts the dose response of the inhibition brought about by wild type Activin A on BMP7-initiated signaling to that obtained by Activin A.Nod.F2TL (Fig. 4B), hence verifying that the reduced inhibition obtained with Activin A.Nod.F2TL is a direct consequence of the inability of Activin A.Nod.F2TL's F2TL region (which is that of Nodal) to bind ACVR1. Using the same criteria, our data shows that Activin A.ΔD406 has partially retained the ability to antagonize BMP7 signaling via ACVR1, making it a less desirable 'agonist-only' Activin A mutein (Fig. S7). Taken together, these results clearly indicate that the remaining ability of Activin A.Nod.F2TL to antagonize BMP7-initiated signaling is indeed through engagement of the type II receptors, whereas its reduced ability to antagonize BMP7 is a property of its inability to engage ACVR1 to form the NSC.

Incidentally, in contrast to what is seen with Activin A.Nod.F2TL and Activin A.ΔD406, inhibition of binding of wild type Activin A to the type II receptor (via REGN2476) does not 'flat-line' the ability of wild type Activin A to inhibit BMP7-initiated signaling (Fig. 4, and Fig. S7). In fact, the response to the two antibodies - H4H10442, which blocks binding to ACVR1 by occupying the Activin A F2TL (Fig. S10), and REGN2476, which blocks binding to the type II receptors (Fig. S9)– is nearly identical when Activin A is the ligand outcompeting BMP7. This indicates that Activin A can engage ACVR1 even in the absence of binding to type II, at least under conditions where ACVR1 is expressed at high levels.

Activin A.Nod.F2TL and Activin A.ΔD406 retain their interactions with their antagonists
As a last step in examining potential differences between Activin A and these two muteins, we tested their interaction with natural Activin inhibitors, Follistatin (FST) and Follistatin-like 3 (FSTL3) (Fig. 5). Both isoforms of FST, differing in the absence (Fs288, Fig. 5*A*) or presence (Fs315, Fig. 5*B*) of c-terminal acidic tail, inhibit Activin A signaling to baseline in Smad2/3

reporter assays. Activin A.Nod.F2TL displays nearly identical inhibition profile compared to wild type, whereas Activin A.ΔD406 shows a slightly decreased ability to be inhibited by both FST isoforms but still can be fully inhibited to baseline (Fig. S11). Contrasting this, both F2TL muteins display decreased ability to be inhibited by FSTL3 (Fig. 5*C*, Fig. S11). We attribute this to a key difference in the way FSTL3 engages Activin A compared to FST. Whereas both FST and FSTL3 engage the Activins and other TGFß ligands in a 2:1 stoichiometric ratio with two Follistatin molecules per one dimeric ligand, when FST engages Activin A, the FSD3 domain reaches around the ligand to non-covalently contact the N-terminal domain of the other FST molecule, essentially fully wrapping around the ligand. (There are relatively few specific contacts between FST and the ligands that it interacts with; this makes FST a quite promiscuous modulator of multiple BMP/TGFß family members.) In contrast, FSTL3 lacks the FSD3 domain and engages Activin A using several specific contacts, which include direct contacts with the finger 2 tip loop. Specifically, the ring nitrogen of His91 in FSTL3 makes a charge contact with D406 of Activin A, and both Ser221 and Arg225 directly interact electrostatically with Activin A Q408 (Cash et al., 2012). Hence, the reduced inhibition of these Activin A muteins by FSTL3 is likely a consequence of their both lacking D406.

Activin A FT2L muteins still activate ACVR1[R206H]

Given that both Activin A.Nod.F2TL and Activin A.ΔD406 display a greatly reduced ability to engage ACVR1, we tested whether they can still activate ACVR1[R206H] (i.e. the most common FOP-causing variant, which utilizes Activins as agonists (Hatsell et al., 2015; Hino et al., 2015)). Both muteins were found to transduce a signal through ACVR1[R206H] (Fig. S12). This indicates that they retain enough binding to ACVR1 to induce signaling, but not to the level required to form a functional NSC. This result is corroborated by the observation that both muteins can still dimerize ACVR1 with ACVR2A to some extent, though not to the degree that Activin A does (Fig. 3B, and S6). In line with these results, structural modeling of Activin A.Nod.F2TL suggests that D405 can shift into the type I interface to compensate for the loss of D406, albeit interacting with only one lysine residue (K78), and hence resulting a weakened interaction (Fig. S1). Furthermore, we found that in this assay Activin A.Nod.F2TL activates ACVR1[R206H] more than wild type Activin A, a finding consistent with the inability of Activin A.Nod.F2TL to form the NSC.

Activin A FT2L mutein results in increased HO in mouse FOP

Given that Activin A.Nod.F2TL escapes recruitment into NSCs yet can activate ACVR1[R206H] (presumably by transient engagement of ACVR1 – see Fig. 3B), we hypothesized that in an FOP setting Activin A.Nod.F2TL will generate more heterotopic bone than wild type Activin A. To test this hypothesis, we induced HO by implantation of Activin A or Activin A.Nod.F2TL adsorbed collagen sponges into *Acvr1[R206H]FlEx/+* (*Acvr1tm2.1Vlcg/+*; MGI:5763014) mice (FOP mice). Consistent with previous observations, FOP mice implanted with sponges containing Activin A develop HO by two weeks following implantation (Hatsell et al., 2015). As would be expected for any Activin mutein that escapes recruitment into the NSC (yet still activates ACVR1[R206H]), we observed more HO in collagen sponges containing Activin A.Nod.F2TL compared to wild type Activin A, in fact approximately six times greater (Fig. 6A, B). These observations were confirmed using fibro/adipogenic progenitors (FAPs), which are the cells that form HO in FOP (Dey et al., 2016; Lees-Shepard, Yamamoto, et al.,

2018). FAPs from FOP mice were transplanted intramuscularly into SCID mice along with either Activin A or with Activin A.Nod.F2TL. When FOP FAPs were co-delivered with Activin A.Nod.F2TL, the resulting HO was also approximately six times the volume of HO obtained from FOP FAPs transplanted in the presence of wild type Activin A (Fig. 6C, D). These results are consistent with a model where Activin A.Nod.F2TL engages the FOP-causing Acvr1[R206H] receptor more frequently than wild type Activin A, simply because the latter can be trapped in the NSC whereas Activin A.Nod.F2TL cannot. This increased frequency in engagement of Acvr1[R206H] by Activin A.Nod.F2TL results in increased signaling in the FAPs, and increased HO. These results are the first conclusive evidence that the NSC is operational *in vivo* where it negatively regulates the signaling that drives HO in FOP. They are also consistent with previous observations where deletion of wild type *Acvr1* results in massively increased heterotopic bone formation in FOP mice (Lees-Shepard, Yamamoto, et al., 2018).

Discussion
The discovery of the ACVR1•Activin A•type II receptor NSC was surprising and unprecedented. Unlike engagement of ligands or receptors by non-signaling or signal-suppressing decoy receptors (Bonecchi, Garlanda, Mantovani, & Riva, 2016; Boyce & Xing, 2007; Mishra, Funair, Cressley, Gittes, & Burns, 2012; Onichtchouk et al., 1999; Palomo, Dietrich, Martin, Palmer, & Gabay, 2015; Sheikh & Fornace, 2000), or inhibitory ligands that naturally form 'dead-end' complexes with signal-transducing receptors (Kokabu et al., 2012; Makanji et al., 2014), the ACVR1•Activin A•type II receptor NSC is unique in that it stoichiometrically mimics signaling complexes utilizing the same type I and type II receptors, albeit with a different (not Activin) ligand class – the BMPs. Even more surprising was the finding that the FOP-causing missense mutations of ACVR1 (which localize in the intracellular domain of this receptor) convert the NSC into a signaling complex, much like that formed by BMPs. The molecular mechanism that underlies this conversion is a matter of debate (Hatsell et al., 2015; Hino et al., 2015). That aside, the realization that the NSC effectively reduces the availability of all the partners involved in that complex (Activin A, ACVR1, and the type II receptors) and hence inhibits or reduces signaling by BMPs via the same receptors, propelled us to ask whether the NSC has biological roles. This question has, thus far, been largely unexplored as the majority of studies on Activin A have focused on its role as a signal-inducing ligand via ACVR1B (Makanji et al., 2014; Namwanje & Brown, 2016).

In order to explore the biological role(s) of the NSC *in vivo,* we engineered agonist-only variants of Activin A (i.e. variants that retain their ability to signal through ACVR1B but fail to form the NSC with ACVR1). To design such muteins we made use of biochemical information regarding receptor utilization by Activin family ligands, prior mutagenesis studies, and structural modeling. We chose Nodal as the main donor for replacements of type I-binding regions of Activin A, a choice driven by the fact that Nodal utilizes the same type II receptors as Activin A along with ACVR1B but does not engage ACVR1. We then focused on three regions for engineering the different chimeric ligands: the pre-helix, the post-helix, and the finger 2 tip loop (F2TL) (see Results section). Of the resulting Activin A-Nodal chimeras, Activin A.Nod.F2TL – a mutein where the finger 2 tip loop was substituted with that of Nodal – displayed the desired properties: it activates ACVR1B nearly to the same degree as wild type Activin A, does not switch to other

type I receptors, and escapes engagement into the NSC. In addition, guided by structural models (Fig. S1), we engineered a single amino acid deletion variant, Activin A.ΔD406. This variant also displayed reduced ability to form the NSC, indicating the importance of D406 as a determinant of the interaction of Activin A with ACVR1.

In parallel with our finding that the F2TL region is required for the formation of the NSC, the importance of the F2TL as a major determinant of type I receptor choice was concurrently demonstrated for GDF11's interaction with TGFBR1 (Goebel et al., 2019). Both findings are further supported by an additional chimeric ligand that we engineered using GDF8 as the 'donor' to generate the corresponding F2TL variant – Activin A.GDF8.F2TL. This variant utilizes TGFBR1 as the type I receptor to a greater degree than Activin A, while retaining its ability to engage ACVR1B (Fig. S13). Therefore, our results lend further credence to the concept that the finger 2 tip loop is a key determinant of type I receptor choice in the Activin class.

Our findings show that the NSC is mediated through very specific contacts between Activin A and ACVR1. Nonetheless, we note that we did not attempt an exhaustive search to define all possible Activin A muteins with the desired properties, and hence it is possible that other such muteins can be engineered. Along these lines, since Activin B and Activin AB are also capable of forming NSCs with ACVR1 and the corresponding type II receptors (Fig. S14), our findings should enable exploration of the formation of the respective complexes by following similar methodology.

Both of the two muteins that we study in detail here deviate from our initial design specifications in that they display a small increase in resistance to inhibition by one of Activin's natural antagonists, FSTL3 (Fig. 5). This property is unavoidable for Activin A variants that bear changes in the finger 2 tip loop, as that region is involved in one of the main contacts between Activin A and FSTL3. Whether this deviation will have physiological consequences, remains to be seen. It should be noted however, that FSTL3 knockout mice have a relatively benign phenotype that has been well characterized and that has been attributed to an increase in Activin A activity (Mukherjee et al., 2007). Given the relatively small reduction in the ability of Activin A.Nod.F2TL to be inhibited by FSTL3 (compared to what would be expected from a complete loss of FSTL3), it is unlikely that this difference will have phenotypic consequences. Nonetheless, the availability of phenotypic data on FSTL3 knockout mice should facilitate the assessment of any differences that might arise in the physiology of mice bearing Activin A.Nod.F2TL as a knock-in into *Inhba*.

The development of an agonist-only Activin A mutein, Activin A.Nod.F2TL, enabled us to directly query the role of the NSC in FOP, and hence build upon the prior discovery of the role of Activin A in this disease. Because of the ability of Activin A.Nod.F2TL to escape recruitment into NSCs, and hence not allow sequestration of both itself and the type II receptors into inactive complexes that render them inaccessible to participate in signaling, we predicted that Activin A.Nod.F2TL should precipitate more HO compared to wild type Activin A. We demonstrate this to be the case in two different yet complimentary settings. These results provide the first evidence for a role of the NSC *in vivo*. Specifically for FOP, the also provide the first direct evidence that the NSC tempers HO in FOP, effectively reducing the amount of heterotopic bone that forms (presumably by reducing the availability of Activin A and type II receptors to form

signaling complexes with FOP-mutant ACVR1). Our data is in agreement with the observation that deletion of wild type *Acvr1* in a mouse FOP setting results in massively increased heterotopic bone (Lees-Shepard, Yamamoto, et al., 2018). The fact that Activin A.Nod.F2TL was successfully employed to query a pathophysiological process *in vivo* sets the stage for future experimentation with the corresponding knock-in mouse line outside of FOP. Given the level of evolutionary conservation in this receptor-ligand system, our data opens the field of investigating whether the ability of forming the NSC was an early or a late event during evolution.

Experimental Procedures

Antibodies and protein reagents
Activin A antibodies H4H10442 and REGN2476 are human monoclonal antibodies specific to Activin A and utilizing an IgG4 constant region. They were generated using Regeneron's VelocImmune® mice (Macdonald et al., 2014; Murphy et al., 2014).

Cloning, expression and purification of Activin A muteins
The mature regions of Activin A or Activin A muteins were cloned downstream of mouse *Inhba* pro-peptide using isothermic Gibson assembly (Gibson et al., 2010) of synthetic gene blocks (Integrated DNA Technologies) into a CMV-based mammalian expression vector. The corresponding expression vectors were introduced into CHO-K1 cells expressing human Furin (CHO-KI.Furin). The resulting conditioned media was concentrated approximately ten-fold using a filtration column with a 3K molecular weight cut off (Pierce) before biochemical characterization of the Activin A muteins in cell-based assays. Following this primary analysis, Activin A and the Activin A F2TL mutants were stably expressed in CHO-K1.Furin cells. The Activin A pro-peptide complex was first purified to homogeneity by heparin chromatography followed by size exclusion chromatography on a preparative scale S200 column. To isolate the mature Activin A mutein dimers, reverse phase chromatography on C4 column was used, and mature Activin A was eluted with an acetonitrile gradient.

Structural modeling and alignments
ClustalW alignment in MacVector was used for amino acid sequence alignments. Structural alignments using known crystal structures as a reference were carried out using STRAP, a web-based alignment program (Gille, Fahling, Weyand, Wieland, & Gille, 2014). All figures with structural models were generated in PyMol (The PyMol Molecular Graphics System, Schrödinger, LLC, New York, NY). The model of ACVR1 was built using Swiss-model (Waterhouse et al., 2018). After manual adjustment that resulted in no clashes and contained realistic contacts, energy minimization of amino acid side chains was performed utilizing YASARA (Krieger et al., 2009).

Binding kinetics measurements
Kinetic binding parameters for the interaction of human ACVR2B.hFc fusion protein with recombinant Activin A and Activin A muteins were determined under neutral pH on Biacore 3000 using dextran-coated (CM5) chips at 25°C. The running buffer was prepared using filtered HBS-ET (10 mM Hepes, 150 mM NaCl, 3.4mM EDTA, 0.05% polysorbate 20, pH 7.4). The capture sensor surface was prepared by covalently immobilizing Protein A (Sigma-Aldrich, St. Louis, MO)

to the chip surface using (1¬Ethyl-3-[3-dimethylaminopropyl]carbodiimide-hydrochloride)/N-hydroxysuccinimide (EDC/NHS) coupling chemistry. Binding kinetics of Activin A and Activin A muteins were measured by flowing 100nM- 0.14nM of ligand, serially diluted three-fold, at 100uL/minute for two minutes and monitored for dissociation for 15 minutes. All capture surfaces were regenerated with one 30-s pulse of 10 mM glycine–HCl (pH 1.5, (GE Healthcare, Marlborough, MA). Kinetic parameters were obtained by globally fitting the data to a 1:1 binding model with mass transport limitation using Scrubber 2.0c Evaluation Software. The equilibrium dissociation constant (KD) was calculated by dividing the dissociation rate constant (kd) by the association rate constant (ka). Data is representative

Cell culture and cell-based assays
Unless otherwise noted, cells were cultured in Dulbecco's modified Eagle's medium containing 10% (v/v) fetal bovine serum (FBS), penicillin/streptomycin (50 U/ml), and 2 mM L-glutamine. Generation of ACVR1 and ACVR1.R206H expressing HEK293 BRE-luciferase reporter cells has been previously described (Hatsell et al., 2015). For this study, a reporter construct with twelve tandem repeats of the $(CAGACagc)_{12}$ response element was introduced into HEK293 cells to engineer the HEK293 CAGA-luciferase reporter line (Latres et al., 2015).

Cell based assays were conducted as previously described (Hatsell et al., 2015). Briefly, HEK293 reporter cells were plated in 96-well plates. After a 16-hour incubation with ligands alone, ligand mixtures or ligand/antagonist mixtures, luciferase expression was measured with Bright-Glo Luciferase Assay System (Promega).

Human U2OS cell lines for receptor dimerization assays were purchased from DiscoverX (Fremont, CA) and used following manufacturers protocols. Briefly, cells were treated with ligands for 16 hours. β-galactosidase enzyme fragment complementation was quantified by a luminescent signal following the addition of the manufacturer-supplied substrate.

In vivo implantation of wild type Activin A- or Activin A.Nod.FT2L mutant-containing collagen sponges
Collagen sponge containing recombinant proteins were implanted as described (Hatsell et al., 2015). Briefly, 4 mm diameter collagen sponges (SpongeCol, Advanced BioMatrix, Carlsbad, CA) were adsorbed with 10 µl suspension of either 20 µg of rhActivin A (R&D Systems, Minneapolis, MN) , 20 µg of Activin A.Nod.FT2L mutant protein, or saline for 30 min at room temperature prior to implantation. Mice were anesthetized with Xylazine and Ketamine. Their right gluteus region was shaved and wiped with alcohol and betadine. Muscle tissue was exposed by making a small incision on the skin and fascia. Muscle fibers were separated using blunt forceps to create a pouch. Collagen sponges were implanted into the muscle pouch and separated muscle fibers were sutured with 5.0 absorbable suture (Visorb, CP Medical, Inc., Norcross, GA). Skin incisions were closed using instant tissue adhesive (Surgi-Lock 2oc, Meridian Animal Health, Omaha, NE). Mice were given two doses of analgesic, Buprenex at a 0.1 mg/kg dose. Two days after implantation, $Acvr1^{[R206H]FlEx/+}$; $Gt(ROSA26)Sor^{CreERT2/+}$ mice were injected with tamoxifen as described (Hatsell et al., 2015)to initiate the model. Heterotopic bone formation was assessed at 2 weeks after implantation by *in vivo* µCT imaging. Mice were assigned to groups without randomization.

Fibro/adipogenic progenitor (FAP) isolation and expansion

Details of skeletal muscle dissection have been previously described (Lees-Shepard, Nicholas, et al., 2018). Dissected muscle was dissociated from tamoxifen treated *Acvr1[R206H]FlEx/+*; *Gt(ROSA26)Sor[CreERT2/+]* mice as described (Hatsell et al., 2015)for tamoxifen regime using the Skeletal Muscle Dissociation Kit (Miltenyi Biotec, Bergisch Gladbach, Germany) and gentleMACS Octo Dissociator with heaters (Miltenyi Biotec), in accordance with manufacturer instructions. Following centrifugation at 300g for 10 min at 4°C, the supernatant was discarded and the pellet was resuspended in growth media (Dulbecco's Modified Eagle Medium (DMEM; Life Technologies, Carlsbad, CA) with 50 U/mL Penicillin and 50 µg/mL Streptomycin (Gibco, Billings, MT) and 20% fetal bovine serum (FBS; Lot# 192K18; Avantor, Randor, PA)). Cells were then plated onto tissue culture flasks (Corning, Corning, NY).

Magnetic-activated cell sorting (MACS) was performed on single cells incubated with anti-mouse Pdgfra APC (clone APA5; eBioscience, San Diego, CA) to label FAPs, followed by magnetic labeling with an Anti-APC MicroBeads kit (Miltenyi Biotec). Cells were then applied to magnetic MS columns (Miltenyi Biotec), in accordance with manufacturer instructions. MACS-isolated FAPs were seeded at a density of 2000 cells/cm^2 onto tissue culture flasks (Corning) in growth media and maintained at 37°C in a humidified atmosphere at 5% CO_2. Media was changed every other day. Prior to transplantation, FAPs were treated with 100nM (*Z*)-4-hydroxytamoxifen (Sigma-Aldrich) for 24 hours to enhance inversion of the R206H-containing exon.

Transplantation and drug treatment

MACS-isolated and expanded FAPs were resuspended at 1×10^7 cells/mL in ice-cold 1X Dulbecco's Phosphate-Buffered Saline (DPBS; Gibco) or Geltrex (LDEV-Free hESC qualified Reduced Growth Factor; Gibco) with or without 2.5µg Activin A or Activin A.Nod.F2TL, as indicated in the corresponding figure. A 50µL volume containing 2×10^5 FAPs was injected into the gastrocnemius or triceps muscle of SCID mice (JAX Stock No: 001913, The Jackson Laboratory, Bar Harbor, ME), as previously described. Mice were assigned to groups without randomization.

µCT imaging analyses and measurements

Heterotopic bone formation was assessed as described previously (Hatsell et al., 2015), Briefly, mice were anesthetized with isoflurane and whole body–scanned, with a field of view at 60 mm × 120 mm, using *in vivo* µCT (Quantum GX, PerkinElmer). The x-ray source was set to a current of 88 µA, voltage of 90 kVp, with a voxel size at 240 mm. The CT imaging was visualized and quantified using Analyze software (Analyze Direct, Overland Park, KS). Whole-body bone tissue was auto segmented by threshold, the heterotopic bone was manually segmented from the original bone, and then bone volume calculated using Analyze. HO volume following transplantation was quantified as previously described (Lees-Shepard, Yamamoto, et al., 2018). Mouse group information was blinded for HO analysis.

Statistical analysis

Statistical analysis was performed using GraphPad Prism (GraphPad, La Jolla, CA). All numerical values are presented as mean ± standard deviation. One-way ANOVA was used to

determine significance, as described in the corresponding figure legends. Differences were considered significant at $p < 0.05$.

Conflict of interest

VI, HK, QZ, AR, RL, XW, JM, JJ, KN, NJR, JBLS, AJM, and ANE are employees of Regeneron Pharmaceuticals and own common stock as well as stock options of Regeneron shares. The remaining authors have no conflicts of interest to declare with respect to the contents of this article.

Author Contributions

VI and ANE conceived of the project. VI, ANE, EJG, KN, JBLS, AJM, and TBT designed experiments and were involved in the interpretation of results and wrote the manuscript. RAC, EJG, CJC, AD, HK, QZ, AR, RL, XW, JJ, NJR, JBLS, and JM performed experiments and contributed data.

References

Allendorph, G. P., Isaacs, M. J., Kawakami, Y., Izpisua Belmonte, J. C., & Choe, S. (2007). BMP-3 and BMP-6 structures illuminate the nature of binding specificity with receptors. *Biochemistry, 46*(43), 12238-12247. doi:10.1021/bi700907k

Allendorph, G. P., Vale, W. W., & Choe, S. (2006). Structure of the ternary signaling complex of a TGF-beta superfamily member. *Proc Natl Acad Sci U S A, 103*(20), 7643-7648. doi:10.1073/pnas.0602558103

Andersson, O., Reissmann, E., & Ibanez, C. F. (2006). Growth differentiation factor 11 signals through the transforming growth factor-beta receptor ALK5 to regionalize the anterior-posterior axis. *EMBO Rep, 7*(8), 831-837. doi:10.1038/sj.embor.7400752

Archambeault, D. R., & Yao, H. H. (2010). Activin A, a product of fetal Leydig cells, is a unique paracrine regulator of Sertoli cell proliferation and fetal testis cord expansion. *Proc Natl Acad Sci U S A, 107*(23), 10526-10531. doi:10.1073/pnas.1000318107

Attisano, Liliana, Cárcamo, Juan, Ventura, Francesc, Weis, Frances M. B., Massagué, Joan, & Wrana, Jeffrey L. (1993). Identification of human activin and TGFβ type I receptors that form heteromeric kinase complexes with type II receptors. *Cell, 75*(4), 671-680. doi:http://dx.doi.org/10.1016/0092-8674(93)90488-C

Aykul, S., Parenti, A., Chu, K. Y., Reske, J., Floer, M., Ralston, A., & Martinez-Hackert, E. (2017). Biochemical and Cellular Analysis Reveals Ligand Binding Specificities, a Molecular Basis for Ligand Recognition, and Membrane Association-dependent Activities of Cripto-1 and Cryptic. *J Biol Chem, 292*(10), 4138-4151. doi:10.1074/jbc.M116.747501

Bonecchi, R., Garlanda, C., Mantovani, A., & Riva, F. (2016). Cytokine decoy and scavenger receptors as key regulators of immunity and inflammation. *Cytokine, 87*, 37-45. doi:10.1016/j.cyto.2016.06.023

Boyce, B. F., & Xing, L. (2007). Biology of RANK, RANKL, and osteoprotegerin. *Arthritis Res Ther, 9 Suppl 1*, S1. doi:10.1186/ar2165

Cash, J. N., Angerman, E. B., Kattamuri, C., Nolan, K., Zhao, H., Sidis, Y., . . . Thompson, T. B. (2012). Structure of myostatin.follistatin-like 3: N-terminal domains of follistatin-type molecules exhibit alternate modes of binding. *J Biol Chem, 287*(2), 1043-1053. doi:10.1074/jbc.M111.270801

Cash, J. N., Rejon, C. A., McPherron, A. C., Bernard, D. J., & Thompson, T. B. (2009). The structure of myostatin:follistatin 288: insights into receptor utilization and heparin binding. *EMBO J, 28*(17), 2662-2676. doi:10.1038/emboj.2009.205

Dey, Devaveena, Bagarova, Jana, Hatsell, Sarah J., Armstrong, Kelli A., Huang, Lily, Ermann, Joerg, . . . Yu, Paul B. (2016). Two tissue-resident progenitor lineages drive distinct phenotypes of heterotopic ossification. *Sci Transl Med, 8*(366), 366ra163-366ra163. doi:10.1126/scitranslmed.aaf1090

Gibson, D. G., Glass, J. I., Lartigue, C., Noskov, V. N., Chuang, R. Y., Algire, M. A., . . . Venter, J. C. (2010). Creation of a bacterial cell controlled by a chemically synthesized genome. *Science, 329*(5987), 52-56. doi:10.1126/science.1190719

Gille, C., Fahling, M., Weyand, B., Wieland, T., & Gille, A. (2014). Alignment-Annotator web server: rendering and annotating sequence alignments. *Nucleic Acids Res, 42*(Web Server issue), W3-6. doi:10.1093/nar/gku400

Goebel, E. J., Corpina, R. A., Hinck, C. S., Czepnik, M., Castonguay, R., Grenha, R., . . . Thompson, T. B. (2019). Structural characterization of an activin class ternary receptor complex reveals a third paradigm for receptor specificity. *Proc Natl Acad Sci U S A, 116*(31), 15505-15513. doi:10.1073/pnas.1906253116

Greenwald, J., Vega, M. E., Allendorph, G. P., Fischer, W. H., Vale, W., & Choe, S. (2004). A flexible activin explains the membrane-dependent cooperative assembly of TGF-beta family receptors. *Mol Cell, 15*(3), 485-489. doi:10.1016/j.molcel.2004.07.011

Harrington, A. E., Morris-Triggs, S. A., Ruotolo, B. T., Robinson, C. V., Ohnuma, S., & Hyvonen, M. (2006). Structural basis for the inhibition of activin signalling by follistatin. *EMBO J, 25*(5), 1035-1045. doi:10.1038/sj.emboj.7601000

Harrison, C. A., Gray, P. C., Fischer, W. H., Donaldson, C., Choe, S., & Vale, W. (2004). An activin mutant with disrupted ALK4 binding blocks signaling via type II receptors. *J Biol Chem, 279*(27), 28036-28044. doi:10.1074/jbc.M402782200

Hatsell, S. J., Idone, V., Wolken, D. M., Huang, L., Kim, H. J., Wang, L., . . . Economides, A. N. (2015). ACVR1R206H receptor mutation causes fibrodysplasia ossificans progressiva by imparting responsiveness to activin A. *Sci Transl Med, 7*(303), 303ra137. doi:10.1126/scitranslmed.aac4358

Hinck, A. P., Mueller, T. D., & Springer, T. A. (2016). Structural Biology and Evolution of the TGF-beta Family. *Cold Spring Harb Perspect Biol, 8*(12). doi:10.1101/cshperspect.a022103

Hino, K., Ikeya, M., Horigome, K., Matsumoto, Y., Ebise, H., Nishio, M., . . . Toguchida, J. (2015). Neofunction of ACVR1 in fibrodysplasia ossificans progressiva. *Proc Natl Acad Sci U S A, 112*(50), 15438-15443. doi:10.1073/pnas.1510540112

Huning, I., & Gillessen-Kaesbach, G. (2014). Fibrodysplasia ossificans progressiva: clinical course, genetic mutations and genotype-phenotype correlation. *Mol Syndromol, 5*(5), 201-211. doi:10.1159/000365770

Katagiri, T., Tsukamoto, S., & Kuratani, M. (2018). Heterotopic bone induction via BMP signaling: Potential therapeutic targets for fibrodysplasia ossificans progressiva. *Bone, 109*, 241-250. doi:10.1016/j.bone.2017.07.024

Kokabu, S., Gamer, L., Cox, K., Lowery, J., Tsuji, K., Raz, R., . . . Rosen, V. (2012). BMP3 suppresses osteoblast differentiation of bone marrow stromal cells via interaction with Acvr2b. *Mol Endocrinol, 26*(1), 87-94. doi:10.1210/me.2011-1168

Korupolu, R. V., Muenster, U., Read, J. D., Vale, W., & Fischer, W. H. (2008). Activin A/bone morphogenetic protein (BMP) chimeras exhibit BMP-like activity and antagonize activin and myostatin. *J Biol Chem, 283*(7), 3782-3790. doi:10.1074/jbc.M704530200

Krieger, E., Joo, K., Lee, J., Lee, J., Raman, S., Thompson, J., . . . Karplus, K. (2009). Improving physical realism, stereochemistry, and side-chain accuracy in homology modeling: Four approaches that performed well in CASP8. *Proteins, 77 Suppl 9*, 114-122. doi:10.1002/prot.22570

Latres, E., Pangilinan, J., Miloscio, L., Bauerlein, R., Na, E., Potocky, T. B., . . . Stitt, T. (2015). Myostatin blockade with a fully human monoclonal antibody induces muscle hypertrophy and reverses muscle atrophy in young and aged mice. *Skelet Muscle, 5*, 34. doi:10.1186/s13395-015-0060-8

Lees-Shepard, J. B., Nicholas, S. E., Stoessel, S. J., Devarakonda, P. M., Schneider, M. J., Yamamoto, M., & Goldhamer, D. J. (2018). Palovarotene reduces heterotopic ossification in juvenile FOP mice but exhibits pronounced skeletal toxicity. *Elife, 7*. doi:10.7554/eLife.40814

Lees-Shepard, J. B., Yamamoto, M., Biswas, A. A., Stoessel, S. J., Nicholas, S. E., Cogswell, C. A., . . . Goldhamer, D. J. (2018). Activin-dependent signaling in fibro/adipogenic progenitors causes fibrodysplasia ossificans progressiva. *Nat Commun, 9*(1), 471. doi:10.1038/s41467-018-02872-2

Macdonald, L. E., Karow, M., Stevens, S., Auerbach, W., Poueymirou, W. T., Yasenchak, J., . . . Murphy, A. J. (2014). Precise and in situ genetic humanization of 6 Mb of mouse immunoglobulin genes. *Proc Natl Acad Sci U S A, 111*(14), 5147-5152. doi:10.1073/pnas.1323896111

Macías-Silva, Marina, Hoodless, Pamela A., Tang, Shao Jun, Buchwald, Manuel, & Wrana, Jeffrey L. (1998). Specific Activation of Smad1 Signaling Pathways by the BMP7 Type I Receptor, ALK2. *Journal of Biological Chemistry, 273*(40), 25628-25636. doi:10.1074/jbc.273.40.25628

Makanji, Y., Zhu, J., Mishra, R., Holmquist, C., Wong, W. P., Schwartz, N. B., . . . Woodruff, T. K. (2014). Inhibin at 90: from discovery to clinical application, a historical review. *Endocr Rev, 35*(5), 747-794. doi:10.1210/er.2014-1003

Matzuk, M. M., Kumar, T. R., Vassalli, A., Bickenbach, J. R., Roop, D. R., Jaenisch, R., & Bradley, A. (1995). Functional analysis of activins during mammalian development. *Nature, 374*(6520), 354-356. doi:10.1038/374354a0

Mishra, S. K., Funair, L., Cressley, A., Gittes, G. K., & Burns, R. C. (2012). High-affinity Dkk1 receptor Kremen1 is internalized by clathrin-mediated endocytosis. *PLoS One, 7*(12), e52190. doi:10.1371/journal.pone.0052190

Mukherjee, A., Sidis, Y., Mahan, A., Raher, M. J., Xia, Y., Rosen, E. D., . . . Schneyer, A. L. (2007). FSTL3 deletion reveals roles for TGF-beta family ligands in glucose and fat homeostasis in adults. *Proc Natl Acad Sci U S A, 104*(4), 1348-1353. doi:10.1073/pnas.0607966104

Murphy, A. J., Macdonald, L. E., Stevens, S., Karow, M., Dore, A. T., Pobursky, K., . . . Yancopoulos, G. D. (2014). Mice with megabase humanization of their immunoglobulin

genes generate antibodies as efficiently as normal mice. *Proc Natl Acad Sci U S A, 111*(14), 5153-5158. doi:10.1073/pnas.1324022111

Namwanje, Maria, & Brown, Chester W. (2016). Activins and Inhibins: Roles in Development, Physiology, and Disease. *Cold Spring Harbor Perspectives in Biology, 8*(7). doi:10.1101/cshperspect.a021881

Olsen, O. E., Wader, K. F., Hella, H., Mylin, A. K., Turesson, I., Nesthus, I., . . . Holien, T. (2015). Activin A inhibits BMP-signaling by binding ACVR2A and ACVR2B. *Cell Commun Signal, 13*, 27. doi:10.1186/s12964-015-0104-z

Onichtchouk, D., Chen, Y. G., Dosch, R., Gawantka, V., Delius, H., Massague, J., & Niehrs, C. (1999). Silencing of TGF-beta signalling by the pseudoreceptor BAMBI. *Nature, 401*(6752), 480-485. doi:10.1038/46794

Palomo, J., Dietrich, D., Martin, P., Palmer, G., & Gabay, C. (2015). The interleukin (IL)-1 cytokine family--Balance between agonists and antagonists in inflammatory diseases. *Cytokine, 76*(1), 25-37. doi:10.1016/j.cyto.2015.06.017

Pangas, S. A., Jorgez, C. J., Tran, M., Agno, J., Li, X., Brown, C. W., . . . Matzuk, M. M. (2007). Intraovarian activins are required for female fertility. *Mol Endocrinol, 21*(10), 2458-2471. doi:10.1210/me.2007-0146

Rebbapragada, A., Benchabane, H., Wrana, J. L., Celeste, A. J., & Attisano, L. (2003). Myostatin signals through a transforming growth factor beta-like signaling pathway to block adipogenesis. *Mol Cell Biol, 23*(20), 7230-7242.

Sheikh, M. S., & Fornace, A. J., Jr. (2000). Death and decoy receptors and p53-mediated apoptosis. *Leukemia, 14*(8), 1509-1513. doi:10.1038/sj.leu.2401865

Stamler, R., Keutmann, H. T., Sidis, Y., Kattamuri, C., Schneyer, A., & Thompson, T. B. (2008). The structure of FSTL3.activin A complex. Differential binding of N-terminal domains influences follistatin-type antagonist specificity. *J Biol Chem, 283*(47), 32831-32838. doi:10.1074/jbc.M801266200

Thompson, T. B., Lerch, T. F., Cook, R. W., Woodruff, T. K., & Jardetzky, T. S. (2005). The structure of the follistatin:activin complex reveals antagonism of both type I and type II receptor binding. *Dev Cell, 9*(4), 535-543. doi:10.1016/j.devcel.2005.09.008

Tsuchida, K., Mathews, L. S., & Vale, W. W. (1993). Cloning and characterization of a transmembrane serine kinase that acts as an activin type I receptor. *Proc Natl Acad Sci U S A, 90*(23), 11242-11246.

Upadhyay, J., Xie, L., Huang, L., Das, N., Stewart, R. C., Lyon, M. C., . . . Hatsell, S. J. (2017). The Expansion of Heterotopic Bone in Fibrodysplasia Ossificans Progressiva Is Activin A-Dependent. *Journal of bone and mineral research : the official journal of the American Society for Bone and Mineral Research, 32*(12), 2489-2499. doi:10.1002/jbmr.3235

Waterhouse, A., Bertoni, M., Bienert, S., Studer, G., Tauriello, G., Gumienny, R., . . . Schwede, T. (2018). SWISS-MODEL: homology modelling of protein structures and complexes. *Nucleic Acids Res, 46*(W1), W296-W303. doi:10.1093/nar/gky427

Yadin, D., Knaus, P., & Mueller, T. D. (2016). Structural insights into BMP receptors: Specificity, activation and inhibition. *Cytokine Growth Factor Rev, 27*, 13-34. doi:10.1016/j.cytogfr.2015.11.005

Figures

Figure 1: Conceptual framework for designing 'agonist-only' Activin A muteins
(A) In a luciferase reporter assay, agonist only Activin A muteins (blue) should retain activation of the Smad2/3 pathway like wild type Activin A (red; left panels), and they should retain ability to be inhibited by the endogenous antagonists Follistatin and FSTL3 (grey; right panels). (B) As a result of a loss of ACVR1 binding, Activin A muteins should be less effective inhibitors of BMP-mediated signaling to the Smad1/5/8 pathway. Therefore, we expect that the agonist only muteins should have reduced antagonism of Smad1/5/8 signaling compared to wild type Activin A. However, antagonism will not be entirely lost, as the agonist-only mutein must still bind to Type II receptor for activation of Smad2/3 signaling.

A

B
```
INHBA: (311)  GLE..CDGKVNICCKKQFFVSFKDIGWNDWIIAPSGYHANYCEGEC PSHIAGTSGSSLS FHSTVINHYRMR GHSPPANLK.S CCVPTKLRPMSMLYY DDGQN IKKDIQNMIVEECGCS (426)
Nodal: (238)  HHLPDR....SQLCRKVKFQVDFNLIGWGSWIIYPKQYNAYRCEGEC PNPVGEEFHPTNH ..AYIQSLLKRY QPHRVP...ST CCAPVKTKPLSMLYV D.NGR LLDHHKDMIVEECGCL (347)
```

Pre-helix Loop   **Post-helix Loop**   **Tip Loop (F2TL)**

C
Type II Interface

90°

Type I Interface

Figure 2: Activin A muteins were engineered with the Activin A pre-helix, post-helix, and F2TL regions replaced with the corresponding regions of Nodal

(A) Ribbon diagram of Activin A. (B) Structural alignment of human Activin A and Nodal that highlights the Activin A pre-helix, post-helix and finger 2 tip loop (F2TL) sequences used to generate agonist-only Activin A muteins. Nodal sequences highlighted in the boxed areas were substituted for the corresponding sequences of Activin A. (C) Crystal structure of FSTL3-bound Activin A (space filled model, PDB 3B4V) with substituted areas colored as follows: pre-helix loop (orange), post-helix loop (red) and F2TL (magenta). Each Activin A monomer is depicted in either light or dark blue. Notice that these substitutions occupy the type I receptor binding site.

Figure 3: Activin A F2TL mutants signal normally through the Smad2/3 pathway, but have lost binding to Acvr1

(A) Activin A F2TL muteins activate Smad2/3 signaling to similar levels as wild type Activin A in HEK293 cells harboring a Smad2/3 reporter construct driving firefly luciferase. (B) U20S cells expressing split beta-galactosidase fusions of corresponding type I and type II receptors were treated with a dose response of BMP7, Activin A, or Activin A.Nod.F2TL. Type 1 receptor binding was measured by luminescence in these receptor dimerization assays. Activin A.Nod.F2TL has reduced ability to dimerize ACVR1:ACVR2A receptors, while retaining wild type capacity to dimerize the ACVR1B:BMPR2 and TGFBR1:ACVR2B receptor pairs. The data presented are representative of at least three independent biological replicates. Three technical replicates were performed per experiment.

Figure 4: The remaining BMP antagonism of Activin A.Nod.F2TL is due to type II receptor binding

(A) Activin A.Nod.F2TL is a less effective inhibitor of BMP7 signaling to Smad1/5/8 than wild type Activin A. HEK293 cells harboring a Smad1/5/8 luciferase reporter construct were treated with varying concentrations of Activin A or Activin A.Nod.F2TL with a constant concentration of BMP7 (12nM) to stimulate Smad1/5/8 signaling. Inhibition of BMP7 is reduced ~60 fold with Activin A.Nod.F2TL compared to Activin A. Using an Activin A antibody that blocks interaction with type II receptor (REGN2476), the remaining inhibition of BMP7 by Activin A.Nod.F2TL is lost. (B) Inhibition of type I receptor binding of Activin A with anti-Activin antibody H4H10442 shows a similar reduction in BMP inhibition to Activin A.Nod.F2TL. (The IC$_{50}$s of Activin A and Activin A.Nod.F2TL are 1.4 x 10^{-9} M and 9.7 x 10^{-8} M, respectively. Insert in Fig. 4A shows a dose response of BMP7 on the HEK293 reporter cells, and the dotted lines represents the Smad1/5/8 signal induced by 12nM BMP7 without inhibition by Activin A.) The data presented are representative of at least three independent biological replicates. Three technical replicates were performed per experiment.

Figure 5: Activin A.Nod.F2TL mutein is inhibited by Follistatin but shows reduced inhibition by FSTL3

Varying concentrations of Follistatin and FSTL3 were preincubated with a constant concentration (10nM) of Activin A or Activin A.Nod.F2TL 'agonist only' mutein. Activity of both Activin A and Activin A.Nod.F2TL were tested in HEK293 cells harboring the Smad2/3 luciferase reporter. Activity of both Activin A and Activin A.Nod.F2TL was blocked by both follistatin-288 (A) and follistatin-315 (B). (C) FSTL3 is a less effective inhibitor of Activin A.Nod.F2TL. The data presented are representative of at least three independent biological replicates. Three technical replicates were performed per experiment.

Figure 6: Activin A.Nod.F2TL mutein is a more potent inducer of HO than wild type Activin A.
(A) Representative µCT images of HO (yellow arrows) from tamoxifen-treated *Acvr1[R206H]FlEx/+; Gt(ROSA26)Sor[CreERT2/+]* mice taken two weeks after implantation of collagen sponges adsorbed with saline, 20 µg wild type Activin A (ActA), or 20 µg Activin A.Nod.F2TL (ActA.Nod.F2TL). (B) Quantification of HO volume 2 weeks post-implantation. Saline, n = 2; ActA, n = 4; ActA.Nod.F2TL, n = 5. Each dot represents a single implantation with group mean (grey bar) and ± standard deviation (error bars) shown. Statistical significance was assessed by one-way ANOVA; **=p ≤ 0.01. (C) Representative µCT images of fore- and hindlimbs from SCID hosts 11 days post-transplantation of FOP FAPs (R206H-FAPs) in Geltrex alone, Geltrex contain 5 µg ActA, or Geltrex containing 5 µg ActA.Nod.F2TL. HO is pseudocolored beige for forelimbs and blue for hindlimbs. (D) Quantification of HO volume 11 days post-transplantation of R206H-FAPs. Geltrex only, n= 8; ActA, n = 14; n = 14; ActA.Nod.F2TL. Each dot represents a single transplantation with group mean (grey bar) and ± standard deviation (error bars) shown. Statistical significance was assessed by one-way ANOVA; ****=p ≤ 0.0001.