Title: Continent-wide effects of urbanization on bird and mammal genetic diversity

Authors: C. Schmidt[1*], M. Domaratzki[2], R.P. Kinnunen[1], J. Bowman[3,4], C.J. Garroway[1*]

Affiliations:

[1]Department Biological Sciences, University of Manitoba, Winnipeg, MB, Canada R3T 2N2
[2]Department of Computer Science, University of Manitoba, Winnipeg, MB, Canada R3T 2N2
[3]Environmental and Life Sciences Graduate Program, Trent University, Peterborough, ON,
Canada K9L 0G2
[4]Wildlife Research and Monitoring Section, Ontario Ministry of Natural Resources and Forestry,
Peterborough, ON, Canada K9J 8M5

*Correspondence to:
Chloé Schmidt
Biological Sciences Building, University of Manitoba
50 Sifton Rd
Winnipeg, MB, Canada R3T 2N2
Email: schmid46@myumanitoba.ca
16
Colin Garroway
Biological Sciences Building, University of Manitoba
50 Sifton Rd
Winnipeg, MB, Canada R3T 2N2
Phone: (204) 474-8267; email: colin.garroway@umanitoba.ca

Abstract: Urbanization and associated environmental changes are causing global declines in

vertebrate populations. In general, population declines of the magnitudes now detected should

lead to reduced effective population sizes for animals living in proximity to humans and

disturbed lands. This is cause for concern because effective population sizes set the rate of

genetic diversity loss due to genetic drift, the rate of increase in inbreeding, and the efficiency

with which selection can act on beneficial alleles. We predicted that the effects of urbanization

should decrease effective population size and genetic diversity, and increase population-level

genetic differentiation. To test for such patterns, we repurposed and reanalyzed publicly archived

genetic data sets for North American birds and mammals. After filtering, we had usable raw

genotype data from 85 studies and 41,023 individuals, sampled from 1,008 locations spanning 41

mammal and 25 bird species. We used census-based urban-rural designations, human population

density, and the Human Footprint Index as measures of urbanization and habitat disturbance. As

predicted, mammals sampled in more disturbed environments had lower effective population

sizes and genetic diversity, and were more genetically differentiated from those in more natural

environments. There were no consistent relationships detectable for birds. This suggests that, in

general, mammal populations living near humans may have less capacity to respond adaptively

to further environmental changes, and be more likely to suffer from effects of inbreeding.

Keywords: data synthesis, urban evolution, habitat fragmentation, human disturbance,

Anthropocene, genetic biodiversity

1. Background

Human activities are among the most prominent and efficient drivers of contemporary evolution (1). In some cases, human-caused evolution in wild populations is well understood and predictable. For instance, we have a well-founded expectation that populations of pests and disease agents will respond adaptively to our attempts at controlling them (1). It is also clear that humans inadvertently alter evolutionary change in wild populations through land use and habitat degradation (2,3). Whether the indirect effects of human activities on evolutionary change can cause predictable evolutionary outcomes is less well understood. We hypothesized that human land use, by limiting population size and fragmenting habitat, reduces effective population size and genetic diversity in wild populations leading to increased genetic differentiation. To investigate this hypothesis, we repurposed and reanalyzed publicly archived raw nuclear genetic data sets for North American birds and mammals to test for general relationships between urbanization and the genetic diversity of populations.

Urbanization is one of the most pervasive causes of habitat fragmentation and general landscape change. In addition to the ~700,000 km^2 occupied by cities (4), nearly 75% of the Earth's land surface has been modified by humans, primarily in support of city dwellers (5). This human-caused degradation of the planet's land surface has consistently reduced its capacity to support wildlife (6). As a result, between 1970 and 2014 vertebrate populations have on average declined in size by ~60% (6). Reductions in population size at this level should decrease genetic diversity by increasing the strength of genetic drift—allele frequency variation due to the random sampling of gametes from one generation to the next. Indeed, there have been general declines in the genetic diversity of populations since the industrial revolution (7). While genetic drift is a

neutral evolutionary process that operates independently of the selective value of alleles, it

reduces the efficiency of deterministic evolutionary processes like selection by causing allele

frequencies to randomly deviate from expected values. When drift is strong relative to selection,

random gamete sampling becomes the predominant cause of allele frequency change. In

addition, increased drift and inbreeding can eventually lead to reduced mean fitness in small

populations. If wildlife populations living in proximity to humans generally experience

reductions in population size and connectivity, and thus increased drift, then they may

systematically become less genetically diverse than those living in less disturbed environments.

By altering a population's genetic composition in this way, human-caused environmental change

could make evolutionary responses to such change less efficient.

The fragmented nature of cities leads to particular expectations about how evolutionary

processes will be altered within them based on population genetic theory (2,8,9) (Fig. 1).

Assuming a finite population of constant size with individuals that randomly mate, die out, and

are completely replaced by their offspring each generation, populations will lose genetic

diversity at a rate inversely proportional to population size. In reality, natural populations always

deviate from these assumptions. Fortunately, we can substitute the concept of effective

population size for census population size and the predictive utility of the theory holds. The

effective population size is the size of an idealized population that conforms to the preceding

assumptions and produces the same rate of drift as observed in the measured population. We can

think of effective population size as a measure of the rate at which genetic drift causes a

population to lose genetic diversity. Nearly all violations of these assumptions cause the effective

population size to be much lower than the census population size, underscoring that drift plays a

more important role in determining genetic diversity and the efficiency of selection than what

might be expected from census population size alone. We predicted that reduced census

population size and gene flow in cities would lead to smaller effective population sizes,

decreased genetic diversity, and increased genetic differentiation in urban populations.

2. Collecting and processing raw publicly archived genetic data

We tested for general relationships between the human modification of terrestrial habitats and

the genetic composition of North American mammals and birds using archived raw

microsatellite data from 85 studies, including 41,023 individuals sampled at 1,008 georeferenced

sample sites, spanning 66 species (Table S1, Table S2). In particular, we studied the effects of

urbanization and the human footprint (10). We conducted a systematic search of online

repositories for all bird and mammal microsatellite data available for North America and applied

a series of filtering steps (see SI Methods, Fig. S5) to build a database of georeferenced neutral

genetic diversity in wild populations. Our approach was made possible due to the accumulation

of data in public data archives, and a still-changing culture of open data in ecological and

evolutionary research. Access to georeferenced raw data originally generated for unrelated

purposes allowed for a particularly powerful synthetic analysis. This was because we could

calculate population genetic parameters and multiple disturbance measures of interest for our

question in a consistent manner, whether or not they were presented in the original publications.

In addition, the fact that these data were collected to address different questions reduces the

likelihood that study system selection—perhaps a tendency to explore evolutionary responses to

humans in systems where such responses are expected—biased our findings.

We chose to analyze data sets that used neutral microsatellite markers because microsatellites

were the most common molecular marker type available in data repositories (11), and because

the evolutionary processes that we are interested in are best measured with neutral markers.

Although the number of loci surveyed in microsatellite studies is often small relative to surveys

of genome-wide markers, the typical number of microsatellites used (~10 loci) in fact estimates

genome-wide diversity well with little gain in accuracy with additional genotyping (12).

Variation in microsatellite loci will likely capture recent, fine-scale changes in population

structure due to their high mutation rates and variability. While questions about adaptive genetic

variation are also interesting, adaptive diversity is currently more difficult to generally define and

interpret than neutral genetic diversity, and there are still relatively few data sets suitable for this

type of multi-population and multi-species analysis.

We tested for effects of urbanization and the human footprint on estimates of four population

genetic parameters calculated for each site: effective population size, gene diversity, allelic

richness, and F_{ST} (196 bird sites, of which 129 sampled non-migratory species and were

reanalyzed separately, and 812 mammal sites, Fig. 2; Tables 1, S1). We estimated contemporary

effective population size of the parental generation using a single sample linkage disequilibrium

method to quantify genetic drift (13–15). Of available methods, this approach is one of the more

accurate and it is relatively robust to departures from underlying assumptions about population

structure (16). Estimators of effective population size perform poorly when sampling error

swamps signals of genetic drift, and this meant that effective population size was not estimable at

some sites, which we excluded from analysis (17 sites excluded for mammals, 1 site excluded for

non-migratory birds; see Table 1 and SI Methods for details). Gene diversity (17) is a measure of

genetic diversity that accounts for the evenness and abundance of alleles, and it is not

significantly affected by sample size or rare alleles (18). We calculated rarefied allelic richness,

the number of alleles per locus corrected for sample size, as a second measure of genetic

diversity (19). To quantify genetic differentiation among sites, we estimated site-specific F_{ST}

(20). At least two sample sites in a given dataset were required to estimate site-specific F_{ST} (20).

3. Modelling strategy

We focused our analyses on the continental United States and Canada due to the historical and

demographic similarities of cities and land-usage in this region (21), and to ensure that species

have had broadly similar exposures to past climate variation (22). We chose three indices of

urbanization and human presence. First, we classified a sample as coming from an urban or rural

site based on United States Census Bureau (23) and Statistics Canada (24) classifications of

urban areas and population centers. Second, we measured human population density at each site,

which may capture aspects of the continuous nature of the effects of human presence that would

not be apparent in the binary urban-rural classification. Lastly, we used the Human Footprint

Index (10) as a measure of human presence because it incorporates data from multiple land use

types including human population density, built-up areas, nighttime lights, land cover, and

human access to coastlines, roads, railways, and navigable rivers.

Current levels of genetic diversity will reflect many past processes in addition to urbanization

and human-caused environmental degradation. Such processes include exposure to Pleistocene

glaciations as well as species-specific life history traits, such as body mass and longevity, each of

which could shape effective population size and thus genetic diversity. Because exposure to past

environments (22,25) and life history trait variation (26) vary spatially, we expect the effects of

such processes to create spatial variation in genetic diversity. We can account for such spatial

patterns by including variables describing spatial patterns in genetic diversity directly in our

models, even when the variables themselves are unmeasured (27). This can be accomplished

with distance-based Moran's Eigenvector Maps, or dbMEMs (27–29). Briefly, dbMEMs are

orthogonal spatially explicit eigenvectors that summarize spatial autocorrelation (Moran's I)

patterns in data across all scales. In our regression models we used dbMEMs that described

spatial variation in our measures of genetic composition to explicitly account for processes

causing spatial patterns in the data (29,30). Neutral genetic diversity also varies with species life

history traits which may lack spatial structure (31). We therefore included species as a random

effect, allowing both slopes and intercepts to vary, in a generalized mixed modeling framework

to capture variation in genetic diversity not already accounted for by dbMEMs (see SI Methods

for details).

We used Bayesian generalized linear mixed models to test for relationships between genetic

diversity and urbanization (32,33). We treated each of our four population genetic parameters

(effective population size, allelic richness, gene diversity, and site-specific F_{ST}) as dependent

variables in a series of regression models. Each genetic parameter was fit to each urbanization

variable (urban-rural, human population density, and Human Footprint) in separate models that

also contained terms for species as a random effect, and spatial variables (dbMEMs) when they

were important descriptors of spatial patterns in genetic data. Finally, we fit a null model to each

population genetic parameter that contained only the random effect for species and spatial

variables. The defining feature of such hierarchical models is that they are models of models –

parameter estimates and intercepts can be estimated for each species, and the distribution of

these species-specific estimates allows us to generalize effects of urbanization across species.

We fit these models for bird and mammal data independently. Migratory behavior in birds may

affect spatial patterns in genetic diversity depending on where samples were taken, and whether

they were sampled during the breeding season. Therefore, we also ran these models separately

for non-migratory birds (7 species, 129 sites; Table S1).

4. Results and Discussion

Relationships between all measures of urbanization and the genetic composition of mammal

populations were consistently in the predicted directions (see the position of parameter estimates

and the breadth of 95% credible intervals in Fig 3a). Effective population size, allelic richness,

and gene diversity tended to be negatively related to the measures of urbanization, and sites

sampled in areas with greater human presence tended to be the most genetically differentiated

(Fig. 3a; Table 1). Contrasting these trends, we found no clear evidence for consistent effects of

urbanization and the human footprint on the genetic composition of non-migratory bird samples

when analyzed alone (Fig. 3b; Table 1), or when migratory and non-migratory species were

combined for analyses (Fig. S1).

To assess model fits we estimated marginal R^2 (R^2_m), the variance explained by the fixed effects,

and conditional R^2 (R^2_c), the variance explained by both fixed and random effects (34). For

mammals, all models containing indices of human disturbance explained more variation in the

genetic composition of populations than null models (Table 1). Human population density

explained the most variation in each measure of the genetic composition of mammal sample sites

(effective population size: R^2_m 0.28; R^2_c 0.25; gene diversity R^2_m 0.12; R^2_c 0.70; allelic richness

R^2_m 0.07; R^2_c 0.70; F_{ST} R^2_m 0.22; R^2_c 0.35) except allelic richness, where explained variance was

similar among all urban predictors (Table 1).

The lack of evidence for genetic effects of urbanization on birds may in part be due to the limited

number of data sets available compared to data availability for mammals. Data from seven

species remained after excluding migratory species: the California scrub jay (*Aphelocoma*

californica), black-capped chickadee (*Poecile atricapillus)*, boreal chickadee (*P. hudsonicus)*,

barn owl (*Tyto alba)*, cactus wren (*Campylorhynchus brunneicapillus)*, spotted owl (*Strix*

occidentalis), and Ridgway's rail (*Rallus obsoletus).* These species have distinctive ecological

and life history traits which vary such that we might not expect to find consistent effects of

human presence for these data (Fig. S3). For example, the first four species are human

commensals whose population sizes may be expected to increase in proximity to humans.

Indeed, exploration of the species-specific effects underlying our mixed model suggests that

genetic diversity increases with urbanization for each of these species except barn owls (Fig. S3).

In contrast, the Ridgway's rail and spotted owl are specialized to California salt marshes and old

growth forests, respectively, and thus may respond negatively to human presence. However, our

results could also be attributable to birds' vagility. Cities and their surrounding areas are

characterized by disjoint patches of habitat interspersed among paved surfaces, buildings, and

grassy or agricultural areas (35). Birds' ability to fly may buffer against the effects of habitat

fragmentation and allow for gene flow from undisturbed populations (36) in situations where

mammal movements would be more restricted. Indeed, a global analysis of 57 mammal species

found that the movements of individuals living in areas with a high Human Footprint Index were

considerably reduced relative to those in less disturbed areas (37), which suggests that

fragmentation could underlie the patterns we detect in mammals.

In addition to an overall pattern of reduced gene flow and stronger drift in mammals, we note

that individual species varied in the strength, and sometimes direction, of their responses to

urbanization in our analyses (Fig. S2). This variation could be due to differences in ecological or

life history traits which render species amenable to or susceptible to urbanization (11).

Surprisingly, we found even synanthropic mammals (e.g. red foxes, *Vulpes vulpes*, and mule

deer, *Odocoileus hemionus*) were negatively affected by urbanization at a genetic level (Fig. S2).

The only other studies that have synthetically reanalyzed raw molecular genetic data from online

repositories to test for effects of human land-use on genetic diversity have used mitochondrial

DNA (mtDNA). Miraldo et al. (38) and Millette et al. (39) reanalyzed raw mtDNA sequence data

at a global scale across multiple taxa to assess spatial patterns of variation in sequence diversity,

and to explore their relationships with measures of human disturbance. These studies arrived at

somewhat contradictory results. Miraldo et al. (38) looked at sequence variation in two

mitochondrial genes in mammals, *cytochrome b* and *cytochrome oxidase subunit I*, and found

that while genetic variation in *cytochrome oxidase subunit I* increased in less disturbed

environments, *cytochrome b* variation was not obviously related to human disturbance. Millette

et al.'s more recent work (39) spanned more taxa, including both birds and mammals, and

examined variation across spatial scales as well as temporal variation in genetic diversity as

measured at *cytochrome oxidase subunit I*. Interestingly, they detected no overarching trend for a

loss of genetic diversity associated with proximity to humans (measured by human population

density), and no systematic decline in diversity through time. They found considerable spatial,

temporal, and taxonomic variation in diversity trends.

How can we explain our results in light of this previous work? Our first thought is that

differences could be due to the general lack of relationship between mtDNA diversity and

population size (40). Habitat fragmentation and reduced population sizes are hypothesized to be

the leading mechanisms causing reduced genetic diversity – if these processes are not captured

well by mtDNA markers, trends may be difficult to detect. Additionally, the studies by Miraldo

et al. (38) and Millette et al. (39) were global in scope. This is certainly a strength of their work,

but underlying spatial differences may be harder to detect and control for at this scale. By

focusing on North America, we attempted to control for variation in the timing and nature of

disturbances that would otherwise be difficult at a global scale. Taken together, it is clear that

there are interesting spatial trends in genetic variation, and the exploration of their underlying

causes warrants further study.

While to our knowledge there have not been other synthetic analyses of raw nuclear genetic data

similar to ours, there have been syntheses based on published measures of genetic variation

(11,41). DiBattista (41) found that allelic richness and heterozygosity (an amalgam of expected

and observed) tended to be lower for mammals described as coming from disturbed populations

relative to those from undisturbed populations and, similar to us, found no differences for bird

species. Miles et al. (11) also found that allelic richness tended to be reduced in urban

populations across a diverse array of taxa. Expected heterozygosity and F_{ST} tended to be lower in

urban areas, but this trend was not universal and not significant. They also noted clear instances

where urbanization appeared to facilitate gene flow and diversity. We detected some positive

effects of urbanization (Figs. S2, S3), but our results more strongly suggest a general negative

effect of urbanization on the genetic composition of mammal populations. We suspect that

differences in effect sizes and consistency between our work and these previous studies are

partly due to our access to raw georeferenced genetic data sets. This meant that we could take a

population-level approach, calculating the same genetic measures for all sites, and comparing

them to consistently measured indices of disturbance across the entire data set. For example, we

could avoid procedures that might add noise to relationships such as combining two measures of

heterozygosity and, by calculating a site-specific F_{ST} metric, not have to take means of pairwise

measures. An advantage of using literature-based data is that the sample of species was higher

than ours for each of the literature-based syntheses. Regardless of approach, these syntheses and

ours paint a consistent picture of the effects of urbanization on genetic diversity.

Urbanization and the broader human footprint are leading causes of the current high rates of

species and population-level biodiversity losses (42,43). The population genetic patterns we

detected reflect patterns in genome-wide nuclear genetic diversity that are ultimately the result of

disturbances related to human presence at the ecosystem, community, and population levels. The

consistent effects for mammals across our three measures of human disturbance suggest that this

pattern is not confined to just urban spaces – human land use is an issue for the genetic diversity

of species in general. This has considerable importance for understanding the current nature and

future consequences of biodiversity loss. While monitoring individuals within populations is the

best tool for detecting population trends and assessing risk, such direct monitoring of many

species is not logistically possible. Our results suggest that calls for genetic monitoring programs

are warranted and feasible (44). A relatively small number of genetic markers reflects genome-

wide diversity well (12) and is capable of detecting the effects of human presence. In fact,

publicly archiving data with publications, originally intended to ensure data posterity, safe-

keeping, and research reproducibility, could be better utilized for this task. Access to raw

molecular data sets will continue to increase for the foreseeable future, and can be used for

monitoring regardless of the original purpose. However, for this to be a useful component of

genetic monitoring, more researchers will need to adhere to the standards and best practices for

data sharing to maximize reusability (45,46). This includes using standardized file and metadata

formats that are clearly communicated in data package metadata, and including all relevant

methodological information. In the data searches presented here, a majority ($192/313 = 61\%$) of

datasets were excluded because they did not meet our study criteria (Data S1). However, an

additional 36 datasets were excluded for reasons associated with difficulty accessing or

interpreting data (Data S1): for example, not being able to download files (e.g., only metadata

was available, or only select datasets were deposited), or unclear methodological detail (e.g., no

species designations, delineation between study groups was unclear, or lack of spatial reference).

We were able to resolve such issues in many cases by contacting the authors, however this might

not always be practical for larger studies and limits the ability to automate the data collection

process.

Relative to populations in more natural environments, mammal populations in proximity to

humans have a reduced capacity to spread beneficial alleles in response to selection pressures,

have reduced genetic diversity which can reduce mean population fitness (47,48), and are more

genetically isolated from natural populations. We are extensively and irreversibly creating

environmental change while simultaneously reducing the capacity of some populations to evolve

in response. Reducing fragmentation and facilitating population connectivity are therefore key to

preserving genetic diversity in mammals. Current estimates suggest that by 2050, just 10% of the

planet's surface will be unaltered by humans (6). Land transformation processes are eroding

genetic diversity in mammals, compounding direct effects of habitat loss in a way that threatens

the long-term existence of populations that persist.

Acknowledgements

CS, RPK, and CJG were supported by a Natural Sciences and Engineering Research Council of

Canada Discovery Grant to CJG. CS and RPK were additionally supported by U. Manitoba

Graduate Fellowships, and a U. Manitoba Graduate Enhancement of Tri-council funding grant to

CJG. We would like to thank Grace Ji for creating the program we used to format raw data. We

also thank Kyle Lefort, Paul O'Brien, Deborah Leigh, the students of the Autumn 2018 section

of Evolutionary Biology (BIOL3300) at U. Manitoba, and two anonymous reviewers for their

comments on earlier drafts of the manuscript.

References

1.   Palumbi, S. R. Humans as the World's Greatest Evolutionary Force. *Science.* **293,** 1786–

1790 (2001).

2.   Johnson, M. T. J. & Munshi-South, J. Evolution of life in urban environments. *Science.*

358, 1–11 (2017).

3.   Otto, S. P. Adaptation, speciation and extinction in the Anthropocene. *Proc. R. Soc. B*

285, 20182047 (2018).

4.   Liu, X. *et al.* High-resolution multi-temporal mapping of global urban land using Landsat

images based on the Google Earth Engine Platform. *Remote Sens. Environ.* **209,** 227–239

(2018).

5.   Barnosky, A. D. *et al.* Approaching a state shift in Earth's biosphere. *Nature* **486,** 52–58

(2012).

6.   WWF. *Living Planet Report - 2018: Aiming Higher.* (WWF, 2018). at

<http://scholar.google.com/scholar?hl=en&btnG=Search&q=intitle:LIVING+PLANET+R

EPORT+2004#0>

7.   Leigh, D. M., Hendry, A. P., Vázquez-Domínguez, E. & Friesen, V. L. Estimated six per

cent loss of genetic variation in wild populations since the industrial revolution. *Evol.*

Appl. (2019). doi:10.1111/eva.12810

8.   Wright, S. Evolution in mendelian populations. *Genetics* **16,** (1930).

9.   Fisher, R. A. XXI.—On the Dominance Ratio. *Proc. R. Soc. Edinburgh* **42,** 321–341

(1923).

10.   Sanderson, E. W. *et al.* The Human Footprint and the Last of the Wild. *Bioscience* **52,** 891

(2002).

11.   Miles, L. S., Rivkin, L. R., Johnson, M. T. J., Munshi☐South, J. & Verrelli, B. C. Gene

flow and genetic drift in urban environments. *Mol. Ecol.* **28,** 4138–4151 (2019).

12.   Mittell, E. A., Nakagawa, S. & Hadfield, J. D. Are molecular markers useful predictors of

adaptive potential? *Ecol. Lett.* **18,** 772–778 (2015).

13.   Waples, R. S. A bias correction for estimates of effective population size based on linkage

disequilibrium at unlinked gene loci. *Conserv. Genet.* **7,** 167–184 (2006).

14.   Hill, W. G. Estimation of effective population size from data on linkage disequilibrium.

Genet. Res. **38,** 209–216 (1981).

15.   Do, C. *et al.* NeEstimator v2: Re-implementation of software for the estimation of

contemporary effective population size (Ne) from genetic data. *Mol. Ecol. Resour.* **14,**

209–214 (2014).

16.   Gilbert, K. J. & Whitlock, M. C. Evaluating methods for estimating local effective

population size with and without migration. *Evolution.* **69,** 2154–2166 (2015).

17.   Nei, M. Analysis of gene diversity in subdivided populations. *Proc. Natl. Acad. Sci. U. S.*

A. **70,** 3321–3323 (1973).

18.   Charlesworth, B. & Charlesworth, D. *Elements of Evolutionary Genetics*. (Roberts &

Company, 2010).

19.   Leberg, P. L. Estimating allelic richness: Effects of sample size and bottlenecks. *Mol.*

Ecol. **11,** 2445–2449 (2002).

20.   Weir, B. S. & Goudet, J. A Unified Characterization of Population Structure. *Genetics*

206, 2085–2103 (2017).

21.   La Sorte, F. A., McKinney, M. L. & Pyšek, P. Compositional similarity among urban

floras within and across continents: Biogeographical consequences of human-mediated

biotic interchange. *Glob. Chang. Biol.* **13,** 913–921 (2007).

22.   Hewitt, G. M. The genetic legacy of the Quaternary ice ages. *Nature* **405,** 907–913 (2000).

23.   U.S. Census Bureau. TIGER/Line Shapefiles: Urban Areas (machinereadable data files).

(2016). at <https://www.census.gov/geo/maps-data/data/tiger-line.html>

24.   Statistics Canada. Population Centres Census Division Cartographic Boundary File. 2016

Census. Statistics Canada Catalogue no. 92-160-X. (2016). at

<https://www12.statcan.gc.ca/census-recensement/2011/geo/bound-limit/bound-limit-

2016-eng.cfm>

25.   Shafer, A. B. A., Cullingham, C. I., Côté, S. D. & Coltman, D. W. Of glaciers and refugia:

A decade of study sheds new light on the phylogeography of northwestern North America.

Mol. Ecol. **19,** 4589–4621 (2010).

26.   Stearns, S. C. *The Evolution of Life Histories*. (Oxford University Press, 1992).

27.   Manel, S., Poncet, B. N., Legendre, P., Gugerli, F. & Holderegger, R. Common factors

drive adaptive genetic variation at different spatial scales in *Arabis alpina*. *Mol. Ecol.* **19,**

3824–3835 (2010).

28.   Borcard, D., Legendre, P., Avois-Jacquet, C. & Tuomisto, H. Dissecting the Spatial

Structure of Ecological Data at Multiple Scales. *Ecology* **85,** 1826–1832 (2004).

29.   Dray, S., Legendre, P. & Peres-Neto, P. R. Spatial modelling: a comprehensive framework

for principal coordinate analysis of neighbour matrices (PCNM). *Ecol. Modell.* **196,** 483–

493 (2006).

30.   Dray, S. *et al.* Community ecology in the age of multivariate multiscale spatial analysis.

Ecol. Monogr. **82,** 257–275 (2012).

31.   Romiguier, J. *et al.* Comparative population genomics in animals uncovers the

determinants of genetic diversity. *Nature* **515,** 261–263 (2014).

32.   Bürkner, P. C. brms: An R package for Bayesian multilevel models using Stan. *J. Stat.*

Softw. **80,** (2017).

33.   Bürkner, P.-C. brms: Bayesian Regression Models using 'Stan'. (2019). at <https://cran.r-

project.org/package=brms>

34.   Nakagawa, S. & Schielzeth, H. A general and simple method for obtaining R^2 from

generalized linear mixed-effects models. *Methods Ecol. Evol.* **4,** 133–142 (2013).

35.   Marzluff, J. M. & Ewing, K. Restoration of fragmented landscapes for the conservation of

birds: A general framework and specific recommendations for urbanizing landscapes.

Restor. Ecol. **9,** 280–292 (2001).

36.   Buchmann, C. M., Schurr, F. M., Nathan, R. & Jeltsch, F. Habitat loss and fragmentation

affecting mammal and bird communities: The role of interspecific competition and

individual space use. *Ecol. Inform.* **14,** 90–98 (2013).

37.   Tucker, M. A. *et al.* Moving in the Anthropocene: Global reductions in terrestrial

mammalian movements. *Science.* **359,** 466–469 (2018).

38.   Miraldo, A. *et al.* An Anthropocene map of genetic diversity. *Science.* **353,** 1532–1535

(2016).

39.   Millette, K. L. *et al.* No consistent effects of humans on animal genetic diversity

worldwide. *Ecol. Lett.* ele.13394 (2019). doi:10.1111/ele.13394

40.   Bazin, E., Glémin, S. & Galtier, N. Population size does not influence mitochondrial

genetic diversity in animals. *Science.* **312,** 570–572 (2006).

41.   DiBattista, J. D. Patterns of genetic variation in anthropogenically impacted populations.

Conserv. Genet. **9,** 141–156 (2008).

42.   Aronson, M. F. J. *et al.* A global analysis of the impacts of urbanization on bird and plant

diversity reveals key anthropogenic drivers. *Proc. R. Soc. B Biol. Sci.* **281,** (2014).

43.   McKinney, M. L. Urbanization as a major cause of biotic homogenization. *Biol. Conserv.*

127, 247–260 (2006).

44.   Mimura, M. *et al.* Understanding and monitoring the consequences of human impacts on

intraspecific variation. *Evol. Appl.* **10,** 121–139 (2017).

45.   Whitlock, M. C. Data archiving in ecology and evolution: Best practices. *Trends Ecol.*

Evol. **26,** 61–65 (2011).

46.   Michener, W. K. Ecological data sharing. *Ecol. Inform.* **29,** 33–44 (2015).

47.   Reed, D. H. & Frankham, R. Correlation between Fitness and Genetic Diversity. *Conserv.*

Biol. **17,** 230–237 (2003).

48.   Szulkin, M., Bierne, N. & David, P. Heterozygosity-fitness correlations: A time for

reappraisal. *Evolution.* **64,** 1202–1217 (2010).

Figure and table legends

Figure 1. Urbanization is expected to cause smaller effective population sizes, lower genetic diversity, and increased population differentiation in comparison to natural habitats (a). As habitats become increasingly urbanized, they experience greater fragmentation (b), resulting in smaller patch sizes with lower connectivity. Smaller patches limit supportable population sizes wherein genetic drift becomes the predominant evolutionary force and movement between patches in urbanized areas (black circles) becomes difficult, reducing gene flow.

Figure 2. Map of 1,008 sample sites for the 66 mammal and bird species native to North America examined in this study. 812 sites were mammals (white points) and 129 birds (orange points). Using microsatellite markers, we calculated effective population size, gene diversity, allelic richness, and population-specific F_{ST} for each site. Sites are overlaid on a map of the Human Footprint Index (HFI) where values range from 0 (wild habitat) to 100 (disturbed habitat). Note HFI resolution was reduced for the purposes of visualization.

Figure 3. GLMM coefficients for fixed urban and human disturbance effects in mammals (top), and non-migratory birds (bottom; for bird results including migratory species, refer to Table 1 and Fig. S1). Open circles represent coefficient estimates, bold lines are 90% credible intervals, and narrow lines are 95% credible intervals. Sample size differed between variables, i.e. sites where effective population size was not calculable were excluded, and calculation of population-specific F_{ST} for all sites within a study required at least two sample sites. Sample sizes are given in Table 1.

469   **Figure 1**

472   **Figure 2**

Figure 3

Table 1. Model summaries for mammals, non-migratory birds, and all birds. Four models were constructed per response variable, each including one of three indices of human disturbance: urban-rural category, human population density (popden), and Human Footprint Index (HFI). The fourth model did not include any measure of disturbance and had only dbMEMs as fixed effects (spatial model), or, where no dbMEMs were selected, a null model. Coefficient of variation, R^2, values are an indicator of model fit; marginal R^2 describes the proportion of variation explained by fixed effects, while conditional R^2 is the variation explained by both fixed and random effects.

Class	variable	sites	Fixed effects		coefficient	95% CI		Marginal R^2	Conditional R^2
			dbMEMS	covariate		lower	upper		
Mammals	effective population size	639	5						
				urban-rural	-0.51	-0.87	-0.16	0.04	0.24
				popden	-1.10	-2.44	-0.28	0.28	0.25
				HFI	-0.27	-0.47	-0.05	0.05	0.25
				none	--	--	--	0.01	0.22
	gene diversity	812	13						
				urban-rural	-0.18	-0.33	-0.04	0.04	0.69
				popden	-0.35	-0.68	-0.12	0.12	0.70
				HFI	-0.12	-0.21	-0.02	0.04	0.70
				none	--	--	--	0.03	0.69
	allelic richness	812	21						
				urban-rural	-0.11	-0.29	0.05	0.07	0.69
				popden	-0.15	-0.33	0.00	0.07	0.70
				HFI	-0.11	-0.18	-0.03	0.07	0.70
				none	--	--	--	0.06	0.69
	F_{ST}	795	10						
				urban-rural	0.25	0.02	0.50	0.10	0.33
				popden	0.47	0.11	0.96	0.22	0.35
				HFI	0.15	0.04	0.28	0.11	0.34
				none	--	--	--	0.09	0.32

Birds (non-migratory)	effective population size	87	0						
				urban-rural	-0.20	-1.27	0.80	0.01	0.20
				popden	-0.21	-1.05	0.49	0.03	0.20
				HFI	-0.37	-0.87	-0.15	0.07	0.22
				none	--	--	--	0.00	0.19
	gene diversity	129	3						
				urban-rural	0.01	-0.38	0.41	0.03	0.80
				popden	0.13	-0.09	0.42	0.03	0.80
				HFI	0.04	-0.18	0.25	0.03	0.80
				none	--	--	--	0.02	0.80
	allelic richness	129	0						
				urban-rural	0.10	-0.77	0.82	0.02	0.24
				popden	-0.09	-0.42	0.23	0.01	0.17
				HFI	0.05	-0.37	0.41	0.02	0.22
				none	--	--	--	0.00	0.16
	F_{ST}	128	2						
				urban-rural	-0.13	-0.70	0.41	0.06	0.13
				popden	-0.05	-0.45	0.31	0.06	0.13
				HFI	-0.06	-0.38	0.24	0.06	0.13
				none	--	--	--	0.04	0.10
Birds (all)	effective population size	125	1						
				urban-rural	-0.18	-0.85	0.50	0.08	0.17
				popden	-0.20	-0.74	0.31	0.10	0.17
				HFI	-0.12	-0.48	0.30	0.08	0.18
				none	--	--	--	0.06	0.15
	gene diversity	196	2						
				urban-rural	-0.03	-0.20	0.15	0.00	0.90
				popden	0.04	-0.07	0.15	0.00	0.90

			HFI	-0.03	-0.05	0.07	0.00	0.90
			none	--	--	--	0.00	0.90
allelic richness	196	1						
			urban-rural	0.20	-0.27	0.62	0.03	0.35
			popden	-0.03	-0.24	0.20	0.02	0.29
			HFI	0.11	-0.12	0.32	0.03	0.33
			none	--	--	--	0.01	0.30
F_{ST}	190	1						
			urban-rural	0.00	-0.35	0.34	0.02	0.05
			popden	0.04	-0.22	0.34	0.02	0.05
			HFI	0.08	-0.12	0.31	0.03	0.07
			none	--	--	--	0.01	0.04

Supplementary Information

Methods

<u>Microsatellite data compilation</u>

Our dataset was comprised of bird and mammal microsatellite data collected from publicly

archived, previously published work (Table S2). To create this dataset, we conducted two

systematic searches of online databases (Figure S3). We obtained a list of species names for 859

birds and 450 mammals native to Canada and the United States from the IUCN Red List

database which includes all species regardless of Red List status. We then queried the Dryad

Digital Repository in February 2018 using a python script with the following search terms:

species name (e.g. "*Branta canadensis*"), "microsat*", "single tandem*", "short tandem*", and

"str". This search yielded 194 unique data packages associated with papers. A second search was

performed in May 2018, this time querying DataOne.org, a network which provides access to

data from multiple repositories such as Dryad, the Knowledge Network for Biocomplexity

(KNB), and the United States Geographic Survey (USGS). This search was conducted in R using

the dataone package (1), a convenient method of querying the DataOne network. Using identical

keywords, 237 unique results were generated, 121 of which overlapped with our first search

(Figure S3, Data S1).

All data sets were then individually screened for suitability, ensuring: location (Canada and the

United States), taxon (native birds and terrestrial mammals), data type (neutral microsatellite

markers), and georeferenced sampling (coordinates, maps, or place names). Studies with other

factors which may have influenced genetic diversity (e.g. island sites, genetic rescue,

translocation, managed or captive populations) were excluded. In total, data from 85 studies were

retained for analysis. In a final step, we assured individual sample sites within datasets adhered

to our study criteria, and removed those which did not. We maintained the same sample site

delineations as in the original work. Criteria for removal from a dataset included island,

managed, or captive populations; sites outside of Canada and the United States; and historical

samples (where identified). Sites for which we were unable to extract geographic information

were also removed, as well as sites with <5 individuals. Any non-neutral microsatellite markers

in the data were removed. We note that we compiled all datasets regardless of their location in

urban or rural areas, because we determined the level of human disturbance for individual sites

using our own criteria which could be consistently applied across all datasets (see "Measures of

urbanization and human presence" below). Finally, unique names were assigned to each site,

and all datasets were formatted as either STRUCTURE or GENEPOP files and read into R

version 3.4.2 (2) using the adegenet package (3).

Geographic site locations

Geographic coordinates provided by the authors were used when available (Table S2). Where

spatial location was available for each individual sampled, coordinates were averaged. If site

names were provided (e.g. "Yellowstone National Park") with no coordinate reference, we

performed a Google Maps search and noted the resulting coordinates. Where applicable,

coordinate information was obtained by searching for site names in the Geographic Names

Information System (GNIS) or GeoNames database, as was the case for a few datasets from the

USGS. In instances where only maps of sampling sites were available, site coordinates were

extracted using a reference map in ArcMap version 10.3.1 (ESRI). When georeferencing map

images, if sampling locations indicated regions rather than single points, centroid coordinates

served as the site location. Centroid coordinates were also calculated as site location for data

accompanied by polygon shapefiles as a spatial reference. All coordinates were recorded using the WGS84 (World Geodetic System 1984) coordinate system in decimal degrees, and transformed from other systems or map projections in ArcMap as needed. Finally, when site locations were offshore (42 sites), points were moved to the nearest terrestrial location using the Generate Near Table tool in ArcGIS. Offshore sites (those located in bodies of water) were moved to avoid generating null values for population density and the Human Footprint Index—both of which are high-resolution terrestrial maps which do not extend far past the coast. In some instances, offshore sites were recorded as thus in the original publication, while other times they were generated during the process of obtaining a single location for a site (e.g. the average location of individual coordinates, or centroid location, was in a body of water). Polar bear sites in the Arctic Archipelago constituted half of all offshore sites, while the remainder were coastal species and species sampled near lakes or oceans.

Genetic diversity estimates

We chose to measure gene diversity (4) and allelic richness for each site as measures of genetic diversity. Gene diversity uses allele frequencies to determine the probability that pairs of alleles drawn at random from a population are different, and accounts for both the number and evenness of alleles. This measure is minimally affected by sample size and rare alleles (5), and thus is convenient to use when sample sizes are variable, as is the case here. Gene diversity was calculated using the adegenet package (3). Allelic richness, the number of alleles per locus, is strongly influenced by sample size and effective population size. To account for differences in sample size, we used rarefaction as employed in the R package hierfstat (6) to standardize allele

counts to the minimum sample size ($n = 5$ individuals) across sites (7). Values were then averaged across loci to obtain a single value per site.

Effective population size estimates

We estimated contemporary effective population sizes at each sites using the linkage disequilibrium method for single samples implemented in the software NeEstimator 2.1 (8). The presence of rare alleles produces an upward bias when estimating effective population size which is especially apparent at small sample sizes (9). We therefore set a conservative exclusion threshold (P_{crit}) of 0.1, meaning estimates were made based only on alleles with frequencies higher than this value, which has been shown to markedly reduce bias (9). Linkage disequilibrium methods work well for estimating effective population sizes in small populations, however are less reliable for large populations (10). An estimate of infinity is returned when sampling error swamps detectable signals of genetic drift—which may be the case if too few individuals or loci were sampled to yield any useful information about effective population size. In these instances, rather than replacing infinity values with arbitrary large values, we chose to exclude all sites for which we were unable to estimate effective population size (Table 1).

Population-specific F_{ST}

To estimate levels of population differentiation in relation to human disturbance, we measured population-specific F_{ST} (11). Population-specific F_{ST} characterizes differentiation using the proportion of pairs of matching alleles within populations (the probability of identity by descent) relative to that of pairs from different populations. It can be interpreted as a measure of how far single populations have diverged from a common ancestor population. This measure differs from

pairwise F_{ST} estimates (12) in that it provides a measure of population differentiation for a single population, as opposed to a single value for population pairs. Using a population-specific estimator of structure allows us to make comparisons between populations of different species. Population-specific F_{ST} was calculated in R using hierfstat (6), and values were averaged across loci. Because population-specific F_{ST} calculations still use comparisons of pairs of alleles between populations, it could only be measured for species with two or more sample sites. Sample size was slightly decreased when this condition was not met (Table 1).

Measures of urbanization and human presence

Urban-rural classification. Our next step was to define urban habitats in North America. The United States Census Bureau and Statistics Canada provide publicly available maps of urban areas and population centers, respectively (13,14). According to the US Census Bureau, an urban area is defined as any densely developed territory with at least 2500 inhabitants. Statistics Canada defines a population center as any area with a minimum population of 1000, and a population density of 400 persons or more per square kilometer. We considered these international designations of urbanization to be comparable. Canadian and American urban area maps were downloaded as polygon GIS layers and merged into a single layer. Site coordinates were transformed from WGS84 to the same projection as the urban area maps (GCS North American 1983) in ArcMap to ensure correct alignment. A spatial join was then performed between sites and the urban area layer in order to classify sample locations as "urban" or "nonurban". The search radius parameter was set to 10 km to encompass the entire urban gradient, and account for sprawl. Periurban landscapes which are adjacent to cities may be less densely inhabited, however often encompass areas highly managed or disturbed by humans

including farmland, parks, and golf courses; in larger cities, periurban landscapes may extend up to 10 km away from the city center (15). Thus, any site located in, or within 10 km of, an urban area was considered "urban" for the purposes of this study.

Human population density. Human population density was used as a proxy of urbanization and human effects on the environment. In contrast to our binary urban-rural designation of sample sites, human population reflects the continuous distribution of the effects of human presence, and thus should indicate the intensity of the effects of human activity on genetic diversity. A raster map of global population density per square kilometer was obtained for the most recent available year (2000) from NASA's Center for Near Earth Object Studies (https://neo.sci.gsfc.nasa.gov/view.php?datasetId=SEDAC_POP). Next, the raster map and shapefile containing sites as point features were read into R (package rgdal and raster; Bivand et al. 2017, Hijmans 2017). Mean population density was calculated within a 10 km buffer zone around each site.

Human Footprint Index. The Global Human Footprint Index (18,19) quantifies human influence on a scale of 0 (most wild) to 100 (most transformed) at a 1 km^2 resolution. It provides a more comprehensive assessment of the effects of humans than urban-rural designations or population density alone because it incorporates data from multiple sources of land use. In particular, it captures human population density, human land use and infrastructure (built-up areas, nighttime lights, land use, and land cover), and human access (coastlines, roads, railways, and navigable rivers). As with the raster map of population density, the Human Footprint Index was imported to R and values per site (within a 10 km buffer zone) calculated using the same method.

Statistical analysis

We modelled birds and mammals separately because we expected them to respond to human disturbance in fundamentally different ways. Within birds, we further classified each species as migratory or non-migratory using information from species accounts in The Birds of North America (20). We then created a separate data subset comprised of only non-migratory species which was analyzed in parallel. Species with a mix of migratory and resident populations were counted as migratory and excluded, as were species with unknown migratory behavior.

Genetic diversity is also affected by regional historical contingencies which would be difficult to specifically identify without detailed knowledge of each species and region in our data set (21). Such events will, however, produce spatial patterns in our genetic measures. These spatial patterns are detectable and can be controlled for—even if their causes are unknown—using distance-based Moran's Eigenvector Maps (dbMEMs) (22–24). The dbMEM analysis we used (R package adespatial (25)) is a type of eigenanalysis based on principal coordinates analysis which produces a set of spatially explicit variables, dbMEMs, that quantify spatial trends at multiple scales. Because they are orthogonal, dbMEMs can subsequently be included in regression analyses to explicitly model spatial patterns (25). In the first steps of dbMEM analysis, a modified matrix of distances between pairs of sites is calculated from site coordinates. The eigenvalues of this matrix are proportional to Moran's I coefficients of spatial autocorrelation (23,26). Importantly, only positive eigenvalues are considered because negative eigenvalues generate complex principal coordinate axes (27). dbMEMs therefore correspond to positive values of Moran's I, and can account for positive spatial autocorrelation present in the data. Positive spatial autocorrelation occurs when sites nearer to each other are more similar than sites further away, and violates the assumption of independence in our statistical tests. Before

undertaking dbMEM, any linear trends in the response variables were removed. Although dbMEM analysis is capable of detecting linear spatial gradients, dbMEMs used to model such trends then cannot be used to recover other, potentially more interesting spatial patterns (22). dbMEM analyses were run in parallel for measures of genetic diversity (gene diversity and allelic richness), population-specific F_{ST}, and effective population size. We were able to calculate gene diversity and allelic richness for all sites, however, removed sites where effective population size was infinite and sites where population-specific F_{ST} could not be computed. To capitalize on available data, we created subsets for genetic diversity, population-specific F_{ST}, and effective population size, omitting rows where the focal variable(s) had null values. For each taxon we thus had 3 data subsets: one for gene diversity and allelic richness, which included all sites; population-specific F_{ST}; and effective population size (Table 1). To select dbMEMs for inclusion in regression analyses, we used forward selection with a p-value criterion (alpha = 0.05) in the SignifReg package (28).

Testing effects of human presence on genetic diversity. To test for the effects of human disturbance on genetic diversity, and to determine whether alternate proxies of urbanization would yield similar results, we constructed four linear mixed models per response variable (effective population size, gene diversity, allelic richness, and population-specific F_{ST}). Three of these models included spatial dbMEMs and a measure of human presence as explanatory variables: (1) urban-rural category, (2) human population density, and (3) Human Footprint Index. The fourth model consisted of dbMEMs only, or, where no dbMEMs were significant, was a null model (Table 1). Species was included as a random effect in all models to account for species-level variation in genetic diversity, effective population size, and population-specific F_{ST}.

The random species effect also accommodated potential variation in the level of species'

responses to human-caused environmental degradation (random slope models). Random effects

account for non-independence of samples within groups and increase the accuracy of parameter

estimation (29). We fit these models in a Bayesian framework using the R package brms (30)

which fits models using Stan. We used default priors (uniform distribution over all real numbers)

for parameter estimates with 4000 iterations after discarding warm-up runs (1000 iterations). In

cases where models did not converge, we first increased the number of iterations or warmup

period (*mammals:* allelic richness ~ population density: 5000 iterations, 5000 warmup; *birds:*

F_{ST} ~ population density, 5000 iterations, 4000 warmup; *non-migratory birds:* allelic richness ~

urban category, 4000 iterations, 2000 warmup; allelic richness ~ population density, 4000

iterations, 4000 warmup; F_{ST} ~ population density, 4000 iterations, 2000 warmup). If

convergence issues persisted we restricted priors to a uniform distribution bounded at -10 and 10

(birds F_{ST} ~ population density and non-migratory, and birds allelic richness ~ population

density). Lastly, we computed marginal and conditional Bayesian R^2 to evaluate and compare

model fits using the performance package (31).

Results

Data Summary

Our final dataset included 1,008 sites consisting of 66 species (41 mammals and 25 birds; Fig. 2,

Table S1). There were 812 mammal sample sites, and 196 bird sample sites (129 non-migratory),

with more rural than urban sites (Table S1). The data included samples from a total of 41,023

individuals. Minimum group size for both classes was set to 5 individuals, and there was a

maximum of 2444 individuals from a single sample site for mammals (median = 26 individuals),

602 individuals (median = 19 individuals) from a single sample site among all birds, and 141

individuals (median = 19 individuals) for the non-migratory bird subset.

The number of loci sampled ranging between 5 – 210 loci with a median of 13. For all

birds (both migratory and non-migratory), the median number of loci sampled was 11 with a

range of 6 – 30; non-migratory birds also had a median number of 11 loci with a range between 6

– 20 loci. Gene diversity (mean ± SD) was slightly higher in mammals (0.72 ± 0.11), compared

to birds (all: 0.63 ± 0.13; non-migratory: 0.67 ± 0.09). Allelic richness (mean ± SD, max) was

similar between mammals (4.79 ± 1.28, 13.70) and birds (all: 4.59 ± 2.19, 22.64; non-migratory:

5.01 ± 2.41, 22.64).

We obtained estimates of effective population size for 639 mammal sites, 125 sites across

all birds, and for 87 non-migratory bird sites (Table 1). Effective population sizes (mean ± SD)

were on average lower in mammals (614.52 ± 8275.46) compared to birds (all: 980.30 ±

9400.30; non-migratory: 1314.92 ± 11267.02). This corresponded to higher average population-

specific F_{ST} among mammals (0.06 ± 0.09) relative to birds (all: 0.04 ± 0.06; non-migratory:

0.04 ± 0.04). F_{ST} was estimated at 796 sites for mammals, 190 for all birds, and 128 for non-

migratory birds.

Effective population size sample sizes

Mammals. We were able to obtain estimates of effective population size for 639 out of 812

mammal sites, with all 41 species represented except for one (moose, *Alces alces*, 2 sites). The

ratio of urban to rural sites for sites with non-infinity estimates was unchanged with respect to

the full mammal subset (0.44). Additionally, the distribution of sites across values of human

population density and the Human Footprint Index did not suggest any bias after removing sites with infinite effective population size estimates (Fig. S4).

Non-migratory Birds. Out of 129 sites for non-migratory birds we had 87 non-infinite values. Again, the ratio of urban to rural sites sampled remained consistent with the overall subset of non-migratory birds (0.74), and we saw no indication of bias with regard to human population density or the Human Footprint Index (Fig. S4).

Spatial autocorrelation

We found spatial patterns underlying the distribution of genetic diversity in both mammals and birds. dbMEMs capture spatial patterns at all scales in the data, starting broadly (dbMEM 1) and progressing towards increasingly finer scales. In general, we noted more spatial patterns, and more patterns at finer scales in mammals for all response variables. In mammals, following stepwise regression, 5 dbMEMs were significantly related to effective population size (dbMEMs 2, 27, 80, 93, 101). Significant patterns were also found for genetic diversity: 13 dbMEMs were significantly related to gene diversity (dbMEMs 2, 4, 5, 11, 22, 30, 31, 32, 47, 49, 102, 143, 193), and 21 to allelic richness (dbMEMs 2, 4, 5, 7, 8, 11, 12, 13, 21, 22, 29, 30, 31, 32, 47, 49, 102, 108, 143, 185, 190). Finally, we found 10 dbMEMs related to site -specific F_{ST} (dbMEMs 2, 10, 14, 27, 48, 70, 125, 127, 170, 197).

Patterns of spatial variation were less apparent in our subset of non-migratory birds. There were no significant dbMEMs for effective population size nor allelic richness. Gene diversity showed the most spatial variation, with 3 significant dbMEMs selected (dbMEMs 3, 6, 18). For site-specific FST, 2 dbMEMs were significant (dbMEMs 3, 6).

Among all birds, only 1 dbMEMs was significant for effective population size (dbMEM 2), 2 were retained for gene diversity (dbMEMs 2, 6), 1 for allelic richness (dbMEM 2) and 1 (dbMEM 6) for site-specific F_{ST}.

All significant dbMEMs were incorporated into later models to account for spatial patterns of genetic diversity measures across North America.

References

1. Jones, M. B. *et al.* dataone: R Interface to the DataONE REST API. (2017). at <https://cran.r-project.org/package=dataone>

2. R Core Team. R: A Language and Environment for Statistical Computing. (2013). at <http://www.r-project.org/>

3. Jombart, T. *et al.* adegenet: Exploratory Analysis of Genetic and Genomic Data. (2017). at <https://cran.r-project.org/package=adegenet>

4. Nei, M. Analysis of gene diversity in subdivided populations. *Proc. Natl. Acad. Sci. U. S. A.* **70,** 3321–3323 (1973).

5. Charlesworth, B. & Charlesworth, D. *Elements of Evolutionary Genetics*. (Roberts & Company, 2010).

6. Goudet, J. & Jombart, T. hierfstat: Estimation and Tests of Hierarchical F-Statistics. (2015). at <https://cran.r-project.org/package=hierfstat>

7. Leberg, P. L. Estimating allelic richness: Effects of sample size and bottlenecks. *Mol. Ecol.* **11,** 2445–2449 (2002).

8. Do, C. *et al.* NeEstimator v2: Re-implementation of software for the estimation of contemporary effective population size (Ne) from genetic data. *Mol. Ecol. Resour.* **14,** 209–214 (2014).

9. Waples, R. S. & Do, C. LDNE: A program for estimating effective population size from data on linkage disequilibrium. *Mol. Ecol. Resour.* **8,** 753–756 (2008).

10. Waples, R. S. & Do, C. Linkage disequilibrium estimates of contemporary N_e using highly variable genetic markers: a largely untapped resource for applied conservation and evolution. *Evol. Appl.* **3,** 244–262 (2010).

11. Weir, B. S. & Goudet, J. A Unified Characterization of Population Structure. *Genetics* **206,** 2085–2103 (2017).

12. Weir, B. S. & Cockerham, C. C. Estimating F-Statistics for the Analysis of Population Structure. *Evolution (N. Y).* **38,** 1358–1370 (1984).

13. U.S. Census Bureau. TIGER/Line Shapefiles: Urban Areas (machinereadable data files). (2016). at <https://www.census.gov/geo/maps-data/data/tiger-line.html>

14. Statistics Canada. Population Centres Census Division Cartographic Boundary File. 2016 Census. Statistics Canada Catalogue no. 92-160-X. (2016). at <https://www12.statcan.gc.ca/census-recensement/2011/geo/bound-limit/bound-limit-2016-eng.cfm>

15. Clergeau, P., Jokimäki, J. & Savard, J. P. L. Are urban bird communities influenced by the bird diversity of adjacent landscapes? *J. Appl. Ecol.* **38,** 1122–1134 (2001).

16. Bivand, R., Keitt, T. & Rowlingson, B. rgdal: Bindings for the Geospatial Data Abstraction Library. (2017). at <https://cran.r-project.org/package=rgdal>

17. Hijmans, R. J. raster: Geographic Data Analysis and Modeling. (2017). at <https://cran.r-project.org/package=raster>

18. Wildlife Conservation Society - WCS & Center for International Earth Science Information Network - CIESIN - Columbia University. Last of the Wild Project, Version 2, 2005 (LWP-2): Global Human Footprint Dataset (Geographic). (2005). doi:10.7927/H4M61H5F

19. Sanderson, E. W. *et al.* The Human Footprint and the Last of the Wild. *Bioscience* **52,** 891 (2002).

20. *The Birds of North America*. (Cornell Laboratory of Ornithology, 2015). at <https://birdsna-org.uml.idm.oclc.org>

21. Shafer, A. B. A., Cullingham, C. I., Côté, S. D. & Coltman, D. W. Of glaciers and refugia: A decade of study sheds new light on the phylogeography of northwestern North America. *Mol. Ecol.* **19,** 4589–4621 (2010).

22. Borcard, D., Legendre, P., Avois-Jacquet, C. & Tuomisto, H. Dissecting the Spatial Structure of Ecological Data at Multiple Scales. *Ecology* **85,** 1826–1832 (2004).

23. Dray, S., Legendre, P. & Peres-Neto, P. R. Spatial modelling: a comprehensive framework for principal coordinate analysis of neighbour matrices (PCNM). *Ecol. Modell.* **196,** 483–493 (2006).

24. Manel, S., Poncet, B. N., Legendre, P., Gugerli, F. & Holderegger, R. Common factors drive adaptive genetic variation at different spatial scales in Arabis alpina. *Mol. Ecol.* **19,** 3824–3835 (2010).

25. Dray, S. *et al.* adespatial: Multivariate Multiscale Spatial Analysis. (2017). at <https://cran.r-project.org/package=adespatial>

26. Moran, P. A. P. The Interpretation of Statistical Maps. *J. R. Stat. Soc. Ser. B* **10,** 243–251 (1948).

27. Borcard, D. & Legendre, P. All-scale spatial analysis of ecological data by means of principal coordinates of neighbour matrices. *Ecol. Modell.* **153,** 51–68 (2002).

28. Kim, J. & Zambom, A. Z. SignifReg: Significant Variable Selection in Linear Regression. (2017). at <https://cran.r-project.org/package=SignifReg>

29. Harrison, X. A. *et al.* A brief introduction to mixed effects modelling and multi-model inference in ecology. *PeerJ* **6,** e4794 (2018).

30. Bürkner, P.-C. brms: Bayesian Regression Models using 'Stan'. (2019). at <https://cran.r-project.org/package=brms>

31. Lüdecke, D., Makowski, D. & Waggoner, P. performance: Assessment of Regression Models Performance. (2019). at <https://cran.r-project.org/package=performance>

Figure S1.

Plotted model coefficients all birds (both migratory and non-migratory species). Open circles represent coefficient estimates, bold lines are 90% credible intervals, and narrow lines are 95% credible intervals. Intervals that overlap zero (dashed vertical line) indicate the disturbance variable has no effect on the response variable. Sample size differed between variables due to limitations estimating effective population size population-specific F_{ST}. Sample sizes for each variable are given in Table 1.

Figure S2.

Species-specific parameter estimates for the effect of human population density (which had the strongest effects for mammals) on mammal gene diversity. Ranges are 90% (thick line) and 95% (thin line) credible intervals.

Figure S3.

Species-specific parameter estimates for the effect of human population density on gene diversity in migratory (blue) and non-migratory birds (black). Ranges are 90% (thick line) and 95% (thin line) credible intervals.

Figure S4.

Histograms showing the number of sites included (non-infinity) or excluded (infinity) from effective population size analyses in mammals (top graphs), non-migratory birds (center graphs), and all birds (bottom graphs). (A, C, E) Distribution of sites across human population density. The x axis represents log-transformed human population density plus a constant (0.1). (B, D, F) Distribution of sites across the Human Footprint Index. Human Footprint is measured on a scale from 0 (most wild) to 100 (most disturbed). Overlapping, similar distributions indicate that excluding sites for which we were unable to estimate effective population size likely did not bias our analyses.

Figure S5. Data pipeline. Schematic shows the number of results from two systematic searches in the DataONE network, and filtering steps taken to arrive at our final dataset. Three microsatellite datasets did not appear in our search results, but were discovered manually and included in our analysis. See Table 2 for a complete list of included work, and Data S1 for raw search results.

Table S1.

Summary of number of species, sites, individuals and loci per taxonomic class (mammals and birds). For dataset-specific information, refer to Table S3.

class	species	number of sites			number of individuals		number of loci	
		total sites	urban sites	rural sites	median	range	median	range
mammals	41	812	250	562	26	5 – 2444	13	5 – 210
birds (all)	25	196	71	125	19	5 – 602	11	6 – 30
birds (non-migratory)	7	129	55	74	19	5 – 141	11	6 – 20

Table S2.

Summary and reference information for data used in this study. **Class**: mammals (M) or birds (B). **Repository**: location of data (USGS = U.S. Geological Survey). **Loci**: number of loci sampled, **populations**: number of populations, **individuals**: total number of individuals summed over all populations. **Coordinates**: method of assigning coordinate locations to each site. C = site coordinates provided in study; AC = coordinates given per sample in study, averaged to obtain site coordinates; GM = site name searched in Google Maps; GNIS = site name searchable in the Geographic Names Information System; M = map provided in study, georeferenced in ArcMap; P = site polygons (shapefile) provided, centroid coordinates taken.

Species	Class	Repository	Loci	Populations	Individuals	Coordinates	Search Date	Reference
Alces alces	M	Dryad	10	2	89	AC	February	(1, 2)
Canis latrans	M	Dryad	10	41	482	GM	February	(3, 4)
Cervus elaphus nannodes	M	Dryad	20	2	54	GM	February	(5, 6)
Lasionycteris noctivagans	M	Dryad	18	1	87	C	February	(7, 8)
Lasiurus cinereus	M	Dryad	19	1	132	C		
Lynx rufus	M	Dryad	10	5	95	GM	February	(9, 10)
Leopardus pardalis	M	Dryad	10	2	70	GM		
Lynx rufus	M	Dryad	9	4	365	AC	February	(11, 12)
Lynx rufus	M	Dryad	15	52	1646	AC	February	(13, 14)
Mephitis mephitis	M	Dryad	9	1	345	AC	February	(15, 16)
Procyon lotor	M	Dryad	10	1	330	AC		
Microdipodops megacephalus	M	Dryad	11	3	184	GM	February	(17, 18)
Microdipodops pallidus	M	Dryad	10	2	105	GM		

Myotis lucifugus	M	Dryad	9	15	735	C	February	(19, 20)
Myotis lucifugus	M	Dryad	11	21	1142	M	February	(21, 22)
Myotis lucifugus	M	Dryad	8	29	1310	C	February	(23, 24)
Myotis septentrionalis	M	Dryad	5	15	896	C		
Odocoileus hemionus	M	Dryad	10	60	1831	C	February	(25, 26)
Oreamnos americanus	M	Dryad	22	1	102	AC	February	(27, 28)
Ovis canadensis	M	Dryad	208	1	276	C	February	(29, 30)
Ovis canadensis	M	Dryad	210	1	216	GM		
Peromyscus leucopus	M	Dryad	10	12	134	C	February	(31, 32)
Peromyscus leucopus	M	Dryad	11	11	367	C	February	(33, 34)
Peromyscus leucopus	M	Dryad	18	13	312	GM	February	(35, 36)
Peromyscus maniculatus	M	Dryad	10	1	31	GM	February	(37, 38)
Puma concolor	M	Dryad	13	1	739	AC	February	(39, 40)
Puma concolor	M	Dryad	18	2	667	AC	February	(41, 42)
Puma concolor	M	Dryad	46	8	354	M	February	(43, 44)
Puma concolor	M	Dryad	10	2	196	M	February	(45, 46)
Rangifer tarandus	M	Dryad	19	18	655	M	February	(47, 48)
Rangifer tarandus	M	Dryad	18	27	606	M	February	(49, 50)
Rangifer tarandus	M	Dryad	21	5	230	M	February	(51, 52)
Rangifer tarandus	M	Dryad	14	27	802	M	February	(53, 54)

Rattus rattus	M	Dryad	9	8	126	GM	February	(55, 56)
Taxidea taxus	M	Dryad	12	1	233	AC	February	(57, 58)
Taxidea taxus	M	Dryad	12	3	917	AC	February	(59, 60)
Taxidea taxus	M	Dryad	20	8	236	M	February	(61, 62)
Ursus americanus	M	Dryad	15	7	269	M	February	(63, 64)
Ursus americanus	M	Dryad	15	4	250	AC	February	(65, 66)
Ursus maritimus	M	Dryad	21	14	2232	AC	February	(67, 68)
Ursus arctos	M	Dryad	8	16	831	M	February	(69, 70)
Ursus maritimus	M	Dryad	8	11	319	M		
Ursus americanus	M	Dryad	8	1	32	M		
Ursus arctos	M	Dryad	20	1	729	GM	February	(71, 72)
Ursus maritimus	M	Dryad	9	2	610	AC	February	(73, 74)
Ursus maritimus	M	Dryad	8	4	402	GM	February	(75, 76)
Vulpes vulpes	M	Dryad	8	5	257	M	February	(77, 78)
Vulpes lagopus	M	Dryad	9	3	78	M		
Vulpes vulpes	M	Dryad	13	11	376	AC	February	(79, 80)
Canis lupus	M	Dryad	12	1	62	C	February	(81, 82)
Canis lycaon	M	Dryad	12	1	62	GM		
Odocoileus_hemionus	M	Dryad	10	2	410	AC	February	(83, 84)
Tamiasciurus hudsonicus	M	Dryad	9	12	199	AC	February	(85, 86)

Tamiasciurus douglasii	M	Dryad	9	14	198	AC		
Lepus americanus	M	Dryad	8	39	853	AC	February	(87, 88)
Martes americana	M	Dryad	12	34	653	C	February	(89, 90)
Agelaius phoeniceus	B	Dryad	10	29	240	C	February	(91, 92)
Aphelocoma californica	B	Dryad	13	7	493	AC	February	(93, 94)
Calidris alpina	B	Dryad	8	30	236	C	February	(95, 96)
Charadrius melodus	B	Dryad	8	3	220	C	February	(97, 98)
Charadrius montanus	B	Dryad	14	7	94	C		
Charadrius nivosus	B	Dryad	15	4	68	C		
Charadrius vociferus	B	Dryad	14	2	49	C		
Laterallus jamaicensis	B	Dryad	15	2	336	M	February	(99, 100)
Selasphorus platycercus	B	Dryad	8	2	58	C	February	(101, 102)
Poecile hudsonicus	B	Dryad	6	2	260	C	February	(103, 104)
Poecile atricapillus	B	Dryad	11	13	913	C	February	(105, 106)
Poecile atricapillus	B	Dryad	14	32	142	C	February	(107, 108)
Setophaga caerulescens	B	Dryad	6	8	401	C	February	(109, 110)
Sialis sialis	B	Dryad	12	1	73	C	February	(111, 112)
Antilocapra americana	M	Dryad	19	4	175	GM	May	(113, 114)
Bison bison	M	Dryad	29	1	188	GM	May	(115, 116)
Odocoileus virginianus	M	Dryad	14	8	2069	C	May	(117, 118)

Otospermophilus beecheyi	M	Dryad	11	64	205	AC	May	(119, 120)
Ovis canadensis nelsoni	M	Dryad	16	3	579	P	May	(121, 122)
Rangifer tarandus	M	Dryad	16	14	480	AC	May	(123, 124)
Sylvilagus transitionalis	M	Dryad	10	5	157	AC	May	(125, 126)
Anser albifrons	B	USGS	8	3	115	C	May	(127)
Campylorhynchus brunneicapillus	B	Dryad	22	4	363	AC	May	(128, 129)
Clangula hyemalis	B	USGS	12	12	109	AC	May	(130)
Falco peregrinus	B	USGS	12	7	112	GNIS	May	(131)
Junco hyemalis	B	Dryad	7	8	602	GM	May	(132, 133)
Strix occidentalis	B	USGS	10	1	423	AC	May	(134)
Rallus obsoletus	B	USGS	9	17	107	C	May	(135)
Vireo atricapilla	B	Dryad	12	7	160	AC	May	(136, 137)
Ursus americanus	M	Dryad	12	1	2444	AC	May	(138, 139)
Lynx canadensis	M	Dryad	15	2	556	M	May	(140, 141)
Lynx rufus	M	Dryad	17	1	106	AC	May	(142, 143)
Odocoileus hemionus	M	Dryad	18	5	135	M	May	(144, 145)
Ursus americanus	M	Dryad	20	4	101	M, AC	May	(146, 147)
Ursus arctos	M	Dryad	20	4	113	M, AC		
Lynx canadensis	M	Dryad	14	2	702	C	February	(148, 149)

Glaucomys volans	M	Dryad	7	2	278	C		
Odocoileus hemionus	M	USGS	15	28	73	AC	May	(150)
Anser albifrons	B	USGS	19	8	10	GM	May	(151)
Branta canadensis	B	USGS	19	1	10	GM		
Branta hutchinsii	B	USGS	18	1	9	GM		
Chen canagica	B	USGS	30	1	27	GM		
Ursus americanus	M	Dryad	15	1	506	AC	May	(152, 153)
Ursus maritimus	M	Dryad	24	3	78	M	May	(154, 155)
Peromyscus maniculatus	M	Dryad	11	10	109	C	May	(156, 157)
Pekania pennanti	M	Dryad	16	28	722	C	NA	(158, 159)
Tyto alba	B	NA	20	4	292	M	NA	(160)
Vireo atricapilla	B	Dryad	9	9	67	M	NA	(161, 162)

References

1. Wilson RE, Farley SD, McDonough TJ, Talbot SL, Barboza PS (2015) A genetic discontinuity in moose (*Alces alces*) in Alaska corresponds with fenced transportation infrastructure. *Conserv Genet* 16(4):791–800.

2. Wilson RE, Farley SD, McDonough TJ, Talbot SL, Barboza PS (2015) Data from: A genetic discontinuity in moose (*Alces alces*) in Alaska corresponds with fenced transportation infrastructure. *Conserv Genet*. doi:doi:10.5061/dryad.bp907.

3. Heppenheimer E, et al. (2017) Demographic history influences spatial patterns of genetic diversityin recently expanded coyote (*Canis latrans*) populations. *Heredity*:1–13.

4. Heppenheimer E, et al. (2017) Data from: Demographic history influences spatial patterns of genetic diversity in recently expanded coyote (*Canis latrans*) populations. *Heredity*. doi:doi:10.5061/dryad.2t965.

5. Sacks BN, Lounsberry ZT, Kalani T, Meredith EP, Langner C (2016) Development and characterization of 15 polymorphic dinucleotide microsatellite markers for tule elk using hiseq3000. *J Hered* 107(7):666–669.

6. Sacks BN, Lounsberry ZT, Kalani T, Meredith EP, Langner C (2016) Data from: Development and characterization of 15 polymorphic di-nucleotide microsatellite markers for tule elk using HiSeq3000. *J Hered*. doi:doi:10.5061/dryad.j2654.

7. Baerwald EF, Barclay RMR (2016) Are migratory behaviours of bats socially transmitted? *R Soc Open Sci* 3(150658). doi:10.1093/jmammal/gyv135.

8. Baerwald EF, Barclay RMR (2016) Data from: Are migratory behaviours of bats socially transmitted? *R Soc Open Sci*. doi:doi:10.5061/dryad.4d3n1.

9. Janecka JE, et al. (2016) Genetic differences in the response to landscape fragmentation by a habitat generalist, the bobcat, and a habitat specialist, the ocelot. *Conserv Genet* 17(5):1093–1108.

10. Janecka JE, et al. (2016) Data from: Genetic differences in the response to landscape fragmentation by a habitat generalist, the bobcat, and a habitat specialist, the ocelot. *Conserv Genet*. doi:doi:10.5061/dryad.5b2k6.

11. Serieys LEK, Lea A, Pollinger JP, Riley SPD, Wayne RK (2015) Disease and freeways drive genetic change in urban bobcat populations. *Evol Appl* 8(1):75–92.

12. Serieys LE, Lea A, Pollinger JP, Riley SPD, Wayne RK (2014) Data from: Disease and freeways drive genetic change in urban bobcat populations. *Evol Appl*. doi:doi:10.5061/dryad.7j8q0.

13. Reding DM, Bronikowski AM, Johnson WE, Clark WR (2012) Pleistocene and ecological effects on continental-scale genetic differentiation in the bobcat (*Lynx rufus*). *Mol Ecol* 21(12):3078–3093.

14. Reding DM, Bronikowski AM, Johnson WE, Clark WR (2012) Data from: Pleistocene and ecological effects on continental-scale genetic differentiation in the bobcat (*Lynx*

rufus). *Mol Ecol*. doi:doi:10.5061/dryad.d3t16pd2.

15. Rioux Paquette S, Talbot B, Garant D, Mainguy J, Pelletier F (2014) Modelling the dispersal of the two main hosts of the raccoon rabies variant in heterogeneous environments with landscape genetics. *Evol Appl* 7(7):734–749.

16. Rioux Paquette S, Talbot B, Garant D, Mainguy J, Pelletier F (2014) Data from: Modelling the dispersal of the two main hosts of the raccoon rabies variant in heterogeneous environments with landscape genetics. *Evol Appl*. doi:doi:10.5061/dryad.jr66k.

17. Andersen JJ, Portnoy DS, Hafner JC, Light JE (2013) Populations at risk: Conservation genetics of kangaroo mice (*Microdipodops*) of the great basin desert. *Ecol Evol* 3(8):2497–2513.

18. Andersen JJ, Portnoy DS, Hafner JC, Light JE (2013) Data from: Populations at risk: conservation genetics of kangaroo mice (*Microdipodops*) of the Great Basin Desert. *Ecol Evol*. doi:doi:10.5061/dryad.r531f.

19. Burns LE, Frasier TR, Broders HG (2014) Genetic connectivity among swarming sites in the wide ranging and recently declining little brown bat (*Myotis lucifugus*). *Ecol Evol* 4(21):4130–4149.

20. Burns LE, Frasier TR, Broders HG (2014) Data from: Genetic connectivity among swarming sites in the wide ranging and recently declining little brown bat (*Myotis lucifugus*). *Ecol Evol*. doi:doi:10.5061/dryad.hs37s.

21. Davy CM, et al. (2017) Prelude to a panzootic: gene flow and immunogenetic variation in northern little brown *Myotis* vulnerable to bat white-nose syndrome. *FACETS* 2:690–714.

22. Davy CM, et al. (2017) Data from: Prelude to a panzootic: gene flow and immunogenetic variation in northern little brown *Myotis* vulnerable to bat white-nose syndrome. *FACETS*. doi:doi:10.5061/dryad.h7n25.

23. Johnson LNL, et al. (2015) Population Genetic Structure Within and among Seasonal Site Types in the Little Brown Bat (*Myotis lucifugus*) and the Northern Long-Eared Bat (M. septentrionalis). *PLoS One* 10(5):1–18.

24. Johnson LNL, et al. (2015) Data from: Population genetic structure within and among seasonal site types in the little brown bat (*Myotis lucifugus*) and the northern long-eared bat (M. septentrionalis). *PLoS One*. doi:doi:10.5061/dryad.47nm0.

25. Latch EK, Reding DM, Heffelfinger JR, Alcalá-Galván CH, Rhodes OE (2014) Range-wide analysis of genetic structure in a widespread, highly mobile species (*Odocoileus hemionus*) reveals the importance of historical biogeography. *Mol Ecol* 23(13):3171–3190.

26. Latch EK, Reding DM, Heffelfinger JR, Alcalá-Galván CH, Rhodes OE (2014) Data from: Range-wide analysis of genetic structure in a widespread, highly mobile species (*Odocoileus hemionus*) reveals the importance of historical biogeography. *Mol Ecol*. doi:doi:10.5061/dryad.ns2jn.

27.   Shafer ABA, et al. (2012) Data from: Habitat selection predicts genetic relatedness in an alpine ungulate. *Ecology*. doi:doi:10.5061/dryad.6b6k8b83.

28.   Shafer ABA, et al. (2012) Habitat selection predicts genetic relatedness in an alpine ungulate. *Ecology* 93(6):1317–1329.

29.   Miller JM, Poissant J, Malenfant RM, Hogg JT, Coltman DW (2015) Temporal dynamics of linkage disequilibrium in two populations of bighorn sheep. *Ecol Evol* 5(16):3401–3412.

30.   Miller JM, Poissant J, Malenfant RM, Hogg JT, Coltman DW (2015) Data from: Temporal dynamics of linkage disequilibrium in two populations of bighorn sheep. *Ecol Evol*. doi:doi:10.5061/dryad.n2cj3.

31.   Taylor ZS, Hoffman SMG (2014) Landscape models for nuclear genetic diversity and genetic structure in white-footed mice (*Peromyscus leucopus*). *Heredity* 112(6):588–595.

32.   Taylor ZS, Hoffman SMG (2014) Data from: Landscape models for nuclear genetic diversity and genetic structure in white-footed mice (*Peromyscus leucopus*). *Heredity*. doi:doi:10.5061/dryad.j272c.

33.   Rogic A, Tessier N, Legendre P, Lapointe F-J, Millien V (2013) Genetic structure of the white-footed mouse in the context of the emergence of Lyme disease in southern Québec. *Ecol Evol* 3(7):2075–2088.

34.   Rogic A, Tessier N, Legendre P, Lapointe F-J, Millien V (2013) Data from: Genetic structure of the white-footed mouse in the context of the emergence of Lyme disease in southern Québec. *Ecol Evol*. doi:doi:10.5061/dryad.43vg5.

35.   Munshi-South J, Kharchenko K (2010) Rapid, pervasive genetic differentiation of urban white-footed mouse (*Peromyscus leucopus*) populations in New York City. *Mol Ecol* 19(19):4242–4254.

36.   Munshi-South J, Kharchenko K (2010) Data from: Rapid, pervasive genetic differentiation of urban white-footed mouse (*Peromyscus leucopus*) populations in New York City. *Mol Ecol*. doi:doi:10.5061/dryad.1893.

37.   Ozer F, Gellerman H, Ashley M V. (2011) Genetic impacts of Anacapa deer mice reintroductions following rat eradication. *Mol Ecol* 20(17):3525–3539.

38.   Ozer F, Gellerman H, MV A (2011) Data from: Genetic impacts of Anacapa deer mice reintroductions following rat eradication. *Mol Ecol*. doi:doi:10.5061/dryad.778c9.

39.   Andreasen AM, Stewart KM, Longland WS, Beckmann JP, Forister ML (2012) Identification of source-sink dynamics in mountain lions of the Great Basin. *Mol Ecol* 21(23):5689–5701.

40.   Andreasen AM, Stewart KM, Longland WS, Beckmann JP, Forister ML (2012) Data from: Identification of source-sink dynamics in mountain lions of the Great Basin. *Mol Ecol*. doi:doi:10.5061/dryad.mj74q.

41.   Warren MJ, Wallin DO, Beausoleil RA, Warheit KI (2016) Forest cover mediates genetic

connectivity of northwestern cougars. *Conserv Genet* 17(5):1011–1024.

42. Warren MJ, Wallin DO, Beausoleil RA, Warheit KI (2016) Data from: Forest cover mediates genetic connectivity of northwestern cougars. *Conserv Genet*. doi:doi:10.5061/dryad.3bn62.

43. Ernest HB, Vickers TW, Morrison SA, Buchalski MR, Boyce WM (2014) Fractured genetic connectivity threatens a Southern California puma (*Puma concolor*) population. *PLoS One* 9(10). doi:10.1371/journal.pone.0107985.

44. Ernest HB, Vickers TW, Morrison SA, Buchalski MR, Boyce WM (2014) Data from: Fractured genetic connectivity threatens a southern California puma (*Puma concolor*) population. *PLoS One*. doi:10.5061/dryad.dp0qj.

45. Holbrook JD, Deyoung RW, Tewes ME, Young JH (2012) Demographic history of an elusive carnivore: Using museums to inform management. *Evol Appl* 5(6):619–628.

46. Holbrook JD, Deyoung RW, Tewes ME, Young JH (2012) Data from: Demographic history of an elusive carnivore: using museums to inform management. *Evol Appl*. doi:doi:10.5061/dryad.k5d9470m.

47. Mager KH, Colson KE, Groves P, Hundertmark KJ (2014) Population structure over a broad spatial scale driven by nonanthropogenic factors in a wide-ranging migratory mammal, Alaskan caribou. *Mol Ecol* 23(24):6045–6057.

48. Mager KH, Colson KE, Groves P, Hundertmark KJ (2014) Data from: Population structure over a broad spatial scale driven by non-anthropogenic factors in a wide-ranging migratory mammal, Alaskan caribou. *Mol Ecol*. doi:doi:10.5061/dryad.3hp5v.

49. Serrouya R, et al. (2012) Population size and major valleys explain microsatellite variation better than taxonomic units for caribou in western Canada. *Mol Ecol* 21(11):2588–2601.

50. Serrouya R, et al. (2012) Data from: Population size and major valleys explain microsatellite variation better than taxonomic units for caribou in western Canada. *Mol Ecol*. doi:doi:10.5061/dryad.250c3s47.

51. Colson KE, Mager KH, Hundertmark KJ (2014) Reindeer introgression and the population genetics of caribou in southwestern Alaska. *J Hered* 105(5):585–596.

52. Colson KE, Mager KH, Hundertmark KJ (2014) Data from: Reindeer introgression and the population genetics of caribou in Southwestern Alaska. *J Hered*. doi:doi:10.5061/dryad.9qh56.

53. Weckworth B V., Musiani M, McDevitt AD, Hebblewhite M, Mariani S (2012) Reconstruction of caribou evolutionary history in Western North America and its implications for conservation. *Mol Ecol* 21(14):3610–3624.

54. Weckworth B V., Musiani M, McDevitt AD, Hebblewhite M, Mariani S (2012) Data from: Reconstruction of caribou evolutionary history in Western North America and its implications for conservation. *Mol Ecol*. doi:doi:10.5061/dryad.gn22271h.

55. Lack JB, et al. (2012) Invasion facilitates hybridization with introgression in the Rattus

rattus species complex. *Mol Ecol* 21(14):3545–3561.

56.   Lack JB, et al. (2012) Data from: Invasion facilitates hybridization with introgression in the Rattus rattus species complex. *Mol Ecol*. doi:doi:10.5061/dryad.sj168561.

57.   Kierepka EM, Latch EK (2016) Fine-scale landscape genetics of the American badger (*Taxidea taxus*): disentangling landscape effects and sampling artifacts in a poorly understood species. *Heredity* 116(1):33–43.

58.   Kierepka EM, Latch EK (2015) Data from: Fine-scale landscape genetics of the American badger (*Taxidea taxus*): disentangling landscape effects and sampling artifacts in a poorly understood species. *Heredity*. doi:doi:10.5061/dryad.mr909.

59.   Kierepka EM, Latch EK (2016) High gene flow in the American badger overrides habitat preferences and limits broadscale genetic structure. *Mol Ecol* 25(24):6055–6076.

60.   Kierepka EM, Latch EK (2016) Data from: High gene flow in the American badger overrides habitat preferences and limits broadscale genetic structure. *Mol Ecol*. doi:doi:10.5061/dryad.3qt42.

61.   Rico Y, et al. (2016) Spatial patterns of immunogenetic and neutral variation underscore the conservation value of small, isolated American badger populations. *Evol Appl* 9(10):1271–1284.

62.   Rico Y, et al. (2016) Data from: Spatial patterns of immunogenetic and neutral variation underscore the conservation value of small, isolated American badger populations. *Evol Appl*. doi:doi:10.5061/dryad.qb87r.

63.   Draper JP, Waits LP, Adams JR, Seals CL, Steury TD (2017) Genetic health and population monitoring of two small black bear (*Ursus americanus*) populations in Alabama, with a regional perspective of genetic diversity and exchange. *PLoS One* 12(11). doi:10.1371/journal.pone.0186701.

64.   Draper JP, Waits LP, Adams JR, Seals CL, Steury TD (2017) Data from: Genetic and population monitoring of two small black bear (*Ursus americanus*) populations in Alabama, within a regional context. *PLoS One*. doi:doi:10.5061/dryad.bj7r3.

65.   Puckett EE, et al. (2014) Influence of drift and admixture on population structure of American black bears (*Ursus americanus*) in the Central Interior Highlands, USA, 50 years after translocation. *Mol Ecol* 23(10):2414–2427.

66.   Puckett EE, et al. (2014) Data from: Influence of drift and admixture on population structure of American black bears (*Ursus americanus*) in the Central Interior Highlands, U.S.A. 50 years after translocation. *Mol Ecol*. doi:doi:10.5061/dryad.q405j.

67.   Peacock E, et al. (2015) Implications of the circumpolar genetic structure of polar bears for their conservation in a rapidly warming Arctic. *PLoS One* 10(1):1–30.

68.   Peacock E, et al. (2015) Data from: Implications of the circumpolar genetic structure of polar bears for their conservation in a rapidly warming Arctic. *PLoS One*. doi:doi:10.5061/dryad.v2j1r.

69. Cronin MA, MacNeil MD (2012) Genetic relationships of extant brown bears (*Ursus arctos*) and polar bears (*Ursus maritimus*). *J Hered* 103(6):873–881.

70. Cronin MA, MacNeil MD (2012) Data from: Genetic relationships of extant brown bears (*Ursus arctos*) and polar bears (*Ursus maritimus*). *J Hered*. doi:doi:10.5061/dryad.q30rt.

71. Kamath PL, et al. (2015) Multiple estimates of effective population size for monitoring a long-lived vertebrate: An application to Yellowstone grizzly bears. *Mol Ecol* 24(22):5507–5521.

72. Kamath PL, et al. (2015) Data from: Multiple estimates of effective population size for monitoring a long-lived vertebrate: an application to Yellowstone grizzly bears. *Mol Ecol*. doi:10.5061/dryad.s6764.

73. Campagna L, et al. (2013) Extensive sampling of polar bears (*Ursus maritimus*) in the Northwest Passage (Canadian Arctic Archipelago) reveals population differentiation across multiple spatial and temporal scales. *Ecol Evol* 3(9):3152–3165.

74. Campagna L, et al. (2013) Data from: Extensive sampling of polar bears (*Ursus maritimus*) in the Northwest Passage (Canadian Arctic Archipelago) reveals population differentiation across multiple spatial and temporal scales. *Ecol Evol*. doi:doi:10.5061/dryad.h1v63.

75. Laidre KL, et al. (2018) Range contraction and increasing isolation of a polar bear subpopulation in an era of sea-ice loss. *Ecol Evol* 8:2062–2075.

76. Laidre KL, et al. (2018) Data from: Range contraction and increasing isolation of a polar bear subpopulation in an era of sea-ice loss. *Ecol Evol*. doi:doi:10.5061/dryad.74v52.

77. Goldsmith EW, et al. (2016) Distribution of Rabies Virus Variants in Alaska. *Mol Ecol* 25(3):675–688.

78. Goldsmith EW, et al. (2015) Data from: Population structure of two rabies hosts relative to the known distribution of rabies virus variants in Alaska. *Mol Ecol*. doi:doi:10.5061/dryad.dc1q8.2.

79. Sacks BN, Brazeal JL, Lewis JC (2016) Landscape genetics of the nonnative red fox of California. *Ecol Evol* 6(14):4775–4791.

80. Sacks BN, Brazeal JL, Lewis JC (2016) Data from: Landscape genetics of the nonnative red fox of California. *Ecol Evol*. doi:doi:10.5061/dryad.hj722.

81. Wheeldon TJ, Rutledge LY, Patterson BR, White BN, Wilson PJ (2013) Y-chromosome evidence supports asymmetric dog introgression into eastern coyotes. *Ecol Evol* 3(9):3005–3020.

82. Wheeldon TJ, Rutledge LY, Patterson BR, White BN, Wilson PJ (2013) Data from: Y-chromosome evidence supports asymmetric dog introgression into eastern coyotes. *Ecol Evol*. doi:doi:10.5061/dryad.72s5k.

83. Latch EK, Kierepka EM, Heffelfinger JR, Rhodes OE (2011) Hybrid swarm between divergent lineages of mule deer (*Odocoileus hemionus*). *Mol Ecol* 20(24):5265–5279.

84. Latch EK, Kierepka EM, Heffelfinger JR, Rhodes OE (2011) Data from: Hybrid swarm between divergent lineages of mule deer (*Odocoileus hemionus*). *Mol Ecol*. doi:doi:10.5061/dryad.hm2047nh.

85. Chavez AS, Saltzberg CJ, Kenagy GJ (2011) Genetic and phenotypic variation across a hybrid zone between ecologically divergent tree squirrels (*Tamiasciurus*). *Mol Ecol* 20(16):3350–3366.

86. Chavez AS, Saltzberg CJ, Kenagy GJ (2011) Data from: Genetic and phenotypic variation across a hybrid zone between ecologically divergent tree squirrels (*Tamiasciurus*). *Mol Ecol*. doi:doi:10.5061/dryad.195qg.

87. Cheng E, Hodges KE, Melo-Ferreira J, Alves PC, Mills LS (2014) Conservation implications of the evolutionary history and genetic diversity hotspots of the snowshoe hare. *Mol Ecol* 23(12):2929–2942.

88. Cheng E, Hodges KE, Melo-Ferreira J, Alves PC, Mills LS (2014) Data from: Conservation implications of the evolutionary history and genetic diversity hotspots of the snowshoe hare. *Mol Ecol*. doi:10.5061/dryad.dh63p.

89. Koen EL, Bowman J, Wilson PJ (2016) Node-based measures of connectivity in genetic networks. *Mol Ecol Resour* 16(1):69–79.

90. Koen EL, Bowman J, Wilson PJ (2015) Data from: Node-based measures of connectivity in genetic networks. *Mol Ecol Resour*. doi:doi:10.5061/dryad.4tg23.

91. Liu IA, et al. (2015) Genetic diversity does not explain variation in extra-pair paternity in multiple populations of a songbird. *J Evol Biol* 28(5):1156–1169.

92. Liu IA, et al. (2015) Data from: Genetic diversity does not explain variation in extra-pair paternity in multiple populations of a songbird. *J Evol Biol*. doi:doi:10.5061/dryad.4nj00.

93. Gowen FC, et al. (2014) Speciation in Western Scrub-Jays, Haldane's rule, and genetic clines in secondary contact. *BMC Evol Biol* 14(1):1–15.

94. Gowen FC, et al. (2014) Data from: Speciation in Western Scrub-Jays, Haldane's rule, and genetic clines in secondary contact. *BMC Evol Biol*. doi:doi:10.5061/dryad.57f48.

95. Miller MP, et al. (2015) Intercontinental genetic structure and gene flow in Dunlin (*Calidris alpina*), a potential vector of avian influenza. *Evol Appl* 8(2):149–171.

96. Miller MP, et al. (2014) Data from: Intercontinental genetic structure and gene flow in Dunlin (*Calidris alpina*), a potential vector of avian influenza. *Evol Appl*. doi:doi:10.5061/dryad.4t806.

97. D'Urban Jackson J, et al. (2017) Polygamy slows down population divergence in shorebirds. *Evolution* 71(5):1313–1326.

98. D'Urban Jackson J, et al. (2017) Data from: Polygamy slows down population divergence in shorebirds. *Evolution*. doi:doi:10.5061/dryad.vn77k.

99. Hall LA, Beissinger SR (2017) Inferring the timing of long-distance dispersal between

Rail metapopulations using genetic and isotopic assignments: *Ecol Appl* 27(1):208–218.

100. Hall LA, Beissinger SR (2016) Data from: Inferring the timing of long-distance dispersal between Rail metapopulations using genetic and isotopic assignments. *Ecol Appl*. doi:10.5061/dryad.27mk2.

101. Malpica A, Ornelas JF (2014) Postglacial northward expansion and genetic differentiation between migratory and sedentary populations of the broad-tailed hummingbird (*Selasphorus platycercus*). *Mol Ecol* 23(2):435–452.

102. Malpica A, Ornelas JF (2014) Data from: Post-glacial northward expansion and genetic differentiation between migratory and sedentary populations of the broad-tailed hummingbird (*Selasphorus platycercus*). *Mol Ecol*. doi:doi:10.5061/dryad.14n6g.

103. Lait LA, Burg TM (2013) When east meets west: Population structure of a high-latitude resident species, the boreal chickadee (*Poecile hudsonicus*). *Heredity* 111(4):321–329.

104. Lait LA, Burg TM (2013) Data from: When east meets west: population structure of a high-latitude resident species, the boreal chickadee (*Poecile hudsonicus*). *Heredity*. doi:doi:10.5061/dryad.82hs7.

105. Adams R V., Burg TM (2015) Influence of ecological and geological features on rangewide patterns of genetic structure in a widespread passerine. *Heredity* 114(2):143–154.

106. Adams R V., Burg TM (2014) Data from: Influence of ecological and geological features on rangewide patterns of genetic structure in a widespread passerine. *Heredity*. doi:doi:10.5061/dryad.k086v.

107. Adams R V., Lazerte SE, Otter KA, Burg TM (2016) Influence of landscape features on the microgeographic genetic structure of a resident songbird. *Heredity* 117(2):63–72.

108. Adams R V., Lazerte SE, Otter KA, Burg TM (2016) Data from: Influence of landscape features on the microgeographic genetic structure of a resident songbird. *Heredity*. doi:doi:10.5061/dryad.gs228.

109. Kaiser SA, et al. (2017) A comparative assessment of SNP and microsatellite markers for assigning parentage in a socially monogamous bird. *Mol Ecol Resour* 17(2):183–193.

110. Kaiser SA, et al. (2016) Data from: A comparative assessment of SNP and microsatellite markers for assigning parentage in a socially monogamous bird. *Mol Ecol Resour*. doi:doi:10.5061/dryad.1vn46.

111. Avery JD, Fonseca DM, Campagne P, Lockwood JL (2013) Cryptic introductions and the interpretation of island biodiversity. *Mol Ecol* 22(8):2313–2324.

112. Avery JD, Fonseca DM, Campagne P, Lockwood JL (2013) Data from: Cryptic introductions and the interpretation of island biodiversity. *Mol Ecol*. doi:doi:10.5061/dryad.1tv93.

113. Dunn SJ, Clancey E, Waits LP, Byers JA (2011) Inbreeding depression in pronghorn (*Antilocapra americana*) fawns. *Mol Ecol* 20(23):4889–4898.

114. Dunn SJ, Clancey E, Waits LP, Byers JA (2011) Data from: Inbreeding depression in pronghorn (*Antilocapra americana*) fawns. *Mol Ecol*. doi:10.5061/dryad.ht46qk45.

115. Cronin MA, et al. (2013) Genetic variation and differentiation of bison (*Bison bison*) subspecies and cattle (*Bos taurus*) breeds and subspecies. *J Hered* 104(4):500–509.

116. Cronin MA, et al. (2013) Data from: Genetic variation and differentiation of extant bison (*Bison bison*) subspecies and cattle (*Bos taurus*) breeds and subspecies. *J Hered*. doi:doi:10.5061/dryad.7k66b.

117. Robinson SJ, Samuel MD, Lopez DL, Shelton P (2012) The walk is never random: Subtle landscape effects shape gene flow in a continuous white-tailed deer population in the Midwestern United States. *Mol Ecol* 21(17):4190–4205.

118. Robinson SJ, Samuel MD, Lopez DL, Shelton P (2012) Data from: The walk is never random: subtle landscape effects shape gene flow in a continuous white-tailed deer population in the Midwestern United States. *Mol Ecol*. doi:doi:10.5061/dryad.p7639.

119. Phuong MA, Lim MCW, Wait DR, Rowe KC, Moritz C (2014) Delimiting species in the genus Otospermophilus (Rodentia: *Sciuridae*), using genetics, ecology, and morphology. *Biol J Linn Soc* 113(4):1136–1151.

120. Phuong MA, Lim MCW, Wait DR, Rowe KC, Moritz C (2014) Data from: Delimiting species in the genus Otospermophilus (Rodentia: *Sciuridae*) using genetics, ecology, and morphology. *Biol J Linn Soc*. doi:doi:10.5061/dryad.s4h2j.

121. Epps CW, Crowhurst RS, Nickerson BS (2018) Assessing changes in functional connectivity in a desert bighorn sheep metapopulation after two generations. *Mol Ecol* 27:2334–2346.

122. Epps CW, Crowhurst RS, Nickerson BS (2018) Data from: Assessing changes in functional connectivity in a desert bighorn sheep metapopulation after two generations. *Mol Ecol*. doi:doi:10.5061/dryad.mp71t50.

123. Yannic G, et al. (2014) Temporally dynamic habitat suitability predicts genetic relatedness among caribou. *Proceedings Biol Sci* 281(2014052). doi:10.1098/rspb.2014.0502.

124. Yannic G, et al. (2014) Data from: Temporally dynamic habitat suitability predicts genetic relatedness among caribou. *Proc R Soc B*. doi:doi:10.5061/dryad.qn1cj.

125. Fenderson LE, et al. (2014) A multiscale analysis of gene flow for the New England cottontail, an imperiled habitat specialist in a fragmented landscape. *Ecol Evol* 4(10):1853–1875.

126. Fenderson LE, et al. (2014) Data from: A multi-scale analysis of gene flow for the New England cottontail, an imperiled habitat specialist in a fragmented landscape. *Ecol Evol*. doi:10.5061/dryad.1s834.

127. Wilson RE (2006) Greater White-Fronted Goose (*Anser albifrons*) Microsatellite DNA Data, Alaska, 1989-2006. doi:10.5066/F71G0JGN.

128. Barr KR, et al. (2015) Habitat fragmentation in coastal southern California disrupts

genetic connectivity in the cactus wren (*Campylorhynchus brunneicapillus*). *Mol Ecol* 24(10):2349–2363.

129. Barr KR, et al. (2015) Data from: Habitat fragmentation in coastal southern California disrupts genetic connectivity in the Cactus Wren (*Campylorhynchus brunneicapillus*). *Mol Ecol*. doi:doi:10.5061/dryad.j5h92.

130. Wilson RE, Talbot SL (2016) Long-tailed Duck (*Clangula hyemalis*) Microsatellite DNA Data; Alaska, Canada, Russia, 1994-2002. doi:10.5066/F7HX19RJ.

131. Pierson BJ (2017) Peregrine Falcon (*Falco peregrinus*) mtDNA and Microsatellite Genetic Data, Alaska, Canada and Russia, 1880-2012. doi:10.5066/F7F18WV0.

132. Liebgold EB, Gerlach NM, Ketterson ED (2013) Similarity in temporal variation in sex-biased dispersal over short and long distances in the dark-eyed junco, *Junco hyemalis*. *Mol Ecol* 22(22):5548–5560.

133. Liebgold EB, Gerlach NM, Ketterson ED (2013) Data from: Similarity in temporal variation in sex-biased dispersal over short and long distances in the dark-eyed junco, *Junco hyemalis*. *Mol Ecol*. doi:doi:10.5061/dryad.1jd40.2.

134. Miller MP, Davis RJ, Mullins TD, Haig SM, Forsman ED (2017) Microsatellite markers, habitat quality, and sample location data for Northern Spotted Owls (*Strix occidentalis caurina*). doi:10.5066/F7J67FVW.

135. Wood DA, et al. (2016) A Century of Landscape Disturbance and Urbanization of the San Francisco Bay Region affects the Present-day Genetic Diversity of the California Ridgway's Rail (*Rallus obsoletus obsoletus*). doi:10.5066/F7HD7SQ0.

136. Athrey G, Lance RF, Leberg PL (2012) How far is too close? restricted, sex-biased dispersal in black-capped vireos. *Mol Ecol* 21(17):4359–4370.

137. Athrey G, Lance RF, Leberg PL (2012) Data from: How far is too close? Restricted, sex-biased dispersal in black-capped vireos. *Mol Ecol*. doi:doi:10.5061/dryad.d2fm8.

138. Moore JA, Xu R, Frank K, Draheim H, Scribner KT (2015) Social network analysis of mating patterns in American black bears (*Ursus americanus*). *Mol Ecol* 24(15):4010–4022.

139. Moore JA, Xu R, Frank K, Draheim H, Scribner KT (2015) Data from: Social network analysis of mating patterns in American black bears (*Ursus americanus*). *Mol Ecol*. doi:doi:10.5061/dryad.c61q0.

140. Prentice MB, et al. (2017) Selection and drift influence genetic differentiation of insular Canada lynx (*Lynx canadensis*) on Newfoundland and Cape Breton Island. *Ecol Evol* 7(9):3281–3294.

141. Prentice MB, et al. (2017) Data from: Selection and drift influence genetic differentiation of insular Canada lynx (*Lynx canadensis*) on Newfoundland and Cape Breton Island. *Ecol Evol*. doi:doi:10.5061/dryad.8ff46.

142. Lee JS, et al. (2012) Gene flow and pathogen transmission among bobcats (*Lynx rufus*) in

a fragmented urban landscape. *Mol Ecol* 21(7):1617–1631.

143. Lee JS, et al. (2012) Data from: Gene flow and pathogen transmission among bobcats (*Lynx rufus*) in a fragmented urban landscape. *Mol Ecol*. doi:doi:10.5061/dryad.mm317th7.

144. Northrup JM, Shafer ABA, Anderson CR, Coltman DW, Wittemyer G (2014) Fine-scale genetic correlates to condition and migration in a wild cervid. *Evol Appl* 7(8):937–948.

145. Northrup JM, Shafer ABA, Anderson CR, Coltman DW, Wittemyer G (2014) Data from: Fine scale genetic correlates to condition and migration in a wild Cervid. *Evol Appl*. doi:doi:10.5061/dryad.3vc1b.

146. Sawaya MA, Kalinowski ST, Clevenger AP (2014) Genetic connectivity for two bear species at wildlife crossing structures in Banff National Park. *Proc R Soc B* 281:1–10.

147. Sawaya MA, Kalinowski ST, Clevenger AP (2014) Data from: Genetic connectivity for two bear species at wildlife crossing structures in Banff National Park. *Proc R Soc B*. doi:doi:10.5061/dryad.5q3b3.

148. Marrotte RR, et al. (2017) Multi-species genetic connectivity in a terrestrial habitat network. *Mov Ecol* 5(1):1–11.

149. Marrotte RR, et al. (2017) Data from: Multi-species genetic connectivity in a terrestrial habitat network. *Mov Ecol*. doi:doi:10.5061/dryad.qn4kq.

150. Vandergast AG, Mitelberg A (2016) DNA fingerprinting of Southern Mule Deer (*Odocoileus hemionus fuliginatus*) to determine movement across California State Route 67. doi:10.5066/F7KW5D32.

151. Talbot SL (2017) DNA Microsatellite and Sex Identification Markers for Emperor Goose (*Chen canagica*) and Cross-Species Amplification of Microsatellites in Select Goose Species, Alaska 2016. doi:10.5066/F71V5C4S.

152. Puckett EE, Etter PD, Johnson EA, Eggert LS (2015) Phylogeographic Analyses of American Black Bears (*Ursus americanus*) Suggest Four Glacial Refugia and Complex Patterns of Postglacial Admixture. *Mol Biol Evol* 32(9):2338–2350.

153. Puckett EE, Etter PD, Johnson EA, Eggert LS (2015) Data from: Phylogeographic analyses of American black bears (*Ursus americanus*) suggest four glacial refugia and complex patterns of post-glacial admixture. *Mol Biol Evol*. doi:doi:10.5061/dryad.dc02b.

154. Malenfant RM, Coltman DW, Davis CS (2015) Design of a 9K illumina BeadChip for polar bears (*Ursus maritimus*) from RAD and transcriptome sequencing. *Mol Ecol Resour* 15(3):587–600.

155. Malenfant RM, Coltman DW, Davis CS (2014) Data from: Design of a 9K SNP chip for polar bears (*Ursus maritimus*) from RAD and transcriptome sequencing. *Mol Ecol Resour*. doi:doi:10.5061/dryad.b35td.

156. Taylor ZS, Hoffman SMG (2012) Microsatellite genetic structure and cytonuclear discordance in naturally fragmented populations of deer mice (*Peromyscus maniculatus*).

J Hered 103(1):71–79.

157. Taylor ZS, Hoffman SMG (2011) Data from: Microsatellite genetic structure and cytonuclear discordance in naturally fragmented populations of deer mice (*Peromyscus maniculatus*). *J Hered*. doi:doi:10.5061/dryad.gs048.

158. Bertrand P, Bowman J, Dyer R, Manseau M, PJ W (2017) Sex-specific graphs: Relating group-specific topology to demographic and landscape data. *Mol Ecol* 26:3898–3912.

159. Bertrand P, Bowman J, Dyer R, Manseau M, PJ W (2017) Data from: Sex-specific graphs: Relating group-specific topology to demographic and landscape data. *Mol Ecol*. doi:doi:10.5061/dryad.167d5.

160. Machado AP, Clément L, Uva V, Goudet J, Roulin A (2018) The Rocky Mountains as a dispersal barrier between barn owl (*Tyto alba*) populations in North America. *J Biogeogr*:1–13.

161. Athrey G, Barr KR, Lance RF, Leberg PL (2012) Birds in space and time: Genetic changes accompanying anthropogenic habitat fragmentation in the endangered black-capped vireo (*Vireo atricapilla*). *Evol Appl* 5(6):540–552.

162. Athrey G, KR B, RF L, PL L (2012) Data from: Birds in space & time: genetic changes accompanying anthropogenic habitat fragmentation in the endangered black-capped vireo (*Vireo atricapilla*). *Evol Appl*. doi:doi:10.5061/dryad.2r030k5p.

Data S1.xlsx

List of all results from systematic searches performed in February and May 2018. Columns in this table are: class (mammal or bird), species name, search date, repository, whether it was included in our study (yes or no), the reason for exclusion if applicable, and a link to the online data. The "overlap" reason for exclusion indicates May results which overlapped with February results, while "duplicate data" indicates a duplicate entry for the same dataset within a search month. See Figure S5 for our data filtering process. Ultimately, 85 data sets were retained for analysis after individual screening to ensure the data met our study criteria, including: location (within the continental United States and Canada), taxon (native birds and terrestrial mammals), data type (neutral microsatellite markers), and georeferenced sampling. Available at: https://doi.org/10.5061/dryad.cz8w9gj0c