Common variants in breast cancer risk loci predispose to distinct tumor subtypes

Thomas U. Ahearn[1*], Haoyu Zhang[1,2*], Kyriaki Michailidou[3,4,5], Roger L. Milne[6,7,8], Manjeet K.

Bolla[3], Joe Dennis[3], Alison M. Dunning[9], Michael Lush[3], Qin Wang[3], Irene L. Andrulis[10], Hoda

Anton-Culver[11], Volker Arndt[12], Kristan J. Aronson[13], Paul L. Auer[14,15], Annelie Augustinsson[16],

Adinda Baten[17], Heiko Becher[18], Sabine Behrens[19], Javier Benitez[20], Marina Bermisheva[21], Carl

Blomqvist[22,23], Stig E. Bojesen[24,25,26,27], Bernardo Bonanni[28], Anne-Lise Børresen-Dale[29,30], Hiltrud

Brauch[31,32,33], Hermann Brenner[12,34,33], Angela Brooks-Wilson[35,36], Thomas Brüning[37], Barbara

Burwinkel[38,39], Federico Canzian[40], Jose E. Castelao[41], Jenny Chang-Claude[19,42], Stephen J.

Chanock[1], Georgia Chenevix-Trench[43], Christine L. Clarke[44], NBCS Collaborators[†,] J.Margriet

Collée[45], Angela Cox[46], Simon S. Cross[47], Kamila Czene[48], Mary B. Daly[49], Peter Devilee[50,51], Thilo

Dörk[52], Miriam Dwek[53], Diana M. Eccles[54], D.Gareth Evans[55,56], Peter A. Fasching[57,58], Jonine

Figueroa[59,60], Giuseppe Floris[17], Manuela Gago-Dominguez[61,62], Susan M. Gapstur[63], José A.

García-Sáenz[64], Mia M. Gaudet[63], Graham G. Giles[6,7,8], Mark S. Goldberg[65,66,67], David E.

Goldgar[68], Anna González-Neira[27], Grethe I. GrenakerAlnæs[29], Mervi Grip[69], Pascal Guénel[70],

Christopher A. Haiman[71], Per Hall[48,72], Ute Torres[73], Elaine F. Harkness[74,75], Bernadette A.M.

Heemskerk-Gerritsen[76], Bernd Holleczek[77], Antoinette Hollestelle[76], Maartje J. Hooning[76],

Robert N. Hoover[1], John L. Hopper[7], Anthony Howell[78], kConFab/AOCS Investigators[†], Milena

Jakimovska[79], Anna Jakubowska[80,81], Esther M. John[82], Michael E. Jones[83], Audrey Jung[19], Rudolf

Kaaks[19], Saila Kauppila[84], Renske Keeman[85], Elza Khusnutdinova[86], Cari M. Kitahara[87], Yon-

Dschun Ko[88], Stella Koutros[1], Vessela N. Kristensen[29,30], Ute Krüger[16], Katerina Kubelka-Sabit[89],

Allison W. Kurian[82], Kyriacos Kyriacou[5,90], Diether Lambrechts[91,92], Derrick G. Lee[93,94], Annika

Lindblom[95,96], Martha Linet[87], Jolanta Lissowska[97], Ana Llaneza[98], Wing-Yee Lo[31,99], Robert J.

MacInnis[6,7], Arto Mannermaa[100,101,102], Mehdi Manoochehri[73], Sara Margolin[72,103], Maria Elena

Martinez[62,104], Catriona McLean[105], Alfons Meindl[106], Usha Menon[107], Heli Nevanlinna[108],

William G. Newman[55,56], Jesse Nodora[109,110], Kenneth Offit[111], Håkan Olsson[16,112], Nick Orr[113],

Tjoung-Won Park-Simon[52], Julian Peto[3,114], Guillermo Pita[115], Dijana Plaseska-Karanfilska[79], Ross

Prentice[14], Kevin Punie[17], Katri Pylkäs[116,117], Paolo Radice[118], Gad Rennert[119], Atocha Romero[120],

Thomas Rüdiger[121], Emmanouil Saloustros[122], Sarah Sampson[123], Dale P. Sandler[124], Elinor J.

Sawyer[125], Rita K. Schmutzler[126], Minouk J. Schoemaker[83], Ben Schöttker[12], Mark E. Sherman[127],

Xiao-Ou Shu[128], Snezhana Smichkoska[129], Melissa C. Southey[8], John J. Spinelli[130,131], Anthony J.

Swerdlow[83,132], Rulla M. Tamimi[133,134,135], William J. Tapper[136], Jack A. Taylor[124,137], MaryBeth

Terry[138], Diana Torres[73,139], Melissa A. Troester[140], Celine M. Vachon[141], Carolien H.M. van

Deurzen[142], Elke M. van Veen[55,56], Philippe Wagner[16], Clarice R. Weinberg[143], Camilla Wendt[103],

Jelle Wesseling[85], Robert Winqvist[51,116,117,144], Alicja Wolk[81,145,146,81], Xiaohong R. Yang[1], Wei

Zheng[128], Fergus J. Couch[147], Jacques Simard[148], Peter Kraft[134,135], Douglas F. Easton[3,9], Paul D.P.

Pharoah[3,9], Marjanka K. Schmidt[85,149], Montserrat García-Closas[1]**, Nilanjan Chatterjee[2,150]**

[1]Division of Cancer Epidemiology and Genetics, National Cancer Institute, National Institutes of

Health, Department of Health and Human Services, Bethesda, MD, USA, [2]Department of

Biostatistics, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD, USA, [3]Centre for

Cancer Genetic Epidemiology, Department of Public Health and Primary Care, University of

Cambridge, Cambridge, UK, [4]Biostatistics Unit, The Cyprus Institute of Neurology and Genetics,

Nicosia, Cyprus, [5]Cyprus School of Molecular Medicine, Nicosia, Cyprus, [6]Cancer Epidemiology

Division, Cancer Council Victoria, Melbourne, Victoria, Australia, [7]Centre for Epidemiology and

Biostatistics, Melbourne School of Population and Global Health, The University of Melbourne,

Melbourne, Victoria, Australia, [8]Precision Medicine, School of Clinical Sciences at Monash

Health, Monash University, Clayton, Victoria, Australia, [9]Centre for Cancer Genetic

Epidemiology, Department of Oncology, University of Cambridge, Cambridge, UK, [10]Fred A.

Litwin Center for Cancer Genetics, Lunenfeld-Tanenbaum Research Institute of Mount Sinai

Hospital, Toronto, ON, Canada, [11]Department of Epidemiology, Genetic Epidemiology Research

Institute, University of California Irvine, Irvine, CA, USA, [12]Division of Clinical Epidemiology and

Aging Research, German Cancer Research Center (DKFZ), Heidelberg, Germany, [13]Department

of Public Health Sciences, and Cancer Research Institute, Queen's University, Kingston, ON,

Canada, [14]Cancer Prevention Program, Fred Hutchinson Cancer Research Center, Seattle, WA,

USA, [15]Zilber School of Public Health, University of Wisconsin-Milwaukee, Milwaukee, WI, USA,

[16]Department of Cancer Epidemiology, Clinical Sciences, Lund University, Lund, Sweden,

[17]Leuven Multidisciplinary Breast Center, Department of Oncology, Leuven Cancer Institute,

University Hospitals Leuven, Leuven, Belgium, [18]Institute of Medical Biometry and

Epidemiology, University of Hamburg, Hamburg, Germany, [19]Division of Cancer Epidemiology,

German Cancer Research Center (DKFZ), Heidelberg, Germany, [20]Centro de Investigación en

Red de Enfermedades Raras (CIBERER), Valencia, Spain, [21]Institute of Biochemistry and

Genetics, Ufa Scientific Center of Russian Academy of Sciences, Ufa, Russia, [22]Department of

Oncology, Helsinki University Hospital, University of Helsinki, Helsinki, Finland, [23]Department of

Oncology, Örebro University Hospital, Örebro, Sweden, [24]Copenhagen General Population

Study, Herlev and Gentofte Hospital, Copenhagen University Hospital, Herlev, Denmark,

[25]Department of Clinical Biochemistry, Herlev and Gentofte Hospital, Copenhagen University

Hospital, Herlev, Denmark, [26]Faculty of Health and Medical Sciences, University of Copenhagen,

Copenhagen, Denmark, [27]Human Cancer Genetics Programme, Spanish National Cancer

Research Centre (CNIO), Madrid, Spain, [28]Division of Cancer Prevention and Genetics, IEO,

European Institute of Oncology IRCCS, Milan, Italy, [29]Department of Cancer Genetics, Institute

for Cancer Research, Oslo University Hospital-Radiumhospitalet, Oslo, Norway, [30]Institute of

Clinical Medicine, Faculty of Medicine, University of Oslo, Oslo, Norway, [31]Dr. Margarete

Fischer-Bosch-Institute of Clinical Pharmacology, Stuttgart, Germany, [32]iFIT-Cluster of

Excellence, University of Tübingen, Tübingen, Germany, [33]German Cancer Consortium (DKTK),

German Cancer Research Center (DKFZ), Heidelberg, Germany, [34]Division of Preventive

Oncology, German Cancer Research Center (DKFZ) and National Center for Tumor Diseases

(NCT), Heidelberg, Germany, [35]Genome Sciences Centre, BC Cancer Agency, Vancouver, BC,

Canada, [36]Department of Biomedical Physiology and Kinesiology, Simon Fraser University,

Burnaby, BC, Canada, [37]Institute for Prevention and Occupational Medicine of the German

Social Accident Insurance, Institute of the Ruhr University Bochum (IPA), Bochum, Germany,

[38]Molecular Epidemiology Group, C080, German Cancer Research Center (DKFZ), Heidelberg,

Germany, [39]Molecular Biology of Breast Cancer, University Womens Clinic Heidelberg,

University of Heidelberg, Heidelberg, Germany, [40]Genomic Epidemiology Group, German

Cancer Research Center (DKFZ), Heidelberg, Germany, [41]Oncology and Genetics Unit, Instituto

de Investigacion Sanitaria Galicia Sur (IISGS), Xerencia de Xestion Integrada de Vigo-SERGAS,

Vigo, Spain, [42]Cancer Epidemiology Group, University Cancer Center Hamburg (UCCH),

University Medical Center Hamburg-Eppendorf, Hamburg, Germany, [43]Department of Genetics

and Computational Biology, QIMR Berghofer Medical Research Institute, Brisbane, Queensland,

Australia, [44]Westmead Institute for Medical Research, University of Sydney, Sydney, New South

Wales, Australia, [45]Department of Clinical Genetics, Erasmus University Medical Center,

Rotterdam, The Netherlands, [46]Sheffield Institute for Nucleic Acids (SInFoNiA), and Weston Park

Cancer Centre, Department of Oncology and Metabolism, University of Sheffield, Sheffield, UK,

[47]Department of Neuroscience, University of Sheffield, Sheffield, UK, [48]Department of Medical

Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden, [49]Department of

Clinical Genetics, Fox Chase Cancer Center, Philadelphia, PA, USA, [50]Department of Pathology,

Leiden University Medical Center, Leiden, The Netherlands, [51]Department of Human Genetics,

Leiden University Medical Center, Leiden, The Netherlands, [52]Gynaecology Research Unit,

Hannover Medical School, Hannover, Germany, [53]Department of Biomedical Sciences, Faculty

of Science and Technology, University of Westminster, London, UK, [54]Cancer Sciences Academic

Unit, Faculty of Medicine, University of Southampton, Southampton, UK, [55]Division of Evolution

and Genomic Medicine, School of Biological Sciences, Faculty of Biology, Medicine and Health,

University of Manchester, Manchester Academic Health Science Centre, Manchester, UK,

[56]Manchester Centre for Genomic Medicine, St Mary's Hospital, Manchester NIHR Biomedical

Research Centre, Manchester University Hospitals NHS, Foundation Trust, Manchester

Academic Health Science Centre, Manchester, UK, [57]David Geffen School of Medicine,

Department of Medicine Division of Hematology and Oncology, University of California at Los

Angeles, Los Angeles, CA, USA, [58]Department of Gynecology and Obstetrics, Comprehensive

Cancer Center ER-EMN, University Hospital Erlangen, Friedrich-Alexander-University Erlangen-

Nuremberg, Erlangen, Germany, [59]Usher Institute of Population Health Sciences and

Informatics, The University of Edinburgh Medical School, Edinburgh, UK, [60]Cancer Research UK

Edinburgh Centre, Edinburgh, UK, [61]Genomic Medicine Group, Galician Foundation of Genomic

Medicine, Instituto de Investigación Sanitaria de Santiago de Compostela (IDIS), Complejo

Hospitalario Universitario de Santiago, SERGAS, Santiago de Compostela, Spain, [62]Moores

Cancer Center, University of California San Diego, La Jolla, CA, USA, [63]Behavioral and

Epidemiology Research Group, American Cancer Society, Atlanta, GA, USA, [64]Medical Oncology

Department, Hospital Clínico San Carlos, Instituto de Investigación Sanitaria San Carlos (IdISSC),

Centro Investigación Biomédica en Red de Cáncer (CIBERONC), Madrid, Spain, [65]Department of

Medicine, McGill University, Montréal, QC, Canada, [66]Division of Clinical Epidemiology, Royal

Victoria Hospital, McGill University, Montréal, QC, Canada, [67]Breast Cancer Research Unit,

Cancer Research Institute, University Malaya Medical Centre, Kuala Lumpur, Malaysia,

[68]Department of Dermatology, Huntsman Cancer Institute, University of Utah School of

Medicine, Salt Lake City, UT, USA, [69]Department of Surgery, Oulu University Hospital, University

of Oulu, Oulu, Finland, [70]Cancer & Environment Group, Center for Research in Epidemiology

and Population Health (CESP), INSERM, University Paris-Sud, University Paris-Saclay, Villejuif,

France, [71]Department of Preventive Medicine, Keck School of Medicine, University of Southern

California, Los Angeles, CA, USA, [72]Department of Oncology, Södersjukhuset, Stockholm,

Sweden, [73]Molecular Genetics of Breast Cancer, German Cancer Research Center (DKFZ),

Heidelberg, Germany, [74]Division of Informatics, Imaging and Data Sciences, Faculty of Biology,

Medicine and Health, University of Manchester, Manchester Academic Health Science Centre,

Manchester, UK, [75]Nightingale Breast Screening Centre, Wythenshawe Hospital, Manchester

University NHS Foundation Trust, Manchester, UK, [76]Department of Medical Oncology, Family

Cancer Clinic, Erasmus MC Cancer Institute, Rotterdam, The Netherlands, [77]Saarland Cancer

Registry, Saarbrücken, Germany, [78]Division of Cancer Sciences, University of Manchester,

Manchester, UK, [79]Research Centre for Genetic Engineering and Biotechnology "Georgi D.

Efremov", MASA, Skopje, Republic of North Macedonia, [80]Department of Genetics and

Pathology, Pomeranian Medical University, Szczecin, Poland, [81]Independent Laboratory of

Molecular Biology and Genetic Diagnostics, Pomeranian Medical University, Szczecin, Poland,

[82]Department of Medicine, Division of Oncology, Stanford Cancer Institute, Stanford University

School of Medicine, Stanford, CA, USA, [83]Division of Genetics and Epidemiology, The Institute of

Cancer Research, London, UK, [84]Department of Pathology, Oulu University Hospital, University

of Oulu, Oulu, Finland, [85]Division of Molecular Pathology, The Netherlands Cancer Institute -

Antoni van Leeuwenhoek Hospital, Amsterdam, The Netherlands, [86]Department of Genetics and

Fundamental Medicine, Bashkir State University, Ufa, Russia, [87]Radiation Epidemiology Branch,

Division of Cancer Epidemiology and Genetics, National Cancer Institute, Bethesda, MD, USA,

[88]Department of Internal Medicine, Evangelische Kliniken Bonn gGmbH, Johanniter

Krankenhaus, Bonn, Germany, [89]Department of Histopathology and Cytology, Clinical Hospital

Acibadem Sistina, Skopje, Republic of North Macedonia, [90]Department of Electron

Microscopy/Molecular Pathology, The Cyprus Institute of Neurology and Genetics, Nicosia,

Cyprus, [91]VIB Center for Cancer Biology, VIB, Leuven, Belgium, [92]Laboratory for Translational

Genetics, Department of Human Genetics, University of Leuven, Leuven, Belgium,

[93]Department of Mathematics and Statistics, St. Francis Xavier University, Antigonish, Canada,

[94]Cancer Control Research, BC Cancer Agency, Vancouver, BC, Canada, [95]Department of

Molecular Medicine and Surgery, Karolinska Institutet, Stockholm, Sweden, [96]Department of

Clinical Genetics, Karolinska University Hospital, Stockholm, Sweden, [97]Department of Cancer

Epidemiology and Prevention, M. Sklodowska-Curie Cancer Center, Oncology Institute, Warsaw,

Poland, [98]General and Gastroenterology Surgery Service, Hospital Universitario Central de

Asturias, Oviedo, Spain, [99]University of Tübingen, Tübingen, Germany, [100]Translational Cancer

Research Area, University of Eastern Finland, Kuopio, Finland, [101]Institute of Clinical Medicine,

Pathology and Forensic Medicine, University of Eastern Finland, Kuopio, Finland, [102]Imaging

Center, Department of Clinical Pathology, Kuopio University Hospital, Kuopio, Finland,

[103]Department of Clinical Science and Education, Södersjukhuset, Karolinska Institutet,

Stockholm, Sweden, [104]Department of Family Medicine and Public Health, University of

California San Diego, La Jolla, CA, USA, [105]Department of Anatomical Pathology, The Alfred

Hospital, Melbourne, Victoria, Australia, [106]Department of Gynecology and Obstetrics, Ludwig

Maximilian University of Munich, Munich, Germany, [107]MRC Clinical Trials Unit at UCL, Institute

of Clinical Trials & Methodology, University College London, London, UK, [108]Department of

Obstetrics and Gynecology, Helsinki University Hospital, University of Helsinki, Helsinki, Finland,

[109]Moores Cancer Center, University of California, San Diego, La Jolla, CA, USA, [110]Department

of Family Medicine and Public Health, School of Medicine, University of California, San Diego, La

Jolla, CA, USA, [111]Clinical Genetics Research Lab, Department of Cancer Biology and Genetics,

Memorial Sloan-Kettering Cancer Center, New York, NY, USA, [112]Clinical Genetics Service,

Department of Medicine, Memorial Sloan-Kettering Cancer Center, New York, NY, USA,

[113]Centre for Cancer Research and Cell Biology, Queen's University Belfast, Belfast, Ireland, UK,

[114]Department of Non-Communicable Disease Epidemiology, London School of Hygiene and

Tropical Medicine, London, UK, [115]Human Genotyping-CEGEN Unit, Human Cancer Genetic

Program, Spanish National Cancer Research Centre, Madrid, Spain, [116]Laboratory of Cancer

Genetics and Tumor Biology, Cancer and Translational Medicine Research Unit, Biocenter Oulu,

University of Oulu, Oulu, Finland, [117]Laboratory of Cancer Genetics and Tumor Biology,

Northern Finland Laboratory Centre Oulu, Oulu, Finland, [118]Unit of Molecular Bases of Genetic

Risk and Genetic Testing, Department of Research, Fondazione IRCCS Istituto Nazionale dei

Tumori (INT), Milan, Italy, [119]Clalit National Cancer Control Center, Carmel Medical Center and

Technion Faculty of Medicine, Haifa, Israel, [120]Medical Oncology Department, Hospital

Universitario Puerta de Hierro, Madrid, Spain, [121]Institute of Pathology, Staedtisches Klinikum

Karlsruhe, Karlsruhe, Germany, [122]Department of Oncology, University Hospital of Larissa,

Larissa, Greece, [123]Prevent Breast Cancer Centre and Nightingale Breast Screening Centre,

Manchester University NHS Foundation Trust, Manchester, UK, [124]Epidemiology Branch,

National Institute of Environmental Health Sciences, NIH, Research Triangle Park, NC, USA,

[125]Research Oncology, Guy's Hospital, King's College London, London, UK, [126]Center for

Hereditary Breast and Ovarian Cancer, Faculty of Medicine and University Hospital Cologne,

University of Cologne, Cologne, Germany, [127]Department of Health Sciences Research, Mayo

Clinic College of Medicine, Jacksonville, FL, USA, [128]Division of Epidemiology, Department of

Medicine, Vanderbilt Epidemiology Center, Vanderbilt-Ingram Cancer Center, Vanderbilt

University School of Medicine, Nashville, TN, USA, [129]Ss. Cyril and Methodius University in

Skopje, Medical Faculty, University Clinic of Radiotherapy and Oncology, Skopje, Republic of

North Macedonia, [130]Population Oncology, BC Cancer, Vancouver, BC, Canada, [131]School of

Population and Public Health, University of British Columbia, Vancouver, BC, Canada, [132]Division

of Breast Cancer Research, The Institute of Cancer Research, London, UK, [133]Channing Division

of Network Medicine, Department of Medicine, Brigham and Women's Hospital and Harvard

Medical School, Boston, MA, USA, [134]Department of Epidemiology, Harvard T.H. Chan School of

Public Health, Boston, MA, USA, [135]Program in Genetic Epidemiology and Statistical Genetics,

Harvard T.H. Chan School of Public Health, Boston, MA, USA, [136]Faculty of Medicine, University

of Southampton, Southampton, UK, [137]Epigenetic and Stem Cell Biology Laboratory, National

Institute of Environmental Health Sciences, NIH, Research Triangle Park, NC, USA,

[138]Department of Epidemiology, Mailman School of Public Health, Columbia University, New

York, NY, USA, [139]Institute of Human Genetics, Pontificia Universidad Javeriana, Bogota,

Colombia, [140]Department of Epidemiology, Gilliungs School of Global Public Health and UNC

Lineberger Comprehensive Cancer Center, University of North Carolina at Chapel Hill, Chapel

Hill, NC, USA, [141]Department of Health Science Research, Division of Epidemiology, Mayo Clinic,

Rochester, MN, USA, [142]Department of Pathology, Erasmus University Medical Center,

Rotterdam, The Netherlands, [143]Biostatistics and Computational Biology Branch, National

Institute of Environmental Health Sciences, NIH, Research Triangle Park, NC, USA,

[144]Department of Molecular Genetics, University of Toronto, Toronto, ON, Canada, [145]Institute

of Environmental Medicine, Karolinska Institutet, Stockholm, Sweden, [146]Department of

Surgical Sciences, Uppsala University, Uppsala, Sweden, [147]Department of Laboratory Medicine

and Pathology, Mayo Clinic, Rochester, MN, USA, [148]Genomics Center, Centre Hospitalier

Universitaire de Québec – Université Laval, Research Center, Québec City, QC, Canada,

[149]Division of Psychosocial Research and Epidemiology, The Netherlands Cancer Institute -

Antoni van Leeuwenhoek hospital, Amsterdam, The Netherlands, [150]Department of Oncology,

School of Medicine, Johns Hopkins University, Baltimore, MD, USA

*These authors contributed equally to this work

**These authors contributed equally to this work

†Lists of participants and their affiliations appear in the Supplementary Information

Conflicts of interest: None to report

Corresponding Author

Montserrat Garcia-Closas

National Cancer Institute

Division of Cancer Epidemiology & Genetics

9609 Medical Center Drive

Rockville, MD 20850

montserrat.garcia-closas@nih.gov

Abstract

Genome-wide association studies (GWAS) have identified over 170 common breast cancer susceptibility variants using standard GWAS methods. Many of these variants have differential associations by estrogen receptor (ER), but how these variants relate with other tumor features and intrinsic molecular subtypes is unclear. We used two-stage polytomous logistic regression models to evaluate variants in relation to multiple tumor features (ER, progesterone receptor (PR), human epidermal growth factor receptor 2 (HER2) and grade) adjusting for each other, and to intrinsic-like subtypes. Eighty-five of 178 variants were associated with at least one tumor feature (false discovery rate <5%), most commonly ER and grade followed by PR and HER2. Case-control comparisons among these 85 variants identified 65 variants strongly or exclusively associated (P<0.05) with luminal-like subtypes, 5 variants associated with all subtypes at differing strengths and 15 variants primarily associated with non-luminal subtypes, especially triple-negative (TN) disease. Five variants were associated with risk of Luminal A-like and TN subtypes in opposite directions. This report demonstrates a high level of complexity in the etiology heterogeneity of breast cancer susceptibility loci and can inform investigations of subtype-specific risk prediction.

Introduction

Breast cancer represents a heterogenous group of diseases with different molecular and clinical features[1]. Clinical assessment of estrogen receptor (ER), progesterone receptor (PR), human epidermal growth factor receptor 2 (HER2) and histological grade are routinely determined to inform treatment strategies and prognostication.[2] Combined, these tumor features define five intrinsic-like subtypes (i.e. luminal A-like, luminal B/HER2-negative-like, luminal B-like, HER2-enriched-like, and basal-like/triple negative) that are correlated with intrinsic subtypes defined by gene expression panels[2]. Most known breast cancer risk or protective factors are related to luminal or hormone receptor (ER or PR; HR) positive tumors, whereas less is known about the etiology of triple-negative (TN) tumors, an aggressive subtype[3, 4].

Breast cancer genome-wide association studies (GWAS) have identified over 170 common susceptibility variants, most of them single nucleotide polymorphisms (SNPs), of which many are differentially associated with ER-positive than ER-negative disease[5, 6, 7]. These include 20 variants that primarily predispose to ER-negative or TN disease[6, 7]. However, few studies have evaluated variant associations with other tumor features, or simultaneously studied multiple, correlated tumor markers to identify source(s) of etiologic heterogeneity[6, 8, 9, 10, 11, 12]. We recently developed a two-stage polytomous logistic regression method that efficiently characterizes etiologic heterogeneity while accounting for tumor marker correlations and missing tumor data[13, 14]. This method can help describe complex relationships between susceptibility variants and multiple tumor features, helping to clarify breast cancer subtype etiologies and increasing the power to generate more accurate risk estimates between

susceptibility variants and less common subtypes. We recently demonstrated the power of this

method in a GWAS to identify novel breast cancer susceptibility accounting for tumor

heterogeneity[14].

In this report, we sought to expand our understanding of etiologic heterogeneity among

breast cancer subtypes, by applying the two-stage polytomous logistic regression methodology

to a large study population from the Breast Cancer Association Consortium (BCAC) for detailed

characterization of risk associations of 178 known risk variants by tumor subtypes defined by

ER, PR, HER2 and tumor grade.

Results

The mean (SD) ages at diagnosis (cases) and enrollment (controls) were 56.6 (12.2) and

56.4 (12.2) years, respectively. Eighty-one percent of tumors were ER-positive, 68% PR-

positive, 83% HER2-negative and 69% grade 1 or 2 (**Table 1; Supplementary Table 1**). The

most common intrinsic-like subtype was luminal A-like (59%), followed by TN (13%), luminal

B/HER2-negative-like (12%), Luminal B-like (12%) and HER2-enriched-like (5%; **Table 1**).

Global test for heterogeneity (primary hypothesis)

The two-stage models fitted for each of the 178 variants included terms for ER, PR, HER2

and grade, and identified 85 of 178 variants (47.7%) with evidence for heterogeneity by at least

one tumor feature (FDR<5%; **Figure 1 and 2; Supplementary Fig. 1**).

Description of sources of heterogeneity by each tumor feature

Marker-specific analyses identified ER and grade as the two features that most often

contributed to the observed heterogeneity (45 and 34 variants respectively had case-case

marker-specific P<0.05), and 30 variants were associated with more than one tumor

characteristic (**Figure 1 and 2; Supplementary Fig. 1)**. Seventeen of these 85 variants showed

no associations in case-case comparisons with any individual tumor marker at P<0.05 in their

corresponding fixed-effect two-stage models (**Supplementary Fig. 1**). Twenty-one variants

showed evidence for having ER as the only source of heterogeneity (**Supplementary Fig. 1**

footnote 'm'), 12 variants showed evidence for having grade as the only source of

heterogeneity (**Supplementary Fig. 1 footnote 'l'**), four variants showed evidence for PR as the

only source of heterogeneity (**Supplementary Fig. 1 footnote 'n'**) and one variant showed

evidence for HER2 as the only source of heterogeneity (**Supplementary Fig. 1 footnote 'o'**).

Description of variant associations with intrinsic-like molecular tumor subtypes

Case-control comparisons for the 85 variants with evidence for global heterogeneity

identified four main clusters of variants according to the p-values for risk associations with each

subtype (**Figure 3 and Supplementary Fig. 2**). Sixty-five variants in cluster 1 (n=3 variants) and

cluster 4 (n=62 variants) showed the strongest evidence for associations with risk for luminal-

like subtypes, cluster 2 (n=5 variants) was associated with risk for all subtypes at varying

strengths and cluster 3 (n=15 variants) with stronger evidence for TN or non-luminal subtype

associations. Five variants (rs67397200, rs6678914 and rs4577244 in cluster 3 and rs7072776

and rs17879961 in cluster 4) were found associated in opposite directions with luminal A-like

and TN subtypes. **Supplementary Table 2** shows the case-control associations between all 178

variants and the intrinsic-like subtypes.

Cluster 1 included two correlated (r^2=0.73) variants at 10q26.13, rs2981578 and

rs35054928, that were strongly associated in case-control analyses with risk of all the luminal-

like subtypes (e.g. OR (95%CI)=1.29 (1.27 to 1.31), P=2.01x10^{-231} and OR (95%CI)=1.35 (1.33 to

1.37), P=5.00x10^{-300} for luminal A-like, respectively), weakly associated with risk of HER2-

enriched-like subtype (OR (95%CI)=1.11 (1.05 to1.16), P=6.32x10^{-05} and OR (95% CI)=1.10 (1.04

to 1.15), P=3.07x10^{-4}, respectively), but not associated (P>0.05) with risk of TN disease (**Figures**

3-4 and Supplementary Fig. 2). In case-case comparisons both rs2981578 and rs35054928

showed the strongest evidence for associations with ER (**Supplementary Fig. 1**). In the

extended two-stage model that additionally included a TN term to test for heterogeneity

between TN vs non-TN subtypes both rs2981578 and rs35054928 were associated with TN

status (**Supplementary Fig. 3**). A third variant in 10q26.13, rs45631563, showed similar

association patterns, but fell into cluster 4 since, unlike rs2981578 and rs35054928, it was not

associated (P>0.05) in case-control analyses with risk of the HER2-enriched-like subtype (**Figure**

3 and Supplementary Figs. S2, S4). Case-case comparisons for rs45631563 showed associations

with ER and grade. A third variant in cluster 1, 16q12.1-rs4784227, was most strongly

associated with risk of luminal-like subtypes (e.g. OR (95%CI)=1.28 (1.26 to 1.30), P=6.68x10^{-176}

for luminal A-like), and the HER2-enriched-like tumors (OR (95%CI)=1.26 (1.19 to 1.33),

P=3.04x10^{-16}) and weaker so with TN disease (OR (95%CI)=1.11 (1.08 to 1.15), P=6.05x10^{-10};

Figures 3-4 and, Supplementary Figs. 2). In case-case analyses, rs4784227 was associated with

all tumor markers, particularly PR (**Supplementary Fig. 1**).

Five variants in cluster 2 were associated in case-control comparisons, to different

extents, with risk of all five intrinsic-like subtypes. 6q25-rs2747652 and 1p36.22-rs616488 were

associated particularly with risk of HER2-positive subtypes (**Figures 3-4**; **Supplementary Fig. 2**),

and in case-case comparisons these variants were associated with HER2 status and grade

(**Supplementary Fig. 1**). Two additional variants in 6q25 showed the strongest evidence for

being associated with risk of TN disease (OR (95%CI)=1.30 (1.24 to 1.38), P=8.26x10^{-23} for

rs9397437 and OR (95%CI)=1.15 (1.12 to 1.19), P=1.24x10^{-19} for rs3757322; **Figures 3-4**;

Supplementary Fig. 2), and in case-case comparisons they were associated with ER and grade

(**Supplementary Fig. 1**). 13q13.1-rs11571833 was associated in case-control comparisons with

risk of all subtypes, but case-case comparisons showed an association only with grade

(**Supplementary Fig. 1**). In case-control comparisons the ORs (95% CIs) for rs11571833 with

grade 3, grade 2 and grade 1 subtypes were: 1.48 (1.36 to 1.62), P=2.3x10^{-19}; 1.27 (1.18 to

1.35), P=5.0x10^{-12}; and 1.08 (0.97-1.20), P=0.15, respectively (**Supplementary Fig. 5**).

Cluster 3 included 15 variants most strongly associated in case-control comparisons with

risk of HR-negative subtypes, including three variants with the strongest evidence for

associations with risk of TN disease: 19p13.11-rs67397200 (OR (95% CI)=1.27 (1.23 to 1.31),

P=1.07x10^{-50}), 5p15.33-rs10069690 (OR (95% CI)=1.27 (1.23 to 1.31), P=3.79x10^{-48}) and 1q32.11-

rs4245739 (OR (95% CI)=1.18 (1.14 to 1.22), P=2.72x10^{-23}). Two variants at 11q22.3, rs11374964

(OR (95% CI)=0.90 (0.88-0.93), P=2.71x10^{-11}) and rs74911261 (OR (95% CI)=0.93 (0.90-0.96),

P=2.71x10^{-11}), were associated (P<0.05) only with TN disease (**Figures 3-4; Supplementary Fig.**

2). In the extended model case-case comparisons these five variants were associated with TN

status (**Supplementary Fig. 3**). The remaining 10 variants in case-control comparisons were all

associated with risk of TN disease (P<0.05; **Figures 3-4; Supplementary Fig. 3**), and in case-case

comparisons seven of these 10 variants showed the evidence for ER as the only source of

heterogeneity. 5p15.33-rs3215401 was associated with both ER and PR, and 2p24.1-

rs12710696 and 19q12-rs113701136 showed no evidence in case-case comparisons for

associations with any of the individual tumor markers (**Supplementary Fig. 1**). Besides

rs11374964 and rs74911261, variants in this cluster were not HR-negative or TN-specific

variants, as they were also associated in case-control comparisons with risk of luminal-like

subtypes. Three variants showed weak associations with luminal A-like disease (OR (95%

CI)=0.98 (0.97 to 1.00); P=0.039 for rs67397200; OR (95% CI)=1.02 (1.00 to 1.03); P=0.024 for

rs6678914; and OR (95% CI)=1.02 (1.00 to 1.04); P=0.02 for rs4577244) in an opposite direction

to their associations with TN disease (OR (95% CI)= 1.27 (1.23 to 1.31); P=1.07×10^{-50} for

rs67397200; OR (95% CI)=0.93 (0.91 to 0.96); P=1.07×10^{-4} for rs6678914 and OR (95% CI)=0.90

(0.86 to 0.93); P=2.99×10^{-9} for rs4577244; **Figures 3-4**; **Supplementary Fig. 3**).

Cluster 4 (n=62 variants) showed evidence in case-control comparisons for associations

with risk of luminal-subtypes, especially luminal A-like disease. The five variants in this cluster

showing the strongest evidence for associations with risk of luminal A-like disease were:

5q11.2-rs62355902 (OR (95% CI)=1.22 (1.20 to 1.25), P=3.94×10^{-85}), 11q13.3-rs75915166 (OR

(95% CI)=1.44 (1.40 to 1.48), P=2.26×10^{-131}), 11q13.3-rs554219 (OR (95% CI)=1.33 (1.30 to 1.37),

P=1.31×10^{-98}), 1p11.2-rs11249433 (OR (95% CI)=1.17 (1.15 to 1.19), P=5.25×10^{-90}) and 5p12-

rs10941679 (OR (95% CI)=1.20 (1.18 to 1.22), P=1.39×10^{-93}; **Figure 3 and Figures S2, S4**). In case-

case comparisons these five variants were associated with grade and with at least one other

tumor marker (**Supplementary Fig. 1**). Eighteen of the 62 variants also showed weaker

evidence in case-control comparisons for associations with non-luminal subtypes, 11 of which

were associated with risk of TN disease (P<0.05; **Figure 3 and Supplementary Figs. S2, S4**).

Notably, rs7072776 and rs17879961 (I157T) were associated in opposite directions with risk of

luminal A-like disease (OR (95% CI)=1.10 (1.08 to 1.11); P=4.96x10^{-27} for rs7072776; and OR

(95% CI)=1.44 (1.31 to 1.59); P=9.26x10^{-14} for rs17879961) and risk of TN disease (OR (95% CI)=

0.96 (0.93 to 0.99); P=0.02 for rs7072776 and OR (95% CI)=0.61 (0.47 to 0.80); P=2.55x10^{-4} for

rs17879961). In case-case comparisons rs7072776 was associated with ER and grade

(**Supplementary Fig. 1**), and rs17879961 was associated with the TN phenotype in the extended

model (**Supplementary Fig. 3**).

Supplementary Fig. 5 shows case-control associations by tumor grade for the 12

variants associated exclusively with grade in case-case comparisons. rs11571833, rs17426269

and rs11820646 showed stronger evidence for predisposing to risk of high-grade subtypes, and

the remaining variants showed stronger evidence for predisposing to risk of low-grade

subtypes.

Discussion

This study demonstrates the extent and complexity of genetic etiologic heterogeneity

among known common breast cancer susceptibility variants by multiple tumor characteristics,

using novel methodology in the largest and the most comprehensive investigation conducted to

date. We found compelling evidence that about half of the investigated breast cancer

susceptibility loci (85 of 178 variants) predispose to tumors of different characteristics. We

identified tumor grade, along with confirming ER and TN status, as important determinants of

etiologic heterogeneity. Associations with individual tumor features translated into differential

associations with the risk of intrinsic-like subtypes defined by their combinations.

Many of the variants showing subtype heterogeneity predisposed to risk of all subtypes,

but with different magnitudes. For example, 21 of 65 variants found predominantly associated

with risk of luminal-like subtypes were also associated with risk of at least one of the non-

luminal subtypes. These include variants identified in early GWAS for overall breast cancer, such

as variants in the loci for *FGFR2* (rs35054928 and rs2981578)[15, 16] and 8q24.21 (rs13281615)[15]

that were associated with luminal-like and HER2-enriched-like subtypes. rs4784227 located

near *TOX3*[15, 17] and rs62355902 located in a *MAP3K1*[15] regulatory element, were associated

with risk of all five subtypes. Of the five variants found associated in opposite directions with

luminal A-like and TN disease, we previously reported rs6678914 and rs4577244 to have

opposite effects between ER-negative and ER-positive tumors[6]. rs17879961 (I157T), a likely

causal[18] mis-sense variant located in a *CHEK2* functional domain that reduces or abolishes

substrate binding[19], was previously reported to have opposite directions of effects on lung

adenocarcinoma and lung squamous cell carcinoma and for lung cancer between smokers and

non-smokers.[20, 21] However, further studies are required to follow-up and clarify the

mechanisms for these apparent cross-over effects.

In prior ER-negative GWAS we identified 20 variants that predispose to ER-negative

disease, of which five variants were only or most strongly associated with risk of TN disease

(rs4245739, rs10069690, rs74911261, rs11374964, and rs67397200)[6, 7]. We further confirmed

these five variants to be most strongly associated with TN disease and that they were

associated in case-case comparisons with TN status in the extended model. The remaining

previously identified 15 variants all showed associations with risk of HR-negative disease, and

for all but four variants (rs17350191, rs200648189, rs6569648, and rs322144) evidence of

global heterogeneity was observed. Among the variants in cluster 3, rs3215401 was the only

variant that was not identified in a prior ER-negative GWAS[6, 7] rs3215401 was identified in a

fine-mapping analysis of *TERT* and, consistent with our findings, reported to be most strongly

associated with risk of ER-negative disease but was also associated with risk of ER-positive

disease[22].

Little is known regarding PR and HER2 as sources of etiologic heterogeneity independent

of ER or TN status. Of the four variants that showed evidence for heterogeneity only according

to PR, rs10759243[5, 23], rs11199914[5] and rs72749841[5] were previously found primarily

associated with risk of ER-positive disease, and rs10816625 was found to be associated with risk

of ER-positive/PR-positive tumors, but not other ER/PR combinations[11]. rs10995201 was the

only variant found in case-case comparisons to be solely associated with HER2 status, although

the evidence was not strong, requiring further confirmation. Previously rs10995201 showed no

evidence of being associated with ER status[24]. Among all variants found with PR or HER2

associations, few have been investigated for PR or HER2 heterogeneity while adjusting for ER[8, 9,

[10, 11, 12]. We previously reported rs10941679 to be associated with PR-status, independent of ER,

and also with grade[9]. We also found suggestive evidence of PR-specific heterogeneity for

16q12-rs3803662[12], which is in high LD ($r^2 = 0.78$) with rs4784227 (*TOX3*), which was strongly

associated with PR status. Our findings for rs2747652 are also consistent with a prior BCAC fine-

mapping analysis across the *ESR1* locus, which found rs2747652 to be associated with risk of

the HER2-enriched subtype and high grade independent of ER[8]. rs2747652 overlaps an

enhancer region and is associated with reduced *ESR1* and *CCDC170* expression[8].

Histologic grade is a composite of multiple tumor characteristics including mitotic count,

nuclear pleomorphism, and degree of tubule or gland formation[25]. Among the 12 variants

identified with evidence of heterogeneity by grade only, rs17426269, rs11820646, and

rs11571833 were found most strongly associated with risk of grade 3 disease. rs11571833 lies

in the *BRCA2* coding region and produces a truncated form of the protein[26] and has been shown

to be associated with both risk of TN disease and risk of serous ovarian tumors, both of which

tend to be high-grade[27]. To our knowledge, rs17426269 and rs11820646 have not been

investigated in relation to grade heterogeneity. The remaining 9 variants were all more strongly

associated with grade 1 or grade 2 disease. Five of these variants were previously reported to

be associated primarily with ER-positive disease[5, 28, 29], highlighting the importance of

accounting for multiple tumor characteristics to better illuminate heterogeneity sources.

For descriptive purposes, we performed clustered analyses that allowed visualization of

common heterogeneity patterns across variants. We identified 17 variants with evidence of

global heterogeneity (FDR<0.05) but no significant (marker-specific P<0.05) associations with

any of the individual tumor characteristic(s). This is likely explained by the fact that the test for

association with specific tumor markers using fixed-effects models are less powerful than

mixed-effects models used to test the primary hypothesis of global heterogeneity by any tumor

marker[13].

A major strength of our study is our large sample size of over 100,000 breast cancer

cases with tumor marker information, and a similar number of controls, making this the largest,

most comprehensive breast cancer heterogeneity investigation. Our application of the two-

stage polytomous logistic regression enabled adjusting for multiple, correlated tumor markers

and accounting for missing tumor marker data. This is a more powerful and efficient modeling

strategy for identifying heterogeneity sources among highly correlated tumor markers,

compared with standard polytomous logistic regression[13, 14]. In simulated and real data

analyses, we have demonstrated that in the presence of heterogenous associations across

subtypes, the two-stage model is more powerful than polytomous logistic regression for

detecting heterogeneity. Moreover, we have demonstrated that in the presence of correlated

markers, the two-stage model, incorporating all markers simultaneously, has much better

ability to distinguish the true source(s) of heterogeneity compared to testing for heterogeneity

by analysis of one marker at a time[13]. In prior analyses, we showed that the two-stage

polytomous regression is a powerful approach to identify susceptibility variants that display

tumor heterogeneity[14].

Our study also had limitations. First, many breast cancer cases from studies included in

this report had missing information on one or more tumor characteristics. To address this

limitation, we implemented an EM algorithm that allowed a powerful analysis to incorporate

cases with missing tumor characteristics under the assumption that tumor characteristics are

missing at random (MAR), i.e. the underlying reason for missing data may depend on observed

tumor markers or/and covariate values, but not on the missing values themselves[30]. If this

assumption is violated it can lead to an inflated type-one error[13]. However, in the context of

genetic association testing, the missingness mechanism would also need related to the genetic

variants under study, which is unlikely. Our study focused on investigating ER, PR, HER2, and

grade as heterogeneity sources and future studies with more detailed tumor characterization

could reveal additional etiologic heterogeneity sources.

In summary, our findings provide insights into the complex etiologic heterogeneity

patterns of common breast cancer susceptibility loci. These findings may inform future studies,

such as fine-mapping and functional analyses to identify the underlying causal variants,

clarifying biological mechanisms that drive genetic predisposition to breast cancer subtypes.

Moreover, these analyses provide precise estimates of relative risk for different intrinsic-like

subtypes that could improve the discriminatory accuracy of subtype-specific polygenic risk

scores[31].

Methods

Study Population and Genotyping

The study population and genotyping are described in previous publications[5, 6] and in

the **Supplemental Methods**. We included invasive cases and controls from 81 BCAC studies

with genotyping data from two Illumina genome-wide custom arrays, the iCOGS and

OncoArray (106,571 cases (OncoArray: 71,788; iCOGS: 34,783) and 95,762 controls

(OncoArray: 58,134; iCOGS: 37,628); **Supplementary Table 1**). We evaluated 178 breast

cancer susceptibility variants that were identified in or replicated by prior BCAC analyses. The

majority of these variants (n=158) were identified because of their association with risk of

overall breast cancer, and a small number of variants (n=20) were identified because of their

association specific to ER-negative breast cancer (**Supplementary Table 3**)[5, 6]. Genotypes for

the variants marking the 178 susceptibility loci were determined by genotyping with the

iCOGS and the OncoArray arrays and imputation to the 1000 Genomes Project (Phase 3)

reference panel.

Statistical Analysis

An overview of the analytic strategy is shown in **Figure** 1 and a detailed discussion of the

statistical methods, including of the two-stage polytomous logistic regression, are provided in

the **Supplementary Methods** and elsewhere[13, 14]. Briefly, our main aim was to identify which of

178 known susceptibility variants showed heterogenous associations by ER, PR, HER2 and

grade. Specifically, for each variant, the global null hypothesis was that there was no difference

in risk of breast cancer associated with the variant genotype across any of the tumor features

being evaluated.

We used a two-stage polytomous regression model to perform global test for

heterogeneity. The two-stage polytomous model allows modelling genetic association of breast

cancer accounting for possible heterogeneity in effects by combination of multiple tumor

markers using a parsimonious decomposition of subtype-specific case-control odds-ratio

parameters in terms of marker-specific case-case odd-ratio parameters. We introduced further

parsimony by using mixed-effect formulation of the model that allows ER-specific case-case

parameters to be treated as fixed and similar parameters for other markers (PR, HER2 and

grade) as random. We used an EM algorithm[32] for parameter estimation under this model to

account for missing data in tumor characteristics. A detailed description of the proposed

method and assessment of performance in simulation and real studies are presented in the

Supplementary Methods and as a separate manuscript[13].

In the main analysis, we accounted for multiple testing (178 tests, one for each of

variant) of the global heterogeneity test using a false discovery rate (FDR) <0.05 under the

Benjamini-Hochberg procedure[33]. For the variants showing evidence for global heterogeneity

after FDR adjustment, we further evaluated which of the tumor features contributed to the

heterogeneity by fitting a fixed-effects two-stage model that simultaneously tested for

associations with each tumor feature (this model was fitted for each variant separately). We

used a threshold of $P<0.05$ for marker-specific tumor heterogeneity tests to describe which

specific tumor marker(s) contributed to the observed heterogeneity, adjusting for the other

tumor markers in the model. This p-value threshold was used only for descriptive purposes, as

the primary hypotheses were tested using the FDR-adjusted global test for heterogeneity

described above.

We conducted various additional analyses to explore for evidence of heterogeneity. We

fitted an extended two-stage model with an additional term for TN status to test for differences

between TN vs non-TN subtypes. We also fitted a two-stage model to estimate case-control ORs

and 95% confidence intervals (CI) between the variants and five intrinsic-like subtypes defined

by combinations of ER, PR, HER2 and grade (**Supplementary Methods**): (1) luminal A-like, (2)

luminal B/HER2-negative-like, (3) luminal B-like, (4) HER2-enriched-like and (5) TN. We

performed sensitivity analyses, in which we estimated the ORs and 95% CI between the variants

and the intrinsic-like subtypes by implementing a standard polytomous model restricted to

cases with complete tumor marker data. For all analyses we analyzed OncoArray and iCOGS

array data separately, adjusting for the first 10 principal components for ancestry-informative

variants, and then meta-analyzed the results. To help describe sources of heterogeneity, we

performed cluster analyses based on Euclidean distance calculated from z-statistics that were

estimated by the individual marker-specific tumor heterogeneity tests and by the case-control

tests between variants and the intrinsic-like subtypes. Clustering was performed in R using the

function Heatmap as implemented by the package "Complex Heatmap" version 3.1[34].

Data Availability

Requests for data can be made to the corresponding author or the Data Access Coordination

Committees (DACCs) of BCAC (see above URL). BCAC DACC approval is required to access data

from the ABCFS, ABCS, ABCS-F, ABCTB, AHS, BBCC, BBCS, BCEES, BCFR-NY, BCFR-PA, BCFR-UT,

BCINIS, BIGGS, BSUCH, CBCS, CECILE, CNIO-BCS, CTS, DIETCOMPLYF, ESTHER, FHRISK, GC-HBOC,

GENICA, GEPARSIXTO, GESBC, HABCS, HCSC, HEBCS, HUBCS, KARBAC, KARMA, KBCP,

KCONFAB/AOCS, LMBC, MABCS, MARIE, MBCSG, MCBCS, MISS, MMHS, MTLGEBCS, NBCS, NC-

BCFR, NCBCS, OBCS, OFBCR, ORIGO, PKARMA, POSH, PREFACE, PROCAS, RBCS, SASBAC, SBCS,

SKKDKFZS, SUCCESSB, SUCCESSC, SZBCS, TNBCC, UCIBCS, UKBGS, UKOPS and USRT studies

(**Supplementary Table 1**).

569
570

Funding and Acknowledgments

This project has been funded in part with Federal funds from the National Cancer

Institute Intramural Research Program, National Institutes of Health. Dr. Nilanjan Chatterjee

was supported by NHGRI (1R01 HG010480-01). OncoArray genotyping was funded by the

government of Canada through Genome Canada and the Canadian Institutes of Health

Research (GPH-129344), the Ministère de l'Économie, de la Science et de l'Innovation du

Québec through Génome Québec, the Quebec Breast Cancer Foundation for the PERSPECTIVE

project, the US National Institutes of Health (NIH) (1 U19 CA 148065 for the Discovery, Biology

and Risk of Inherited Variants in Breast Cancer (DRIVE) project and X01HG007492 to the Center

for Inherited Disease Research (CIDR) under contract HHSN268201200008I), Cancer Research

UK (C1287/A16563), the Odense University Hospital Research Foundation (Denmark), the

National R&D Program for Cancer Control–Ministry of Health and Welfare (Republic of Korea)

(1420190), the Italian Association for Cancer Research (AIRC; IG16933), the Breast Cancer

Research Foundation, the National Health and Medical Research Council (Australia) and

German Cancer Aid (110837).

iCOGS genotyping was funded by the European Union (HEALTH-F2-2009-223175),

Cancer Research UK (C1287/A10710, C1287/A10118 and C12292/A11174]), NIH grants

(CA128978, CA116167 and CA176785) and the Post-Cancer GWAS initiative (1U19 CA148537,

1U19 CA148065 and 1U19 CA148112 (GAME-ON initiative)), an NCI Specialized Program of

Research Excellence (SPORE) in Breast Cancer (CA116201), the Canadian Institutes of Health

Research (CIHR) for the CIHR Team in Familial Risks of Breast Cancer, the Ministère de

l'Économie, Innovation et Exportation du Québec (PSR-SIIRI-701), the Komen Foundation for

the Cure, the Breast Cancer Research Foundation and the Ovarian Cancer Research Fund.

A full description of the funding and acknowledgments is provided in the

Supplementary Information.

References

1.   Cancer Genome Atlas N. Comprehensive molecular portraits of human breast tumours. *Nature*
490, 61-70 (2012).

2.   Curigliano G*, et al.* De-escalating and escalating treatments for early-stage breast cancer: the St.
Gallen International Expert Consensus Conference on the Primary Therapy of Early Breast
Cancer 2017. *Ann Oncol* **28**, 1700-1712 (2017).

3.   Barnard ME, Boeke CE, Tamimi RM. Established breast cancer risk factors and risk of intrinsic
tumor subtypes. *Biochim Biophys Acta* **1856**, 73-85 (2015).

4.   Yang XR*, et al.* Associations of breast cancer risk factors with tumor subtypes: a pooled analysis
from the Breast Cancer Association Consortium studies. *Journal of the National Cancer Institute*
103, 250-263 (2011).

5.   Michailidou K*, et al.* Association analysis identifies 65 new breast cancer risk loci. *Nature* **551**,
92-94 (2017).

6.   Milne RL*, et al.* Identification of ten variants associated with risk of estrogen-receptor-negative
breast cancer. *Nature genetics* **49**, 1767-1778 (2017).

7.   Garcia-Closas M*, et al.* Genome-wide association studies identify four ER negative-specific breast
cancer risk loci. *Nature genetics* **45**, 392-398, 398e391-392 (2013).

8.   Dunning AM*, et al.* Breast cancer risk variants at 6q25 display different phenotype associations
and regulate ESR1, RMND1 and CCDC170. *Nature genetics* **48**, 374-386 (2016).

9.   Milne RL*, et al.* Confirmation of 5p12 as a susceptibility locus for progesterone-receptor-
positive, lower grade breast cancer. *Cancer epidemiology, biomarkers & prevention : a*
publication of the American Association for Cancer Research, cosponsored by the American
Society of Preventive Oncology **20**, 2222-2231 (2011).

10.   Figueroa JD*, et al.* Associations of common variants at 1p11.2 and 14q24.1 (RAD51L1) with
breast cancer risk and heterogeneity by tumor subtype: findings from the Breast Cancer
Association Consortium. *Human molecular genetics* **20**, 4693-4706 (2011).

11.   Orr N*, et al.* Fine-mapping identifies two additional breast cancer susceptibility loci at 9q31.2.
Human molecular genetics **24**, 2966-2984 (2015).

12.   Broeks A*, et al.* Low penetrance breast cancer susceptibility loci are associated with specific
breast tumor subtypes: findings from the Breast Cancer Association Consortium. *Human*
molecular genetics **20**, 3289-3303 (2011).

639
640   13.   Zhang H, Zhao N, Ahearn TU, Wheeler W, García-Closas M, Chatterjee N. A mixed-model
641   approach for powerful testing of genetic associations with cancer risk incorporating tumor
642   characteristics. *Biostatistics*, (2020).

643
644   14.   Zhang H*, et al.* Genome-wide association study identifies 32 novel breast cancer susceptibility
645   loci from overall and subtype-specific analyses. *bioRxiv*, 778605 (2019).

646
647   15.   Easton DF*, et al.* Genome-wide association study identifies novel breast cancer susceptibility
648   loci. *Nature* **447**, 1087-1093 (2007).

649
650   16.   Hunter DJ*, et al.* A genome-wide association study identifies alleles in FGFR2 associated with risk
651   of sporadic postmenopausal breast cancer. *Nature genetics* **39**, 870-874 (2007).

652
653   17.   Stacey SN*, et al.* Common variants on chromosomes 2q35 and 16q12 confer susceptibility to
654   estrogen receptor-positive breast cancer. *Nature genetics* **39**, 865-869 (2007).

655
656   18.   Fachal L*, et al.* Fine-mapping of 150 breast cancer risk regions identifies 191 likely target genes.
657   *Nature genetics* **52**, 56-73 (2020).

658
659   19.   Li J*, et al.* Structural and functional versatility of the FHA domain in DNA-damage signaling by the
660   tumor suppressor kinase Chk2. *Mol Cell* **9**, 1045-1054 (2002).

661
662   20.   McKay JD*, et al.* Large-scale association analysis identifies new lung cancer susceptibility loci and
663   heterogeneity in genetic susceptibility across histological subtypes. *Nature genetics* **49**, 1126-
664   1132 (2017).

665
666   21.   Wang Y*, et al.* Rare variants of large effect in BRCA2 and CHEK2 affect risk of lung cancer. *Nature*
667   *genetics* **46**, 736-741 (2014).

668
669   22.   Bojesen SE*, et al.* Multiple independent variants at the TERT locus are associated with telomere
670   length and risks of breast and ovarian cancer. *Nature genetics* **45**, 371-384, 384e371-372 (2013).

671
672   23.   Li X*, et al.* Association of multiple genetic variants with breast cancer susceptibility in the Han
673   Chinese population. *Oncotarget* **7**, 85483-85491 (2016).

674
24.   Darabi H*, et al.* Polymorphisms in a Putative Enhancer at the 10q21.2 Breast Cancer Risk Locus
Regulate NRBF2 Expression. *American journal of human genetics* **97**, 22-34 (2015).

25. Elston CW, Ellis IO. Pathological prognostic factors in breast cancer. I. The value of histological
grade in breast cancer: experience from a large study with long-term follow-up. *Histopathology*
19, 403-410 (1991).

26. Mazoyer S*, et al.* A polymorphic stop codon in BRCA2. *Nature genetics* **14**, 253-254 (1996).

27. Meeks HD*, et al.* BRCA2 Polymorphic Stop Codon K3326X and the Risk of Breast, Prostate, and
Ovarian Cancers. *Journal of the National Cancer Institute* **108**, (2016).

28. Darabi H*, et al.* Fine scale mapping of the 17q22 breast cancer locus using dense SNPs,
genotyped within the Collaborative Oncological Gene-Environment Study (COGs). *Scientific*
reports **6**, 32512 (2016).

29. Michailidou K*, et al.* Large-scale genotyping identifies 41 new loci associated with breast cancer
risk. *Nature genetics* **45**, 353-361, 361e351-352 (2013).

30. Little RJA, Rubin DB. Statistical analysis with missing data. In: *Wiley series in probability and*
statistics). Third edition edn. Wiley, (2019).

31. Mavaddat N*, et al.* Polygenic Risk Scores for Prediction of Breast Cancer and Breast Cancer
Subtypes. *American journal of human genetics* **104**, 21-34 (2019).

32. Dempster AP, Laird NM, Rubin DB. Maximum Likelihood from Incomplete Data Via Em
Algorithm. *J Roy Stat Soc B Met* **39**, 1-38 (1977).

33. Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and Powerful
Approach to Multiple Testing. *Journal of the Royal Statistical Society Series B (Methodological)*
57, 289-300 (1995).

34. Gu Z, Eils R, Schlesner M. Complex heatmaps reveal patterns and correlations in
multidimensional genomic data. *Bioinformatics* **32**, 2847-2849 (2016).

Figure 1. Overview of the analytic strategy and results from the investigation of 178 known breast cancer susceptibility variants for evidence of heterogeneity according to the estrogen receptor (ER), progesterone receptor (PR), human epidermal growth factor receptor 2 (HER2), and grade

a=We evaluated 178 variants reported by Michailidou k, et al Nature 2017 and Milne RL, et al Nature Genetics 2017 that were associated with breast cancer at a p-value of $<10^{-4}$

b=Model 1 (Primary analyses): Mixed-effect two-stage polycotomos model (ER as fixed-effect, and PR, HER2 and grade as random-effects) to test for global heterogeneity between each individual susceptibility variant and any of the tumor features (separate models were fit for each variant).

c=Model 2: Fixed effect two-stage polycotomos model for marker-specific tumor heterogeneity tests between each individual variant and each of the tumor features (ER, PR, HER2, and grade) mutually adjusted for each other (separate models were fit for each variant).

d=Model 3: Fixed effect two-stage polycotomos model for intrinsic-like subtypes: luminal A-like, luminal B/HER2-negative-like, luminal B-like, HER2-enriched-like, and basal-like/triple negative.

Note: In extended analyses, we fitted similar mixed and fixed effects models to evaluate heterogeneity according to ER, PR, HER2, grade and/or triple-negative status (TN; **Supplementary Fig. 3**)

Figure 2. Heatmap and clustering of the P-values from the two-stage model's case-case marker-specific heterogeneity test for associations between each of the 178 breast cancer susceptibility variants and estrogen receptor (ER), progesterone receptor (PR), human epidermal growth factor receptor 2 (HER2) or grade, adjusting for principal components and each tumor marker. Columns represent individual variants. For more detailed information on the context of figure see **Supplementary Fig. 1.**

Figure 3. Heatmap and clustering of case-control P-values and odds ratios (OR) from two-stage models for intrinsic-like subtypes, fitted for each of the 85 breast cancer susceptibility variants with evidence for global heterogeneity. Columns are labeled according to variants and loci. (**A**) Clustering based on case-control p-value of the associations between susceptibility variants and breast cancer intrinsic-like subtypes[a,b]. (**B**) Heatmap of ORs between susceptibility variants and

intrinsic-like subtypes. For more detailed information on the context of figure see **Supplementary Fig. 2.**

a = luminal A-like (ER+ and/or PR+, HER2-, grade 1 & 2); luminal B/HER2-negative-like (ER+ and/or PR+, HER2-, grade 3); luminal B-like (ER+ and/or PR+, HER2+); HER2-enriched-like (ER- and PR-, HER2+); TN (ER-, PR-, HER2-)

b = Clusters: 1 and 4: strongest associations with hormone receptor (HR)-positive subtypes; Cluster 2: Associated with all subtypes at varying strengths, and Cluster 3: strongest associations with TN or non-luminal subtypes

c = 32 variants associated ($p<0.05$) with risk of the TN subtype

d = variants previously identified as primarily predisposing to ER-negative or TN disease as reported in Milne RL, et al. Nat Genet 2017;49(12):1767-1778

e = 21 variants with evidence for having ER as the only source of heterogeneity

f = 12 variants with evidence for having grade as the only source of heterogeneity

g = 1 variants with evidence for having HER2 as the only source of heterogeneity

h = 4 variants with evidence for having PR as the only source of heterogeneity

Figure 4. Associations[a] between breast cancer susceptibility variants with evidence of heterogeneity in the two-stage model[b] and intrinsic-like subtypes. variants presented in order as shown in **figure 3**, clusters 1, 2 and 3.

[a] = Per-minor allele odds ratio (95% confidence limits).

[b] = Two-stage model testing for heterogeneity according to estrogen receptor (ER), progesterone receptor (PR), human epidermal growth factor receptor 2 (HER2) and grade

[c] = luminal A-like (ER+ and/or PR+, HER2-, grade 1 & 2); luminal B/HER2-negative-like (ER+ and/or PR+, HER2-, grade 3); luminal B-like (ER+ and/or PR+, HER2+); (4) HER2-enriched-like (ER- and PR-, HER2+), and triple-negative (ER-, PR-, HER2-)

[d] = variants previously identified as primarily predisposing to ER-negative or TN disease as reported in Milne RL, et al. Nat Genet 2017;49(12):1767-1778.

Table 1. Distribution of estrogen receptor (ER), progesterone receptor (PR), human epidermal growth factor receptor 2 (HER2), and grade and the intrinsic-like subtypes [a,b] among invasive cases of breast cancer in studies from the Breast Cancer Consortium Association.

Tumor Marker	N (%)
ER	
Negative	16,900 (19)
Positive	70,030 (81)
Unknown	19,641
PR	
Negative	24,283 (32)
Positive	51,603 (68)
Unknown	30,685
HER2	
Negative	47,693 (83)
Positive	9,529 (17)
Unknown	49,349
Grade	
1	15,583 (20)
2	37,568 (49)
3	24,382 (31)
Unknown	29,038
Intrinsic-like subtypes	
Luminal A-like	33,083 (59)
Luminal B/Her2-negative-like	6,804 (12)
Luminal B-like	6,511 (12)
HER2-enriched-like	2,797 (5)
Triple-negative	7,178 (13)

a=luminal A-like (ER+ and/or PR+, HER2-, grade 1 & 2); luminal B/HER2-negative-like (ER+ and/or PR+, HER2-, grade 3); luminal B-like (ER+ and/or PR+, HER2+); HER2-enriched-like (ER- and PR-, HER2+); TN (ER-, PR-, HER2-)
b=Intrinsic subtypes defined among 56,373 cases with available tumor marker data