A quantitative method for proteome reallocation using minimal regulatory interventions

Lastiri-Pancardo Gustavo[1], Mercado-Hernandez J.S[1], Kim Juhyun [2], Jiménez José I.[2], Utrilla José[1*.]

1. Systems and Synthetic Biology Program, Centro de Ciencias Genómicas, Universidad Nacional Autónoma de México.
2. Faculty of Health and Medical Sciences, University of Surrey, Guildford, UK

*Corresponding author:
Name: José Utrilla
Address: UNAM-Centro de Ciencias Genómicas. Av. Universidad s/n Col. Chamilpa. Cuernavaca, Morelos Mexico. 62210
utrilla@ccg.unam.mx

Keywords: proteome minimization, regulatory network, Escherichia coli, rational design, synthetic biology, systems biology

Highlights

- Proteome reduction with minimal genetic intervention as design principle
- Regulatory and proteomic data integration to identify transcription factor activated proteome
- Deletion of the TF combination that reduces the greater proteomic load
- Regulatory interventions are highly specific
- Designed strains show less burden, improved protein and violacein production

Abstract

Engineering resource allocation in biological systems for synthetic biology applications is an ongoing challenge. Wild type organisms allocate abundant cellular resources for ensuring survival in changing environments, reducing the productivity of engineered functions. Here we present a novel approach for engineering the resource allocation of *Escherichia coli* by rationally modifying the transcriptional regulatory network of the bacterium. Our method (ReProMin) identifies the minimal set of genetic interventions that maximise the savings in cell resources that would normally be used to express non-essential genes. To this end we categorize Transcription Factors (TFs) according to the essentiality of the genes they regulate and we use available proteomic data to rank them based on its proteomic balance, defined as the net proteomic charge they release. Using a combinatorial approach, we design the removal of TFs that maximise the release of the proteomic charge and we validate the model predictions experimentally. Expression profiling of the resulting strain shows that our designed regulatory interventions are highly specific. We show that our resulting engineered strain containing only three mutations, theoretically releasing 0.5% of their proteome, has higher proteome budget and show increased production yield of a molecule of interest obtained from a recombinant metabolic pathway. This approach shows that combining whole-cell proteomic and regulatory data is an effective way of optimizing strains in a predictable way using conventional molecular methods.

Importance

Biological regulatory mechanisms are complex and occur in hierarchical layers such as transcription, translation and post-translational mechanisms. We foresee the use of regulatory mechanism as a control layer that will aid in the design of cellular phenotypes. Our ability to engineer biological systems will be dependent on the understanding of how cells sense and respond to their environment at a system level. Few studies have tackled this issue and none of them in a rational way. By developing a workflow of engineering resource allocation based on our current knowledge of *E. coli*'s regulatory network, we pursue the objective of minimizing cell proteome using a minimal genetic intervention principle. We developed a method to rationally design a set of genetic interventions that reduce the hedging proteome allocation. Using available datasets of a model bacterium we were able to reallocate parts of the unused proteome in laboratory conditions to the production of an engineered task. We show that we are able to reduce the unused proteome (theoretically 0.5%) with only three regulatory mutations designed in a rational way, which

66   results in strains with increased capabilities for recombinant expression of pathways of
67   interest.

Introduction

The removal of accessory non-essential functions is one of the strategies used to engineer microbial phenotypes. This approach relies on the assumption that cellular resources for gene expression are limited and, therefore, by removing unneeded genes in a certain environment, the cell is capable of allocating resources to other functions (e.g. expression of recombinant genes). These minimization approaches are mostly done by reducing genome size and gene number including performing random deletions[1,2], however, the precise way in which the resource allocation takes place after the genetic intervention is not considered.

Organisms respond to the environment by cellular signalling encoded in regulatory networks[3]. The intricacy of the lifestyle of an organism is generally translated into signalling complexity[4]. Biological regulatory networks are robust[5] and evolvable[6] to cope with environmental and lifestyle perturbations, however this robustness involves intrinsic trade-offs, such as resource allocation strategies. It has been shown that cellular states are naturally "primed" for typical upcoming changes. Bacteria anticipate to fluctuations in the environment[7,8], draining resources from functions that are mostly performed in relatively stable conditions. The expression of anticipation functions, also called hedging functions, is encoded in the regulatory network and it has a proteomic cost[9]. Genome scale models along with experimental data sets enable the calculation of the minimal theoretical proteome needed to sustain growth in a defined condition[10]. Therefore, comparing minimal theoretical proteomes with measured proteomes reveal the costs of the hedging proteome allocation. Proteome econometric approaches can facilitate the engineering of cellular states or phenotypes aimed at displaying an engineered function. Recent studies have focused in the host-construct interactions for increasing predictability of synthetic constructs[11–13]. In addition to these approaches, the rational design of the host used for expression following econometric models can be adopted to improve the performance of synthetic constructs, including production phenotypes for molecules of added value. Among other benefits, streamlined organisms obtained this way are less likely to develop undesired emerging behaviours[11].

In this work we develop a new top- down cell engineering strategy for *Escherichia coli* using the Transcriptional Regulatory Network (TRN) as a control layer for proteome allocation. By combining high-throughput proteomic information, regulatory network interactions and gene essentiality observations, we develop a method capable of finding the minimal set of genetic

interventions required to divert the resources invested in superfluous hedging into increased

biosynthetic potential. The resulting strain exhibits an increased availability of cellular

resources to express engineered functions.

Results

Combining gene essentiality and TRN analysis identifies dispensable TFs for

proteome reduction in a defined condition

The genome scale model of Metabolism and gene Expression (ME model) computes the

minimal theoretical proteome and allows calculating the cost of expressing hedging

functions. It can be used to simulate different scenarios of expression of the hedging

proteome (as unused protein fraction coefficient, see methods)[14]. These simulations allow

us to calculate the costs and the benefits of different interventions, e.g. by modulating the

expression of the hedging proteome, expressed in terms of growth, the size of both essential

and recombinant proteome sectors (Supp. fig. 1).

We build on ME-models to design strains containing the minimal genetic interventions that

reduce the greatest amount of proteomic resources not required to grow in a specific

condition. Our method uses Transcription Factors (TFs) as the key dials controlling the

allocation of the hedging proteome in a pre-defined specific environment. We begin by

establishing batch growth in minimal media (M9) supplemented with glucose as the sole

carbon source as the defined environment for the first case of this study. Then, by compiling

experimental and genome-scale model generated essential gene sets, we generated a list

of essential genes for growth in this specific environment (Figure 1A, Supp. Table 1, see

methods). Once the case specific gene essentiality is defined, we analysed the TF-gene

interactions compiled in RegulonDB [15]. After determining gene essentiality and TF-gene

regulatory interactions, we analyse the sub-network of interactions of each TF (Figure 1B)

looking for dispensable TFs, defined as those that do not activate the expression of any

essential gene. According to our analysis, 156 from the 200 TFs contained in the regulatory

network can be eliminated (Figure 1C). Since our goal is to reduce the hedging proteome,

out of the 156 dispensable TFs we select as candidates for non-essential function reduction

those 34 TF´s with at least one unique (meaning it is not activated by any other TF) positive

regulated gene (Supp. Table 2) (See methods); this gives the certainty of silencing at least

one gene.

Figure 1 Gene essentiality and transcriptional regulatory network analysis in the pre-defined

condition. A) Essentiality profile of *E. coli* genome under selected growth condition. The essential

fraction of the genome is divided into core (always needed) and conditional (M9-glucose needed). **B)**

Graphical representation of the sub-network of interactions considered for the classification of the

TFs; grey squares represent essential TFs, light green squares dispensable TFs, green squares

candidate TFs, dark blue circles essential genes, light blue circles dispensable genes and arrows

positive interactions. **C)** Essentiality profile of the TFs contained in RegulonDB. Total *E. coli* genes

according to Ecocyc[16].

Integration of proteomic data and the TF-Gene regulatory network

We determine the proteome associated to each non-essential TF in our network integrating

a quantitative proteomic data set[17], that provides protein copy number per cell under 22

different growth conditions with 95% of proteome coverage (by mass). Here we define two

emerging properties derived from the quantitative proteomics data integration: the Proteomic

Load of a gene (P_L) in fg of protein per cell (Figure 2A) and the Proteomic Balance (P_B) of a

TF resulting from the summation of the P_L of the genes that would result silenced or activated

by the elimination of a TF (Figure 2B). P_B is conceptually important to rank the TFs according

to the size of the proteome they control, since it takes into account the net addition of protein

mass (in fg of protein/cell) liberated when removing a TF. A graphic representation of the 34

candidate TF subnetwork illustrating the P_L all the possible targets to affect is shown in

Figure 4A.

$$P_L = \frac{MW_P * C_{cell}}{N_A}$$

$$P_B = \sum SilencedP_L - \sum InducedP_L$$

Figure 2 Emerging properties from proteomics data integration. A) Proteomic load of a gene

(P_L), this property is defined as the molecular weight of the protein (MW_P) multiplied by the number of

copies per cell (C_{cell}) divided by Avogadro's number (N_A) (6.022×10^8 fg equivalent), the more

expressed the gene is, the more proteomic load it generates. **B)** Proteomic Balance of a TF (P_B)

which is defined as the sum of the P_L of the silenced genes minus the sum of the P_L of the induced

genes. **C)** Schematic of a simple case of shared regulation in which removing both TFs silences all

genes but this is not the case when the TFs are silenced individually.

Computational search of minimal TF eliminations for the release of the maximal

hedging proteome

Even though *E. coli* is one of the most studied organisms and its TRN has been widely

investigated, only half of its genes have regulatory information (RegulonDB). In order to

prevent detrimental effects on gene expression due to our incomplete knowledge of the

regulatory network, we searched the smallest combination of TFs that liberate the greatest

amount of resources. We observed that many TFs have shared target genes (Figure 2C), in

fact, many of them are part of a simplified version of a Dense Overlapping Regulon (DOR)

network motif[18], meaning that a particular combination of TFs is needed in order to "fully"

silence these targets. Due to the size of the landscape of potential phenotypes resulting

from the combinatorial TFs deletions, we developed a computational tool to assist with the

design of mutant strains. We called our tool ReProMin (**Re**gulation based **Pro**teome

Minimization). ReProMin tool uses the previously described TF-Gene interaction network;

quantitative proteomic data and a list of candidate non-essential TF to find the n-combination

of mutations that silences the higher proteomic load (see methods).

In order to test our method (ReProMin), we first performed calculations using data from the

glucose minimal media condition on which we defined gene and TF essentiality. Calculations

were made considering two cases, the Shared Target (ST) case: considering all non-
essential TFs with positive P_B (132 TF), which takes into account some TFs with no unique
regulated genes. And the Unique Target (UT) case: considering candidate TFs with positive
P_B (20 TF). Our computational tool was able to solve up to triple TF combinations for ST
case and up to 20 TF combinations for UT case (Figure 3A and B).

For the ST case, calculations revealed that the elimination of all the non-essential TFs
theoretically would liberate up to 1.06% of total proteome. However, up to 0.53% of total
proteome can be released by removing a top combination of three TFs. For the UT case,
the elimination of all 20 candidate TFs would liberate up to 0.72% of the proteome, and our
simulations show that there is not a significant improvement in resource release after the
elimination of 8 TFs.

We tested the accuracy of ReProMin predictions in other conditions for which proteomic
data is available, such as growth on galactose and acetate minimal media. In this case we
used the rich media essentiality gene set (see methods) and for the environment specific
genes we performed essentiality simulations with a genome scale metabolic model in the
corresponding growth condition (see methods). As a result, we obtained 164 and 166 non-
essential TFs for galactose and acetate respectively and found that most of the identified
TFs are shared among the three evaluated (glucose, galactose, acetate minimal media)
meaning that they are all non-essential for minimal media growth with those carbon sources
(Supp. fig. 2). In both cases we identified 23 candidate TFs that belong to the UT case and
have a positive P_B. Proteome liberation calculations were made using these subsets of
candidate TFs. Our predictions show that we can release 0.88% and 0.81% of the total
proteome in galactose and acetate, respectively, with the deletion of all these TFs (Figure
3C and D).

Figure 3 Proteome liberation calculations using ReProMin. A) Potential optimization landscape corresponding to the ST case; the solved region is shown in blue while the unsolved region in red, PYC mutant location in the landscape is shown with a green circle. Solved optimization landscape for UT case in B) glucose, PFC mutant location in the landscape is shown with a green circle, C) galactose and D) acetate.

Generation of combinatorial strains

Based on our ReProMin predictions, two triple knockout strains were generated, for the ST case: PYC ($\Delta phoB$ - phosphate scavenging system, $\Delta yedW$ - unknown gene, $\Delta cusR$ - copper efflux system) with a P_B of 1.3 fg representing 0.53% of the total proteome in glucose (Figure 3A), is a particular case of shared regulation where most of the target genes are only silenced by the deletion of all the three TFs together (a graphical representation of its TF-gene network is presented in Figure 4B). For the UT case: PFC ($\Delta phoB$ - phosphate scavenging system, $\Delta flhC$ - flagella master regulator, $\Delta cueR$- copper efflux system) with a P_B of 1.08 fg representing 0.44% of the total proteome in glucose (Figure 3B), has a higher

grade of confidence in the design than PYC due to simpler regulatory subnetwork (Figure
4C). We also generated a strain, using an intuitive approach, in which we eliminated three
TFs that regulate non-growth related functions. The resulting strain is called FOG (Δ*fliA* -
flagella sigma factor, Δ*oxyR* - oxidative stress master regulator, Δ*gadE* - acid resistance
regulator). The FOG strain was not generated by our design pipeline; therefore the
regulatory interventions may affect some important functions and it was used as a control to
compare to our designed strains.

Figure 4 TF-Gene network and P$_L$ representation. A) Subnetwork of interactions corresponding to
the 34 candidate TF representing all the potentially affected targets; green squares represent
candidate TF, while light blue circles represent dispensable genes, dark blue circles are essential
genes and light green squares are dispensable TF, circle size is proportional to the P$_L$ of the gene.
Subnetwork of predicted regulated targets of the PYC **(B)** and PFC **(C)** mutants. In both cases, green
circles represent predicted silenced targets, red circles predicted induced targets and yellow circles
genes with no proteomic coverage; the size of the circles is proportional to the P$_L$ of the target. High
resolution versions of the sub networks are available in the supplementary material (Supp. figs. 3 -
5).

RNA-seq analysis confirms the high specificity of introduced mutations

The predictive power of ReProMin depends on the accuracy of the interactions compiled in
the *E. coli* TRN. We validated the predictions for the PFC mutant strain by comparing its
transcriptome profile obtained by RNA-seq to that of the wild-type (WT). This experiment
aims at determining the degree of success in gene silencing at the transcriptional level, and
at assessing other possible transcriptional perturbations resulting from our regulatory

modifications. Results show that no transcripts corresponding to the three deleted TFs were
detected in PFC (Figure 5A). By mapping the fold change obtained in the analysis to the
predictions of the computational tool, it is possible to visualize the impact at the
transcriptional level of the missing regulators on their targets (Figure 5C). Four targets
associated to *flhC*, corresponding to genes forming the flagella (*flgB, flgC, flgE* and *flgG*)
were completely silenced; furthermore, all the other flagella-related genes also registered a
decrease on their expression. Regarding *phoB,* two targets were successfully silenced (*phnI*
and *phnL*), both genes belong to an operon that is induced under phosphate starvation and
is required for use of phosphonate and phosphite as phosphorous sources [19], many other
targets related to this operon also reduced their expression. On the contrary, *phnK* present
in the same operon was surprisingly overexpressed. We were unable to map any transcripts
to six genes belonging to the previously mentioned operon, which may not be entirely
expressed in the absence of phosphate starvation. Furthermore, *phoR* (part of the *phoB-*
phoR two-component system) also reduced its expression. Finally, both targets of *cueR*
(copA and *cueO*) also decreased drastically their abundance.
Regarding the accuracy of our ReProMin predictions, 28 genes of 47 predicted silenced
genes were silenced at different levels, whereas 11 predicted silenced genes presented
higher expression values than the WT strain. Finally, transcripts of 8 predicted targets were
not found in either strain. These observations show that in 72% (28 of 39 measured genes)
of the cases the predictions of the computational tool were accurate (Supp. fig. 7).
Additionally to the designed transcriptional changes, we found 17 genes differentially
expressed (8 down regulated genes and 9 up regulated (log2 Fold Change ≥ 1 or ≤ -1 and
p-value ≤ 0.05) (Figure 5B) (Supp. Table 3). This RNA-seq analysis shows that besides the
intended transcriptional changes, few off-target effects were identified at the transcriptomic
level in the PFC strain.

Figure 5 Transcriptomic analysis of the designed strain. A) Correlation plot for PFC and WT strains transcripts. Blue dots represent the three deleted TFs. **B)** Volcano plot showing differential gene expression. In both cases, statistically significant genes are highlighted. **C)** Integration of transcriptomics with computational tool predictions. The size of the circle corresponds to the fold change of each target (the largest circles represent fully silenced genes), in all cases green circles represent targets releasing resources (down regulated), red circles represent targets generating burden (up regulated) and yellow circles targets that were not expressed. High resolution version of the sub network is available in supplementary material (Supp. fig. 6).

Phenotypic evaluation revealed reduced burden, increased cell yield and production budget for designed strains

Our three ReProMin generated mutants (ST case, UT case, and control) were evaluated in rich (LB) as well as in minimal media containing three different carbon sources (acetate, galactose, glucose). The computationally designed strains (PYC and PFC) showed neither

growth defects nor increase in the growth rate or biomass yield in any of the four conditions

tested. On the contrary, the control strain (FOG) showed growth defects in all growth

condition tested (Supp. fig. 8). For glucose minimal media, we also evaluated the effect of

recombinant protein production using a plasmid expressing a genetic circuit with two

fluorescent reporters (Figure 6A) [20]. The burden caused by carrying a plasmid is reflected

as a decrease in the growth rate in all tested strains (Supp. fig. 9A); this decrease is higher

when the plasmid is expressing the genetic circuit, however the burden displayed by both

ReProMin designed strains is lower compared to the wild-type counterpart. Additionally, the

PFC strain also showed a higher final biomass production (Supp. fig. 9B). It has been

described that the expression levels of two protein reporters encoded on the same plasmid

but without a regulatory connection between them is captured by a linear relationship which

can be interpreted as an isocost line, a concept used in microeconomics to describe how

two products can be bought with a limited budget, so the more is used on one, the less can

be used on the other. These lines represent the boundary of the production budget of a

given strain and condition (Figure 6B) [20]. We obtained the isocost lines at balanced growth

(~5 h.) determined by two different methods: mean plate reader fluorescence and mean

fluorescence measured by flow cytometry. The line corresponding to PFC strain show a

parallel upward shift compared to the WT strain, which represents an increase of 9% in

absolute fluorescence ($p < 0.01$) (Figure 6C) and 12% in mean fluorescence per cell ($p <$

0.01) (Supp. fig. 10), this difference is increased at the stationary phase of the culture (~24

h.) were higher maximal biomass is achieved and the quantity of recombinant protein is

increased up to 18% (Figure 6D).

Figure 6 Synthetic circuit characterization. A) Schematic of the gene circuit evaluated, which encodes the fluorescent reporters GFP and RFP: the first is constitutively expressed, while the latter is under the control of an N-acyl-homoserine lactone (AHL) inducible promoter. **B)** When plotting the expression of one protein against the other at different levels of induction, an isocost line is obtained. The size of the area below the line represents the total proteome budget dedicated to the circuit, a parallel upward shift in the line represent an increase in the budget. Isocost lines of the designed strains showing absolute fluorescence during **C)** balanced growth and **D)** stationary phase. Each point represent the red reporter (x axis) plotted against the green reporter (y axis) in an increasing inductor concentration (1.25, 2.5, 5, 10, 20 nM AHL) and represents the mean of three replicates across three different experiments. A linear regression was used to fit the points to a line.

Expression of an engineered metabolic pathway: violacein production.

We tested the ability of our engineered strain for synthesizing the molecule violacein as a proof-of-concept for applications of our method in metabolic engineering. Violacein is a pigment from *Chromobacterium violaceum* endowed with many biological activities (antibacterial, antiviral, anti-parasite) and has recently gained importance in the industrial field especially for applications in cosmetics, medicines and fabrics [21]. Violacein is synthesized in a five-step metabolic pathway using tryptophan as a precursor. Here we used the violacein pathway plasmid reported by Darlington *et al.*, 2018, where the five genes for violacein biosynthesis are arranged in two operons, one consisting of *vioA* constitutively

expressed, while the rest of the pathway encoded by the *vioBCDE* genes is under the control

of an AHL inducible promoter [13] (Figure 7A). This construction follows the same principle as

the previous circuit so the more of one module is produced, the less of the other is expressed

due to the competition for limited resources for gene expression. However, in this case the

number of genes in each module is different and code for actively metabolic enzymes with

different kinetic properties, which results in differential violacein biosynthesis.

We evaluated violacein production after 24 hours in the wild-type and PFC strains using M9

glucose medium supplemented with tryptophan (2.0 g/L) and AHL (1.25, 2.5, 5, 10, 20 nM)

for induction. PFC showed a mean increase in violacein production of 18% (Figure 7B),

additionally we found that the maximum production is achieved with just a minimum quantity

of inducer (1.25 nM) indicating that is crucial to have a balanced expression of the pathway

with the right amount of each module to maximize the synthesis of the final product. Similarly

to our observations on fluorescent protein production, the increase in violacein production

shows that our approach can be harnessed to increase the production of metabolites from

costly heterologous metabolic pathways.

Figure 7 Violacein Pathway Evaluation. A) Schematic of the pathway of violacein biosynthesis. **B)** Total violacein production using 2 g/L tryptophan after 24 h in the presence of increasing inducer concentrations; each value represents the mean of three replicates across three different experiments.

Discussion

Gene regulatory networks are robust and can be severely rewired with interesting phenotypic outcomes[22], thus they are a perfect rational engineering target for synthetic biology applications. In this work we prove that the definition of an essential gene set together with regulatory network information allows the identification of TFs whose elimination leads directly to silencing proteome fractions that are not used in a particular condition. We show that by eliminating hedging proteome activators we can release resources and increase cellular capacity for engineered functions. In agreement with the presented ME-model simulations reducing the unused protein fraction, our designed strain shows higher proteomic budget, measured by the isocost lines, and a higher capacity to produce a metabolite from a heterologous pathway. Furthermore, by comparing our strains with an intuitive control strain, we show that inaccurate TF elimination results in detrimental effects on growth, maximal biomass and protein production (Supp. figs. 9 and 11). These findings indicate that the elimination of a combination of TFs is not a trivial process; it may affect essential functions and introduce phenotypic defects. Our method shows good accuracy in terms of the obtained gene expression changes measured by RNAseq, despite our limited knowledge of the regulatory networks. In addition, the regulatory data available is condition dependent, what limits the predictive power of our method, since we need to assume that regulatory interactions are present at all times. We anticipate that developments in high throughput technologies (such as Chip-seq) combined with novel computational approaches[23–25] will enable the fast generation of complete regulatory networks and the application of our method to even non-model organisms. Several approaches have been applied for resource allocation optimization in bacterial host engineering. Genome minimization has been mainly done by large scale genetic interventions whose outcomes are difficult to predict and do not show greater genome stability [26,27]. Adaptive Laboratory Evolution (ALE) has showed great success, especially to identify functions not related to growth [28], however, it selects for fast growing strains which not necessarily result in the best production phenotypes. Moreover, the underlying selection mechanisms in ALE are normally not known therefore its effects are not predictable [29]. Genome scale models, such as the ME-model, may also be used to find the proteomic cost and fitness benefit of gene expression, thus to aid in the design of proteome allocation, however kinetic data of each protein is needed [30] and its scope focuses on growth related functions. There are only a few reports describing regulatory approaches to improve production phenotypes, such as the global Transcriptional Machinery Engineering (gTME)

[31], but none of them followed a rational approach. The methodology presented in this work

is a novel strategy for proteome optimization with minimal genetic interventions which

overcomes the serious limitations of deleting large regions of the genome; it is a flexible

pipeline which can be applied to other growth and production conditions and also to different

organisms where sufficient information is available. This work shows the potential of rational

design of biological systems over the predominantly used trial and error approaches.

Materials and Methods

1) ME-Model Simulations

All simulations were done using model iJL1678b ME [14]. The corresponding transcription and

translation reactions for recombinant protein (GFP) production were manually added to the

model using standard methods. Unused protein fraction and flux through the recombinant

protein production are changeable variables in the ME-Model that affect predicted growth

rate and proteome composition, the values of these two variables were systematically

changed in the ME-Model to assess their effect on growth rate (UPF = 0.36, 0.30, 0.25; Flux

= 0, 0.001, 0.002, 0.0025, 0.0030, 0.0035, 0.0040), all other model parameters were set as

default. Proteome sectors were classified according to O'Brien *et al.*, 2016.

2) Definition of the essential gene list

To compile the essential gene list in the glucose minimal media condition we combined five

different datasets from different sources. Three of them were experimentally generated

using different methods of gene disruption: A) random transposon mutagenesis using M9

with glucose as growth condition (Tn-seq)[32], B) removing large fragments of the

chromosome using a homologous recombination system in rich medium (LB) [33] and C) the

updated list of the mutants of the Keio collection that are lethal, the collection was generated

using rich medium [34,35]. Two gene lists were generated *in silico* using simulations of genome

scale metabolic and expression models capable of predicting gene expression needs in a

particular condition: D) genes that are essential for growth in M9 with glucose using

iOL1554-ME model [36] and E) genes that are essential for growth in the metabolic model

iJO1366 and also experimentally in M9 with glucose [37]. Within the compiled list, genes

exclusively belonging to the Tn-seq and the glucose minimal media ME-model simulations

gene lists were considered conditionally essential, as these gene lists were originally

generated using M9 with glucose as the growth condition, while the rest of the genes were

classified as core essential for our purposes.

For the cases of galactose and acetate minimal media conditions we performed gene

essentially analysis with the iML1515 model [38] in COBRApy [39].

3) Identification of candidate regulators and combinatorial analysis

We sorted the TF/gene interactions from RegulonDB (version 8, regulondb.ccg.unam.mx),

discarding all the sigma factors-gene interactions. Next, we classified as essential all TFs

that activate at least one essential gene (from our M9 glucose condition gene list) and as
non-essential all TF´s that do not activate any essential genes. Then we analysed the sub-
network of interactions of each non-essential TF by numerically analysing the output level
of each TF (TF_{OUT}), which is classified into positive and negative output (TF_{OUT+} and TF_{OUT-}
) representing positive and negative regulated genes respectively and the degree of entry
of each regulated gene ($GENE_{IN}$) in turn also divided into positive and negative ($GENE_{IN+}$
and $GENE_{IN-}$). We defined as candidate for proteome reduction all those TFs that activate
at least one unique gene, which numerically meet the following condition and represent a
simplified version of a SIM:

$$TF_{OUT+} \geq 1 \quad \wedge \quad GENE_{n_{IN+}} = 1 \quad \wedge \quad GENE_{n_{IN-}} \geq 0$$

439   Where *GENE_n,* represents any gene activated by TF.

On the other hand, the proteomic dataset previously described [17] was used to calculate the
Proteomic Load (P_L) of each gene and Proteomic Balance (P_B) of each TF according to the
equations in Figure 2.
The combinatorial analysis was achieved as follows, given a list of TFs, we created and
tested all the possible N combinations. Next, the total number of silenced and induced genes
for each combination was determined following the next criteria: for every gene involved in
the combination, we subtracted one from the value of $GENE_{IN+}$ for each TF that regulates
the target gene positively and one to the value of $GENE_{IN-}$ for each TF that regulates the
target gene negatively. At the end of this process a gene was considered silenced if:

$$GENE_{IN} = GENE_{IN+} = 0$$

or induced if:

$$GENE_{IN} = GENE_{IN-} = 0$$

Finally, the P_B of each combination tested was calculated and ranked. The full computational
set of tools coded in Python and datasets used in the analysis are available in the following

repository (https://github.com/utrillalab/repromin). Cytoscape software version 3.7 [40] was
used to plot the network representation of the data.

4) Generation of combinatorial knock-out strains
The combinatorial mutants were generated by sequential P1 phage transduction from the
individual knock-out strains of the Keio collection according to the protocol described by
Miller (1992) [41]. The removal of the kanamycin resistance cassette before each transduction
was done using the pE-FLP plasmid (Addgene plasmid #45978), pE-FLP was a gift from
Drew Endy & Keith Shearwinand. Each knock-out strain was confirmed by PCR using
primers flanking each gene. In all experiments *Escherichia coli* BW25113 was used as the
WT background. The characteristics of the strains, plasmids and primers used in this study
are described in the supplementary material (Supp. Table 4).

5) RNA sample extraction and sequencing
Strains were grown in 50 mL of M9 media with glucose (4 g/L) M9 media in 250 mL
Erlenmeyer flasks cultures in an orbital incubator at 37°C (250 rpm). Cells were harvested
in mid-log phase using the Qiagen's RNAprotect bacteria reagent according to the
manufacturer's specifications. Cell pellets were incubated with lysozyme, SuperaseIn and
protease K for 10 min at 37°C. Total RNA was isolated and purified using Zymo Research's
Quick-RNA kit according to the manufacturer's specifications. All samples' quality was
inspected in a bioanalyzer RNA chip (Agilent). Starting with 10ug of total RNA of each
sample, the removal of ribosomal RNA was done with the Ribominus kit by Invitrogen. For
the construction of the libraries, the TruSeq Stranded mRNA Sample Prep Kit by Illumina
was used, following the HT protocol. For sequencing a NextSeq 500 v2 was used, with a
configuration of 2 x 75 paired-end read and 10 million reads per sample.
Reads were mapped to reference genome *E. coli* MG1655 (RefSeq: NC_000913.3) using
aligner Bowtie2 (http://bowtie-bio.sourceforge.net/bowtie2). Final differential analysis was
made using the Cufflinks library (http://cole-trapnell-lab.github.io/cufflinks). Genes with log2
Fold Change ≥ 1 were considered up regulated and ≤ -1 down regulated, considering a p-
value ≤ 0.01.

6) Growth phenotype characterization
For the evaluation of growth in different carbon sources, the following conditions were used:
glucose M9 medium (4 g/L), galactose M9 medium (3.2 g/L), acetate M9 medium (2.5 g/L)

and LB. Cells were cultured overnight in the corresponding media. The next day the strains
were diluted to an OD_{600} of 0.05 in fresh medium and 150 µl of the fresh culture were
transferred to a transparent 96-well plate (Corning) and incubated at 37 °C with fast linear
shaking in a microplate reader (Synergy 2.0, BioTek) for 24 hours, taking measurements for
OD_{600} every 20 min. In all trials three replicates were included and the experiment was
repeated independently on three different days.
The characterization of the growth kinetics was conducted using the algorithm developed by
Swain *et al.*, [42] with the default parameters. In all cases, when comparing the parameters
obtained from the mutant strains against the wild type, the p-value was determined using a
two-tailed Student t-test as the statistical significance reliability index.
503

7) Isocost circuit evaluation
Strains were inoculated into glucose M9 medium with gentamicin (20 ug/ml), and grown
overnight. Next day strains were diluted to an OD_{600} of 0.05 in fresh glucose M9 medium
containing *N*-acyl homoserine lactone (AHL, Sigma-Aldrich, St. Louis, MO, USA, final
concentrations of 1.25, 2.5, 5, 10, 20 nM), then 150 µl of the fresh culture were transferred
to a 96-well black plate with transparent bottom (Corning) and incubated as described
above, taking measurements for OD_{600}, GFP (ex., 485 nm, em., 528 nm) and RFP (ex., 590
nm, em., 645 nm). The characterization of the production kinetics of GFP and RFP was also
done using the algorithm described above.
513

8) Flow cytometry measurements
For flow cytometry measurements, cell cultures were prepared as described above, but later
grown in 24-well plates using 1 ml of medium. Every hour 50 µL aliquots were taken from
each well and mixed with 150 µL of PBS, the volume of the wells was kept constant by
adding fresh medium. Cell suspension was loaded into an Attune NxT Flow Cytometer
(ThermoFisher, Waltham, MA, USA) and analysed for GFP (excitation 488 nm; emission
525/50 nm) and RFP (excitation 561 nm; emission 620/15 nm). For each sample 20,000
events were analysed and population means were estimated using the default software of
the instrument.
523

9) Characterization of violacein-producing strains.
The strains were inoculated into glucose M9 medium with gentamicin (20 ug/ml), and grown
overnight. Next day strains were diluted to an OD_{600} of 0.05 in fresh glucose M9 medium

containing AHL (1.25, 2.5, 5, 10, 20 nM) and tryptophan (0, 0.5, 1.0 and 2.0 g/L), then 150

μl of the fresh culture were transferred to a 96-well plate and incubated as described above.

After 24 h the plate was centrifuged (13,000×g, 10 min), and the supernatant of each well

was discarded. Violacein was extracted by suspending the pellet in each well in 200 μl

absolute ethanol and incubating the plate at 95 °C for 10 min followed by pelleting cell debris

(13,000×g, 10 min). Violacein present in the extract was determined spectrophotometrically

at 575 nm in a microplate reader (Synergy 2.0, BioTek).

Acknowledgements: Elisa Marquez-Zavala and Colton Lloyd for ME-Model simulations

support. Yared Castillo-Franco for computational support and Georgina Hernandez-Chavez

for technical support.

Funding: UNAM–DGAPA-PAPIIT projects IA200716 and IA201518. Newton advanced

Fellowship Project NA 160328. JJ, and JK acknowledge the support received from the

Biology and Biotechnology Research Council (BBSRC) (grant BB/M009769/1) and

European Union's Horizon 2020 research and innovation programme for the project P4SB

(grant agreement no. 633962).

Author contribution: JU and GLP designed ReProMin; GLP developed computational

methods and performed data analysis; GLP, JSMH carried out experiments; GLP, JK and

JIJ analysed flow cytometry experiments, isocost lines and violacein production; JU

supervised the study; JU, GLP and JIJ wrote the manuscript.

Data availability: RNA-seq data from this study have been deposited in NCBI's Gene

Expression Omnibus (GSE 134335). The code to run ReProMin can be found at:

https://github.com/utrillalab/repromin

Conflict of interest: JU and GLP are inventors in a MX patent application filled by UNAM.

References

1.   Hutchison, C. A. *et al.* Design and synthesis of a minimal bacterial genome. *Science
(80-.).* **351**, aad6253–aad6253 (2016).

2.   Balikó, G. *et al.* Rational efforts to streamline the escherichia coli genome. in
Synthetic Biology: Parts, Devices and Applications 49–80 (2018).

3.   Ishihama, A. Prokaryotic genome regulation: Multifactor promoters, multitarget
regulators and hierarchic networks. *FEMS Microbiology Reviews* **34**, 628–645
(2010).

4.   Ulrich, L. E. & Zhulin, I. B. The MiST2 database: a comprehensive genomics
resource on microbial signal transduction. *Nucleic Acids Res.* **38**, D401-7 (2010).

5.   Kitano, H. Biological robustness. *Nat. Rev. Genet.* **5**, 826–837 (2004).

6.   Isalan, M. *et al.* Evolvability and hierarchy in rewired bacterial gene networks.
Nature **452**, 840–845 (2008).

7.   Mitchell, A. *et al.* Adaptive prediction of environmental changes by microorganisms.
Nature **460**, 220–224 (2009).

8.   Tagkopoulos, I., Liu, Y.-C. & Tavazoie, S. Predictive behavior within microbial
genetic networks. *Science* **320**, 1313–1317 (2008).

9.   Utrilla, J. *et al.* Global Rebalancing of Cellular Resources by Pleiotropic Point
Mutations Illustrates a Multi-scale Mechanism of Adaptive Evolution. *Cell Syst.* **2**,
260–71 (2016).

10.   O'Brien, E. J., Utrilla, J. & Palsson, B. O. Quantification and Classification of E. coli
Proteome Utilization and Unused Protein Costs across Environments. *PLoS*
Comput. Biol. **12**, e1004998 (2016).

11.   Nikolados, E.-M., Weiße, A. Y., Ceroni, F. & Oyarzún, D. A. Growth Defects and
Loss-of-Function in Synthetic Gene Circuits. *ACS Synth. Biol.* acssynbio.8b00531
(2019).

12.   Ceroni, F. *et al.* Burden-driven feedback control of gene expression. *Nat. Methods*
(2018).

13.   Darlington, A. P. S., Kim, J., Jiménez, J. I. & Bates, D. G. Dynamic allocation of
orthogonal ribosomes facilitates uncoupling of co-expressed genes. *Nat. Commun.*
9, 695 (2018).

14.   Lloyd, C. J. *et al.* COBRAme: A computational framework for genome-scale models
of metabolism and gene expression. *PLOS Comput. Biol.* **14**, e1006302 (2018).

15.   Santos-Zavaleta, A. *et al.* RegulonDB v 10.5: tackling challenges to unify classic
and high throughput knowledge of gene regulation in E. coli K-12. *Nucleic Acids*
Res. **47**, D212–D220 (2019).

16.   Keseler, I. M. *et al.* EcoCyc: a comprehensive view of Escherichia coli biology.
Nucleic Acids Res. **37**, D464-70 (2009).

17.   Schmidt, A. *et al.* The quantitative and condition-dependent Escherichia coli
proteome. *Nat. Biotechnol.* **34**, 104–110 (2015).

18.   Shen-Orr, S. S., Milo, R., Mangan, S. & Alon, U. Network motifs in the
transcriptional regulation network of Escherichia coli. *Nat. Genet.* **31**, 64–68 (2002).

19.   Yakovleva, G. M., Kim, S. K. & Wanner, B. L. Phosphate-independent expression of
the carbon-phosphorus lyase activity of Escherichia coli. *Appl. Microbiol. Biotechnol.*
49, 573–8 (1998).

20.   Gyorgy, A. *et al.* Isocost Lines Describe the Cellular Economy of Genetic Circuits.
Biophys. J. **109**, 639–46 (2015).

21.   Durán, N. *et al.* Advances in Chromobacterium violaceum and properties of
violacein-Its main secondary metabolite: A review. *Biotechnol. Adv.* **34**, 1030–1045
(2016).

22.   Baumstark, R. *et al.* The propagation of perturbations in rewired bacterial gene
602   networks. *Nat. Commun.* **6**, 10105 (2015).

603   23.   Fang, X. *et al.* Global transcriptional regulatory network for *Escherichia coli* robustly
604   connects gene expression to transcription factor activities. *Proc. Natl. Acad. Sci.*
605   201702581 (2017).

606   24.   Ibarra-Arellano, M. A., Campos-González, A. I., Treviño-Quintanilla, L. G., Tauch, A.
607   & Freyre-González, J. A. Abasy Atlas: a comprehensive inventory of systems,
608   global network properties and systems-level elements across bacteria. *Database*
609   **2016**, baw089 (2016).

610   25.   Sastry, A. V. *et al.* The Escherichia coli Transcriptome Consists of Independently
611   Regulated Modules. *bioRxiv* 620799 (2019).

612   26.   Couto, J. M., McGarrity, A., Russell, J. & Sloan, W. T. The effect of metabolic stress
613   on genome stability of a synthetic biology chassis Escherichia coli K12 strain.
614   *Microb. Cell Fact.* **17**, 8 (2018).

615   27.   Choe, D. *et al.* Adaptive laboratory evolution of a genome-reduced Escherichia coli.
616   *Nat. Commun.* **10**, 935 (2019).

617   28.   Yang, L. *et al.* Principles of proteome allocation are revealed using proteomic data
618   and genome-scale models. *Sci. Rep.* **6**, 36734 (2016).

619   29.   McCloskey, D. *et al.* Evolution of gene knockout strains of E. coli reveal regulatory
620   architectures governed by metabolism. *Nat. Commun.* **9**, 3796 (2018).

621   30.   Nilsson, A., Nielsen, J. & Palsson, B. O. Metabolic Models of Protein Allocation Call
622   for the Kinetome. *Cell Syst.* **5**, 538–541 (2017).

623   31.   Klein-Marcuschamer, D., Santos, C. N. S., Yu, H. & Stephanopoulos, G.
624   Mutagenesis of the bacterial RNA polymerase alpha subunit for improvement of
625   complex phenotypes. *Appl. Environ. Microbiol.* **75**, 2705–11 (2009).

626   32.   Yang, L. *et al.* Systems biology definition of the core proteome of metabolism and
627   expression is consistent with high-throughput data. *Proc. Natl. Acad. Sci.* **112**,
628   10810–10815 (2015).

629   33.   Kato, J.-I. & Hashimoto, M. Construction of consecutive deletions of the Escherichia
630   coli chromosome. *Mol. Syst. Biol.* **3**, 132 (2007).

631   34.   Baba, T. *et al.* Construction of Escherichia coli K-12 in-frame, single-gene knockout
632   mutants: the Keio collection. *Mol. Syst. Biol.* **2**, 2006.0008 (2006).

633   35.   Yamamoto, N. *et al.* Update on the Keio collection of Escherichia coli single-gene
634   deletion mutants. *Mol. Syst. Biol.* **5**, 335 (2009).

635   36.   O'Brien, E. J., Lerman, J. a, Chang, R. L., Hyduke, D. R. & Palsson, B. Ø. Genome-
636   scale models of metabolism and gene expression extend and refine growth
637   phenotype prediction. *Mol. Syst. Biol.* **9**, 693 (2013).

638   37.   Orth, J. D. & Palsson, B. Gap-filling analysis of the iJO1366 Escherichia coli
639   metabolic network reconstruction for discovery of metabolic functions. *BMC Syst*
640   *Biol* **6**, 30 (2012).

641   38.   Monk, J. M. *et al.* iML1515, a knowledgebase that computes Escherichia coli traits.
642   *Nat. Biotechnol.* **35**, 904–908 (2017).

39.   Ebrahim, A., Lerman, J. A., Palsson, B. O. & Hyduke, D. R. COBRApy: COnstraints-
Based Reconstruction and Analysis for Python. *BMC Syst. Biol.* **7**, 74 (2013).

40.   Shannon, P. *et al.* Cytoscape: a software environment for integrated models of
biomolecular interaction networks. *Genome Res.* **13**, 2498–504 (2003).

41.   Miller, J. H. *A short course in bacterial genetics : a laboratory manual and handbook*
for Escherichia coli and related bacteria. (Cold Spring Harbor Laboratory Press,
1992).

42.   Swain, P. S. *et al.* Inferring time derivatives including cell growth rates using
Gaussian processes. *Nat. Commun.* **7**, 13766 (2016).

43.   Pirt, S. J. The Maintenance Energy of Bacteria in Growing Cultures. *Biological*
Sciences **163**, (1965).

Supplementary Figures

Supplementary Figure 1 ME-Model simulations and proteome sector response. The expression of
hedging functions has a proteomic cost that impacts growth and resource availability for the
recombinant sector [9, 10]. Genome scale models of metabolism and gene expression can be used to
analyse the amount of resources devoted to those functions and predict engineering outcomes.
Similar to the maintenance energy coefficient [43], the hedging proteome and other non-growth related
(thus not modelled) functions are accounted for in ME-models as a part of the unused protein fraction
(UPF). This proteome sector is comprised by functions not directly related to growth in the simulated
environment; therefore those functions are not included in the model. The ME-model *i*JL1678b-ME [14]
was used to simulate the effect of the reduction of the UPF and different expression levels of an
unused recombinant model protein (GFP). The simulation shows an increased availability of cellular
resources for recombinant protein production by reducing the UPF.

Supplementary Figure 2. Non-essential TF distribution for the UT case across three growth

conditions tested in this study.

680   **Supplementary Figure 3** Subnetwork of interactions corresponding to the 34 candidate TF
681   representing all the potentially affected targets; green squares represent candidate TF, while light
682   blue circles represent dispensable genes, dark blue circles essential genes and light green squares
683   dispensable TF, circle size is proportional to the P_L of the gene.

688   **Supplementary Figure 4** Regulatory subnetwork of predicted gene targets of the PYC mutant; green
689   circles represent predicted silenced targets, red circles predicted induced targets and yellow circles
690   genes with no proteomic coverage; size of the circles is proportional to the P_L of the target.

Supplementary Figure 5 Regulatory subnetwork of predicted gene targets of the PFC mutant; green circles represent predicted silenced targets, red circles predicted induced targets and yellow circles genes with no proteomic coverage; size of the circles is proportional to the P_L of the target.

702   **Supplementary Figure 6** Integration of transcriptomics with computational tool predictions, the size
703   of the circle corresponds to the fold change of each target (biggest circles represent fully silenced
704   genes), in all cases green represent targets liberating resources (down regulated), red circles
705   represent targets generating burden (up regulated) and yellow circles are targets not expressed.

712
713
Supplementary Figure 7 Accuracy of computational tool predictions (PFC) based on RNAseq data.
Red circles represent wrong predictions, green circles represent accurate predictions and yellow
circles represent unmapped predictions (expression was not detected).

Supplementary Figure 8 Phenotypic evaluation of strains on different carbon source
supplemented M9 media and rich media (LB). (**A-D**) shows max growth rate and (**E-H**)
shows max O.D.

Supplementary Figure 9 Metabolic burden while carrying empty, circuit plasmid and induced circuit plasmid, **A)** shows max growth rate and **B)** shows max O.D.

740   **Supplementary Figure 10** Isocost lines showing mean fluorescence per cell measured by

741   flow cytometry during balanced growth (~5 h).

747   **Supplementary Figure 11** Isocost lines of the FOG mutant compared to the WT strain

748   during balanced growth (~5 h).