Flies exploit predictable perspectives and backgrounds to enhance iridescent signal salience and mating success

Thomas E. White[1,3], Nina Vogel-Ghibely[1], Nathan J. Butterworth[2]

[1]School of Life and Environmental Sciences, The University of Sydney, Sydney, Australia 2106

[2]Centre for Sustainable Ecosystem Solutions, School of Earth, Atmospheric, and Life Sciences, University of Wollongong, Wollongong, NSW, 2522, Australia

[3] Corresponding author.

E-mail: thomas.white@sydney.edu.au

Keywords: communication, structural colour, sexual selection, vision, Muscidae

Word count (excluding abstract): 3766

Number of figures: 4

Number of tables: 3

Supplementary information: Supplementary figure

Abstract

Communication requires both the encoding of information and its effective transmission and reception. The latter need is most acute in 'noisy' environments which set limits on the salience of signals, but little is known about the features of displays that work primarily in the service of efficacy. Here we examined the elaborate courtships of the muscid fly *Lispe cana*, a cursorial species that lives and mates in heterogeneous foreshores. In a field-based assay we quantified the broad structure of male colour signals, display behaviours, and signalling environments, and tested the prediction that males should seek to enhance signal salience — and consequent fitness — through the flexible choice of display locations. We show that courting males gain direct access to the field of view of females by straddling them and holding their wings closed, before presenting their structurally coloured faces in ritualised dances. During these displays males preferentially present their UV-white facial signals against darker visual backgrounds, and the magnitude of signal/background contrast predicts the duration of displays, which in turn predicts mating success. The indirect influence of signal contrast on mating success suggests it captures female attention which, combined with the motion and retinal sweep of the display, serves to enhance signal salience. Our results demonstrate a striking interplay between the physical and attentional manipulation of receivers during sexual communication, and reveal innovative routes to the enhancement of signal efficacy in noisy environments.

Introduction

Communication requires the effective transmission and reception of information in complex natural environments. Selection has favoured diverse solutions to this basic challenge, which are showcased among the visual ornaments and displays of animals (Maia et al. 2013; Girard et al. 2015; White 2018). The functions of traits involved in sexual communication are twofold. Namely; to encode information relevant to mate choice and assessment (i.e. signal content), and ensure its effective transmission and reception (i.e. signal efficacy). Extensive work on content has shown how colour signals and (to a lesser extent) behaviours can encode information on benefits to potential mates, and are the targets of pre-copulatory choice (Brooks and Endler 2001; Kemp 2008a; Kemp 2008b; Barry et al. 2015). This is well illustrated in the literature on pigmentary colour expression and individual condition. The production of carotenoid-based signals, for example, can be physiologically tied to past or present body condition, reproductive and parental quality, and/or immune function, and so offer an honest guide to aspects of mate quality (Weaver et al. 2018). Less is known, however, about the features of displays that act primarily in the service of efficacy.

Most signalling traits are multifunctional (Candolin 2003; Bro-Jørgensen 2010). In addition to encoding information, colour patterns and displays may capture attention (Ord and Stamps 2008), amplify the conspicuousness of other traits (Smith et al. 2009), or modify the content of co-expressed signals (Endler et al. 2014). Traits that serve such purposes are not the target of choice, but are nonetheless key determinants of attractiveness. This is particularly true in heterogeneous environments which will modify, and set limits on, the salience of signals, and the outcomes of selection for efficacy under such conditions are broadly predictable. In evolutionary terms, signal forms that best cut through noise to stimulate receivers will be favoured (Dusenbery 1992). Structural colours, which arise from the selective reflection of light by nano-scale structures, offer one such solution since they are capable of generating uniquely broad, rich, and dynamic colour palettes (Greenewalt et al. 1960; Vigneron and Simonis 2010; White et al. 2012; Maia et al. 2013), and are evolutionarily labile (Maia et al. 2013; Wasik et al. 2014). Of particular interest to this study is their potential for extreme chromaticity and brightness, which are valuable for maximising basic features of efficacy such as signal conspicuousness (Schultz et al. 2008) and detectability (Schultz and Fincke 2009) in the wild. In ecological terms, environmental complexity can drive flexible behavioural solutions that improve the signal-to-noise ratio. These include the precise behavioural delivery of signals (White et al. 2015; Simpson and McGraw 2018), varying the timing and duration of displays (Poesel et al. 2006), and selecting optimal locations for courtship (Endler and Thery 1996).

Flies possess a suite of colour-producing mechanisms and display behaviours and so offer excellent,

albeit underutilised, opportunities for exploring questions of signal evolution (Marshall 2012). The genus *Lispe* is a cursorial group of muscids that spend much of their life on or near the ground in littoral habitats (Pont 2019). This includes courtship, during which males pursue and physically straddle walking females before presenting their iridescent faces and wings in ritualised 'dances' (Frantsevich and Gorb 2006; Pont 2019). Though excellent work continues to document the structure of these and related displays (Spofford and Kurczewski 1985; Frantsevich and Gorb 2006; Jones et al. 2017; Butterworth et al. 2019), it is not known whether or how such colours and behaviours serve to effectively transmit information to mates. Given that the visual structure of seaweed-dominated foreshores will vary over short temporal and spatial scales (e.g. with tides), theory predicts functional links between signal structure and display behaviours to enhance signal salience (Lythgoe 1979; Dusenbery 1992). Specifically, selection should favour male colour traits that are reliably conspicuous to conspecifics in their surrounds, and/or which are delivered via flexible behaviours for exploiting locally optimal conditions.

Here we examine the courtship displays of the fly *Lispe cana* with a view to testing how signal structure and display behaviour mediate communication efficacy and mating success. *Lispe cana* is a species endemic to Australia that inhabits supralittoral (foreshore) habitats spanning the east coast. Males and females have striking structurally-coloured 'white' and 'yellow' faces (Fig. 1), respectively, that are weakly iridescent across lateral viewing angles and more strongly so in the dorseventral plane (unpublished data). They are active predators of other ground-dwelling invertebrates and they live, hunt, and mate on shorelines (ca. 0-5 m from the waterline) which are often populated by patchy distributions of seaweed and detritus (Fig. 1). The casual observation of courtship in this species showed that the smaller males (5.5 - 7 mm body length) approach and 'straddle' larger females (6.5 - 8 mm), before moving to present their faces and wings in ritualised displays (supplementary video). This presents ecologically tractable opportunities to examine the predicted links between signal structure and signalling behaviour, since the initial straddling of mates implies the potential for males to optimise the timing and location of displays. In a field-based assay, we thus quantified the visual structure of signals and signalling environments, and tested the prediction that males should seek to maximise the conspicuousness of their facial signals within their dynamic visual habitats to enhance mating success.

Methods

Sampling courtships, flies, and visual backgrounds

We recorded 42 independent courtships of *L. cana* using a GoPro Hero 6 at either 30 or 60 fps in the

field. From these we extracted the duration of the straddling and display phase of each courtship, which are readily identified by eye. We also estimated the male-female alignment and elevation of straddling males in three haphazardly selected frames from each courtship sequence, which we averaged. Alignment was taken as the angle between by male and female midlines (e.g. with 0 indicating complete alignment) as measured from points on the centre of their heads and abdomens from top-down video. Similarly, male elevation was taken as the angle between male and female midlines from a lateral view. We were unable to extract complete data for nine courtship sequences due to a lack of suitable viewing geometries.

All courtships were recorded within a 6 m^2 region of the supralittoral zone of Toowoon Bay, NSW, Australia, on clear days between 1100 and 1300. Files typically dispersed outside this focal area at the conclusion of a display, minimising the risk of repeated recordings, and the few that did remain were not intentionally observed again. Following each display we noted whether the interaction resulted in copulation, and collected a sample of the nearest piece of background material within 150 mm directly behind the courting male. If no such material was present, we took sand to be the relevant visual background. At the conclusion of the experiment we also gathered all material within the 6 m^2 observation area as a representative sample of the all possible visual backgrounds available to males. From these we collected reflectance measurements at three evenly-spaced points along the length of each piece of material using the general method described below, and averaged them. By systematically collecting and measuring all material within the area of observation, we also approximately accounted for the relative abundance of potential background material in our analyses of background selection and bias (detailed below).

We also collected 17 male and 20 female flies, separate from those observed during courtship, and recorded the reflectance of their facial colouration. We used an OceanOptics Jaz UV-VIS spectrometer with a pulsed xenon light source, set to an integration time of 50 ms with a boxcar width of five. We used a 400 um bifurcated probe oriented normal to the plane of each fly's face to capture the entirety of its ca. 5 mm^2 area, and recorded and averaged two measurements per individual. When measuring background material we instead used a measurement geometry of 0° illumination and 45° collection to minimise specular reflections from damp surfaces. A spectralon 99% diffuse reflector (Labsphere, New Hampshire) and black velvet served as our light and dark standards, respectively, and we recalibrated between each measurement. Finally, we binned the resulting spectra at 1 nm intervals and applied some minor locally weighted scatterplot (LOESS) smoothing ($\alpha = 0.15$) prior to analysis.

Visual modelling

We used an early-stage (retinal) model of muscid colour vision to estimate the subjective colour

and luminance contrasts of male *Lispe* faces against presentation backgrounds. We drew on the receptor sensitivities of *Musca domestica* as the closest available relative of *Lispe*, and assumed the involvement of R7p, R8p, R7y, and R8y photoreceptors in chromatic processing, and R1-6 receptors in achromatic processing (Hardie 1986; Troje 1993). For chromatic contrasts, we drew on the model of Troje (Troje 1993) and estimated receptor quantum catches as the integrated product of receptor sensitivity, stimulus reflectance, and a standard-daylight illuminant. We then calculated the difference in relative stimulation between R7y-R8y and R7p-R8p receptors as putative opponent channels that define the location of a given stimulus in dipteran colourspace (Fig. 1, right side), and chromatic contrasts were taken as the Euclidean distance between stimuli in this space. We estimated luminance contrast as the Weber contrast of fly faces against their backgrounds, with quantum catches calculated as above albeit using the R1-6 receptor absorbance. All spectral processing, analysis, and visual modelling was carried out using the package package 'pavo' v2.2 in R v3.5.2 (Maia and White 2018; R Core Team 2018; Maia et al. 2019).

Statistical analyses

We first tested whether males flies display against visual backgrounds non-randomly with respect to their signal contrast. We used a bootstrap test which, for a given run, entailed drawing 42 backgrounds with replacement from the total pool of background material collected from the observation area, before calculating the mean of the chromatic and achromatic contrasts of the average male face against each item. We repeated this 5000 times, thereby generating a null distribution of contrasts that represent the probability of observing a given mean chromatic or achromatic contrast value under the assumption that males display against backgrounds at random. We then calculated the probability of attaining our observed chromatic and achromatic contrast values given this null expectation, and used Cohen's d (the difference between sample and null means, divided by the pooled standard deviation) as an estimate of the magnitude of any difference between observed and null-distributed contrast values.

We then examined the effects of signal contrast and display duration on mating success using generalised linear models in a restricted maximum likelihood information-theoretic framework (Anderson and Burnham 2004). We specified a global model with mating success as a binomial response (with logit link function), and included display duration, and chromatic and achromatic signal contrast as main effects. We examined all models containing linear combinations of these predictors along with an intercept-only null model, and ranked them according to Akaike's Information Criterion (AICc, adjusted for small sample sizes). We also calculated the R^2 for the most parsimonious models, as approximated by a ΔAICc of < 2 (Anderson and Burnham 2004). We also examined the effect of signal contrasts and straddling duration on the duration of the display phase. We included the display duration as a Gaussian response (with identity link function),

and included chromatic and achromatic signal contrasts and the duration of the straddling phase as main effects. We visually confirmed the assumptions of normality among residuals and homogeneous variance structures for all models, and used the package 'MuMIn' v1.43.6 (Bartoń 2013) for all model selection in R (R Core Team 2018)

Results

During courtship male *L. cana* pursue lone females before straddling them and holding their wings closed with their forelegs. While in this position, males are slightly elevated ($29.4 \pm 7.5^{\mathrm{o}}$) and closely aligned with the body axis of females ($2.1 \pm 1.4^{\mathrm{o}}$), and are afforded direct access to the female's field of view (Fig. 2). This straddling phase lasted 1.58 - 16.53 seconds, after which females were released by males who then repeatedly ran in close semi-circles in front of females as part of a ritualised display (supplementary video; also see Frantsevich and Gorb 2006; Butterworth et al. 2019 for comparable examples). During these display phases which lasted 1.58 - 16.53 seconds, male *L. cana* preferentially presented themselves against backgrounds that generated strong achromatic ($p < 0.001$, d = 3.363) and chromatic (bootstrap p = 0.047, d = 1.744) contrast with their facial colouration (Fig. S1). That is, their faces appear much brighter and slightly more colourful to conspecifics than would be expected if they were displaying at random within their visual environments.

The most parsimonious model of mating success (12/42 courtships) indicated a strong positive contribution of display duration alone (Table 1; Table 2), and was more informative than the second best model that included the additional effect of achromatic contrast (ΔAICc = 0.049, $w_1/w_2 = 1.28$). Both models clearly outperformed the null ($(w_1 + w_2)/w_{\mathrm{null}} > 700$). The individual effect of display duration was much stronger than luminance contrast (Fig. 4; Table 2) and this, taken with the minimal change in log-likelihood and model fit between the first and second top models despite the inclusion of luminance (Table 2), suggest a more direct role for male display duration in mediating mating success. This is further supported by the predictive relationship between achromatic, but not chromatic, signal contrast and display duration itself (Table 3; Fig. 4). The effects of signal/background luminance contrast on mating success is thus almost entirely driven by its influence on male display duration. That is, males presenting brighter signals were able to display longer, which improved mating success. Finally, we found no effect of straddle duration on display duration (Table 4), indicating that the duration of the two courtship phases is unrelated.

Discussion

Selection has generated myriad solutions to the challenge of communicating amidst noise, with animals varying the location (Endler and Thery 1996), timing (Poesel et al. 2006), and behavioural delivery (White 2017; Simpson and McGraw 2018) of signals to ensure their effective reception. Here we reveal novel links between a structural colour signal and its behavioural presentation, which in part relies on the exploitation of the receiver's predictable field of view (Fig. 2). Male *Lispe cana* straddle and align themselves with females during courtship, before releasing and displaying their conspicuously coloured faces in a ritualised dance. Males preferentially display against darker backgrounds which enhances visual contrast (Fig. S1), leading to longer display times (Fig. 4) and, in turn, enhanced mating success (Fig. 3). Our results suggest that males are flexibly exploiting females' perspectives and visual environments to enhance the attractiveness of their complex displays and, hence, mating success.

The courtships of *L. cana* reveal a striking interplay between the physical and attentional manipulation of receivers. When straddling, males physically impede females from flying, and the independence of straddling and display durations suggests that it proceeds until either (the consistently larger) females forcefully disengage males, or males release to display (Table 1). Whether males have any control over the mobility of walking females, or present an encumbrance at all, is unclear, though the near-constant movement of females (pers. observation) means that diverse display locations may be passively sampled within a short time. Once males do release then females are physically free to leave, which necessitates the capturing of female attention during signalling to ensure further receptivity and mating. The predictive relationships between facial contrast and display duration (Fig. 3), and display duration and mating success (Fig. 4), suggest a central role for male signals and presentation behaviours in maintaining female attention. The structurally generated UV-white colour of male faces will, by definition, maximise luminance contrast among the desaturated hues of surrounding seaweed (Fig. 1; though less so among sand), and their flexible display behaviours, combined with predictable knowledge of viewer perspectives, allow for the exploitation of locally optimal conditions. Thus males presenting greater luminance contrast are afforded longer displays — an aspect of courtship that is otherwise under female control — with tangible benefits for mating success.

Several aspects of male displays are likely to further improve salience. For one, the rapid movement of males from outside the female field of view to within it between straddling and display phases will generate strong temporal contrast. This may be amplified by the adaptation of female receptors to visual backgrounds immediately prior to seeing the male display, since the fastest component of receptor adaptation takes less than a second (Smirnakis et al. 1997; Shevell 2001; Baccus and Meister 2002). The continual semi-circular

movement of males during displays will also sweep out a larger area of the female's retina (supplementary video; unpublished data), further enhancing its salience. Analogous display features are described in male great bowerbirds *Ptilonorhynchus nuchalis*, which offer an illustrative comparison. Males construct bowers and exploit the fixed perspective of females to induce a visual illusion, the quality of which is predictive of male mating success (Kelley and Endler 2012). In addition to this focal display males use foraged materials to colour the walls of their avenues and also actively flash coloured objects across the female's field of view. Though not associated with mating success directly, these latter features serve to increase communication efficacy through the capturing of receiver attention and enhancement of other signals (Kelley and Endler 2012; Endler et al. 2014).

The strong effect of display time on mating success suggests that females may be assessing content-rich features of the male display, since the informational load of a task increases decision times (Chernev et al. 2015; Hemingway et al. 2019). Though not directly explored here, the structurally coloured faces of male *L. cana* are theoretically well-suited conduits of information on mate quality. This colouration is primarily a consequence of light scattering by flattened bristles (Frantsevich and Gorb 2006), which demand close developmental control during metamorphosis for optimal expression (Ghiradella and Butler 2009). Variation in the resulting signal may thus inform potential mates about foraging ability, developmental stability, and/or aspects of genetic 'quality' (Kemp 2008b; Barry et al. 2015). The primacy of luminance over chromatic contrast in our tests (Table 1; Table 2) suggests a role for signal brightness and variation therein as the more salient channel. It is also consistent with the relatively under-developed colour sense of flies, which instead tend to draw on luminance to guide the identification and categorisation of stimuli (Troje 1993; Lunau 2014; Lunau 2014). Though luminance is a less reliable cue than hue in natural environments, it is a known channel of sexual communication in several species (Kemp 2008a; Kemp 2008b; Barry et al. 2015). Well-described examples include the mantid *Pseudomantis albofimbriata* in which the brightness of females abdomens is tied to their current condition and is the focus of male choice (Barry et al. 2015), and the butterfly *Eurema hecabe*, wherein the intensity of male UV wing colouration encodes information on larval resource acquisition (Kemp 2008b; Kemp 2008a).

Despite the plausibility of *L. cana*'s faces as informative signals, we found no direct relationship between visual contrast and mating success as might be predicted for traits under selection for such a purpose. There are several reasons for this. For one, the role of any given trait may be to hold attention, as our results show is at least partly the case (Fig. 4), with others acting as more direct indicators of mate quality. The iridescent wing interference patterns of male *L. cana* that are actively presented during displays offer one such candidate trait, and emerging evidence suggests these patterns are both widespread among

flies and may be subject to sexual selection (Katayama et al. 2014; Hawkes et al. 2019). A closely related possibility is that male facial colouration is an amplifier (Hasson 1991; Byers et al. 2010), which serves to make the direct targets of choice easier to discern or assess. Alternately, if male facial colouration is instead a signal of species or sex recognition, then female preferences may be expressed as a simple threshold function. That is, the trait may have to be above a certain value to be effective, but variation beyond that will be irrelevant. Finally, and more specific to this study, is the fact that we averaged facial reflectance across a sample of flies that were not observed in the focal courtships themselves, which prevents us from assessing the direct contribution of individual-level signal variation (in addition to behavioural variation) to mating success.

Environments frequently modify, and set limits on, the salience of sexual displays. Here we describe an innovative solution to this problem that relies on the physical and attentional manipulation of viewers. By adopting the perspective of receivers during courtship, male flies are able to select locally optimal display locations to enhance the salience of their sexual signals and consequent mating success. More broadly, our results suggest that conspicuous signals and flexible display behaviours can arise in response to environmental heterogeneity. Display site properties will vary between habitats, however, which may mediate local adaptation and sexual isolation. This will be most pronounced between habitats that vary acutely in relevant aspects of structural complexity, such as the reliably seaweed-laden foreshore of our focal population, and the entirely barren beaches of nearby populations. This presents intriguing opportunities for illuminating the role of sexual communication in diversification, for which tractable groups such as *Lispe* hold excellent promise.

Acknowledgements

TEW thanks Elizabeth Mulvenna and Cormac White for their support. TEW also thanks Hannah Rowland for the salient observation of parallels between *Lispe* and bowerbirds, which sparked closer consideration of a fly's view of courtship.

Funding

No funding to report.

Data availability

All data and code will be made persistently available upon acceptance.

References

Anderson, D., and K. Burnham. 2004. Model selection and multi-model inference. Second. NY: Springer-Verlag 63.

Baccus, S. A., and M. Meister. 2002. Fast and slow contrast adaptation in retinal circuitry. Neuron 36:909–919.

Barry, K. L., T. E. White, D. N. Rathnayake, S. A. Fabricant, and M. E. Herberstein. 2015. Sexual signals for the colour-blind: Cryptic female mantids signal quality through brightness. Functional Ecology 29:531–539.

Bartoń, K. 2013. {MuMIn}: Multi-model inference, {r} package version 1.9.13.

Bro-Jørgensen, J. 2010. Dynamics of multiple signalling systems: Animal communication in a world in flux. Trends in Ecology & Evolution 25:292–300.

Brooks, R., and J. A. Endler. 2001. Direct and indirect sexual selection and quantitative genetics of male traits in guppies (poecilia reticulata). Evolution 55:1002–1015.

Butterworth, N. J., P. G. Byrne, and J. F. Wallman. 2019. The blow fly waltz: Field and laboratory observations of novel and complex dipteran courtship behavior. Journal of Insect Behavior 1–11.

Byers, J., E. Hebets, and J. Podos. 2010. Female mate choice based upon male motor performance. Animal Behaviour 79:771–778.

Candolin, U. 2003. The use of multiple cues in mate choice. Biological reviews 78:575–595.

Chernev, A., U. Böckenholt, and J. Goodman. 2015. Choice overload: A conceptual review and meta-analysis. Journal of Consumer Psychology 25:333–358.

Dusenbery, D. B. 1992. Sensory ecology: How organisms acquire and respond to information. WH Freeman New York.

Endler, J. A., J. Gaburro, and L. A. Kelley. 2014. Visual effects in great bowerbird sexual displays and their implications for signal design 281:20140235.

Endler, J. A., and M. Thery. 1996. Interacting effects of lek placement, display behavior, ambient light, and color patterns in three neotropical forest-dwelling birds. American Naturalist 421–452.

Frantsevich, L., and S. Gorb. 2006. Courtship dances in the flies of the genus lispe (diptera: Muscidae): From the fly's viewpoint. Archives of Insect Biochemistry and Physiology: Published in Collaboration with the Entomological Society of America 62:26–42.

Ghiradella, H. T., and M. W. Butler. 2009. Many variations on a few themes: A broader look at development of iridescent scales (and feathers). Journal of the Royal Society Interface rsif–2008.

Girard, M. B., D. O. Elias, and M. M. Kasumovic. 2015. Female preference for multi-modal courtship: Multiple signals are important for male mating success in peacock spiders. Proceedings of the Royal Society B: Biological Sciences 282:20152222.

Greenewalt, C. H., W. Brandt, and D. D. Friel. 1960. Iridescent colors of hummingbird feathers. JOSA 50:1005–1013.

Hardie, R. C. 1986. The photoreceptor array of the dipteran retina. Trends in Neurosciences 9:419–423.

Hasson, O. 1991. Sexual displays as amplifiers: Practical examples with an emphasis on feather decorations. Behavioral Ecology 2:189–197.

Hawkes, M., E. Duffy, R. Joag, A. Skeats, J. Radwan, N. Wedell, M. Sharma, et al. 2019. Sexual selection drives the evolution of male wing interference patterns. Proceedings of the Royal Society B 286:20182850.

Hemingway, C. T., A. M. Lea, R. A. Page, and M. J. Ryan. 2019. Effects of information load on response times in frogs and bats: Mate choice vs. Prey choice. Behavioral Ecology and Sociobiology 73:111.

Jones, S. D., P. G. Byrne, and J. F. Wallman. 2017. Exploring the influence of individual courtship behaviors on male mating success in a blow fly. Journal of insect behavior 30:528–543.

Katayama, N., J. K. Abbott, J. Kjærandsen, Y. Takahashi, and E. I. Svensson. 2014. Sexual selection on wing interference patterns in drosophila melanogaster. Proceedings of the National Academy of Sciences 111:15144–15148.

Kelley, L. A., and J. A. Endler. 2012. Illusions promote mating success in great bowerbirds. Science 335:335–338.

Kemp, D. J. 2008a. Female mating biases for bright ultraviolet iridescence in the butterfly eurema hecabe (pieridae). Behav. Ecol. 19:1–8.

Kemp, D. J. 2008b. Resource-mediated condition dependance in sexually dichromatic butterfly wing coloration. Evolution 62:2346–2358.

Lunau, K. 2014. Visual ecology of flies with particular reference to colour vision and colour preferences. Journal of Comparative Physiology A 1–16.

Lythgoe, J. N. 1979. Ecology of vision. Clarendon Press; Oxford University Press.

Maia, R., H. Gruson, J. A. Endler, and T. E. White. 2019. Pavo 2: New tools for the spectral and spatial analysis of colour in r. Methods in Ecology and Evolution 10:1097–1107.

Maia, R., D. R. Rubenstein, and M. D. Shawkey. 2013. Key ornamental innovations facilitate diversification in an avian radiation. Proceedings of the National Academy of Sciences 110:10687–10692.

Maia, R., and T. E. White. 2018. Comparing colors using visual models. Behavioral Ecology 29:649–659.

Marshall, S. A. 2012. Fliesthe natural history & diversity of diptera.

Ord, T. J., and J. A. Stamps. 2008. Alert signals enhance animal communication in "noisy" environments. Proceedings of the National Academy of Sciences 105:18830–18835.

Poesel, A., H. P. Kunc, K. Foerster, A. Johnsen, and B. Kempenaers. 2006. Early birds are sexy: Male age, dawn song and extrapair paternity in blue tits, cyanistes (formerly parus) caeruleus. Animal Behaviour 72:531–538.

Pont, A. C. 2019. Studies on the australian muscidae (diptera). VIII. The genus lispe latreille, 1797. Zootaxa 4557:1–232.

R Core Team. 2018. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.

Schultz, T. D., C. N. Anderson, and L. B. Symes. 2008. The conspicuousness of colour cues in male pond damselflies depends on ambient light and visual system. Animal Behaviour 76:1357–1364.

Schultz, T. D., and O. M. Fincke. 2009. Structural colours create a flashing cue for sexual recognition and male quality in a neotropical giant damselfly. Functional Ecology 23:724–732.

Shevell, S. K. 2001. The time course of chromatic adaptation. Color Research & Application: Endorsed by Inter-Society Color Council, The Colour Group (Great Britain), Canadian Society for Color, Color Science Association of Japan, Dutch Society for the Study of Color, The Swedish Colour Centre Foundation, Colour Society of Australia, Centre Français de la Couleur 26:S170–S173.

Simpson, R. K., and K. J. McGraw. 2018. Two ways to display: Male hummingbirds show different color-display tactics based on sun orientation. Behavioral ecology 29:637–648.

Smirnakis, S. M., M. J. Berry, D. K. Warland, W. Bialek, and M. Meister. 1997. Adaptation of retinal processing to image contrast and spatial scale. Nature 386:69.

Smith, C. L., D. A. Van Dyk, P. W. Taylor, and C. S. Evans. 2009. On the function of an enigmatic ornament: Wattles increase the conspicuousness of visual displays in male fowl. Animal Behaviour 78:1433–1440.

Spofford, M., and F. Kurczewski. 1985. Courtship and mating behavior of phrosinella aurifacies downes (diptera: Sarcophagidae: Miltogramminae). Proceedings-Entomological Society of Washington (USA).

Troje, N. 1993. Spectral categories in the learning behaviour of blowflies. Zeitschrift fur Naturforschung C 48:96–96.

Vigneron, J.-P., and P. Simonis. 2010. Structural colours. In Advances in insect physiology (Vol. 38, pp. 181–218). Elsevier.

Wasik, B. R., S. F. Liew, D. A. Lilien, A. J. Dinwiddie, H. Noh, H. Cao, and A. Monteiro. 2014. Artificial selection for structural color on butterfly wings and comparison with natural evolution. Proceedings of the National Academy of Sciences 111:12109–12114.

Weaver, R. J., E. S. Santos, A. M. Tucker, A. E. Wilson, and G. E. Hill. 2018. Carotenoid metabolism strengthens the link between feather coloration and individual quality. Nature communications 9:73.

White, T. E. 2017. Jewelled spiders manipulate colour-lure geometry to deceive prey. Biology letters 13:20170027.

White, T. E. 2018. Illuminating the evolution of iridescence. Trends in ecology & evolution 33:374–375.

White, T. E., J. Macedonia, D. Birch, J. Dawes, and D. J. Kemp. 2012. The nanoanatomical basis of sexual dimorphism in iridescent butterfly colouration. Australian journal of zoology 60:101–107.

White, T. E., J. Zeil, and D. J. Kemp. 2015. Signal design and courtship presentation coincide for highly biased delivery of an iridescent butterfly mating signal. Evolution 69:14–25.

 Figures and Tables

Figure 1: Male and female *Lispe cana* and the material sampled from their foreshore habitats (left; photo credits NB & TEW), the reflectance spectra of fly faces and background material (centre), and the spectra as represented in a muscid colourspace (right).

Figure 2: During courtship male *Lispe cana* straddle and align themselves (2.1 ± 1.4º offset, 29.4 ± 7.5º elevation) with females, thus effectively sharing a field of view. Males then preferentially display against backgrounds that enhance the luminance contrast of their UV-white faces, which leads to longer display times and enhanced mating success.

Figure 3: The relationship between display duration and mating success in *L. cana*. Circles represent the presence or absence of mounting following courtship, with the line indicating model fit as estimated by a binomial GLM (Table 2).

Figure 4: The relationship between chromatic and achromatic face/background contrast and courtship display duration in male *L. cana*. Solid and dashed lines indicate the best fit ± 95% CI as estimated by a GLM (Table 1).

Table 1: Full model selection table, detailing the relative strength of candidate binomial GLM's for the relationship between mating success and one or more linear combinations of courtship display duration (duration), and the chromatic (ΔS) and achromatic (ΔL) contrast of male facial colouration against their visual backgrounds.

model	d.f.	LL	AICc	ΔAICc	w
duration	2	-14.756	33.8	0.00	0.417
duration + ΔL	3	-13.837	34.3	0.49	0.327
duration + ΔS	3	-14.665	36.0	2.14	0.143
duration + ΔS + ΔL	4	-13.774	36.6	2.81	0.102
ΔL	2	-18.640	41.6	7.77	0.009
ΔS + ΔL	3	-18.634	43.9	10.08	0.003
null (intercept only)	1	-25.127	52.4	18.54	0.000
ΔS	2	-24.850	54.0	20.19	0.000

Table 2: Full results of the leading candidate models of mating success, drawn from a global binomial GLM that included display duration, chromatic contrast (ΔS), and achromatic contrast (ΔL) as predictors (Table x).

model	parameter	estimate	std. err.	z	R^2
mated ~ duration					0.413
	intercept	-5.715	1.689	-3.382	
	duration	0.830	0.267	3.112	
mated ~ duration + ΔL					0.449
	intercept	-6.247	1.808	-3.454	
	duration	0.684	0.283	2.422	
	dL	0.065	0.050	1.293	

Table 3: Results of the GLM estimating the effect of chromatic (dS) and achromatic (dL) face/background contrasts, and the duration of the 'straddling' courtship phase, on the duration of male *L. cana* courtship displays.

	estimate	std. err.	t	p	R^2
intercept	3.00	1.302	2.308	0.027	0.334
ΔS	-0.657	3.813	-0.173	0.864	
ΔL	0.129	0.030	4.204	<0.001	
straddle duration	-0.055	0.086	-0.643	0.524	