BASAL RESISTANCE AGAINST A PATHOGEN IS MORE BENEFICIAL THAN IMMUNE

PRIMING RESPONSES IN FLOUR BEETLES

Arun Prakash[1], Deepa Agashe[1]* & Imroze Khan[2]*

[1]National Centre for Biological Sciences

GKVK Campus, Bellary Road

Bangalore 560065, India

[2]Ashoka University

Plot No. 2, Rajiv Gandhi Education City, National Capital Region

P.O. Rai, Sonepat, Haryana-131029, India

Correspondence

dagashe@ncbs.res.in

imroze.khan@ashoka.edu.in

KEYWORDS

Bacillus thuringiensis, evolutionary costs, immune priming, pathogen selection, *Tribolium castaneum*

ABSTRACT

Insects exhibit various forms of immune responses, including basal resistance to pathogens and a form of immune memory ("priming") that can act within or across generations. The evolutionary drivers of such diverse immune functions remain poorly understood. Previously, we found that in the beetle *Tribolium castaneum*, both resistance and priming evolved as mutually exclusive strategies against the pathogen *Bacillus thuringiensis*. However, since evolved resistance improved survival far more than priming, the evolution of priming in some populations was puzzling. Was resistance more costly in these populations, or did priming provide added benefits? To test this, we revisited our evolved beetles and analyzed the costs and benefits of evolved priming vs. resistance. Surprisingly, resistant beetles increased reproduction after infection, with no measurable costs. In contrast, mounting a priming response reduced offspring early survival, development rate and reproduction. Even added trans-generational survival benefits of evolved priming could not tilt the balance in favor of priming. Hence, resistance is consistently more beneficial than priming; and the evolution and persistence of costly priming rather than resistance remains a mystery. Nevertheless, our work provides the first detailed comparison of the complex fitness consequences of distinct insect immune strategies, opening new questions about their evolutionary dynamics.

INTRODUCTION

Until recently, it was assumed that insect immunity is nonspecific and cannot build immune memory against previously encountered pathogens, since insects lack the immune cells responsible for adaptive immunity in vertebrates (Cooper & Eleftherianos 2017). Now, growing evidence contradicts this traditional view: priming with a sub-lethal exposure to a pathogen protects against a subsequent exposure to the same pathogen. This survival benefit of priming is observed both in later life stages of primed individuals ("within-generation immune priming"; henceforth WGIP), and in their offspring ("trans-generational immune priming"; henceforth TGIP), in a range of insect species (reviewed in Milutinović et al. 2016) including Dipterans (Pham et al. 2007; Ramirez et al. 2015, 2017), Coleopterans (Roth et al. 2009, 2010; Khan et al. 2016), Lepidopterans, and Hymenopterans (Sadd & Schmid-Hempel 2006). Theoretical studies also highlight the importance of priming in reducing infection prevalence and regulating population size, stability and age structure during infection (Tate & Rudolf 2012; Best et al. 2013). Thus, it appears that under pathogen pressure, priming should be selectively favored. Recently, we directly demonstrated this adaptive value of WGIP and showed that it is a distinct immune startegy that can evolve independently of basal pathogen resistance in the flour beetle *Tribolium castaneum* (Khan et al. 2017a). However, a striking result of this study was that although the net survival benefit of evolved resistance was higher than that of priming (80% vs. 50% survival after infection; Khan *et al.* 2017a), resistance against pathogens did not evolve in all populations (**Fig. 1**).

One reason for this observation could be that resistance imposes higher costs than priming. For instance, several studies suggest that resistance is associated with overexpression of fast acting immune responses that impose large physiological costs (e.g. Sadd and Siva-Jothy 2006; Khan et al. 2017b). A general mathematical model predicts that such costs of constitutively expressed basal resistance can be outweighed by its benefit only under frequent lethal pathogenic infections, maximising the population's growth rate (Mayer et al. 2016). However, the cost of resistance may be larger when pathogens are encountered infrequently. This is perhaps one reason why our beetle populations infected with a single large dose of infection every generation evolved priming, whereas resistance could evolve only in populations that were exposed repeatedly to the pathogen (primary exposure with heat-killed bacteria followed by live bacterial infection) (Khan et al. 2017a). However, few studies have actually measured the fitness consequences of evolved resistance, and these were equivocal: while some found significant costs (Ma et al. 2012; Ye et al. 2009), several others did not (Faria et al. 2015; Gupta et al. 2016). Costs of pathogen resistance may also manifest as widespread tradeoffs with other life-history parameters (Reviewed in Sheldon & Verhulst 1996; Lochmiller & Deerenberg 2000; Norris & Evans 2000; Rolff & Siva-Jothy 2003). In contrast, the impact of immune priming on multiple various fitness parameters has only recently been tested: primed mosquitoes

(Contreras-Garduño *et al.* 2014), tobacco hornworms (Trauer & Hilker 2013) and flour beetles (Khan et al.

2019) show reduced fecundity, and primed mealworm beetle mothers produced progeny that develop

slowly (Zanchi et al. 2011) and have reduced competitive ability (Koella & Boete 2016). Although these

experiments tested for correlations between immune priming and fitness related traits, the direct costs of

evolved priming in response to pathogen pressure have not been measured. Hence, it remains unclear

whether a larger cost of evolved resistance could explain the evolution of priming under infrequent

pathogen exposure.

A second possibility of why resistance did not evolve in all populations is that evolved priming may confer

added survival benefits that manifest across generations (i.e. TGIP), enhancing its net fitness impacts and

facilitating its spread in populations. Although no direct experiments have tested whether such trans-

generational benefits evolve simultaneously with WGIP, theory offers some important clues. A model by

Tidbury and coworkers suggests that since TGIP has a lower ability to reduce infection prevalence,

selection should favor WGIP (Tidbury et al. 2012). On the other hand, Tate and Rudolf suggested that the

stage-specific effects of infection are important: TGIP is more beneficial when an infection affects juvenile

stages, whereas WGIP is more effective if adults incur higher infection costs than larvae (Tate & Rudolf

2012). The model also predicts that selection can strongly favor both WGIP and TGIP when the pathogen

affects larvae and adults equally (Tate & Rudolf 2012). Our previous experimental results suggest that this

hypothesis is relevant at least for flour beetles: both WGIP and TGIP were equally beneficial in beetles

infected with the general insect pathogen *Bacillus thuringiensis* (Bt), which imposed similar infection costs

in both life stages (Khan et al. 2016). Although these results represent an interesting correlation, the causal

link between the pathogen's impact on the host and its role in determining relative investment in different

priming responses is not yet confirmed.

Thus, our understanding of the selective pressures and fitness effects that directly impact the evolution of

diverse priming responses vs. basal resistance is incomplete. To fill these gaps, we used previously

described, evolved replicate populations of the red flour beetle *T. castaneum* that were infected in each

generation with Bt, either with or without the opportunity of priming with heat-killed Bt cells (see C, P, PI,

I populations; Khan et al. 2017a). Previously, we had analyzed evolved immune responses of these

populations after 11 generations of evolution (Khan et al. 2017a). Here, we re-tested the same populations

after a further 3 generations of evolution. We first confirmed that populations (I) where unprimed beetles

were injected directly with a high dose of live Bt still retained a strong WGIP response, whereas beetle

populations (PI) that were both primed and infected every generation showed evolved basal resistance.

Subsequently, to disentangle their respective fitness costs and adaptive benefits, we compared the fitness

effects of evolved immune strategies for critical fitness related traits of such as offspring development, early

reproduction and early survival. We also measured the impact of evolved immune functions on beetle lifespan under starvation and normal conditions. Although these traits not directly relevant for our specific selection lines (since the imposed generation time was much shorter than the beetles' expected lifespan), these traits are known predictors of body condition in the wild, and often trade off with immunity (Hoang 2001; Jacot *et al.* 2004). Astonishingly, despite the higher survival benefits, resistance did not impose any costs, contradicting our expectation that it would show strong fitness trade-offs. Instead, we found that the maintenance and deployment of priming was costly, reducing multiple fitness parameters of I beetles. We also found that WGIP in I populations was associated with evolved trans-generational priming (TGIP); but the combined benefit of evolved priming was still lower than that of increased resistance. We were thus unable to explain why priming was favored in I populations. Nevertheless, our present work provides the first systematic analysis of the evolutionary cost and benefit structure influencing parallelly evolved, divergent insect immune responses.

MATERIALS AND METHODS

Experimental evolution

We used laboratory-adapted populations of *T. castaneum* to initiate four distinct selection regimes: control (C; untreated), priming only (P), primed and infected (PI) and infection only (I), each with 4 independent replicate populations (Khan et al. 2017a). In the present study, for logistical reasons, we only analyzed three replicates from each selection regime (C 1, 2 & 4; P 1, 2 & 4; PI 1, 2 & 4; I 1, 2 & 4). On different days, we handled only one replicate population from each selection regime together – e.g. C1, P1, PI1, I1 or C2, P2, PI2, I2 or C4, P4, PI4, I4). The detailed protocol for the experimental evolution is described in Khan et al. (2017a). Briefly, every generation, we first primed 10-day-old virgin P and PI adults from each replicate population with heat-killed bacterial slurry (see supplementary information for priming protocol). Simultaneously, we also pricked virgin C and I beetles with sterile insect Ringer solution (mock priming). Six days later, we challenged individuals from I and PI regimes with live Bt, whereas C and P beetles were pricked with sterile insect ringer solution (mock challenge) (see supplementary information for infection protocol). We thus created two infection regimes where populations were challenged with a high dose of infection, with (PI) or without (I) the opportunity of priming; and two control regimes where beetles were either pricked with Ringer (C) or heat-killed bacteria (P), but never exposed to live infection. Following the priming and infection treatments, we randomly isolated 60 pairs of live virgin males and females from each replicate population and provided them with 300g wheat to mate and oviposit for 5 days to initiate the next generation. After 14 generations of continuous selection, we isolated a subset of individuals from each replicate population to maintain them under relaxed conditions for two generations without priming or

138   infection (unhandled). The relaxed selection is expected to generate standardized experimental beetles with
139   minimum non-genetic parental effects.

Joint assays of evolved priming and resistance, and their impacts on reproduction

We designed our experimental framework to compare survival benefits and reproductive effects of evolved
priming vs. resistance (see **Fig. 2** for experimental design). Besides measuring survival after priming and
infection, we measured female reproductive output both before and after infection. This allowed us to
estimate the direct impact of experimental evolution with pathogens vs. the actual impact of inducing each
type of immune response. Simultaneously, we also tested for the evolution of TGIP, to compare relative
survival and reproductive effects of different priming responses.

To this end, we first collected pupae from each standardized population and isolated them into 96-well
microplate wells with ~0.25g wheat flour, for eclosion. We randomly assigned 10-day old virgin males and
virgin females from each population to one of the following primary exposure treatments: (a) naïve (or
unhandled) (b) primed (injected with heat-killed Bt) and (c) unprimed (i.e. injected with Ringer). After 24
hours of primary exposure, we formed mating pairs using males and females from each population and
treatment combination in 1.5ml micro-centrifuge tubes with 1g of wheat flour (n = 12 mating-pairs per
replicate population per selection regime). We allowed them to mate for 48 hours and then isolated the 12-
day-old females to oviposit for another 48 hours in 5 g whole wheat flour (oviposition plate), whereas males
were returned to 96-well microplates. After oviposition, we also returned the 14-day-old females to 96-well
microplates. Two days later (total six days after primary exposure), we infected males and females with
live Bt. We recorded male survival every 6 hours for 1 day and then every 24 hours for 7 days post-infection
(same as the selection window during experimental evolution; Khan et al 2017a). We tracked female
survival similarly, except that a day later, we again allowed 48-hour oviposition to estimate the impact of
infection and induction of any priming responses on reproduction. Here, we note that since bacterial
infection imposed significant mortality across regimes, the replicate size for our fitness assays was lower
than expected. Although more beetles were alive in PI regime during the experimental window of
reproductive assay, we did not find any significant difference in proportion of live beetles that reproduced
and assayed across different treatments and selection regime (**Table** S1). We also conducted mock
challenge for a subset of unprimed beetles as a procedural control for survival assay, but not for reproductive
output. We did not find any mortality in uninfected beetles within the experimental window of 7 days.

We allowed eggs laid by naïve, unprimed and primed females (both before and after infection) to develop
for 21 days at 34ºC and counted the total number of progeny (mostly pupae). We retained the offspring
from the first round of oviposition (without infection). At this time, most offspring were pupae, and the few

adults we observed had pale body coloration indicating that they were not sexually mature and hence,
unlikely to be mated (Sokoloff 1977). We isolated these pupae and adults in 96-well plates with ~0.2g flour,
to obtain virgin beetles for future assays to measure trans-generational priming and offspring reproduction.
We only included offspring from mothers that produced more than 8 female and 4 male offspring (n= 8-10
mothers/ priming / replicate population/ selection regime), enabling us to sample enough beetles to test for
a correlation between offspring post-infection survival (a proxy of trans-generational priming) and
reproduction of each parental pair, as described below.

After 10 days, we allowed a subset of female offspring from each parental pair (n=4 offspring/ parental
pair/ replicate population/ selection regime) to mate with 10 day old virgin males from standard laboratory
stock population into a single mating pair for 48 hours and then allowed to oviposit as described before.
This procedure enabled us to measure the impact of parental priming on offspring reproductive output
across populations. On day 16, we infected females and then again assayed their reproductive output as
described above. On the same day, we also infected the remaining 16 day old virgin male and female
offspring from each parental pair with live Bt (n= 4 offspring/ sex/ parental pair) and noted their survival
every 6 hours for 2 days and then every 24 hours until all of them were dead. This experimental design not
only allowed us to jointly estimate the survival and reproductive effects of WGIP vs. TGIP for each parental
pair, but also analyze the impact of each immune response relative to evolved resistance. We did not find
any mortality in sham infected offspring within the experimental window.

We calculated the survival benefit of within-generation priming as the estimated hazard ratio of unprimed
infected versus primed infected groups, using Cox proportional hazard survival analysis conducted
separately for males and females from each standardized replicate population (with priming treatment as a
fixed factor). We noted individuals that were still alive at the end of the survival experiment as censored
values. A hazard ratio significantly greater than one indicates higher risk of mortality in the unprimed group
relative to primed individuals; hence, a significant survival benefit of within-generation priming.
Separately, we also estimated the hazard ratio of naïve infected beetles from P, PI or I regime versus naïve
infected C beetles to quantify evolved resistance. A hazard ratio significantly lesser than one indicates lower
risk of mortality, or increased resistance relative to C beetles.

To measure TGIP, we recorded survival of 4 male and 4 female offspring from each parental mating pair
assayed earlier for within-generation priming. We first calculated their mean lifespan as the unit of analysis
and then compared group means using a mixed model ANOVA with selection regime, parental priming
status and offspring sex as fixed factors across replicate populations. We noted that residuals of mean
lifespan data were not normally distributed (verified with Shapiro-Wilk tests). Therefore, we first
transformed the data into their square root values that fit a normal distribution. Since we noted a significant

main effect of replicate population identity, we then separately analyzed selection regimes that were
handled together using a 3-way ANOVA with selection regime, parental priming status and offspring sex
as fixed factors. We tested for pairwise differences between selection regimes and treatments after
correcting for multiple comparisons using Tukey's HSD.

To compare the relative survival benefits of TGIP versus WGIP, we also analyzed group mean male and
female offspring survival data using Cox proportional survival analysis to calculate the estimated hazard
ratio of offspring from unprimed parents versus primed parents. Subsequently, we used non-parametric
Wilcoxon Rank Sum tests compare hazard ratios from TGIP versus WGIP for each population.

We noted that the residuals of pre-infection reproductive output data of both parents and offspring were
non-normally distributed, and could not be transformed to a normal distribution. We therefore used non-
parametric Wilcoxon Rank Sum tests to analyze the impact of selection regime and priming treatment (for
replicate populations of C, P, PI and I that were handled together). We also used Wilcoxon tests to analyze
the impact of bacterial infection on the reproductive output of parents and offspring, separately for each
replicate population across selection regimes and treatments. Since residuals of reproductive output data
after infection were normally distributed, we analyzed these data using a 3-way ANOVA with selection
regime and treatment as fixed factors crossed with replicate populations, providing an overall estimate of
each effect. Further, to disentangle the effects of each type of evolved immune response (TGIP, WGIP and
resistance), we compared reproductive data from each selection regime separately with that of control
beetles. We used Tukey's HSD to test for pairwise differences between selection regimes and treatments,
as described above.

Quantifying development and survival under starvation and with food, in evolved lines

In separate experiments, we measured the direct impacts of evolved priming responses and resistance on
other fitness components of naïve beetles.

(1) <u>Impact on lifespan under starvation and with food:</u> We first isolated 10 day old naïve virgin males and
females from each population in 96-well microplate wells without food (n = 20 beetles/ sex/
population). We noted mortality every 12 hours (10 am & 10pm ±1 hour) for the next 12 days until all
beetles died. In a separate experiment, we similarly distributed naïve virgin females into 96-well
microplates, but with access to food. We noted their survival every 5 days for 95 days to estimate the
long-term survival costs of evolved immune responses. We did not assay males for long-term survival
costs due to logistical challenges. We analysed survival data under starvation and with food for each
replicate population and sex separately, using Cox proportional hazard test with the original selection
regime as a fixed factor.

(2) <u>Quantifying early survival, development and viability costs in evolved lines:</u> We next estimated the
impact of evolved immune responses on aspects of early survival and development. We allowed 12 day
old mated females from each population (n = 60) to oviposit in 150g of doubly sifted flour (using sieves
with pore size of 50μ to remove large flour particles; Diager USA) for 24 hours. We discarded the
females, and isolated 96 randomly chosen eggs into 96-well microplate wells with ~0.2 g flour. This
method is designed to minimize competition during larval development. After 10 days, we sifted the
flour from each microplate to count live larvae and measure egg hatchability. Following this, we
returned the live larvae to 96-well plates and provided fresh flour. In our standard stock beetle
populations, pupation and adult emergence begins around 3-4 weeks after oviposition. Therefore, we
estimated the proportion of pupae and adults after 3 and 4 weeks post-egg collection respectively, as
proxies for time to pupation and adult emergence. We repeated this experiment three times. We did not
assay P beetles due to logistical challenges. We analysed data using a 2-way ANOVA with selection
regime and replicate experiments as fixed factors, and tested for pairwise differences using Tukey's
HSD.
249

RESULTS

Our previous work demonstrated that lethal Bt infection can rapidly select for divergent immune strategies
in PI and I beetles, within 11 generations (Khan et al. 2017a). Populations (I regime) that were directly
infected with a single large dose of Bt evolved within-generation priming, whereas PI populations where
beetles were injected first with heat-killed and then live Bt evolved high resistance. We also found that
resistance provides higher survival benefits than priming, and yet I populations evolved priming instead of
resistance.

Here, we reanalyzed the same beetle populations after 14 generations of experimental evolution to directly
test whether higher costs of evolving resistance could explain this surprising pattern of evolved immune
responses. As observed after 11 generations (Khan et al. 2017a), we found evolved priming responses only
in males and females from I populations (~3-fold increase in their survival relative to control beetles) (**Fig.**
3A & S1, **Table** S2-S3); whereas PI beetles had higher basal resistance (3 to 28-fold increase in the survival
of naïve PI beetles relative to control beetles) (**Fig.** 3A, **Table** S4). We also found that whereas the survival
of I beetles after Bt infection was still 50%, PI beetle survival had increased to ~85% (**Fig.** S2). Replicate
populations from the C or P regimes where beetles were not exposed to live infection did not evolve any
priming ability or higher resistance to infection.

Evolved immune responses do not incur reproductive costs

We first measured the impact of evolved immune responses on beetle reproduction, and found complex
fitness effects that varied substantially with priming type and infection. Evolved priming or resistance had
no impact on the reproduction of naïve unhandled beetles or uninfected beetles pricked with Ringer solution
or heat-killed bacteria (**Fig.** 3B, naïve treatment before infection; **Table** S5). Thus, the maintenance of
priming or resistance does not impose a reproductive cost. However, infection with live pathogen reduced
beetle reproduction in most populations, except PI beetles (with evolved resistance) where the impact of
infection was inconsistent across treatments and replicate populations (**Fig.** 3B, naïve treatment after
infection; **Table** S6). Only a few PI populations showed reduced reproductive output after infection,
whereas others showed no impact (**Table** S6). Overall, the average post-infection reproductive cost of
evolved resistance was lower than that of evolved priming (compare PI vs. I populations in **Fig** 3B, naïve
treatment after infection; **Table** S6).

Subsequently, we analyzed the impact of experimental priming (mimicking selection regimes during
experimental evolution) on the reproductive output of infected beetles. We expected that after infection,
beetles in the priming treatment would reflect reproduction of PI beetles during experimental evolution;
when compared to beetles from the C (control) regime, these data would inform about the impact of evolved
resistance on reproduction. Similarly, after infection, beetles in the unprimed treatment would mimic I
beetles during experimental evolution, and in comparison to C beetles, provide an estimate of the
reproductive cost (or benefit) of evolved WGIP. A mixed model ANOVA followed by separate
comparisons with control beetles (e.g. PI vs. C; I vs. C; P vs. C) revealed main effects of both priming
treatment and original selection regime, as well their interaction, in each case (**Table** S7). Evolved
resistance were beneficial for reproduction, but only in naïve or unprimed beetles (compare PI and P
regimes vs. C regime after infection, **Fig.** 3B). However, experimental priming also increased the
reproduction of C beetles, revealing that I beetles (with evolved priming) pay a relative reproductive cost
compared to PI and C beetles (compare primed beetles after infection, **Fig.** 3B). Overall, this suggests that
I lines (which evolved priming) paid a reproductive cost of their increased survival benefits after mounting
within-generation priming responses; but PI lines (which evolved resistance) could alleviate this
reproductive cost. Thus, evolved resistance is better than priming not only in terms of their survival benefit,
but also in terms of reproduction.

Evolved priming reduces early survival and extends development time

In separate experiments, we tested the direct impacts of evolved priming and resistance on other fitness
traits such as survival under starvation or normal condition and features of early survival such as egg
hatchability and total number of viable offspring at various developmental stages. We also measured the
proportion of pupae and adults at week 3 and 4, as proxies of development rate. An analysis of survival

data under starvation using Cox proportional hazard test (**Table** S8) revealed that males and females across all selection regimes had similar lifespan under starvation (**Fig.** S3). Similarly, we also analyzed long-term survival data of naïve females under normal condition up to 95 days from all the selection treatments. None of the selection treatments had any consistent impact on long-term survival (**Fig** S4, **Table** S9).

In contrast, we found significant effects of selection regime on egg hatchability, total number of viable offspring and proportion of adult offspring at week 4 (but not on the proportion of pupae at week 3) (**Fig.** 4A-D). Since we also observed significant impacts of replicate experiments, we analyzed each replicate experiment separately. In all replicate experiments, we found that the number of viable offspring at week 4 was drastically reduced in beetles from the I regime (**Fig.** 4D, **Table** S10). This is perhaps due to significant early mortality during egg to larval development in I beetles: while ~75% C, P and PI eggs hatched into larvae, only 55% I eggs survived (**Fig.** 4A, **Table** S10). In addition, the proportion of adults at week 4 was lowest in I regime, suggesting delayed development (**Fig.** 4C, **Table** S10). Overall, these results suggest that maintenance of priming imposed considerable costs of reduced early survival and slower development in I beetles. In contrast, evolved basal resistance did not appear to impose a substantial cost with respect to these traits.

Evolved within-generation priming (WGIP) is associated with trans-generational priming (TGIP)

Finally, we asked whether evolved priming conferred added trans-generational benefits, increasing its overall fitness impacts. To do this, we used a mixed model ANOVA (randomized across replicate populations) to analyze the mean post-infection survival of offspring from beetles assayed above as a function of selection regime, parental priming status and offspring sex (**Table** S11). Both selection regime and parental priming status had significant impacts, but offspring sex did not affect survival. Here too, we found that overall, offspring of PI beetles had the highest survival, though they did not show effects of parental priming. In contrast, parental priming increased offspring survival in the I regime, suggesting that TGIP benefits are solely restricted to I beetles. Since we also observed a significant impact of replicate population identity, we next separately analyzed selection regimes that were handled together (**Table** S12). Parental priming increased female offspring's post-infection survival in all I populations (I1, I2 & I4), whereas male offspring had longer lifespan only in replication populations I1 and I2 (**Fig.** 5A). Male offspring from primed I4 parents also appeared to survive longer than offspring of unprimed parents, but the difference was not statistically significant (P>0.05). We also tested whether the relative survival benefits of TGIP were equal to that of WGIP. We used Cox proportional hazard analysis of the grouped mean offspring survival data for each parental mating pair from I populations, and calculated the strength of evolved TGIP as the estimated hazard ratio for offspring from unprimed vs. primed parents. We found a significant TGIP response in offspring from replicate populations I1 and I2 (**Table** S13). In contrast, primed

and unprimed offspring from replicate population 4 had similar survival. Interestingly, the survival benefit

of TGIP and WGIP was also similar across replicate populations (p>0.05; **Fig**. S5A, **Table** S14), supporting

the hypothesis that Bt-imposed selection favors the evolution of both types of priming to a similar extent

(**Fig**. S5B, **Table** S13).

As found with mothers (above), evolved priming and resistance did not consistently affect the reproductive

output of naïve or uninfected offspring (**Fig** 5B, **Table** S14), but infection generally reduced offspring

reproductive output in all selection regimes except PI beetles (**Table** S15). A full factorial mixed model

ANOVA revealed significant main effects of only selection regime, whereas priming and replicate

populations had no impact (**Table** S16). Offspring of PI beetles again reproduced more than other beetles,

regardless of their parental priming status; whereas TGIP had no impact on the reproduction of I offspring.

Overall, it is surprising that although multiple forms of priming jointly evolved in I populations, their

combined effects were still not as high as resistance, and I beetles (without priming) were still highly

susceptible to infection, suffering a large relative fitness loss each generation.

DISCUSSION

Previously, we showed that priming and resistance against *B. thuringiensis* infection evolve as mutually

exclusive strategies in flour beetles (Khan et al. 2017a). However, since evolved resistance conferred a

greater survival benefit than priming, it was puzzling why some populations evolved priming instead of

resistance. We had speculated that resistance might incur hidden fitness costs that we had not been able to

measure. Here, we revisited our beetle lines to systematically test this hypothesis. Conversely, we also

asked whether priming confers additional, trans-generational fitness benefits that may facilitate its fixation.

To our surprise, we did not find any evidence for a cost of evolved resistance: it did not impact development,

reproduction, or survival during starvation and normal conditions, contradicting the traditional view of

immunity-fitness trade-offs (Ye *et al.* 2009; Ma *et al.* 2012). Instead, our data add to the growing body of

work that suggest only a weak role for life-history trade-offs during the evolution of pathogen resistance

(Faria *et al.* 2015; Gupta *et al.* 2016). Interestingly, we also found that WGIP (within-generation immune

priming) was associated with the evolution of TGIP (trans-generation immune priming) in females from all

replicate populations, and in males from two of the three replicate populations that we tested. However, the

combined benefit of these two forms of priming (~50% survival after Bt infection) was still lower than that

conferred by increased baseline resistance to Bt (~85% survival). Hence, the peculiar patterns of the

evolution of various immune responses remain a mystery.

Most surprisingly, we found that although infection reduced reproduction in all regimes, the effect was less

pronounced in PI beetles (which had evolved increased resistance), and hence, evolved basal resistance was

also associated with a relative reproductive advantage. Interestingly, P (priming only) beetles also had
higher reproduction than control beetles after infection, which is counterintuitive because these beetles
never experienced live infection during experimental evolution. Note that this relative reproductive
advantage would be important during experimental evolution, since beetles reproduced for 5 days after
infection in each generation (see methods). How do we interpret these apparent reproductive fitness benefits
in PI and P beetles? First, the reduced cost of infection in these beetles might represent evolved tolerance,
whereby beetles do not invest in directly clearing pathogens via canonical resistance mechanisms, allowing
greater reproductive investment during an infection (Ayres and Schneider 2012). Second, these results
could reflect a trade-off between early vs. late reproduction. In other words, increased reproduction might
represent terminal investment in P and PI populations, whereas C and I populations instead suppress
immediate reproduction after infection to maintain survival and somatic maintenance later in life (Luu and
Tate 2017). Although we could not test these hypotheses here, our results suggest that divergent immune
responses can have important consequences for reproductive success, and deserve further attention.

Our results also contradict our prior hypothesis that at a low pathogen frequency (experienced by I beetles),
priming may be more favorable than resistance due to its low maintenance costs (Khan et al. 2017a).
Instead, we found that overall maintenance of priming responses is costly. Although evolved priming did
not affect lifespan or survival under starvation, it directly reduced egg hatchability, offspring viability and
development rate in naïve I beetles compared to control beetles. However, priming had variable effects on
reproduction. For instance, mounting a within-generation priming response helped C beetles to increase
their reproduction after infection; whereas infected I beetles, despite evolving survival benefits, could not
improve their reproduction. These results mirror our recent observations with wild-caught populations,
where primed and infected females with increased post-infection lifespan produced fewer offspring (Khan
et al. 2019) and vice versa. We thus speculate that a hidden trade-off with reproduction might constrain the
survival benefits of within-generation priming responses at a much lower level than resistance. Mounting
trans-generational priming responses, on the other hand, had no effect on offspring reproduction, suggesting
that fitness effects are not uniform across different priming responses. Our results broadly corroborate other
work showing the negative effects of priming on various fitness parameters (Trauer & Hilker 2013;
Contreras-Garduño *et al.* 2014). However, these studies primarily used phenotypic manipulations within a
single generation, whereas ours is the first study to directly measure the complex fitness costs associated
with evolved priming across multiple generations of pathogen exposure.

Our experiments provide the first empirical evidence that insects can evolve multiple priming responses
simultaneously. Interestingly, both transgenerational and within-generation priming provided almost
equivalent fitness benefits, corroborating our prior work showing similar benefits of WGIP and TGIP across

10 distinct wild-caught beetle populations (Khan et al. 2016). Such parallel results from natural and

laboratory-evolved populations indicate that pathogens such as Bt may serve as a potent source of selection

favoring the evolution of diverse immune responses in insects. As discussed earlier, Bt reduces the survival

of flour beetle larvae and adults equally (Khan et al 2016), which should favor the simultaneous evolution

of WGIP and TGIP (Tate and Rudolf 2012). However, during experimental evolution we only infected

adult beetles, which should have restricted host-pathogen interaction to adults. It is possible that infected

adults directly transmitted Bt to eggs, imposing selection favoring TGIP. Alternatively, infected adults

could have transmitted Bt (or antigen) to larvae via the flour, either through infected beetle cadavers (~10-

15% mortality during oviposition period in I beetles) or excreta (Argôlo-filho & Loguercio 2014). Another

possibility is that ancestral beetle populations may have already coevolved with Bt in their natural habitat

before they were brought into the lab. Consequently, despite being infected only as adults during

experimental evolution, the beetle immune system could perhaps readily recognize Bt as a risk across life

stages, due to their shared evolutionary history. Finally, if WGIP and TGIP involve shared molecular

pathways, direct pathogen pressure on adults could result in simultaneous evolution of both types of

priming. While the molecular details responsible for immune priming are still unclear (Cooper &

Eleftherianos 2017), recent data hint at shared immune pathways between different priming types. For

instance, both within- (Pham et al. 2007) and trans-generationally primed honeybees (Barribeau et al. 2016)

show increased expression of Toll signaling pathways. Further experiments to carefully compare the

molecules underlying different immune responses can help distinguish between the above hypotheses.

In closing, we note that the relative importance of priming vs. general resistance has long been debated,

primarily because it was unclear whether (a) diverse priming types (within- vs. trans-generational) together

constitute distinct strategies, separate from basal resistance (b) their costs vs. benefits differ substantially,

and (c) they involve different or overlapping sets of immune pathways. Our work represents one of the first

steps to address the first two problems, demonstrating distinct costs and benefits of multiple priming

responses vs. resistance evolving simultaneously in response to selection imposed by the same pathogen

(also see Khan et al. 2017a). While these results highlight the remarkable diversity and flexibility of insect

innate immune adaptation against infections, they also suggest that the early survival vs. reproductive costs

of priming can constrain their adaptive evolution, much more so than resistance. However, it remains a

mystery why putative resistance alleles either did not arise or failed to outcompete putative priming alleles,

despite their large selective advantage in I beetles. We hope that our results will motivate further

experiments to address this problem. Specifically, we look forward to detailed mechanistic studies to test

whether host-pathogen interactions at low frequency of infection not only favor the evolution of priming,

but involve immune pathways that mechanistically preclude more beneficial resistance alleles from fixing

in host populations.

AUTHOR CONTRIBUTIONS

IK conceived experiments; IK, AP and DA designed experiments; AP carried out experiments; IK and AP analyzed data; IK and DA acquired funding; IK and DA wrote the manuscript with inputs from AP. All authors gave final approval for publication.

ACKNOWLEDGEMENTS

We are grateful to Devshuvam Banerjee, Laasya Samhita, Srijan Seal and Shaym Budh for feedback on the manuscript. We thank Kunal Ankola, Sunidhi Thakur and Shyam Buddh for their help during experiments.

FUNDING

We acknowledge funding and support from Ashoka University and the National Centre for Biological Sciences, India.

COMPETING INTERESTS

We have no competing interests

REFERENCES

Argôlo-filho, R. C., and L. L. Loguercio. 2014. *Bacillus thuringiensis* is an environmental pathogen and host-specificity has developed as an adaptation to human-generated ecological niches. Insects. 5:62–91.

Ayres, J. S., and D. S. Schneider. 2012. Tolerance of infections. Annu. Rev. Immunol. 30:271-294.

Barribeau, S. M., P. Schmid-Hempel, and B. M. Sadd. 2016. Royal decree: gene expression in transgenerationally immune primed bumblebee workers mimics a primary immune response. PLoS one. 1(7):p.e0159635.

Best, A., H. Tidbury, A. White, and M. Boots. 2013. The evolutionary dynamics of within-generation immune priming in invertebrate hosts. J. R. Soc. Interface. 10:20120887.

Contreras-Garduño, J., M. C. Rodríguez, M. H. Rodríguez, A. Alvarado-Delgado, and Lanz-H. Mendoza. 2014. Cost of immune priming within generations: Trade-off between infection and reproduction. Microbes Infect. 16:261–267.

Cooper, D., and I. Eleftherianos. 2017. Memory and specificity in the insect immune system: Current perspectives and future challenges. Front. Immunol. 8:539

Faria, G., N. E. Martins, and S. Magalh. 2015. Evolution of *Drosophila* resistance against different

pathogens and infection routes. Evolution. 69:2799–2809.

Gupta, V., S. Venkatesan, M. Chatterjee, Z. A. Syed, V. Nivsarkar, amd N. G. Prasad. 2016. No apparent
cost of evolved immune response in *Drosophila melanogaster*. Evolution. 70:934–943.

Hoang, A. 2001. Immune Response To Parasitism Reduces Resistance of *Drosophila Melanogaster*.
Evolution. 55:2353–2358.

Jacot, A., H. Scheuber, and M. W. G. Brinkhof. 2004. Costs of an induced immune response on sexual
display and longevity in field crickets. Evolution. 58:2280–2286.

Khan, I., A. Prakash, and D. Agashe. 2016. Divergent immune priming responses across flour beetle life
stages and populations. Ecol. Evol. 6:7847–7855.

Khan, I., A. Prakash, and D. Agashe. 2017a. Experimental evolution of insect immune memory versus
pathogen resistance. Proc. Biol. Sci. 284:20171583.

Khan, I., D. Agashe, and J. Rolff. 2017b. Early-life inflammation, immune response and ageing. Proc. Biol.
Sci. 284(1850):20170125.

Khan, I., A. Prakash, and D. Agashe. 2019. Pathogen susceptibility and fitness costs explain variation in
immune priming across natural populations of flour beetles. J. Anim. Ecol,
https://doi.org/10.1111/1365-2656.13030.

Koella, J. C., and C. Boete. 2016. A genetic correlation between age at pupation and melanization immune
response of the yellow fever mosquito *Aedes aegypti*. Evolutio. 56:1074–1079.

Lochmiller, R. L., and C. Deerenberg. 2000. Trade-Offs in Evolutionary Immunology: Just What Is the
Cost of Immunity? Oikos. 88:87–98.

Luu, H., and A. T. Tate. 2017. Recovery and immune priming modulate the evolutionary trajectory of
infection-induced reproductive strategies. J. Evol. Biol. 30:1748–1762.

Ma, J., A. K. Benson, S. D. Kachman, Z. Hu, and L. G. Harshman. 2012. *Drosophila melanogaster* selection
for survival of *Bacillus cereus* infection: Life history trait indirect responses. Int. J. Evol. Biol.
2012:935970.

Mayer A., T. Mora, O. Rivoire, and A. M. Walczak. 2016 Diversity of immune strategies explained by
adaptation to pathogen statistics. Proc. Natl. Acad. Sci. 113:8630-8635.

Milutinović, B., Peuß, R., Ferro, K. & Kurtz, J. 2016. Immune priming in arthropods: an update focusing
on the red flour beetle. Zoology. 119: 254-261.

Norris, K., and M. R. Evans. 2000. Ecological immunology : life history trade-offs and immune defense in birds. Behav. Ecol. 1:19–26.

Pham, L.N., M. S. Dionne, M. Shirasu-Hiza, and D. S. Schneider. 2007. A specific primed immune response in *Drosophila* is dependent on phagocytes. PLoS Pathog. 3:e26.

Ramirez, J. L., G. de Almeida Oliveira, E. Calvo, J. Dalli, R. A. Colas, C. N. Serhan, *et al.* 2015. A mosquito lipoxin/lipocalin complex mediates innate immune priming in *Anopheles gambiae.* Nat. Commun. 6:7403.

Ramirez, J. L., A. B. F.Barletta, and C. V. Barillas-Mury. 2017. Molecular mechanisms mediating immune priming in *Anopheles gambiae* mosquitoes. Arth. Vec. Cont. Dis. Transm. Vol. 1:pp91-100. Elsevier Inc.

Reaney, L. T., and R. J. Knell. 2009. Immune activation but not male quality affects female current reproductive investment in a dung beetle. Behav. Ecol. 21:1367-1372.

Rolff, J., and M. T. Siva-Jothy. 2003. Invertebrate ecological immunology. Science. 301:472–5.

Roth, O., G. Joop, H. Eggert, J. Hilbert, J. Daniel, P. Schmid-Hempel, *et al.* 2010. Paternally derived immune priming for offspring in the red flour beetle, Tribolium castaneum. J. Anim. Ecol. 79:403–13.

Roth, O., B. M. Sadd, P. Schmid-Hempel, and J. Kurtz. 2009. Strain-specific priming of resistance in the red flour beetle, Tribolium castaneum. Proc. Biol. Sci. 276:145–51.

Sadd, B.M., and P. Schmid-Hempel. 2006. Insect immunity shows specificity in protection upon secondary pathogen exposure. Curr. Biol. 16:1206–10.

Sadd, B.M., and M. Siva-Jothy. 2006. physiological costs of immunity

Sheldon, B. C., and S. Verhulst. 1996. Ecological immunology - costly parasite defenses and trade- offs in evolutionary ecology. Trends Ecol. Evol. 11:317–321.

Sokoloff A. 1977. The biology of *Tribolium* with special emphasis on genetic aspects. Volume 3. Oxford, UK: Clarendon.

Tate, A.T., and V. H. W. Rudolf. 2012. Impact of life stage specific immune priming on invertebrate disease dynamics. Oikos. 121:1083–1092.

Tidbury, H. J., A. Best, and M. Boots. 2012. The epidemiological consequences of immune priming. Proc. Biol. Sci. 279:4505–12.

Trauer, U., and M. Hilker. 2013. Parental legacy in insects : Variation of transgenerational immune priming

during offspring development. PLoS One. 8:e63392

Ye, Y. H., S. F. Chenoweth, and E. A. McGraw. 2009. Effective but costly, evolved mechanisms of defense

against a virulent opportunistic pathogen in *Drosophila melanogaster*. PLoS Pathog. 5:e1000385.

Zanchi, C., J. Troussard, and G. Martinaud. 2011. Differential expression and costs between maternally and

paternally derived immune priming for offspring in an insect. J. Anim. Ecol. 1174–1183.

FIGURES

Figure 1. Summary of the design and outcome of experimental evolution of *Tribolium castaneum* flour
beetles against the bacterial pathogen *Bacillus thuringiensis*, previously described in Khan et al 2017a. The
schematic indicates beetle survival before and after 11 generations of experimental evolution, as well as the
evolved immune response (resistance or priming) observed in all populations of each regime. Every
generation, 10-day-old virgin beetles were either injected with heat-killed bacterial slurry (P & PI) or sterile
insect Ringer solution (C & I) (primary exposure). After six days, individuals from I and PI regimes were
challenged with live Bt, whereas C and P beetles were pricked with sterile insect ringer solution (secondary
exposure). Each selection regime included 4 independent replicate populations. In the current study, we
analyzed 3 replicate populations from each regime.

542 **Figure 2.** Design of joint experiments to assay evolved immune responses and their impacts on beetle

543 reproduction.

Figure 3. (A) Survival curves for within-generation priming and resistance in females (n= 12 females/treatment/selection regime/replicate population) after 14 generations of selection. Asterisks and the numbers in parentheses for I beetles denote the hazard ratios calculated from survival curves for priming that are significantly greater than 1 (p<0.05; a greater hazard ratio indicates higher benefit of priming) (B) Impact of evolved within-generation priming (WGIP) and resistance on female reproductive output, both before (n=12 females/treatment/selection regime/replicate population) and after bacterial infection (n=5-11 females/treatment/selection regime/replicate population). Alphabets indicate significant changes in C beetles' post-infection reproduction after mounting a within-generation priming response.

Figure 4. Impact of evolved immune responses on (A) total number of eggs that hatched into larvae (egg
hatchability); proportion of (B) pupae at week 3 and (C) adults at week 4 as proxies for developmental rate;
(D) total number of viable offspring, including larvae, pupae and adults, at week 4; (n=3 females/selection
regime/replicate experiment). P values for the impact of selection regime are reported in each panel.
Significantly different groups are indicated by distinct alphabets, based on Tukey's HSD.

Figure 5. (A) Offspring survival after trans-generational immune priming (TGIP) and infection (group mean survival of 4 offspring from 8-11 parental pairs/ treatment/ selection regime/ offspring sex). TGIP increased offspring survival only in I regime, indicated by distinct alphabets, based on Tukey's HSD. Asterisks indicate significant increase in post-infection survival (resistance) of naïve PI beetles compared to naïve C beetles. (B) Impact of evolved trans-generational priming on offspring's reproductive output, both with and without infection, for replicate populations that were handled together (group mean survival of 2-4 offspring from 8-11 parental pair/ treatment/ selection regime/ offspring sex). P values for the impact of selection regime on post-infection reproductive output are reported in each panel.

571